**Sexual imprinting and speciation in two *Peromyscus* species**

E.K. Delaney[1,2] and H.E. Hoekstra[1]

[1]Department of Organismic and Evolutionary Biology, Department of Molecular and Cellular

Biology, Museum of Comparative Zoology, Howard Hughes Medical Institute, Cambridge MA

02138 USA   Email: hoekstra@oeb.harvard.edu

[2]Current address: Department of Evolution and Ecology, University of California, Davis CA

95616 USA

KEYWORDS: deer mice, mate choice, mosaic sympatry, reproductive isolation, sexual isolation

**ABSTRACT**

Sexual isolation, a reproductive barrier, can prevent interbreeding between diverging

populations or species. Sexual isolation can have a clear genetic basis; however, it may also

result from learned mate preferences that form via sexual imprinting. Here, we demonstrate that

two sympatric sister species of mice—the white-footed mouse (*Peromyscus leucopus*) and its

closest relative, the cotton mouse (*P. gossypinus*)—hybridize only rarely in the wild despite co-

occurring in the same habitat and lack of any measurable intrinsic postzygotic barriers in

laboratory crosses. We present evidence that strong conspecific mate preferences in each species

form significant sexual isolation. We find that these mating preferences are learned in one

species but may be genetic in the other: *P. gossypinus* sexually imprints on its parents, but innate

biases or social learning affects mating preferences in *P. leucopus*. Our study demonstrates that

sexually imprinting contributes to reproductive isolation that reduces hybridization between

otherwise inter-fertile species, supporting a previously underappreciated role for learning in

mammalian speciation.

## INTRODUCTION

Sexual isolation, when divergent mating preferences limit or prevent interbreeding between individuals from different populations, is a prevalent premating reproductive barrier that may be vital to early stages of speciation. Comparative studies in fruit flies (Coyne and Orr 1989, 1997) and darter fish (Mendelson 2003) have shown that sexual isolation accumulates rapidly among young species relative to postzygotic reproductive barriers (e.g., hybrid sterility and inviability). Sexual isolation also can significantly reduce interbreeding among incipient sympatric species pairs (Coyne and Orr 1997; Noor 1997; Ramsey et al. 2003; Boughman et al. 2005; Nosil 2007; Matsubayashi and Katakura 2009), suggesting that it either allows species to co-exist or is selected via the process of reinforcement. In a few cases, sexual isolation is the sole reproductive barrier preventing hybridization between sympatric species, indicating that sexual isolation alone can be strong enough to inhibit hybridization and maintain genetic differentiation (e.g., Seehausen 1997; Fisher et al. 2006). Yet, despite the important role that sexual isolation can play among incipient species, its mechanistic basis—whether mating preference is genetic or learned—is often unknown.

Sexual isolation forms when mating traits and preferences diverge among populations, suggesting that sexual isolation should evolve when genetic loci for mating traits and preferences become linked (Felsenstein 1981). This scenario is a "two allele mechanism" of reproductive isolation, in which divergent alleles for trait and preference loci must fix between a pair of populations. As this mode of sexual isolation can break down due to recombination, linkage disequilibrium (particularly when caused by physical linkage) can promote sexual reproductive barriers. For example, plumage and plumage preferences are both sex-linked in finches and flycatchers and thus reduce hybridization between alternate color morphs (Pryke 2010) and

congener species (Saether et al. 2007), respectively. Alternatively, the issue of recombination

could be circumvented if the loci for mating traits and preferences were under the same genetic

control (i.e., pleiotropy). Pleiotropic genes undergoing divergent natural selection can also cause

non-random mating and thus behave as "magic traits". Although several examples of magic traits

(e.g., Jiggins et al. 2001; Mckinnon et al. 2004; Bradshaw & Schemske 2003) suggest they may

not be as rare as previously thought (Servedio et al. 2011), the putative "magic" genes remain

elusive. In either of these scenarios, a minimum of two alleles must be present and fixed in each

population, requiring selection or drift in opposite directions in each species (Felsenstein 1981).

Sexual imprinting, the process of learning to prefer parental traits at a young age, is an

alternate mechanism for establishing sexual isolation that arguably can be more efficient at

establishing reproductive isolation than the above-mentioned genetic mechanisms. Sexual

imprinting is immune to genetic recombination because learned mating preferences are

automatically "inherited" with a given trait locus. Thus, sexual imprinting is considered a "one-

allele mechanism" of reproductive isolation because the same "sexual imprinting allele" (e.g., an

ability to learn parental traits) could result in assortative mating that reduces interbreeding

between divergent populations, thereby reducing the number of steps required to achieve sexual

isolation. An example of one-allele assortative mating locus has been localized in the *Drosophila*

genome (Ortíz-Barrientos and Noor 2005). In addition to needing only one allele at a learning

locus, several theoretical models have shown that learned mating preferences will maintain

sexual isolation much longer in populations experiencing gene flow than if mating preferences

had a genetic basis (Laland 1994; Verzijden et al. 2005). Often, sexual imprinting will maintain

reproductive isolation in the face of gene flow because it lowers the amount of divergent natural

selection needed to isolate groups (Verzijden et al. 2005). Sexual imprinting may also enhance

sexual isolation in sympatry through reinforcement (Servedio et al. 2009). Finally, if sexual

imprinting is asymmetric—cases in which offspring prefer even more extreme versions of the

traits on which they imprinted—it may even facilitate divergence by creating a bias and thereby

selection for more divergent mating traits (ten Cate et al. 2006).

While sexual imprinting has long been recognized as a phenomenon that occurs within

species, its impact on speciation has become appreciated only recently (Irwin and Price 1999). It

is a phenomenon that occurs in species with parental care, and has been documented in over 15

orders of birds (ten Cate and Vos 1999) as well as some mammals (Kendrick et al. 1998) and

fish (Verzijden and ten Cate 2007; Kozak and Boughman 2009; Verzijden and Rosenthal 2011).

A few empirical studies have explicitly tested for a connection between sexual imprinting and

sexual isolation between closely related populations or species. For example, benthic and

limnetic sticklebacks sexually imprint paternal cues under ecologically divergent selection,

which results in significant sexual isolation between the two morphs (Kozak et al. 2011). Other

studies in cichlids (Verzijden and ten Cate 2007), great and blue tits (Slagsvold et al. 2002), and

Darwin's finches (Grant and Grant 1997) have demonstrated that sexual imprinting can maintain

sexual isolation. Therefore, sexual imprinting seems to be a key, but underexplored, avenue to

speciation.

Here we assess the role of sexual imprinting in generating reproductive isolation

between two mammalian species, the white-footed mouse (*Peromyscus leucopus*) and the cotton

mouse (*P. gossypinus*), which are thought to have diverged during the Pleistocene (Blair 1950).

*P. leucopus* is distributed across the Midwest and eastern United States, whereas *P. gossypinus* is

restricted to the Southeast (Figure 1); their ranges overlap in the Gulf Coast states, from Texas to

Virginia. These species show some level of sexual isolation: *P. leucopus* and *P. gossypinus*

hybridize and produce viable, fertile offspring in the lab (this study, Dice 1937, 1940); however,

when multiple mice of each species are placed in large arenas, both species mate with

conspecifics (Bradshaw 1965, 1968). While assortative mating in laboratory studies is potentially

strong, there is mixed evidence as to whether it is strong enough to prevent hybridization in wild

sympatric populations (Dice 1940, Howell 1921, McCarley 1954).

In this study, we used genomic data to first assess hybridization in the wild and

conclusively found that the two species remain genetically distinct in sympatry despite rare

hybridization events. We then examined the degree of sexual isolation between *P. leucopus* and

*P. gossypinus,* and tested if it had a learned or genetic basis. Our results show that sexual

imprinting produces strong sexual isolation between these sister species, and we suggest that

learning disproportionately contributes to the total reproductive isolation we have observed in

between two inter-fertile, sympatric species.

**METHODS**

**Detection of hybrids in sympatric populations**

*Wild samples*

During April 2008 and January-February of 2010 and 2011, we collected mice from seven

allopatric locations and 13 sympatric locations (Figure 1). At each location, we placed up to 300

Sherman traps every 20 feet in transects of 50 traps per line. From each mouse captured, we took

liver or tail tissue and stored tissues in 100% ethanol for subsequent DNA extraction. We

augmented our own sampling with tissues collected at additional sites from museum specimens

at the Harvard Museum of Comparative Zoology, Oklahoma State University Collection of

Vertebrates, Sam Noble Museum of Natural History, and the Museum of Texas Tech University.

Precise collecting locations and sample sizes for all animals included in this study are provided

in Supplementary Table 1.

*Lab strains*

We obtained *P. leucopus* animals from the *Peromyscus* Genetic Stock Center (University

of South Carolina). The line was originally established with 38 founders brought into the lab

during 1982-1985. In 2009, we established a breeding colony of *P. gossypinus* animals with 18

founders caught in Jackson and Washington counties in Florida. In captivity, both breeding

colonies have been have been deliberately outbred to preserve genetic diversity.

All animals were housed in standard mouse cages in either mate pairs (one female and

one male) or in same sex cages with a maximum of five adults. Offspring were weaned into same

sex cages 23 days after birth. We set the light cycle to 14 hours of light and 10 hours of dark and

maintained a room temperature between 70 and 77 degrees Fahrenheit. All mice were fed a

regular Purina diet (Purina Iso Pro 5P76) *ad libitum*.

In addition to maintaining these two species strains, we also bred hybrids in the

laboratory. First generation hybrids (F1) were generated from both *P. gossypinus* female x *P.*

*leucopus* male matings as well as the reciprocal cross. Next, these F1 hybrids were backcrossed

to either *P. gossypinus* or *P. leucopus*.

*ddRADseq library construction and genotyping*

We extracted genomic DNA from 374 wild-caught individuals and two lab-raised hybrids

using an Autogen kit and AutoGenprep 965 instrument. We prepared double digest restriction-

associated DNA tag (ddRAD) libraries from each individual following the protocol described in

Peterson et al. (2012). Briefly, we digested 100-200 ng of DNA from every individual with two

restriction enzymes, EcoRI-HF and MspI (New England Biolabs) and purified the reactions with

AMPure XP beads (Beckman Coulter Genomics). After quantifying the cleaned and digested

product on a spectrophotometer plate reader (SpectraMax Gemini XS Plate Reader), we ligated

approximately 50 ng of digested DNA to uniquely barcoded EcoRI adapters and MspI adapters

in a 40 μl reaction volume with T4 DNA ligase (New England Biolabs). We pooled equal

amounts of 48 ligated samples and used two rounds of AMPure XP bead purification to reduce

the total pooled volume of up to 30 μl. We loaded each ligation pool onto a 2% agarose Pippin

Prep cassette (Sage Science) and selected fragments with a size of $300 \pm 35$ bp. We ran five

replicate Phusion PCRs according to the Finnzyme kit directions (Thermo Scientific) for 12

cycles with 5 μl of eluted Pippin Prep product as template. Each PCR was indexed using a

unique reverse primer (primer and index sequences from Peterson et al. 2012). Following PCR,

we pooled all replicate reactions and purified them with AMPure XP beads to concentrate each

ddRAD library. We multiplexed ddRAD libraries in equimolar ratios and sequenced 50 bp single

reads on an Illumina Genome Analzyer II or HiSeq (2000 or 2500).

We demultiplexed reads and aligned them by sample to the draft genome sequence of

*Peromyscus maniculatus* (NCBI: GCA_000500345.1) with STAMPY run in hybrid mode using

the BWA mem algorithm with default parameters (Lunter and Goodson 2011). We identified and

removed adapter sequences with Picard-tools 1.100 (http://picard.sourceforge.net). We realigned

potential indels with the Genome Analysis Tool Kit v. 3.2-2 IndelRealigner (GATK; McKenna

et al. 2010) and performed SNP discovery across all samples simultaneously using the GATK

UnifiedGenotyper (DePristo et al. 2011). We filtered alignments, keeping regions with 100 or

more total reads and an average base quality greater than 20. We retained bi-allelic SNPs with a

minimum mapping quality of 30 that were present in at least 90% of our individuals at a depth of

10 or greater. To reduce linkage among SNPs in our dataset, we identified "clusters" of SNPs

within 100 bp of each other and more than 100 bp from another SNP and we randomly selected

one SNP per cluster. Our final dataset contained 3,707 SNPs and 321 mice that had over 90% of

genotypes present at these SNPs. Of these mice, 21 *P. leucopus* were caught at allopatric sites or

lab-raised, 54 *P. gossypinus* were caught in allopatric sites or lab-raised, two individuals were

hybrids from our colonies, and 244 individuals were of unknown ancestry from the predicted

sympatric species range. Short read data were deposited in GenBank (accession number:

SRPXXXXXX).

*Identification of hybrids*

We used genetic principal component analysis (PCA) to evaluate admixture between *P.*

*leucopus* and *P. gossypinus*. We implemented genetic PCA using smartpca from the Eigensoft

6.0.1 package (Patterson et al. 2006) and output the first ten eigenvectors. After excluding outlier

individuals and SNPs, our final dataset contained 288 individuals and 2,528 SNPs. We expected

the first principal component to separate genetic differences between species and included

known individuals from each species (allopatric and identified museum specimens) to identify

PC1 values corresponding to each species. We assessed the eigenvector significance using

Tracy-Widom statistics (Patterson et al. 2006) implemented using twstats in Eigensoft 6.0.1.

**Measurement of sexual isolation between species**

Using lab-based assays, we first tested for intrinsic postzygotic isolation and then

estimated sexual isolation without mate choice. We could then compare our sexual isolation

estimate from no-choice assays to those with mate choice to quantify the contribution of mating

preferences to reproductive isolation between *P. leucopus* and *P. gossypinus*.

*Intrinsic postzygotic isolation and sexual isolation without choice*

We tested for intrinsic postzygotic sexual isolation and sexual isolation between lab-

raised *P. leucopus* and *P. gossypinus* using no-choice trials. We set up 20 crosses for each

conspecific and heterospecific pairing: L♀ x L♂, G♀ x G♂, L♀ x G♂, and G♀ x L♂ (in which

"L" represents *P. leucopus* and "G" represents *P. gossypinus*). When F1 offspring were

produced, we used these mice as F1 hybrids in additional no-choice trials in backcross mating

pairs: F1♀ x L♂, F1♀ x G♂, L♀ x F1♂, and G♀ x F1♂. We avoided any sib-sib or sib-parent

pairings.

We set up mating pairs by adding a sexually receptive virgin female to the cage of a

virgin, sexually mature male. We determined female sexual receptivity through vaginal lavage

and considered a female to be receptive between proestrus and estrus stages. We gave pairs 60

days to produce a litter, which is approximately 12 estrous cycles (mean estrous cycle length for

both species is 5-6 days; Dewsbury et al. 1977) or opportunities for successful reproduction. We

considered the production of offspring as a successful mating event and inferred the latency to

the first successful mating by subtracting the average gestation period—23 days in both species

(Pournelle 1952; Wolfe and Linzey 1977; Lackey et al. 1985)—from the total number of days

until a litter was born. Although our metric for mating success is conservative because it is

confounded with any fertility differences that might exist among individuals or between the

species, our assay nonetheless captures hybridization between these species.

We first used the no-choice assays to test hybrid viability and fertility in our laboratory

strains of *P. leucopus* and *P. gossypinus*. We scored offspring survival to reproductive age in

heterospecific crosses (L♀ x G♂, G♀ x L♂), and then used these F1 hybrids in backcrosses to

look for evidence of reduced fertility relative to conspecific crosses. To compare the proportion

of successful mating events between conspecific and heterospecific crosses, we used a logistic

regression to quantify the effects of the female species, male species, or the interaction between

female and male species. We then selected the best-fit model based on the lowest Akaike

Information Criterion (AIC). We compared the 95% confidence intervals for the mean mating

success among backcross pairs (F1♀ x L♂, F1♀ x G♂, L♀ x F1♂, G♀ x F1♂) to those of

conspecific crosses. Together, these no-choice data provide an estimate of hybrid viability and

relative fertility.

We next tested for differences in mating latency between conspecific, heterospecific, and

backcross mating pairs using a non-parametric Kruskal-Wallis rank sum test followed by

pairwise Wilcoxon tests with adjusted Bonferroni-corrected p-values. To quantify sexual

isolation, we counted the number of successful mating events to estimate a joint isolation index,

$I_{PSI}$ (Rolán-Alvarez and Caballero 2000), which compares observed to expected mating events

(assuming random mating among individuals) among conspecific and heterospecific pairs. This

index ranges from -1 (all mating occurred between species) to +1 (all mating occurred within

species), with a value of 0 indicating equal mating among pair types. We used the number of

conspecific and heterospecific pairs that produced litters to estimate $I_{PSI}$ in JMATING v. 1.0.8

(Carvajal-Rodriguez and Rolán-Alvarez 2006). We used 10,000 bootstrap replicates to estimate

the isolation indices, their standard deviation, and to test the hypothesis that our estimates of the

joint isolation deviated significantly from a null hypothesis of random mating.

*Sexual isolation with choice*

We contrasted our estimate of the sexual isolation index ($I_{PSI}$) from no-choice assays to

the sexual isolation index estimated from two-way choice assays. We measured conspecific

mating preferences in a two-way electronically-controlled gated mate choice apparatus that

consisted of three collinear rat cages, with each pair of cages separated by two RFID antennae

and gates (FBI Science Gmbh; Figure 3A). Each pair of gates was programed to allow passage

depending on the identity of the mouse. Specifically, for each trial we implanted three mice with

small transponders (1.4 mm x 9 mm, ISO FDX-B, Planet ID Gmbh) in the interscapular region

using a sterile hypodermic implanter and programmed the gates to allow the designated

"chooser" mouse (i.e. the individual whose preference we tested) to pass freely through all cages

while constraining each "stimulus" mouse to the left or right cage, respectively.

With this apparatus, we tested mate preferences of males and females of each species for

conspecific and heterospecific stimuli of the opposite sex. We allowed the chooser mouse—

either a sexually receptive virgin female (in proestrus or estrus as determined by vaginal lavage)

or a sexually mature virgin male—to acclimate to the apparatus for one day, adding food, water,

used nesting material, and a hut from each stimulus mouse's colony housing cage to the flanking

cages of the apparatus. Approximately 24 hours later, we returned the chooser mouse to the

center cage if it had not already nested there, closed all gates, and added stimulus mice to the two

flanking cages to allow them two to four hours to acclimate to their new environment. At lights

out (4:00 pm; 14:10 hour light:dark cycle), we re-opened the gates and recorded RFID readings

at all antennae as well as webcam video streams from each flanking cage for two nights (~44

hours; camera model: DLINK DCS-942L). Each chooser mouse was tested once.

At the end of each trial, we parsed a log file of RFID readings and calculated chooser

preference for a stimulus as the proportion of time spent with that stimulus divided by the time

spent with both stimuli (i.e. a ratio of association time between the stimuli). We analyzed only

trials in which the chooser mouse investigated both cages during the acclimation, spent at least

10 minutes investigating at least one stimulus during the trial, and the stimuli mice were in their

cages at least 75% of the trial period (we discarded 15% of trials that did not meet these criteria).

We compared the preferences of 8-11 adults (at 9-14 weeks of age) of each species and

sex for conspecific and heterospecific stimuli of the opposite sex. For female-choice trials, we

tested virgin female preferences for either: (1) pairs of sexually experienced males that had

successfully sired offspring with a conspecific female prior to use in the two-way choice trials

(*P. leucopus*, $N = 5$ trials; *P. gossypinus*, $N = 7$ trials), or (2) pairs of virgin males as stimuli (*P.*

*leucopus*, $N = 6$ trials; *P. gossypinus*, $N = 4$ trials). Because we did not detect a significant

difference in female preference based on male stimulus sexual experience (two-sided Wilcoxon

rank sum test, *P. gossypinus* females: $W = 9$, $p = 0.41$; *P. leucopus* females: $W = 15$, $p = 1$), we

combined female preference data from trials with sexually experienced and virgin male stimuli.

For male-choice trials, we used only virgin females as stimuli.

We estimated $I_{PSI}$ for each sex separately in JMATING v. 1.0.8 (Carvajal-Rodriguez and

Rolán-Alvarez 2006) because behavior of the stimuli may not be similar across male- and

female-choice trials. We estimated $I_{PSI}$ by considering the chooser and its most preferred

stimulus as a "mated" pair; when we observed no mating, we replaced zero values with a 1 to

allow for bootstrapping with resampling. We used 10,000 bootstrap replicates to estimate the

isolation indices and test for deviation from random mating ($I_{PSI} = 0$).

## Testing for sexual imprinting

To determine whether conspecific mating preferences are learned in the nest, we

measured the preferences of mice from each species after they had been cross-fostered—raised

from birth until weaning—by parents of the opposite species. We swapped whole litters at birth

between breeding pairs of *P. leucopus* and *P. gossypinus*, reducing litters to the same number of

offspring if litters differed in number of pups. All cross-fostering attempts were successful,

indicating that parents readily attended to unrelated offspring. We allowed cross-fostered

offspring to remain with their foster parents until weaning (23 days after birth), when we

separated offspring into same sex cages; this matches the life cycle of all other mice in our study.

As a control, we also cross-fostered offspring within species (i.e. swapped litters between

conspecific families) to partition the effects of litter transfer and foster parent species on mating

preference. Although there is mixed (or incomplete) information for whether fathers contribute

parental care in *P. leucopus* and *P. gossypinus* (Hartung and Dewsbury 1979; Schug et al. 1992),

we cross-fostered offspring to both parents because we maintain male-female breeding pairs in

our laboratory colonies of *P. leucopus* and *P. gossypinus* and aimed to compare preferences of

mice from cross-fostered and non-cross-fostered trials.

We tested the mating preferences of all cross-fostered mice in the two-way gated choice

assay described above. We predicted that if young mice sexually imprint on their parents, cross-

fostered mice raised with the opposite species should prefer heterospecific stimuli and exhibit a

weaker preference for conspecifics compared to individuals raised by their biological parents or

other unrelated conspecific parents. We evaluated the effects of chooser sex and cross-fostering

treatment on preferences for *P. leucopus* in each species, using linear modeling after applying an

arcsin transformation to the proportion of time spent with *P. leucopus*. To test for the possibility

that the sexes within each species might react differently to cross-fostering, we considered

models with and without an interaction between chooser sex and cross-fostering treatment and

selected the best-fit models based on the lowest AIC.

**Assessment of two-way choice assay**

We confirmed that our two-way mate choice assay accurately predicts mating preference

by measuring whether the most preferred stimulus corresponded to mating events in a subset of

trials in which mating occurred. We identified trials with successful mating events by either the

presence of sperm in a female reproductive tract at the end of a trial or the birth of a litter three

weeks later. If a female-choice trial resulted in offspring, we determined the identity of the father

by genotyping both the male stimuli and the pups at two to three microsatellite markers (loci 14,

35, and 80 from Weber et al. 2010) following the protocol described in Weber et al. 2010 ($N =$

15 trials) or screening video data for copulation events ($N = 5$ trials). We tested whether the most

preferred individual (as determined by the greatest proportion of association time) predicted

mating success using a linear regression. We applied an arcsin transformation to association time

proportions. This analysis allowed us to determine that association time is an accurate predictor

of mating, and thus reflects mating preference.

**RESULTS**

**Hybridization is rare in sympatric populations**

Using thousands of markers across the genome summarized in a genetic PCA, we tested

for evidence of hybridization between *P. leucopus* and *P. gossypinus* in sympatric populations.

We identified six significant principal components by Tracy-Widom statistics with the following

eigenvalues: (1) 37.855, (2) 4.352, (3) 3.627, (4) 3.161, (5) 3.054, and (6) 2.941. Based on

clustering with known allopatric and previously identified *P. leucopus* and *P. gossypinus*

individuals, the first eigenvector clearly separates *P. leucopus* (negative values) and *P.*

*gossypinus* (positive values) (Figure 1A). As expected, a control lab-generated F1 hybrid falls at

the midpoint along the first eigenvector and a lab backcross mouse (F1 x *P. gossypinus*) falls half

way between the F1 hybrid and the mean value of *P. gossypinus* values (Figure 1A). Of the 244

sympatric mice we collected, all could be easily assigned to either the *P. leucopus* or *P.*

*gossypinus* species, with only two exceptions: two mice from Big Lake Wildlife Management

Area, Louisiana had intermediate values along eigenvector 1 (Figure 1A). These admixed

individuals showed greater *P. leucopus* ancestry, similar to a F1 backcross or advanced

backcross to *P. leucopus*, indicating that there may be biased gene flow from *P. gossypinus* into

*P. leucopus*.

The second eigenvector revealed two genetically distinct *P. gossypinus* subgroups. These

likely reflect genetic differences between *P. gossypinus* subspecies, *P. gossypinus gossypinus*

and *P. gossypinus megacephalus*. Specifically, higher PC2 values corresponded to mice caught

east of the Mississippi River—which are more likely to be *P. g. gossypinus*—whereas lower PC2

values corresponded to mice caught west of the river—which are more likely to be *P. g.*

*megacephalus* (Wolfe and Linzey 1977). The Mississippi river is a known biogeographic barrier

for many species (Soltis et al. 2006), and our data suggest that this may also be the case for *P.*

*gossypinus*. Only one individual from the Tunica Hills wildlife management area population in

Louisiana failed to fit this pattern (Figure 1A): this individual occurred to the east of the

Mississippi River but it clustered with individuals from the western group. We did not find any

evidence to suggest a similar barrier to gene flow in *P. leucopus*, but we also did not have the

equivalent population-level sampling on both sides of the river. The remaining four eigenvectors

(3, 4, 5, and 6) identified population structure within *P. leucopus* (Supplemental Figure 1).

### *P. leucopus* and *P. gossypinus* co-occur in mosaic sympatry

340       Using cluster assignments based on the genetic PCA, we identified six of 13 sympatric

341 sites that contained both species (Figure 1B). However, the other seven sites contained only a

342 single species, highlighting the patchy distribution of both species within their broadly sympatric

343 range from Texas and Virginia. Thus, our data confirm that the two species occur in mosaic

344 sympatry.

### No evidence for intrinsic postzygotic isolation

Previous studies have suggested that there is no measurable intrinsic postzygotic isolation

in laboratory crosses of *P. leucopus* and *P. gossypinus* (Dice 1937). We confirmed this result in

our independent lines (i.e., different spatial and temporal origin) of these two species. We first

measured reproductive success within and between species in no-choice assays. The proportion

of crosses that produced offspring was determined largely by the female (logistic regression: $\beta$ =

1.25, SE = 0.47, $p$ = 0.008), with *P. leucopus* females showing greater mean mating success than

*P. gossypinus* (Supplemental Figure 2). Importantly, this means that *P. leucopus* females had

greater reproductive success with *P. gossypinus* males (12/20 pairs had offspring) than the

reciprocal cross between *P. gossypinus* females and *P. leucopus* males (6/20 pairs had offspring),

indicating some asymmetry in either mate preferences, copulation attempts, or female fertility.

Successful heterospecific crosses confirmed that we can produce viable F1 hybrids, which

survive until reproductive age. In addition, we compared the mating successes of backcrosses to

conspecific and heterospecific crosses. We found that these F1 hybrids are as fertile in

backcrosses (i.e., had similar frequency of litter production) as either conspecific or

heterospecific crosses, and that all backcross offspring are also viable (Supplemental Figure 2).

**<u>Mate choice causes significant sexual isolation</u>**

We examined whether mating preferences lead to significant sexual isolation between the

species in a laboratory environment (Supplemental Table 2). In no-choice assays, heterospecific

pairs hybridized and produced viable offspring, indicating no measurable sexual isolation exists

in the absence of mate choice ($I_{PSI} = 0.00$, $SD = 0.19$, $p = 0.960$; Figure 2). However,

conspecific, heterospecific, and backcross mating pairs had significantly different latencies to

produce offspring (Figure 2; Kruskal-Wallis: $\chi^2 = 6.7626$, df = 2, $p = 0.034$). Pairwise

comparisons between mating pairs revealed significance differences in latency to mating only

between conspecific and heterospecific mating pairs ($W = 69$, $p_{Bonferroni} = 0.010$). Heterospecific

pairs took an average of 5.4 days longer to produce litters than conspecific pairs, indicative of

either delayed heterospecific mating or longer hybrid gestation times. This delay is roughly

equivalent to one estrus cycle in *Peromyscus* (Dewsbury et al. 1977).

By contrast, we detected significant sexual isolation between the species in two-way

choice assays (Supplemental Table 2). Sexual isolation estimates were similar in female- and

male-choice trials: *P. leucopus* and *P. gossypinus* females strongly preferred conspecific mates

(Figure 3B; $I_{PSI} = 0.75$, $SD = 0.14$, $p < 0.01$) as did *P. leucopus* and *P. gossypinus* males (Figure

3B; $I_{PSI} = 0.75$, $SD = 0.15$, $p < 0.01$). More generally, there were strong preferences for

conspecific mates in both species, regardless of sex.

**Sexual imprinting contributes to sexual isolation in at least one species**

We next investigated whether mating preferences in these species had a learned or

genetic basis using a series of cross-fostering experiments and found that cross-fostering had

different effects on mating preference in the two focal species. In *P. leucopus,* the AIC selected a

model in which cross-fostering, sex, and their interaction was significant ($F = 5.09$ on 3 and 25

df, $p = 0.007$). When raised with their own parents, *P. leucopus* of both sexes preferred *P.*

*leucopus* stimuli (Figure 3B; female *P. leucopus* preference $\hat{p} = 0.689$; male *P. leucopus*

preference $\hat{p} = 0.959$). *P. leucopus* males that were cross-fostered significantly changed their

preference (Figure 3C; *P. leucopus* preference $\hat{p} = 0.184$; t $= 3.853$, $p_{Bonferroni} = 0.001$), whereas

cross-fostering did not significantly change female preferences (Figure 3; *P. leucopus* preference

$\hat{p} = 0.764$; t $= -0.390$, $p_{Bonferroni} = 1$). Thus, *P. leucopus* females always preferred *P. leucopus* to

*P. gossypinus* mates, whereas males spent more time with the species by whom it was raised

In *P. gossypinus,* the model selected by AIC showed a strong cross-fostering effect but no

significant sex effects or interactions between cross-fostering and sex ($F = 51.31$ on 1 and 33 df,

$p < 0.001$). When raised with their own parents, *P. gossypinus* of both sexes preferred *P.*

*gossypinus* stimuli (Figure 3B; *P. leucopus* preference $\hat{p} = 0.069$), whereas *P. gossypinus* raised

with *P. leucopus* preferred *P. leucopus* stimuli (Figure 3C; *P. leucopus* preference $\hat{p} = 0.781$).

To confirm that the preference reversal in *P. gossypinus* was not an effect of transferring

litters but rather an effect of the foster parent species, we collected an additional control dataset

for *P. gossypinus.* We cross-fostered *P. gossypinus* to unrelated *P. gossypinus* foster parents

(females: $N = 4$, males: $N = 7$) and found that foster species, and not the litter transfer itself,

affected *P. gossypinus* preferences (Supplemental Figure 3). Pairwise t-tests on arcsin-

transformed proportion of time spent with *P. leucopus* revealed no significant differences

between *P. gossypinus* raised with their own parents or unrelated conspecific parents ($t$ = -0.72,

df = 15.38, $p_{Bonferroni}$ = 1). Thus, *P. gossypinus* males and females preferred *P. gossypinus* to *P.*

*leucopus* mates, independent of if they were related to their conspecific parents.

To examine the effects of sexual imprinting on sexual isolation, we calculated the sexual

isolation index assuming the most preferred stimulus from each heterospecifically cross-fostered

choice trial (Figure 3C) as a "successful mating". Cross-fostering eliminated sexual isolation in

female-choice trials ($I_{PSI}$ = 0.25, $SD$ = 0.34, $p$ = 0.57) and male-choice trials ($I_{PSI}$ = -0.29, $SD$ =

0.42, $p$ = 0.32). Thus, our cross-fostering results confirm that sexual isolation between *P.*

*leucopus* and *P. gossypinus* is the result of sexual imprinting.

**<u>Two-way choice test accurately measures preference</u>**

To confirm that the time spent with either stimulus mouse was an accurate predictor of

mate preference and hence mate choice, we recorded 20 mating events in our two-way choice

assays: 12 mating events occurred in trials where choosers were raised with their own parents

and 8 trials where choosers were raised with heterospecific foster parents. In 19 out of 20 trials,

choosers mated with the stimulus individual with which they spent the most time (Figure 4).

Mating outcome (with conspecific or heterospecific stimulus) was predicted by proportion of

time spent with the conspecific stimulus (logistic regression: $\beta$ = 10.06, SE = 4.86, $p$ = 0.04),

indicating that our two-way choice assay accurately detects mating preferences.

**DISCUSSION**

Sexual imprinting could be a powerful generator of sexual isolation because it quickly

and effectively associates preferences with traits in populations. Furthermore, sexual imprinting

has been documented in a diversity of taxa—e.g. birds, fish, mammals, amphibians, and

insects—suggesting it could be a broadly important driver of speciation (Immelmann 1975). Our

study shows that sexually imprinted mate-choice has maintained and contributed to strong sexual

reproductive isolation in a classic mammalian system.

**Rare hybridization in sympatry indicates a high degree of reproductive isolation**

To test the strength of reproductive isolation between *P. leucopus* and *P. gossypinus* in

nature, we collected 374 mice from across their range overlap and generated genomic data to

measure the frequency of hybridization between these species in sympatry. Classic studies by

mammalogists in the mid 1900's reported mixed evidence as to the extent of interspecific

hybridization in sympatric populations. In Louisiana, Alabama, and southern Illinois, Howell

(1921) and later McCarley (1954) identified a few intermediate individuals resembling hybrids

based on morphology and allozyme genotypes. By contrast, Dice (1940) found no evidence of

morphological intermediates in his studies in Virginia. Thus, it was unclear how frequently these

species hybridize in the wild.

In our study, we found less than 1% of individuals collected from sites in which both

species co-occurred were admixed. Our genomic analysis thus supports the conclusions of

previous studies that suggested *P. leucopus* and *P. gossypinus* may remain genetically distinct in

nature (Dice 1940; Price and Kennedy 1980; Robbins et al. 1985) in spite of rare hybrids

(Howell 1921; McCarley 1954b; Barko and Feldhamer 2002). The two hybrids that we found

were both from Big Lake Wildlife Management Area (Louisiana), and they had greater

proportions of *P. leucopus* ancestry. At this site, *P. gossypinus* were less common than *P.*

*leucopus*, potentially accounting for the biased gene flow into *P. leucopus*. Previous behavioral

data also support the possibility of biased gene flow from *P. gossypinus* into *P. leucopus*. For

example, using population cages to test interactions among pairs of male and female *P. leucopus*

and *P. gossypinus*, Bradshaw (1965) found that *P. leucopus* females were far more tolerant of *P.*

*gossypinus* males than vice versa. Although no hybrid offspring were produced in this study, if

heterospecific matings were to occur, one would predict they would be in the direction of a *P.*

*leucopus* female with a *P. gosyspinus* male.

Although we detected some admixture and possible biased gene flow from *P. gossypinus*

into *P. leucopus*, the overall lack of pervasive hybridization between these otherwise inter-fertile

species suggests that they are strongly reproductively isolated in nature. Our genomic data

revealed that *P. leucopus* and *P. gosssypinus* are distributed in a mosaic sympatry, with many

sites containing only one species (seven of thirteen sampling sites). This patchiness could be

driven by differences in microhabitat use: *P. leucopus* often occupy upland habitat and use more

arboreal nest sites while *P. gossypinus* often occupy swamps and bottomland habitat and use

more ground nest sites when they co-occur (McCarley 1954c, 1963; Taylor and Mccarley 1963).

However, these habitat differences are not enough to exclude contact in sympatry because both

species can be trapped in the same patch of forest, especially where these habitat types abut

(Dice 1940; Calhoun 1941; Price and Kennedy 1980; Roehrs et al. 2012). In fact, we often

caught both species in the same trap line. Similarly, there do not appear to be any significant

differences in breeding seasons: the two species have overlapping peak reproductive activities in

the winter months, but adults from both species can also be caught in reproductive condition

throughout the year in Texas, Louisiana, and Alabama (Pournelle 1952; McCarley 1954a; Wolfe

and Linzey 1977). Thus, the distributions, habitat preferences, and breeding seasons do not

appear to form complete or even strong reproductive barriers, suggesting that behavioral

differences may be the most important cause of reproductive isolation.

**_P. leucopus_ and _P. gossypinus_ remain genetically distinct due to learned sexual isolation**

We tested for evidence of sexual isolation, as previous studies suggested that mating

preferences might explain the lack of hybridization in the wild. Using no-choice and choice trials

to examine _P. leucopus_ and _P. gossypinus_ mating preferences, we found that conspecific

preferences form a significant sexual reproductive barrier between the two species. Without a

choice of mates, _P. leucopus_ and _P. gossypinus_ did not show significant sexual isolation,

although there was an increase in latency to mate in heterospecific relative to conspecific pairs.

However, when given a choice of mates, the species mated assortatively, and we estimated the

average joint sexual isolation index ($I_{PSI}$) between the species to be 0.651. While sexual isolation

is not yet complete ($I_{PSI} < 1$) between these species, the amount of sexual isolation we have

observed is far greater than what has been detected among cactophilic ($I_{PSI} = 0.12$; Etges and

Tripodi 2008) or Caribbean _Drosophila_ ($I_{PSI} = 0.159-0.282$; Yukilevich and True 2008), walking

stick insect populations ($I_{PSI} = 0.24-0.53$; Nosil et al. 2013), or gold and normal Nicaraguan

cichlid color morphs ($I_{PSI} = 0.39$ and $0.86$; Elmer et al. 2009), placing _P. leucopus_ and _P._

_gossypinus_ quite far along a speciation continuum.

We cannot determine if sexual isolation resulted because of strong female preferences

and weak male preferences (where males select mates based on female acceptance or rejection)

or the reverse, weak female preferences and strong male preferences (where females select mates

based on male courtship). However, because our test apparatus provides the opportunity for

choosers to avoid either stimulus, we interpret a pattern of sexual isolation in female- and male-

choice trials to be the result of active conspecific preferences in both sexes.

Our two-way choice results, compared to our no-choice results, indicate that mating

preferences increase sexual isolation (Coyne et al. 2005). Testing the mating preferences of

heterospecifically cross-fostered mice revealed that both species are affected by sexual

imprinting, but that the degree of imprinting differed by species and sex. For one, both male and

female *P. gossypinus* strongly sexually imprinted on their foster parent species, indicating that

mating preferences in *P. gossypinus* are entirely learned. By contrast, we found that *P. leucopus*

also sexually imprint on their parents, though weakly. Some *P. leucopus* males had a reduced

preference for conspecifics when raised with heterospecific parents, whereas all *P. leucopus*

females appeared unaffected by cross-fostering. In other words, *P. leucopus* females showed an

own species bias, suggesting that their mate preferences could be genetic. *P. leucopus* showed a

similar sexual difference in imprinting in a study that examined *P. leucopus* preferences for

soiled bedding after cross-fostering to grasshopper mice, *Onychomys torridus* (McCarty and

Southwick 1977a): male and female *P. leucopus* raised with *O. torridus* parents had decreased

preference for conspecific soiled bedding, but the effect was more dramatic in males than

females. Thus, both *P. leucopus* and *P. gossypinus* appear to learn mating preferences, but the

degree of sexual imprinting varies between the two species, particularly in cross-fostered *P.*

*leucopus* females which show a bias towards mates of their own species.

**Genetic preference or preference reversal after socialization may account for the lack of**

**sexual imprinting observed in *P. leucopus* females**

Sex differences in sexual imprinting are most likely caused by sex-specific genetic (or

epigenetic) differences in learning, or learning in both sexes followed by preference reversal in

one sex after social interaction. Sex-specific genetic differences that affect the length of the

sensitive learning period or how the learned preference is internalized could create a sex bias in

learning. In some species such as sticklebacks and cichlids, only one sex (females) sexually

imprints (Verzijden and ten Cate 2007; Verzijden et al. 2009; Kozak et al. 2011); in other species

such as zebra finches, both sexes sexually imprint but on different parental cues (Vos 1995).

Learning in these species may be affected by sex-specific differences, but genetic loci that cause

these differences have yet to be identified. Alternatively, conspecific mating preferences could

be epigenetically-determined by parental behaviors or experiences. Some behavioral traits such

as pup licking in rats (Francis et al. 1999), fear of odors in mice (Dias and Ressler 2013), anxiety

in stickleback fish (McGhee and Bell 2014), have been linked to epigenetic inheritance. We

cannot rule out that conspecific preferences in female *P. leucopus* were determined

epigenetically, but suspect that social interactions most likely explain the own species bias in *P.*

*leucopus* females.

Social interactions can also influence sexual imprinting, and may be more likely to

account for sex differences. Irwin & Price (1999) describe a learning model in which offspring

might initially develop a generalized behavioral response to a range of traits resembling their

parents that later contracts after experience with heterospecifics. In cross-fostering studies, this

model would be supported by instances in which individuals sexually imprint on their foster

parents but subsequently alter their preferences in the presence of conspecifics. For example, a

longitudinal study of the effects of cross-fostering between sheep and goats found that females

initially preferred males of their foster species but later preferred conspecifics after a year of

socialization with conspecifics (Kendrick et al. 1998). Such preference reversals might be driven

because of courtship behavior.

Species in which females show an own species bias after cross-fostering treatments with

heterospecific parents have often been shown to be affected by differences in the courtship

activity of male stimuli. For example, female zebra finches raised with heterospecific Bengalese

foster parents spent more time with Bengalese males but directed more sexually receptive tail

quivering behavior to conspecific males (ten Cate and Mug 1984); this own species bias was

shown to be the result of greater conspecific male courtship activity. Male zebra finches sang

more vigorously and frequently than Bengalese males, biasing female sexual behavior toward

conspecifics (ten Cate and Mug 1984). Male courtship activity has also been shown to modify

sexually imprinted preferences in female mallards (Bossema and Kruijt 1982; Kruijt et al. 1982).

In our trials, male *P. leucopus* stimuli could have directed more copulatory behavior toward *P.*

*leucopus* females, whereas male *P. gossypinus* stimuli may have been antagonistic, causing *P.*

*leucopus* females to develop stronger preferences for conspecific males. Females could have

been responsive to differences in ultrasonic vocalizations which help attract mates (Pomerantz et

al. 1983; Musolf et al. 2010), male mounting attempts, or aggression. Because our choice assay

permits physical interaction between the stimuli and the chooser mice, we cannot rule out the

very likely possibility that *P. leucopus* females did sexually imprint but are affected by the

behavior of stimuli in our two-way choice assay.

Further studies designed to test for sexual imprinting as a function of species-directed

male courtship may be able to determine whether *P. leucopus* females sexually imprint but alter

their preferences based on conspecific courtship activity, or if they truly show an innate

preference for males of their own species. Whether *P. leucopus* females show biased conspecific

preferences because of genetic differences or subsequence social interactions, the asymmetry in

*P. leucopus* female and *P. gossypinus* female responses suggest that the species differ

meaningfully in either female learning or male courtship behaviors.

**<u>Sexual imprinting cues</u>**

Although we do not know the precise imprinting cues (e.g., odors, vocalizations) that *P.*

*leucopus* and *P. gossypinus* learn in the nest, they most likely learn olfactory cues as other

rodents are known to imprint on nest odors (Mainardi et al. 1965; Marr and Gardner 1965; Carter

and Marr 1970; Quadagno and Banks 1970; McCarty and Southwick 1977a; Porter et al. 1983).

If *P. leucopus* and *P. gossypinus* diverged in odor sources containing species, sex, or individual

information such as saliva (Gray et al. 1984; Smith and Block 1991; Talley et al. 2001), urine

(Doty 1973; Smadja and Ganem 1998; Pillay 2000; Hurst et al. 2001), scent marks (Johnston and

Brenner 1982; Becker et al. 2012), or major histocompatibility complex alleles (Yamazaki et al.

1979; Brown et al. 1989), these could serve as imprinting cues (e.g. Penn & Potts 1998). *P.*

*leucopus* and *P. gossypinus* have diverged in small urinary proteins (Cain et al. 1992) and *P.*

*leucopus* have been shown to sexually imprint on olfactory information (McCarty and Southwick

1977b), suggesting that sexual imprinting could be olfactory-based in this species pair.

Natural olfactory signals, which are affected by diet, could have been obscured by the

common Purina laboratory diet that we fed our *P. leucopus* and *P. gossypinus* stocks. Diet

influences odors in guinea pigs (Beauchamp 1976), mice (Schellinck et al. 1992), and voles

(Ferkin et al. 1997), and *P. gossypinus* and *P. leucopus* consume different diets in nature. *P.*

*gossypinus* is mainly carnivorous, with over two thirds its stomach contents containing animal

matter (insects and gastropods) compared to *P. leucopus*, which is primarily herbivorous, with

greater than two thirds of its stomach contents containing plant matter (Calhoun 1941). If

species-specific olfactory signals (e.g., urine, feces, scent marks) were affected by different diets,

imprinting cues and thus sexual isolation may be much more pronounced between natural

populations of *P. leucopus* and *P. gossypinus* than what we have detected in the laboratory.

**<u>Sexual imprinting may limit the effects of rare hybridization</u>**

When gene flow occurs, theoretical models have shown that sexual imprinting can create

substantial reproductive isolation and facilitate sympatric speciation more often than genetic

preferences (Verzijden et al. 2005). When hybrid fitness is high, as is the case for *P. leucopus*

and *P. gossypinus*, mating preferences formed by sexual imprinting may be more effective at

producing reproductive isolation than if the preferences were genetically controlled on

autosomes or sex chromosomes (Servedio et al. 2009).

We suspect that occasional hybridization between *P. leucopus* and *P. gossypinus* might

be tolerated because sexual imprinting would prevent the formation of a hybrid swarm. The two

hybrids that we found showed greater *P. leucopus* ancestry, indicating asymmetrical gene flow

from *P. gossypinus* to *P. leucopus*. While we did not have mitochondrial or Y chromosome

markers in our dataset that would have allowed us to identify which type of heterospecific

mating pair produced these hybrids, our no-choice data indicate that *P. leucopus* females have

greater reproductive success with *P. gossypinus* males than *P. gossypinus* females have with *P.*

*leucopus* males. We therefore predict that the few hybrids we found are likely offspring from an

initial cross between a *P. leucopus* mother and a *P. gossypinus* father. Futhermore, *P. leucopus*

females are far more tolerant of *P. gossypinus* males than the reciprocal direction (Bradshaw

1965). If *Peromyscus* offspring sexually imprint on their mothers as they do in other mammals,

we would expect F1 hybrids to preferentially backcross with *P. leucopus* mates, facilitating

asymmetrical gene flow while simultaneously preventing rampant hybridization. This type of

sexual imprinting analysis in F1 hybrids is seldom done (but see Albert 2005), but could reveal

how sexual imprinting might bias gene flow. In a limited number of unpublished trials, we found

evidence that *Peromyscus* mice sexually imprint on mothers (i.e., sexual imprinting still occurs

when no fathers are present). Although more rigorous testing is necessary to determine if

*Peromyscus* imprint on one or both parents, we predict that maternal imprinting leads to

introgression but prevents complete admixture, allowing the species to remain genetically

distinct in sympatry even with a small amount of hybridization.

Sexual imprinting might also prevent rampant gene flow between inter-fertile sympatric

species if it becomes reinforced (Irwin and Price 1999; Servedio et al. 2009). There is some

evidence for reinforcement between *P. leuocopus* and *P. gossypinus*. In a nesting assay,

sympatric *P. gossypinus* males and females and *P. leucopus* females preferred to spend more

time near conspecific individuals, while allopatric mice from both species showed no significant

preference for conspecifics (McCarley 1964). Similar patterns of increased species recognition in

sympatry have also been observed between *P. eremicus* and *P. californicus* (Smith 1965; Carter

and Brand 1986). However, for reinforcement to occur, hybrids must have reduced fitness. We

did not find evidence of hybrid inviability or sterility in our laboratory study, but we did not

quantify the degree of hybrid fertility which can vary in severity in hybrid zones (e.g. Turner et

al. 2011). In contrast to fertility-related traits, there is some evidence of postzygotic behavioral

sterility. A previous study found that *P. leucopus* and *P. gossypinus* reciprocal hybrids have

copulatory behaviors like each parental species but that they initiate copulation less frequently

than either *P. leucopus* or *P. gossypinus* (Lovecky et al. 1979). Similarly, both reciprocal hybrids

have similar exploratory behavior to both parental species but spent more time freezing in open-

field exploratory behavior tests (Wilson et al. 1976). In nature, any reduced mating success or

exploratory behavior would reduce hybrid fitness relative to their parents. Finally, hybrids might

also be behaviorally sterile if they have intermediate mating traits. For example, hybrids between

*M. m. musuculus* and *M. m. domesticus* have intermediate urinary signals that are sexually

selected against by each subspecies (Latour et al. 2014). The potential for hybrid behavioral

sterility, coupled with the fact that moderate sexual imprinting induces sexual isolation in our

allopatric lab stocks, suggests that it may be possible for reinforcement to boost reproductive

isolation in sympatry, helping explain the paucity of hybrids we have observed in our study.

**CONCLUSION**

Our study supports an emerging view that sexual imprinting could be vital to the

generation and maintenance of sexual reproductive barriers. Pending divergent natural selection

on an imprintable trait, a species that learns mating preferences may develop significant sexual

isolation that might mitigate the effects of hybridization. Our demonstration of sexual imprinting

in *Peromyscus leucopus* and *P. gossypinus*, sympatric sister species that have few other

reproductive barriers between them, indicates that sexual imprinting may disproportionately

contribute to their total reproductive isolation. Sexual imprinting may sculpt reproductive

isolation in subspecies (e.g. benthic and limnetic sticklebacks) undergoing initial morphological

and behavioral divergence, or help preserve reproductive isolation between already divergent

species, as we have shown to be the case in *P. leucopus* and *P. gossypinus*. Examining the role of

sexual imprinting in similar cases of speciation driven by sexual reproductive barriers will

continue to expand our understanding of the role of behavior in speciation.

**Acknowledgements**: We thank members of the Hoekstra lab who helped with field work: G. Gonçalves, V. Domingues, H. Fisher, E. Jacobs-Palmer, E. Kingsley, C. Linnen, H. Metz, B. Peterson, J. Weber, and several undergraduates, A. Chiu, K. Lin, and E. zu Ermgassen; we also thank E. Pivorun, R. Sikes, and their students for assistance with field work. J. Chupasko and M. Omura from the MCZ Mammal Department helped with specimen preparation. Nokuse Plantation granted us permission to trap on their property. The following museums donated tissue samples: Museum of Comparative Zoology, Oklahoma State University Collection of Vertebrates, Sam Noble Museum of Natural History, and the Museum of Texas Tech University. We also wish to thank N. Delaney, K. Ferris, and J. Mallet for helpful discussions and comments on this manuscript. This work was funded by: NSF GRFP, NSF DDIG, and grants from the American Society of Mammaologists, Animal Behavior Society, and the Harvard University Mind, Brain & Behavior Initiative to EKD. HEH is an Investigator of the Howard Hughes Medical Institute. The authors have no conflicts of interest.

**Author contributions**: EKD and HEH conceived the project. EKD collected and analyzed the data. EKD and HEH wrote the paper.

**References**

Albert, A. Y. K. 2005. Mate choice, sexual imprinting, and speciation: a test of a one-allele isolating mechanism in sympatric sticklebacks. Evolution 59:927–931.

Barko, V. A., and G. A. Feldhamer. 2002. Cotton mice (*Peromyscus gossypinus*) in southern Illinois: evidence for hybridization with white-footed mice (*Peromyscus leucopus*). Am.

Midl. Nat. 147:109–115.

Beauchamp, G. K. 1976. Diet influences attractiveness of urine in guinea pigs. Nature 263:587–
588.

Becker, E. A., S. Petruno, and C. A. Marler. 2012. A comparison of scent marking between a
monogamous and promiscuous species of *Peromyscus*: pair bonded males do not advertise
to novel females. PLoS One 7:e32002.

Blair, W. F. 1950. Ecological factors in speciation of *Peromyscus*. Evolution 4:253–275.

Bossema, I., and J. P. Kruijt.1982. Male activity and female mate acceptance in the mallard
(*Anas platyrhynchos*). Behaviour 79:313–324.

Boughman, J. W., H. D. Rundle, and D. Schluter. 2005. Parallel evolution of sexual isolation in
sticklebacks. Evolution 59:361–373.

Bradshaw, H. D., and D. W. Schemske. 2003. Allele substitution at a flower colour locus
produces a pollinator shift in monkeyflowers. Nature 426:176–178.

Bradshaw, W. N. 1968. Progeny from experimental mating tests with mice of the *Peromyscus*
*leucopus* group. J. Mammal. 49:475–480.

Bradshaw, W. N. 1965. Species discrimination in the *Peromyscus leucopus* group of mice. Texas
J. Sci. 17:278–293.

Brown, R. E., B. Roser, and P. B. Singh. 1989. Class I and class II regions of the major
histocompatibility complex both contribute to individual odors in congenic inbred strains of
rats. Behav. Genet. 19:659–674.

Cain, K. A., T. A. Burns, and D. T. Stalling. 1992. Urinary proteins in four rodent species.
Comp. Biochem. Physiol. 101B:199–204.

Calhoun, J. B. 1941. Distribution and food habits of mammals in the vicinity of the Reelfoot

Lake Biological Station. J. Tennessee Acad. Sci. 16:177–187.

Carter, C. S., and J. N. Marr. 1970. Olfactory imprinting and age variables in the guinea-pig,

*Cavia porcellus*. Anim. Behav. 18:238–244.

Carter, R. L., and L. R. Brand. 1986. Species recognition in wild-caught, laboratory-reared and

cross-fostered *Peromyscus californicus* and *Peromyscus eremicus* (Rodentia, Cricetidae).

Anim. Behav. 34:998–1006.

Coyne, J. A., S. Elwyn, and E. Rolán-Alvarez. 2005. Impact of experimental design on

*Drosophila* sexual isolation studies: direct effects and comparison to field hybridization

data. Evolution 59:2588–2601.

Coyne, J. A., and H. A. Orr. 1989. Patterns of speciation in *Drosophila*. Evolution 43:362–381.

Coyne, J. A., and H. A. Orr. 1997. "Patterns of speciation in *Drosophila*" revisited. Evolution

51:295–303.

DePristo, M. A., E. Banks, R. Poplin, K. V Garimella, J. R. Maguire, C. Hartl, A. A. Philippakis,

G. del Angel, M. A. Rivas, M. Hanna, A. McKenna, T. J. Fennell, A. M. Kernytsky, A. Y.

Sivachenko, K. Cibulskis, S. B. Gabriel, D. Altshuler, and M. J. Daly. 2011. A framework

for variation discovery and genotyping using next-generation DNA sequencing data. Nat.

Genet. 43:491–498.

Dewsbury, D. A., D. Q. Estep, and D. L. Lanier. 1977. Estrous cycles of nine species of muroid

rodents. J. Mammal. 58:89–92.

Dias, B. G., and K. J. Ressler. 2014. Parental olfactory experience influences behavior and neural

structure in subsequent generations. Nat. Neurosci. 17:89-96.

Dice, L. R. 1937. Fertility relations in the *Peromyscus leucopus* group of mice. Contrib. Lab.

Vert. Genet. 4:1–3.

Dice, L. R. 1940. Relationships between the wood-mouse and the cotton-mouse in eastern

Virginia. J. Mammal. 21:14–23.

Doty, R. L. 1973. Reactions of deer mice (*Peromyscus maniculatus*) and white-footed mice

(*Peromyscus leucopus*) to homospecific and heterospecific urine odors. J. Comp. Physiol.

Psychol. 84:296–303.

Elmer, K. R., T. K. Lehtonen, and A. Meyer. 2009. Color assortative mating contributes to

sympatric divergence of neotropical cichlid fish. Evolution 63:2750–2757.

Etges, W. J., and D. A. Tripodi. 2008. Premating isolation is determined by larval rearing

substrates in cactophilic *Drosophila mojavensis*. VIII. Mating success mediated by

epicuticular hydrocarbons within and between isolated populations. J. Evol. Biol. 21:1641–

1652.

Felsenstein, J. 1981. Skepticism towards Santa Rosalia, or why are there so few kinds of

animals? Evolution 35:124–138.

Ferkin, M. H., E. S. Sorokin, R. E. Johnston, and C. J. Lee. 1997. Attractiveness of scents varies

with protein content of the diet in meadow voles. Anim. Behav. 53:133–141.

Fisher, H. S., B. B. M. Wong, and G. G. Rosenthal. 2006. Alteration of the chemical

environment disrupts communication in a freshwater fish. Proc. R. Soc. B. 273:1187–1193.

Francis, D., J. Diorio, D. Liu, and M. J. Meaney. 1999. Nongenomic transmission across

generations of maternal behavior and stress responses in the rat. Science 286:1155–1158.

Grant, P. R., and B. R. Grant. 1997. Hybridization, sexual imprinting, and mate choice. Am. Nat.

149:1–28.

Gray, B., R. B. Fischer, and G. F. Meunier. 1984. Preferences for salivary odor cues by female

hamsters. Horm. Behav. 18:451–456.

Hall, E. R. 1981. The mammals of North America. Second edition. John Wiley & Sons, New

York.

Hall, E. R. and K. R. Kelson. 1959. The mammals of North America. Ronald Press, New York.

Hartung, T. G., and D. A. Dewsbury. 1979. Paternal behavior in six species of muroid rodents.

Behav. Neural Biol. 26:466–478.

Howell, A. H. 1921. A biological survey of Alabama. North Am. Fauna 45:1–89.

Hurst, J. L., C. E. Payne, C. M. Nevison, A. D. Marie, R. E. Humphries, D. H. L. Robertson, A.

Cavaggioni, and R. J. Beynon. 2001. Individual recognition in mice mediated by major

urinary proteins. Nature 414:631–634.

Immelmann, K. 1975. Ecological significance of imprinting and early learning. Annu. Rev. Ecol.

Evol. Syst. 6:15–37.

Irwin, D. E., and T. Price. 1999. Sexual imprinting, learning and speciation. Heredity 82:347–

354.

Jiggins, C. D., R. E. Naisbit, R. L. Coe, and J. Mallet. 2001. Reproductive isolation caused by

colour pattern mimicry. Nature 411:302–305.

Johnston, R. E., and D. Brenner. 1982. Species-specificity of scent marking in hamsters. Behav.

Neural Biol. 35:46–55.

Kendrick, K. M., M. R. Hinton, K. Atkins, M. A. Haupt, and J. D. Skinner. 1998. Mothers

determine sexual preferences. Nature 395:229–230.

Kozak, G. M., and J. W. Boughman. 2009. Learned conspecific mate preference in a species pair

of sticklebacks. Behav. Ecol. 20:1282–1288.

Kozak, G. M., M. L. Head, and J. W. Boughman. 2011. Sexual imprinting on ecologically

divergent traits leads to sexual isolation in sticklebacks. Proc. R. Soc. B. 278:2604–2610.

Kruijt, J. P., I. Bossema, and G. J. Lammers. 1982. Effects of early experience and male activity

on mate choice in mallard females (*Anas platyrhynchos*). Behaviour 80:32–43.

Lackey, J. A., D. G. Huckaby, and B. G. Ormiston. 1985. *Peromyscus leucopus*. Mamm. Species

247:1–10.

Laland, K. N. 1994. On the evolutionary consequences of sexual imprinting. Evolution 48:477–

489.

Latour, Y., M. Perriat-Sanguinet, P. Caminade, P. Boursot, C. M. Smadja, and G. Ganem. 2014.

Sexual selection against natural hybrids may contribute to reinforcement in a house mouse

hybrid zone. Proc. R. Soc. B. 281:20132733.

Lovecky, D. V., D. Q. Estep, and D. A. Dewsbury. 1979. Copulatory behavior of cotton mice

(*Peromyscus gossypinus*) and their reciprocal hybrids with white-footed mice (*P. leucopus*).

Anim. Behav. 27:371–375.

Lunter, G., and M. Goodson. 2011. Stampy: a statistical algorithm for sensitive and fast mapping

of Illumina sequence reads. Genome Res. 21:936–939.

Mainardi, D., M. Marsan, and A. Pasquali. 1965. Causation of sexual preferences of the house

mouse. The behavior of mice reared by parents whose odor was artificially altered. Atti.

Soc. Ital. Sci. Nat. 104:325–338.

Marr, J. N., and L. E. J. Gardner. 1965. Early olfactory experience and later social behaviour in

the rat: preference, sexual responsiveness, and care of young. J. Genet. Pyschology

107:167–174.

Matsubayashi, K. W., and H. Katakura. 2009. Contribution of multiple isolating barriers to

reproductive isolation between a pair of phytophagous ladybird beetles. Evolution 63:2563–

2580.

McCarley, H. 1963. Distributional relationships of sympatric populations of *Peromyscus*

*leucopus* and *P. gossypinus*. Ecology 44:784–788.

McCarley, H. 1964. Ethological isolation in the cenospecies *Peromyscus leucopus*. Evolution

18:331–332.

McCarley, W. H. 1954a. Fluctuations and structure of *Peromyscus gossypinus* populations in

eastern Texas. J. Mammal. 35:526–532.

McCarley, W. H. 1954b. Natural hybridization in the *Peromyscus leucopus* species group of

mice. Evolution 8:314–323.

McCarley, W. H. 1954c. The ecological distribution of the *Peromyscus leucopus* species group

in eastern Texas. Ecology 35:375–379.

McCarty, R., and C. H. Southwick. 1977a. Cross-species fostering: effects on the olfactory

preference of *Onychomys torridus* and *Peromyscus leucopus*. Behav. Biol. 19:255–260.

McCarty, R., and C. H. Southwick. 1977b. Patterns of parental care in two Cricetid rodents,

*Onychomys torridus* and *Peromyscus leucopus*. Anim. Behav. 25:945–948.

McGhee, K. E., and A. M. Bell. 2014. Paternal care in a fish: epigenetics and fitness enhancing

effects on offspring anxiety. Proc. R. Soc. B 281:20141146.

McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernytsky, K. Garimella, D.

Altshuler, S. Gabriel, M. Daly, and M. A. DePristo. 2010. The Genome Analysis Toolkit: a

MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res.

20:1297–1303.

Mckinnon, J. S., S. Mori, B. K. Blackman, L. David, D. M. Kingsley, L. Jamieson, J. Chou, and

D. Schluter. 2004. Evidence for ecology's role in speciation. Nature 429:294–298.

Mendelson, T. C. 2003. Sexual isolation evolves faster than hybrid inviability in a diverse and

sexually dimorphic genus of fish (Percidae: *Etheostoma*). Evolution 57:317–327.

Musolf, K., F. Hoffmann, and D. J. Penn. 2010. Ultrasonic courtship vocalizations in wild house

mice, *Mus musculus musculus*. Anim. Behav. 79:757–764.

Noor, M. A. F. 1997. How often does sympatry affect sexual isolation in *Drosophila*? Am. Nat.

149:1156–1163.

Nosil, P. 2007. Divergent host plant adaptation and reproductive isolation between ecotypes of

*Timema cristinae* walking sticks. Am. Nat. 169:151–162.

Nosil, P., R. Riesch, and M. Muschick. 2013. Climate affects geographic variation in host-plant

but not mating preferences of *Timema cristinae* stick-insect populations. Evol. Ecol. Res.

15:1–16.

Ortíz-Barrientos, D., and M. A. F. Noor. 2005. Evidence for a one-allele assortative mating

locus. Science 310:1467.

Patterson, N., A. L. Price, and D. Reich. 2006. Population structure and eigenanalysis. PLoS

Genet. 2:2074–2093.

Penn, D., and W. Potts. 1998. MHC-disassortative mating preferences reversed by cross-

fostering. Proc. R. Soc. B. 265:1299–306.

Peterson, B. K., J. N. Weber, E. H. Kay, H. S. Fisher, and H. E. Hoekstra. 2012. Double digest

RADseq: an inexpensive method for *de novo* SNP discovery and genotyping in model and

non-model species. PLoS One 7:e37135.

Pillay, N. 2000. Female mate preference and reproductive isolation in populations of the striped

mouse *Rhabdomys pumilio*. Behaviour 137:1431–1441.

Pomerantz, S. M., A. A. Nunez, and N. J. Bean. 1983. Female behavior is affected by male

ultrasonic vocalizations in house mice. Physiol. Behav. 31:91–96.

Porter, R. H., J. A. Matochik, and J. W. Makin. 1983. Evidence for phenotype matching in spiny

mice (*Acomys Cahirinus*). Anim. Behav. 31:978–984.

Pournelle, G. H. 1952. Reproduction and early post-natal development of the cotton mouse,

*Peromyscus gossypinus*. J. Mammal. 33:1–20.

Price, P. K., and M. L. Kennedy. 1980. Genic relationships in the white-footed mouse,

*Peromyscus leucopus*, and the cotton mouse, *Peromyscus gossypinus*. Am. Midl. Nat.

103:73–82.

Pryke, S. R. 2010. Sex chromosome linkage of mate preference and color signal maintains

assortative mating between interbreeding finch morphs. Evolution 64:1301–1310.

Quadagno, B. Y. D. M., and E. M. Banks. 1970. The effect of reciprocal cross fostering on the

behaviour of two species of rodents, *Mus musculus* and *Baiomys taylori ater*. Anim. Behav.

18:379–390.

Ramsey, J., H. D. Bradshaw, and D. W. Schemske. 2003. Components of reproductive isolation

between the monkeyflowers *Mimulus lewisii* and *M. cardinalis* (Phrymaceae). Evolution

57:1520–1534.

Robbins, L. W., M. H. Smith, M. C. Wooten, and R. K. Selander. 1985. Biochemical

polymorphism and its relationship to chromosomal and morphological variation in

*Peromyscus leucopus* and *Peromyscus gossypinus*. J. Mammal. 66:498–510.

Roehrs, Z. P., J. B. Lack, C. E. S. Jr., C. J. Seiden, R. Bastarache, W. D. Arbour, D. M. L. Jr.,

and R. A. Van Den Bussche. 2012. Mammals of Red Slough Wildlife Management Area,

with comments on McCurtain county, Oklahoma. Texas Tech Univ. Occas. Pap. 309:1-24.

Rolán-Alvarez, E., and A. Caballero. 2000. Estimating sexual selection and sexual isolation

effects from mating frequencies. Evolution 54:30–36.

Sæther, S. A., G.-P. Sætre, T. Borge, C. Wiley, N. Svedin, G. Andersson, T. Veen, J. Haavie, M.

R. Servedio, S. Bures, M. Král, M. B. Hjernquist, L. Gustafsson, J. Träff, and A.

Qvarnström. 2007. Sex chromosome-linked species recognition and evolution of

reproductive isolation in flycatchers. Science 318:95–97.

Schellinck, H. M., A. M. West, and R. E. Brown. 1992. Rats can discriminate between the urine

odors of genetically identical mice maintained on different diets. Physiol. Behav. 51:1079–

1082.

Schug, M. D., S. H. Vessey, and E. M. Underwood. 1992. Paternal behavior in a natural

population of mice (*Peromyscus leucopus*). Am. Midl. Nat. 127:373–380.

Seehausen, O., J. M. van A. Jacques, and F. Witte. 1997. Cichlid fish diversity threatened by

eutrophication that curbs sexual selection. Science 277:1808–1811.

Servedio, M. R., S. a. Sæther, and G.-P. Sætre. 2009. Reinforcement and learning. Evol. Ecol.

23:109–123.

Servedio, M. R., G. S. Van Doorn, M. Kopp, A. M. Frame, and P. Nosil. 2011. Magic traits in

speciation: "magic" but not rare? Trends Ecol. Evol. 26:389–397.

Slagsvold, T., B. T. Hansen, L. E. Johannessen, and J. T. Lifjeld. 2002. Mate choice and

imprinting in birds studied by cross-fostering in the wild. Proc. R. Soc. B. 269:1449–1455.

Smadja, C., and G. Ganem. 1998. Subspecies recognition in the house mouse: a study of two

populations from the border of a hybrid zone. Behav. Ecol. 13:312–320.

Smith, B. A., and M. L. Block. 1991. Male saliva cues and female social choice in Mongolian

gerbils. Physiol. Behav. 50:379–384.

Smith, M. H. 1965. Behavioral discrimination shown by allopatric and sympatric males of

*Peromyscus eremicus* and *Peromyscus californicus* between females of the same two

884 species. Evolution 19:430–435.

885 Soltis, D. E., A. B. Morris, J. S. McLachlan, P. S. Manos, and P. S. Soltis. 2006. Comparative

886  phylogeography of unglaciated eastern North America. Mol. Ecol. 15:4261–4293.

887 Talley, H. M., C. M. Laukaitis, and R. C. Karn. 2001. Female preference for male saliva:

888  implications for sexual isolation of *Mus musculus* subspecies. Evolution 55:631–634.

889 Taylor, R. J., and H. McCarley. 1963. Vertical distribution of *Peromyscus leucopus* and *P.*

890  *gossypinus* under experimental conditions. Southwest. Assoc. Nat. 8:107–115.

891 ten Cate, C., and G. Mug. 1984. The development of mate choice in zebra finch females.

892  Behaviour 90:125–150.

893 ten Cate, C., M. N. Verzijden, and E. Etman. 2006. Sexual imprinting can induce sexual

894  preferences for exaggerated parental traits. Curr. Biol. 16:1128–1132.

895 ten Cate, C., and D. R. Vos. 1999. Sexual imprinting and evolutionary processes in birds: a

896  reassessment. Pp. 1–31 in P. J. Slater, J. S. Rosenblatt, C. T. Snowdon, and T. J. Roper, eds.

897  Mate Choice. Academic Press.

898 Turner, L. M., D. J. Schwahn, and B. Harr. 2011. Reduced male fertility is common but highly

899  variable in form and severity in a natural mouse hybrid zone. Evolution 66:443–458.

900 Verzijden, M. N., R. F. Lachlan, and M. R. Servedio. 2005. Female mate-choice behavior and

901  sympatric speciation. Evolution 59:2097–2108.

902 Verzijden, M. N., and G. G. Rosenthal. 2011. Effects of sensory modality on learned mate

903  preferences in female swordtails. Anim. Behav. 82:557–562.

904 Verzijden, M. N., and C. ten Cate. 2007. Early learning influences species assortative mating

905  preferences in Lake Victoria cichlid fish. Biol. Lett. 3:134–136.

906 Verzijden, M. N., J. Zwinkels, and C. ten Cate. 2009. Cross-fostering does not influence the mate

preferences and territorial behaviour of males in Lake Victoria cichlids. Ethology 115:39–

48.

Vos, D. R. 1995. Sexual imprinting in zebra-finch females: do females develop a preference for

males that look like their father? Ethology 99:252–262.

Weber, J. N., M. B. Peters, O. V Tsyusko, C. R. Linnen, C. Hagen, N. A. Schable, T. D.

Tuberville, A. M. McKee, S. L. Lance, K. L. Jones, H. S. Fisher, M. J. Dewey, H. E.

Hoekstra, and T. C. Glenn. 2010. Five hundred microsatellite loci for *Peromyscus*. Conserv.

Genet. 11:1243–1246.

Wilson, R. C., T. Vacek, D. L. Lanier, and D. A. Dewsbury. 1976. Open-field behavior in

muroid rodents. Behav. Biol. 506:495–506.

Wolfe, J. L., and A. V Linzey. 1977. *Peromyscus gossypinus*. Mamm. Species 70:1–5.

Yamazaki, K., M. Yamaguchi, L. Baranoski, J. Bard, E. A. Boyse, and L. Thomas. 1979.

Recognition among mice. Evidence from the use of a Y-maze differentially scented by

congenic mice of different major histocompatibility types. J. Exp. Med. 150:755–760.

Yukilevich, R., and J. R. True. 2008. Incipient sexual isolation among cosmopolitan *Drosophila*

*melanogaster* populations. Evolution 62:2112–2121.

**Figure 1.** Hybridization is extremely rare between sympatric *P. leucopus* and *P. gossypinus* mice. **(A)** Genetic PCA discriminates between species. The first eigenvector strongly separates species based on the values of known allopatric *P. leucopus* (green dots) and *P. gossypinus* (blue dots) mice. The second eigenvector detects population structure among *P. gossypinus* populations that could correspond to mice collected east (higher values) and west (lower values) of the Mississippi river. Known lab-generated F1 and backcross (F1 x *P. gossypinus*) hybrids (cyan dots) fall intermediate along the first eigenvetor. Mice collected from the sympatric range overlap (grey dots) cluster discretely with *P. leucopus* or *P. gossypinus* with the exception of two mice that may be hybrids (arrows), but showing greater *P. leucopus* ancestry. **(B)** Range map of the two species: *P. leucopus* (green) and *P. gossypinus* (blue) adapted from (Hall and Kelson 1959; Hall 1981), showing areas of allopatry and sympatry. Pie diagrams show collecting locations and frequencies of each species scaled in size to represent the number of mice sampled at each site. Mice were classified as *P. leucopus* (green dots), *P. gossypinus* (blue dots), or potential hybrids (cyan dots) based on the genetic PCA (shown in **A**). The two possible hybrids were collected at "Big Lake" wildlife management area in Louisiana.

Figure 2

**Figure 2.** Latency to mating between *P. leucopus* (L), *P. gossypinus* (G) and their hybrids (F1). Estimated days since copulation are shown for conspecific, heterospecific, and backcross mating pairs that produced offspring (sample size in parentheses) in no-choice assays. F1 hybrids were generated with both LxG and GxL crosses. In all pairs, female individual is listed first. ** p = 0.01.

Figure 3

**A**

**B**

**C**

**Figure 3**. Mating preferences in two-way choice trials. **(A)** Photograph of the mate-choice apparatus. Center chamber is connected to two test chambers, each housing a "stimulus" animal, separated by gated doors activated by only the "chooser" animal. **(B)** Mating preferences for mice raised by their own parents. *P. leucopus* spent greater time with P. leucopus stimuli than both *P. gossypinus* sexes. **(C)** Mating preferences for mice raised by heterospecific foster parents. *P. leucopus* males were strongly affected cross-fostering (p = 0.004), whereas *P. leucopus* females were not (asterisks). Both *P. gossypinus* sexes spent significantly more time with the heterospecific stimulus than when raised by their own parents (p < 0.001).

Figure 4

**Figure 4.** The proportion of time spent with a stimulus predicts mating outcome in trials when mating occurred. Mating occurred in 12 trials where choosers were raised with their own parents (gray dots) and 8 trials in where choosers were raised with heterospecific parents (black dots). Dotted line indicates the predicted probability for mate choice (conspecific versus heterspecific) given the proportion of time a chooser spent with a conspecific individual. The proportion of time a chooser individual spent with the conspecific stimulus strongly predicts the mating partner (p = 0.038). With the exception of one *P. leucopus* female raised with her own parents, all mice spent more time with their preferred mate.