Transcriptome and histone epigenome of *Plasmodium vivax* salivary-gland sporozoites point to tight regulatory control and potential mechanisms for liver-stage differentiation.

Vivax Sporozoite Consortium[*] (Ivo Muller[1,2,3], Aaron R. Jex[1,3,4], Stefan H. I. Kappe[5], Sebastian A. Mikolajczak[5], Jetsumon Sattabongkot[7], Rapatbhorn Patrapuvich[6], Scott Lindner[8], Erika L. Flannery[5], Cristian Koepfli[1], Brendan Ansell[4], Anita Lerch[1], Samantha J Emery-Corbin[1], Sarah Charnaud[1], Jeffrey Smith[1], Nicolas Merrienne[2], Kristian E. Swearingen[5], Robert L. Moritz[9], Michaela Petter[10,11], Michael Duffy[10], Vorada Chuenchob[5]).

[*]**Group authorship – all authors are equal contributors (order per author contributions section below).**

1. Population Health and Immunity Division, The Walter and Eliza Hall Institute for Medical Research, 1G Royal Parade, Parkville, Victoria, 3052, Australia.
2. Malaria: Parasites & Hosts Unit, Institut Pasteur, 28 Rue de Dr. Roux, 75015, Paris, France.
3. Department of Medical Biology, The University of Melbourne, Victoria, 3010, Australia.
4. Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Corner of Park and Flemington Road, Parkville, Victoria, 3010, Australia.
5. Center for Infectious Disease Research, 307 Westlake Avenue North, Suite 500, Seattle, WA 98109, USA;
6. Department of Global Health, University of Washington, Seattle, WA 98195, USA.
7. Mahidol *Vivax* Research Center, Faculty of Tropical Medicine, Mahidol University, Bangkok 10400, Thailand.
8. Department of Biochemistry and Molecular Biology, Center for Malaria Research, Pennsylvania State University, University Park, PA 16802, USA.
9. Institute for Systems Biology, Seattle, WA, 98109, USA.
10. Department of Medicine Royal Melbourne Hospital, The Peter Doherty Institute, The University of Melbourne, 792 Elizabeth Street, Melbourne, Victoria 3000, Australia.
11. Institute of Microbiology, University Hospital Erlangen, Erlangen 91054, Germany

ABSTRACT

Plasmodium vivax is the key obstacle to malaria elimination in Asia and Latin America, largely attributed to its ability to form resilient hypnozoites (sleeper-cells) in the host liver that escape treatment and cause relapsing infections. The decision to form hypnozoites is made early in the liver infection and may already be set in sporozoites prior to invasion. To better understand these early stages of infection, we undertook a comprehensive transcriptomic and histone epigenetic characterization of *P. vivax* sporozoites. The salivary-gland sporozoite transcriptome is heavily composed of transcripts associated with functions needed for early infection of the vertebrate host and development within hepatocytes. Through comparisons to recently published proteome data for the *P. vivax* sporozoite, our study finds that although highly transcribed, these transcripts are not detectable as proteins and may be regulated through translational repression; a finding we test for a small subset of transcripts and proteins through immunofluorescent microscopy of sporozoites and liver stages in humanized mice. We identify differential transcription between the sporozoite and published transcriptomes of asexual blood-stages and mixed versus hypnozoite-enriched liver stages. These comparisons point to multiple layers of transcriptional, post-transcriptional and post-translational control that appear active in sporozoites and to a lesser extent hypnozoites, but largely absent in replicating liver schizonts or mixed blood-stages. Common transcripts up-regulated in sporozoites and hypnozoites compared to mixed (i.e., schizont) liver-stages identify genes linked to dormancy/persistence in bacteria, amoebae and plants. We also characterise histone epigenetic modifications in the *P. vivax* sporozoite and explore their role in regulating transcription. Collectively, these data support the hypothesis that the sporozoite as a tightly programmed stage primed to infect the human host and identifies potential mechanisms for hypnozoite-formation that may be further explored in liver stage models.

INTRODUCTION

Malaria is among the most significant infectious diseases impacting humans globally, with 3.3 billion people at risk of infection, 381 million suspected clinical cases and up to ~660,000 deaths attributed to malaria in 2014 [1]. Two major parasitic species contribute to the vast majority of human malaria, *Plasmodium falciparum* and *P. vivax*. Historically, *P. falciparum* has attracted the majority of global attention, due to its higher contribution to morbidity and

mortality. However, *P. vivax* is broadly distributed, more pathogenic than previously thought, and is recognised as the key obstacle to malaria elimination in the Asia-Pacific and Americas [2]. Unlike *P. falciparum*, *P. vivax* can establish long-lasting 'sleeper-cells' (= hypnozoites) in the host liver that emerge weeks, months or years after the primary infection (= relapsing malaria) [3]. Primaquine is the only approved drug that prevents relapse. However, the short half-life, long dosage regimens and incompatibility of primaquine with glucose-6-phosphate-dehydrogenase deficiency (which requires pre-screening of recipients [4]) makes it unsuitable for widespread use. As a consequence, *P. vivax* is overtaking *P. falciparum* as the primary cause of malaria in a number of co-endemic regions [5]. Developing new tools to diagnose, treat and/or prevent hypnozoite infections is considered one of the highest priorities in the malaria elimination research agenda [6].

When *Plasmodium* sporozoites are deposited by an infected mosquito, they likely traverse the skin cells, enter the blood-stream and are trafficked to the host liver, as has been shown in rodents [7]. The sporozoites' journey from skin deposition to hepatocytes takes less than a few minutes [8]. Upon reaching the liver, sporozoites traverse Kupffer and endothelial cells to reach the parenchyma, moving through several hepatocytes before invading a final hepatocyte suitable for development [7, 9]. Within hepatocytes, these parasites replicate, and undergo further development and differentiation to produce merozoites that emerge from the liver and infect red blood cells. However, *P. vivax* sporozoites are able to commit to two distinct developmental fates within the hepatocyte: they either immediately continue development as replicating schizonts and establish a blood infection, or delay replication and persist as hypnozoites. Regulation of this major developmental fate decision is not understood and this represents a key gap in current knowledge of *P. vivax* biology and control.

Sporozoites prepare for mammalian host infection while still residing in the mosquito salivary glands. It has been hypothesized that *P. vivax* sporozoites exist within an inoculum as replicating 'tachysporozoites' and relapsing 'bradysporozoites' [10] and that these subpopulations may have distinct developmental fates as schizonts or hypnozoites, thus contributing to their relapse phenotype [10-12]. This observation is supported by the stability of different hypnozoite phenotypes (ratios of hypnozoite to schizont formation) in *P. vivax* infections of liver-chimeric mouse models [13]. To determine fates in the sporozoite stage control of protein expression must take place. Studies using rodent malaria parasites have identified genes [14] that are transcribed in sporozoites but translationally repressed (i.e., present as transcript but un- or under-represented as protein), via RNA-binding proteins [15], and ready for immediate translation after the parasites' infection of the mammalian host cell [13, 16]. It is therefore also possible that translational repression (i.e., the blocking of translation of present and retained transcripts) and other mechanisms of epigenetic control may contribute to the *P. vivax* sporozoite fate decision and hypnozoite formation, persistence and activation. Supporting this hypothesis, histone methyltransferase inhibitors stimulate increased activation of *P. cynomolgi* hypnozoites to become schizonts in macaque hepatocytes [17, 18]. Epigenetic control of stage development is further evidenced in *Plasmodium* through chromatin structure controlling expression of PfAP2-G, a specific transcription factor that, in turn, regulates gametocyte (dimorphic sexual stages) development in blood-stages [19]. It is well documented that *P. vivax* hypnozoite activation patterns stratify with climate and geography [11] and recent modelling suggests transmission potential selects for hypnozoite phenotype [20]. Clearly the ability for *P. vivax* to dynamically regulate hypnozoite formation and relapse phenotypes in response to high or low transmission periods in different climate conditions would confer a significant evolutionary advantage.

Unfortunately, despite recent advances [21] current approaches for *in vitro P. vivax* culture do not support routine maintenance in the laboratory and tools to directly perturb gene function are not established. This renders studies on *P. vivax*, particularly its sporozoites and liver stages, exceedingly difficult. Although *in-vitro* liver stage assays and humanised mouse models are being developed [13], 'omics analysis of *P. vivax* liver stage dormancy has until recently [22] been impossible and even now is in its early stages. Recent characterization [23] of liver-stage (hypnozoites and schizonts) of *P. cynomolgi* (a related and relapsing parasite in macaques) provides valuable insight, but investigations in *P. vivax* directly are clearly

needed. The systems analysis of *P. vivax* sporozoites that reside in the mosquito salivary
glands and are poised for transmission and liver infection offer a key opportunity to gain
insight into *P. vivax* infection. *Plasmodium vivax* sporozoites have been explored previously
by microarray [24] and most recently, in a single RNA-seq replicate [25] and a study on
sporozoite activation {Roth, 2018 #66}. Epigenetic regulation in sporozoites has only been
explored in *P. falciparum* [26, 27]. Here, we present a detailed characterization of the *P. vivax*
sporozoite transcriptome and histone epigenome and use these data to better understand this
key infective stage and the role of sporozoite programming in invasion and infection of the
human host, and development within the host liver.
123
RESULTS AND DISCUSSION
Mosquito infections were generated by membrane feeding of blood samples taken from *P.*
vivax infected patients in western Thailand (n = 9). Approximately 3-15 million *P. vivax*
sporozoites were harvested per isolate from *Anopheles dirus* salivary glands. Using RNA-seq,
we detected transcription for 5,714 *P. vivax* genes (based on the *P. vivax* P01 gene models:
[28]) and obtained a high degree of coverage (4,930 with a mean counts per million (CPM) $\geq$
1.0; Figure S1 and Table S1 and S2). Among the most highly transcribed genes in the
infectious sporozoite stage are *csp* (c̲ircums̲porozoite p̲rotein), five *etramps* (e̲arly t̲ranscribed
m̲embrane p̲roteins), including *uis3* (u̲p-regulated in i̲nfective s̲porozoites), *uis4* and *lsap-1*
(l̲iver s̲tage a̲ssociated p̲rotein 1), a variety of genes involved in cell transversal and initiation
of invasion, including *celtos* (c̲el̲l t̲raversal protein for o̲okinetes and s̲porozoites), *gest*
(g̲amete e̲gress and s̲porozoite t̲raversal protein), *spect1* (s̲porozoite p̲rotein e̲ssential for c̲ell
t̲raversal) and *siap-1* (s̲porozoite i̲nvasion a̲ssociated p̲rotein), and genes associated with
translational repression (*alba1*, *alba4* and *Puf2*). Collectively, these genes account for >1/3rd
of all transcripts in the sporozoite. Although we found only moderate agreement (R^2 = 0.35;
Figure S2) between our RNA-seq data and previous microarray data for *P. vivax* sporozoites
and blood-stages [24], improved transcript detection and quantitation is expected with the
increased technical resolution of RNA-seq over microarray. Supporting this, we find higher
correlation between RNA-seq data from *P. vivax* and *P. falciparum* (single replicate
sequenced herein for comparative purposes) sporozoite datasets (R^2 = 0.42), compared to
either species relative to published microarray data (Figure S2 and Table S3).
Although microarray supports the high transcription in sporozoites of genes such as
uis4, *csp*, *celtos* and several other *etramps*, 27% and 16% of the most abundant 1% of
transcribed genes in our sporozoite RNA-seq data are absent from the top decile or quartile
respectively in the existing *P. vivax* sporozoite microarray data [24]. Among these are genes
involved in early invasion/hepatocyte development, such as *lsap-1*, *celtos*, *gest* and *siap-1*, or
translational repression (e.g., *alba-1* and *alba-4*); orthologs of these genes are also in the top
percentile of transcripts in RNA-seq (see [26, 29]) and previous microarray data [30, 31] for
human-infecting *P. falciparum* and murine-infecting *P. yoelii* sporozoites, suggesting many
are indeed more abundant than previously characterized. A subset of representative
transcripts, including *Pv_AP2-X* (PVP01_0733100), *d13*, *gest*, *g10* (PVP01_1011100), 40S
ribosomal protein S27 (PVP01_1409300), *puf-2*, *zipco* and 14-3-3, were tested by qPCR for
their transcript abundance relative to *celtos* and *sera* (Figure 1A and Table S4). This
representative set differed markedly in their relative abundance between our RNAseq and
previous microarray data [24]. To control for batch effects introduced by collection of the
sporozoites used here for RNAseq, this testing was conducted in an additional six sample
replicates representing four additional clinical *P. vivax* isolates (PvSPZ-Thai13-16; with
PvSPZ-Thai16 tested in technical triplicate). The qPCR results agreed with the RNAseq data
for these transcripts (Table S4).
163
***Transcription in* P. vivax *relative to other plasmodia* sporozoites.** To gain insight into
species-specific aspects of the *P.vivax* transcriptome, we qualitatively compared these data
with available data for *P. falciparum* [27] and *P. yoelii* sporozoites (single replicate only) for
4,067 single-copy orthologs (SCO) (transcribed at $\geq$ 1 TPM in *P. vivax* infectious
sporozoites) shared with *P. falciparum* and *P. yoelli* (Table S5). Genes highly transcribed in

salivary-gland sporozoites of all three species include *celtos*, *gest*, *trap*, *siap1*, *spect1* and
puf2. There are 696 *P. vivax* genes shared as orthologs between *P. vivax* P01 and *P. vivax*
Sal1 lacking a defined SCO in *P. falciparum* or *P. yoelli* transcribed at a mean of ≥ 1 TPM in
P. vivax salivary-gland sporozoites (Table S6). Prominent among these are *vir* (n=25) and *Pv-*
fam (41 fam-e, 16 fam-b, 14 fam-a, 8 fam-d and 3 fam-h) genes, as well as hypothetical
proteins or proteins of unknown function (n=212) and, interestingly, a number of 'merozoite
surface protein' 3 and 7 homologs (n=5 of each). Both *msp3* and *msp7* have undergone
significant expansion in *P. vivax* relative to *P. falciparum* and *P. yoelii* [32] and may have
repurposed functions in sporozoites. In addition, there are 69 *P. vivax* P01 genes lacking a
defined ortholog in *P. vivax* Sal1, *P. falciparum* or *P. yoelli* transcribed at ≥ 1 TPM in
infectious *P. vivax* sporozoites; most of which are *Plasmodium* interspersed repeat (PIR)
genes [32] found in telomeric regions of the P01 assembly and likely absent from the Sal1
assembly but present in the Sal1 genome, indicating the improved coverage of telomeric
regions in P01 relative to Sal1.

***P. vivax* sporozoites transcriptome compared with proteome.** We compared relative
protein abundance presented in a recently published *P. vivax* sporozoite proteome [33] to
relative transcript abundance from the current study (Figure 1B and Table S7). The proteome
study incorporated data from the same PvSPZ-Thai1 and PvSPZ-Thai5 isolates tested by
RNAseq here. We identified 2,402 *P. vivax* genes transcribed in the sporozoite (CPM > 1) for
which no protein expression was detected. Although many of these are lowly transcribed and
likely below the detection sensitivity of LC-MS proteomics, others are among the most highly
transcribed genes in the sporozoite, indicating these may be under translational repression.
Translational repression, the mechanism through which transcripts are held in stasis
by RNA binding proteins, has been demonstrated to have important functional roles in the
transition of *Plasmodium* spp. between the vertebrate to invertebrate host. More than 700
genes have been identified as translationally repressed in *Plasmodium berghei* ('rodent
malaria') gametocytes based on DOZI (DEAD box RNA helicase "<u>d</u>evelopment <u>o</u>f <u>z</u>ygote
<u>i</u>nhibited") pulldowns [34]. Translational repression mechanisms mediated through Puf-2
have been explored in sporozoites of several *Plasmodium* species and regulate some of the
most abundant transcripts in the sporozoite, such as *uis-3* and *uis-4*. UIS3 and UIS4 are the
best characterized proteins under translational repression by Puf-2 in sporozoites [35] and are
essential for liver-stage development [14].
In considering genes that may be translationally repressed (i.e., transcribed but not
translated) in the *P. vivax* sporozoite, we confine our observations to those transcripts
representing the top decile of transcript abundance to ensure their lack of detection as proteins
was not due to limitations in the detection sensitivity of the proteomic dataset. Approximately
1/3rd of transcripts in the top decile of transcriptional abundance (n = 170 of 558) in *P. vivax*
sporozoites were not detectable as peptides in multiple replicates (**Figure 1B** and Table S7).
Of these 170 putatively repressed transcripts, 156 and 154 have orthologs in *P. falciparum*
and *P. yoelii* respectively, with 89 and 118 of these also not detected as proteins in *P.*
falciparum and *P. yoelii* salivary-gland sporozoites [36] despite being highly transcribed in
these stages (see [26, 29]; Tables S3-S5), and 133 (78.2%) having no detectable sporozoite
expression (>1 unique peptide count) in LC-MS data deposited for any species in PlasmoDB
(Table S8). In contrast, 106 of these putatively repressed transcripts with orthologs in other
Plasmodium species (Table S8) for which proteomic data is available in PlasmoDB, are
detectable (>1 unique peptide count) by LC-MS methods in at least one other life-cycle stage,
indicating against a technical issue (e.g., inability to be trypsin-digested) preventing their
detection in the *P. vivax* sporozoite proteome [33]. In addition to *uis3* and *uis4*, genes
involved in liver stage development and detectable as transcripts but not proteins in the *P.*
vivax sporozoites include *lsap1* (<u>l</u>iver <u>s</u>tage <u>a</u>ssociated <u>p</u>rotein 1), *zipco* (<u>ZIP</u> domain-
<u>c</u>ontaining protein), several other *etramps* (PVP01_1271000, PVP01_0422600,
PVP01_0504800 and PVP01_0734800), *pv1* (<u>p</u>arasitophorous <u>v</u>acuole protein 1) and *lisp1*
and *lisp2* (PVP01_1330800 and PVP01_0304700). Also notable among genes detectable as
transcripts but not proteins in sporozoites is a putative 'Yippee' homolog (PVP01_0724100).

Yippee is a DNA-binding protein that, in humans (YPEL3), suppresses cell growth [37] and is regulated through histone acetylation [38], making it noteworthy in the context of *P. vivax* hypnozoite developmental arrest.

Although verifying each putatively repressed transcript will require further empirical data, our system level approach is supported by immunofluorescent microscopy (Figure 1C) of UIS4, LISP1, EXP1 and ACP (PVP01_0416300). These represent one known and three putative (i.e., newly proposed here) translationally repressed genes in *P. vivax* sporozoites, and are compared to TRAP and BiP (which are both transcribed and expressed as protein in the *P. vivax* sporozoite; Table S8). The *lisp1* gene is an interesting find. In *P. berghei*, *lisp1* is essential for rupture of the PVM during liver stage development allowing release of the merozoite into the host blood stream. *Pv-lisp1* is ~350-fold and ~1,350-fold more highly transcribed in *P. vivax* sporozoites compared to sporozoites of either *P. falciparum* or *P. yoelii* (see Table S5). IFAs using LISP1 specific mAbs (Figure 1C) show that this protein is undetectable in sporozoites but clearly expressed at 7 days post-infection in liver schizonts.

Up-regulated transcripts in *P. vivax* sporozoites relative to other life-cycle stages. Recently completed studies of the transcriptome of *P. vivax* for sporozoite activation [39], as well as, liver [22] and asexual blood-stages [40] support comparative transcriptomic study of sporozoites, their biology and transcriptional regulation over the *P. vivax* life-cycle. Recently published data for activate sporozoites from Roth et al [39] was significantly lower depth coverage, with ~0.03 to 0.6M reads mapped the *P. vivax* P01 coding domains; compared with 0.7 to 15.3 M, 2.4 to 10.6 M and 18.7 to 57.6M mapped reads for salivary sporozoite, liver-stages [22] and asexual blood stages [40] respectively. This lower coverage could not be compensated for through data normalization and therefore data from Roth et al [39] was not included in our quantitative analyses, although qualitatively, many of the highly transcribed genes in Roth et al [39] sporozoites were among the highly transcribed genes in salivary sporozoites from the present study. The remaining RNAseq data presents an analytical challenge in that each (sporozoites, liver-stages and blood stages) is produced in a separate study and may be influenced by technical batch effects that cannot be differentiated from biologically meaningful changes. To address this, we first examined *P. vivax* transcripts in a previous microarray study of multiple *P. vivax* life-cycle stages [24], including sporozoites and several blood-stages, to identify genes that may be transcriptionally stable across the life-cycle. We identified ~160 genes with low transcriptional variability between sporozoites and blood-stages that covered the breadth of transcript abundance levels in Westenberger et al [24]. These include genes typically associated with "house-keeping" functions, such as ribosomal proteins, histones, translation initiation complex proteins and various chaperones (see Figure S3 and Table S8). We assessed transcription of these 160 genes among the current and recently published RNA-seq data for *P. vivax* and all were of similarly low variability (Figure S4). This suggests that any batch effect between the studies is sufficiently lower than the biological differences between each life-cycle stage, allowing informative comparisons. We then combined all published RNAseq-based, transcriptomic data available for *P. vivax* [22, 39, 40] with the salivary sporozoite data generated here.

P. vivax *sporozoite relative to blood-stage transcriptome.* To identify transcripts up-regulated in sporozoites, we first compared the *P. vivax* sporozoite transcriptome to RNA-seq data for *P. vivax* blood-stages [40] (Figure 2 and Figures S7-S10). We identified 1,672 up (Table S9); Interactive Glimma Plot - Supplementary Data 1) and 1,958 down-regulated (Table S9); Interactive Glimma Plot - Supplementary Data 1) transcripts (FDR $\leq$ 0.05; minimum 2-fold change in Counts per Million (CPM)) and explored patterns among these differentially transcribed genes (DTGs) by protein family (Figure 2C and Table S10) and Gene Ontology (GO) classifications (Table S11). RNA recognition motifs (RRM-1 and RRM-6) and helicase domains (Helicase-C and DEAD box helicases) are over-represented (p-value <0.05) among transcripts up-regulated in sporozoites, consistent with translational repression through ribonucleoprotein (RNP) granules [41]. Transcripts encoding nucleic acid binding domains, such as bromodomains (PF00439; which can also bind lysine-acetylated

proteins), zinc fingers (PF13923) and EF hand domains (PF13499) are also enriched in sporozoites. Included among these proteins are a putative ApiAP2 transcription factor (PVP01_1211900) and a homologue of the *Drosophila* zinc-binding protein 'Yippee' (PVP01_0724100). Thrombospondin-1 like repeats (TSR: PF00090) and von Willebrand factor type A domains (PF00092) are enriched in sporozoites as well. In *P. falciparum* sporozoites, genes enriched in TSR domains are important in invasion of the mosquito salivary gland (e.g., *trap*) and secretory vesicles released by sporozoites upon entering the vertebrate host (e.g., *csp*) [42]. By comparison, genes up-regulated in blood-stages are enriched for *vir* gene domains (PF09687 and PF05796), Tryptophan-Threonine-rich *Plasmodium* antigens (PF12319; which are associated with merozoites [43]), markers of cell-division (PF02493; [44]) protein production/degradation (PF00112, PF10584, PF00152, PF09688 and PF00227) and ATP metabolism (PF08238 and PF12774). 47 of the 343 transcripts unique to *P. vivax* sporozoites relative to *P. falciparum* or *P. yoelii* are up-regulated in sporozoites compared to *P. vivax* blood stages. Nine of these are in the top decile of transcription, and include a Pv-fam-e (PVP01_0525200), a Pf-fam-b homolog (PVP01_0602000) and 7 proteins of unknown function. A further nine have an ortholog in *P. cynomolgi* (which also forms hypnozoites) but not the closely related *P. knowlesi* (which does not form hypnozoites) and include '*msp7*'-like (PVP01_1219600, PVP01_1220300 and PVP01_1219900), '*msp3*'-like (PVP01_1031300), Pv-fam-e genes (PVP01_0302100, PVP01_0524500 and PVP01_0523400), a serine-threonine protein kinase (PVP01_0207300) and a RecQ1 helicase homolog (PVP01_0717000). Notably, the *P. cynomolgi* ortholog of PVP01_0207300, PCYB_021650, is transcriptionally up-regulated in hypnozoites relative to replicating schizonts [23], indicating a target of significant interest when considering hypnozoite formation and/or biology and suggesting that the list here may contain other genes important in hypnozoite biology.

P. vivax *sporozoites are enriched in translational repressors.* In *Plasmodium*, translational repression regulates key life-cycle transitions coinciding with switching between the mosquito and the mammalian host (either as sporozoites or gametocytes) [41]. For example, although *uis4* is the most abundant transcript in the infectious sporozoite ([24, 31]; Table S2), UIS4 is translationally repressed in this stage [15] and only expressed after hepatocyte invasion [45]. In sporozoites, it is thought that PUF2 binds to mRNA transcripts and prevents their translation [29], and SAP1 stabilises the repressed transcripts and prevents their degradation [45]. Consistent with this, *Puf2* and *SAP1* (PVP01_0947600) are up-regulated in the sporozoite relative to blood-stages. Indeed, *Puf2* (PVP01_0526500) is among the top percentile of transcripts in salivary sporozoites and expressed at high levels in the proteome [33]. However, our data implicate other genes that may act in translational repression in *P. vivax* sporozoites, many of which are already known to be involved in translational repression in other *Plasmodium* stages and other protists [41]. Among these are *alba-2* and *alba-4*, both of which are among the top 2% of genes transcribed in sporozoites and ~14 to 20-fold more highly transcribed in sporozoites relative to blood-stages; ALBA-2 is in the top 100 most abundant proteins in the *P. vivax* sporozoite proteome [33]. In addition, *P. vivax* sporozoites are enriched for genes encoding RRM-6 RNA helicase domains. Intriguing among these are HoMu (homolog of Musashi; top decile of sporozoite proteins by abundance [33]) and ptbp (polypyrimidine tract binding protein). Musashi regulates eukaryotic stem cell differentiation through translational repression [46] and HoMu localizes with DOZI and CITH in *Plasmodium* gametocytes [47]. PTBP is linked to mRNA stability, splice regulation and translational initiation [48] and may perform a complementary role to SAP1.

P. vivax *sporozoite relative to Plasmodium spp. liver stage transcriptomes.* New advances in *P. vivax* liver culture has allowed recent publication of mixed stage and hypnozoite-enriched transcriptomes [22]. This is an early, yet highly valuable, study and, due no doubt to the difficulty in generating the material, is limited to biological duplicates. Noting this, although we undertake differential transcriptomic studies of this dataset here, we recognize that additional biological replication is needed and have used a higher burden of significance

(FDR ≤ 0.01 and ≥ 2-fold change) than used with blood-stages. Nevertheless, these
comparisons identified 1,015 and 856 sporozoite up-regulated transcripts relative to mLS and
HPZs respectively and 1,007 and 1,079 transcripts up-regulated in mLS and HPZs relative to
sporozoites respectively (Figures 3 and S11-S13, Table S12 and S13 and Interactive Glimma
Plot - Supplementary Data 1).

Compared to mLS transcriptomes, sporozoites are enriched for many of the
transcripts similarly up-regulated in comparison to blood-stages (e.g., *uis4*, *celtos*, *puf2*, *siap1*
and *plp-1*). More broadly, SPZ up-regulated transcripts over-represent (p-value ≤ 0.05) Pfam
domains (Figure 3A) associated with transcriptional regulation (PF00176, PF01096, PF01661
and PF08711), translational repression/regulation (PF00076, PF00279, PF01008, PF01873,
PF01917, PF02847, PF02854, PF13893 and PF14259), DNA/RNA binding (PF0097,
PF13445, PF13639, PF14570 and PF15247) and chromatin regulation (PF00271, PF00850
and PF13489). In contrast, the mLS transcriptome is enriched in genes involved in replication
and merozoite formation [n = 14; including PVP01_0728900 (*msp1*), PVP01_0010670
(*msp3*) and PVP01_1446800 (*msp9*)], rhoptry function [n = 9; including PVP01_1469200
(*rnp3*), PVP01_1255000 (*rnp2*) and PVP01_1338500 (*rap1*)] and reticulocyte binding [n=10
including PVP01_0534300 (rbp2c), PVP01_1402400 (rbp2a), PVP01_0701100 (rbp1b) and
PVP01_0800700 (rbp2b)]. These data are further enriched for Pfam domains associated with
cell division (PF00493), merozoite formation (PF07133 and PF12984), proteasome function
(PF00227, PF00400, PF00656, PF01344, PF01398 and PF03981), protein export / vesicle
function (PF00350 and PF00996), membrane proteins (PF01105, PF03011, PF05424 and
PF12139) and metabolism (PF00085, PF00118, PF00268, PF01066, PF01214 and PF01214).
Collectively, in addition to markers consistent with sporozoite or merozoite formation, these
data point towards the sporozoite stage as being highly regulated and controlled at
transcriptional, translational and chromatin levels, with the mLS stages representing a release
of this control allowing replication, protein turn-over, reconfiguration of the proteins on the
plasma membrane and metabolic activity.

Comparison of sporozoites with HPZs does not indicate a similar release of control,
or at least that any release is more specific than for mLS. The sporozoite is enriched, relative
to HPZs, in genes such as PVP01_1258000 (*gest*), PVP01_0418000 (*sera*), PVP01_1435400
(*celtos*), PVP01_0835600 (*csp*) and PVP01_0602100 (*uis4*). At a broad level, sporozoite
enriched Pfam domains include a smaller number associated with translational
repression/regulation (PF00076) or DNA/RNA binding (PF01428 and PF12756).
Interestingly, sporozoites are enriched in Pfam domains specifically associated with
heterochromatin (H3K9me3) reading/interaction (PF02463, PF00628, PF13831 and
PF13865). Ours (see below) and previous epigenetic studies of *Plasmodium* sporozoites [27]
find dense heterochromatin in the telomeric to subtelomeric regions of the chromosome,
which is more transcriptionally active in blood-stages [49]. Others have noted an up-
regulation of methyl/acetyltransferases in *P. cynomolgi* HPZs [23] and/or shown
methyltransferase inhibitors stimulate hypnozoite activation *in vitro* [17]. The potential that
histone epigenetics of sporozoites has a role in or changes with liver-stage development and
the formation of liver schizonts or HPZs is intriguing but requires detailed study of the
chromatin of liver-stage parasites, which is not presently available for *P. vivax*. In contrast,
HPZs were enriched, relative to sporozoites, for genes including histone proteins
(PVP01_1138700, PVP01_1131700 and PVP01_0905900) and classic markers of metabolism
(PVP01_MITO3300 and PVP01_MITO3400) and *lisp2*. Pfam data indicated– largely similar
domain enrichment trends as were seen for the mLS stage relative to sporozoites, including a
number of proteosomal (PF00227, PF00112, PF03981), vesicular transport (PF00996) and
metabolic (PF00118, PF00268, PF01066, PF01214 a) associated functions. This supports
HPZs being an arrested, rather than classically 'dormant', stage with active metabolism and
protein turn-over. HPZs are also enriched for Pfams associated with mRNA/tRNA regulation
and turnover (PF04857, PF01612, PF00009 and PF01138) and glycine metabolism (PF01571
and PF00464) and acetyl-CoA production (PF02779 and PF00676).

Finally, although not the focus of this study, we looked at differential transcription
between mLS and HPZ stages using the Gural et al [22] data, but using the same approaches

as employed here. In particular, we were interested in what these comparisons might provide
in terms of sporozoite differentiation or development into liver schizonts or HPZs (Table
S14). Among mLS up-regulated transcripts are genes associated with rhoptry function (n =
11; including PVP01_0107500, PVP01_1469200 and PVP01_1469200), cytoadherence to
red-cells (PVP01_1401400 and PVP01_0734500), merozoite formation (PVP01_0728900
and PVP01_0612400) and exported proteins (n = 6; including PVP01_0504000,
PVP01_0119200 and PVP01_0801600). Consistent with *P. cynomolgi* [23], HPZ up-
regulated transcripts include several key sporozoite transcripts, specifically *uis4*
(PVP01_0602100), *puf1* (PVP01_1015000) and *speld* (PVP01_0938800). At the Pfam
domain level, mLS is enriched for metabolic (PF00317) and proteosomal (PF00112) domains
also enriched in mLS or HPZs relative to sporozoites above, as well as domains associated
with merozoite formation (PF12948, PF07462), rhoptry function (PF0712), DNA/RNA
binding (PF12756, PF10601 and PF02151) and cell division, development and DNA
replication (PF06705, PF00533, PF00488, PF02460, PF07034, PF02181). In contrast, HPZs
are enriched in Pfam domains that overlap notably with key sporozoite markers, including
etramps (PF09716) and *puf* proteins (PF00806), as well as domains associated with calcium
(PF08683) and nucleotide metabolism (PF06437). These data largely indicate that the
hypnozoite bears similarity both to the sporozoite and liver schizonts consistent with a stalled
stage on the path to schizont development regulated by checkpoint signals that halt/restart
normal schizont development, which has been proposed previously for this species [24].
With this is mind, we looked at transcripts that are differentially transcribed in mLS,
but not HPZs, relative to SPZs. There are 107 transcripts down-regulated in mLS relative to
SPZs that are transcribed at roughly similar levels in both SPZs and HPZs (Figure 3B). A
common theme among many of these genes are their role in transcriptional, post-
transcriptional, translational or post-translational regulation. Among transcriptional regulators
are transcription factors including AP2-SP2 (PVP01_0303400) and three non-AP2-like
transcription factors (PVP01_0306600, PVP01_0204300 and PVP01_1415800). Post-
transcriptional controllers include several DNA/RNA-binding proteins (PVP01_1011000.
PVP01_0932900, PVP01_0715300, PVP_1242600 and PVP01_0605200), RNA helicases
(PVP01_1403600 and PVP01_1329800) and mRNA processing (PVP01_1443100 and
PVP01_1458200) genes. Translational control includes several key regulators of translation
initiation (PVP01_1467700), tRNA processing (PVP01_0318700 and PVP01_1017700) or
ribosomal function/biogenesis (PVP01_1443700, PVP01_0421400, PVP01_1117200 and
PVP01_0215100). Post-translational control includes two methyltransferases
(PVP01_1428800 and PVP01_1465200), including CARM1, which methylates of H3R17
and, in mice, prevents differentiation in embryonic stem cells [50], and a putative histone
methylation reading enzyme, EEML2 (PVP01_1014100). The remaining genes in this group
have three noteworthy and largely overlapping themes: (1) an association with calcium
binding, metabolism or signalling, (2) a role in organellar metabolism and (3) homologs in
other organisms, including a variety of prokaryotes and eukaroytes, with key roles in
germination, dormancy and persistent non-replicating stages. The latter most function is
clearly intriguing in the context of HPZ formation and activation. These genes include a
homolog of dihydrolipoamide acyltransferase (aka 'sucB'), which is essential for growth in
Mycobacterium tuberculosis [51] and a key regulator in persistent *Escherichia coli* stages
[52]. Another example is gamete fusion factor HAP2, which, despite the name, has been
shown to regulate dormancy in eukaryotes ranging from plants [53, 54] to amoebae [55].
In addition to data for *P. vivax*, two transcriptomic studies are now available for *P.
cynomolgi* [27, 56] that compare mixed/schizont stage parasites with small-form
"hypnozoites". In comparing *P. cynomolgi* liver-stage RNA-seq and *P. vivax* liver-stage
microarray data, Cubi et al [23] noted a moderate to good level of agreement (R^2 = 0.50) as
evidence of *P. cynomolgi* being predictive and representative of *P. vivax*. However, Voorberg
van der Wel et al [56] explored congruence between their and the Cubi et al [23] studies and
found generally good agreement among schizonts and overall relatively poor agreement
among hypnozoites from each study. This highlights the complexity of these datasets and
indicates caution in comparing the current data to *P. vivax*. ApiAP2 transcription factors

feature prominently in each liver-stage transcriptomic study for *P. cynomolgi* [23, 56] and *P. vivax* [22]. Cubi et al [23] noted an ApiAP2 (dubbed "AP2-Q"; PCYB_102390) as transcriptionally up-regulated in *P. cynomolgi* hypnozoites and proposed this as a potential hypnozoite marker. We note that the *P. vivax* ortholog of *Pc*-AP2-Q (PVP01_1016100) is among the genes detectable as a transcript but not protein in *P. vivax* sporozoites. This may point to a translationally repressed signal in sporozoites to regulate hypnozoite formation. However, as *Pv*-AP2-Q is transcribed at an abundance (~50 TPM) at or below which ~50% of *P. vivax* genes are detectable as transcripts but not as proteins (Figure 1B), this could as likely result from LC-MS detection sensitivity. Further, although AP2-Q was reported as specific to hypnozoite forming *Plasmodium* species [23], it is indeed found in non-hypnozoite producing species, such as *P. knowlesi*, *P. gallinaceum* and *P. inui* [56]. Up-regulation of AP2-Q transcripts is not observed for hypnozoites in subsequent transcriptomic studies of *P. cynomolgi* [56] or *P. vivax* [22], nor do we see such an up-regulation here. Voorberg van der Wel et al [56] note transcription of a range of AP2s in *P. cynomolgi* liver stages, but do not find any to be up-regulated in hypnozoites. AP2s also feature among transcribed genes in *P. vivax* liver stages, with one, PVP01_0916300, significantly up-regulated in hypnozoites. We note that PVP01_0916300 is up-regulated in *P. vivax* sporozoites relative to blood-stages and found in the top quartile of transcripts by abundance (TPM = 104).

Chromatin epigenetics in *P. vivax* sporozoites. As noted above, transcriptomic data for sporozoites, and their comparison with liver and blood-stages, implicate histone epigenetics as having an important role in sporozoite biology and liver stage differentiation. This concept has been alluded to in recent liver-stage studies of *P. cynomolgi* [17, 23] that propose methyltransferases as having a potential role in hypnozoite formation. No epigenetic data are currently available for any *P. vivax* life-cycle stage. Studies of *P. falciparum* blood-stages have identified the importance of histone modifications as a primary epigenetic regulator [57, 58] and characterized key markers of heterochromatin (H3K9me3) and euchromatin/transcriptional activation (H3K4me3 and H3K9ac). Recently, these marks have been explored with the maturation of *P. falciparum* sporozoites in the mosquito [26]. Here, we characterize major histone marks in *P. vivax* sporozoites and assess their relationship to transcript abundance.

*Histone modifications in **P. vivax** sporozoites.* Using ChIP-seq, we identified 1,506, 1,999 and 5,262 ChIP-seq peaks stably represented in multiple *P. vivax* sporozoite replicates and associated with H3K9me3, H3K9ac and H3K4me3 histone marks respectively (Figure 4 and S14-S19). Peak width, spacing and stability differed with histone mark type (Figures S15 and S16). H3K4me3 covered the greatest breadth of the genome (36.0% of all bases) and was the most stable among replicates, with ~84% of bases associated with an H3K4me3 found in multiple biological replicates. By comparison H3K9me3 marks were least stable, with 46% of bases associated with this mark found in just one replicate. Consistent with observations in *P. falciparum*, H3K9me3 'heterochromatin' marks primarily clustered in telomeric and subtelometric regions (**Figure 4**). In contrast, the 'euchromatin' / transcriptionally open histone marks, H3K4me3 and H3K9ac, were distributed in chromosome central regions and did not overlap with regions under H3K9me3 suppression. Both H3K9me3 and H3K4me3 marks were reasonably uniformly distributed (mean peak spacing ~500bp for each) within their respective regions of the genome. In contrast, H3K9ac peaks were spaced further apart (mean: ~2kb), but also with a greater variability in spacing (likely reflecting their association with promoter regions [59]; Figure S17 and S19). The instability of H3K9me3 may reflect its use in *Plasmodium* for regulating expression of contingency genes from multigene families, whose members have overlapping and redundant functions [49] and confer phenotypic plasticity [60].

Genes under histone regulation. We explored an association between these histone marks and the transcriptional behaviour of protein coding genes (Figure 4 and S19 and Tables S15-S20). 485 coding genes stably intersected with an H3K9me3 mark; all are located near the

ends of the chromosomal scaffolds (i.e., are (sub)telomeric). On average, these genes are transcribed at ~30 fold lower levels (mean 0.7 TPM) than genes not stably intersected by H3K9me3 marks. These data clearly support the function of this mark in transcriptional silencing. This is largely consistent with observations in *P. falciparum* sporozoites [26], however, in contrast to *P. falciparum* sporozoites where a single *var* gene was described to lack heterochromatin structure [27] we observe no genes within heterochromatin dense region that lacked a stable H3K9me3 signal or were transcribed at notable levels (i.e., above ~5 TPM). Whether this relates to differences in epigenetic control between the species is not clear. We note that (sub)telomeric genes are overall transcriptionally silent in *P. vivax* sporozoites relative to blood-stages (Figure 4 and Tables S21 and S22). Consistent with observations in *P. falciparum* [57], the bulk of these genes include complex protein families, such as *vir* and *Pv-fam* genes, which are so far described to function primarily in blood-stages. Also notable among the genes are several reticulocyte-binding proteins, including RBP2, 2a, 2b and 2c. This transcriptional silence in telomeric and subtelomeric regions was recently observed in *P. falciparum* sporozoites [27].

Outside of the telomeres and subtelomeres, H3K4me3 marks are stably associated with the intergenic regions of 3,676 genes. H3K9ac marks are also identified within 1kb of the transcriptional start site (TSS) of 1,284 coding genes, with 1085 of these stably marked also by H3K4me3 (Figure 4B). The average transcription of these genes is 50, 112 and 112 TPMs respectively (72, 160 and 160-fold higher than H3K9me3 marked genes). Gene-by-gene observations show that H3K9ac and H3K4me3 marks cluster densely in the 1000kb up and down-stream of the start and stop codon respectively of transcribed genes, but are much less dense within coding regions of these genes (Figure S15). This pattern directly correlates with transcription and contrasts H3K9me3 marks, which are distributed across the length of the gene at even density and are correlated with a lack of transcription. These data support the role of H3K9ac and H3K4me3 in transcriptional activation in *P. vivax*. The lower transcriptional abundance of H3K4me3 marked, compared with H3K9ac or H3K9ac and H3K4me3 marked genes suggest these marks work synergistically and that H3K9ac is possibly the better of the two, as a single mark indicator of transcriptional activity in *P. vivax*. This is consistent with recent observations in *P. falciparum* sporozoites [26].

Interestingly, H3K9ac-marked genes ranged in transcriptional activity from the most abundantly transcribed genes to many in the lower 50% and even lowest decile of transcription. This suggests more contributes to transcriptional activation in *P. vivax* sporozoites than, simply, gene accessibility through chromatin regulation. Specific activation by a transcription factor (e.g., ApiAP2s [61]) is the obvious candidate. To explore this, we compared upstream regions (within 1kb of the TSS or up to the 3' end of the next gene upstream, whichever was less) of highly (top 10%) and lowly (bottom 10%) transcribed H3K9ac marked genes for over-represented sequence motifs in the highly expressed genes that might coincide with known ApiAP2 transcription factor binding sites [62]. We identified these based on the location of the nearest stable H3K9ac peak relative to the transcription start site for each gene (Figure S12). In most instances, these peaks were within 100bp of the TSS and, consistent with data from *P. falciparum* [59], *P. vivax* promoters appear to be no more than a few hundred to a maximum of 1000 bp upstream of the TSS. Exploring these regions, we identified two over-represented motifs: TGTACMA (e-value $2.7e^{-2}$) and ATATTTH (e-value $3.3e^{-3}$) (Fig. 2D). TGTAC is consistent with the known binding site for *Pf*-AP2-G, which regulates sexual differentiation in gametocytes [63], but its *P. vivax* ortholog (PVP01_1418100) is neither highly transcribed nor expressed in sporozoites. ATATTTH is similar to the binding motif for *Pf*-AP2-L (AATTCC), a transcription factor that is important for liver stage development in *P. berghei* [64]. In contrast to AP2-G, *Pv*-AP2-L (PVX_081180) is in the top 10% of transcription and expression in *P. vivax* sporozoites and up-regulated relative to blood-stages. In *P. vivax* sporozoites, the ATATTTH motif is associated with a number of highly transcribed genes, including *lisp1* and *uis2-4*, known to be regulated by AP2-L in *P. berghei* [64] as well as many of the most highly transcribed, H3K9ac marked genes, including two *etramps* (PVP01_0734800 and PVP01_0504800), several RNA-binding proteins, including *Puf2*, *ddx5*, a putative ATP-

dependent RNA helicase DBP1 (PVP01_1429700), and a putative *bax1* inhibitor (PVP01_1465600). Interestingly, a number of highly transcribed and translationally repressed genes associated with the ATATTTH motif, including *uis4*, *siap2* and *pv1*, are not stably marked by H3K9ac in all replicates (i.e., there is significant variation in the placement of the H3K9ac peak or their presence/absence among replicates for these genes). It may be that additional histone modifications, for example H3K27me, H3R17me3 or H2A or H4 modifications, are involved in regulating transcription of these genes. Certainly the incorporation of the H2A.Z histone variant, which is present in intergenic regions of *P. falciparum (*Petter et al 2011*)*, and controls temperature responses in plants [65] is intriguing as a potential mark regulating sporozoite fate in *P. vivax* considering the association between hypnozoite activation rate and climate [11], as is H3R17me3 in consideration of the enrichment of markers/readers of this modification in HPZs noted above and the role of this mark in cell fate progression in other species [50].

CONCLUSIONS

We provide the first comprehensive study of the transcriptome and epigenome of mature *Plasmodium vivax* sporozoites and undertake detailed comparisons with recently published proteomic data for *P. vivax* sporozoites [33] and transcriptomic data for *P. vivax* mixed and hypnozoite-enriched liver-stages [22] and mixed blood-stages [40]. These data support the proposal that the sporozoite is a highly-programmed stage that is primed for invasion of and development in the host hepatocyte. Cellular regulation, including at transcription, translational and epigenetic levels, appears to play a major role in shaping this stage (which continues on in some form in hypnozoites), and many of the genes proposed here as being under translational repression are involved in hepatocyte infection and early liver-stage development (Figure 5). We highlight a major role for RNA-binding proteins, including PUF2, ALBA2/4 and, intriguingly, 'Homologue of Musashi' (HoMu). We find that transcriptionally, the hypnozoite appears to be a transition point between the sporozoite and replicating schizonts, having many of the dominant sporozoite transcripts and retaining high transcription of a number of key regulatory pathways involved in transcription, translation and chromatin configuration (including histone arginine methylation). A consistent theme in the study is the prominence of a number of genes that have a role in numerous eukaryotic systems in cell fate determination and differentiation (e.g., HoMu, Yippee and CARM1) and overlap with dormancy and/or persistent cell states in bacteria, protists or higher eukaryotes (e.g., bacterial sucB and gamete fusion protein HAP2). These data do not point to one single programming switch for dormancy or liver developmental fate in *P. vivax*, but present a number of intriguing avenues for exploration in subsequent studies, particularly in model species such as *P. cynomolgi*. Our study contributes to understanding the early stages of hepatocyte infection and the developmental switch between liver trophozoite and hypnozoite formation. We also identify potential avenues for rationally prioritizing targets underpinning liver-stage differentiation for functional evaluation in humanized mouse and simian models for relapsing *Plasmodium* species and identifying novel avenues to understand and eradicate liver-stage infections.

MATERIALS AND METHODS

Ethics Statement. Collection of venous blood from human patients with naturally acquired vivax infection for the current study was approved by the Ethical Review Committee of the Faculty of Tropical Medicine, Mahidol University (Human Subjects Protocol number TMEC 11-033) with the informed written consent of each donor individual. All mouse tissue used in the current study was from preserved infected tissues generated previously [13]. All mouse infection work in [13] was carried out at the Centre for Infectious Diseases Research (CIDR) in Seattle, USA, under direct approval of the CIDR Institutional Animal Care and Use Committee (IACUC) and performed in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health, USA. The Centre for Infectious Disease Research Biomedical Research Institute has an Assurance

608 from the Public Health Service (PHS Assurance number is A3640-01) through the Office of
609 Laboratory Animal Welfare (OLAW) for work approved by its IACUC.
610
Material collection, isolation and preparation. Nine field isolates (PvSpz-Thai 1 to 9),
representing symptomatic blood-stage malaria infections were collected as venous blood (20
mL) from patients presenting at malaria clinics in Tak and Ubon Ratchatani provinces in
Thailand. Each isolate was used to establish, infections in *Anopheles dirus* colonized at
Mahidol University (Bangkok) by membrane feeding [13], after 14-16 days post blood
feeding, ~3-15 million sporozoites were harvested per field isolate from the salivary glands of
up to 1,000 of these mosquitoes as per [66] and shipped in preservative (trizol (RNA/DNA) or
1% paraformaldehyde (DNA for ChiP-seq) to the Walter and Eliza Hall Institute (WEHI).
619
Transcriptomics sequencing and differential analysis. Upon arrival at WEHI, messenger
RNAs were purified from an aliquot (~0.5-1 million sporozoites) of each *P. vivax* field isolate
as per [40] and subjected to RNA-seq on Illumina NextSeq using TruSeq library construction
chemistry as per the manufacturer's instructions. Raw reads for each RNA-seq replicate are
available through the Sequence Read Archive (XXX-XXX). Sequencing adaptors were
removed and low quality reads trimmed and filtered using Trimmomatic v. 0.36 [67]. To
remove host contaminants, processed reads were aligned, as single-end reads, to the
Anopholes dirus wrari2 genome (VectorBase version W1) using Bowtie2 [68] (--very-
sensitive preset). All non-host reads were then aligned to the manually curated transcripts of
the *P. vivax* P01 genome (http://www.genedb.org/Homepage/P*vivax*P01; [28]) using RSEM
[69] (pertinent settings: --bowtie2 --bowtie2-sensitivity-level very_sensitive --calc-ci --ci-
memory 10240 --estimate-rspd --paired-end). Transcript abundance for each gene in each
replicate was calculated by RSEM as raw count, posterior mean estimate expected counts
(pme-EC) and transcripts per million (TPM).
Transcriptional abundance in *P. vivax* sporozoites was compared qualitatively (by
ranked abundance) with previously published microarray data for *P. vivax* salivary-gland
sporozoites [24]. As a further quality control, these RNA-seq data were compared also with
previously published microarray data for *P. falciparum* salivary-gland sporozoites [30], as
well as RNA-seq data from salivary-gland sporozoites generated here for *P. falciparum*
(single replicate generated from *P. falciparum* 3D7 lab cultures isolated from *Anopholes*
stephensi and processed as above) and previously published for *P. yoelii* [29]. RNA-seq data
from these additional *Plasmodium* species were (re)analysed from raw reads and
transcriptional abundance for each species was determined (raw counts and pme-EC and TPM
data) as described above using gene models current as of 04-10-2016 (PlasmoDB release
v29). Interspecific transcriptional behaviour was qualitatively compared by relative ranked
abundance in each species using TPM data for single copy orthologs (SCOs; defined in
PlasmoDB) only, shared between *P. vivax* and *P. faliciparum* or shared among *P. vivax*, *P.*
falciparum and *P. yoelii*.
To define transcripts that were up-regulated in sporozoites, we remapped raw reads
representing early (18-24 hours post-infection (HPI)), mid (30-40 HPI) and late (42-46 HPI)
P. vivax blood-stage infections recently published by Zhu *et al* [40] to the *P. vivax* P01
transcripts using RSEM as above. All replicate data was assessed for mapping metrics,
transcript saturation and other standard QC metrics using QualiMap v 2.1.3 [70]. Differential
transcription between *P. vivax* salivary-gland sporozoites and mixed blood-stages [40] was
assessed using pme-EC data in EdgeR [71] and limma [72] (differential transcription cut-off:
$\geq$ 2-fold change in counts per million (CPM) and a False Discovery Rate (FDR) $\leq$ 0.05).
Pearson Chi squared tests were used to detect over-represented Pfam domains and Gene
Ontology (GO) terms among differentially transcribed genes in sporozoites (Bonferroni-
corrected $p < 0.05$), based on gene annotations in PlasmoDB (release v29).
We also compared transcription of the sporozoite stages to recently published liver-
stage data from Gural et al [22] as per the sporozoite to blood-stage comparisons above, with
the following modifcations: (1) EC values were normalized using the 'upper quartile' method
instead of TMM, (2) differential transcription was assessed using a quasi-likelihood

generalize linear model (instead of a linear model) and (3) an FDR threshold for significance
of $\leq$ 0.01 was used instead of $\leq$ 0.05. These differences related to specific attributes of the
liver-stage dataset, particularly the small number of replicates (n = 2) per experiment
condition. Data visualization and interactive R-shiny plots were produced in R using the
ggplot[73], ggplot2 [74], gplots(heatmap.2) [75] and Glimma [76] packages.
668
Assessment of Sporozoite RNA-seq transcriptome by selective RT-qPCR: Extracted RNA
was DNase treated (Sigma D5307) as per manufacturer recommendations. RNA was
quantified using the TapeStation High Sensitivity RNA kit (Agilent). Two intron-spanning
primer pairs were designed per gene of interest using Primer3 and BLAST. Primer pairs were
tested in two concentrations (0.75ng and 2.83ng per reaction) to determine efficiency and
specificity. Product was run on a 1% agarose gel with ethidium bromide. Primer pairs
indicating non-specific priming were removed. The resulting 11 primer pairs were used on
four sporozoite samples; VUBR06, VUNL23, VUBR24, VTTY84. RNA was reverse
transcribed (Sensifast, Bioline) and used at 0.75ng per reaction, run on a Roche LightCycler
480 II. Melt curves were assessed and products were run on a gel to ensure specificity again.
Cp threshold was set automatically. ΔCp value was calculated as target gene – comparator
gene (SERA and CelTOS were used). Data were log transformed and fold change calculated.
681
682 RT-qPCR Primers were as follows:

Name	Gene	Forward Primer	Reverse Primer
RPS27	PVX_122245	ACCACCTTGTTTAGCCATGC	TAATTTGCACTTTCCACCCGTT
D13	PVX_089510	CTGTACACGCACGAGCTGGC	CAGCTCCTTGACGCCACTG
G10	PVX_080110	ACGAGCTGTACTACAAGCGGA	TTTCTCCTGCACCAGGTAGTC
AP2	PVX_086995*	GCCCCACTGGAAGTTTTGGA	CGTTCAGCCGCTGGTAGTAT
SERA	PVX_003790	CTGAAGACCTCCAGGGACAAG	TTTCTGCCTCTCCAGTGATATCTTT
CelTOS	PVX_123510*	CCCCCAAAGGCAAAATGAACAA	CGCTCTTTCCCCTCAAGGAC
GEST	PVX_118040	GACATATCAAGCAGTGAGGGA	CATGTTGTGGCCTTTATATGCTG
ALBA4	PVX_083270	TATCAACGGAGCCTTTGCCG	GGACTTGATTTCCTCGTCGG
PUF2	PVX_089945	ATCATAGAGAACGTCGACAAGCTTA	CTACGTTTCCAGGTTGCTGATC
14-3-3	PVX_089505	GACAACTTGACCTTGTGGACGTC	TACTCGAGGCCTTCATCCTTCGATT
ZIPCO	PVX_001980	TTAGCTCAATTGCTTGTGGCTTTTT	TGCCACTAACTCCAAGGAAATAACT

683
684 * denotes single exon gene
685
Salivary-gland sporozoite and liver-stage immunofluorescence assays (IFAs). IFAs were
performed as per [13] using preserved, vivax infected mouse liver tissue generated previously
for that study. In [13], female FRG [fumarylacetoacetate hydrolase (F), recombination
activation gene 2 (R), interleukin-2 receptor subunit gamma (G)] triple KO mice engrafted
with human hepatocytes (FRG KO huHep) were purchased from Yecuris Corporation
(Oregon, USA). Mice were infected through intravenous injection into the tail with $3.5 \times$
10^5 to 1×10^6 sporozoites isolated from the salivary glands of infected mosquitoes in 100 µl
of RPMI media. Liver stages for the current study were obtained from 10µm formalin fixed
paraffin embedded day 7 liver stages generated previously [13] from FRG knockout huHep
mice;[13] these were deparaffinized prior to staining. Fresh salivary-gland sporozoites were
fixed in acetone per [13]. All cells were incubated twice for 3 minutes in Xylene, then 100%
Ethanol, and finally once for 3 minutes each in 95%, 70%, and 50% Ethanol. The cells were
rinsed in DI water and permeabilized immediately in 1XTBS, containing Triton X-100 and
30% hydrogen peroxide. The cells were blocked in 5% milk in 1XTBS. The hepatocytes were
stained overnight with a rabbit polyclonal LISP1 antibody (A), a rabbit polyclonal UIS4

antibody (B), and a rabbit polyclonal BIP antibody (C) in blocking buffer. The cells were
washed with 1XTBS and the primary antibodies were detected with goat anti-rabbit Alexa
Fluor 488 antibody (Life Technologies). The cells were washed in 1XTBS. The hepatocytes
were rinsed in KMNO4 and washed in 1XTBS. The cells were incubated in DAPI for 5
minutes.
706
Histone ChIP sequencing and analysis. Aliquots of 2 – 6 million freshly isolated
sporozoites were fixed with 1% paraformaldehyde for 10 min at 37°C and the reaction
subsequently quenched by adding glycine to a final concentration of 125 mM. After three
washes with PBS, sporozoite pellets were stored at -80°C and shipped to Australia. Nuclei
were released from the sporozoites by dounce homogenization in lysis buffer (10 mM Hepes
pH 7.9, 10 mM KCl, 0.1 mM EDTA, 0.1 mM EDTA, 1 mM DTT, 1x EDTA-free protease
inhibitor cocktail (Roche), 0.25% NP40). Nuclei were pelleted by centrifugation at 21,000 g
for 10 min at 4°C and resuspended in SDS lysis buffer (1% SDS, 10 mM EDTA, 50 mM Tris
pH 8.1, 1x EDTA-free protease inhibitor cocktail). Chromatin was sheared into 200–1000 bp
fragments by sonication for 16 cycles in 30 sec intervals (on/off, high setting) using a
Bioruptor (Diagenode) and diluted 1:10 in ChIP dilution buffer (0.01% SDS, 1.1% Triton X-
100, 1.2 mM EDTA, 16.7 mM Tris pH 8.1, 150 mM NaCl). Chromatin was precleared for 1
hour with protein A/G sepharose (4FastFlow, GE Healthcare) equilibrated in 0.1% BSA
(Sigma-Aldrich, USA) in ChIP dilution buffer. Chromatin from 3×10^5 nuclei was taken
aside as input material. Chromatin from approximately 3×10^6 sporozoite nuclei was used for
each ChIP. ChIP was carried out over night at 4°C with 5 µg of antibody (H3K9me3 (Active
Motif), H3K4me3 (Abcam), H3K9ac (Upstate), H4K16ac (Abcam)) and 10 µl each of
equilibrated protein A and G sepharose beads (4FastFlow, GE Healthcare). After washes in
low-salt, high-salt, LiCl, and TE buffers (EZ-ChIP Kit, Millipore), precipitated complexes
were eluted in 1% SDS, 0.1 M NaHCO3. Cross-linking of the immune complexes and input
material was reversed for 6 hours at 45°C after addition of 500 mM NaCl and 20 µg/ml of
proteinase K (NEB). DNA was purified using the MinElute® PCR purification kit (Qiagen)
and paired-end sequenced on Illumina NextSeq using TruSeq library construction chemistry
as per the manufacturer's instructions. Raw reads for each ChIP-seq replicate are available
through the Sequence Read Archive (XXX-XXX).
Fastq   files   were   checked   for   quality   using   fastqc
(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and adapter sequences were
trimmed using cutadapt [77]. Paired end reads were mapped to the *P. vivax* P01 strain
genome annotation using Bowtie2 [68]. The alignment files were converted to Bam format,
sorted and indexed using Samtools [78]. ChIP peaks were called relative to input using
MACS2[79] in paired end mode with a q value less than or equal to 0.01. Peaks and peak
summits were converted to sorted BED files. Bedtools intersect[80] was used to identify
genes that intersected H3K9me3 peaks and Bedtools closest was used to identify genes that
were closest to and downstream of H3K9ac and H3K4me3 peak summits.
741
Sequence motif analysis. Conserved sequence motifs were identified using the program
DREME [81]. Only genes in the top decile of transcription showing no evidence of protein
expression in multiple salivary-gland sporozoite replicates were considered as putatively
translationally repressed (n = 170). We queried coding regions and regions upstream of the
transcriptional start site (TSS) for each gene, defined by Zhu *et al* [40] and/or predicted here
from all RNA-seq data using the Tuxedo suite [82], for enriched sequence motifs in
comparison to 170 genes found to be in the top decile of both transcriptional and expressional
abundance in the same sporozoite replicates. In searching for motifs associated with highly
transcribed genes with stable H3K9ac marks within 1kb of the TSS (or up to the 3' end of the
next gene upstream), we compared H3K9ac marked genes in the top decile of transcription to
the same number of H3K9ac marked genes in the bottom decile of transcription. In both
instances, an e-value threshold of 0.05 was considered the minimum threshold for statistical
significance.
755

References

1. Organization WH. World Malaria Report 2015. WHO, Geneva. 2015.

2. Feachem RG, Phillips AA, Hwang J, Cotter C, Wielgosz B, Greenwood BM, et al. Shrinking the malaria map: progress and prospects. Lancet. 2010;376 9752:1566-78. doi:10.1016/S0140-6736(10)61270-6.

3. Price RN, Douglas NM and Anstey NM. New developments in *Plasmodium vivax* malaria: severe disease and the rise of chloroquine resistance. Curr Opin Infect Dis. 2009;22 5:430-5. doi:10.1097/QCO.0b013e32832f14c1.

4. Baird KJ. Malaria caused by *Plasmodium vivax*: recurrent, difficult to treat, disabling, and threatening to life - averting the infectious bite preempts these hazards. Pathogens and global health. 2013;107:475-9.

5. Sattabongkot J, Tsuboi T, Zollner GE, Sirichaisinthop J and Cui L. *Plasmodium vivax* transmission: chances for control? Trends in parasitology. 2004;20 4:192-8.

6. Mueller I, Galinski MR, Baird JK, Carlton JM, Kochar DK, Alonso PL, et al. Key gaps in the knowledge of *Plasmodium vivax*, a neglected human malaria parasite. Lancet Infect Dis. 2009;9 9:555-66.

7. Lindner SE, Miller JL and Kappe SH. Malaria parasite pre-erythrocytic infection: preparation meets opportunity. Cell Microbiol. 2012;14 3:316-24.

8. Shin SC, Vanderberg JP and Terzakis JA. Direct infection of hepatocytes by sporozoites of *Plasmodium berghei*. J Protozool. 1982;29 3:448-54.

9. Mota MM, Pradel G, Vanderberg JP, Hafalla JC, Frevert U, Nussenzweig RS, et al. Migration of *Plasmodium* sporozoites through cells before infection. Science. 2001;291 5501:141-4.

10. Lysenko AJ, Beljaev A and Rybalka V. Population studies of *Plasmodium vivax*: 1. The theory of polymorphism of sporozoites and epidemiological phenomena of tertian malaria. Bull WHO. 1977;55 5:541.

11. White NJ. Determinants of relapse periodicity in *Plasmodium vivax* malaria. Malar J. 2011;10:297.

12. Price RN, Tjitra E, Guerra CA, Yeung S, White NJ and Anstey NM. Vivax malaria: neglected and not benign. Amer J Trop Med Hyg. 2007;77 6 Suppl:79-87.

13. Mikolajczak SA, Vaughan AM, Kangwanrangsan N, Roobsoong W, Fishbaugher M, Yimamnuaychok N, et al. *Plasmodium vivax* liver stage development and hypnozoite persistence in human liver-chimeric mice. Cell Host Microbe. 2015;17 4:526-35.

14. Mueller A-K, Camargo N, Kaiser K, Andorfer C, Frevert U, Matuschewski K, et al. *Plasmodium* liver stage developmental arrest by depletion of a protein at the parasite–host interface. Proc Nat'l Acad Sci USA. 2005;102 8:3022-7.

15. Silvie O, Briquet S, Muller K, Manzoni G and Matuschewski K. Post-transcriptional silencing of UIS4 in *Plasmodium berghei* sporozoites is important for host switch. Mol Microbiol. 2014;91 6:1200-13.

16. Mackellar DC, O'Neill MT, Aly AS, Sacci JB, Jr., Cowman AF and Kappe SH. *Plasmodium falciparum* PF10_0164 (ETRAMP10.3) is an essential parasitophorous vacuole and exported protein in blood stages. Eukaryot Cell. 2010;9 5:784-94.

17. Dembele L, Franetich JF, Lorthiois A, Gego A, Zeeman AM, Kocken CH, et al. Persistence and activation of malaria hypnozoites in long-term primary hepatocyte cultures. Nat Med. 2014;20 3:307-12.

18. Malmquist NA, Moss TA, Mecheri S, Scherf A and Fuchter MJ. Small-molecule histone methyltransferase inhibitors display rapid antimalarial activity against all blood stage forms in *Plasmodium falciparum*. Proc Nat'l Acad Sci USA. 2012;109 41:16708-13.

19. Josling GA and Llinas M. Sexual development in *Plasmodium* parasites: knowing when it's time to commit. Nat Rev Microbiol. 2015;13 9:573-87.

20. White MT, Karl S, Battle KE, Hay SI, Mueller I and Ghani AC. Modelling the
contribution of the hypnozoite reservoir to *Plasmodium vivax* transmission. Elife.
2014;3.
21. Roobsoong W, Tharinjaroen CS, Rachaphaew N, Chobson P, Schofield L, Cui L, et
al. Improvement of culture conditions for long-term in vitro culture of
Plasmodium vivax. Malaria J. 2015;14 1:1.
22. Gural N, Mancio-Silva L, Miller AB, Galstian A, Butty VL, Levine SS, et al. In vitro
culture, drug sensitivity, and transcriptome of *Plasmodium vivax* hypnozoites.
Cell Host Microbe. 2018;23 3:395-406 e4.
23. Cubi R, Vembar SS, Biton A, Franetich JF, Bordessoulles M, Sossau D, et al. Laser
capture microdissection enables transcriptomic analysis of dividing and
quiescent liver stages of *Plasmodium* relapsing species. Cell Microbiol. 2017.
24. Westenberger SJ, McClean CM, Chattopadhyay R, Dharia NV, Carlton JM,
Barnwell JW, et al. A systems-based analysis of *Plasmodium vivax* lifecycle
transcription from human to mosquito. PLoS Negl Trop Dis. 2010;4 4:e653.
25. Kim A, Popovici J, Vantaux A, Samreth R, Bin S, Kim S, et al. Characterization of P.
vivax blood stage transcriptomes from field isolates reveals similarities among
infections and complex gene isoforms. Sci Rep. 2017;7 1:7761.
26. Gomez-Diaz E, Yerbanga RS, Lefevre T, Cohuet A, Rowley MJ, Ouedraogo JB, et al.
Epigenetic regulation of *Plasmodium falciparum* clonally variant gene expression
during development in *Anopheles gambiae*. Sci Rep. 2017;7:40655.
27. Zanghi G, Vembar SS, Baumgarten S, Ding S, Guizetti J, Bryant JM, et al. A specific
PfEMP1 is expressed in *P. falciparum* sporozoites and plays a role in hepatocyte
infection. Cell Rep. 2018;22 11:2951-63.
28. Auburn S, Bohme U, Steinbiss S, Trimarsanto H, Hostetler J, Sanders M, et al. A
new *Plasmodium vivax* reference sequence with improved assembly of the
subtelomeres reveals an abundance of pir genes. Wellcome Open Res. 2016;1:4.
29. Lindner SE, Mikolajczak SA, Vaughan AM, Moon W, Joyce BR, Sullivan WJ, Jr., et
al. Perturbations of *Plasmodium* Puf2 expression and RNA-seq of Puf2-deficient
sporozoites reveal a critical role in maintaining RNA homeostasis and parasite
transmissibility. Cell Microbiol. 2013;15 7:1266-83.
30. Le Roch KG, Johnson JR, Florens L, Zhou Y, Santrosyan A, Grainger M, et al. Global
analysis of transcript and protein levels across the *Plasmodium falciparum* life
cycle. Genome research. 2004;14 11:2308-18.
31. Mikolajczak SA, Silva-Rivera H, Peng X, Tarun AS, Camargo N, Jacobs-Lorena V, et
al. Distinct malaria parasite sporozoites reveal transcriptional changes that
cause differential tissue infection competence in the mosquito vector and
mammalian host. Mol Cell Biol. 2008;28 20:6196-207.
32. Carlton JM, Adams JH, Silva JC, Bidwell SL, Lorenzi H, Caler E, et al. Comparative
genomics of the neglected human malaria parasite *Plasmodium vivax*. Nature.
2008;455 7214:757-63.
33. Swearingen KE, Lindner SE, Flannery EL, Vaughan AM, Morrison RD,
Patrapuvich R, et al. Proteogenomic analysis of the total and surface-exposed
proteomes of *Plasmodium vivax* salivary gland sporozoites. PLoS Negl Trop Dis.
2017;11 7:e0005791.
34. Guerreiro A, Deligianni E, Santos JM, Silva PA, Louis C, Pain A, et al. Genome-wide
RIP-Chip analysis of translational repressor-bound mRNAs in the *Plasmodium*
gametocyte. Genome Biol. 2014;15 11:493.
35. Silvie O, Briquet S, Müller K, Manzoni G and Matuschewski K.
Post-transcriptional silencing of UIS4 in Plasmodium berghei sporozoites is
important for host switch. Molecular microbiology. 2014;91 6:1200-13.
36. Lindner SE, Swearingen KE, Harupa A, Vaughan AM, Sinnis P, Moritz RL, et al.
Total and putative surface proteomics of malaria parasite salivary gland
sporozoites. Mol Cell Proteomics : MCP. 2013;12 5:1127-43.

37. Kelley KD, Miller KR, Todd A, Kelley AR, Tuttle R and Berberich SJ. YPEL3, a p53-
regulated gene that induces cellular senescence. Cancer Res. 2010;70 9:3566-75.
38. Tuttle R, Simon M, Hitch DC, Maiorano JN, Hellan M, Ouellette J, et al. Senescence-
associated gene YPEL3 is downregulated in human colon tumors. Ann Surg
Oncol. 2011;18 6:1791-6.
39. Roth A, Adapa SR, Zhang M, Liao X, Saxena V, Goffe R, et al. Unraveling the
Plasmodium vivax sporozoite transcriptional journey from mosquito vector to
human host. Sci Rep. 2018;8 1:12183.
40. Zhu L, Mok S, Imwong M, Jaidee A, Russell B, Nosten F, et al. New insights into
the *Plasmodium vivax* transcriptome using RNA-Seq. Sci Rep. 2016;6:20498.
41. Kramer S. RNA in development: how ribonucleoprotein granules regulate the life
cycles of pathogenic protozoa. WIR: RNA. 2014;5 2:263-84.
42. Tucker RP. The thrombospondin type 1 repeat superfamily. Int J Biochem Cell
Biol. 2004;36 6:969-74.
43. Ntumngia FB, Bouyou-Akotet MK, Uhlemann AC, Mordmuller B, Kremsner PG
and Kun JF. Characterisation of a tryptophan-rich *Plasmodium falciparum*
antigen associated with merozoites. Mol Biochem Parasitol. 2004;137 2:349-53.
44. Gubbels MJ, Vaishnava S, Boot N, Dubremetz JF and Striepen B. A MORN-repeat
protein is a dynamic component of the *Toxoplasma gondii* cell division
apparatus. J Cell Sci. 2006;119 Pt 11:2236-45.
45. Aly AS, Lindner SE, MacKellar DC, Peng X and Kappe SH. SAP1 is a critical post-
transcriptional regulator of infectivity in malaria parasite sporozoite stages. Mol
Microbiol. 2011;79 4:929-39.
46. Okano H, Imai T and Okabe M. Musashi: a translational regulator of cell fate.
Journal of Cell Science. 2002;115 7:1355-9.
47. Cui L, Lindner S and Miao J. Translational regulation during stage transitions in
malaria parasites. Annals N Y Acad Sci. 2015;1342 1:1-9.
48. Lasko P. Gene regulation at the RNA layer: RNA binding proteins in intercellular
signaling networks. Sci STKE. 2003;179:RE6.
49. Guizetti J and Scherf A. Silence, activate, poise and switch! Mechanisms of
antigenic variation in *Plasmodium falciparum.* Cell microbiol. 2013;15 5:718-26.
50. Wu Q, Bruce AW, Jedrusik A, Ellis PD, Andrews RM, Langford CF, et al. CARM1 is
required in embryonic stem cells to maintain pluripotency and resist
differentiation. Stem Cells. 2009;27 11:2637-45.
51. Shi S and Ehrt S. Dihydrolipoamide acyltransferase is critical for *Mycobacterium*
tuberculosis pathogenesis. Infect Immun. 2006;74 1:56-63.
52. Ma C, Sim S, Shi W, Du L, Xing D and Zhang Y. Energy production genes sucB and
ubiF are involved in persister survival and tolerance to multiple antibiotics and
stresses in Escherichia coli. FEMS Microbiol Lett. 2010;303 1:33-40.
53. Schrader J, Moyle R, Bhalerao R, Hertzberg M, Lundeberg J, Nilsson P, et al.
Cambial meristem dormancy in trees involves extensive remodelling of the
transcriptome. Plant J. 2004;40 2:173-87.
54. Yazawa K and Kamada H. Identification and characterization of carrot HAP
factors that form a complex with the embryo-specific transcription factor C-
LEC1. J Exp Bot. 2007;58 13:3819-28.
55. Wood FC, Heidari A and Tekle YI. Genetic Evidence for Sexuality in
Cochliopodium (Amoebozoa). J Hered. 2017;108 7:769-79.
56. Voorberg-van der Wel A, Roma G, Gupta DK, Schuierer S, Nigsch F, Carbone W, et
al. A comparative transcriptomic analysis of replicating and dormant liver stages
of the relapsing malaria parasite *Plasmodium cynomolgi.* Elife. 2017;6
57. Lopez-Rubio J-J, Mancio-Silva L and Scherf A. Genome-wide analysis of
heterochromatin associates clonally variant gene regulation with perinuclear
repressive centers in malaria parasites. Cell host & microbe. 2009;5 2:179-90.

58.   Duffy MF, Selvarajah SA, Josling GA and Petter M. Epigenetic regulation of the
918   *Plasmodium falciparum* genome. Brief Funct Genomics. 2014;13 3:203-16.

919   59.   Cui L, Miao J, Furuya T, Li X, Su XZ and Cui L. PfGCN5-mediated histone H3
920   acetylation plays a key role in gene expression in *Plasmodium falciparum*.
921   Eukaryot Cell. 2007;6 7:1219-27.

922   60.   Rovira-Graells N, Gupta AP, Planet E, Crowley VM, Mok S, de Pouplana LR, et al.
923   Transcriptional variation in the malaria parasite *Plasmodium falciparum*.
924   Genome research. 2012;22 5:925-38.

925   61.   De Silva EK, Gehrke AR, Olszewski K, León I, Chahal JS, Bulyk ML, et al. Specific
926   DNA-binding by apicomplexan AP2 transcription factors. Proc Nat'l Acad Sci.
927   2008;105 24:8393-8.

928   62.   Painter HJ, Campbell TL and Llinás M. The Apicomplexan AP2 family: integral
929   factors regulating *Plasmodium* development. Mol Biochem Parasitol. 2011;176
930   1:1-7.

931   63.   Kafsack BF, Rovira-Graells N, Clark TG, Bancells C, Crowley VM, Campino SG, et
932   al. A transcriptional switch underlies commitment to sexual development in
933   human malaria parasites. Nature. 2014;507 7491:248.

934   64.   Iwanaga S, Kaneko I, Kato T and Yuda M. Identification of an AP2-family protein
935   that is critical for malaria liver stage development. PLoS One. 2012;7 11:e47557.

936   65.   Boden SA, Kavanova M, Finnegan EJ and Wigge PA. Thermal stress effects on
937   grain yield in Brachypodium distachyon occur via H2A.Z-nucleosomes. Genome
938   Biol. 2013;14 6:R65.

939   66.   Kennedy M, Fishbaugher ME, Vaughan AM, Patrapuvich R, Boonhok R,
940   Yimamnuaychok N, et al. A rapid and scalable density gradient purification
941   method for *Plasmodium* sporozoites. Malar J. 2012;11:421.

942   67.   Bolger AM, Lohse M and Usadel B. Trimmomatic: a flexible trimmer for Illumina
943   sequence data. Bioinformatics (Oxford, England). 2014;30 15:2114-20.

944   68.   Langmead B and Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat
945   Methods. 2012;9 4:357-9.

946   69.   Li B and Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data
947   with or without a reference genome. BMC bioinformatics. 2011;12 1:323.

948   70.   Okonechnikov K, Conesa A and Garcia-Alcalde F. Qualimap 2: advanced multi-
949   sample quality control for high-throughput sequencing data. Bioinformatics
950   (Oxford, England). 2016;32 2:292-4.

951   71.   Nikolayeva O and Robinson MD. edgeR for differential RNA-seq and ChIP-seq
952   analysis: an application to stem cell biology. Methods Mol Biol. 2014;1150:45-
953   79.

954   72.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers
955   differential expression analyses for RNA-sequencing and microarray studies.
956   Nucleic Acids Res. 2015;43 7:e47-e.

957   73.   Wickham H. ggplot: An implementation of the Grammar of Graphics in R, 2006. R
958   package version 04 0.

959   74.   Wickham H and Chang W. ggplot2: An implementation of the Grammar of
960   Graphics. R package version 07, URL: http://CRAN R-project org/package=
961   ggplot2. 2008.

962   75.   Warnes GR, Bolker B, Bonebakker L, Gentleman R, Huber W, Liaw A, et al. gplots:
963   Various R programming tools for plotting data. R package version. 2009;2 4:1.

964   76.   Law CW, Alhamdoosh M, Su S, Smyth GK and Ritchie ME. RNA-seq analysis is
965   easy as 1-2-3 with limma, Glimma and edgeR. F1000Research. 2016;5.

966   77.   Martin M. Cutadapt removes adapter sequences from high-throughput
967   sequencing reads. EMBnet journal. 2011;17 1:pp. 10-2.

968   78.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
969   Alignment/Map format and SAMtools. Bioinformatics (Oxford, England).
970   2009;25 16:2078-9.

79. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-
based analysis of ChIP-Seq (MACS). Genome Biol. 2008;9 9:R137.
80. Quinlan AR and Hall IM. BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics (Oxford, England). 2010;26 6:841-2.
81. Bailey TL. DREME: motif discovery in transcription factor ChIP-seq data.
Bioinformatics (Oxford, England). 2011;27 12:1653-9.
82. Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, et al. Differential gene
and transcript expression analysis of RNA-seq experiments with TopHat and
Cufflinks. Nat Protoc. 2012;7 3:562-78.

Figure Legends:

Fig. 1 Transcriptional activity of the *P. vivax* sporozoite and evidence for translational
repression. **a** Relative transcript abundance of key marker genes for sporozoites inferred by
RNA-seq and qPCR (here) relative to previously published microarray data [24]; **b** Relative
proportion of genes detectable as transcripts and proteins or transcripts only in RNA-seq and
previously published proteomic data. Dashed line shows cut-off used in the current study for
putatively repressed transcripts. Immunofluorescent staining of select proteins either known
(UIS4) or predicted here (LISP1, EXP1 and ACP) to be translationally repressed in
sporozoites in **c** sporozoite stages. CSP, mTIP as known positive controls and TRAP and BIP
as exerpimental positive controls and **d** liver stages (schizonts) at 7 days post-infection in
HuHep mice. Liver expression of EXP1 and ACP has been demonstrated by IFA in
Mikolajczak et al [13], using the same antibodies as used here.

Fig. 2 Differential transcription between *Plasmodium vivax* salivary-gland sporozoites and
blood-stages. **A** BCV plot showing separation between blood-stage (black) and salivary-gland
sporozoite (red) biological replicates. **B** Volcano plot of distribution of fold-changes (FC) in
transcription between blood-stages and salivary-gland sporozoites relative to statistical
significance threshold (False Discovery Rate (FDR) $\leq$ 0.05). Positive FC represents up-
regulated transcription in the sporozoite stage. **C** Mirror plot showing pFam domains
statistically significantly (FDR $\leq$ 0.05) over-represented in salivary-gland sporozoite up-
regulated (red) or blood-stage up-regulated (black) transcripts. Scale bar truncated for
presentation. * - 55 PRESAN domains are in this dataset. ** - 99 Vir domains are in this
dataset.

Fig. 3 Differential transcription between *P. vivax* sporozoites (SPZ), mixed (mLS) and
hypnozoite (HPZ) enriched liver stages (liver-stage data from Gural et al [22]). **A** Heatmap
comparisons showing summed transcription of enriched Pfam domains in HPZ vs mLS (left),
SPZ vs HPZ (top middle) and SPZ vs mLS (top right) comparisons. All Pfam domains
statistically significantly enriched at p-value 0.05). All transcript data for stage up-regulated
genes at FDR 0.01). **B** Violin box-plot showing relative fold-change differences between SPZ
and HPZ compared with SPZ and mLS for genes down-regulated in mLS compared to SPZ,
but not down-regulated in HPZ compared to SPZ. **C** Ternary heatmap summarizing relative
transcript abundance in each of SPZ, mLS and HPZ stages.

Fig. 4 Histone epigenetics relative to transcriptional behaviour in salivary-gland sporozoites.
a Representative H3K9me3, H3K4me3 and H3K9ac ChIP-seq data (grey) from a
representative chromosome (*P. vivax* P01 Chr5) relative to mRNA transcription in salivary-
gland sporozoites (black) and blood-stages (black). Small numbers to top left of each row
show data range. **b** Salivary-gland sporozoite transcription relative to nearest stable histone
epigenetic marks. Numbers at the top of the figure represent total genes included in each
category. Numbers within in box plot represent mean transcription in transcripts per million
(TPM). **c** Sequence motifs enriched within 1kb upstream of the Transcription Start Site of
highly transcribed (top 10%) relative to lowly transcribed genes associated with H3K9ac
marks in salivary-gland sporozoites. **d** Relative transcription of (sub)telomeric genes in *P.*
vivax salivary-gland sporozoites and blood-stages categorized by gene sets up-regulated in

blood-stages (blue), salivary sporozoites (red) or not stage enriched (grey). Numbers in each box show mean transcription in TPM.

Fig. 5 Schematic of potential mechanisms underpinning development in differentiation of *P. vivax* sporozoites during liver-stage infection as hypnozoites and schizonts. We suggest differentiation programming at different points in development; first, schizont or hypnozoite fate possibly encoded in the sporozoite as epigenetic signals or translationally repressed transcripts; secondly, suppression signals that halt progression of the hypnozoite to schizont stage and support persistence; and finally activation signals signified by a release in chromatin, (post)transcriptional and (post)translational control leading to up-regulation of replication, metabolic and protein export pathways.

DIFFERENTIATION
- **Translational repressors**
  e.g., Puf2, Alba2/4, HOMU, Yippee, Zipco
- **Invasion**
  e.g., Celtos, Gest
- **Liver development**
  e.g., LISP1, LISP2, ETRAMPS, TRAP
- **Transcription factors**
  e.g., AP2-SP2, AP2-L, AP2-Q?

SUPPRESSION
- **Metabolic/replication suppressors**
  e.g., sucB, HAP2, MAK16
- **Histone arginine methylation**
  e.g., CARM1, EEML2
- **Protein translation**
  e.g., eIF-3H, Puf1

ACTIVATION
- **Merozoite development**
  e.g., MSP1, MSP3, MSP9
- **Rhoptry function**
  e.g., RAP1, RNP2, RNP3
- **Reticuocytye binding**
  e.g., RBP2a, RBP2b, RBP2C
- **Exported Proteins**
  e.g., PHISTs