Transcriptome and histone epigenome of *Plasmodium vivax* salivary-gland sporozoites point to tight regulatory control and potential mechanisms for liver-stage differentiation.

Vivax Sporozoite Consortium[*] (Ivo Muller[1,2,3], Aaron R. Jex[1,3,4], Stefan H. I. Kappe[5], Sebastian A. Mikolajczak[5], Jetsumon Sattabongkot[7], Rapatbhorn Patrapuvich[6], Scott Lindner[8], Erika L. Flannery[5], Cristian Koepfli[1], Brendan Ansell[4], Anita Lerch[1], Samantha J Emery-Corbin[1], Sarah Charnaud[1], Jeffrey Smith[1], Nicolas Merrienne[2], Kristian E. Swearingen[5], Robert L. Moritz[9], Michaela Petter[10,11], Michael Duffy[10], Vorada Chuenchob[5]).

[*]**Group authorship – all authors are equal contributors (order per author contributions section below).**

1. Population Health and Immunity Division, The Walter and Eliza Hall Institute for Medical Research, 1G Royal Parade, Parkville, Victoria, 3052, Australia.
2. Malaria: Parasites & Hosts Unit, Institut Pasteur, 28 Rue de Dr. Roux, 75015, Paris, France.
3. Department of Medical Biology, The University of Melbourne, Victoria, 3010, Australia.
4. Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Corner of Park and Flemington Road, Parkville, Victoria, 3010, Australia.
5. Center for Infectious Disease Research, 307 Westlake Avenue North, Suite 500, Seattle, WA 98109, USA;
6. Department of Global Health, University of Washington, Seattle, WA 98195, USA.
7. Mahidol *Vivax* Research Center, Faculty of Tropical Medicine, Mahidol University, Bangkok 10400, Thailand.
8. Department of Biochemistry and Molecular Biology, Center for Malaria Research, Pennsylvania State University, University Park, PA 16802, USA.
9. Institute for Systems Biology, Seattle, WA, 98109, USA.
10. Department of Medicine Royal Melbourne Hospital, The Peter Doherty Institute, The University of Melbourne, 792 Elizabeth Street, Melbourne, Victoria 3000, Australia.
11. Institute of Microbiology, University Hospital Erlangen, Erlangen 91054, Germany

ABSTRACT

Plasmodium vivax is the key obstacle to malaria elimination in Asia and Latin America, largely attributed to its ability to form resilient 'hypnozoites' (sleeper-cells) in the host liver that cause relapsing infections. The decision to form hypnozoites is made early in the liver infection and may be set in sporozoites prior to invasion. To better understand these early stages of infection, we undertook transcriptomic and histone epigenetic characterization of *P. vivax* sporozoites. Through comparisons to recently published proteomic data for the *P. vivax* sporozoite, our study highlights the loading of the salivary-gland sporozoite with proteins required for cell traversal and invasion and transcripts for infection of and development within hepatocytes. Though highly transcribed, these transcripts are not detectable as proteins, indicating they may be regulated in the sporozoite through translational repression. We also undertook differential transcriptomic studies comparing the sporozoite with newly published mixed blood-stage and mixed versus hypnozoite-enriched liver stage transcriptomes. These comparisons indicate multiple layers of transcriptional, post-transcriptional and post-translational control active in the sporozoite stage and to a lesser extent hypnozoites, but largely absent in mixed liver or blood-stages. Common transcripts up-regulated in sporozoites and hypnozoites compared to mixed stages include several AP2 transcription factors, translational repressors and histone epigenetic regulators, as well as genes linked to dormancy/persistence in bacteria, amoebae and plants. We characterise histone epigenetic modifications in the *P. vivax* sporozoite and explore their role in regulating transcription. This work shows a close correlation between H3K9ac marks and transcriptional activity, with H3K4me3 and H3K9me3 appearing to act as general markers of euchromatin and heterochromatin respectively. We also show little evidence of transcriptional activity in the (sub)telomeres in sporozoites and discuss potential roles of AP2 transcription factors, including ApiAP2-L in regulating this stage. Collectively, these data indicate the sporozoite as a tightly programmed stage primed to infect the human host and identifies key targets to be further explored in liver stage models.

Author summary

Our study is the first to use RNA-seq and ChIP-seq technologies to comprehensively characterize the *P. vivax* sporozoite transcriptome and histone epigenome, and the first to

integrate these technologies with proteomic studies to explore translational repression in the sporozoite of any *Plasmodium* spp. Our study improves on previous work in this field and is supported by recent publication of the transcriptome/histone epigenome of *P. falciparum* and the liver-stage transcriptomes of a related simian parasite (*P. cynomolgi*) that also forms dormant liver stages. Collectively, these data characterize the infectious sporozoite as a highly programmed and tightly regulated, 'torpedo'-like stage primed to infect the human host and suggests that hypnozoite formation is mediated both by differential signals in the sporozoite and by developmental arrest regulators that suppress further development to the schizont stage. This work reveals the primary importance of translational repression and chromatin epigenetics in this process. Our study provides a foundation for exploring the genetic differences between sporozoites derived from parasites with broadly differing hypnozoite formation and relapse phenotypes and highlights genes of major importance in liver stage development that may now be rationally investigated (including as potential drug targets) in relevant models.

INTRODUCTION
Malaria is among the most significant infectious diseases impacting humans globally, with 3.3 billion people at risk of infection, 381 million suspected clinical cases and up to ~660,000 deaths attributed to malaria in 2014 [1]. Two major parasitic species contribute to the vast majority of human malaria, *Plasmodium falciparum* and *P. vivax*. Historically, *P. falciparum* has attracted the majority of global attention, due to its higher contribution to morbidity and mortality. However, *P. vivax* is broadly distributed, more pathogenic than previously thought, and is recognised as the key obstacle to malaria elimination in the Asia-Pacific and Americas [2]. Unlike *P. falciparum*, *P. vivax* can establish long-lasting 'sleeper-cells' (= hypnozoites) in the host liver that emerge weeks, months or years after the primary infection (= relapsing malaria) [3]. Primaquine is the only approved drug that prevents relapse. However, the short half-life, long dosage regimens and incompatibility of primaquine with glucose-6-phosphate-dehydrogenase deficiency (which requires pre-screening of recipients [4]) makes it unsuitable for widespread use. As a consequence, *P. vivax* is overtaking *P. falciparum* as the primary cause of malaria in a number of co-endemic regions [5]. Developing new tools to diagnose, treat and/or prevent hypnozoite infections is considered one of the highest priorities in the malaria elimination research agenda [6].

When *Plasmodium* sporozoites are deposited by an infected mosquito, they likely traverse the skin cells, enter the blood-stream and are trafficked to the host liver, as has been shown in rodents [7]. The sporozoites' journey from skin deposition to hepatocytes takes less than a few minutes [8]. Upon reaching the liver, sporozoites traverse Kupffer and endothelial cells to reach the parenchyma, moving through several hepatocytes before invading a final hepatocyte suitable for development [7, 9]. Within hepatocytes, these parasites replicate, and undergo further development and differentiation to produce merozoites that emerge from the liver and infect red blood cells. However, *P. vivax* sporozoites are able to commit to two distinct developmental fates within the hepatocyte: they either immediately continue development as replicating schizonts and establish a blood infection, or delay replication and persist as hypnozoites. Regulation of this major developmental fate decision is not understood and this represents a key gap in current knowledge of *P. vivax* biology and control.

Sporozoites prepare for mammalian host infection while still residing in the mosquito salivary glands. It has been hypothesized that *P. vivax* sporozoites exist within an inoculum as replicating 'tachysporozoites' and relapsing 'bradysporozoites' [10] and that these subpopulations may have distinct developmental fates as schizonts or hypnozoites, thus contributing to their relapse phenotype [10-12]. This observation is supported by the stability of different hypnozoite phenotypes (ratios of hypnozoite to schizont formation) in *P. vivax* infections of liver-chimeric mouse models [13]. To determine fates in the sporozoite stage, control of protein expression must take place. Studies using rodent malaria parasites have identified genes [14] that are transcribed in sporozoites but translationally repressed (i.e., present as transcript but un- or under-represented as protein), via RNA-binding proteins [15], and ready for immediate translation after the parasites' infection of the mammalian host cell

[13, 16]. It is therefore also possible that translational repression (i.e., the blocking of
translation of present and retained transcripts) and other mechanisms of epigenetic control
may contribute to the *P. vivax* sporozoite fate decision and hypnozoite formation, persistence
and activation. Supporting this hypothesis, histone methyltransferase inhibitors stimulate
increased activation of *P. cynomolgi* hypnozoites to become schizonts in macaque
hepatocytes [17, 18]. Epigenetic control of stage development is further evidenced in
Plasmodium through chromatin structure controlling expression of PfAP2-G, a specific
transcription factor that, in turn, regulates gametocyte (dimorphic sexual stages) development
in blood-stages [19]. It is well documented that *P. vivax* hypnozoite activation patterns
stratify with climate and geography [11] and recent modelling suggests transmission potential
selects for hypnozoite phenotype [20]. Clearly the ability for *P. vivax* to dynamically regulate
hypnozoite formation and relapse phenotypes in response to high or low transmission periods
in different climate conditions would confer a significant evolutionary advantage.
Unfortunately, despite recent advances [21] current approaches for *in vitro P. vivax*
culture do not support routine maintenance in the laboratory and tools to directly perturb gene
function are not established. This renders studies on *P. vivax*, particularly its sporozoites and
liver stages, exceedingly difficult. Although *in-vitro* liver stage assays and humanised mouse
models are being developed [13], 'omics analysis of *P. vivax* liver stage dormancy has until
recently [22] been impossible and even now is in its early stages. Recent characterization [23]
of liver-stage (hypnozoites and schizonts) of *P. cynomolgi* (a related and relapsing parasite in
macaques) provides valuable insight, but further study of *P. vivax* directly is needed. The
systems analysis of *P. vivax* sporozoites that reside in the mosquito salivary glands and are
poised for transmission and liver infection offer a key opportunity to gain insight into *P. vivax*
infection. To date, such characterization of *Plasmodium vivax* sporozoites is limited to
microarray data [24], and epigenetic regulation in sporozoites has only been explored in *P.*
falciparum [25, 26]. Here, we present a detailed characterization of the *P. vivax* sporozoite
transcriptome and histone epigenome and use these data to better understand this key
infective stage and the role of sporozoite programming in invasion and infection of the human
host, and development within the host liver.

Fig. 1 *Transcriptional activity of the* P. vivax *sporozoite and evidence for translational*
repression. ***a*** *Relative transcript abundance of key marker genes for sporozoites inferred by*
RNA-seq and qPCR (here) relative to previously published microarray data [24]; ***b*** *Relative*
proportion of genes detectable as transcripts and proteins or transcripts only in RNA-seq and
previously published proteomic data. Dashed line shows cut-off used in the current study for
putatively repressed transcripts. Immunofluorescent staining of select proteins either known
(UIS4) or predicted here (LISP1, EXP1 and ACP) to be translationally repressed in
sporozoites in ***c*** *sporozoite stages. CSP, mTIP as known positive controls and TRAP and BIP*
as exerpimental positive controls and ***d*** *liver stages (schizonts) at 7 days post-infection in*
HuHep mice. Liver expression of EXP1 and ACP has been demonstrated by IFA in
Mikolajczak et al [13], using the same antibodies as used here.

RESULTS AND DISCUSSION
Mosquito infections were generated by membrane feeding of blood samples taken from *P.*
vivax infected patients in western Thailand (n = 9). Approximately 3-15 million *P. vivax*
sporozoites were harvested per isolate from *Anopheles dirus* salivary glands. Using RNA-seq,
we detected transcription for 5,714 *P. vivax* genes (based on the *P. vivax* P01 gene models:
[27]) and obtained a high degree of coverage (4,991 with a mean transcript per million (TPM)
count ≥ 1.0) at a mean estimated abundance of 175.1 TPM (Figure S1 and Table S1 and S2).
Among the most highly transcribed genes in the infectious sporozoite stage are *csp*
(circumsporozoite protein), five *etramps* (early transcribed membrane proteins), including
uis3 (up-regulated in infective sporozoites), *uis4* and *lsap-1* (liver stage associated protein 1),
a variety of genes involved in cell transversal and initiation of invasion, including *celtos* (cell
traversal protein for ookinetes and sporozoites), *gest* (gamete egress and sporozoite traversal
protein), *spect1* (sporozoite protein essential for cell traversal) and *siap-1* (sporozoite invasion

associated protein), and genes associated with translational repression (*alba1*, *alba4* and
Puf2). Collectively, these genes account for >1/3rd of all transcripts in the sporozoite. While
we found only moderate agreement (R^2 = 0.35; Figure S2) between our RNA-seq data and
previous microarray data for *P. vivax* sporozoites and blood-stages [24], improved transcript
detection and quantitation is expected with the increased technical resolution of RNA-seq
over microarray. Supporting this, we find higher correlation between RNA-seq data from *P.*
vivax and *P. falciparum* (single replicate sequenced herein for comparative purposes)
sporozoite datasets (R^2 = 0.42), compared to either species relative to published microarray
data (Figure S2 and Table S3). Although microarray supports the high transcription in
sporozoites of genes such as *uis4*, *csp*, *celtos* and several other *etramps*, 27% and 16% of the
most abundant 1% of transcribed genes in our sporozoite RNA-seq data are absent from the
top decile or quartile respectively in the existing *P. vivax* sporozoite microarray data [24].
Among these are genes involved in early invasion/hepatocyte development, such as *lsap-1*,
celtos, *gest* and *siap-1*, or translational repression (e.g., *alba-1* and *alba-4*); orthologs of these
genes are also in the top percentile of transcripts in RNA-seq (see [25] and Table S4) and [28]
and Table S5) and previous microarray data [29, 30] for human-infecting *P. falciparum* and
murine-infecting *P. yoelii* sporozoites, suggesting many are indeed more abundant than
previously characterized. A subset of representative transcripts, including a *Pv_AP2*
transcription factor (PVP01_0733100), *d13*, *gest*, *g10* (PVP01_1011100), 40S ribosomal
protein S27 (PVP01_1409300), *puf-2*, *zipco* and 14-3-3, were tested by qPCR for their
transcript abundance relative to *celtos* and *sera* (**Fig. 1A**). This representative set differed
markedly in their relative abundance between our RNAseq and previous microarray data [24].
To control for batch effects introduced by collection of the sporozoites used here for RNAseq,
this testing was conducted in an additional six sample replicates representing four additional
clinical isolates *P. vivax* isolates (PvSPZ-Thai13-16; with PvSPZ-Thai16 tested in technical
triplicate). The qPCR results overwhelmingly agreed with the RNAseq data for these
transcripts.

***Transcription in* P. *vivax relative to other plasmodia* sporozoites.** To gain insight into
species-specific aspects of the *P.vivax* transcriptome, we qualitatively compared these data
with available data for *P. falciparum* [26] and *P. yoelii* sporozoites (single replicate only) for
4,220 and 4,067 single-copy orthologs (SCO) (transcribed at ≥ 1 TPM in *P. vivax* infectious
sporozoites) shared with *P. falciparum* (Table S4) and with both *P. falciparum* and *P. yoelli*
(Table S5) respectively. Genes highly transcribed in salivary-gland sporozoites of all three
species include *celtos*, *gest*, *trap*, *siap1*, *spect1* and *puf2*. There are 696 *P. vivax* genes shared
as orthologs between *P. vivax* P01 and *P. vivax* Sal1 lacking a defined SCO in *P. falciparum*
or *P. yoelli* transcribed at a mean of ≥ 1 TPM in *P. vivax* salivary-gland sporozoites (Table
S6). Prominent among these are *vir* (n=25) and *Pv-fam* (41 fam-e, 16 fam-b, 14 fam-a, 8 fam-
d and 3 fam-h) genes, as well as, hypothetical proteins or proteins of unknown function
(n=212) and, interestingly, a number of 'merozoite surface protein' 3 and 7 homologs (n=5 of
each). Both *msp3* and *msp7* have undergone significant expansion in *P. vivax* relative to *P.*
falciparum and *P. yoelii* [31] and may have repurposed functions in sporozoites. In addition,
there are 69 *P. vivax* P01 genes lacking a defined ortholog in *P. vivax* Sal1, *P. falciparum* or
P. yoelli transcribed at ≥ 1 TPM in infectious *P. vivax* sporozoites; most of which are
Plasmodium interspersed repeat (PIR) genes [31] found in telomeric regions of the P01
assembly and likely absent from the Sal1 assembly but present in the Sal1 genome, indicating
the improved coverage of telomeric regions in P01 relative to Sal1.

P. *vivax* sporozoites transcriptome compared with proteome. We compared relative
protein abundance presented in a recently published *P. vivax* sporozoite proteome [32] to
relative transcript abundance from the current study (**Fig. 1B**, Table S7). The proteome study
incorporated data from the same PvSPZ-Thai1 and PvSPZ-Thai5 isolates tested by RNAseq
here. We identified 2,402 *P. vivax* genes transcribed in the sporozoite (CPM > 1) for which
no protein expression was detected. Although many of these are lowly transcribed and likely

below the detection sensitivity of LC-MS proteomics, others are among the most highly transcribed genes in the sporozoite, indicating these may be under translational repression.

Translational repression, the mechanism through which transcripts are held in stasis by RNA binding proteins, has been demonstrated to have important functional roles in the transition of *Plasmodium* spp. from the vertebrate to invertebrate host and back again. More than 700 genes have been identified as being translationally repressed in *Plasmodium berghei* ('rodent malaria') gametocytes based on DOZI pulldowns [33]. Translational repression mechanisms mediated through Puf-2 have been explored in sporozoites of several *Plasmodium* species and regulates some of the most abundant transcripts in the sporozoite, such as *uis-3* and *uis-4*. UIS3 and UIS4 are the best characterized proteins under translational repression by Puf-2/SAP1 in sporozoites [34] and are essential for liver stage development [14].

In considering genes that may be translationally repressed in the *P. vivax* sporozoite, we confine our observations to those transcripts representing the top decile of transcript abundance to ensure their lack of detection as proteins was not due to limitations in the detection sensitivity of the proteomic dataset. Approximately $1/3^{rd}$ of transcripts in the top decile of transcriptional abundance (n = 170 of 558) in *P. vivax* sporozoites were not detectable as peptides in multiple replicates (**Fig. 1B** and Table S7). Of these 170 putatively repressed transcripts, 156 and 154 have orthologs in *P. falciparum* and *P. yoelii* respectively, with 89 and 118 of these also not detected as proteins in *P. falciparum* and *P. yoelii* salivary-gland sporozoites [35] despite being transcribed in these stages (see [25, 28]; Tables S3-S5), and 133 (78.2%) having no detectable expression (>1 unique peptide count) in LC-MS data deposited for any species in PlasmoDB (Table S8). In contrast, 106 of the 170 putatively repressed transcripts with orthologs in other *Plasmodium* species (62.3%; Table S8) for which proteomic data is available in PlasmoDB, are detectable (>1 unique peptide count) by LC-MS methods in at least one other life-cycle stage. This indicates against a technical issue (e.g., inability to be trypsin-digested) preventing their detection in the *P. vivax* sporozoite proteome [32]. In addition to *uis3* and *uis4*, genes involved in liver stage development and detectable as transcripts but not proteins in the *P. vivax* sporozoites include *lsap1* (liver stage associated protein 1), *zipco* (ZIP domain-containing protein), several other *etramps* (PVP01_1271000, PVP01_0422600, PVP01_0504800 and PVP01_0734800), *pv1* (parasitophorous vacuole protein 1) and *lisp1* and *lisp2* (PVP01_1330800 and PVP01_0304700). Also notable among genes detectable as transcripts but not proteins in sporozoites is a putative 'Yippee' homolog (PVP01_0724100). Yippee is a DNA-binding protein that, in humans (YPEL3), suppresses cell growth [36] and is regulated through histone acetylation [37], making it noteworthy in the context of *P. vivax* hypnozoite developmental arrest.

Although verifying each putatively repressed transcript will require further empirical data, our system level approach is supported by immunofluorescent imaging (**Fig. 1C**) of UIS4, LISP1, EXP1 and ACP (PVP01_0416300). These represent one known and three putative translationally repressed genes in *P. vivax* sporozoites, and are compared to TRAP and BiP (which are both transcribed and expressed as protein in the *P. vivax* sporozoite; Table S7). The *lisp1* gene is an interesting find. In *P. berghei*, *lisp1* is essential for rupture of the PVM during liver stage development allowing release of the merozoite into the host blood stream. *Pv-lisp1* is ~350-fold and ~1,350-fold more highly transcribed in *P. vivax* sporozoites compared to sporozoites of either *P. falciparum* or *P. yoelii* (see Table S5). IFAs using LISP1 specific mAbs (**Fig. 1C**) show that this protein is undetectable in sporozoites but clearly expressed at 7 days post-infection in liver schizonts.

Up-regulated transcripts in *P. vivax* sporozoites relative to other life-cycle stages. Recently completed studies of the transcriptome of *P. vivax* in the liver [22] and in blood-cells [38] support comparative transcriptomic study of sporozoites, their biology and transcriptional regulation over the *P. vivax* life-cycle. These data present an analytical challenge in that each (sporozoites, liver-stages and blood stages) is produced in a separate study and may be influenced by technical batch effects that cannot be differentiated from biologically meaningful changes. To address this, we first examined *P. vivax* transcripts in a

previous microarray study of multiple *P. vivax* life-cycle stages [24], including sporozoites and several blood-stages, to identify housekeeping genes that are transcriptionally stable across the life-cycle. We identified ~160 genes with low transcriptional variability across the *P. vivax* life-cycle, that covered the breadth of transcript abundance levels in Westenberger et al [24]. These genes include functions typically associated with housekeeping genes, such as ribosomal proteins, histones, translation initiation complex proteins and various chaperones (see Figure S3 and S4 and Table S8). We assessed transcription of these 160 'housekeeping' genes among the current and recently published RNA-seq data for *P. vivax* and all were of similarly low variability (Figure S6). This suggests that any batch effect between the studies is sufficiently lower than the biological differences between each life-cycle stage, allowing informative comparisons.

P. vivax *sporozoite relative to blood-stage transcriptome.* To identify transcripts up-regulated in sporozoites, we first compared the *P. vivax* sporozoite transcriptome to RNA-seq data for *P. vivax* blood-stages [38] (**Fig. 2** and Figures S5-S9). We identified 1,672 up (Table S9); Interactive Glimma Plot - Supplementary Data 1) and 1,958 down-regulated (Table S9); Interactive Glimma Plot - Supplementary Data 1) transcripts (FDR ≤ 0.05; minimum 2-fold change in Counts per Million (CPM)) and explored patterns among these differentially transcribed genes by protein family (**Fig. 2C** and Table S10) and Gene Ontology (GO) classifications (Table S11). RNA recognition motifs (RRM-1 and RRM-6) and helicase domains (Helicase-C and DEAD box helicases) are over-represented (p-value <0.05) among transcripts up-regulated in sporozoites, consistent with translational repression through ribonucleoprotein (RNP) granules [39]. Transcripts encoding nucleic acid binding domains, such as bromodomains (PF00439; which can also bind lysine-acetylated proteins), zinc fingers (PF13923) and EF hand domains (PF13499) are also enriched in sporozoites. Included among these proteins are a putative ApiAP2 transcription factor (PVP01_1211900) and a homologue of the *Drosophila* zinc-binding protein 'Yippee' (PVP01_0724100). Thrombospondin-1 like repeats (TSR: PF00090) and von Willebrand factor type A domains (PF00092) are enriched in sporozoites as well. In *P. falciparum* sporozoites, genes enriched in TSR domains are important in invasion of the mosquito salivary gland (e.g., *trap*) and secretory vesicles released by sporozoites upon entering the vertebrate host (e.g., *csp*) [40]. By comparison, genes up-regulated in blood-stages are enriched for *vir* gene domains (PF09687 and PF05796), Tryptophan-Threonine-rich *Plasmodium* antigens (PF12319; which are associated with merozoites [41]), markers of cell-division (PF02493; [42]) protein production/degradation (PF00112, PF10584, PF00152, PF09688 and PF00227) and ATP metabolism (PF08238 and PF12774). 47 of the 343 transcripts unique to *P. vivax* sporozoites relative to *P. falciparum* or *P. yoelii* are up-regulated in sporozoites compared to *P. vivax* blood stages. Nine of these are in the top decile of transcription, and include a Pv-fam-e (PVP01_0525200), a Pf-fam-b homolog (PVP01_0602000) and 7 proteins of unknown function. A further nine have an ortholog in *P. cynomolgi* (which also forms hypnozoites) but not the closely related *P. knowlesi* (which does not form hypnozoites) and include '*msp7*'-like (PVP01_1219600, PVP01_1220300 and PVP01_1219900), '*msp3*'-like (PVP01_1031300), Pv-fam-e genes (PVP01_0302100, PVP01_0524500 and PVP01_0523400), a serine-threonine protein kinase (PVP01_0207300) and a RecQ1 helicase homolog (PVP01_0717000). Notably, the *P. cynomolgi* ortholog of PVP01_0207300, PCYB_021650, is transcriptionally up-regulated in hypnozoites relative to replicating schizonts [23], indicating a target of significant interest when considering hypnozoite formation and/or biology and suggesting that our list here may contain other genes important in hypnozoite biology.

Fig. 2 *Differential transcription between Plasmodium vivax salivary-gland sporozoites and blood-stages. **A** BCV plot showing separation between blood-stage (black) and salivary-gland sporozoite (red) biological replicates. **B** Volcano plot of distribution of fold-changes (FC) in transcription between blood-stages and salivary-gland sporozoites relative to statistical significance threshold (False Discovery Rate (FDR) ≤ 0.05). Positive FC*

P. vivax *sporozoites are enriched in translational repressors.* In *Plasmodium*, translational repression regulates key life-cycle transitions coinciding with switching between the mosquito and the mammalian host (either as sporozoites or gametocytes) [39]. For example, although *uis4* is the most abundant transcript in the infectious sporozoite ([24, 30]; Table S2), UIS4 is translationally repressed in this stage [15] and only expressed after hepatocyte invasion [43]. In sporozoites, it is thought that PUF2 binds to mRNA transcripts and prevents their translation [28], and SAP1 stabilises the repressed transcripts and prevents their degradation [43]. Consistent with this, *Puf2* and *SAP1* are among the more abundant *P. vivax* transcripts up-regulated in the sporozoite relative to blood-stages. Indeed, *Puf2* is among the top percentile of transcripts in infectious sporozoites and expressed at high levels in the proteome [32]. However, our data implicate other genes that may act in translational repression in *P. vivax* sporozoites, many of which are already known to be involved in translational repression in other *Plasmodium* stages and other protists [39]. Among these are *alba-2* and *alba-4*, both of which are among the top 2% of genes transcribed in sporozoites and ~14 to 20-fold more highly transcribed in sporozoites relative to blood-stages. In addition, *P. vivax* sporozoites are enriched for genes encoding RRM-6 RNA helicase domains. Intriguing among these are HoMu (homolog of Musashi) and ptbp (polypyrimidine tract binding protein). Musashi is a master regulator of eukaryotic stem cell differentiation through translational repression [44] and HoMu localizes with DOZI and CITH in *Plasmodium* gametocytes [45]. PTBP is linked to mRNA stability, splice regulation and translational initiation [46] and may perform a complementary role to SAP1.

*Fig. 3 Differential transcription between P. vivax sporozoites (SPZ), mixed (mLS) and hypnozoite (HPZ) enriched liver stages (liver-stage data from Gural et al [22]). **A** Heatmap comparisons showing summed transcription of enriched Pfam domains in HPZ vs mLS (left), SPZ vs HPZ (top middle) and SPZ vs mLS (top right) comparisons. All Pfam domains statistically significantly enriched at p-value 0.05). All transcript data for stage up-regulated genes at FDR 0.01). **B** Violin box-plot showing relative fold-change differences between SPZ and HPZ compared with SPZ and mLS for genes down-regulated in mLS compared to SPZ, but not down-regulated in HPZ compared to SPZ. **C** Ternary heatmap summarizing relative transcript abundance in each of SPZ, mLS and HPZ stages.*

P. vivax *sporozoite relative to Plasmodium spp. liver stage transcriptomes.* New advances in *P. vivax* in vitro liver stage culture has allowed recent publication of mixed liver-stage (mLS) and hypnozoite-enriched (HPZs) transcriptomes [22]. This is an early study and, due no doubt to the difficulty in generating the material, is limited to biological duplicates. Noting this, although we undertake differential transcriptomic studies of this dataset here, we recognize that additional biological replication is needed and have used a higher burden of significance (FDR ≤ 0.01 and ≥ 2-fold change) than used with blood-stages. Nevertheless, these comparisons identified 1,015 and 856 sporozoite up-regulated transcripts relative to mLS and HPZs respectively and 1,007 and 1,079 transcripts up-regulated in mLS and HPZs relative to sporozoites respectively (**Fig. 3**; Figures S10-S12, Table S12 and S13 and Interactive Glimma Plot - Supplementary Data 1).

Compared to mLS transcriptomes, sporozoites are enriched for many of the transcripts similarly up-regulated in comparison to blood-stages (e.g., *uis4*, *celtos*, *puf2*, *siap1* and *plp-1*). More broadly, SPZ up-regulated transcripts over-represent (p-value ≤ 0.05) Pfam domains (**Fig. 3A**) associated with transcriptional regulation (PF00176, PF01096, PF01661 and PF08711), translational repression/regulation (PF00076, PF00279, PF01008, PF01873, PF01917, PF02847, PF02854, PF13893 and PF14259), DNA/RNA binding (PF0097,

PF13445, PF13639, PF14570 and PF15247) and chromatin regulation (PF00271, PF00850 and PF13489). In contrast, the mLS transcriptome is enriched in genes involved in replication and merozoite formation [n = 14; including PVP01_0728900 (*msp1*), PVP01_0010670 (*msp3*) and PVP01_1446800 (*msp9*)], rhoptry function [n = 9; including PVP01_1469200 (*rnp3*), PVP01_1255000 (*rnp2*) and PVP01_1338500 (*rap1*)] and reticulocyte binding [n=10 including PVP01_0534300 (rbp2c), PVP01_1402400 (rbp2a), PVP01_0701100 (rbp1b) and PVP01_0800700 (rbp2b)]. These data are further enriched for Pfam domains associated with cell division (PF00493), merozoite formation (PF07133 and PF12984), proteasome function (PF00227, PF00400, PF00656, PF01344, PF01398 and PF03981), protein export / vesicle function (PF00350 and PF00996), membrane proteins (PF01105, PF03011, PF05424 and PF12139) and metabolism (PF00085, PF00118, PF00268, PF01066, PF01214 and PF01214). Collectively, in addition to markers consistent with sporozoite or merozoite formation, these data point towards the sporozoite stage as being highly regulated and controlled at transcriptional, translational and chromatin levels, with the mLS representing a release of this control allowing replication, protein turn-over, reconfiguration of the proteins on the plasma membrane and metabolic activity.

Comparison of sporozoites with HPZs does not indicate a similar release of control, or at least that any release is more specific than for mLS. The sporozoite is enriched, relative to HPZs, in genes such as PVP_1258000 (*gest*), PVP01_0418000 (*sera*), PVP01_1435400 (*celtos*), PVP01_0835600 (*csp*) and PVP01_0602100 (*uis4*). At a broad level, sporozoite enriched Pfam domains include a smaller number associated with translational repression/regulation (PF00076) or DNA/RNA binding (PF01428 and PF12756). Interestingly, sporozoites are enriched in Pfam domains specifically associated with heterochromatin (H3K9me^3) reading/interaction (PF02463, PF00628, PF13831 and PF13865). Ours (see below) and previous epigenetic studies of *Plasmodium* sporozoites [26] find dense heterochromatin in the telomeric to subtelomeric regions of the chromosome, which opens up and becomes transcriptionally active in blood-stages [47]. Others have noted an up-regulation of methyl/acetyltransferases in *P. cynomolgi* HPZs [23] and/or shown methyltransferase inhibitors stimulate hypnozoite activation *in vitro* [17]. The potential that histone epigenetics of sporozoites has a role in or changes with liver-stage development and the formation of liver schizonts or HPZs is intriguing but requires detailed study of the chromatin of liver-stage parasites, which is not presently available for *P. vivax*. In contrast, HPZs were enriched, relative to sporozoites, for genes including histone proteins (PVP01_1138700, PVP01_1131700 and PVP01_0905900) and classic markers of metabolism (PVP01_MITO3300 and PVP01_MITO3400) and *lisp2*. Pfam data indicated, largely similar domain enrichment trends as were seen of the mLS relative to sporozoites, including a number of proteosomal (PF00227, PF00112, PF03981), vesicular transport (PF00996) and metabolic (PF00118, PF00268, PF01066, PF01214 a) associated functions. This supports HPZs being an arrested, rather than classically 'dormant', stage with active metabolism and protein turn-over. HPZs are also enriched for Pfams associated with mRNA/tRNA regulation and turnover (PF04857, PF01612, PF00009 and PF01138) and glycine metabolism (PF01571 and PF00464) and acetyl-CoA production (PF02779 and PF00676).

Finally, although not the focus of this study, we looked at differential transcription between mLS and HPZ stages using the Gural et al [22] data, but using the same approaches as employed here. In particular, we were interested in what insight these comparisons might provide in terms of sporozoite differentiation or development into liver schizonts or HPZs. Among mLS up-regulated transcripts are genes associated with rhoptry function (n = 11; including PVP01_0107500, PVP01_1469200 and PVP01_1469200), cytoadherence to red-cells (PVP01_1401400 and PVP01_0734500), merozoite formation (PVP01_0728900 and PVP01_0612400) and exported proteins (n = 6; including PVP01_0504000, PVP01_0119200 and PVP01_0801600). Consistent with *P. cynomolgi* [23], HPZ up-regulated transcripts include several key sporozoite transcripts, specifically *uis4* (PVP01_0602100), *puf1* (PVP01_1015000) and *speld* (PVP01_0938800). At the Pfam domain level, mLS is enriched for metabolic (PF00317) and proteosomal (PF00112) domains also enriched in mLS or HPZs relative to sporozoites above, as well as domains associated with merozoite formation

(PF12948, PF07462), rhoptry function (PF0712), DNA/RNA binding (PF12756, PF10601
and PF02151) and cell division, development and DNA replication (PF06705, PF00533,
PF00488, PF02460, PF07034, PF02181). In contrast, HPZs are enriched in Pfam domains
that overlap notably with key sporozoite markers, including *etramps* (PF09716) and *puf*
proteins (PF00806), as well as domains associated with calcium metabolism (PF08683) and
nucleotide metabolism (PF06437). These data largely indicate that the hypnozoite bears
similarity both to the sporozoite and liver schizonts consistent with a stalled stage on the path
to schizont development regulated by checkpoint signals that halt/restart normal schizont
development, which has been proposed previously for this species [24].
With this is mind, we looked at transcripts that are differentially transcribed in mLS,
but not HPZs, relative to SPZs. There are 107 transcripts down-regulated in mLS relative to
SPZs that are transcribed at roughly similar levels in both SPZs and HPZs (**Fig 3B** and Table
S14). A common theme among many of these genes are their role in transcriptional, post-
transcriptional, translational or post-translational regulation. Among transcriptional regulators
are transcription factors including AP2-SP2 (PVP01_0303400) and three non-AP2-like
transcription factors (PVP01_0306600, PVP01_0204300 and PVP01_1415800). Post-
transcriptional controllers include several DNA/RNA-binding proteins (PVP01_1011000.
PVP01_0932900, PVP01_0715300, PVP_1242600 and PVP01_0605200), RNA helicases
(PVP01_1403600 and PVP01_1329800) and mRNA processing (PVP01_1443100 and
PVP01_1458200) genes. Translational control includes several key regulators of translation
initiation (PVP01_1467700), tRNA processing (PVP01_0318700 and PVP01_1017700) or
ribosomal function/biogenesis (PVP01_1443700, PVP01_0421400, PVP01_1117200 and
PVP01_0215100).   Post-translational   control   includes   two   methyltransferases
(PVP01_1428800 and PVP01_1465200), including CARM1, which methylates of H3R17
and, in mice, prevents differentiation in embryonic stem cells [48], and a putative histone
methylation reading enzyme, EEML2 (PVP01_1014100). The remaining genes in this group
have three noteworthy and largely overlapping themes: (1) an association with calcium
binding, metabolism or signalling, (2) a role in organellar metabolism and (3) homologs in
other organisms, including a variety of prokaryotes and eukaryotes, with key roles in
germination, dormancy and persistent non-replicating stages. The latter most function is
clearly intriguing in the context of HPZ formation and activation. These genes include a
homolog of dihydrolipoamide acyltransferase (aka 'sucB'), which is essential for growth in
Mycobacterium tuberculosis [49] and a key regulator in persistent *Escherichia coli* stages
[50]. Another key example is gamete fusion factor HAP2, which, despite the name, has been
shown to regulate dormancy in eukaryotes ranging from plants [51, 52] to amoebae [53].
In addition to data for *P. vivax*, two transcriptomic studies are now available for *P.
cynomolgi* [26, 54] that compare schizont stage parasites with small-form "hypnozoites". In
comparing *P. cynomolgi* liver-stage RNA-seq and *P. vivax* liver-stage microarray data [24],
Cubi et al [23] noted a moderate to good level of agreement (R^2 = 0.50) as evidence of *P.
cynomolgi* being predictive and representative of *P. vivax*. However, it should be noted also,
that Voorberg van der Wel et al [54] explored congruence between their's and the Cubi et al
[23] studies and found generally good agreement among schizonts and overall relatively poor
agreement among hypnozoites from each study. This highlights the complexity of these
datasets and indicates caution in comparing the current data to *P. vivax*. ApiAP2 transcription
factors feature prominently in each liver-stage transcriptomic studies for *P. cynomolgi* [23,
54] and *P. vivax* [22]. Cubi et al [23] noted an ApiAP2 (dubbed "AP2-Q"; PCYB_102390) as
transcriptionally up-regulated in *P. cynomolgi* hypnozoites and proposed this as a potential
hypnozoite marker. We note that The *P. vivax* ortholog of *Pc*-AP2-Q (PVP01_1016100) is
among the genes detectable as a transcript but not protein in *P. vivax* sporozoites. This may
point to a translationally repressed signal in sporozoites to regulate hypnozoite formation.
However, as *Pv*-AP2-Q is transcribed at an abundance (~50 TPM) at or below which ~50% of
P. vivax genes are detectable as transcripts but not as proteins (**Fig 1B**), this could as likely
result from LC-MS detection sensitivity. Further, although AP2-Q was reported as absent
from non-hypnozoite forming *Plasmodium* species [23], it is indeed found broadly across the
genus, including among several non-hypnozoite producing *Plasmodium* species, such as *P.

knowlesi, *P. gallinaceum* and *P. inui* [54]. Up-regulation of AP2-Q transcripts is not observed
for hypnozoites in subsequent transcriptomic studies of *P. cynomolgi* [54] or *P. vivax* [22],
nor do we see such an up-regulation here. Voorberg van der Wel et al [54] note transcription
of a range of AP2s in *P. cynomolgi* liver stages, but do not find any to be up-regulated in
hypnozoites. AP2s also feature among transcribed genes in *P. vivax* liver stages, with one,
PVP01_0916300, significantly up-regulated in hypnozoites. We note that PVP01_0916300 is
up-regulated in *P. vivax* sporozoites relative to blood-stages and found in the top quartile of
transcripts by abundance (TPM = 104).
506
Chromatin epigenetics in *P. vivax* sporozoites. As noted above, transcriptomic data for
sporozoites, and their comparison with liver and blood-stages implicate histone epigenetics as
having an important role in sporozoite biology and differentiation into liver stages. This
concept has been alluded to also in recent liver-stage studies of *P. cynomolgi* [17, 23], which
propose methyltransferases as having a potential role in hypnozoite formation. No epigenetic
data are currently available for any *P. vivax* life-cycle stage. Studies of *P. falciparum* blood-
stages have identified the importance of histone modifications as a primary epigenetic
regulator [55, 56] and characterized key markers of heterochromatin (H3K9me^3) and
euchromatin/transcriptional activation (H3K4me^3 and H3K9ac). Recently, these marks have
been explored with the maturation of *P. falciparum* sporozoites in the mosquito [25]. Here,
we characterize major histone marks in *P. vivax* sporozoites and assess their relationship to
transcript abundance.
519
Histone modifications in* P. vivax *sporozoites. Using ChIP-seq, we identified 1,506, 1,999
and 5,262 ChIP-seq peaks stably represented in multiple *P. vivax* sporozoite replicates and
associated with H3K9me^3, H3K9ac and H3K4me^3 histone marks respectively (**Figures 4** and
S13). Peak width, spacing and stability differed with histone mark type (Figures S14 and
S15). H3K4me^3 peaks were significantly broader (mean width: 1,985 bp) than H3K9 peaks,
and covered the greatest breadth of the genome; 36.0% of all bases were stably associated
with H3K4me^3 marks. This mark was also most stable among replicates, with just ~16% of
bases associated with an H3K4me^3 not supported by more than one biological replicate. By
comparison H3K9me^3 marks were narrowest (mean width: 796 bp) and least stable, with 46%
of bases associated with this mark supported by just one replicate. Consistent with
observations in *P. falciparum* H3K9me^3 'heterochromatin' marks primarily clustered in
telomeric and subtelometric regions (**Figure 4**). In contrast, the 'euchromatin' /
transcriptionally open histone marks, H3K4me^3 and H3K9ac, clustered around genic regions
and did not overlap with regions under H3K9me^3 suppression. Both H3K9me^3 and H3K4me^3
marks were reasonably uniformly distributed (mean peak spacing ~500bp for each) within
their respective regions of the genome. In contrast, H3K9ac peaks were spaced further apart
(mean: ~2kb), but also with a greater variability in spacing (likely reflecting their association
with promoter regions [57]). The instability of H3K9me^3 may reflect its use in *Plasmodium*
for regulating variegated expression of contingency genes from multigene families whose
members have overlapping and redundant functions [47] and confer phenotypic plasticity
[58].
541
Fig. 4 Histone epigenetics relative to transcriptional behaviour in salivary-gland sporozoites.
a Representative H3K9me^3, H3K4me^3 and H3K9ac ChIP-seq data (grey) from a
representative chromosome (P. vivax P01 Chr5) relative to mRNA transcription in salivary-
gland sporozoites (black) and blood-stages (black). Small numbers to top left of each row
show data range. b Salivary-gland sporozoite transcription relative to nearest stable histone
epigenetic marks. Numbers at the top of the figure represent total genes included in each
category. Numbers within in box plot represent mean transcription in transcripts per million
(TPM). c Sequence motifs enriched within 1kb upstream of the Transcription Start Site of
highly transcribed (top 10%) relative to lowly transcribed genes associated with H3K9ac
marks in salivary-gland sporozoites. d Relative transcription of (sub)telomeric genes in P.
vivax salivary-gland sporozoites and blood-stages categorized by gene sets up-regulated in

Genes under histone regulation. We explored an association between these histone marks and the transcriptional behaviour of protein coding genes (**Fig. 4** and Tables S15-S20). 485 coding genes stably intersected with an H3K9me[3] mark; all are located near the ends of the chromosomal scaffolds (i.e., are (sub)telomeric). On average, these genes are transcribed at ~30 fold lower levels (mean 0.7 TPM) than genes not stably intersected by H3K9me[3] marks. These data clearly support the function of this mark in transcriptional silencing. This is largely consistent with observations in *P. falciparum* sporozoites ([25]; Figure S16), however, we observe no genes within heterochromatin dense region that lacked a stably marked by H3K9me[3] signal or were transcribed at notable levels (i.e., above ~5 TPM). Whether this relates to differences in epigenetic control between the species is not clear. We note that (sub)telomeric genes are overall transcriptionally silent in *P. vivax* sporozoites relative to blood-stages (**Fig. 4a and 4b** and Tables S21 and S22). Consistent with observations in *P. falciparum* [55], the bulk of these genes include complex protein families, such as *vir* and *Pv-fam* genes, which function primarily in blood-stages. Also notable among the genes are several reticulocyte-binding proteins, including RBP2, 2a, 2b and 2c. This transcriptional silence in telomeric and subtelomeric regions was recently observed in *P. falciparum* sporozoites [26].

Outside of the telomeres and subtelomeres, H3K4me[3] marks are stably associated with the Transcription Start Site (TSS) and/or 5' UTRs of 3,677 genes. H3K9ac marks are also identified within 1kb of the TSS of 1,284 coding genes, with 179 of these stably marked also by H3K4me[3] (Figure S17). The average transcription of these genes is 50, 112 and 112 TPMs respectively (72, 160 and 160-fold higher than H3K9me[3] marked genes). Gene-by-gene observations show that H3K9ac and H3K4me[3] marks cluster densely in the 1000kb up and down-stream of the start and stop codon respectively of transcribed genes, but are much less dense within coding regions of these genes (Figure S17). This pattern directly correlates with transcription and contrasts H3K9me[3] marks, which are distributed across the length of the gene at even density and are correlated with a lack of transcription. These data support the role of these marks in transcriptional activation, the lower abundance of H3K4me[3] marker, compared with H3K9ac or H3K9ac and H3K4me[3] marked genes suggest these marks work synergistically and that H3K9ac is possibly the better single mark indicator of transcriptional activity in *P. vivax*. This is consistent with recent observations in *P. falciparum* sporozoites [25].

Interestingly, H3K9ac-marked genes ranged in transcriptional activity from the most abundantly transcribed genes to many in the lower 50% and even lowest decile of transcription. This suggests more contributes to transcriptional activation in *P. vivax* sporozoites than, simply, gene accessibility through chromatin regulation. Specific activation by a transcription factor (e.g., ApiAP2s [59]) is the obvious candidate. To explore this, we compared upstream regions (within 1kb of the TSS or up to the 3' end of the next gene upstream, whichever was less) of highly (top 10%) and lowly (bottom 10%) transcribed H3K9ac marked genes for over-represented sequence motifs that might coincide with known ApiAP2 transcription factor binding sites [60]. We identified these based on the location of the nearest stable H3K9ac peak relative to the transcription start site for each gene (Figure S12). In most instances, these peaks were within 100bp of the TSS and, consistent with data from *P. falciparum* [57], *P. vivax* promoters appear to be no more than a few hundred to a maximum of 1000 bp upstream of the TSS. Exploring these regions, we identified two over-represented motifs: TGTACMA (e-value $2.7e^{-2}$) and ATATTTH (e-value $3.3e^{-3}$) (Fig. 2D). TGTAC is consistent with the known binding site for *Pf*-AP2-G, which regulates sexual differentiation in gametocytes [61], but its *P. vivax* ortholog (PVP01_1418100) is neither highly transcribed nor expressed in sporozoites. ATATTTH is similar to the binding motif for *Pf*-AP2-L (AATTTCC), a transcription factor that is important for liver stage development in *P. berghei* [62]. In contrast to AP2-G, *Pv*-AP2-L (PVX_081180) is in the top 10% of transcription and expression in *P. vivax* sporozoites and up-regulated relative to blood-stages.

In *P. vivax* sporozoites, the ATATTTH motif is associated with a number of highly
transcribed genes, including *lisp1* and *uis2-4*, known to be regulated by AP2-L in *P. berghei*
[62] as well as many of the most highly transcribed, H3K9ac marked genes, including two
etramps (PVP01_0734800 and PVP01_0504800), several RNA-binding proteins, including
Puf2, *ddx5*, a putative ATP-dependent RNA helicase DBP1 (PVP01_1429700), and a putative
bax1 inhibitor (PVP01_1465600). Interestingly, a number of highly transcribed and
translationally repressed genes associated with the ATATTTH motif, including *uis4*, *siap2*
and *pv1*, are not stably marked by H3K9ac in all replicates (i.e., there is significant variation
in the placement of the H3K9ac peak or their presence/absence among replicates for these
genes). It may be that additional histone modifications, for example H3K27me, H3R17me^3 or
H2 or H4 modifications, are involved in regulating transcription of these genes. Certainly the
H2A.Z modification, which is present in *P. falciparum*, and controls temperature responses in
plants [63] is intriguing as a potential mark regulating sporozoite fate in *P. vivax* considering
the association between hypnozoite activation rate and climate [11], as is H3R17me^3 in
consideration of the enrichment of markers/readers of this modification in HPZs noted above
and the role of this mark in cell fate progression in other species [48].
624
Fig. 5 *Schematic of potential mechanisms underpinning development in differentiation of* P.
vivax *sporozoites during liver-stage infection as hypnozoites and schizonts. We suggest*
differentiation programming at different points in development; first, schizont or hypnozoite
fate possibly encoded in the sporozoite as epigenetic signals or translationally repressed
transcripts; secondly, suppression signals that halt progression of the hypnozoite to schizont
stage and support persistence; and finally activation signals signified by a release in
chromatin, (post)transcriptional and (post)translational control leading to up-regulation of
replication, metabolic and protein export pathways.
633
CONCLUSIONS
We provide the first comprehensive study of the transcriptome and epigenome of mature
Plasmodium vivax sporozoites and undertake detailed comparisons with recently published
proteomic data for *P. vivax* sporozoites [32] and transcriptomic data for *P. vivax* mixed and
hypnozoite-enriched liver-stages [22] and mixed blood-stages [38]. These data support the
proposal that the sporozoite is a highly-programmed stage that is primed for invasion of and
development in the host hepatocyte. Cellular regulation, including at transcription,
translational and epigenetic levels, appears to play a major role in shaping this stage (which
continues on in some form in hypnozoites), and many of the genes proposed here as being
under translational repression are involved in hepatocyte infection and early liver-stage
development (**Fig. 5**). We highlight a major role for RNA-binding proteins, including PUF2,
ALBA2/4 and, intriguingly, 'Homologue of Musashi' (HoMu). We find that transcriptionally,
the hypnozoite appears to be a transition point between the sporozoite and replicating
schizonts, having many of the dominant sporozoite transcripts and retaining high transcription
of a number of key regulatory pathways involved in transcription, translation and chromatin
configuration (including histone arginine methylation). Our data support recent findings in
other *Plasmodium* sporozoites and liver-stages for an important role in AP2 transcription
factors, including AP2-SP2 and AP2-L. The data for AP2-Q, recently described as a marker
for hypnozoites in *P. cynomolgi* [23, 54] is less clear and merits further study. A consistent
theme in the study is the prominence of a number of genes that have a role in numerous
eukaryotic systems in cell fate determination and differentiation (e.g., HoMu, Yippee and
CARM1) and overlap with dormancy and/or persistent cell states in bacteria, protists or
higher eukaryotes (e.g., bacterial sucB and gamete fusion protein HAP2). These data do not
point to one single programming switch for dormancy or liver developmental fate in *P. vivax*
but present a number of intriguing avenues for exploration in subsequent studies, particularly
in model species such as *P. cynomolgi*. Our study provides a key foundation for
understanding the early stages of hepatocyte infection and the developmental switch between
liver trophozoite and hypnozoite formation. Importantly, it is a major first step in rationally
prioritizing targets underpinning liver-stage differentiation for functional evaluation in

humanized mouse and simian models for relapsing *Plasmodium* species and identifying novel
avenues to understand and eradicate liver-stage infections.
665
\
MATERIALS AND METHODS
Ethics Statement. Collection of venous blood from human patients with naturally acquired
vivax infection for the current study was approved by the Ethical Review Committee of the
Faculty of Tropical Medicine, Mahidol University (Human Subjects Protocol number TMEC
11-033) with the informed written consent of each donor individual. All mouse tissue used in
the current study was from preserved infected tissues generated previously [13]. All mouse
infection work in [13] was carried out at the Centre for Infectious Diseases Research (CIDR)
in Seattle, USA, under direct approval of the CIDR Institutional Animal Care and Use
Committee (IACUC) and performed in strict accordance with the recommendations in the
Guide for the Care and Use of Laboratory Animals of the National Institutes of Health, USA.
The Centre for Infectious Disease Research Biomedical Research Institute has an Assurance
from the Public Health Service (PHS Assurance number is A3640-01) through the Office of
Laboratory Animal Welfare (OLAW) for work approved by its IACUC.
680
Material collection, isolation and preparation. Nine field isolates (PvSpz-Thai 1 to 9),
representing symptomatic blood-stage malaria infections were collected as venous blood (20
mL) from patients presenting at malaria clinics in Tak and Ubon Ratchatani provinces in
Thailand. Each isolate was used to establish, infections in *Anopheles dirus* colonized at
Mahidol University (Bangkok) by membrane feeding [13], after 14-16 days post blood
feeding, ~3-15 million sporozoites were harvested per field isolate from the salivary glands of
up to 1,000 of these mosquitoes as per [64] and shipped in preservative (trizol (RNA/DNA) or
1% paraformaldehyde (DNA for ChiP-seq) to the Walter and Eliza Hall Institute (WEHI).
689
Transcriptomics sequencing and differential analysis. Upon arrival at WEHI, messenger
RNAs were purified from an aliquot (~0.5-1 million sporozoites) of each *P. vivax* field isolate
as per [38] and subjected to RNA-seq on Illumina NextSeq using TruSeq library construction
chemistry as per the manufacturer's instructions. Raw reads for each RNA-seq replicate are
available through the Sequence Read Archive (XXX-XXX). Sequencing adaptors were
removed and low quality reads trimmed and filtered using Trimmomatic v. 0.36 [65]. To
remove host contaminants, processed reads were aligned, as single-end reads, to the
Anopholes dirus wrari2 genome (VectorBase version W1) using Bowtie2 [66] (--very-
sensitive preset). All non-host reads were then aligned to the manually curated transcripts of
the *P. vivax* P01 genome (http://www.genedb.org/Homepage/P*vivax*P01; [27]) using RSEM
[67] (pertinent settings: --bowtie2 --bowtie2-sensitivity-level very_sensitive --calc-ci --ci-
memory 10240 --estimate-rspd --paired-end). Transcript abundance for each gene in each
replicate was calculated by RSEM as raw count, posterior mean estimate expected counts
(pme-EC) and transcripts per million (TPM).
Transcriptional abundance in *P. vivax* sporozoites was compared qualitatively (by
ranked abundance) with previously published microarray data for *P. vivax* salivary-gland
sporozoites [24]. As a further quality control, these RNA-seq data were compared also with
previously published microarray data for *P. falciparum* salivary-gland sporozoites [29], as
well as RNA-seq data from salivary-gland sporozoites generated here for *P. falciparum*
(single replicate generated from *P. falciparum* 3D7 lab cultures isolated from *Anopholes*
stephensi and processed as above) and previously published for *P. yoelii* [28]. RNA-seq data
from these additional *Plasmodium* species were (re)analysed from raw reads and
transcriptional abundance for each species was determined (raw counts and pme-EC and TPM
data) as described above using gene models current as of 04-10-2016 (PlasmoDB release
v29). Interspecific transcriptional behaviour was qualitatively compared by relative ranked
abundance in each species using TPM data for single copy orthologs (SCOs; defined in
PlasmoDB) only, shared between *P. vivax* and *P. faliciparum* or shared among *P. vivax*, *P.*
falciparum and *P. yoelii*.

To define transcripts that were up-regulated in sporozoites, we remapped raw reads representing early (18-24 hours post-infection (HPI)), mid (30-40 HPI) and late (42-46 HPI) *P. vivax* blood-stage infections recently published by Zhu *et al* [38] to the *P. vivax* P01 transcripts using RSEM as above. All replicate data was assessed for mapping metrics, transcript saturation and other standard QC metrics using QualiMap v 2.1.3 [68]. Differential transcription between *P. vivax* salivary-gland sporozoites and mixed blood-stages [38] was assessed using pme-EC data in EdgeR [69] and limma [70] (differential transcription cut-off: $\geq$ 2-fold change in counts per million (CPM) and a False Discovery Rate (FDR) ≤ 0.05). Pearson Chi squared tests were used to detect over-represented Pfam domains and Gene Ontology (GO) terms among differentially transcribed genes in sporozoites (Bonferroni-corrected $p < 0.05$), based on gene annotations in PlasmoDB (release v29).

We also compared transcription of the sporozoite stages to recently published liver-stage data from Gural et al (ref) as per the sporozoite to blood-stage comparisons above, with the following modifcations: (1) EC values were normalized using the 'upper quartile' method instead of TMM, (2) differential transcription was assessed using a quasi-likelihood generalize linear model (instead of a linear model) and (3) an FDR threshold for significance of $\leq$ 0.01 was used instead of $\leq$ 0.05. These differences related to specific attributes of the liver-stage dataset, particularly the small number of replicates (n = 2) per experiment condition. Data visualization and interactive R-shiny plots were produced in R using the ggplot[71], ggplot2 [72], gplots(heatmap.2) [73] and Glimma [74] packages.

Assessment of Sporozoite RNA-seq transcriptome by selective RT-qPCR: Extracted RNA was DNase treated (Sigma D5307) as per manufacturer recommendations. RNA was quantified using the TapeStation High Sensitivity RNA kit (Agilent). Two intron-spanning primer pairs were designed per gene of interest using Primer3 and BLAST. Primer pairs were tested in two concentrations (0.75ng and 2.83ng per reaction) to determine efficiency and specificity. Product was run on a 1% agarose gel with ethidium bromide. Primer pairs indicating non-specific priming were removed. The resulting 11 primer pairs were used on four sporozoite samples; VUBR06, VUNL23, VUBR24, VTTY84. RNA was reverse transcribed (Sensifast, Bioline) and used at 0.75ng per reaction, run on a Roche LightCycler 480 II. Melt curves were assessed and products were run on a gel to ensure specificity again. Cp threshold was set automatically. ΔCp value was calculated as target gene – comparator gene (SERA and CelTOS were used). Data were log transformed and fold change calculated.

RT-qPCR Primers were as follows:

Name	Gene	Forward Primer	Reverse Primer
RPS27	PVX_122245	ACCACCTTGTTTAGCCATGC	TAATTTGCACTTTCCACCCGTT
D13	PVX_089510	CTGTACACGCACGAGCTGGC	CAGCTCCTTGACGCCACTG
G10	PVX_080110	ACGAGCTGTACTACAAGCGGA	TTTCTCCTGCACCAGGTAGTC
AP2	PVX_086995*	GCCCCACTGGAAGTTTTGGA	CGTTCAGCCGCTGGTAGTAT
SERA	PVX_003790	CTGAAGACCTCCAGGGACAAG	TTTCTGCCTCTCCAGTGATATCTTT
CelTOS	PVX_123510*	CCCCCAAAGGCAAAATGAACAA	CGCTCTTTCCCCTCAAGGAC
GEST	PVX_118040	GACATATCAAGCAGTGAGGGA	CATGTTGTGGCCTTTATATGCTG
ALBA4	PVX_083270	TATCAACGGAGCCTTTGCCG	GGACTTGATTTCCTCGTCGG
PUF2	PVX_089945	ATCATAGAGAACGTCGACAAGCTTA	CTACGTTTCCAGGTTGCTGATC
14-3-3	PVX_089505	GACAACTTGACCTTGTGGACGTC	TACTCGAGGCCTTCATCCTTCGATT
ZIPCO	PVX_001980	TTAGCTCAATTGCTTGTGGCTTTTT	TGCCACTAACTCCAAGGAAATAACT

* denotes single exon gene

Salivary-gland sporozoite and liver-stage immunofluorescence assays (IFAs). IFAs were
performed as per [13] using preserved, vivax infected mouse liver tissue generated previously
for that study. In [13], female FRG [fumarylacetoacetate hydrolase (F), recombination
activation gene 2 (R), interleukin-2 receptor subunit gamma (G)] triple KO mice engrafted
with human hepatocytes (FRG KO huHep) were purchased from Yecuris Corporation
(Oregon, USA). Mice were infected through intravenous injection into the tail with $3.5 \times$
10^5 to 1×10^6 sporozoites isolated from the salivary glands of infected mosquitoes in 100 µl
of RPMI media. Liver stages for the current study were obtained from 10µm formalin fixed
paraffin embedded day 7 liver stages generated previously [13] from FRG knockout huHep
mice;[13] these were deparaffinized prior to staining. Fresh salivary-gland sporozoites were
fixed in acetone per [13]. All cells were incubated twice for 3 minutes in Xylene, then 100%
Ethanol, and finally once for 3 minutes each in 95%, 70%, and 50% Ethanol. The cells were
rinsed in DI water and permeabilized immediately in 1XTBS, containing Triton X-100 and
30% hydrogen peroxide. The cells were blocked in 5% milk in 1XTBS. The hepatocytes were
stained overnight with a rabbit polyclonal LISP1 antibody (A), a rabbit polyclonal UIS4
antibody (B), and a rabbit polyclonal BIP antibody (C) in blocking buffer. The cells were
washed with 1XTBS and the primary antibodies were detected with goat anti-rabbit Alexa
Fluor 488 antibody (Life Technologies). The cells were washed in 1XTBS. The hepatocytes
were rinsed in KMNO4 and washed in 1XTBS. The cells were incubated in DAPI for 5
minutes.
776
Histone ChIP sequencing and analysis. Aliquots of 2 – 6 million freshly isolated
sporozoites were fixed with 1% paraformaldehyde for 10 min at 37°C and the reaction
subsequently quenched by adding glycine to a final concentration of 125 mM. After three
washes with PBS, sporozoite pellets were stored at -80°C and shipped to Australia. Nuclei
were released from the sporozoites by dounce homogenization in lysis buffer (10 mM Hepes
pH 7.9, 10 mM KCl, 0.1 mM EDTA, 0.1 mM EDTA, 1 mM DTT, 1x EDTA-free protease
inhibitor cocktail (Roche), 0.25% NP40). Nuclei were pelleted by centrifugation at 21,000 g
for 10 min at 4°C and resuspended in SDS lysis buffer (1% SDS, 10 mM EDTA, 50 mM Tris
pH 8.1, 1x EDTA-free protease inhibitor cocktail). Chromatin was sheared into 200–1000 bp
fragments by sonication for 16 cycles in 30 sec intervals (on/off, high setting) using a
Bioruptor (Diagenode) and diluted 1:10 in ChIP dilution buffer (0.01% SDS, 1.1% Triton X-
100, 1.2 mM EDTA, 16.7 mM Tris pH 8.1, 150 mM NaCl). Chromatin was precleared for 1
hour with protein A/G sepharose (4FastFlow, GE Healthcare) equilibrated in 0.1% BSA
(Sigma-Aldrich, USA) in ChIP dilution buffer. Chromatin from 3×10^5 nuclei was taken
aside as input material. Chromatin from approximately 3×10^6 sporozoite nuclei was used for
each ChIP. ChIP was carried out over night at 4°C with 5 µg of antibody (H3K9me3 (Active
Motif), H3K4me3 (Abcam), H3K9ac (Upstate), H4K16ac (Abcam)) and 10 µl each of
equilibrated protein A and G sepharose beads (4FastFlow, GE Healthcare). After washes in
low-salt, high-salt, LiCl, and TE buffers (EZ-ChIP Kit, Millipore), precipitated complexes
were eluted in 1% SDS, 0.1 M NaHCO3. Cross-linking of the immune complexes and input
material was reversed for 6 hours at 45°C after addition of 500 mM NaCl and 20 µg/ml of
proteinase K (NEB). DNA was purified using the MinElute® PCR purification kit (Qiagen)
and paired-end sequenced on Illumina NextSeq using TruSeq library construction chemistry
as per the manufacturer's instructions. Raw reads for each ChIP-seq replicate are available
through the Sequence Read Archive (XXX-XXX).
Fastq files were checked for quality using fastqc
(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and adapter sequences were
trimmed using cutadapt [75]. Paired end reads were mapped to the *P. vivax* P01 strain
genome annotation using Bowtie2 [66]. The alignment files were converted to Bam format,
sorted and indexed using Samtools [76]. ChIP peaks were called relative to input using
MACS2[77] in paired end mode with a q value less than or equal to 0.01. Peaks and peak
summits were converted to sorted BED files. Bedtools intersect[78] was used to identify
genes that intersected H3K9me3 peaks and Bedtools closest was used to identify genes that
were closest to and downstream of H3K9ac and H3K4me3 peak summits.

811
Sequence motif analysis. Conserved sequence motifs were identified using the program
DREME [79]. Only genes in the top decile of transcription showing no evidence of protein
expression in multiple salivary-gland sporozoite replicates were considered as putatively
translationally repressed (n = 170). We queried coding regions and regions upstream of the
transcriptional start site (TSS) for each gene, defined by Zhu *et al* [38] and/or predicted here
from all RNA-seq data using the Tuxedo suite [80], for enriched sequence motifs in
comparison to 170 genes found to be in the top decile of both transcriptional and expressional
abundance in the same sporozoite replicates. In searching for motifs associated with highly
transcribed genes with stable H3K9ac marks within 1kb of the TSS (or up to the 3' end of the
next gene upstream), we compared H3K9ac marked genes in the top decile of transcription to
the same number of H3K9ac marked genes in the bottom decile of transcription. In both
instances, an e-value threshold of 0.05 was considered the minimum threshold for statistical
significance.
825

References
1.   Organization WH. World Malaria Report 2015. WHO, Geneva. 2015.
2.   Feachem RG, Phillips AA, Hwang J, Cotter C, Wielgosz B, Greenwood BM, et al.
Shrinking the malaria map: progress and prospects. Lancet. 2010;376(9752):1566-78.
doi: 10.1016/S0140-6736(10)61270-6.
3.   Price RN, Douglas NM, Anstey NM. New developments in *Plasmodium vivax*
malaria: severe disease and the rise of chloroquine resistance. Curr Opin Infect Dis.
2009;22(5):430-5. doi: 10.1097/QCO.0b013e32832f14c1.
4.   Baird KJ. Malaria caused by *Plasmodium vivax*: recurrent, difficult to treat,
disabling, and threatening to life - averting the infectious bite preempts these hazards.
Pathogens and global health. 2013;107:475-9.
5.   Sattabongkot J, Tsuboi T, Zollner GE, Sirichaisinthop J, Cui L. *Plasmodium vivax*
transmission: chances for control? Trends Parasitol. 2004;20(4):192-8.
6.   Mueller I, Galinski MR, Baird JK, Carlton JM, Kochar DK, Alonso PL, et al. Key gaps
in the knowledge of *Plasmodium vivax*, a neglected human malaria parasite. Lancet
Infect Dis. 2009;9(9):555-66. doi: 10.1016/S1473-3099(09)70177-X.
7.   Lindner SE, Miller JL, Kappe SH. Malaria parasite pre-erythrocytic infection:
preparation meets opportunity. Cell Microbiol. 2012;14(3):316-24. doi: 10.1111/j.1462-
5822.2011.01734.x.
8.   Shin SC, Vanderberg JP, Terzakis JA. Direct infection of hepatocytes by
sporozoites of *Plasmodium berghei*. J Protozool. 1982;29(3):448-54.
9.   Mota MM, Pradel G, Vanderberg JP, Hafalla JC, Frevert U, Nussenzweig RS, et al.
Migration of *Plasmodium* sporozoites through cells before infection. Science.
2001;291(5501):141-4. doi: 10.1126/science.291.5501.141.
10.   Lysenko AJ, Beljaev A, Rybalka V. Population studies of *Plasmodium vivax*: 1. The
theory of polymorphism of sporozoites and epidemiological phenomena of tertian
malaria. Bulletin of the World Health Organization. 1977;55(5):541.
11.   White NJ. Determinants of relapse periodicity in *Plasmodium vivax* malaria.
Malar J. 2011;10:297. doi: 10.1186/1475-2875-10-297.
12.   Price RN, Tjitra E, Guerra CA, Yeung S, White NJ, Anstey NM. Vivax malaria:
neglected and not benign. Amer J Trop Med Hyg. 2007;77(6 Suppl):79-87.
13.   Mikolajczak SA, Vaughan AM, Kangwanrangsan N, Roobsoong W, Fishbaugher M,
Yimamnuaychok N, et al. *Plasmodium vivax* liver stage development and hypnozoite
persistence in human liver-chimeric mice. Cell Host Microbe. 2015;17(4):526-35. doi:
10.1016/j.chom.2015.02.011.
14.   Mueller A-K, Camargo N, Kaiser K, Andorfer C, Frevert U, Matuschewski K, et al.
Plasmodium liver stage developmental arrest by depletion of a protein at the parasite–
host interface. Proc Natl Acad Sci USA. 2005;102(8):3022-7.

15.   Silvie O, Briquet S, Muller K, Manzoni G, Matuschewski K. Post-transcriptional
silencing of UIS4 in *Plasmodium berghei* sporozoites is important for host switch. Mol
Microbiol. 2014;91(6):1200-13. doi: 10.1111/mmi.12528.
16.   Mackellar DC, O'Neill MT, Aly AS, Sacci JB, Jr., Cowman AF, Kappe SH.
Plasmodium falciparum PF10_0164 (ETRAMP10.3) is an essential parasitophorous
vacuole and exported protein in blood stages. Eukaryot Cell. 2010;9(5):784-94. doi:
10.1128/EC.00336-09.
17.   Dembele L, Franetich JF, Lorthiois A, Gego A, Zeeman AM, Kocken CH, et al.
Persistence and activation of malaria hypnozoites in long-term primary hepatocyte
cultures. Nat Med. 2014;20(3):307-12. doi: 10.1038/nm.3461.
18.   Malmquist NA, Moss TA, Mecheri S, Scherf A, Fuchter MJ. Small-molecule histone
methyltransferase inhibitors display rapid antimalarial activity against all blood stage
forms in *Plasmodium falciparum*. Proc Natl Acad Sci U S A. 2012;109(41):16708-13. doi:
10.1073/pnas.1205414109.
19.   Josling GA, Llinas M. Sexual development in Plasmodium parasites: knowing
when it's time to commit. Nat Rev Microbiol. 2015;13(9):573-87. doi:
10.1038/nrmicro3519.
20.   White MT, Karl S, Battle KE, Hay SI, Mueller I, Ghani AC. Modelling the
contribution of the hypnozoite reservoir to *Plasmodium vivax* transmission. Elife.
2014;3. doi: 10.7554/eLife.04692.
21.   Roobsoong W, Tharinjaroen CS, Rachaphaew N, Chobson P, Schofield L, Cui L, et
al. Improvement of culture conditions for long-term in vitro culture of *Plasmodium vivax*.
Malaria journal. 2015;14(1):1.
22.   Gural N, Mancio-Silva L, Miller AB, Galstian A, Butty VL, Levine SS, et al. In vitro
culture, drug sensitivity, and transcriptome of *Plasmodium vivax* hypnozoites. Cell Host
Microbe. 2018;23(3):395-406 e4.
23.   Cubi R, Vembar SS, Biton A, Franetich JF, Bordessoulles M, Sossau D, et al. Laser
capture microdissection enables transcriptomic analysis of dividing and quiescent liver
stages of *Plasmodium* relapsing species. Cell Microbiol. 2017. doi: 10.1111/cmi.12735.
24.   Westenberger SJ, McClean CM, Chattopadhyay R, Dharia NV, Carlton JM,
Barnwell JW, et al. A systems-based analysis of *Plasmodium vivax* lifecycle transcription
from human to mosquito. PLoS Negl Trop Dis. 2010;4(4):e653. doi:
10.1371/journal.pntd.0000653.
25.   Gomez-Diaz E, Yerbanga RS, Lefevre T, Cohuet A, Rowley MJ, Ouedraogo JB, et al.
Epigenetic regulation of *Plasmodium falciparum* clonally variant gene expression during
development in Anopheles gambiae. Sci Rep. 2017;7:40655. doi: 10.1038/srep40655.
26.   Zanghi G, Vembar SS, Baumgarten S, Ding S, Guizetti J, Bryant JM, et al. A specific
PfEMP1 is expressed in *P. falciparum* sporozoites and plays a role in hepatocyte
infection. Cell Rep. 2018;22(11):2951-63.
27.   Auburn S, Bohme U, Steinbiss S, Trimarsanto H, Hostetler J, Sanders M, et al. A
new Plasmodium vivax reference sequence with improved assembly of the
subtelomeres reveals an abundance of pir genes. Wellcome Open Res. 2016;1:4. Epub
2016/12/23. doi: 10.12688/wellcomeopenres.9876.1.
28.   Lindner SE, Mikolajczak SA, Vaughan AM, Moon W, Joyce BR, Sullivan WJ, Jr., et
al. Perturbations of *Plasmodium* Puf2 expression and RNA-seq of Puf2-deficient
sporozoites reveal a critical role in maintaining RNA homeostasis and parasite
transmissibility. Cell Microbiol. 2013;15(7):1266-83. doi: 10.1111/cmi.12116.
29.   Le Roch KG, Johnson JR, Florens L, Zhou Y, Santrosyan A, Grainger M, et al. Global
analysis of transcript and protein levels across the *Plasmodium falciparum* life cycle.
Genome research. 2004;14(11):2308-18.
30.   Mikolajczak SA, Silva-Rivera H, Peng X, Tarun AS, Camargo N, Jacobs-Lorena V, et
al. Distinct malaria parasite sporozoites reveal transcriptional changes that cause
differential tissue infection competence in the mosquito vector and mammalian host.
Molecular and cellular biology. 2008;28(20):6196-207.

31. Carlton JM, Adams JH, Silva JC, Bidwell SL, Lorenzi H, Caler E, et al. Comparative
genomics of the neglected human malaria parasite *Plasmodium vivax*. Nature.
2008;455(7214):757-63. doi: 10.1038/nature07327.
32. Swearingen KE, Lindner SE, Flannery EL, Vaughan AM, Morrison RD,
Patrapuvich R, et al. Proteogenomic analysis of the total and surface-exposed proteomes
of Plasmodium vivax salivary gland sporozoites. PLoS Negl Trop Dis.
2017;11(7):e0005791. Epub 2017/08/02. doi: 10.1371/journal.pntd.0005791.
33. Guerreiro A, Deligianni E, Santos JM, Silva PA, Louis C, Pain A, et al. Genome-wide
RIP-Chip analysis of translational repressor-bound mRNAs in the *Plasmodium*
gametocyte. Genome Biol. 2014;15(11):493. doi: 10.1186/s13059-014-0493-0.
34. Silvie O, Briquet S, Müller K, Manzoni G, Matuschewski K. Post-transcriptional
silencing of UIS4 in *Plasmodium berghei* sporozoites is important for host switch.
Molecular microbiology. 2014;91(6):1200-13.
35. Lindner SE, Swearingen KE, Harupa A, Vaughan AM, Sinnis P, Moritz RL, et al.
Total and putative surface proteomics of malaria parasite salivary gland sporozoites.
Mol Cell Proteomics. 2013;12(5):1127-43. doi: 10.1074/mcp.M112.024505.
36. Kelley KD, Miller KR, Todd A, Kelley AR, Tuttle R, Berberich SJ. YPEL3, a p53-
regulated gene that induces cellular senescence. Cancer Res. 2010;70(9):3566-75. doi:
10.1158/0008-5472.CAN-09-3219.
37. Tuttle R, Simon M, Hitch DC, Maiorano JN, Hellan M, Ouellette J, et al. Senescence-
associated gene YPEL3 is downregulated in human colon tumors. Ann Surg Oncol.
2011;18(6):1791-6. doi: 10.1245/s10434-011-1558-x.
38. Zhu L, Mok S, Imwong M, Jaidee A, Russell B, Nosten F, et al. New insights into
the *Plasmodium vivax* transcriptome using RNA-Seq. Sci Rep. 2016;6:20498. doi:
10.1038/srep20498.
39. Kramer S. RNA in development: how ribonucleoprotein granules regulate the life
cycles of pathogenic protozoa. WIR: RNA. 2014;5(2):263-84.
40. Tucker RP. The thrombospondin type 1 repeat superfamily. Int J Biochem Cell
Biol. 2004;36(6):969-74. doi: 10.1016/j.biocel.2003.12.011.
41. Ntumngia FB, Bouyou-Akotet MK, Uhlemann AC, Mordmuller B, Kremsner PG,
Kun JF. Characterisation of a tryptophan-rich *Plasmodium falciparum* antigen associated
with merozoites. Mol Biochem Parasitol. 2004;137(2):349-53. doi:
10.1016/j.molbiopara.2004.06.008.
42. Gubbels MJ, Vaishnava S, Boot N, Dubremetz JF, Striepen B. A MORN-repeat
protein is a dynamic component of the *Toxoplasma gondii* cell division apparatus. J Cell
Sci. 2006;119(Pt 11):2236-45. doi: 10.1242/jcs.02949.
43. Aly AS, Lindner SE, MacKellar DC, Peng X, Kappe SH. SAP1 is a critical post-
transcriptional regulator of infectivity in malaria parasite sporozoite stages. Mol
Microbiol. 2011;79(4):929-39. doi: 10.1111/j.1365-2958.2010.07497.x.
44. Okano H, Imai T, Okabe M. Musashi: a translational regulator of cell fate. Journal
of Cell Science. 2002;115(7):1355-9.
45. Cui L, Lindner S, Miao J. Translational regulation during stage transitions in
malaria parasites. Annals N Y Acad Sci. 2015;1342(1):1-9.
46. Lasko P. Gene regulation at the RNA layer: RNA binding proteins in intercellular
signaling networks. Sci STKE. 2003;179:RE6.
47. Guizetti J, Scherf A. Silence, activate, poise and switch! Mechanisms of antigenic
variation in *Plasmodium falciparum*. Cell microbiol. 2013;15(5):718-26.
48. Wu Q, Bruce AW, Jedrusik A, Ellis PD, Andrews RM, Langford CF, et al. CARM1 is
required in embryonic stem cells to maintain pluripotency and resist differentiation.
Stem Cells. 2009;27(11):2637-45. Epub 2009/06/23. doi: 10.1002/stem.131.
49. Shi S, Ehrt S. Dihydrolipoamide acyltransferase is critical for *Mycobacterium*
tuberculosis pathogenesis. Infect Immun. 2006;74(1):56-63. Epub 2005/12/22. doi:
10.1128/IAI.74.1.56-63.2006.

50.   Ma C, Sim S, Shi W, Du L, Xing D, Zhang Y. Energy production genes sucB and
ubiF are involved in persister survival and tolerance to multiple antibiotics and stresses
in Escherichia coli. FEMS Microbiol Lett. 2010;303(1):33-40. Epub 2010/01/01. doi:
10.1111/j.1574-6968.2009.01857.x.
51.   Schrader J, Moyle R, Bhalerao R, Hertzberg M, Lundeberg J, Nilsson P, et al.
Cambial meristem dormancy in trees involves extensive remodelling of the
transcriptome. Plant J. 2004;40(2):173-87. Epub 2004/09/28. doi: 10.1111/j.1365-
313X.2004.02199.x.
52.   Yazawa K, Kamada H. Identification and characterization of carrot HAP factors
that form a complex with the embryo-specific transcription factor C-LEC1. J Exp Bot.
2007;58(13):3819-28. Epub 2007/12/07. doi: 10.1093/jxb/erm238.
53.   Wood FC, Heidari A, Tekle YI. Genetic Evidence for Sexuality in *Cochliopodium*
(Amoebozoa). J Hered. 2017;108(7):769-79. Epub 2017/10/17. doi:
10.1093/jhered/esx078.
54.   Voorberg-van der Wel A, Roma G, Gupta DK, Schuierer S, Nigsch F, Carbone W, et
al. A comparative transcriptomic analysis of replicating and dormant liver stages of the
relapsing malaria parasite *Plasmodium cynomolgi*. Elife. 2017;6. Epub 2017/12/08. doi:
10.7554/eLife.29605.
55.   Lopez-Rubio J-J, Mancio-Silva L, Scherf A. Genome-wide analysis of
heterochromatin associates clonally variant gene regulation with perinuclear repressive
centers in malaria parasites. Cell host & microbe. 2009;5(2):179-90.
56.   Duffy MF, Selvarajah SA, Josling GA, Petter M. Epigenetic regulation of the
Plasmodium falciparum genome. Brief Funct Genomics. 2014;13(3):203-16. doi:
10.1093/bfgp/elt047.
57.   Cui L, Miao J, Furuya T, Li X, Su XZ, Cui L. PfGCN5-mediated histone H3
acetylation plays a key role in gene expression in *Plasmodium falciparum*. Eukaryot Cell.
2007;6(7):1219-27. doi: 10.1128/EC.00062-07.
58.   Rovira-Graells N, Gupta AP, Planet E, Crowley VM, Mok S, de Pouplana LR, et al.
Transcriptional variation in the malaria parasite *Plasmodium falciparum*. Genome
research. 2012;22(5):925-38.
59.   De Silva EK, Gehrke AR, Olszewski K, León I, Chahal JS, Bulyk ML, et al. Specific
DNA-binding by apicomplexan AP2 transcription factors. Proc Natl Aacd Sci.
2008;105(24):8393-8.
60.   Painter HJ, Campbell TL, Llinás M. The Apicomplexan AP2 family: integral factors
regulating *Plasmodium* development. Molecular and biochemical parasitology.
2011;176(1):1-7.
61.   Kafsack BF, Rovira-Graells N, Clark TG, Bancells C, Crowley VM, Campino SG, et
al. A transcriptional switch underlies commitment to sexual development in human
malaria parasites. Nature. 2014;507(7491):248.
62.   Iwanaga S, Kaneko I, Kato T, Yuda M. Identification of an AP2-family protein that
is critical for malaria liver stage development. PLoS One. 2012;7(11):e47557.
63.   Boden SA, Kavanova M, Finnegan EJ, Wigge PA. Thermal stress effects on grain
yield in *Brachypodium distachyon* occur via H2A.Z-nucleosomes. Genome Biol.
2013;14(6):R65. doi: 10.1186/gb-2013-14-6-r65.
64.   Kennedy M, Fishbaugher ME, Vaughan AM, Patrapuvich R, Boonhok R,
Yimamnuaychok N, et al. A rapid and scalable density gradient purification method for
Plasmodium sporozoites. Malar J. 2012;11:421. doi: 10.1186/1475-2875-11-421.
65.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 2014;30(15):2114-20. doi:
10.1093/bioinformatics/btu170.
66.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat
Methods. 2012;9(4):357-9. doi: 10.1038/nmeth.1923.
67.   Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. BMC Bioinformatics. 2011;12(1):323.

68.	Okonechnikov K, Conesa A, Garcia-Alcalde F. Qualimap 2: advanced multi-
sample quality control for high-throughput sequencing data. Bioinformatics.
2016;32(2):292-4. doi: 10.1093/bioinformatics/btv566.
69.	Nikolayeva O, Robinson MD. edgeR for differential RNA-seq and ChIP-seq
analysis: an application to stem cell biology. Methods Mol Biol. 2014;1150:45-79. doi:
10.1007/978-1-4939-0512-6_3.
70.	Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers
differential expression analyses for RNA-sequencing and microarray studies. Nucleic
acids research. 2015;43(7):e47-e.
71.	Wickham H. ggplot: An implementation of the Grammar of Graphics in R, 2006. R
package version 04 0.
72.	Wickham H, Chang W. ggplot2: An implementation of the Grammar of Graphics.
R package version 07, URL: http://CRAN R-project org/package= ggplot2. 2008.
73.	Warnes GR, Bolker B, Bonebakker L, Gentleman R, Huber W, Liaw A, et al. gplots:
Various R programming tools for plotting data. R package version. 2009;2(4):1.
74.	Law CW, Alhamdoosh M, Su S, Smyth GK, Ritchie ME. RNA-seq analysis is easy as
1-2-3 with limma, Glimma and edgeR. F1000Research. 2016;5.
75.	Martin M. Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet journal. 2011;17(1):pp. 10-2.
76.	Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9. doi:
10.1093/bioinformatics/btp352.
77.	Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-
based analysis of ChIP-Seq (MACS). Genome Biol. 2008;9(9):R137. doi: 10.1186/gb-
2008-9-9-r137.
78.	Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. 2010;26(6):841-2. doi: 10.1093/bioinformatics/btq033.
79.	Bailey TL. DREME: motif discovery in transcription factor ChIP-seq data.
Bioinformatics. 2011;27(12):1653-9.
80.	Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, et al. Differential gene
and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks.
Nat Protoc. 2012;7(3):562-78. doi: 10.1038/nprot.2012.016.
1057

DIFFERENTIATION

- **Translational repressors**
  e.g., Puf2, Alba2/4, HOMU, Yippee, Zipco
- **Invasion**
  e.g., Celtos, Gest
- **Liver development**
  e.g., LISP1, LISP2, ETRAMPS, TRAP
- **Transcription factors**
  e.g., AP2-SP2, AP2-L, AP2-Q?

SUPPRESSION

- **Metabolic/replication suppressors**
  e.g., sucB, HAP2, MAK16
- **Histone arginine methylation**
  e.g., CARM1, EEML2
- **Protein translation**
  e.g., eIF-3H, Puf1

ACTIVATION

- **Merozoite development**
  e.g., MSP1, MSP3, MSP9
- **Rhoptry function**
  e.g., RAP1, RNP2, RNP3
- **Reticuocytye binding**
  e.g., RBP2a, RBP2b, RBP2C
- **Exported Proteins**
  e.g., PHISTs