A Nasal Brush-based Classifier of Asthma Identified by Machine Learning Analysis of

Nasal RNA Sequence Data

Authors: Gaurav Pandey[1], Om P. Pandey[1], Angela J. Rogers[2], Mehmet E. Ahsen[1], Gabriel E.

Hoffman[1], Benjamin A. Raby[3], Scott T. Weiss[3], Eric E. Schadt[1], Supinda Bunyavanich[1,4]*

Affiliations:

[1] Icahn Institute for Genomics and Multiscale Biology and Department of Genetics and Genomic

Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[2] Division of Pulmonary and Critical Care Medicine, Department of Medicine, Stanford University

School of Medicine, Stanford, CA, USA.

[3] Channing Division of Network Medicine and Division of Pulmonary and Critical Care Medicine,

Brigham & Women's Hospital, and Harvard Medical School, Boston, MA, USA.

[4] Division of Allergy & Immunology, Department of Pediatrics, Icahn School of Medicine at Mount

Sinai, New York, NY, USA.

*To whom correspondence should be addressed: Supinda Bunyavanich, MD, MPH, Icahn

School of Medicine at Mount Sinai, 1425 Madison Avenue #1498, New York, NY 10029, USA,

Tel. +1 212 659 8262, Fax +1 212 426 1902, supinda@post.harvard.edu

ABSTRACT:

Asthma is a common, under-diagnosed disease affecting all ages. We sought to identify a nasal

brush-based classifier of mild/moderate asthma. One hundred ninety subjects with

mild/moderate asthma and controls underwent nasal brushing and RNA sequencing of nasal

samples. A machine learning-based pipeline, comprised of feature selection, classification, and

statistical analyses, identified a classifier of asthma consisting of 90 nasally expressed genes

interpreted via an L2-regularized logistic regression classification model. This nasal brush-

based classifier performed with strong predictive value and sensitivity across eight test sets,

including (1) a test set of independent asthmatic and non-asthmatic subjects profiled by RNA

sequencing (positive and negative predictive values of 1.00 and 0.96, respectively; AUC of

0.994), (2) two independent case-control cohorts of asthma profiled by microarray, and (3) five

independent cohorts of subjects with other respiratory conditions (allergic rhinitis, upper

respiratory infection, cystic fibrosis, smoking), where the classifier had a low to zero rate of

misclassification. Following validation in large, prospective cohorts, this classifier could be

developed into a nasal brush-based biomarker of asthma.

Introduction

Asthma is a chronic respiratory disease that affects 8.6% of children and 7.4% of adults in the United States [1]. Its true prevalence may be higher [2]. The fluctuating airflow obstruction, bronchial hyper-responsiveness, and airway inflammation that characterize mild to moderate asthma can be difficult to detect in busy, routine clinical settings [3]. In one study of US middle school children, 11% reported physician-diagnosed asthma with current symptoms, while an additional 17% reported active asthma-like symptoms without a diagnosis of asthma [2]. Undiagnosed asthma leads to missed school and work, restricted activity, emergency department visits, and hospitalizations [2, 4]. Given the high prevalence of asthma and consequences of missed diagnosis, there is high potential impact of improved biomarkers for asthma [5].

National and international guidelines recommend that the diagnosis of asthma be based on a history of typical symptoms *and objective findings* of variable expiratory airflow limitation [6, 7]. However, obtaining such objective findings can be challenging given currently available tools. Pulmonary function tests (PFTs) require equipment, expertise, and experience to execute well [8, 9]. Results are unreliable if the procedure is done with poor technique [8]. PFTs are usually not immediately available in primary care settings. Despite the published guidelines, PFTs are not done in over half of patients suspected of having asthma [8]. Induced sputum and exhaled nitric oxide have been explored as asthma biomarkers, but their implementation requires technical expertise and does not yield better clinical results than physician-guided management alone [10]. Given the above, the reality is that most asthma is still clinically diagnosed and managed based on self-report [8, 9]. This is problematic because most patients with asthma are frequently asymptomatic at the time of exam and under-perceive as well as under-report symptoms [11].

A nasal biomarker of asthma is of high interest given the accessibility of the nose and shared airway biology between the upper and lower respiratory tracts [12-15]. The easily

accessible nasal passages are directly connected to the lungs and exposed to common

environmental factors. Here we describe first steps toward the development of a nasal

biomarker of asthma by reporting our identification of an asthma classifier using nasal gene

expression data and machine learning (**Figure 1**). Specifically, we used RNA sequencing

(RNAseq) to comprehensively profile gene expression from nasal brushings collected from

subjects with mild to moderate asthma and controls, creating the largest nasal RNAseq data set

in asthma to date. We focused on mild to moderate asthma because the waxing and waning

nature of non-severe asthma render it relatively difficult to diagnose. Using a robust machine

learning-based pipeline comprised of feature selection [16], classification [17], and statistical

analyses [18], we identified an asthma classifier that accurately differentiates between subjects

with and without mild-moderate asthma based on nasal brushings. We evaluated the

classification performance of this asthma classifier on eight test sets of independent subjects

with asthma and other respiratory conditions, finding that it performed with high accuracy,

sensitivity, and specificity for asthma. Although this study's focus is asthma, the pipeline

described could potentially be used to develop classifiers for other phenotypes with high-

dimensional data.

Results

Study population and baseline characteristics

We performed nasal brushing on 190 subjects for this study, including 66 subjects with

well-defined mild to moderate persistent asthma (based on symptoms, medication need, and

demonstrated airway hyper-responsiveness by methacholine challenge) and 124 subjects

without asthma (based on no personal or family history of asthma, normal spirometry, and no

bronchodilator response). The definitional criteria we used for mild-moderate asthma are

consistent with US National Heart Lung Blood Institute guidelines for the diagnosis of asthma

[7], and are the same criteria used in the longest NIH-sponsored study of mild-moderate asthma

[19, 20].

From these 190 subjects, a random selection of 150 subjects were *a priori* assigned as

the development set (to be used for asthma classifier development), and the remaining 40

subjects were *a priori* assigned as the RNAseq test set (to be used as one of 8 test sets for

evaluation of the asthma classifier identified from the development set).

The baseline characteristics of the subjects in the development set (n=150) are shown in

the left section of **Table 1**. The mean age of subjects with asthma was somewhat lower than

subjects without asthma, with slightly more male subjects with asthma and more female

subjects without asthma. Caucasians were more prevalent in subjects without asthma, which

was expected based on the inclusion criteria. Consistent with reversible airway obstruction that

characterizes asthma [3], subjects with asthma had significantly greater bronchodilator

response than control subjects (T-test P = 1.4×10^{-5}). Allergic rhinitis was more prevalent in

subjects with asthma (Fisher's exact test P = 0.005), consistent with known comorbidity

between allergic rhinitis and asthma [21]. Rates of smoking between subjects with and without

asthma were not significantly different.

RNA isolated from nasal brushings from the subjects was of good quality, with mean RIN

7.8 (±1.1). The median number of paired-end reads per sample from RNA sequencing was 36.3

million. Following pre-processing (normalization and filtering) of the raw RNAseq data, 11,587

genes were used for statistical and machine learning analysis. variancePartition analysis [22],

which is designed to analyze the contribution of technical and biological factors to variation in

gene expression, showed that age, race, and sex contributed minimally to total gene expression

variance (**Supplementary Figure 1**). For this reason, we did not adjust the pre-processed

RNAseq data for these factors.

Differential gene expression analysis by DeSeq2 [23] showed that 1613 and 1259 genes

were respectively over- and under-expressed in asthma cases versus controls (false discovery

rate (FDR) ≤0.05) (**Supplementary Table 1**). These genes were enriched for disease-relevant

pathways in the Molecular Signature Database [24], including immune system (fold change=3.6,

FDR=1.07 x 10^{-22}), adaptive immune system (fold change=3.91, FDR=1.46 x 10^{-15}), and innate

immune system (fold change=4.1, FDR=4.47 x 10^{-9}) (**Supplementary Table 1**).

Identifying a nasal brush-based asthma classifier

To identify a nasal brush-based asthma classifier using the RNAseq data generated, we

developed a machine learning pipeline that combined feature (gene) selection [16] and

classification techniques [17] that was applied to the development set (**Materials and Methods**

and **Supplementary Figure 2**). This pipeline was designed with a systems biology-based

perspective that a set of genes, even ones with marginal effects, can collectively classify

phenotypes (here asthma) more accurately than individual genes [25].

Feature selection [16] is the process of identifying a subset of features (e.g. genes) from

a much larger subset in an automated data-driven fashion. In our pipeline, this process involved

a cross validation-based protocol [26] using the well-established Recursive Feature Elimination

(RFE) algorithm [16] combined with L_2-regularized Logistic Regression (LR or Logistic) and

Support Vector Machine (SVM-Linear (kernel)) algorithms [17] (combinations referred to as LR-

RFE and SVM-RFE respectively) (**Supplementary Figure 3**). Classification analysis was then

performed by applying four global classification algorithms (SVM-Linear, AdaBoost, Random

Forest, and Logistic) [17] to the expression profiles of the gene sets identified by feature

selection. To reduce the potential adverse effect of overfitting, this process (feature selection

and classification) was repeated 100 times on 100 random splits of the development set into

training and holdout sets. The final classifier was selected by statistically comparing the models

in terms of both classification performance and parsimony, i.e., the number of genes included in

the model [18] (**Supplementary Figure 4)**.

Due to the imbalance of the two classes (asthma and controls) in our cohort (consistent

with imbalances in the general population for asthma and most disease states), we used F-

measure as the main evaluation metric in our study [27, 28]. This class-specific measure is a

conservative mean of precision (predictive value) and recall (same as sensitivity), and is

described in detail in **Box 1** and **Supplementary Figure 5**. F-measure can range from 0 to 1,

with higher values indicating superior classification performance. An F-measure value of 0.5

does not represent a random model. To provide context for our performance assessments, we

also computed commonly used evaluation measures, including positive and negative predictive

values (PPVs and NPVs) and Area Under the Receiver Operating Characteristic (ROC) Curve

(AUC) scores (**Box 1** and **Supplementary Figure 5)**.

____

Box 1: Evaluation measures for classifiers

Many measures exist for evaluating the performance of classifiers. The most commonly

used evaluation measures in biology and medicine are the positive and negative predictive

values (PPV and NPV respectively; **Supplementary Figure 5**), and Area Under the Receiver

Operating Characteristic (ROC) Curve (AUC score) [27]. However, these measures have

several limitations. PPV and NPV ignore the critical dimension of sensitivity [27]. For instance, a

classifier may predict perfectly for only one asthma sample in a cohort and make no predictions

for all other asthma samples. This will yield a PPV of 1, but poor sensitivity, since none of the

other asthma samples were identified by the classifier. ROC curves and their AUC scores do

not accurately reflect performance when the number of cases and controls in a sample are

imbalanced [27, 28], which is frequently the case in biomedical studies. For such situations,

precision, recall, and F-measure (**Supplementary Figure 5**) are considered more meaningful

performance measures for classifier evaluation [28]. Note that precision for cases (e.g. asthma)

is equivalent to PPV, and precision for controls (e.g. no asthma) is equivalent to NPV

(**Supplementary Figure 5**). Recall is the same as sensitivity. F-measure is the harmonic

(conservative) mean of precision and recall that is computed separately for each class, and thus

provides a more comprehensive and reliable assessment of model performance for cohorts with

unbalanced class distributions. For the above reasons, we consider F-measure as the primary

evaluation measure in our study, although we also provide PPV, NPV and AUC measures for

context. Like PPV, NPV and AUC, F-measure ranges from 0 to 1, with higher values indicating

superior classification performance, but a value of 0.5 for F-measure does not represent a

random model and could in some cases indicate superior performance over random.

The best performing and most parsimonious combination of feature selection and

classification algorithm identified by our machine learning pipeline was LR-RFE & Logistic

Regression (**Supplementary Figure 4**). The classifier inferred using this combination was built

on 90 predictive genes and will be henceforth referred to as the *asthma classifier*. We

emphasize that the expression values of the classifier's 90 genes must be used in combination

with the Logistic classifier and the model's optimal classification threshold (i.e. predicted

label=asthma if classifier's probability output≥0.76, else predicted label=no asthma) to be used

effectively for asthma classification.

Evaluation of the asthma classifier in an RNAseq test set of independent subjects

Our next step was to evaluate the asthma classifier in an RNAseq test set of

independent subjects, for which we used the test set (n=40) of nasal RNAseq data from

independent subjects. The baseline characteristics of the subjects in this test set are shown in

the right section of **Table 1**. Subjects in the development and test sets were generally similar,

except for a lower prevalence of allergic rhinitis among those without asthma in the test set.

The asthma classifier performed with high accuracy in the RNAseq test set's

independent subjects, achieving AUC = 0.994 (**Figure 2**), PPV = 1.00, and NPV = 0.96

(**Figures 3B and 3D, left most bar**). In terms of the F-measure metric, the classifier achieved F

= 0.98 and 0.96 for classifying asthma and no asthma, respectively (**Figures 3A and 3C, left

most bar**). For comparison, the much lower performance of permutation-based random models

is shown in **Supplementary Figure 6**.

Our machine learning pipeline evaluated models from several combinations of feature

selection and classification algorithms to select the most predictive classifier. Potentially

predictive genes can also be identified from differential expression analysis and results from

prior asthma-related studies. **Figure 4** shows the performance of the asthma classifier in the

RNAseq test set next to alternative classifiers trained on the development set using: (1) other

classifiers tested in our machine learning pipeline, (2) all genes in our data set (11587 genes

after filtering), (3) all differentially expressed genes in the development set (2872 genes)

(**Supplementary Table 1**), (4) genes associated with asthma from prior studies [29] (70 genes)

(**Supplementary Table 2**), and (5) a commonly used one-step classification model (L1-Logistic)

[30] (243 genes). The asthma classifier identified by our pipeline outperformed all these

alternative classifiers despite its reliance on a small number of genes.

We emphasize that our classifier produced more accurate predictions than models using

all genes, all differentially expressed genes, and all known asthma genes. This supports that

data-driven methods can build more effective classifiers than those built exclusively on

traditional statistical methods (which do not necessarily target classification), and current

domain knowledge (which may be incomplete and subject to investigation bias). Our classifier

also outperformed and was more parsimonious than the model learned using the commonly

used L1-Logistic method, which combined feature selection and classification into a single step.

The fact that our asthma classifier performed well in an independent RNAseq test set while also

outperforming alternative models lends confidence to its classification ability.

Evaluation of the asthma classifier in external asthma cohorts

To assess the performance of our asthma classifier in other populations and profiling

platforms, we applied the classifier to nasal gene expression data generated from independent

cohorts of asthmatics and controls profiled by microarrays: Asthma1 (GEO GSE19187) [31] and

Asthma2 (GEO GSE46171) [32]. To the best of our knowledge, there are no publicly available

nasal RNAseq data sets for cohorts with asthma (prior to our study). **Supplementary Table 3**

summarizes the characteristics of these external, independent case-control cohorts. In general,

RNAseq-based predictive models are not expected to translate well to microarray-profiled

samples [33, 34]. A major reason is that gene mappings do not perfectly correspond between

RNAseq and microarray due to disparities between array annotations and RNAseq gene models

[34]. Our goal was to assess the performance of our asthma classifier despite discordances in

study designs, sample collections, and gene expression profiling platforms.

The asthma classifier performed relatively well (**Figure 3 middle bars**) and consistently

better than permutation-based random models (**Supplementary Figure 6**) in classifying asthma

and no asthma in both the Asthma1 and Asthma2 microarray-based test sets. The classifier

achieved similar F-measures in the two test sets (**Figures 3A and 3C middle bars**), although

the PPV and NPV measures were more dissimilar for Asthma2 (PPV 0.93, NPV 0.31) than for

Asthma1 (PPV 0.61, NPV 0.67) (**Figure 3B and 3D middle bars**). The classifier's performance

was better than its random counterparts for both these test sets, although the difference in this

performance was smaller for Asthma2. This occurred partially because Asthma2 includes many

more asthma cases than controls (23 vs. 5), which is counter to the expected distribution in the

general population. In such a skewed data set, it is possible for a random model to yield an

artificially high F-measure for asthma by predicting every sample as asthmatic. We verified that

this occurred with the random models tested on Asthma2.

To assess how the asthma classifier might perform in a larger external test set, we

combined samples from Asthma1 and Asthma2 and performed the evaluation on this combined

set. We chose this approach because no single large, external dataset of nasal gene expression

in asthma exists, and combining cohorts could yield a joint test set with heterogeneity that

partially reflects real-life heterogeneity of asthma. As expected, all the performance measures

for this combined test set were intermediate to those for Asthma1 and Asthma2 (**Figure 3 right**

most bars). These results supported that our classifier also performs reasonably well in a larger

and more heterogeneous cohort.

Overall, despite the discordance of gene expression profiling platforms, study designs,

and sample collection methods, our asthma classifier performed reasonably well in these

external test sets, supporting a degree of generalizability of the classifier across platforms and

cohorts.

Specificity of the asthma classifier: testing in external cohorts with non-asthma respiratory

conditions

To assess the specificity of our asthma classifier, we next sought to determine if it would

misclassify as asthma other respiratory conditions with symptoms that overlap with asthma. To

this end, we evaluated the performance of the asthma classifier on nasal gene expression data

derived from case-control cohorts with allergic rhinitis (GSE43523) [35], upper respiratory

infection (GSE46171) [32], cystic fibrosis (GSE40445) [36], and smoking (GSE8987) [12].

Supplementary Table 4 details the characteristics for these external cohorts with non-asthma

respiratory conditions. In three of these five non-asthma cohorts (Allergic Rhinitis, Cystic

Fibrosis and Smoking), the classifier appropriately produced one-sided classifications, i.e.,

samples were all appropriately classified as "no asthma." This is shown by the zero F-measure

for the positive (asthma) class (**Figure 5A**) and perfect F-measure for the negative (no asthma)

class (**Figure 5C**) obtained by the classifier in these cohorts. In other words, the precision for

the asthma class (PPV) of our classifier was exactly and appropriately zero (**Figure 5B**), and

NPV was perfectly 1.00 for these cohorts with non-asthma conditions (**Figures 5D**). The URI

day 2 and 6 cohorts were slight deviations from these trends, where the classifier achieved

perfect NPVs of 1.00 (**Figure 5D**), but marginally lower F-measure for the "no asthma" class

(**Figure 5C**) due to slightly lower than perfect sensitivity. This may have been influenced by

common inflammatory pathways underlying early viral inflammation and asthma [37].

Nonetheless, consistent with the other non-asthma test sets, the classifier's misclassification of

URI as asthma was rare and substantially less than its random counterpart classifiers

(**Supplementary Figure 7**).

To assess the asthma classifier's performance if presented with a large, heterogeneous

collection of non-asthma respiratory conditions reflective of real clinical settings, we aggregated

the non-asthma cohorts into a "Combined non-asthma" test set and applied the asthma

classifier. The results included an appropriately zero F-measure for asthma and zero PPV, and

an F-measure of 0.97 for no asthma, and NPV of 1.00 (**Figure 5, right most bars**). Results

from the individual and combined non-asthma test sets collectively support that the asthma

classifier would rarely misclassify other respiratory diseases as asthma.

Statistical and Pathway Examination of Genes in the Asthma Classifier

282   An interesting question to ask for a disease classifier is how does its predictive ability

relate to the individual differential expression status of the genes constituting the classifier? We

found that 46 of the 90 genes included in our classifier were differentially expressed (FDR

≤0.05), with 22 and 24 genes over- and under-expressed in asthma respectively (**Figure 6,**

Supplementary Table 1). More generally, the genes in our classifier had lower differential

expression FDR values than other genes (Kolmogorov-Smirnov statistic=0.289, P-

value=2.73×10^{-37}) (**Supplementary Figure 8**).

289   In terms of biological function, pathway enrichment analysis of our classifier's 90 genes,

though statistically limited by the small number of genes, yielded enrichment for pathways

including defense response (fold change=2.86, FDR=0.006) and response to external stimulus

(fold change=2.50, FDR=0.012). Only four (*C3, DEFB1, CYFIP2* and *GSTT1*) of the 90 genes

are known asthma genes and are functionally involved in complement activation, microbicidal

activity, T-cell differentiation, and oxidative stress, respectively [29]. These results suggest that

our machine learning pipeline was able to extract information beyond individually differentially

expressed or previously known disease-related genes, allowing for the identification of a

parsimonious set of genes that collectively enabled accurate disease classification.

Discussion

Using RNAseq data generated from our cohorts, combined with a systematic machine

learning analysis approach, we identified a nasal brush-based classifier that accurately

distinguishes subjects with mild/moderate asthma from controls. This asthma classifier,

consisting of the expression profiles of 90 genes interpreted via a logistic regression

classification model, performed with high precision (PPV=1.00 and NPV=0.96) and recall for

classifying asthma (AUC=0.994). The performance of the asthma classifier across independent

test sets demonstrates potential for the classifier's generalizability across study populations and

two major modalities of gene expression profiling (RNAseq and microarray). Additionally, the

classifier's low to zero rate of misclassification on external cohorts with non-asthma respiratory

conditions supports the specificity of this asthma classifier. Our results represent the first steps

toward the development of a nasal biomarker of asthma.

Our nasal brush-based asthma classifier is based on the common biology of the upper

and lower airway, a concept supported by clinical practice and previous findings [12-15].

Clinicians often rely on the united airway by screening for lower airway infections (e.g. influenza,

methicillin-resistant Staphylococcus aureus) with nasal swabs [38]. Sridhar et al. found that

gene expression consequences of tobacco smoking in bronchial epithelial cells were reflected in

nasal epithelium [12]. Wagener et al. compared gene expression in the nasal and bronchial

epithelia from 17 subjects, finding that 99% of the 33,000 genes tested exhibited no differential

expression between the nasal and bronchial epithelia in those with airway disease [13]. In a

study of 30 children, Guajardo et al. identified gene clusters with differential expression in nasal

epithelium between subjects with exacerbated asthma vs. controls [14]. The above studies were

done with small sample sizes and microarray technology. More recently, Poole et al. compared

RNAseq profiles of nasal brushings from 10 asthmatic and 10 control subjects to publicly

available bronchial transcriptional data, finding correlation ($\rho = 0.87$) between nasal and

bronchial transcripts, as well as correlation ($\rho=0.77$) between nasal differential expression and

previously observed bronchial differential expression in asthmatics [15]. To the best of our

knowledge, our study has generated the largest nasal RNAseq data set in asthma to date and is

the first to identify a nasal brush-based classifier of asthma.

Although based on only 90 genes, our asthma classifier classified asthma with greater

accuracy than models based on all genes, all differentially expressed genes, and known asthma

genes (**Figure 4**). Its superior performance supports that our machine learning pipeline

successfully selected a parsimonious set of informative genes that (1) captures more actionable

knowledge than traditional differential expression and genetic association analyses, and (2) cuts

through the potential noise of genes irrelevant to asthma. These results illustrate that data-

driven methods can build more effective classifiers than those built exclusively on current

domain knowledge. This is likely true not just for asthma but for other phenotypes as well. About

half the genes in our asthma classifier were not differentially expressed at FDR ≤ 0.05, and as

such would not have been examined with greater interest had we only performed traditional

differential expression analysis, which is the main analytic approach of virtually all studies of

gene expression in asthma [12-15, 39, 40]. Consistent with motivations underlying systems

biology and genomic approaches [25, 41], our study demonstrated that the asthma classifier

captures signal from differential expression as well as genes below traditional significance

thresholds that may still have a contributory role to asthma classification.

With prospective validation in large cohorts, our asthma classifier could lead to the

development of a minimally invasive biomarker to aid asthma diagnosis at clinical frontlines,

where time and resources often preclude pulmonary function testing (PFT). Nasal brushing can

be performed quickly, does not require machinery for collection, and implementation of our

classification model yields a straightforward, binary result of asthma or no asthma. Because it

takes seconds for nasal brushing and bioinformatic interpretation could be automated, an

asthma classifier such as ours may be attractive to time-strapped clinicians, particularly primary

care providers at the frontlines of asthma diagnosis. Asthma is frequently diagnosed and treated

in the primary care setting [42] where access to PFTs is often not immediately available. Gene

expression-based diagnostic classifiers are being successfully used in other disease areas, with

prominent examples including the commercially available MammaPrint [43] and Oncotype DX

[44] for diagnosing breast cancer phenotypes, leading to better outcomes. These examples from

the cancer field demonstrate an existing path for moving a classifier such as ours to clinical use.

We recognize that our asthma classifier did not perform quite as well in the microarray-

based vs. RNAseq-based asthma test sets, which was to be expected due to differences in

study design and technological factors between RNAseq and microarray profiling. First, the

baseline characteristics and phenotyping of the subjects differed. Subjects in the RNAseq test

set were adults who were classified as mild/moderate asthmatic or healthy using the same strict

criteria as the development set, which required subjects with asthma to have an objective

measure of obstructive airway disease (i.e. positive methacholine challenge response). In

contrast, subjects in the Asthma1 microarray test set were all children (i.e. not adults) with nasal

pathology, as entry criteria included dust mite allergic rhinitis specifically [31] (**Supplementary**

Table 3). Subjects from the Asthma2 cohort were adults who were classified as having asthma

or healthy based on history. As mentioned, the diagnosis of asthma based on history alone

without objective lung function testing can be inaccurate [45]. The phenotypic differences

between these test sets alone could explain differences in performance of our asthma classifier

in these test sets. Second, the differential performance may be due to the difference in profiling

approach. Gene mappings do not perfectly correspond between RNAseq and microarray due to

disparities between array annotations and RNAseq gene models [34]. Compared to

microarrays, RNAseq quantifies more RNA species and captures a wider range of signal [39].

Prior studies have shown that microarray-derived models can reliably predict phenotypes based

on samples' RNAseq profiles, but the converse does not often hold [34]. Despite the above

limitations, our asthma classifier performed with reasonable accuracy in classifying asthma in

these independent microarray-based test sets. These results support a degree of

generalizability of our classifier to asthma populations that may be phenotyped or profiled

differently.

An effective clinical classifier should have good positive and negative predictive value

[46]. This was indeed the case with our asthma classifier, which achieved high positive and

negative predictive values of 1.00 and 0.96 respectively in the RNAseq test set. We also tested

our asthma classifier on independent tests sets of subjects with allergic rhinitis, upper

respiratory infection, cystic fibrosis, and smoking, and showed that the classifier had a low to

zero rate of misclassifying other respiratory conditions as asthma (**Figure 5**). These results

were particularly notable for allergic rhinitis, a predominantly nasal condition. Although our

classifier is based on nasal gene expression, and asthma and allergic rhinitis frequently co-

occur [21], our classifier did not misclassify allergic rhinitis as asthma. Although these

conclusions are based on relatively small test sets due to the scarcity of nasal gene expression

data in the public domain, the performance of our classifier gives hope that it has the potential to

be generalizable and specific in much larger cohorts as well.

Although we have generated one of the largest nasal RNAseq data set in asthma to

date, a future direction of this study is to recruit additional cohorts for nasal gene expression

profiling and extend validation of our findings in a prospective manner. We recognize that our

development set was from a single center and its baseline characteristics do not characterize all

populations. However, we find it reassuring that the classifier performed reasonably well in

multiple external data sets spanning children and adults of varied racial distributions, and with

asthma and other respiratory conditions defined by heterogeneous criteria. Subjects with

asthma in our development cohort were not all symptomatic at the time of sampling. The fact

that the performance of our asthma classifier does not rely on symptomatic asthma is a

strength, as many mild/moderate asthmatics are only sporadically symptomatic given the

fluctuating nature of the disease.

In summary, we demonstrated an innovative application of RNA sequencing and

machine learning to identify a classifier consisting of genes expressed in nasal brushings that

accurately classifies subjects with mild/moderate asthma from controls. This asthma classifier

performed with accuracy across independent and external test sets, indicating reasonable

generalizability across study populations and gene expression profiling modality, as well as

specificity to asthma. This asthma classifier could potentially lead to the development of a nasal

biomarker of asthma.

Materials and Methods

Study design and subjects

Subjects with mild/moderate asthma were a subset of participants of the Childhood

Asthma Management Program (CAMP), a multicenter North American study of 1041 subjects

with mild to moderate persistent asthma [19, 20]. Findings from the CAMP cohort have defined

current practice and guidelines for asthma care and research [20]. Asthma was defined by

symptoms $\geq$2 times per week, use of an inhaled bronchodilator $\geq$ twice weekly or use of daily

medication for asthma, and increased airway responsiveness to methacholine ($PC_{20} \leq 12.5$

mg/ml). The subset of subjects included in this study were CAMP participants who presented for

a visit between July 2011 and June 2012 at Brigham and Women's Hospital (Boston, MA), one

of the eight study centers for CAMP.

Subjects with "no asthma" were recruited during the same time period by advertisement

at Brigham & Women's Hospital. Selection criteria were no personal history of asthma, no family

history of asthma in first-degree relatives, and self-described Caucasian ethnicity. Participation

was limited to Caucasian individuals because a concurrent independent study was planned that

would compare these same subjects to 968 Caucasian CAMP subjects who participated in the

CAMP Genetics Ancillary study [47]. Subjects underwent pre- and post-bronchodilator

spirometry according to American Thoracic Society guidelines. Only those meeting selection

criteria and with demonstrated normal lung function without bronchodilator response were

considered to have "no asthma."

Nasal brushing and RNA sequencing

Nasal brushing was performed with a cytology brush. Brushes were immediately placed

in RNALater (ThermoFisher Scientific, Waltham, MA) and then stored at 40^0C until RNA

extraction. RNA extraction was performed with Qiagen RNeasy Mini Kit (Valencia, CA).

Samples were assessed for yield and quality using the 2100 Bioanalyzer (Agilent Technologies,

Santa Clara, CA) and Qubit fluorometry (Thermo Fisher Scientific, Grand Island, NY).

Of the 190 subjects who underwent nasal brushing (66 with mild/moderate asthma, 124

with no asthma), a random selection of 150 subjects were *a priori* assigned as the development

set (for classification model development), with the 40 remaining subjects earmarked to serve

as a test set of independent subjects (for testing the classification model). To minimize potential

batch effects, all samples were submitted together for RNA sequencing (RNAseq). Staff at the

Mount Sinai genomics core were blinded to the assignment of samples as development or test

set. The sequencing library was prepared with the standard TruSeq RNA Sample Prep Kit v2

protocol (Illumina). The mRNA libraries were sequenced on the Illumina HiSeq 2500 platform

with a per-sample target of 40-50 million 100 bp paired-end reads. The data were put through

Mount Sinai's standard mapping pipeline[48] (using Bowtie [49] and TopHat [50], and

assembled into gene- and transcription-level summaries using Cufflinks [51]). Mapped data

were subjected to quality control with FastQC and RNA-SeQC [52]. Data were pre-processed

separately for the development and test sets to avoid leakage of information across the two data

sets and maintain fairness of the machine learning procedures as much as possible. Genes with

fewer than 100 counts in at least half the samples were dropped to reduce the potentially

adverse effects of noise. DESeq2 [23] was used to normalize the data sets using its variance

stabilizing transformation method.

VariancePartition Analysis of Potential Confounders

Given differences in age, race, and sex distributions between the asthma and "no

asthma" classes, we used the variancePartition method [22] to assess the degree to which

these variables influenced gene expression and potentially confounded the target phenotype

(asthma status). The total variance in gene expression was partitioned into the variance

attributable to age, race, and sex using a linear mixed model implemented in variancePartition

v1.0.0 [22]. Age (continuous variable) was modeled as a fixed effect, while race and sex

(categorical variables) were modeled as random effects. The results showed that age, race, and

sex accounted for minimal contributions to total gene expression variance (**Supplementary**

Figure 1). Downstream analyses were therefore performed with gene expression data

unadjusted for these variables.

Differential gene expression and pathway enrichment analysis

DESeq2 [23] was used to identify differentially expressed genes in the development set.

Genes with FDR ≤ 0.05 were deemed differentially expressed, with fold change <1 implying

under-expression and vice versa. To identify the functions underlying these genes, pathway

enrichment analysis was performed using the Gene Set Enrichment Analysis method applied to

the Molecular Signature Database (MSigDB) [24].

Identification of the Asthma Classifier by Machine Learning Analyses of the RNAseq

Development Set

To identify gene expression-based classifiers that predict asthma status, we applied a

rigorous machine learning pipeline that combined feature (gene) selection [16], classification

[17], and statistical analyses of classification performance [18] to the development set

(**Supplementary Figure 2**). The pipeline was implemented in Python using the scikit-learn

package [53]. Feature selection and classification were applied to a training set comprised of

120 randomly selected samples from the development set (n=150) as described below. For an

independent evaluation of the candidate classifiers generated from the training set by this

process, they were then evaluated on the remaining 30 samples (holdout set). Finally, to reduce

the dependence of the finally chosen classifier on a specific training-holdout split, this process

was repeated 100 times on 100 random splits of the development set into training and holdout

sets. The details of the overall process as well as the individual components are as follows.

Feature selection: The purpose of the feature selection component was to identify

subsets of the full set of genes in the development set, whose expression profiles could be used

to predict the asthma status as accurately as possible. The two main computations constituting

this component were (i) the optimal number of features that should be selected, and (ii) the

identification of this number of genes from the full gene set. To reduce the likelihood of

overfitting when conducting both these computations on the entire training set, we used a 5x5

nested (outer and inner) cross-validation (CV) setup [26] for selecting features from the training

set (**Supplementary Figure 3**). The inner CV round was used to determine the optimal number

of genes to be selected, and the outer CV round was used to select the set of predictive genes

based on this number, thus separating the samples on which these decisions are made. The

supervised Recursive Feature Elimination (RFE) algorithm [54] was executed on the inner CV

training split to determine the optimal number of features. The use of RFE within this setting

enabled us to identify groups of features that are collectively, but not necessarily individually,

predictive. Specifically, we used the L2-regularized Logistic Regression (LR or Logistic) [55] and

SVM-Linear (kernel) [56] classification algorithms in conjunction with RFE (combinations

henceforth referred to as LR-RFE and SVM-RFE respectively). For this, for a given inner CV

training split, all the features (genes) were ranked using the absolute values of the weights

assigned to them by an inner classification model, trained using the LR or SVM algorithm, over

this split. Next, for each of the conjunctions, the set of top-k ranked features, with k starting with

11587 (all filtered genes) and being reduced by 10% in each iteration until k=1, was considered.

The discriminative strength of feature sets consisting of the top k features as per this ranking

was assessed by evaluating the performance of the LR or SVM classifier based on them over all

the inner CV training-test splits. The optimal number of features to be selected was determined

as the value of k that produces the best performance. Next, a ranking of features was derived

from the outer CV training split using exactly the same procedure as applied to the inner CV

training split. The optimal number of features determined above was selected from the top of

this ranking to determine the optimal set of predictive features for this outer CV training split.

Executing this process over all the five outer CV training splits created from the development set

identified five such sets. Finally, the set of features (genes) that was common to all these sets

(i.e. in their intersection/overlap), which is expected to yield a more robust feature set than the

individual outer CV splits, was selected as the predictive gene set for this training set. One such

set was identified for each application of LR-RFE and SVM-RFE to the training set.

517   *Classification analyses*: Once predictive gene sets had been selected from feature

selection, four global classification algorithms (L2-regularized Logistic Regression (LR or

Logistic) [55], SVM-Linear [56], AdaBoost [57], and Random Forest (RF) [58]) were used to

learn *intermediate classification models* over the training set. These intermediate models were

then applied to the corresponding holdout set to generate probabilistic asthma predictions for

the samples. An optimal threshold for converting these probabilistic predictions into binary ones

(higher than threshold=asthma, lower than threshold=no asthma) was then computed as the

threshold that yielded the highest classification performance on the holdout set. This

optimization resulted in the *proposed classification models*.

526   *Statistical analyses of classification performance*: After the above components were run

on 100 training-holdout splits of the development set, we obtained 100 proposed classification

models for each of eight feature selection-global classification combinations (two feature

selection algorithms (LR-RFE and SVM-RFE) and four global classification algorithms Logistic,

SVM-Linear, AdaBoost and RF). The next step of our pipeline was to determine the best

performing combination. Instead of making this determination just based on the highest

evaluation score, as is typically done in machine learning studies, we utilized this large

population of models and their optimized holdout evaluation scores to conduct a statistical

comparison to make this determination. Specifically, we applied the Friedman test followed by

the Nemenyi test [18, 59] to this population of modules and their evaluation scores. These tests,

which account for multiple hypothesis testing, assessed the statistical significance of the relative

difference of performance of the combinations in terms of their relative ranks across the 100

splits.

Optimization for parsimony: For a phenotype classifier, it is essential to consider

parsimony in model selection (i.e. minimize number of features (i.e. genes)) to enhance its

clinical utility and acceptability. To enforce this for our asthma classifier, an adapted

performance measure, defined as the absolute performance measure (F-measure) divided by

the number of genes in that model, was used for the above statistical comparison, i.e. as input

to the Friedman-Nemenyi tests. In terms of this measure, a model that does not obtain the best

performance measure among all models, but uses much fewer genes than the others, may be

judged to be the best model. The result of the statistical comparison using this adapted measure

was visualized as a Critical Difference plot [18] (**Supplementary Figure 4**), and enabled us to

identify the best combination of feature selection and classification method as the left-most entry

in this plot.

Final model development: The final step in our pipeline was to determine the

representative model out of the 100 learned the above best combination by finding which of

these models yielded the highest evaluation measure (F-measure; **Box 1** and **Supplementary**

Figure 5). In case of ties among multiple candidates, the gene set that produced the best

average asthma classification F-measure across all four global classification algorithms was

chosen as the gene set constituting the representative model for that combination. This analysis

yielded the representative gene set, global classification algorithm, and the optimized asthma

classification threshold. The asthma classifier was built by training the global classification

algorithm to the expression profiles of the representative gene set, and using the optimized

threshold for classifying samples positive/negative for asthma.

Evaluation of the Asthma Classifier in an RNAseq test set of independent subjects

The asthma classifier identified by our machine learning pipeline was then tested on the

RNAseq test set (n=40) to assess its performance in independent subjects. F-measure was

used as the primary measure for classification performance, as described in **Box 1** and

Supplementary Figure 5. AUC, PPV and NPV were additionally calculated for context.

Comparison of Performance to Alternative Classification Models

Although our classifier was identified using a rigorous machine learning methodology,

the pipeline explored several other models from all combinations of feature selection and global

classification methods. Thus, we compared the performance of our classifier with all these other

possible classifiers.

Also, our methodology was not the only way to develop gene expression-based

classifiers. Thus, we also compared the classifier's performance with several other valid

methods by applying our machine learning pipeline with the feature (gene) selection step

replaced with the following alternatively determined gene sets: (1) all filtered RNAseq genes, (2)

all differentially expressed genes, and (3) known asthma genes from a recent review of asthma

genetics [29]. To maintain consistency with the machine learning pipeline-derived models, each

of these gene sets was run through the same pipeline (**Supplementary Figure 2** with the

feature selection component turned off) to identify the best performing global classification

algorithm and the optimal asthma classification threshold for this predetermined set of features.

The algorithm and threshold were used to train each of these gene sets' representative

classification model over the entire development set, and the resulting models for each of these

gene sets were then evaluated on the RNAseq test set. Finally, as a baseline representative of

alternative sparse classification algorithms, which represent a one-step option for doing feature

selection and classification simultaneously, we also trained an L1-regularized logistic regression

model (L1-Logistic) [30] on the development set and evaluated it on the RNAseq test set.

Comparison of Performance to Permutation-based Random Models

To determine the extent to which the performance of all the above classification models

could have been due to chance, we compared their performance with that of their random

counterpart models (**Supplementary Figures 6 and 7**). These counterparts were obtained by

randomly permuting the labels of the samples in the development set and executing each of the

above model training procedures on these randomized data sets in the same way as for the real

development set. These random models were then applied to each of the test sets considered

in our study, and their performances were also evaluated in terms of the same measures. For

each of real models tested in our study, 100 corresponding random models were learned and

evaluated as above, and the performance of the real models was compared with the average

performance of the corresponding random models.

Evaluation of the asthma classifier in external independent asthma cohorts

To assess the generalizability of the asthma classifier to other populations, microarray-

profiled data sets of nasal gene expression from two external asthma cohorts-- Asthma1

(GSE19187) [31] and Asthma2 (GSE46171) [32] (**Supplementary Table 3**)-- were obtained

from NCBI Gene Expression Omnibus (GEO) [60]. For each of these data sets, we obtained

their probe-level normalized versions from GEO, and then obtained gene-level expression

profiles by averaging the normalized expression of all the probes corresponding to the same

gene. The probe-to-gene mappings were obtained from the microarray platform (GPL) files also

available from GEO. The asthma classifier was then applied to these gene-level data sets and

its performance evaluated on these external asthma cohorts.

Evaluation of the asthma classifier in external cohorts with other respiratory conditions

To assess the classifier's ability to distinguish asthma from respiratory conditions that

can have overlapping symptoms with asthma, i.e. its specificity to asthma, microarray-profiled

data sets of nasal gene expression were also obtained for five external cohorts with allergic

rhinitis (GSE43523) [35], upper respiratory infection (GSE46171) [32], cystic fibrosis

(GSE40445) [36], and smoking (GSE8987) [12] (**Supplementary Table 4**). Gene-level

expression data sets were obtained for these cohorts using the same methodology as described

above. The asthma classifier was then applied to these data sets, and its performance

evaluated on these external cohorts with non-asthma respiratory conditions.

Data availability

Data and code for this study (doi:10.7303/syn9878922) will be made publically available upon

publication via Synapse, a software platform for open, reproducible data-driven science, at

https://www.synapse.org/#!Synapse:syn9878922/files/

Declarations

Ethics approval and consent to participate: The institutional review boards of Brigham &

Women's Hospital and the Icahn School of Medicine at Mount Sinai approved the study

protocols. Written informed consent was obtained from all subjects.

Funding: This study was supported by the US National Institutes of Health (NIH R01AI118833,

K08AI093538, R01GM114434) and the Icahn Institute for Genomics and Multiscale Biology,

including computational resources provided by Scientific Computing at the Icahn School of

Medicine at Mount Sinai.

Author contributions: SB directed the study. SB and AJR directed the recruitment of subjects

and sample collection. BAR and STW provided guidance for access to subjects. EES advised

on sequencing strategy. SB curated the clinical data. SB, GP, EES, OPP and MEA designed

and performed the statistical and computational analyses. SB and GP wrote the manuscript. SB,

GP, OPP, AJR, MEA, GEH, BAR, STW, and EES edited the manuscript. All authors contributed

significantly to the work presented in this paper.

Competing interests: SB, GP, and EES have filed a patent application related to the findings of

this manuscript. The remaining authors declare that they have no competing interests.

Acknowledgments: We thank Kathryn Paul, Laura Ting, Anne Plunkett, Nancy Madden, Ann

Fuhlbrigge, Kelan Tantisira, Dan Cossette, Aimee Garciano, and Roxanne Kelly for their

assistance and support with recruitment, specimen collection, and sample processing. We thank

Robert Griffin and Ana Stanescu for critically reviewing the paper.

References

1. **Current Asthma Prevalence Percents by Age, Sex, and Race/Ethnicity, United States, 2015. Asthma Surveillance Data.** *National Health Interview Survey, National Center for Health Statistics, Centers for Disease Control and Prevention wwwcdcgov/asthma/asthmadatahtm, downloaded 6/12/2017*.

2. Yeatts K, Shy C, Sotir M, Music S, Herget C: **Health consequences for children with undiagnosed asthma-like symptoms.** *Arch Pediatr Adolesc Med* 2003, **157:**540-544.

3. Fanta CH: **Asthma.** *N Engl J Med* 2009, **360:**1002-1014.

4. Stempel DA, Spahn JD, Stanford RH, Rosenzweig JR, McLaughlin TP: **The economic impact of children dispensed asthma medications without an asthma diagnosis.** *J Pediatr* 2006, **148:**819-823.

5. Szefler SJ, Wenzel S, Brown R, Erzurum SC, Fahy JV, Hamilton RG, Hunt JF, Kita H, Liu AH, Panettieri Jr RA, et al: **Asthma outcomes: Biomarkers.** *Journal of Allergy and Clinical Immunology* 2012, **129:**S9-S23.

6. Reddel HK, Bateman ED, Becker A, Boulet LP, Cruz AA, Drazen JM, Haahtela T, Hurd SS, Inoue H, de Jongste JC, et al: **A summary of the new GINA strategy: a roadmap to asthma control.** *Eur Respir J* 2015, **46:**622-639.

7. **Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma.** Washington DC: National Heart Lung and Blood Institute and National Asthma Education and Prevention Program; 2007.

8. Gershon AS, Victor JC, Guan J, Aaron SD, To T: **Pulmonary function testing in the diagnosis of asthma: a population study.** *Chest* 2012, **141:**1190-1196.

9. Sokol KC, Sharma G, Lin YL, Goldblum RM: **Choosing wisely: adherence by physicians to recommended use of spirometry in the diagnosis and management of adult asthma.** *Am J Med* 2015, **128:**502-508.

10.   Petsky HL, Cates CJ, Lasserson TJ, Li AM, Turner C, Kynaston JA, Chang AB: **A systematic review and meta-analysis: tailoring asthma treatment on eosinophilic markers (exhaled nitric oxide or sputum eosinophils).** *Thorax* 2012, **67:**199-208.

11.   van Schayck CP, van Der Heijden FM, van Den Boom G, Tirimanna PR, van Herwaarden CL: **Underdiagnosis of asthma: is the doctor or the patient to blame? The DIMCA project.** *Thorax* 2000, **55:**562-565.

12.   Sridhar S, Schembri F, Zeskind J, Shah V, Gustafson AM, Steiling K, Liu G, Dumas YM, Zhang X, Brody JS, et al: **Smoking-induced gene expression changes in the bronchial airway are reflected in nasal and buccal epithelium.** *BMC Genomics* 2008, **9:**259.

13.   Wagener AH, Zwinderman AH, Luiten S, Fokkens WJ, Bel EH, Sterk PJ, van Drunen CM: **The impact of allergic rhinitis and asthma on human nasal and bronchial epithelial gene expression.** *PLoS One* 2013, **8:**e80257.

14.   Guajardo JR, Schleifer KW, Daines MO, Ruddy RM, Aronow BJ, Wills-Karp M, Hershey GK: **Altered gene expression profiles in nasal respiratory epithelium reflect stable versus acute childhood asthma.** *J Allergy Clin Immunol* 2005, **115:**243-251.

15.   Poole A, Urbanek C, Eng C, Schageman J, Jacobson S, O'Connor BP, Galanter JM, Gignoux CR, Roth LA, Kumar R, et al: **Dissecting childhood asthma with nasal transcriptomics distinguishes subphenotypes of disease.** *J Allergy Clin Immunol* 2014, **133:**670-678 e612.

16.   Saeys Y, Inza I, Larranaga P: **A review of feature selection techniques in bioinformatics.** *Bioinformatics* 2007, **23:**2507-2517.

17.   Witten IH, Frank E, Hall MA: *Data mining : practical machine learning tools and techniques.* 3rd edn. Burlington, MA: Morgan Kaufmann; 2011.

18.   Demsar J: **Statistical Comparisons of Classifiers over Multiple Data Sets.** *J Mach Learn Res* 2006, **7:**1-30.

19. **The Childhood Asthma Management Program (CAMP): design, rationale, and methods. Childhood Asthma Management Program Research Group.** *Control Clin Trials* 1999, **20:**91-120.

20. Covar RA, Fuhlbrigge AL, Williams P, Kelly HW, the Childhood Asthma Management Program Research G: **The Childhood Asthma Management Program (CAMP): Contributions to the Understanding of Therapy and the Natural History of Childhood Asthma.** *Curr Respir Care Rep* 2012, **1:**243-250.

21. Egan M, Bunyavanich S: **Allergic rhinitis: the "Ghost Diagnosis" in patients with asthma.** *Asthma Research and Practie* 2015, **1:**DOI: 10.1186/s40733-40015-40008-40730.

22. Hoffman GE, Schadt EE: **variancePartition: Quantifying and interpreting drivers of variation in complex gene expression studies.** *BMC Bioinformatics* 2016, **17:**483.

23. Love MI, Huber W, Anders S: **Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.** *Genome Biol* 2014, **15:**550.

24. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP: **Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles.** *Proc Natl Acad Sci U S A* 2005, **102:**15545-15550.

25. Schadt EE, Friend SH, Shaywitz DA: **A network view of disease and compound screening.** *Nat Rev Drug Discov* 2009, **8:**286-295.

26. Whalen S, Pandey OP, Pandey G: **Predicting protein function and other biomedical characteristics with heterogeneous ensembles.** *Methods* 2016, **93:**92-102.

27. Lever J, Krzywinski M, Altman N: **Points of Significance: Classification Evaluation.** *Nature Methods* 2016, **13:**603-604.

28. Saito T, Rehmsmeier M: **The precision-recall plot is more informative than the ROC plot when evaluating binary classifiers on imbalanced datasets.** *PLoS One* 2015, **10:**e0118432.

29. Mathias RA: **Introduction to genetics and genomics in asthma: genetics of asthma.** *Adv Exp Med Biol* 2014, **795:**125-155.

30. Vidaurre D, Bielza C, Larrañaga P: **A Survey of L1 Regression.** *International Statistical Review* 2013, **81:**361-387.

31. Giovannini-Chami L, Marcet B, Moreilhon C, Chevalier B, Illie MI, Lebrigand K, Robbe-Sermesant K, Bourrier T, Michiels JF, Mari B, et al: **Distinct epithelial gene expression phenotypes in childhood respiratory allergy.** *Eur Respir J* 2012, **39:**1197-1205.

32. McErlean P, Berdnikovs S, Favoreto S, Jr., Shen J, Biyasheva A, Barbeau R, Eisley C, Barczak A, Ward T, Schleimer RP, et al: **Asthmatics with exacerbation during acute respiratory illness exhibit unique transcriptional signatures within the nasal mucosa.** *Genome Med* 2014, **6:**1.

33. Zhang W, Yu Y, Hertwig F, Thierry-Mieg J, Zhang W, Thierry-Mieg D, Wang J, Furlanello C, Devanarayan V, Cheng J, et al: **Comparison of RNA-seq and microarray-based models for clinical endpoint prediction.** *Genome Biol* 2015, **16:**133.

34. Su Z, Fang H, Hong H, Shi L, Zhang W, Zhang W, Zhang Y, Dong Z, Lancashire LJ, Bessarabova M, et al: **An investigation of biomarkers derived from legacy microarray data for their utility in the RNA-seq era.** *Genome Biol* 2014, **15:**523.

35. Imoto Y, Tokunaga T, Matsumoto Y, Hamada Y, Ono M, Yamada T, Ito Y, Arinami T, Okano M, Noguchi E, Fujieda S: **Cystatin SN upregulation in patients with seasonal allergic rhinitis.** *PLoS One* 2013, **8:**e67057.

36. Clarke LA, Sousa L, Barreto C, Amaral MD: **Changes in transcriptome of native nasal epithelium expressing F508del-CFTR and intersecting data from comparable studies.** *Respir Res* 2013, **14:**38.

37. Oliver BG, Robinson P, Peters M, Black J: **Viral infections and asthma: an inflammatory interface?** *Eur Respir J* 2014, **44:**1666-1681.

38. Cowling BJ, Chan KH, Fang VJ, Lau LL, So HC, Fung RO, Ma ES, Kwong AS, Chan CW, Tsui WW, et al: **Comparative epidemiology of pandemic and seasonal influenza A in households.** *N Engl J Med* 2010, **362:**2175-2184.

39. Bunyavanich S, Schadt EE: **Systems biology of asthma and allergic diseases: A multiscale approach.** *J Allergy Clin Immunol* 2014.

40. Sordillo J, Raby BA: **Gene expression profiling in asthma.** *Adv Exp Med Biol* 2014, **795:**157-181.

41. Libbrecht MW, Noble WS: **Machine learning applications in genetics and genomics.** *Nat Rev Genet* 2015, **16:**321-332.

42. Wechsler ME: **Managing asthma in primary care: putting new guideline recommendations into context.** *Mayo Clin Proc* 2009, **84:**707-717.

43. Cardoso F, van't Veer LJ, Bogaerts J, Slaets L, Viale G, Delaloge S, Pierga JY, Brain E, Causeret S, DeLorenzi M, et al: **70-Gene Signature as an Aid to Treatment Decisions in Early-Stage Breast Cancer.** *N Engl J Med* 2016, **375:**717-729.

44. Paik S, Shak S, Tang G, Kim C, Baker J, Cronin M, Baehner FL, Walker MG, Watson D, Park T, et al: **A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer.** *N Engl J Med* 2004, **351:**2817-2826.

45. Jain VV, Allison DR, Andrews S, Mejia J, Mills PK, Peterson MW: **Misdiagnosis Among Frequent Exacerbators of Clinically Diagnosed Asthma and COPD in Absence of Confirmation of Airflow Obstruction.** *Lung* 2015, **193:**505-512.

46. Brower V: **Biomarkers: Portents of malignancy.** *Nature* 2011, **471:**S19-21.

47. Himes BE, Hunninghake GM, Baurley JW, Rafaels NM, Sleiman P, Strachan DP, Wilk JB, Willis-Owen SA, Klanderman B, Lasky-Su J, et al: **Genome-wide association**

analysis identifies PDE4D as an asthma-susceptibility gene. *Am J Hum Genet* 2009, **84:**581-593.

48.   Fromer M, Roussos P, Sieberts SK, Johnson JS, Kavanagh DH, Perumal TM, Ruderfer DM, Oh EC, Topol A, Shah HR, et al: **Gene expression elucidates functional impact of polygenic risk for schizophrenia.** *Nat Neurosci* 2016.

49.   Langmead B, Trapnell C, Pop M, Salzberg SL: **Ultrafast and memory-efficient alignment of short DNA sequences to the human genome.** *Genome Biol* 2009, **10:**R25.

50.   Trapnell C, Pachter L, Salzberg SL: **TopHat: discovering splice junctions with RNA-Seq.** *Bioinformatics* 2009, **25:**1105-1111.

51.   Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ, Pachter L: **Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation.** *Nat Biotechnol* 2010, **28:**511-515.

52.   DeLuca DS, Levin JZ, Sivachenko A, Fennell T, Nazaire MD, Williams C, Reich M, Winckler W, Getz G: **RNA-SeQC: RNA-seq metrics for quality control and process optimization.** *Bioinformatics* 2012, **28:**1530-1532.

53.   Pedregosa F, Varoquaux Ge, Gramfort A, Michel V, Thirion B, others: **Scikit-learn: Machine Learning in Python.** *Journal of Machine Learning Research* 2011, **12:**2825-2830.

54.   Guyon I, Weston, J, Barnhill, S, Vapnik, V: **Gene selection for cancer classification using support vector machines.** *Machine Learning* 2002, **46:**389-422.

55.   Bewick V, Cheek L, Ball J: **Statistics review 14: Logistic regression.** *Crit Care* 2005, **9:**112-118.

56.   Burges CJ: **A tutorial on support vector machines for pattern recognition.** *Data mining and knowledge discovery* 1998, **2:**121-167.

57.   Freund Y, Schapire RE: **A Decision-Theoretic Generalization of On-Line Learning and an Application to Boosting.** *J Comput Syst Sci* 1997, **55:**119-139.

58.   Breiman L: **Random Forests.** *Machine Learning* 2001, **45:**5-32.

59.   Hollander M, Wolfe DA, Chicken E: *Nonparametric statistical methods.* John Wiley & Sons; 2013.

60.   Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, Marshall KA, Phillippy KH, Sherman PM, Holko M, et al: **NCBI GEO: archive for functional genomics data sets--update.** *Nucleic Acids Res* 2013, **41:**D991-995.

Figure Legends

Figure 1: Study flow for the identification of a nasal brush-based classifier of asthma by machine learning analysis of RNAseq data. One hundred and ninety subjects with mild/moderate asthma and controls without asthma were recruited for phenotyping, nasal brushing, and RNA sequencing of nasal brushings. The RNAseq data generated were then *a priori* split into development and test sets. The development set was used for differential expression analysis and machine learning (involving feature selection, classification, and statistical analyses of classification performance) to identify an asthma classifier that can classify asthma from no asthma as accurately as possible. The asthma classifier was then evaluated on eight test sets, including (1) the RNAseq test set of independent subjects with and without asthma, (2) two external test sets of subjects with and without asthma with nasal gene expression profiled by microarray, and (3) five external test sets of subjects with non-asthma respiratory conditions (allergic rhinitis, upper respiratory infection, cystic fibrosis, and smoking) and nasal gene expression profiled by microarray.

Figure 2: Receiver operating characteristic (ROC) curve of the predictions generated by applying the asthma classifier to the RNAseq test set of independent subjects (n=40). The ROC curve for a random model is shown for reference. The curve and its corresponding AUC score show that the classifier performs well for both asthma and no asthma (control) samples in this test set.

Figure 3: **Evaluation of the asthma classifier on test sets of independent subjects with asthma.** Performance of the asthma classifier in classifying asthma (A) and no asthma (C) in terms of F-measure, a conservative mean of precision and sensitivity. F-measure ranges from 0 to 1, with higher values indicating superior classification performance. The classifier was applied

to an RNAseq test set of independent subjects with and without asthma, two external microarray

data sets from subjects with and without asthma (Asthma1 and Asthma2), and combined data

from Asthma1 and Asthma2. Positive (B) and negative (D) predictive values are also provided

for context.

Figure 4: **Comparative performance of the asthma classifier and other classification**

models in the RNAseq test set. Performances of the asthma classifier and other classification

models in classifying asthma (left panel) and no asthma (right panel) are shown in terms of F-

measure, with individual measures shown in the bars. The number of genes in each model is

shown in parentheses within the bars. The asthma classifier is labeled in red and classification

models learned from the machine learning pipeline using other combinations of feature selection

and classification are labeled in black. These other classification models were combinations of

two feature selection algorithms (LR-RFE and SVM-RFE) and four global classification

algorithms (Logistic Regression, SVM-Linear, AdaBoost and Random Forest). For context,

alternative classification models (labeled in blue) are also shown and include: (1) a model

derived from an alternative, single-step classification approach (sparse classification model

learned using the L1-Logistic regression algorithm), and (2) models substituting feature

selection with each of 3 pre-selected gene sets (all genes after filtering, all differentially

expressed genes in the development set, and known asthma genes [29]) with their respective

best performing global classification algorithms. These results show the superior performance of

the asthma classifier compared to all other models, in terms of classification performance and

model parsimony (number of genes included). LR = Logistic Regression. SVM = Support Vector

Machine. RFE = Recursive Feature Elimination.

Figure 5: **Evaluation of the asthma classifier on test sets of independent subjects with**

non-asthma respiratory conditions. Performance statistics of the classifier when applied to

external microarray-generated data sets of nasal gene expression derived from case/control cohorts with non-asthma respiratory conditions. Performance is shown in terms of F-measure (A and C), a conservative mean of precision and sensitivity, as well as positive (B) and negative predictive values (D). The classifier had a low to zero rate of misclassifying other respiratory conditions as asthma, supporting that the classifier is specific to asthma and would not misclassify other respiratory conditions as asthma.

Figure 6: **Heatmap showing expression profiles of the 90 genes constituting the asthma classifier.** Columns shaded pink at the top denote asthma samples, while samples from subjects without asthma are denoted by columns shaded grey. 22 and 24 of these genes were over- and under-expressed in asthma samples (DESeq2 FDR ≤ 0.05), denoted by orange and purple groups of rows, respectively. The four genes in this set that have been previously associated with asthma [29] are marked in blue. The classifier's inclusion of genes not previously known to be associated with asthma as well as genes not differentially expressed in asthma (beige group of rows) demonstrates the ability of a machine learning methodology to move beyond traditional analyses of differential expression and current domain knowledge.

Table 1: Baseline characteristics of subjects in the RNAseq development and test sets

	RNAseq Development set			RNAseq Test Set			Development vs. Test Set P value[B]
	All (n=150)	Asthma (n=53)	No Asthma (n=97)	All (n=40)	Asthma (n=13)	No Asthma (n=27)	
Age: years	26.9 (5.4)	25.7 (2.0)	27.6 (6.5)	26.2 (5.1)	25.3 (2.1)	26.6 (6.1)	0.47
Sex: female	89 (59.3%)	24 (45.3%)	65 (67.0%)	21 (52.5%)	2 (15.3%)	19 (70.4%)	0.40
Race							0.60
Caucasian	116 (77.3%)	21 (40.4%)	96 (99.0%)	32 (80.0%)	5 (38.5%)	27 (100.0%)	
African American	24 (16.0%)	23 (43.4%)	1 (1.0%)	5 (12.5%)	5 (38.5%)	0 (0.0%)	
Latino	5 (3.3%)	5 (9.4%)	0 (0.0%)	3 (7.5%)	3 (23.1%)	0 (0.0%)	
Other	5 (3.3%)	4 (7.5%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
FEV1[A]: % predicted	94.7 (10.0)	94.6% (10.9)	94.8 (9.7)	94.5 (11.4)	94.4 (12.0)	94.6 (11.3)	0.90
FEV1/FVC[A]: %	82.5 (6.4)	81.5 (6.7)	83.1 (6.3)	82.7 (5.5)	84.8 (4.4)	81.6 (5.8)	0.91
Bronchodilator response: %	5.6 (6.0)	8.7 (6.4)	3.9 (5.1)	4.5 (5.4)	7.0 (6.1)	3.3 (4.7)	0.29
Age asthma onset: years		3.2 (2.7)	n/a		3.4 (2.0)		0.78
Allergic rhinitis	60 (40.0%)	29 (54.7%)	31 (32.0%)	7 (17.5%)	7 (53.8%)	0 (0.0%)	0.009
Nasal steroids	14 (9.3%)	9 (17.0%)	5 (5.2%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0.07
Smoking	7 (4.7%)	1 (1.9%)	6 (6.2%)	1 (2.5%)	0 (0.0%)	1 (3.7%)	1.0

Mean (SD) or Number (%) provided

[A]pre-bronchodilator measures. FEV1 = forced expiratory flow volume in 1 second, FVC = forced vital capacity

[B]Fisher's Exact test for categorical variables and t-test for continuous variables

Supplementary Materials

Supplementary Figure 1: variancePartition analysis of the RNAseq development set.

Supplementary Figure 2: Visual description of the machine learning pipeline used to select predictive features (genes) and develop classification models based on them in the RNAseq development set.

Supplementary Figure 3: Visual description of the feature (gene) selection component of the machine learning pipeline.

Supplementary Figure 4: Critical Difference plots demonstrating results of the statistical comparison of the performance of 100 asthma classification models obtained by various combinations of feature selection and global classification algorithms in terms of the classification performance and parsimony (numbers of genes included) of the models.

Supplementary Figure 5: Evaluation measures for classification models[27].

Supplementary Figure 6: Performance of permutation-based random classification models in test sets of independent subjects with asthma and controls.

Supplementary Figure 7: Performance of permutation-based random classification models in test sets of independent subjects with non-asthma respiratory conditions and controls.

Supplementary Figure 8: Distribution of DESeq2 FDR values of differential expression in the asthma classifier (blue bars) vs. other genes in the RNAseq development set (coral bars).

Supplementary Table 1: Lists of over- and under-expressed genes and pathways in asthma cases compared to controls (in different tabs of this file). Differentially expressed genes were identified using DESeq2 [23] applied to the development set, and enriched pathways were identified from the Molecular Signature Database [24], both using an upper FDR threshold of 0.05.

Supplementary Table 2: List of known asthma-associated genes from a recent review of asthma genetics [29] that overlap with genes in our RNAseq data sets.

Supplementary Table 3: Characteristics of the external asthma cohorts used in the evaluation of the asthma classifier.

Supplementary Table 4: Characteristics of the external cohorts with non-asthma respiratory conditions and controls used in the evaluation of the asthma classifier.

Figure 5

Pandey et al.

Figure 6

**Color Key
and Histogram**

Count

3000

-5   0   5

Row Z-Score

ANKRD9
C3
CDHR3
CHCHD10
CROCC
CROCCP2
DLEC1
DNAI1
GNAL
LOC388564
LPHN1
MS4A8B
NWD1
PCP4L1
PCSK6
SDK1
SEMA5A
SERPINB11
SERPINF1
TEKT2
TMEM190
TXNDC5
C12orf36
CLEC2B
CLEC7A
DEFB1
EPPK1
ESR1
LPAR6
LTBP1
LY6D
NMU
PTER
RPH3AL
SCD
SCGB1A1
SERPINE2
SLC44A5
SLC5A3
SORD
SPRR1A
SYNM
TFPI
TM4SF1
XIST
ZNF117
ALOX15B
MMP9
IER3
RHCG
TPSAB1
CDH26
NCRNA00152
SLC35E2
C6orf105
CEBPD
CRABP2
LOC286002
ALDH1A3
ARL4D
NR1D1
SC4MOL
SLC5A8
CPAMD8
CYFIP2
GSTT1
TPSB2
DUSP1
S100A7
S100A8
AKR1B15
LOC90784
SPRR2D
PTAFR
EPDR1
CYP1B1
SLC26A4
HLA.C
CPA3
FOS
IGF1
SLC9A3
CDKN1A
ERAP2
SERPINB3
FCGBP
KRT24
CD177
B3GNT6
HIPK2