Gene Regulatory Divergence Between Locally Adapted Ecotypes in Their Native Habitats

Billie A. Gould[a], Yani Chen[a], David B. Lowry[a,b,c,*]

[a]Department of Plant Biology, Michigan State University, East Lansing, MI, USA

[b]Program in Ecology, Evolutionary Biology and Behavior, Michigan State University, East

Lansing, MI 48824, USA

[c]Plant Resilience Institute, Michigan State University, East Lansing, MI 48824, USA

Key words: *Mimulus*, local adaptation, allele-specific expression, ecotype, heterosis, salt

tolerance, drought, chromosome inversion

[*]To whom correspondence should be addressed: David B. Lowry

Department of Plant Biology, Michigan State University, Plant Biology Laboratories, 612

Wilson Road, Room 166, East Lansing MI 48824, USA

Tel: (517) 432-4883

E-mail: dlowry@msu.edu

Running title: Gene Expression Divergence in Nature

ABSTRACT

Local adaptation is a key driver of ecological specialization and the formation of new species. Despite its importance, the evolution of gene regulatory divergence among locally-adapted populations is poorly understood, especially how that divergence manifests in nature. Here, we evaluate gene expression divergence and allele-specific gene expression responses for locally-adapted coastal perennial and inland annual accessions of the yellow monkeyflower, *Mimulus guttatus*, in a field reciprocal transplant experiment. Overall, 6765 (73%) of surveyed genes were differentially expressed between coastal and inland habitats, while 7213 (77%) were differentially expressed between the coastal perennial and inland annual accessions. Further, 18% of transcripts had significant genotype x site (GxE) effects. Habitat-specific differential expression was found for 62% of the GxE transcripts (differential expression in one habitat, but not the other), while only 94 (~5%) GxE transcripts had crossing reaction norms. *Cis*-regulatory variation was pervasive, affecting 79% (5532) of differentially expressed genes. We detected *trans* effects for 52% (3611) of differentially expressed genes. Consistent with the supergene hypothesis of chromosome inversion evolution, a locally adaptive inversion was enriched for *cis*-regulatory divergence. Above and below mid-parent heterosis for gene expression was prevalent across habitats and heterotic genes were enriched for growth and development gene ontologies. These results provide multiple new insights into the evolution of transcriptome-wide gene regulatory divergence and plasticity among locally adapted populations.

INTRODUCTION

One of the most important forces driving the evolution ecological specialization and the origin of biodiversity is local adaptation (Clausen 1951; Colosimo et al. 2005; Sobel et al. 2010). Local adaptation is characterized by reciprocal home-site advantage, whereby populations perform best in their home habitat while performing poorly in foreign habitats (Kawecki & Ebert 2004; Hoekstra et al. 2006; Hereford 2009; Wadgymar et al. 2017). Case-studies, such as the evolution of beak size in Darwin's Finches (Abzhanov et al. 2004) and coat color of Beach Mice (Hoekstra et al. 2006), suggest that local adaptation should involve fitness trade-offs at the level of individual genes. However, studies that have combined reciprocal transplant field experiments with quantitative trait locus (QTL) mapping have observed that the majority of loci which confer substantial fitness benefits in one habitat result in little or no cost in alternative habitats (Verhoeven et al. 2008; Gardner & Latta 2006; Lowry et al. 2009; Lowry & Willis 2010; Anderson et al. 2011, 2013; Leinonen et al. 2013; Agren et al. 2013; Wadgymar et al. 2017). While multiple studies have now evaluated the effects of individual loci on fitness across habitats, the patterns gene regulatory divergence that might explain habitat-dependent fitness effects in those systems have not been evaluated under natural field conditions.

One very promising route for understanding how gene expression regulation evolves between locally adaptive populations is to conduct allele specific expression (ASE) analyses in nature (Tung et al. 2009; Lovell et al. 2016; Wang et al. 2017; Lohman et al. 2017). In contrast to coarse QTL mapping, conducting ASE analyses in the field allows for the evaluation of the patterns of gene expression responses to habitat context at the individual allele level. ASE analysis is often conducted in F1 hybrids, where parental alleles are evaluated for significant

differences in expression (de Meaux et al. 2005; Guo et al. 2008; Zhang & Borevitz 2009;

Cubillos et al. 2014; Coolon et al. 2014; Steige et al. 2015, 2017; Lovell et al. 2016; Aguilar-

Rangel et al. 2017). ASE relies on there being nucleotide polymorphisms between parental

alleles, so that those alleles can be individually counted in the F1 transcript pool. *Cis*-regulatory

allelic expression differences are expected to persist regardless of whether alleles are expressed

separately in parental genotypes or together in an F1 hybrid. In this way, ASE allows for the

identification of *cis* expression Quantitative Trait Loci (eQTL), which can be evaluated for their

responses to different habitats. Further, *trans* effects on gene expression are indicated by a

change in allelic effects between the parental generation and the F1 hybrids (Cowles et al. 2002;

Wittkopp et al. 2004). Despite its great potential, we are unaware of any field reciprocal

transplant experiment that has measured ASE to evaluate how gene expression divergence

manifests across the habitats known to be responsible for local adaptation.

In this study, we evaluated gene regulatory divergence in nature between locally adapted coastal

and inland population accessions of the yellow monkeyflower, *Mimulus guttatus* (*syn.*

Erythranthe guttata (Fisch. ex DC.) G.L. Nesom). *M. guttatus* is a major study system for

understanding the genetic underpinnings of fundamental patterns in evolutionary ecology

(reviewed in Wu et al. 2008; Twyford et al. 2015). Inland populations of *M. guttatus* have

evolved an early-flowering annual life-history to escape from the seasonal summer drought that

occurs in the Mediterranean climate of the California Coastal Mountain Range. In contrast,

coastal populations of *M. guttatus* are sheltered from the summer drought by pervasive oceanic

fog, which supplies soil moisture and reduces transpiration rates (Corbin et al. 2005). As a result,

these populations have evolved a late-flowering perennial life-history (Hall & Willis 2006;

Lowry et al. 2008). While coastal habitats are benign in terms of soil water availability, these

populations must contend with toxic oceanic salt spray, in response to which they have evolved

high salt tolerance (Lowry et al. 2008, 2009; Selby et al. 2014). Across previous reciprocal

transplant field experiments, inland plants have survived to flower at 23.9 times the rate of

coastal plants in inland habitat, while coastal plants have survived at 2.5 times the rate of inland

plants in coastal habitat (Hall & Willis 2006; Lowry et al. 2008, 2009; Hall et al. 2010).

Population structure analysis has repeatedly demonstrated that coastal populations collectively

constitute a single ecotype, which is distinct genetically from inland annual populations (Lowry

et al. 2008; Twyford & Friedman 2015). The individual genes responsible for the divergence of

coastal and inland ecotypes are unknown, but it is clear that a large paracentric chromosomal

inversion on chromosome 8 is responsible for much of the divergence in traits and fitness

between coastal and inland ecotypes (Lowry & Willis 2010; Friedman 2014; Oneal et al. 2014;

Gould et al. 2017). Smaller effect loci across the genome also account for divergence between

the coastal and inland ecotypes (Hall et al. 2006, 2010; Lowry et al. 2009; Friedman et al. 2015).

In addition, significant heterosis occurs for multiple traits (fitness, flowering time, internode

length) in F1 hybrids between coastal and inland accessions (Lowry et al. 2008; Lowry & Willis

2010). The extent to which gene expression also shows patterns of heterosis in this system is

unknown.

The primary goal of our study was to characterize gene regulatory divergence between coastal

perennial and inland annual *M. guttatus* in the natural habitats that drove the evolution of these

distinct coastal and inland ecotypes. In particular, our study aimed to 1) determine the prevalence

of genotype x environment (GxE) interactions affecting the expression of native and non-native

alleles across the divergent habitats, 2) characterize the relative contributions of *cis* and/or *trans*

regulatory variation in expression divergence, 3) establish the role of genome structure (e.g.

inversions) in the evolution of locally adaptive regulatory variation, 4) quantify gene expression

heterosis across habitats, and 5) identify candidate genes underlying local adaptation in this

system. The results of our study provide new insights into the gene regulatory evolution and

suggest that coupling allele-specific expression with reciprocal transplant experiments is a

promising approach for modern local adaptation research.

MATERIALS AND METHODS

Germplasm

Original field collections of the seeds used in this experiment were made in the Spring of 2005

from the SWB (N39.0359', W123.6904') and LMC (N38.8640', W123.0839') populations

located in Mendocino County, California. The SWB and LMC populations have been the subject

of extensive recent research (Lowry et al. 2008; Lowry & Willis 2010; Wu et al. 2010; Friedman

& Willis 2013; Friedman 2014; Gould et al. 2017). Seeds from these populations were inbred for

multiple generations in the Duke University Greenhouses. The plants used for the field

experiment included the following: 1) An inbred (7 generations self-fertilized) inland line, LMC-

L1-2, henceforth referred to as LL; 2) An inbred (13 generation self-fertilized) coastal line,

SWB-S1-2, henceforth referred to as SS; and 3) An F1 hybrid between the two inbred lines,

henceforth referred to as F1 or SL. The F1s used in this experiment were made in the Michigan

State University (MSU) Growth Chamber Facilities through a cross between the two parental

inbred lines.

Field Reciprocal Transplant Experiment

We conducted a field reciprocal transplant experiment in tandem with allele-specific expression

analysis to understand the causes (*cis*, *trans*) of gene expression divergence between locally

adapted coastal and inland populations of *M. guttatus* (Fig. 1). The reciprocal transplant

experiment was conducted in 2016 using one coastal (Bodega Marine Reserve; N38.3157',

W123.0687') and one inland (Pepperwood Preserve; N38.5755', W122.7008') field site located

in Sonoma County, CA. SS, LL, and F1 seeds were originally sown in Sun Gro Metromix 838

soil on February 3. The seeds were then stratified in the dark at 4°C until February 9, when they

were removed from the cold and germinated in the UC Berkeley Oxford Track Greenhouses.

Seedlings were moved to the Bodega Marine Reserve Greenhouses on February 21 for

acclimation. On February 22, the seedlings were transferred to native soil collected from the field

sites. Seedlings were planted in the field at Bodega on March 7 and at Pepperwood on March 8.

~90 plants per type (SS, LL, F1) were planted at each field site. Plants were evenly distributed

among three blocks per each field site and randomized within block. Tissue for RNA

sequencings was collected from Bodega on April 23 and Pepperwood on April 24. To avoid

effects of circadian rhythm on gene expression, all tissue was collected between 12:00pm and

1:00pm. Both collection days had mostly clear skies, with mean local temperatures during

collection periods of 14° C at Bodega and 17° C at Pepperwood. Plants had progressed to

different developmental stages at time of collection, with some plants approaching flowering and

others not. To best standardize tissue collection across genotypes, only the two youngest pairs of

fully expanded leaves were collected from each plant. Tissue was immediately flash frozen on

liquid nitrogen in the field. Samples were then shipped on dry ice to Michigan State University

for RNA extractions.

RNA-sequencing

Total RNA was extracted from 80 leaf tissue samples with the Spectrum Plant Total RNA Kit

(Sigma-Aldrich, St. Louis, MO, USA). RNA was extracted from 12 replicates of both SS and LL

parental lines from the coastal field site and 13 replicates of both parental lines from the inland

field site. RNA from 15 F1 hybrids from each site was also extracted. Total RNA was submitted

for library preparation and sequencing at the MSU Genome Core

(https://rtsf.natsci.msu.edu/genomics/). RNA-Seq libraries were prepared using the Illumina

TruSeq Stranded mRNA Library Preparation Kit on a Perkin Elmer (Waltham, MA) Sciclone

NGS robot. Completed libraries were quality controlled and quantified using a combination of

Qubit dsDNA HS (Thermo Fisher Scientific, Waltham, MA), Caliper LabChipGX HS DNA

(Perkin Elmer, Waltham, MA), and Kapa Biosystems Illumina Library Quantification qPCR

(Wilmington, MA) assay kits. All libraries were barcoded and library concentrations were

normalized and pooled in equimolar amounts: 8 pools each consisting of 10 randomly chosen

libraries were made. Each of these pools was loaded on one lane of an Illumina HiSeq 2500 High

Output flow cell (v4) and sequenced in the 1x100bp (SE100) format using HiSeq SBS reagents

(also v4). Base calling was done by Illumina Real Time Analysis (RTA) v1.18.64 and output of

RTA was demultiplexed and converted to Fastq format with Illumina Bcl2fastq v1.8.4.

Reference Transcriptome Assembly

Alignment of RNA-seq to a single reference genome can lead to major biases in ASE studies

(Stevenson et al. 2013). To overcome reference bias, we assembled reference transcriptomes for

the coastal perennial (SS) and inland annual (LL) parents. We chose this method over

construction of a pseudo-transcriptome reference (Shen et al. 2013) to avoid generating bias

caused by the reference genome and transcriptome for *M. guttatus* (IM62; Hellsten et al. 2013),

which was generated from an inland annual line. Reference bias is of particular concern in *M.*

guttatus because pairwise polymorphism is generally high (π_S = 0.033-0.065; Brandvain et al.

2014; Gould et al. 2017; Puzey et al. 2017). To generate a reference transcriptome assembly that

contained allele-specific sequences for each parental genotype, we sequenced and conducted *de*

novo assembly using RNA-sequencing. Plants of each of the coastal (SS) and inland (LL) lines

were grown in the MSU greenhouses until both genotypes were flowering. Floral bud, leaf, and

root tissue were collected from each of the SS and LL lines, flash frozen with liquid nitrogen,

and stored at -80° C. Total RNA was extracted from each sample using the Spectrum Plant Total

RNA Kit and submitted to the MSU Genome Core for library construction and sequencing.

Strand-specific Tru-Seq Illumina RNA-seq libraries were prepared and sequenced along with 4

libraries constructed from field collected leaf tissue from the reciprocal transplant experiment (1

sample per parental line per field site). These libraries were included in the transcriptome

assembly pipeline to capture sequences from genes that may only be expressed under field

conditions. Completed libraries were quality controlled and quantified using a combination of

Qubit dsDNA HS, Caliper LabChipGX HS DNA, and Kapa Biosystems Illumina Library

Quantification qPCR assay kits. A total of 10 libraries were barcoded with dual index barcodes

and pooled together. The entire pool was sequenced using 125 bp paired-end reads on a single

lane of the Illumina GA 2500 platform.

Raw output reads were quality checked using FastQC v.0.11.3 (Andrews 2015). Residual adapter

sequences were removed and reads were quality trimmed using Trimmomatic v.0.33 (Bolger et

al. 2014), discarding sequences less than 50 bp in length. Paired and un-paired reads from each

ecotype were combined and assembled into two separate *de novo* (non-reference guided)

assemblies using Trinity v20140413p1 (Grabherr et al. 2011) with *in silico* read normalization.

To check the quality of our two transcriptomes, we compared our *de novo* assemblies to the set

of primary transcripts from the IM62 reference transcriptome using Trinity utility scripts. We

found that 74.3% of primary transcripts represented at least 50% sequence coverage in the inland

(L) transcriptome and 74.1% in the coastal (S) transcriptome. We also aligned the filtered input

reads back to the finished assemblies using Bowtie2 (Langmead et al. 2012). A total of 98.98%

and 99.26% of input reads aligned back to the inland (L) and coastal (S) assemblies, respectively.

We identified possible non-plant contaminant sequences in each assembly through comparison to

the NCBI nt database using blastn. We eliminated transcriptome sequences with top blast

sequence matches (e-value cutoff of 1e-20) to non-plant organisms ($N = 16243$ and $N = 1222$

eliminated sequences for the inland and coastal transcriptomes, respectively).

To construct a combined reference transcriptome that contained pairs of allelic sequences from

both lines, we used a reciprocal best blast strategy. Each transcriptome was compared against the

other (blastn) at an e-value cutoff of 1e-5 and reciprocal best hits were returned using a custom

Python (v. 2.7.2) script (https://github.com/lowrylab/Mimulus_Gene_Expression). We tested a

range of e-value cutoffs in this analysis and found that the choice of e-value between 1e-5 and

1e-50 had little effect on the total output number of best blast pairs (see Fig. S1). We considered

two sequences best blast hits if total high-scoring pair (HSP) coverage was 75% or greater

between sequences. We included only aligned sequence regions of the longest HSP in the

combined transcriptome.

To match transcripts in the combined *de novo* reference transcriptome with their corresponding

Mimulus genes, we compared *de novo* assembled alleles against the reference (IM62) set of

primary transcripts (blastn) at an e-value cut-off of 1e-5. In rare cases, where two alleles from a

de novo assembled transcript matched different *Mimulus* genes, the gene with the lowest e-value

alignment was chosen as a match for the allele pair. Note that with this annotation method, each

de novo assembled transcript has only one matching gene and thus, can be unambiguously

assigned a specific location in the genome. However, multiple transcripts can be assigned to the

same genomic location, which is typical of isoforms stemming from the same gene.

Differential Expression Analyses

Raw reads from the 80 field RNA-seq libraries were de-multiplexed, quality filtered, and

trimmed using FastQC and Trimmomatic, as described above for the transcriptome assembly.

Filtered reads were aligned to the *de novo* combined reference transcriptome (described above)

using the un-gapped aligner Bowtie v1.0.0 (Langmead et al. 2009). We outputted only unique

alignments with zero mismatches. In this way, we analyzed only read count data for reads that

aligned uniquely and perfectly to only one parental allele in the combined reference

transcriptome. Using these criteria, on average 28.6% of library reads per individual were

aligned (range 23.7 – 30.7%). For confirmation, some alignments were visually checked using

IGV v.2.3.65 (Thorvaldsdóttir et al. 2013).

Read alignments to each transcript were quantified using a custom Python (v.2.7.2) script. Using

the raw count data, we identified allele pairs in the transcriptome with a high degree of residual

heterozygosity in the parental inbred lines. This is potentially indicated where a significant

number of reads from a homozygous parent align to the opposite parental allele in the

transcriptome. We eliminated from further analysis any pair of alleles where the average number

of reads aligning to the correct parental allele was less than 5 times the average number of reads

aligning to the alternate allele ($N = 2255$ eliminated transcripts). We also used the initial count

data to test for mapping bias (Fig. S2). For both field sites mapping bias was present but well

controlled. There was a small but statistically significant relationship between transcript

nucleotide divergence and differential expression (equivalent to 0.001 increase in the absolute

value of LFC between genotypes, per SNP). Count data was normalized for library size using

Trimmed Mean of M-values (TMM) normalization to counts per million, which reduces biases

generated by the presence of very highly and lowly expressed transcripts (Robinson et al. 2010).

We eliminated any transcript without at least one library containing $>= 10$ counts per million

reads. The remaining sequences constitute the 'expressed' set of transcripts ($N = 10,122$). The

mean expressed library size for samples from the coast and inland sites was not significantly

different (*t*-tests: all libraries, $P = 0.60$; parental libraries only, $P = 0.22$, Fig. S3A), and is thus,

unlikely to strongly affect statistical comparisons between the two sites. Only 2 transcripts were

expressed at only one site and only 7 transcripts were expressed by only one parental line.

We analyzed the expressed transcript data using the limma (Ritchie et al. 2015) and edgeR

(Robinson et al. 2010) packages in R v.3.2.1 (R. Core Team 2012). Variance stabilizing

normalization of the data was performed using the voom (Law et al. 2014) function in limma,

with a design matrix corresponding to the particular linear model being used for analysis. Model

design matrices were coded using the default treatment-contrast parameterization scheme (see

Smyth et al. (2015), section 9.5.4) and the significance of model effects were calculated using

voom precision weights and the eBayes function. Model effects were judged significant if they

fell below a Benjamini-Hoechberg false-discovery rate corrected *P*-value of 0.05 using the

decideTests function.

We used several different linear models to analyze the count data for expressed transcripts,

modified from the method of Lovell et al. (2016). For this approach, it is important that the size

of the L and S sub-transcriptome in each F1 is approximately equal to produce unbiased results,

which we verified independently (Fig. S4). Each linear model had one or more fixed factors:

allele (L/S), environment (coastal field site vs. inland field site), generation (F0/F1), and

interaction terms. We analyzed the parental plants (F0 generation) separately, the F1 plants

separately, and both generations together in different models. We avoided analyzing models with

more than one categorical factor, except for the purposes of measuring interaction effects

because effect significance testing for crossing categorical factors only captures differences

present in the first level of each factor.

For data from the F0 plants, we ran models examining the allele (homozygous genotype) effect

within each environment separately. We also tested for environmental effects separately within

each parental genotype. We used a model with both genotype and environment to test the

significance of the genotype x environment interactions. Because the inland site had one extra

plant per genotype compared with the coastal site, we analyzed data from 12 randomly chosen

parental lines of each genotype to equalize the power to detect effects between sites. A *post-hoc*

power analysis was conducted using the R package *RnaSeqSampleSize* (Bi & Liu 2016).

We used data from all plants together (both F0 and F1) to test for *cis* and *trans* regulation. *Cis*-

regulatory allelic expression differences are expected to persist regardless of whether alleles are

expressed separately in parental genotypes or together in an F1 hybrid. However, in the F1

genotype, *trans* acting factors from both parental genotypes will influence the expression of an

allele, so *trans* regulation is indicated if allelic patterns of differential expression are different in

the parents vs. the F1 plants. In our model, a significant effect of allele is indicative of a *cis*-

regulatory difference in expression. If a transcript has a significant allele x generation interaction

effect, this indicates *trans* regulation. Note that use of linear models is limited in its ability to

detect *cis* and *trans* effects when those effects are antagonistic (Lovell et al. 2016). In such cases,

significant allelic differential expression can be detected between parental genotypes, but

becomes non-significant in the F1 generation. Within each environment, we tested a model with

the following factors: allele (genotype), generation (F0/F1), and the interaction term. Any

transcript with any *cis* regulation had a significant allele effect (measured in the F1 generation

only). Any transcript with only *trans* regulation had a significant difference between generations,

but a non-significant allele effect in the F1 generation. Any transcript with both *cis* and *trans*

regulation had a significant generational effect and a significant allele effect in the F1.

To gauge the significance of *cis* x environment interactions, we tested the interaction term of a

model using only F1 data and factors allele, environment, and interaction. To gauge the

significance of trans x environment interactions we modeled all the data (F0 and F1) and

measured the significance of a three-way interaction term: allele x generation x environment.

Gene Ontology Enrichment and Expression Networks

We tested for enrichment of gene-ontology (GO) terms among subsets of genes with particular

expression patterns (significant model effects). To annotate GO terms, all IM62 reference

Mimulus proteins were paired with their best *Arabidopsis* protein matches (blastp) at an e-value

cutoff of 10^{-3}. The set of expressed transcripts in our study was matched to the IM62 reference

genome genes, as described above, and corresponding GO terms were matched to the

Arabidopsis BLAST hits. Subsequent GO enrichment tests were done using the R

BIOCONDUCTOR package topGO (Alexa & Rahnenführer 2015). We used the Bioconductor

database 'org.At.tair.db' for annotation and the algorithm 'classic' (statistic = 'fisher') for

statistical tests. Because *P*-value distributions of GO enrichment tests do not typically conform to

a distribution amenable to Benjamini-Hoechberg FDR control, and because such control methods

can be overly conservative causing loss of valuable functional information (Alexa &

Rahnenführer 2015), we report GO terms with uncorrected enrichment *P*-values <= 0.01 (Table

S1). Enrichment tests were conducted at the level of genes rather than transcripts. For each gene

in the IM62 reference transcriptome, we determined the model effects for all matching expressed

transcripts in our data set. A model effect for a gene was considered significant if at least one

matching transcript had a significant effect. GO enrichment was then tested among all expressed

genes ($N = 9326$) with a significant effect vs. all expressed genes without a significant effect.

Enrichments for GxE effects were measured against all genes where either environmental or

allelic DE could be detected. Reaction norm plots were generated using the LSMean expression

level of all transcripts for each gene in each environment.

We conducted clustering analysis of gene expression within each parental genotype using the R

package WGCNA (Langfelder et al. 2008). Gene expression was quantified as the expression

level of the longest matching transcript for each gene. For all analyses, we calculated

dissimilarity of expression using a soft thresholding power of 6 and used the dynamicTreeCut

function to define modules with a minimum number of 30 genes. We calculated the correlation

of each cluster (eigengene) with habitat (Fig. S5). Gene ontology term enrichment was calculated

for all modules with a correlation value of r > 0.75 (Table S2).

Chromosomal Inversion Analyses

Two inverted regions of the genome, one on chromosome 5 and one on chromosome 8, have

been linked to adaptive differences between inland and coastal ecotypes in previous studies

(Lowry & Willis 2010; Oneal et al. 2014; Twyford & Friedman 2015; Gould et al. 2017). When

examining genes in putatively inverted regions of the genome, we included reference genome

v1.1 scaffolds that are inferred to be in the inverted regions from a previous linkage mapping

study (Holeski et al. 2014). For the inversion on chromosome 5, we included v1.1 scaffolds 36,

149, 158, 170, 197, 266, 281, 288, 327, and 368. For the inversion on chromosome 8, we

included v1.1 scaffolds 11, 59, 76, 155 233, 604, and 1093. We tested whether genes in these

regions of the genome are enriched for any model effects. Enrichment tests of model effects

among annotated inversion and non-inversion genes were carried out using Fisher's Exact tests.

The magnitude of model effects among genes with significant effects were compared using

Wilcox rank sum tests in R. It should be noted that the presence of alternative orientations of

chromosome 8 inversion in the SS and LL parental lines has been confirmed by previous crosses

(Lowry & Willis 2010) while the chromosome 5 inversion was originally identified in a QTL

cross between different inland and coastal lines (Holeski et al. 2014). High differentiation

between the SS and LL parental lines in both inversion regions has been demonstrated by an

allele frequency outlier analysis (Gould et al. 2017).

Expression Heterosis

We analyzed heterosis of gene expression separately for each field site using a model with a

single factor for genotype (SS/LL/SL). Data from the inland site was down-sampled to 12 lines

for each parental genotype to equalize power to detect effects between sites. Specific contrasts of

the F1 genotype (SL) were made to each parental genotype (LL/SS) using makeContrasts in

limma. Contrasts were judged significant if they fell below a Benjamini-Hoechberg false-

discovery rate corrected P-value of 0.05. Heterosis of gene expression for each transcript was

classified into 7 categories depending on contrasts between the parental genotype with lower

expression (LP), parental genotype with higher expression (HP), and the F1:

i.   additive expression (no heterosis): LP<F1<HP

ii.   low parent heterosis: F1<LP<HP

iii.   below mid-parent heterosis: F1=LP<HP

iv.   above mid-parent heterosis: LP<HP=F1

v.   high parent heterosis: LP<HP<F1

vi.   ambiguous: LP<HP, but LP=F1=HP

vii.   no differential expression: LP=HP

Transcripts with ambiguous expression patterns were excluded from enrichment tests (Table S1).

Patterns of Candidate Gene Expression

In a study of species-wide genomic variation, Gould et al. (2017) identified a set of 667

candidate adaptation genes. These genes had unusually high differentiation in allele frequencies

between a set of 47 coastal and 50 inland *Mimulus* populations. High differentiation was detected

in either the genic or 1 kb upstream promoter regions of the genes, or both. We tested for

enrichment of differential expression patterns and heterosis among the expressed candidate genes

in the current study. Enrichment was tested against all expressed non-candidate genes using

Fisher's Exact tests. The magnitude of model effects among genes with significant effects were

compared using Wilcox rank sum tests in R.

RESULTS

Reference Transcriptome Assembly

From the tissues of the two *M. guttatus* parental lines we were able to reconstruct both parental

alleles for 25,893 transcripts, of which 708 pairs were removed because they contained no

polymorphism between lines. Eighty-two percent of transcripts could be matched back to

previously annotated *M. guttatus* genes (www.phytozome.org). Our transcriptome reconstructed

54.0% of *M. guttatus* primary transcripts at 50% coverage or greater. On average, there were 43

polymorphisms (SNPs or indels) between each pair of alleles in the reference (Fig. S3B).

Differential Expression Analyses

RNA-sequencing of plants from the field experiments produced approximately 25.9 million raw

reads per library (range 9.2-35.9 million reads). We were able to evaluate gene expression for

10,122 transcripts corresponding to 9,326 genes, after transcriptome reconstruction and filtering.

Significant expression differences (FDR = 0.05) between the inland and coastal field sites were

common (6765 genes; 73% of genes). Expression differences between parental genotypes within

field sites were also common (7213 genes; 77%; Table 1 and 2). We found 54% more genes

were differentially expressed between parental lines at the inland site than at the coastal site

(Table 2). We can attribute only part of this discrepancy to differences in power at the two sites.

Average power to detect DE genes was only slightly higher at the inland site than the coastal site

(75% vs. 68% for a minimum fold-change of 2X).

Genotype x site (GxE) interactions were relatively common. In total, 1837 transcripts

(corresponding to 19% of expressed genes, Table 1, Table S3) had significant GxE interactions,

where the magnitude of expression differences between parental ecotypes depended on the field

site in which they were grown (Fig. 2A). Only 94 transcripts had crossing reaction norms (green

points, Fig. 2B). More often, expression differences between genotypes were significant at one

site and non-significant at the other (1137 transcripts, 11%, blue and red points, Fig. 2).

Most of the expression differences between the coastal and inland lines could be attributed to *cis*-

regulatory variation: 79% (5532 genes) of differentially expressed genes had *cis* regulatory

differences, while about half (3611 genes) of the differentially expressed genes had detectable

trans effects (Fig. 3A-C, Table 1). At both coastal and inland field sites, the magnitudes of *cis*

and *trans* regulatory differences were similar (Table S4). However, we detected far more

significant *trans* effects in the inland field site than the coastal field site (Fig. 3A-C, Table S4).

Overall, there was only moderate overlap between the sets of *cis*-regulated transcripts from the

two environments (Fig. S6). The same was true for the sets of *trans*-regulated transcripts from

the two environments.

Gene Ontology Enrichment and Expression Networks

We explored the functional significance of genes with differential expression between

environments and between parental ecotypes. Differentially expressed genes in the coastal (SS)

parent between sites were enriched for glycosinolate/glucosinolate/S-glycoside biosynthesis and

metabolism, response to starvation, and others (Table S1). Differentially expressed genes in the

inland parent (LL) between sites were enriched for GO categories related to cuticle development,

meristem and floral development, and pigment production. Differentially expressed genes

between the parental lines (across sites) were enriched for genes involved with cell wall

formation, meristem initiation, and responses to viral pathogens. Genes with genotype x site

(GxE) interactions had the most significantly enriched GO terms ($N = 69$), including

brassinosteroid biosynthesis/metabolism, cell wall formation, regulation of development, and

pigment accumulation.

We used network clustering analysis to examine correlated expression of gene clusters (modules)

within habitats for the coastal and inland lines (Pearson's correlation, r > 0.75, Fig. S5). The

coastal perennial line (SS) preferentially expressed modules related to root and epidermal

development at its home site, while oxidative stress response, chlorophyll metabolism, and

circadian rhythm modules were highly expressed in inland habitat (Table S2). The inland annual

line preferentially expressed gene clusters related to nucleoside catabolism, sugar and starch

metabolism, pigment accumulation, and photoperiodism at the inland field site, but had no

modules strongly associated with the coastal field site.

Chromosomal Inversion Analyses

Chromosomal inversions are thought to act as adaptation supergenes that hold together long

haplotypes containing multiple adaptive polymorphisms through suppressed recombination

(Kirkpatrick 2010; Joron et al. 2011; Thompson & Jiggins 2014). We were thus interested in

evaluating the hypothesis of whether the known inversions in this system hold together

haplotypes of differentially expressed genes. There were significantly more differentially

expressed genes inside the chromosome 5 inversion than expected by chance and there were

significantly more GxE interactions for genes in this inversion (Table 1). The adaptive (Lowry &

Willis 2010) chromosome 8 inversion also had significantly more differentially expressed genes,

but only at the coastal site (Table 2). Interestingly, the chromosome 8 inversion had a significant

enrichment of *cis*-eQTLs (Table 1), suggesting that it may maintain a higher level regulatory

divergence caused by the evolution of *cis*-regulatory elements.

Expression Heterosis

Heterosis is commonly found in crosses among locally-adapted populations and can potentially

facilitate gene flow between those populations (Lowry et al. 2008; Charlesworth & Willis 2009;

Baack et al. 2016). Heterosis is typically quantified at the level of an individual trait, including

expression traits by comparing the trait value of hybrids to the trait values of the parents

(Springer & Stupar 2007a,b; Niemi et al. 2015; Lovell et al. 2016). Consistent with previous

studies (Lowry et al. 2008; Lowry & Willis 2010), we have found robust heterosis for multiple

phenotypic traits in F1 hybrids between coastal and inland lines (Fig. 1F). In the current study,

we also found nearly 73% of differentially expressed transcripts had heterotic (non-additive)

expression in the F1 (Fig. 3D, Table S5). The percentage of transcripts with heterotic expression

did not vary strongly by site (Fig 3D), although only 54% of heterotic genes were shared

between sites. Heterotic genes were enriched for functional terms associated with multiple

growth and development terms at both field sites (Table S6).

Patterns of Candidate Gene Expression

To identify genes putatively linked to local adaptation, we cross-referenced our results with a

recent genomic outlier analysis, which identified 667 genes with a high level of allelic

differentiation between coastal and inland populations (Lowry et al. 2010). We found that the

magnitude of differential expression between lines and between field sites was significantly

elevated for candidate genes with highly differentiated SNPs (from Gould et al. 2017) in their

promoter regions (Table S7).

We were able to examine gene expression in seven top candidate genes highlighted by the outlier

analysis and/or previous QTL mapping studies (Fig. 3E-G, Fig. S7). The candidate gene *ABA1*

mediates production of abscisic acid, which controls responses to osmotic stress, including

drought (Xiong et al. 2002). An *ABA1* homolog, Migut.H00431, is located within the

chromosome 8 inversion and may contribute to inland ecotype's adaptation to low soil water

availability. *ABA1* was always more highly expressed in the inland line and was upregulated in

both lines at the inland field site (Fig. 3E). Previously, we showed that coastal *M. guttatus*

populations have evolved leaf tissue tolerance to oceanic salt spray (Lowry et al. 2009). Leaf salt

tolerance is generally mediated by a Na^+/H^+ antiporter (*SOS1*; Qiu et al. 2002), which is

activated by a calcium sensor (*CBL10*; Kim et al. 2007). Both *SOS1* (Migut.E00570) and *CBL10*

(Migut.A00138) were more highly expressed in the coastal line than the inland line across

habitats (Fig. 3G, F), a pattern consistent with elevated salt tolerance of coastal populations.

DISCUSSION

Overall, our results identified multiple key patterns regarding the relationship between gene

expression and the evolution of local adaptation. Differential expression is substantial between

locally-adapted accessions in nature. Expression plasticity of alleles across habitats (genotype x

environment interactions) appear to be common and may minimize fitness trade-offs at

individual loci that contribute to the overall pattern of local adaptation. A large portion of that

differential expression is due to *cis*-regulatory divergence. Chromosomal inversions appear to act

as supergenes by holding together haplotypes of differentially expressed genes, but this pattern

depends on habitat context. Heterosis of gene expression is pervasive and heterotic genes are

enriched for growth and developmental functions. We discuss each of these major findings as

well as the limitations and caveats associated with our study in detail below.

Sources of Variation in Gene Expression

One of the major outstanding questions in local adaptation research is why the fitness effects of

individual loci on local adaptation are so often asymmetric across habitats. Theoretical models

(Levene 1953; Hedrick 1976, 1986; Dieckmann & Doebeli 1999) and studies in key model

systems (Abzhanov et al. 2004; Hoekstra et al. 2006; Barrett et al. 2009) have suggested trade-

offs at individual loci (e.g. antagonistic pleiotropy) to be common. However, field reciprocal

transplant experiments and experimental evolution studies have often found a pattern of

conditional neutrality, where loci have a strong effect on fitness in one environment, but have an

undetectable effect on fitness in alternative environments (reviewed in Bono et al. 2017;

Wadgymar et al. 2017). While a failure to statistically detect a trade-off across habitats does not

rule out the possibility of a small trade-off for an individual locus, it does suggest that adaptive

loci are often strongly asymmetric in their fitness effects across habitats.

Adaptive phenotypic plasticity mediated through gene regulatory divergence between locally

adapted populations is one possible mechanism by which trade-offs could be minimized across

habitats for individual loci (Rutherford 2003; Ghalambor et al. 2007; Des Marais et al. 2013,

2017; Dayan et al. 2015; Lohman et al. 2017). Adaptations of organisms to maintain homeostasis

when faced by environmental challenges can be caused by genotypic specific changes in gene

expression (Gibson 2008; Des Marais et al. 2017). However, such environmentally dependent

changes in gene expression would not be expected to result in a cost if gene expression did not

differ among genotypes in habitats that lacked particular environmental stresses. In our study, we

identified 1137 transcripts that were differentially expressed between the coastal perennial and

inland annual lines in one habitat (coast or inland), but were not differentially expressed in the

alternative habitat. We cannot definitively determine whether any of these genes are responsible

for asymmetries in fitness effects across habitats. Further, some of the genotype specific changes

in gene expression could potentially be maladaptive (Ghalambor et al. 2007). However, our

results do clearly show that habitat-specific differential gene expression is common for locally

adapted ecotypes. Future studies should focus on making the critical linkages between patterns of

gene expression and fitness in the field to determine whether asymmetries of fitness effects

across habitats are the result of habitat-specific differential gene expression.

Overall, *cis*-regulatory evolution was responsible for at least 79% of the differentially expressed

genes. This is about twice the percentage of *cis*-regulated differentially expressed genes for

crosses both within and between *Drosophilia* species (Coolon et al. 2014). However, the

relatively high proportion of *cis*-regulatory divergence underlying differentially expressed genes

is not too surprising, given the high level of polymorphism among *M. guttatus* accessions

(Brandvain et al. 2014; Gould et al. 2017; Puzey et al. 2017). Indeed, the degree of allele

frequency differentiation in promoter regions between the coastal and inland ecotypes (Gould et

al. 2017) was predictive of differential expression in this study. Further, in terms of the

proportion of *cis* regulated genes out of all genes surveyed (59% for our study), our results are

not very different from other study systems. Mack et al. (2016) found significant *cis* effects for

68% of surveyed genes in crosses between house mice subspecies (*Mus musculus musculus* and

M. m. domesticus). Steige et al. (2015) found that 33-39% of genes surveyed had significant

allele-specific expression effects for hybrids between accessions of the plant *Capsella*

grandiflora.

Gene Regulatory Evolution and Chromosomal Inversions

Chromosomal inversions have been thought to be involved in evolutionary adaptations since

classic studies in *Drosophila* found that inversion polymorphisms are frequently correlated with

environmental conditions over geographic space and through time (Dobzhansky 1951, 1970;

Kirpatrick & Barton 2006; Hoffman & Rieseberg 2008; Cheng et al. 2012; Ayala et al. 2013,

2014; Adrion et al. 2015). Inversions strongly suppress genetic recombination in heterokaryotic

(inversion heterozygotes) individuals because recombinant gametes are unbalanced (Dobzhansky

1970; Rieseberg 2001). Researchers have thus hypothesized that inversions could act as

supergenes by holding together long haplotypes containing multiple adaptive polymorphisms

through suppressed recombination (Dobzhansky 1970; Joron et al. 2011; Thompson & Jiggins

2014; Kirkpatrick & Barton 2006; Kirkpatrick & Barrett 2015).

The supergene hypothesis is a major reason why inversions are thought to frequently contribute

to the evolution of adaptations (Dobzhansky 1970; Kirkpatrick & Barton 2006; Hoffman &

Rieseberg 2008; Faria & Navarro 2009; Thompson & Jiggins 2014). Early researchers,

especially Dobzhansky (1970), argued that inversions could hold together co-adapted epistatic

complexes of genes through suppressed recombination. However, theoreticians have recently

shown that inversions can spread due to purely additive effects, when at least two locally

adaptive loci are captured within the inverted region (Kirkpatrick & Barton 2006; Kirkpatrick &

Barrett 2015). Regardless of the ultimate genetic mechanisms, these models all predict that

suppression of recombination by inversions maintains adaptive divergence at multiple linked

loci.

Global gene expression analyses offer one way to test whether inversions hold together

haplotypes of genetic variation potentially responsible for phenotypic divergence between locally

adapted populations. Research in a few systems has found evidence of increased levels of

expression divergence in inversions (Marquès-Bonet et al. 2004; Cassone et al. 2011; Fuller et al.

2016). For example, Marquès-Bonet et al. (2004) found 8.9% greater divergence (fold change) in

human versus chimp brain gene expression for chromosomal inversions when compared to the

rest of the genome. We did not observe a significant difference in fold change between inverted

and non-inverted regions in our study. However, genes within the region of the chromosome 5

inversion were 12.5% more likely to be differentially expressed. Genes within the adaptive

chromosome 8 inversion were 8.3% more likely to have a *cis*-eQTL.

The enrichment of gene expression differences in our study is consistent with patterns found in

inversions between chimpanzees and humans (Marquès-Bonet et al. 2004) and *Drosophila*

(Fuller et al. 2016), but contrasts with results for the plant *Boechera stricta* (Lee et al. 2017).

While the finding of an enrichment of differential gene expression for inversions in this study

and others is consistent with the supergene hypothesis of inversion evolution, it is by no means

definitive proof. Demonstrating that an inversion has evolved as an adaptation supergene

requires confirmation that multiple genes within the inversion have effects on adaptive

phenotypes (Hoffman & Rieseberg 2008; Kirkpatrick 2010; Kirkpatrick & Kern 2012). There is

currently no study that we are aware of which has experimentally confirmed that two or more

genes within an inversion contribute additively or epistatically to adaptive phenotypic

differences between locally adapted populations, ecotypes, or species (but see Kunte et al. 2014).

Heterosis of Gene Expression

We have previously found that heterosis is common in F1 hybrids between coastal perennial and

inland annual accessions (Lowry et al. 2008; Lowry & Willis 2010). In this study, we observed a

high level of above mid-parent heterosis for gene expression. However, only a small percentage

(<1%) of genes had above high-parent or below low-parent heterosis. Heterosis often causes

increased growth and vigor (reviewed in Hochholdinger & Hoecker 2007; Charlesworth & Willis

2009). Consistent with this pattern, heterotic genes were enriched for functional terms associated

with growth and development at both field sites (Table S1). The enrichment of heterotic genes

involved with growth and development likely reflects the downstream responses of the ultimate

genetic causes of heterosis.

Even though heterosis is a well-documented phenomenon occurring in hybrids, the ultimate

causes of heterosis are still not well characterized (Hochholdinger & Hoecker 2007;

Charlesworth & Willis 2009). The dominance hypothesis postulates that heterosis is the result of

masking of deleterious recessive loci through complementation genome-wide in hybrids.

Alternatively, the overdominance hypothesis proposes that heterosis is the result of heterozygote

advantage at multiple loci across the genome. A third major alternative hypothesis is that

heterosis is conferred by epistatic interactions.

Some readers may question our use of the term heterosis to describe the observation of non-

additive gene expression in F1 hybrid individuals. Many in the field have used other terms to

describe non-additive gene expression in allele-specific expression studies, often labeling mid-

parent heterosis as "overdominance" and below mid-parent heterosis as "underdominance"

(Gibson et al. 2004; Swanson-Wagner et al. 2006; Rockman & Kruglyak 2006; Bell et al. 2013).

However, the use of overdominance in this way conflates the phenomena of heterosis with one of

the hypotheses for its ultimate cause (i.e. heterosis is conflated with the overdomiance

hypothesis; Hochholdinger & Hoecker 2007; Niemi et al. 2015). Further, it is important to

recognize that the quantification of heterosis is not limited to fitness and yield traits. In the plant

biology and crop literature, heterosis is routinely quantified for any trait measured in F1 hybrids,

including leaf shape, root morphology, and flowering time (Blum et al. 1977; Barth et al. 2003).

Gene expression is also a trait. Thus, heterosis can be quantified as transcript levels for

individual genes, as has been done in previous studies (Huang et al. 2006; Stupar & Springer

2006; Ji et al. 2014; Niemi et al. 2015; Lovell et al. 2016).

Caveats, Limitations, and Future Directions

There are important caveats and concerns to consider in the interpretation of the results of our

study. One of the foremost caveats of the study is its narrow scope. We only evaluated gene

expression differences between one inbred line per ecotype, at two field sites, and for gene

expression at one time point. Despite these drawbacks, this is the first study that we are aware of

to evaluate allele-specific expression in the field between known locally adapted populations

(Voelckel et al. 2017), which allowed us to detect novel *cis* x habitat and *trans* x habitat

interaction (Fig. 3; Table S8). The major outstanding questions include how variable gene

expression is across populations, field sites, and tissue types? In the future, it will be critical to

evaluate gene expression at different time points and across tissue types to gain a more

comprehensive understanding of how ecotypic differences in gene expression manifest

throughout development under natural field conditions.

Perhaps, the greatest challenge for interpreting gene expression results from reciprocal transplant

experiments is understanding how patterns of expression actually contribute to phenotypes and

fitness across habitats. Fitness trade-offs that are associated with local adaptation could

potentially be caused by any type of gene expression pattern. Constitutive expression differences

between ecotypes could cause stable phenotypic differences that are beneficial in one habitat but

deleterious in another. For example, in our study, expression of *SOS1* does not appear to be

plastic in either ecotype and it may be a major contributor to the fitness trade-offs observed

between habitats (Fig. 3F). Gene expression characterized by genotype x environment

interactions could cause fitness trade-offs, but just as likely could mitigate trade-offs by helping

ecotypes maintain homeostasis across habitats.

Until recently, studies of gene regulation have for the most part been limited to the laboratory

setting, and coupling those studies with reciprocal transplant experiments to better understand

local adaptation is still rare (Kenkel & Matz 2016; Lohman et al. 2017; Voelckel et al. 2017).

Allele-specific expression is clearly an important tool for understanding the evolution of

regulatory differences between locally adapted populations, ecotypes, and species. However we

hope that future reciprocal transplant experiments will go beyond RNA-sequencing by also

examining genotypic differences in genome-wide patterns of chromatin remodeling and protein

translation across divergent habitats (Rodgers-Melnick et al. 2016; Voelckel et al. 2017).

ACKNOWLEDGMENTS

We thank D. Des Marais, J. Tung, J. Friedman, R. Guerrero, and J. Lasky for comments that

helped to improve our manuscript. We thank K. Childs and J. Lovell for guidance with

expression analyses, B. Blackman and E. Patterson for starting seeds for us at UC Berkeley, and

D. Popovic for assistance in field planting. The Bodega Marine Reserve and the Pepperwood

Preserve kindly allowed us to conduct this reciprocal transplant field experiment. Funding was

provided by Michigan State University. The authors declare no conflicts of interest.

REFERENCES

Abzhanov A, Protas M, Grant BR *et al.* (2004) Bmp4 and morphological variation of beaks in

Darwin's finches. *Science,* **305,** 1462-1465.

Adrion JR, Hahn MW, Cooper BS (2015) Revisiting classic clines in *Drosophila melanogaster*

in the age of genomics. *Trends in Genetics*, **31**, 434-444.

Aguilar-Rangel MR, Montes RA, González-Segovia E *et al.* (2017) Allele specific expression

analysis identifies regulatory variation associated with stress-related genes in the Mexican

highland maize landrace Palomero Toluqueño. *PeerJ,* **23**, 5e3737.

Ågren, J, Oakley CG, McKay JK *et al.* (2013) Genetic mapping of adaptation reveals fitness

trade-offs in *Arabidopsis thaliana. Proceedings of the National Academy of Sciences,* **110**,

21077-21087.

Alexa A, Rahnenführer J (2015) Gene set enrichment analysis with topGO. http://www.mpi-

sb.mpg.de/~alexa

Anderson JT, Willis JH, Mitchell-Olds T (2011) Evolutionary genetics of plant adaptation.

Trends in Genetics, **27**, 258-266.

Anderson JT, Lee CR, Rushworth C *et al.* (2013) Genetic tradeoffs and conditional neutrality

contribute to local adaptation. *Molecular Ecology*, **22**, 699-708.

Andrews S (2015) FastQC v0.11.3. A quality control tool for high throughput sequence data.

http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Ayala D, Guerrero RF, Kirkpatrick M (2013) Reproductive isolation and local adaptation

quantified for a chromosome inversion in a malaria mosquito. *Evolution,* **67**, 946-958.

Baack E, Melo MC, Rieseberg LH, *et al.* (2016) The origins of reproductive isolation in plants.

New Phytol, **207,** 207, 968-984.

Barrett RD, Rogers SM, Schluter D (2009) Environment specific pleiotropy facilitates

divergence at the Ectodysplasin locus in threespine stickleback. *Evolution*, **63**, 2831-2837.

Barth S, Busimi AK, Utz HF, Melchinger AE (2003) Heterosis for biomass yield and related

traits in five hybrids of *Arabidopsis thaliana* L. Heynh. *Heredity,* **1**, 36-42.

Bi R, Liu P (2016) Sample size calculation while controlling false discovery rate for differential

expression analysis with RNA-sequencing experiments. BMC Bioinformatics, **17**, 146.

Bell GD, Kane NC, Rieseberg LH, Adams KL (2013) RNA-seq analysis of allele-specific

expression, hybrid effects, and regulatory divergence in hybrids compared with their parents

from natural populations. *Genome Biology and Evolution,* **5**, 1309-1323.

Blum A, Jordan WR, Arkin GF (1977) Sorghum root morphogenesis and growth. II.

Manifestation of heterosis. *Crop Science*, **17**, 153-157.

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics,* **30**, 2114-2120.

Bono LM, Smith LB, Pfennig DW, Burch CL (2017) The emergence of performance trade-offs

during local adaptation: insights from experimental evolution. *Molecular Ecology*, **26**, 1720-

1733.

Brandvian Y, Kenney AM, Flagel L *et al.* (2014) Speciation and introgression between *Mimulus*

nasutus and *Mimulus guttatus*. *PLoS Genetics,* **10**, e1004410.

Cassone BJ, Molloy MJ, Cheng C *et al.* (2011) Divergent transcriptional response to thermal

stress by *Anopheles gambiae* larvae carrying alternative arrangements of inversion 2La.

Molecular Ecology, **20**, 2567-2580.

Charlesworth D, Willis JH (2009) The genetics of inbreeding depression. *Nature Review*

Genetics, **10**, 783-796.

Cheng C, White BJ, Kamdem C *et al.* (2012) Ecological genomics of *Anopheles gambiae* along a

latitudinal cline: a population-resequencing approach. *Genetics,* **190**, 1417-1432.

Colosimo PF, Hosemann KE, Balabhadra S *et al.* (2005) Widespread parallel evolution in

sticklebacks by repeated fixation of ectodysplasin alleles. *Science,* **307**, 1928-1933.

Coolon JD, McManus CJ, Stevenson KR *et al.* (2014) Tempo and mode of regulatory evolution

in Drosophila. *Genome Research,* **24**, 797-808.

Clausen J (1951) Stages in the evolution of plant species. *Cornell University Press*. Ithaca NY,

USA.

Corbin JD, Thomsen MA, Dawson TE, D'Antonio CM (2005) Summer water use by California

coastal prairie grasses: fog, drought, and community composition. *Oecologia*, **145**, 511–521.

Cowles CR, Hirschhorn JN, Altshuler D *et al.* (2002) Detection of regulatory variation in mouse

genes. *Nature Genetics,* **32**, 432-437.

Cubillos FA, Stegle O, Grondin C *et al.* (2014) Extensive *cis*-regulatory variation robust to

environmental perturbation in *Arabidopsis*. *Plant Cell,* **26**, 4298-4310.

Dayan DI, Crawford DL, Oleksiak MF (2015) Phenotypic plasticity in gene expression

contributes to divergence of locally adapted populations of *Fundulus heteroclitus*. *Molecular*

Ecology, **24**, 3345-3359.

de Meaux J, Goebel U, Pop A *et al.* (2005) Allele-specific assay reveals functional variation in

the chalcone synthase promoter of *Arabidopsis thaliana* that is compatible with neutral

evolution. *Plant Cell,* **17**, 676-690.

Des Marais DL, Hernandez KM, Juenger TE (2013) Genotype-by-environment interaction and

plasticity: exploring genomic responses of plants to the abiotic environment. *Annual Review of*

Ecology, Evolution, and Systematics, **44**, 5-29.

Des Marais DL, Guerrero RF, Lasky JR, Scarpino SV (2017) Topological features of a gene co-

expression network predict patterns of natural diversity in environmental response. *Proceedings*

of the Royal Society B, **284**, 20170914.

Dieckmann U, Doebeli M (1999) On the origin of species by sympatric speciation. *Nature*, **400**,

354–357.

Dobzhansky T. 1951. Genetics and the Origin of Species (3rd ed.). *Columbia University Press.*

New York, NY, USA.

Dobzhansky T. 1970. Genetics of the Evolutionary Process. *Columbia University Press*. New

York, NY, USA.

Faria R, Navarro A (2010) Chromosomal speciation revisited: rearranging theory with pieces of

evidence. *Trends in Ecology & Evolution,* **25**, 660-669.

Fuller ZL, Haynes GD, Richards S *et al.* (2016) Genomics of natural populations: how

differentially expressed genes shape the evolution of chromosomal inversions in *Drosophila*

pseudoobscura. *Genetics,* **204**, 287-301.

Friedman J (2014) Genetic determinants and epistasis for life history trait differences in the

common monkeyflower, *Mimulus guttatus*. *Journal of Heredity,* **105**, 816-827.

Friedman J, Twyford AD, Willis JH *et al.* (2015) The extent and genetic basis of phenotypic

divergence in life history traits in *Mimulus guttatus*. *Molecular Ecology*, **24**, 111-122.

Gardner KM, Latta RG (2006) Identifying loci under selection across contrasting environments

in *Avena barbata* using quantitative trait locus mapping. *Molecular Ecology,* **15,** 1321-1333.

Ghalambor CK, McKay JK, Carroll SP *et al.* (2007) Adaptive versus non-adaptive phenotypic

plasticity and the potential for contemporary adaptation in new environments. *Functional*

Ecology, **21**, 394-407.

Gibson G (2008) The environmental contribution to gene expression profiles. *Nature Reviews*

Genetics, **9**, 575-581.

Gibson G, Riley-Berger R, Harshman L *et al.* (2004) Extensive sex-specific nonadditivity of

gene expression in *Drosophila melanogaster*. *Genetics,* **167**, 1791-1799.

Gould BA, Chen Y, Lowry DB (2017) Pooled ecotype sequencing reveals candidate genetic

mechanisms for adaptive differentiation and reproductive isolation. *Molecular Ecology,* **26**, 163-

177.

Grabherr MG, Haas BJ, Yassour M *et al.* (2011) Full-length transcriptome assembly from RNA-

Seq data without a reference genome. *Nature Biotechnology,* **29**, 644-652.

Hall MC, Willis JH (2006) Divergent selection on flowering time contributes to local adaptation

in *Mimulus guttatus* populations. *Evolution,* **60**, 2466-2477.

Hall MC, Basten C, Willis JH (2006) Pleiotropic quantitative trait loci contribute to population

divergence in traits associated with life-history variation in *Mimulus guttatus*. *Genetics,* **172**,

1829-1844.

Hall MC, Lowry DB, Willis JH (2010) Is local adaptation in *Mimulus guttatus* caused by

tradeoffs at individual loci? *Molecular Ecology* **19**, 2739-2753.

Hedrick PW, Ginevan ME, Ewing EP. 1976. Genetic polymorphism in heterogeneous

environments. *Annual Review of Ecology and Systematics*, **7**, 1-32.

Hedrick PW (1986) Genetic polymorphism in heterogeneous environments: a decade later.

Annual Review of Ecology and Systematics, **17**, 535-566.

Hellsten U, Wright KM, Jenkins J *et al.* (2013) Fine-scale variation in meiotic recombination in

Mimulus inferred from population shotgun sequencing. *Proceedings of the National Academy of*

Sciences, **110**, 19478-19482.

Hereford J (2009) A quantitative survey of local adaptation and fitness trade-offs. *American*

Society of Naturalists, **173**, 579-588.

Hoekstra HE, Hirschmann RJ, Bundey RA *et al.* (2006) A single amino acid mutation

contributes to adaptive beach mouse color pattern. *Science,* **313**, 101-104.

Hochholdinger F, Hoecker N (2007) Towards the molecular basis of heterosis. *Trends in Plant*

Science, **12**, 427-432.

Holeski L, Monnahan P, Koseva B *et al.* (2014) A high-resolution genetic map of yellow

monkeyflower identifies chemical defense QTLs and recombination rate variation. *G3: Genes,*

Genomes, Genetics, **4**, 813-821.

Hoffmann AA, Rieseberg LH (2008) Revisiting the impact of inversions in evolution: From

population genetic markers to drivers of adaptive shifts and speciation? *Annual Review of*

Ecology, Evolution, and Systematics, **39**, 21-42.

Huang Y, Zhang L, Zhang J *et al.* (2006) Heterosis and polymorphisms of gene expression in an

elite rice hybrid as revealed by a microarray analysis of 9198 unique ESTs. *Plant Molecular*

Biology, **62**, 579-591.

Ji H, Pardo JM, Batelli G *et al.* (2013) The Salt Overly Sensitive (SOS) pathway: established and

emerging roles. *Molecular Plant,* **6**, 275-286.

Joron M, Frezal L, Jones RT *et al.* (2011) Chromosomal rearrangements maintain a polymorphic

supergene controlling butterfly mimicry. *Nature,* **477**, 203-206.

Kawecki TJ, Ebert D (2004) Conceptual issues in local adaptation. *Ecology Letters,* **7**, 1225-

1241.

Kenkel CD, Matz MV (2016) Gene expression under chronic heat stress in populations of the

mustard hill coral (*Porites astreoides*) from different thermal environments. *Nature Ecology and*

Evolution, **1,** 0014.

Kim BG, Waadt R, Cheong YH, Pandey GK, Dominguez-Solis JR, Schültke S, Lee SC, Kudla J,

Luan S. (2007) The calcium sensor CBL10 mediates salt tolerance by regulating ion homeostasis

in *Arabidopsis*. *The Plant Journal*, **52**, 473-484.

Kirkpatrick M, Barton NH (2006) Chromosome inversions, local adaptation and speciation.

Genetics, **173**, 419-434.

Kirkpatrick M, Barrett B (2015) Chromosome inversions, adaptive cassettes and the evolution of

species' ranges. *Molecular Ecology,* **24**, 2046-2055.

Kirkpatrick M, Kern A (2012) Where's the money? Inversions, genes, and the hunt for genomic

targets of selection. *Genetics*, **190**, 1153-1155.

Kirkpatrick M (2010) How and why chromosome inversions evolve. *PLOS Biology*, **8**, e1000501

Langfelder P, Horvath S (2008) WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics,* **9**, 559.

Kunte K, Zhang W, Tenger-Trolander A et al. (2014) Doublesex is a mimicry supergene. *Nature*,

507, 229-232.

Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nature Methods,* **9**,

357-359.

Langmead B, Trapnell C, Pop M *et al.* (2009) Ultrafast and memory-efficient alignment of short

DNA sequences to the human genome. *Genome Biology,* **10**, R25.

Law CW, Chen Y, Shi W *et al.* (2014) Precision weights unlock linear model analysis tools for

RNA-seq read counts. *Genome Biology,* **15**, R29.

Lee CR, Wang B, Mojica JP *et al.* (2017) Young inversion with multiple linked QTLs under

selection in a hybrid zone. *Nature Ecology and Evolution,* **1**, 0119.

Leinonen PH, Remington DL, Leppala *et al.* (2013) Genetic basis of local adaptation and

flowering time variation in *Arabidopsis lyrata*. *Molecular Ecology,* **22,** 709-723.

Levene H (1953) Genetic equilibrium when more than one ecological niche is available. *The*

American Naturalist, **87**, 331-333.

Lohman B, Stutz W, Bolnick D (2017) Gene expression stasis and plasticity following migration

into a foreign environment. *bioRxiv*, 121608.

Ji T, Liu P, Nettleton D (2014) Estimation and testing of gene expression heterosis. *Journal of*

Agricultural, Biological, and Environmental Statistics, **19**, 319.

Lowry DB, Rockwood JH, Willis JH (2008) Ecological reproductive isolation of coast and

inland races of *Mimulus guttatus*. *Evolution,* **62**, 2196-2214.

Lowry DB, Hall MC, Salt DE *et al.* (2009) Genetic and physiological basis of adaptive salt

tolerance divergence between coastal and inland *Mimulus guttatus*. *New Phytol,* **183**, 776-788.

Lowry DB, Willis JH (2010) A widespread chromosomal inversion polymorphism contributes to

a major life-history transition, local adaptation, and reproductive isolation. *PLOS Biology,* **8**,

e1000500.

Lovell JT, Schwartz S, Lowry DB *et al.* (2016) Drought responsive gene expression regulatory

divergence between upland and lowland ecotypes of a perennial C4 grass. *Genome Research,* **26**,

510-518.

Mack KL, Campbell P, Nachman MW (2016) Gene regulation and speciation in house mice.

Genome Research, **26**, 451-461.

Marquès-Bonet T, Cáceres M, Bertranpetit J *et al.* (2004) Chromosomal rearrangements and the

genomic distribution of gene-expression divergence in humans and chimpanzees. *Trends in*

Genetics, **20**, 524-529.

Niemi J, Mittman E, Landau W *et al.* (2015) Empirical Bayes analysis of RNA-seq data for

detection of gene expression heterosis. *Journal of Agricultural, Biological, and Environmental*

Statistics, **20**, 614-628.

Oneal E, Lowry DB, Wright KM *et al.* (2014) Divergent population structure and climate

associations of a chromosomal inversion polymorphism across the *Mimulus guttatus* species

complex. *Molecular Ecology,* **23**, 2844-2860.

Puzey JR, Willis JH, Kelly JK (2017) Population structure and local selection yield high

genomic variation in *Mimulus guttatus*. *Molecular Ecology*, **26**, 519-535.

Qiu QS, Guo Y, Dietrich MA, Schumaker KS, Zhu JK (2002) Regulation of SOS1, a plasma

membrane Na+/H+ exchanger in *Arabidopsis thaliana*, by SOS2 and SOS3. *Proceedings of the*

National Academy of Sciences, **99**, 8436-8441.

R. Core Team (2012) R: A language and environment for statistical computing. (available at

http://www.r-project.org/.le).

Rieseberg LH (2001) Chromosomal rearrangements and speciation. *Trends in Ecology &*

Evolution, **16**, 351-358.

Ritchie ME, Phipson B, Wu D *et al.* (2015) limma powers differential expression analyses for

RNA-sequencing and microarray studies. *Nucleic Acids Research,* **43**, e47.

Robinson M, Oshlack A (2010) A scaling normalization method for differential expression

analysis of RNA-seq data. *Genome Biology,* **11**, R25.

Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. *Bioinformatics,* **26**, 139-140.

Rockman MV, Kruglyak L (2006) Genetics of global gene expression. *Nature Reviews Genetics,*

7, 862-872.

Rodgers-Melnick E, Vera DL, Bass HW, Buckler ES (2016) Open chromatin reveals the

functional maize genome. *Proceedings of the National Academy of Sciences*, **113**, E3177-3184.

Rutherford SL (2003) Between genotype and phenotype: protein chaperones and evolvability.

Nature Review Genetics, **4**, 263-274.

Selby JP, Jeong AL, Toll K, Wright KM, Lowry DB (2014) Methods and discoveries in the

pursuit of understanding the genetic basis of adaptation to harsh environments in *Mimulus*. In

Plant Ecology and Evolution in Harsh Environments (pp. 243-265). *NOVA Science Publishers*.

Hauppauge, NY, USA.

Shen Y, Garcia T, Pabuwal V, et al. (2013) Alternative strategies for development of a reference

transcriptome for quantification of allele specific expression in organisms having sparse genomic

resources. *Comp. Biochem. Physiol. - Part D Genomics Proteomics*, **8**, 11–16.

1003

1004 Sobel JM, Chen GF, Watt LR *et al.* (2010) The biology of speciation. *Evolution,* **64**, 295-315.

1005

1006 Smyth GK, Ritchie M, Thorne N (2015) Linear models for microarray and RNA-Seq data user's

1007 guide. R, doi:10.1093/nar/gkv007.

1008

1009 Stupar RM, Springer NM (2006) *Cis*-transcriptional variation in maize inbred lines B73 and

1010 Mo17 leads to additive expression patterns in the F1 hybrid. *Genetics*, **173**, 2199-2210.

1011

1012 Springer NM, Stupar RM (2007a) Allelic variation and heterosis in maize: how do two halves

1013 make more than a whole? *Plant Cell,* **19**, 2391-2402.

1014

1015 Springer NM, Stupar RM (2007b) Allele-specific expression patterns reveal biases and embryo-

1016 specific parent-of-origin effects in hybrid maize. *Plant Cell,* **19**, 2391-2402.

1017

1018 Steige KA, Reimegård J, Koenig D *et al.* (2015) *Cis*-regulatory changes associated with a recent

1019 mating system shift and floral adaptation in *Capsella*. *Molecular Biology and Evolution,* **32**,

1020 2501-2514.

1021

1022   Steige KA, Laenen B, Reimegård J (2017) Genomic analysis reveals major determinants of cis-

1023   regulatory variation in *Capsella grandiflora*. *Proceedings of the National Academy of Sciences,*

1024   **114**, 1087-1092.

1025

1026   Stevenson KR, Coolon JD, Wittkopp PJ (2013) Sources of bias in measures of allele-specific

1027   expression derived from RNA-seq data aligned to a single reference genome. *BMC genomics,* **14**,

1028   536.

1029

1030   Swanson-Wagner R, Jia Y, De Cook R *et al.* (2006) All possible modes of gene action are

1031   observed in a global comparison of gene expression in a maize f1 hybrid and its inbred parents.

1032   *Proceedings of the National Academy of Sciences* **103,** 6805–6810.

1033

1034   Thompson MJ, Jiggins CD 2014. Supergenes and their role in evolution. *Heredity*, **113**, 1–8.

1035

1036   Thorvaldsdóttir H, Robinson JT, Mesirov JP (2015) Integrative Genomics Viewer (IGV): high-

1037   performance genomics data visualization and exploration. *Briefings in Bioinformatics,* **14**, 1476-

1038   1486.

1039

1040   Tung J, Primus A, Bouley AJ, Severson TF, Alberts SC, Wray GA (2009) Evolution of a malaria

1041   resistance gene in wild primates. *Nature*, **460**, 388-391.

1042

1043   Twyford AD, Friedman J (2015) Adaptive divergence in the monkey flower *Mimulus guttatus* is

1044   maintained by a chromosomal inversion. *Evolution,* **69**, 1476-1486.

1045

1046   Twyford AD, Streisfeld MA, Lowry DB *et al.* Genomic studies on the nature of species:

1047   adaptation and speciation in *Mimulus*. *Molecular Ecology,* **24**, 2601-2609.

1048

1049   Verhoeven KJF, Poorter H, Nevo E *et al.* (2008) Habitat-specific natural selection at a

1050   flowering-time QTL is a main driver of local adaptation in two wild barley populations.

1051   *Molecular Ecology,* **17,** 3416-3424.

1052

1053   Voelckel C, Gruenheit N, Lockhart P (2017) Evolutionary transcriptomics and proteomics:

1054   Insight into plant adaptation. *Trends in Plant Science*, **22**, 462-471.

1055

1056   Wadgymar S, Lowry DB, Gould BA *et al.* (2017) Identifying targets and agents of selection:

1057   innovative methods to evaluate the processes that contribute to local adaptation. *Methods in*

1058   *Ecology and Evolution*, **8**, 738-749.

1059

1060   Wang M, Uebbing S, Ellegren H (2017) Bayesian inference of allele-specific gene expression

1061   indicates abundant *cis*-regulatory variation in natural flycatcher populations. *Genome Biology*

1062   *and Evolution,* **9**, 1266-1279.

1063

1064   Wittkopp PJ, Haerum BK, Clark AG (2004) Evolutionary changes in *cis* and *trans* gene

1065   regulation. *Nature*, **430**, 85-88.

1066

1067   Wu CA, Lowry DB, Nutter LI *et al.* (2010) Natural variation for drought-response traits in the

1068   *Mimulus guttatus* species complex. *Oecologia,* **162**, 23-33.

1069

1070   Xiong L, Lee H, Ishitani M *et al.* (2002) Regulation of osmotic stress-responsive gene

1071   expression by the LOS6/ABA1 Locus in *Arabidopsis. The Journal of Biological Chemistry,* **277**,

1072   8588–8596.

1073

1074   Zhang X, Borevitz JO (2009) Global analysis of allele-specific expression in *Arabidopsis*

1075   *thaliana*. *Genetics,* **182**, 943-954.

1076

1077   **DATA ACCESSIBILITY**

1078   RNA-sequence data are deposited are available through the NCBI Sequence Read Archive under

1079   project identifier XX.

1080

1081   **AUTHOR CONTRIBUTIONS**

1082   DBL designed and conducted the field experiment. YC conducted the laboratory work. BAG

1083   analyzed the data. DBL and BAG wrote the manuscript.

1084

1085

1086

1087

1088

1089

 TABLES

1091

1092   Table 1: Summary of model effects on expressed genes.

	Inversion 5 (246)		Inversion 8 (311)		Non-Inversion (8805)		total (9326)
Differential Expression (model 1):	no.	LFC	no.	LFC	no.	LFC	no.
Genotypic DE	201	0.893	239	0.932	6773	0.86	7213
Environmental DE	185	0.667	230	0.808	6350	0.729	6765
Genotype by Environment	**60***	0.19	61	0.213	1659	0.196	1780
Regulation effects (model 2):							
total DE genes	**204***		232		6490		6962
Cis regulation	158	1.023	**198***	1.04	5176	1.04	5532
Trans regulation	106	1.707	109	1.628	3396	1.229	3611

1093   *significantly greater percentage than non-inversion regions at p<0.05, **p<0.01.
1094   LFC: average magnitude (abs) log2 fold-change of genes with significant effects
1095

1096

1097

1098   Table 2: Number and average effect size of differentially expressed genes within each habitat.

	Coast		Inland	
	no.	LFC	no.	LFC
Inversion 5 (246)	119	1.136	184	0.932
Inversion 8 (311)	**161***	1.092	210	0.993
Non-Inversion (8805)	3997	1.084	6195	0.897
Total (9326)	4277	1.085	**6589***	**0.901***

1099   *significantly greater than non-inversion regions at p<0.05
1100   *** significantly greater than the coastal field site, p<0.001
1101   LFC: average magnitude (abs) log2 fold-change of genes with significant effects
1102

1103

Figure 1. A) Map of Northern California with the location of two field reciprocal transplant sites. Colors represent mean temperature of the warmest quarter in °C x 10. Inset: left, coastal ecotype; right, inland ecotype. B) The coastal field site. C) The inland field site. D) SS parental gene expression at two field sites. E) LL parental gene expression at two field sites. F) Survival of parental and F1 (SL) plants at the two field sites.

Figure 2. A) Average log2 fold-change (LFC) differences between parental genotypes for expressed transcripts at two field sites. B) Expression reaction norms corresponding to colored points in A and B. LFC has been restricted to <3 and <-3 for graphing purposes. Note, 28 transcripts with ambiguous expression patterns are not included.

Figure 3. *Cis* and *trans* regulated transcripts detected at the coastal (A) and inland (B) field sites. Average log2-fold change (LFC) ratios are between inland and coastal alleles in the parental (y-axis) and F1 (x-axis) genetic backgrounds. C) The frequency of significant model effects among expressed genes. D) The frequency of expression heterosis in F1 plants at the inland and coastal field sites. LP, low parent; HP, high parent. E-G) Reaction norms of expression at the two field sites for three candidate adaptation genes. Red, inland parent; Blue, coastal parent; Pink, F1 .