The Gene Regulatory Landscape of Local Adaptation

Billie A. Gould[a,*], Yani Chen[a], David B. Lowry[a,b,c]

[a]Department of Plant Biology, Michigan State University, East Lansing, MI, USA

[b]Program in Ecology, Evolutionary Biology and Behavior, Michigan State University, East

Lansing, MI 48824, USA

[c]Plant Resilience Institute, Michigan State University, East Lansing, MI 48824, USA

Local adaptation is a key driver of ecological specialization. Despite its importance, the

evolution of gene regulatory divergence among locally-adapted populations is poorly

understood. Here, we evaluate allele-specific gene expression responses for locally-adapted

coastal and inland accessions of the plant *Mimulus guttatus* in a field reciprocal transplant

experiment. We found 18% of transcripts were characterized by habitat-specific differential

expression, a pattern consistent with the minimization of fitness trade-offs through expression

plasticity. *Cis*-regulatory variation was pervasive, affecting 92% of differentially expressed

genes. Chromosomal inversions were enriched for genes with expression divergence between

ecotypes in a habitat-dependent manner, and gene expression heterosis was prevalent across

habitats. These results provide multiple new insights into the evolution of transcriptome-wide

gene regulatory divergence and plasticity among locally adapted populations.

One of the most important forces driving the evolution ecological specialization and the origin of

biodiversity on Earth is local adaptation (*1-3*). Local adaptation is characterized by reciprocal

home-site advantage, whereby populations perform best in their home habitat while performing

poorly in foreign habitats (*4-7*). Some case-studies, such as the evolution of beak size in

Darwin's Finches (*8*) and coat color of Beach Mice (*6*), suggest that local adaptation should

involve fitness trade-offs at the level of individual genes. However, studies that have combined

reciprocal transplant field experiments with quantitative trait locus (QTL) mapping have

observed that trade-offs at the individual locus level are rare (*7*). Despite recent progress, we still

know little about how transcriptome-wide gene regulatory divergence among locally adapted

populations contributes to fitness trade-offs in nature. To understand this important evolutionary

process, it is crucial to determine the prevalence of genotype x environment (GxE) interactions

affecting the expression of native and non-native alleles across the divergent habitats,

characterize the relative contributions of *cis* and/or *trans* regulatory variation to those

interactions, and establish the role of genome structure (e.g. inversions) in the evolution of

locally adaptive regulatory variation.

We conducted a field reciprocal transplant experiment in tandem with allele-specific expression

analysis to understand how the individual alleles of locally adapted coastal and inland

populations of the yellow monkeyflower, *Mimulus guttatus*, respond to divergent habitats (Fig.

1). There is a long history of local adaptation research in *M. guttatus.* Across previous reciprocal

transplant field experiments, inland plants have survived to flower at 23.9 times the rate of

coastal plants in inland habitat, while coastal plants have survived at 2.5 times the rate of inland

plants in coastal habitat (*9-11*). Many inland populations have evolved an early flowering annual

life-history to escape the harsh summer drought in inland habitats. In contrast, coastal

populations experience year-round soil moisture due to summer sea fog, but have had to adapt to

toxic oceanic salt spray (*10, 11*). We have previously shown that several regions of the genome,

including a chromosomal inversion polymorphism on chromosome 8, contribute to local

adaptation of coastal and inland populations (*12*). An additional inversion polymorphism on

chromosome 5 also appears to be differentiated between coastal and inland populations (*13*).

To evaluate allele-specific gene expression of native and foreign transplants across divergent

habitats, we planted replicates of an inbred coastal line (SS genotype), an inland line (LL

genotype), and their F1 hybrid (SL genotype) into one coastal and one inland field site in the

spring of 2016 (Fig. 1). We planted both gardens in early March and collected leaf tissue for

RNA sequencing at the end of April, just prior to the summer drought in inland habitat. We

aligned reads from field sampled individuals to reference transcriptomes constructed individually

for both inbred parental lines. To understand the effect of genotype and environment on gene

expression, we partitioned parental expression variation into genotype, site (habitat), and GxE

effects using linear models. We exploited allelic polymorphisms between parental lines to

quantify allele-specific expression of coastal and inland alleles in the F1 hybrids. This allowed us

to partition *cis* and *trans* effects and the interactions of those effects with habitat (site) (*14*).

Overall, we were able to evaluate gene expression for 10,122 transcripts corresponding to 9,326

genes, after transcriptome reconstruction and filtering. Significant expression differences (FDR =

0.05) between the inland and coastal field sites (6170 genes; 66%) and between parental

genotypes (6460 genes; 69%) were common (Table S1, S2). Most of the expression differences

between the coastal and inland lines could be attributed to *cis*-regulatory variation: 92% of DE

transcripts had *cis* regulatory differences, while only 39% had detectable trans effects (Fig. 2A

and 2B). On average, the magnitude of *cis*-regulatory effects (LFC = 0.291) were larger than

trans effects (LFC = 0.107; *P* <0.0001).

Across the transcriptome we found 7700 (76%) transcripts were differentially expressed (DE)

between parental ecotypes in either one or both habitats (Fig. 3). A total of 1809 (18%)

transcripts had significant GxE interactions affecting expression (Fig. 3B). Only 94 transcripts

had crossing reaction norms (green points, Fig. 3). More often, expression differences between

genotypes were significant at one site and non-significant at the other (1137 transcripts, 11%,

blue and red points, Fig 3). When there is a cost to gene expression, GxE interactions may allow

genotypes a fitness advantage in one habitat without incurring a fitness trade-off in alternative

habitats. Similarly, a recent study of corals found that plasticity of gene regulation is important

for local adaptation in the field (*15*).

We used network clustering analysis to examine correlated expression of gene clusters (modules)

within habitats for the coastal and inland lines (cor > 0.75, Fig. S1). The coastal line

preferentially expressed modules related to root and epidermal development at its home site,

while oxidative stress response, chlorophyll metabolism, and circadian rhythm modules were

highly expressed in inland habitat (Table S3). The inland line preferentially expressed gene

clusters related to nucleoside catabolism, sugar and starch metabolism, pigment accumulation,

and photoperiodism at its home site, but had no modules strongly associated with the coastal site.

Chromosomal inversions are thought to act as adaptation supergenes that hold together long

haplotypes containing multiple adaptive polymorphisms through suppressed recombination (*16,*

17). We thus tested the hypothesis that inversions hold together haplotypes of DE genes. There

were significantly more DE genes inside the chromosome 5 inversion than expected by chance

and there were significantly more GxE interactions for genes in this inversion (Tables S1). The

chromosome 8 inversion also had significantly more DE genes but only when expression was

measured at the coastal site (Table S4). The enrichment of gene expression differences is

consistent with patterns found in inversions between chimpanzees and humans (*18*) and

Drosophila (*19*), but contrasts with results for the plant *Boechera stricta* (*20*). Our results are

unique among these studies, as they demonstrate supergene associated expression divergence

that is dependent on habitat context (GxE).

Heterosis (hybrid vigor) is commonly found in crosses among locally-adapted populations and

can potentially facilitate gene flow between those populations (*21, 22*). Heterosis is typically

quantified at the level of an individual trait, including expression traits (15, 23), by comparing

the trait value of hybrids to the trait values of the parents (*23*). In this and a previous study, we

have found robust heterosis for multiple phenotypic traits in F1 hybrids between coastal and

inland lines (Fig. 1F; Lowry et al. 2008). In the current study we also found nearly 73% of DE

transcripts had heterotic (non-additive) expression in the F1 (Fig. 2D, Table S5). The percentage

of transcripts with heterotic expression did not vary strongly by site (Fig 2D), although only 54%

of heterotic genes were shared between sites. Heterosis often causes increased growth and vigor

(21). Consistent with this, heterotic genes were enriched for functional terms associated with

growth and development at both field sites (Table S1).

To identify genes putatively linked to local adaptation, we cross-referenced our results with a

recent genomic outlier analysis, which identified 667 genes with a high level of allelic

differentiation between coastal and inland populations (*13*). We found that the magnitude of

differential expression between lines and between field sites was significantly elevated for

candidate genes with highly differentiated SNPs in their promoter regions (Table S6). We were

able to examine gene expression in seven top candidate genes highlighted by the outlier analysis

and/or previous QTL mapping studies (Fig. 2E-G, Fig. S2). The candidate gene *ABA1* mediates

production of abscisic acid, which controls responses to osmotic stress, including drought (*24*).

An *ABA1* homolog, Migut.H00431, is located within the chromosome 8 inversion and may

contribute to inland drought adaptation. *ABA1* was always more highly expressed in the inland

line, and also upregulated in both lines at the inland field site (Fig. 2E). Previously, we showed

that coastal *M. guttatus* populations have evolved leaf tissue tolerance to oceanic salt spray (*14*).

Leaf salt tolerance is generally mediated by a Na^+/H^+ antiporter (*SOS1*), which is activated by a

calcium sensor (*CBL10*; *25*). Both *SOS1* (Migut.E00570) and *CBL10* (Migut.A00138) were more

highly expressed in the coastal line than the inland line across habitats (Fig. 2G, F), a pattern

consistent with elevated salt tolerance of coastal populations (14). However, because expression

of *SOS1* does not appear to be plastic in either ecotype it may be a major contributor to the

fitness trade-offs observed between habitats (Fig. 2F).

In summary, our results demonstrated multiple key patterns regarding the relationship between

gene expression and the evolution of local adaptation. Differential expression is substantial

between locally-adapted populations in nature and the vast majority of that differential

expression is due to *cis*-regulatory divergence. Chromosomal inversions appear to act as

supergenes by holding together haplotypes of DE genes, but this pattern depends on habitat

context. Heterosis of gene expression is pervasive and heterotic genes are enriched for growth

and developmental functions. Expression plasticity of alleles across habitats (GxE) appears to be

common and may minimize fitness trade-offs at individual loci that contribute to the overall

pattern of local adaptation.

REFERENCES AND NOTES

1. J. Clausen (Cornell University Press, Ithaca, NY, 1951).

2. P. F. Colosimo *et al.*, *Science* **307**, 1928-1933 (2005).

3. J. M. Sobel, G. F. Chen, L. R. Watt, D. W. Schemske, *Evolution* **64**, 295-315 (2010).

4. T. J. Kawecki, D. Ebert, *Ecol. Lett.* **7**, 1225-1241 (2004).

5. J. Hereford, *Am. Nat.* **173**, 579-588 (2009).

6. H. E. Hoekstra, R. J. Hirschmann, R. A. Bundey, P.A. Insel, J. P. Crossland, *Science* **313**, 101-

104 (2006).

7. S. Wadgymar *et al.*, *Methods Ecol. Evol. In press* (2017).

8. A. Abzhanov, M. Protas, B. R. Grant, P. R. Grant, C. J. Tabin, *Science* **305**, 1462-1465

(2004).

9. M. C. Hall, J. H. Willis, *Evolution* **60**, 2466-2477 (2006).

10. D. B. Lowry, R. C. Rockwood, J. H. Willis, *Evolution* **62**, 2196-2214 (2008).

11. D. B. Lowry, M. C. Hall, D. E. Salt, J. H. Willis, *New Phytol.* **183**, 776-788 (2009).

12. D. B. Lowry, J. H. Willis, *PLOS Biol.* **8**, e1000500 (2010).

13. B. A. Gould, Y. Chen, D. B. Lowry, *Mol. Ecol.* **26**, 163-177 (2017).

14. J. T. Lovell *et al.*, *Genome Res.* **26**, 510-518 (2016).

15. C. D. Kenkel, M. V. Matz, *Nat. Ecol. Evol.* **1**, 0014 (2016).

16. M. Kirkpatrick, *PLOS Biol.* **8**, e1000501 (2010).

17. M. Joron *et al. Nature* **477**, 203-206 (2011).

18. T. Marquès-Bonet *et al.*, *Trends. Genet.* **20**, 524-529 (2004).

19. Z. L. Fuller, G. D. Haynes, S. Richards, S. W. Schaeffer, *Genetics* **204**, 287-301 (2016).

20. C. R. Lee *et al.*, *Nat. Ecol. Evol.* **1**, 0119 (2017).

21. D. Charlesworth, J. H. Willis, *Nat. Rev. Genet.* **10**, 783-796 (2009).

22. E. Baack, M. C. Melo, L. H. Rieseberg, D. Ortiz-Barrientos, *New Phytol.* **207**, 968-984

(2016).

23. N. M. Springer, R. M. Stupar, *Plant Cell* **19**, 2391-2402 (2007).

24. L. Xiong, H. Lee, M. Ishitani, J. K. Zhu, *J. Biol. Chem.* **277**, 8588–8596 (2002).

25. H. Ji *et al.*, *Mol. Plant.* **6**, 275-286 (2013).

ACKNOWLEDGMENTS

We thank D. Des Marais, J. Tung, J. Friedman, R. Guerrero for comments that helped to improve

our manuscript. We thank K. Childs for guidance with expression analyses, B. Blackman and E.

Patterson for starting seeds for us at UC Berkeley, and D. Popovic for assistance in field

planting. The Bodega Marine Reserve and the Pepperwood Preserve kindly allowed us to

conduct this reciprocal transplant experiment. Funding was provided by Michigan State

University. Sequence data are deposited in the Short Read Archive (XX). The authors declare no

conflicts of interest.

SUPPLEMENTARY MATERIALS

Materials and Methods

Figs. S1 to S9

Tables S1 to S6

References (26-46)

Fig. 1 A) Map of Northern California with the location of the two field transplant sites. Colors represent mean temperature of the warmest quarter in degrees C x 10. Inset: left, coastal ecotype; right, inland ecotype. B) The coastal field site. C) The inland field site. D) SS parental gene expression at two field sites. E) LL parental gene expression at two field sites.F) Survival of parental and F1 (SL) plants at the two field sites.

Fig.2. *Cis* and *trans* regulated transcripts detected at the coastal (A) and inland (B) field sites. Average log2-fold change (LFC) ratios are between inland and coastal alleles in the parental (y-axis) and F1 (x-axis) genetic backgrounds. C) The frequency of significant model effects among expressed genes. D) The frequency of expression heterosis in F1 plants at the inland and coastal field sites. LP, low parent; HP, high parent. E-G) Reaction norms of expression at the two field sites for three candidate adaptation genes. Red, inland parent; Blue, coastal parent; Pink, F1 .

Fig. 3 A) Average log2 fold-change (LFC) differences between parental genotypes for expressed transcripts at two field sites. B) LFC differences only for transcripts with significant GxE interactions. C) Expression reaction norms corresponding to colored points in A and B. Row 3 shows the sign of the allelic effect of the L (inland) allele in the coastal and inland habitats. Row 4 lists the number of transcripts with significant allelic affects within sites in the specified direction. Row 5 lists the number of transcripts with significant GxE.