Design Principle of Lysis/Lysogeny Decision vis-a-vis Multiplicity of Infection

Dinkar Wadhwa*

Department of Chemical Engineering

Indian Institute of Technology Bombay, Mumbai - 400 076, India

dinkar.wadhwa84@gmail.com

*The author is not currently affiliated with the institution.

Abstract

Bacteriophage lambda possesses dual strategy of replication. Upon infecting its host, *Escherichia coli*, it can either choose lytic pathway, in which the host undergoes lysis, releasing hundreds of progeny viruses, or opt for lysogeny, in which the viral genome exists as part of bacterial chromosome known as prophage. Classic and molecular studies have shown that the lysis/lysogeny decision depends upon the number of coinfecting phages, viz. the multiplicity of infection (MoI): lysis at low MoI; lysogeny at high MoI. Here, by constructing an expression for quality of the lysis/lysogeny minimalist two-protein switch which, beside another thing, demands high equilibrium concentration of Cro-like protein (Lyt) and low equilibrium concentration of CI-like protein (Lys) - that is, lytic development - at MoI of 1, and vice versa - that is, lysogeny development - at MoI of 2, I demonstrate that positive feedback loop formed by activation of *cI*'s transcription by its own product in a cooperative manner underlies the switch's design. The minimalist two-protein model, in which Lys performs exactly the same function as CI does in lambda phage's genetic regulatory network (GRN), is justified by showing its analogy with the GRN responsible for lysis/lysogeny decision. Existence of another stable state at MoI of 1 is argued to be responsible for lysogen stability. Further, by comparing the minimalist model and its variants, possessing the positive feedback loop, with other models, without having the positive feedback loop, such as the mutual repression model, it is shown why lysis/lysogeny switch involving positive autoregulation of *cI* is evolved instead of one without it. A three-protein simplified version of lambda switch is shown to be equivalent to a close variant of the two-protein minimalist switch. Finally, only a fraction of parameter sets that produced switch deterministically were able to do so, if at all, under stochastic simulations more than 95% of the time. Additionally, another stable state at MoI of 1 was not found during stochastic simulation.

Keywords: Bacteriophage λ, switch, positive feedback, bistablity

Introduction

Virulent bacteriophages possesses only one method of replication; that is, lytic strategy. However,
other bacteriophages have a dual perpetuation strategy, viz. lytic and lysogeny. In lytic strategy,
phage injects its genetic material into the host bacterium, viral genes are transcribed, m-RNAs,
thus produced, are translated, and phage's genetic material is replicated. Finally, the host
bacterium undergoes lysis, releasing progeny particles. In lysogeny, lytic pathway is repressed,
the viral genome is integrated into that of the host bacterium, and thus, it exists in a latent
form known as prophage. As the teleological explanation goes, lytic strategy leads to fast
multiplication, but its risky, as viral progenies have to find new hosts which don't already
contain lysogenized phages. On the other hand, a lysogenized phage replicates along with its
host, and therefore, reproduces by a slower process as compared to lytic strategy, but this way
phage safeguards its survival. Should a phage infect a bacterium containing lysogenized phage,
lambda repressors (CI) present in the cytosol will not allow expression from *pR*. Thus, the
newly entered phage would remain inert and, ultimately, get digested by the host's nucleases.

Classic [1] and molecular studies [2] have shown that the lysis/lysogeny decision depends
upon MoI. Avlund et al. analysed [3] Kourilsky's data [1,4] and determined the probability of
lysogeny at MoI of 1 to be almost zero, at MoI of 2 to be around 0.6960, and at all higher
MoIs to be around 0.9886. This ability of phage to choose between lysis and lysogeny based
upon multiplicity of infection is but a form of quorum sensing occurring inside a bacterium.
As described in sections below, a minimalist two-protein model, which was analogous to
lambda's GRN, and many other models were constructed. The models were evaluated on the
quality of switch they generated, by solving their defining equations using parameters, which
were searched in two steps (see Methods), and few sets of Hill coefficients. It is shown that
positive feedback loop formed by CI activating transcription of its own gene is the essence of
lysis/lysogeny switch's model. Lastly, a three-protein simplified version of lambda switch is
constructed in which the roles of Lyt and Lys are identical to those of Cro and CI in the latter,
respectively, and the function of CII-like protein is fairly similar to that of CII in the latter.

Result and discussion

Minimalist two-protein lysis/lysogeny switch

The promoter of *lyt* gene is constitutive; whereas, that of *lys* gene is positively regulated as they are in lambda phage's GRN. The role of Lys in the minimalist two-protein model; that is, binding cooperatively to the intergenic region, activating transcription of its own gene, and inhibiting transcription of *lyt* gene, is identical to that of CI in lambda phage's GRN. The role of Lyt was conceptualized from first principle in the following way. At MoI of 2, equilibrium concentrations of Lyt and Lys should be much lower and higher, respectively, as compared to those at MoI of 1. However, if Lyt did not bind to *lys* promoter, assuming no basal expression of *lys* (which is weak promoter anyway), equilibrium concentration of Lyt at MoI 2 would be even higher, let alone much lower, than that at MoI of 1. And equilibrium concentration of Lys would be very low, instead of being high enough to repress *lyt*, at MoI of 2. Since the only protein present to actuate any process is Lyt, it was argued that Lyt should engender lysogeny and inhibit lytic pathway at MoI of 2.

Thus, Lyt activates transcription of *lys* (whose product causes lysogeny development), represses transcription of its own gene, thereby suppressing lytic development (though, as shown below, the last interaction is dispensable), and activates imaginary downstream pathway which leads to lytic development. This seemingly paradoxical role of Lyt, as explained below, is due to it being proxy for CII, which causes lysogeny, and anti-termination factor Q, which enables transcription of lytic genes. The positive feedback loop constituted by transcriptional activation of *lys* by its own protein causes Lys to accumulate to low concentration at MoI of 1 and high concentration at MoI of 2. Thus, at MoI of 1 Lyt's equilibrium concentration is high because it is constitutively produced and Lys' equilibrium concentration is not high enough to repress its production. On the other hand, at MoI of 2 Lyt's equilibrium concentration is low because of repression by Lys, which is present in high concentration.

GRN underlying lysis/lysogeny decision is much more complex than the minimalist two-protein

model proposed here, because MoI is but one of many signals taken into account by the

phage to decide between lysis and lysogeny. Since the expression for quality of lysis/lysogeny

switch (the switching quotient) takes equilibrium values into account, the values of degradation

constants of X (concentration of Lyt) and Y (concentration of Lys), viz. k_2 and k_5, respectively,

can be subsumed into k_1, k_3, and k_4. Hence, they are taken to be unity for all two-protein

models. This model would henceforth be referred to as 1A_Lyt_Lys.

1A_Cro_CI:

$$\frac{dX}{dt} = \frac{mk_1}{1 + \dfrac{X^a}{K_{D1}} + \dfrac{Y^b}{K_{D2}}} - k_2 X \tag{1}$$

$$\frac{dY}{dt} = \frac{m(k_3 \dfrac{X^a}{K_{D1}} + k_4 \dfrac{Y^b}{K_{D2}})}{1 + \dfrac{X^a}{K_{D1}} + \dfrac{Y^b}{K_{D2}}} - k_5 Y \tag{2}$$

96 where, m is multiplicity of infection, k_1 is basal expression rate of *lys*, k_3 and k_4 are rate

97 constants for transcriptional activation of *lys* by Lyt and Lys, respectively, K_{D1} and K_{D2} are

98 the "combined" dissociation constants of Lyt and Lys, respectively (see Methods). In those

99 models where *lys* has basal expression, k_3 represents basal expression rate. Exponents a and b

100 are Hill coefficients for binding of Lyt and Lys, respectively.

Analogy between the minimalist two-protein model (1A_Lyt_Lys) and lambda phages GRN

103 Upon infection, RNA polymerase transcribes from the constitutive promoters, *pL* and *pR*, till it

104 encounters transcription terminators *tL1* and *tR1*, respectively. *N* and *cro* genes are transcribed

105 by *pL* and *pR*, respectively. The product of *N* is an anti-termination factor that modifies

106 subsequent RNAPs initiating at *pL* and *pR* so that they move past their respective terminators

107 and transcribe *cIII* and *cII* genes, respectively. Such an RNAP from *pR* is also able to transcribe

Figure 1: Various two-protein models, and three-protein model. (**A**) The minimalist model or 1A_Lyt_Lys. (**B**) Previous model with self-repression of *lyt* removed or 1B_Lyt_Lys. (**C**) Mutual repression or 2_Lyt_Lys. (**D**) 3_Lyt_Lys. (**E**) 4_Lyt_Lys. (**F**) 5_Lyt_Lys. (**G**) 6_Lyt_Lys. (**H**) A three-protein simplified version of lambda switch or Lyt_Lys_CII. Lower arrowhead represents basal expression.

through another terminator, *tR2*, present upstream of gene *Q* (see Figure 2). Up to this point,

the pathway for lytic and lysogeny are identical. Lytic pathway is chosen when the extended

transcription from *pR* also causes gene *Q* to be transcribed. Q, being an anti-termination factor,

causes transcription of *pR'* to not terminate, as it would otherwise do, at *tR'*, which is present

at about 200 bases away from the beginning, thereby allowing transcription of the lytic genes

downstream of *Q*. Once this happens, the cell is committed to lysis. CIII protein has an indirect

role in establishing lysogeny. It prevents the degradation of CII by inhibiting bacterial protease

HflB [5,6]. As the current paper focuses on the design principle of lysis/lysogeny switch, the

(indrect) role of cIII will not be taken into consideration.

In lambda's GRN, *cII* and *Q* are under the control of promoter *pR*. Since in 1A_Lyt_Lys

lyt is transcribed from *pR*, Lyt protein should be functionally equivalent to CII and Q. That

is, on the whole, CII and Q should carry out three actions: activate transcription from *lys*,

inhibit transcription from *lyt* gene, and engender lytic development. When CII accumulates

in sufficient concentration, it activates transcription from three promoters: *pI*, *pRE*, and *pAQ*

[10,11]. Promoter *pI* transcribes *int* gene, required for the integration of phage genome into

that of the host bacterium. Transcript produced from *pRE* contains orf for *cI*; hence, activation

of this promoter leads to production of CI. Thus, the action of CII on promoters *pI* and *pRE*

is functionally equivalent to Lyt protein activating transcription of *lys*. Notably, while the role

of Cro in lambda's GRN is to inhibit the expression of *lys*, Cro-like protein (Lyt) activates the

expression of *lys* in the 1A_Lyt_Lys.

CII inhibits lytic development by activating transcription from *pAQ*, which is located

within *Q* gene in the opposite polarity. The transcript, thus produced, being antisense to (a part

of) *Q* mRNA hybridizes with the latter, thereby preventing the translation of *Q* m-RNA, which

is essential for lytic development [2]. Thus, the action of CII on promoter *pAQ* is functionally

equivalent to Lyt protein inhibiting transcription of its own gene. If CII is not produced in

sufficient amount, *Q* m-RNA is translated and anti-terminator Q, thus produced, causes lysis.

Figure 2: GRN and transcription map of lambda (adapted from Figure 1 of [8]). Transcripts that are produced earliest, viz. from *pL* and *pR* promoters, are depicted as green arrows. The late transcript, viz. from *pR'*, is a black arrow. Transcripts from CII-activated promoters, viz. *pI*, *pRE*, and *pAQ*, are shown as blue arrows. Transcript from *pRM*, which is activated by CI, is shown as red arrow. Transcription terminators, namely *tL1*, *tR1*, and *tR2*, are depicted in red.

Variants of 1A_Lyt_Lys and mutual repression model

In order to better demonstrate that the positive feedback underlies lysis/lysogeny switch, I considered variants of 1A_Lyt_Lys, mutual repression model, which doesn't have positive feedback loop, and its variants, and a model having the features of 1A_Lyt_Lys and mutual repression model. Since two features, viz. constitutive expression of *lyt* and its inhibition by Lys, are common, they would not be mentioned in the description of the models below. Since *cI* gene is positively regulated in lambda's GRN, *lys* has to have either basal expression or be activated by Lyt. All of these models can be categorized in terms of three factors, as shown in the Table 1. First column shows whether *lys* possesses basal expression or is activated by Lyt. Second column shows if positive feedback, constituted by transcriptional activation of *lys* by its own product, is present. Third column shows if inhibition of *lys* by Lyt is present. Inhibition of *lys* by Lyt can only be present when *lys* possesses basal expression. Thus, for *lys* having basal expression, there are four models; and where it gets activated by Lyt, there are two models.

1B_Lyt_Lys: This model differs from 1A_Lyt_Lys only in not having self-inhibition of Lyt. The inhibition of *lyt*, required at MoI of 2, by its own product is dispensable, as Lys performs the

Table 1: Classification of additional two-protein models.

Model	Basal expression of *lys*/ Activation of *lys* by Lyt	Activation of *lys* by Lys	Inhibition of *lys* by Lyt
1B_Lyt_Lys	Activation	Yes	N/A
5_Lyt_Lys	Activation	No	N/A
3_Lyt_Lys	Basal	Yes	No
6_Lyt_Lys	Basal	Yes	Yes
4_Lyt_Lys	Basal	No	No
2_Lyt_Lys	Basal	No	Yes

150 same function, and more so, because at MoI of 2 Lyt's concentration is required to be much

151 lower than that of Lys in order for switch to be of good quality. In terms of lambda's GRN, this

152 would mean CI, instead of CII, activating transcription from *pAQ*.

$$\frac{dX}{dt} = \frac{mk_1(1 + \frac{X^a}{KD_1})}{1 + \frac{X^a}{K_{D1}} + \frac{Y^b}{K_{D2}}} - k_2 X \tag{3}$$

$$\frac{dY}{dt} = \frac{m(k_3 \frac{X^a}{K_{D1}} + k_4 \frac{Y^b}{K_{D2}})}{1 + \frac{X^a}{K_{D1}} + \frac{Y^b}{K_{D2}}} - k_5 Y \tag{4}$$

153 **2_Lyt_Lys** (Mutual repression): Lyt represses *lys*, which has basal expression.

$$\frac{dX}{dt} = \frac{mk_1(1 + \frac{X^a}{K_{D1}})}{1 + \frac{X^a}{K_{D1}} + \frac{Y^b}{K_{D2}}} - k_2 X \tag{5}$$

$$\frac{dY}{dt} = \frac{mk_3(1 + \frac{Y^b}{K_{D2}})}{1 + \frac{X^a}{K_{D1}} + \frac{Y^b}{K_{D2}}} - k_5 Y \tag{6}$$

154 **3_Lyt_Lys**: *lys* has basal expression and is activated by Lys cooperatively.

$$\frac{dX}{dt} = \frac{mk_1}{1 + \frac{Y^b}{K_{D2}}} - k_2 X \tag{7}$$

$$\frac{dY}{dt} = \frac{m(k_3 + k_4 \frac{Y^b}{K_{D2}})}{1 + \frac{Y^b}{K_{D2}}} - k_5 Y \tag{8}$$

155 **4_Lyt_Lys**: *lys* has basal expression.

$$\frac{dX}{dt} = \frac{mk_1}{1 + \frac{Y^b}{K_{D2}}} - k_2 X \tag{9}$$

$$\frac{dY}{dt} = mk_3 - k_5 Y \tag{10}$$

156 **5_Lyt_Lys**: *lys* is activated by Lyt.

$$\frac{dX}{dt} = \frac{mk_1(1 + \frac{X^a}{K_{D1}})}{1 + \frac{X^a}{K_{D1}} + \frac{Y^b}{K_{D2}}} - k_2 X \tag{11}$$

$$\frac{dY}{dt} = \frac{mk_3 \frac{X^a}{K_{D1}}}{1 + \frac{X^a}{K_{D1}} + \frac{Y^b}{K_{D2}}} - k_5 Y \tag{12}$$

157 **6_Lyt_Lys**: *lys* has basal expression, is activated by Lys, and inhibited by Lyt.

$$\frac{dX}{dt} = \frac{mk_1(1 + \frac{X^a}{K_{D1}})}{1 + \frac{X^a}{K_{D1}} + \frac{Y^b}{K_{D2}}} - k_2 X \tag{13}$$

$$\frac{dY}{dt} = \frac{m(k_3 + k_4 \frac{Y^b}{K_{D2}})}{1 + \frac{X^a}{K_{D1}} + \frac{Y^b}{K_{D2}}} - k_5 Y \qquad (14)$$

Deterministic simulation

Since Cro forms dimer, Hill coefficient for Lyt's binding is considered to be 2; whereas, since CI forms tetramer, Hill coefficient for Lys' binding was taken to be 4. However, in the interest of completeness, another set of Hill coefficients, viz. a=2, b=2, was also considered. The rate constants and dissociation constants of equations defining a given model were searched (see Methods) in two stages: order search and linear search (as they are called here). For a given model and set of Hill coefficients (a and b), a set of rate constants and dissociation constants would henceforth be referred to as a parameter set (That is, Hill coefficients are not part of parameter set). Parameter sets were selected on the basis of quality of switch, viz. switch quotient (as it is called here), they generated. Switch quotient was initially considered to be determined by the expression

$$SQ = \frac{(S_1 - S_2)}{S_1}$$

$S_1 = \min\{\text{Lyt at MoI of 1, Lys at MoI of 2}\}$

$S_2 = \max\{\text{Lys at MoI of 1, Lyt at MoI of 2}\}$

The expression, however, selected parameter sets which gave unequal equilibrium values of Lyt at MoI of 1 and Lys at MoI of 2. From the perspective of simplicity, I believe that the difference between the two should be minimal; therefore, the previous expression is multiplied by ratio of S_1 to S_3 in order to penalize the difference between S_3 and S_1.

$$SQ = \frac{(S_1 - S_2)}{S_1} \cdot \frac{S_1}{S_3} = \frac{(S_1 - S_2)}{S_3}$$

$S_3 = \max\{\text{Lyt at MoI of 1, Lys at MoI of 2}\}$

This expression (like the older one) varies between 0 and 1. Only those parameter sets were

177 selected whose corresponding switch quotients (SQ) were positive.

Figure 3: Schematic of switch's profile, viz. equilibrium concentrations of Lyt and Lys at the two MoIs.

As Table 2 shows, all of the models possessing the positive feedback loop have average

SQ of more than 0.97 for both sets of Hill coefficients (lowest SQ among all the models in

this category was 0.9270). Mutual repression model for Hill coefficients' set of a=2, b=2 have

average SQ of 0.5283 (highest SQ was 0.6666); and, for that of a=2, b=4 all SQs were more

than 0.9 except for one parameter set, whose SQ was 0.5. 4_Lyt_Lys for Hill coefficients' set

of a=2, b=2 gives SQs of 0.4794 and 0.4707; and, for that of a=2, b=4 both SQs were almost

0.5. Thus, if we exclude 2_Lyt_Lys for Hill coefficients' set of a=2, b=4 from the analysis, the

lowest SQ among models with the positive feedback loop, viz. 1A_Lyt_Lys, 1A_Lyt(1)_Lys,

1B_Lyt_Lys, 3_Lyt_Lys, and 6_Lyt_Lys, was much higher than the highest SQ among models

without it, viz. 2_Lyt_Lys and 4_Lyt_Lys.

In the former set, Lys activating its own gene lets the value of Lys at MoI of 1 to be

disproportionately lower for its desired particular value at MoI of 2. On the other hand,

in 4_Lyt_Lys, since increase in genome copy number leads to proportional increase in the

equilibrium activity of *lys*' promoter, value of Lys at MoI of 1 would be half its value at MoI of

2. However, mutual repression model does generate many parameter sets with SQ greater than

Table 2: Average and SD of SQs under deterministic and stochastic conditions for various thresholds of stochastic success rate.

Model	Deterministic AVG SQ (SD)		Deterministic AVG SQ (SD) (SSR[a] ⩾ 95)		Stochastic AVG SQ (SD) (SSR ⩾ 95)	
	a=2, b=2	a=2, b=4	a=2, b=2	a=2, b=4	a=2, b=2	a=2, b=4
1A_Lyt_Lys	0.9917 (0.0106)	0.9896 (0.0049)	0.9898 (N/A)	0.9905 (0.0043)	0.7997 (N/A)	0.6204 (0.0624)
1A_Lyt(1)_Lys	0.9950 (0.0053)	0.9923 (0.0045)	none	none	none	none
1B_Lyt_Lys	0.9806 (0.0236)	0.9769 (0.0277)	0.9971 (N/A)	0.9270 (N/A)	0.7995 (N/A)	0.7679 (N/A)
2_Lyt_Lys	0.5283 (0.0696)	0.8917 (0.1766)	none	0.5001 (N/A)	none	0.2725 (N/A)
3_Lyt_Lys	0.9938 (0.0078)	0.9873 (0.0157)	none	none	none	none
4_Lyt_Lys	0.4751 (0.0043)	0.4956 (0.0006)	none	0.4956 (0.0006)	none	0.2983 (0.0102)
6_Lyt_Lys	0.9988 (0.0004)	0.9876 (0.0135)	none	none	none	none
Lyt_Lys_CII	0.9855 (0.0155)	N/A	0.9573 (N/A)	N/A	0.7526 (N/A)	N/A
Lyt_Lys_CII(1)	0.9801 (0.0151)	N/A	0.9718 (N/A)	N/A	0.7595 (N/A)	N/A
Lyt(1)_Lys_CII(1)	0.9894 (0.0134)	N/A	none	N/A	none	N/A

[a] SSR = Stochastic Success Rate

0.9 for Hill coefficients' set of a=2, b=4. Since this model exhibits very different behaviour in

the stochastic simulations, it will be discussed further in the section for stochastic simulations.

The model 5_Lyt_Lys did not generate any parameter set. The reason is that in the absence

of the positive feedback loop, *lyt* needs to have strong basal expression in order to sustain high

concentration of Lys, whose gene is activated by Lyt, at MoI of 2. Equivalently, the desired

high concentration of Lyt at MoI of 1, also leads to excessive production of Lys at the same

MoI. Thus, both proteins are present in similar amounts at both MoIs. In hindsight, one notes

that the equations for Lyt and Lys are almost identical for this model.

In order to examine the significance of cooperativity in positive feedback here, another set

of Hill coefficients, viz. a=2, b=1, was also considered for 1A_Lyt_Lys. However, parameter

sets generated by this set gave SQs which were almost equal to zero. For models having

the positive feedback loop, average SQ of parameter sets was very slightly, almost negligibly,

greater for Hill coefficients' set of a=2, b=2 than that for set of a=2, b=4.

Closer to lambda's GRN: the three-protein model

In order to further verify if 1A_Lyt_Lys represents reduced form of lambda's GRN, I consider a

three-protein simplified version of lambda switch and show that it is equivalent to a two-protein

model possessing the positive feedback loop: 1B_Lyt_Lys. A CII-like protein is added to

1A_Lyt_Lys beside extending the role of Lyt. Since genes *lyt* and *cII* are under the control of

same promoter, in order to allow for potentially different rates of translation of their corresponding

cistrons during stochastic simulations, their mRNAs are considered explicitly. The role of Lyt

in this model is identical to that of Cro in lambda phage's GRN. That is, now Lyt represses

transcription of *lys*, in addition to repressing that of its own gene. The role of CII in the

three-protein model is to activate transcription of *lys*. This corresponds to CII's activation of

pRE promoter, leading to synthesis of mRNA which contains orf for *cI*. The three-protein

model considered here is different from that in [7], in which CII activates transcription of *cI*

from a distinct (*pRE*) promoter. Since in the three-protein model, CII has to compete with Lyt,

Figure 4: Deterministic and stochastic simulations of the minimalist two-protein model (1A_Lyt_Lys) and three-protein model (Lyt_Lys_CII). Lyt, Lys, and CII are represented by green, red, and blue, respectively. For deterministic simulations, concentrations of proteins at MoI of 1 and 2 are depicted by solid curve and dashed curve, respectively. For stochastic simulations, solid curve and dotted curve, respectively, represent average and standard deviation of number of protein molecules from 500 simulations. For a given model, the parameter set which had maximum stochastic success rate was used for simulation. The stochastic simulation trajectories shown here are qualitatively similar to those of all other models for parameter sets with high stochastic success rate; whereas, the deterministic simulation trajectories were so, irrespective of stochastic success rate. In stochastic simulation graphs, the original abscissa, which had unequally spaced time intervals, was converted to one with equally spaced time intervals. Each (arb.) unit of abscissa was divided into 10000 intervals. For the tiny fraction of intervals which still contained more than one event, their last events were defined to be their only events. (**A**) Deterministic simulations of 1A_Lyt_Lys. At MoI of 2, initially, the concentration of Lyt becomes more than its equilibrium concentration at MoI of 1 but then comes back to very low level. It is due to double initial rate of production of Lyt at MoI of 2 as compared to that at MoI of 1; however, as Lyt's concentration increases, *lys'* transcription becomes stronger, leading to production of Lys, which in turn represses *lyt*. (**B-C**) Stochastic simulations of 1A_Lyt_Lys for MoI of 1 and 2, respectively. (**D**) Deterministic simulations of Lyt_Lys_CII. At MoI of 2, initially, concentrations of Lyt and CII become more than their respective equilibrium concentrations at MoI of 1 but then come back to very low levels. This was also observed for a three-protein model, which is very similar to that of this paper, in a theoretical study [7]. Analogous to the two-protein model, it's due to heightened initial rate of production of CII at MoI of 2 as compared to that at MoI of 1; however, as CII's concentration increases, transcription of *lys* becomes stronger, leading to production of Lys, which represses *lyt* and *cI*. (**E-F**) Stochastic simulations of Lyt_Lys_CII for MoI of 1 and 2, respectively. Bell-shaped curve for CII at MoI of 6 was reported by an experimental study [2].

which represses transcription of *lys*, for binding to the intergenic region, the demonstration of

equivalence of the three-protein model (Lyt_Lys_CII) with 1A_Lyt_Lys, or any of its variants,

gets more challenging. The degradation constants for *xz* (concentration of *lyt-cII* mRNA) ,

X (concentration of Lyt), *Z* (concentration of CII), and *Y* (concentration of Lys), viz. k_6,

k_7, k_9, k_8, respectively, are taken to be unity for the same reason why degradation constants

for two-protein models were set equal to 1. Since for 1A_Lyt_Lys SQs generated by Hill

coefficients' set of a=2, b=2 were as high as SQs generated by that of a=2, b=4, applying

occam's razor, Hill coefficients for binding of Lyt and Lys are taken to be 2 and 2, respectively,

not 2 and 4. Further, taking lead from here, Hill coefficient for CII's binding is considered to

be 2, even though it has been shown to exist as tetramer in solution [14] and in crystallized free

and DNA-bound state [15].

Model equations for three-protein model are as follows.

Transcription of *lyt-cII* genes:
$$\frac{dxz}{dt} = \frac{mk_1(1 + \frac{Z^c}{K_{D3}})}{1 + \frac{X^a}{K_{D1}} + \frac{Y^b}{K_{D2}} + \frac{Z^c}{K_{D3}}} - k_6 xz \qquad (15)$$

235 Translation of *lyt*:
$$\frac{dX}{dt} = k_2 xz - k_7 X \qquad (16)$$

236 Translation of *cII*:
$$\frac{dZ}{dt} = k_4 xz - k_9 Z \qquad (17)$$

237 Production of Lys:
$$\frac{dY}{dt} = \frac{m(k_5 \frac{Y^b}{K_{D2}} + k_3 \frac{Z^c}{K_{D3}})}{1 + \frac{X^a}{K_{D1}} + \frac{Y^b}{K_{D2}} + \frac{Z^c}{K_{D3}}} - k_8 Y \qquad (18)$$

238 where, c is the Hill coefficient of CII's binding, k_1 is basal expression rate of *lyt-cII* genes, K_{D3}

239 is the "combined" dissociation constant of CII (see Methods), k_2 and k_4 are translation rates of

240 *lyt* and *cII*, respectively. k_5 and k_3 are rate constants for transcriptional activation of *lys* by Lys

241 and CII, respectively.

242 Equilibrium values of xz, X, Z, and Y are

$$k_6\overline{xz} = \frac{mk_1(1 + \dfrac{\overline{Z}^a}{K_{D3}})}{1 + \dfrac{\overline{X}^a}{K_{D1}} + \dfrac{\overline{Y}^b}{K_{D2}} + \dfrac{\overline{Z}^a}{K_{D3}}} \tag{19}$$

$$k_7\overline{X} = k_2\overline{xz} \tag{20}$$

$$k_9\overline{Z} = k_4\overline{xz} \tag{21}$$

$$k_8\overline{Y} = \frac{m(k_5\dfrac{\overline{Y}^b}{K_{D2}} + k_3\dfrac{\overline{Z}^a}{K_{D3}})}{1 + \dfrac{\overline{X}^a}{K_{D1}} + \dfrac{\overline{Y}^b}{K_{D2}} + \dfrac{\overline{Z}^a}{K_{D3}}} \tag{22}$$

243 From (20) and (21), it can be seen that equilibrium value of CII is in constant proportion to that
244 of Lyt. Hence, CII can be written in terms of Lyt

$$\overline{Z} = p\overline{X} \tag{23}$$

245 where

$$p = \frac{k_4k_7}{k_2k_9}$$

246 Using (20) and (23), (19) and (22) can be written as

$$\overline{X} = \frac{m\dfrac{k_1k_2}{k_6k_7}(1 + \dfrac{(p\overline{X})^a}{K_{D3}})}{1 + \overline{X}^a(\dfrac{1}{K_{D1}} + \dfrac{p^a}{K_{D3}}) + \dfrac{\overline{Y}^b}{K_{D2}}} \tag{24}$$

$$\overline{Y} = \frac{m\dfrac{1}{k_8}\left(k_5\dfrac{\overline{Y}^b}{K_{D2}} + k_3\dfrac{(p\overline{X})^a}{K_{D3}}\right)}{1 + \overline{X}^a\left(\dfrac{1}{K_{D1}} + \dfrac{p^a}{K_{D3}}\right) + \dfrac{\overline{Y}^b}{K_{D2}}} \tag{25}$$

The equivalence of equations (24) and (25) to the defining equations of 1B_Lyt_Lys which

have reached equilibrium validates two-protein model. Two-protein model being sufficient for

producing lysis/lysogeny switch constitutes an argument that *cro* in lambda's GRN is expendable.

Mathematically, the reason for Cro being expendable lies in its equilibrium concentration being

proportional to that of CII.

Kobiler et al. [2] showed that infection with lambda lacking cro gene (λcro⁻) leads to

production of CII to level sufficient to cause lysogeny even at MoI of 1. This, however, does

not mean that Cro, per se, is required to engender lytic development. Cro represses *pL* and

pR by fourfold and twofold, respectively [12]. Thus, the absence of Cro increases the level

of CII in two ways: first, by allowing transcription of *cII*, which is under the control of *pR*,

and *cIII*, which is under the control of *pL* and whose product prevents degradation of CII by

protease HflB. In the wild type strain, parameters associated with transcription rates of *cII* and

cIII, translation and degradation rates of their respective mRNAs, and degradation rates of CII

and CIII are such that enough CII is produced, despite Cro's repression of *pL* and *pR*, at higher

MoIs so as to sufficiently activate *pRE* promoter, leading to production of CI to level which

is enough to cause lysogeny. However, when *cro* is deleted, CI produced even at MoI of 1 is

enough to engender lysogeny. With appropriate changes in the aforementioned parameters, it

would be possible to model λcro⁻ strain which behaves like its wild type counterpart.

As stated above, there are experimental evidences for CII present as tetramer in solution

[14] and in crystallized free and DNA-bound state [15]. Additionally, as Figure 4 in [10] shows,

the binding curve of CII to *pAQ* has appreciable lag phase, indicating that it binds as a multimer.

However, Figure 2c in [2] shows that curve of *pRE*'s activity with respect to CII levels is not

sigmoidal as expected from multimeric binding, but hyperbolic as seen in monomeric binding.

Therefore, another model was considered where Hill coefficient for CII binding was taken to

be 1 (Lyt_Lys_CII(1)). Additionally, one more model was considered where Hill coefficient

for Lyt too was taken to be 1 (Lyt(1)_Lys_CII(1)). This made the current author go back to

two-protein models and consider 1A_Lyt_Lys model too with Hill coefficients' set of a=1, b=2

and a=1, b=4, named 1A_Lyt(1)_Lys. SQs generated by all new variants were similar in values

to those generated from their counterparts, where Hill coefficient of either Lyt or CII, or both,

were taken to be 2. Specifically, for 1A_Lyt(1)_Lys all SQs were more than 0.98 for both sets

of Hill coefficients. For all of the three protein models, all SQs were greater than 0.95. Just like

the Hill coefficients' set of a=2, b=1, parameter sets generated by the set of a=1, b=1 gave SQs

which were almost equal to zero.

Stochastic simulation

Since gene expression is stochastic [17,18], the true validity of results obtained in the deterministic

simulations lie in their being replicated in the stochastic simulations. Thus, stochastic simulations

were performed, using Gillespie algorithm [19], for parameter sets obtained in the deterministic

simulations.

For both two-protein and three-protein models, for any given parameter set, SQ generated

in the stochastic simulation, or stochastic switch quotient (SSQ), was less than its deterministic

counterpart, or deterministic switch quotient (DSQ). No parameter set was able to produce

switch in every run during stochastic simulation. That is because either the SSQ was negative

($S_1 < S_2$) or, rarely, S_3 was zero. Percentage of runs that produce finite, positive SQs during

stochastic simulation for a given parameter set and set of Hill coefficients would henceforth be

referred to as stochastic success rate.

An interesting property was observed for mutual repression model for Hill coefficients' set

of a=2, b=4. It was the only set of Hill coefficients for any model lacking the positive feedback

that produced a DSQ more than 0.9 (highest SQ for the same model for Hill coefficients' set of

a=2, b=2 was 0.6666). As aforementioned, all of the parameter sets for Hill coefficients' set of

a=2, b=4 produced DSQ of more than 0.9 except one, whose DSQ was 0.5. Notably, this is the

Table 3: Number of parameter sets for various ranges of stochastic success rate.

Model	SSR[a] ⩾ 95		95 > SSR ⩾ 90		90 > SSR ⩾ 80		Total no. of parameter sets	
	a=2 b=2	a=2 b=4	a=2 b=2	a=2 b=4	a=2 b=2	a=2 b=4	a=2 b=2	a=2 b=4
1A_Lyt_Lys	1	4	5	3	10	3	21	17
1A_Lyt(1)_Lys	0	0	1	1	3	9	17	15
1B_Lyt_Lys	1	1	1	5	2	2	11	11
2_Lyt_Lys	0	1	0	0	0	0	6	6
3_Lyt_Lys	0	0	0	0	1	4	8	10
4_Lyt_Lys	0	2	0	0	0	0	2	2
6_Lyt_Lys	0	0	0	0	0	0	12	12
Lyt_Lys_CII	1	N/A	2	N/A	1	N/A	9	N/A
Lyt_Lys_CII(1)	1	N/A	1	N/A	5	N/A	9	N/A
Lyt(1)_Lys_CII(1)	0	N/A	1	N/A	1	N/A	9	N/A

[a] SSR = Stochastic Success Rate

parameter set which had very high stochastic success rate, viz. that of 97%; while, maximum stochastic success rate among other parameter sets was 50%. This peculiar result for mutual repression has been reported earlier also.

Avlund et al. showed that various two-protein models, based upon mutual repression model, which were able to produce switch in a noise-less environment, did not function when noise was introduced [9]. However, additional CII-like protein conferred robustness to noise in 8% of the parameter sets that produced switch deterministically. The different behaviour of mutual repression model in deterministic simulations with respect to stochastic simulations warrants theoretical investigation. Notably, one of their rare two-protein models (i.e., b of Figure 2) which did produce switch even in the presence of noise (though with much lower success as compared to their three-protein models) is model 6_Lyt_Lys in the current paper.

Thus, taking into account the stochastic success rate of at least 95%, two-protein models can be divided into two sets based upon DSQs or SSQs. One set comprises of two models with the positive feedback loop, viz. 1A_Lyt_Lys and 1B_Lyt_Lys, and another without it, viz. 2_Lyt_Lys and 4_Lyt_Lys. The one with the positive feedback loop has appreciably higher DSQs and SSQs than the one without it.

However, for the same stochastic success rate cut-off, the lowest SSQ among parameter

sets with Hill coefficients' set of a=2, b=2 was greater than the highest SSQ among those
with Hill coefficients' set of a=2, b=4, for any given model (data not shown). This trend gets
confirmed if one considered more parameter sets, viz. by relaxing the cut-off of stochastic
success rate from 95% to 90%. The relaxation lets the inclusion of 1A_Lyt(1)_Lys in the
analysis. This result is against one's expectation: since Lys activating transcription of its own
gene in a cooperative manner is crux of the switch, increasing Hill coefficient of Lys should
have, if at all, increased the SSQ. This comparison could not be made in models without the
positive feedback loop because none of their parameter sets with Hill coefficients' set of a=2,
b=2 had stochastic success rate of at least 90%.

Table 4: Maximum stochastic success rate.

Model	Hill coefficients' set	Maximum stochastic success rate
1A_Lyt_Lys	a=2, b=4	96.8
1A_Lyt(1)_Lys	a=1, b=2	91
1B_Lyt_Lys	a=2, b=4	97.2
2_Lyt_Lys	a=2, b=4	97
3_Lyt_Lys	a=2, b=4	87
4_Lyt_Lys	a=2, b=4	98.8
6_Lyt_Lys	a=2, b=4	73
Lyt_Lys_CII	a=2, b=2, c=2	95.5
Lyt_Lys_CII(1)	a=2, b=2, c=1	97
Lyt(1)_Lys_CII(1)	a=1, b=2, c=1	93.5

Bistability at MoI of 1 and lysogen stability

In this study, parameter sets were searched for their ability to cause lysis at MoI of 1 and
lysogeny at MoI of 2. However, if only one of the phage genomes gets integrated into the
bacterial chromosome, it would not be able to maintain lysogeny, and lysis would ensue, if
only one stable state existed at MoI of 1. In the deterministic simulations, all of the two-protein
models possessing the positive feedback exhibited bistability at MoI of 1 for all of the parameter
sets, except one (for 1B_Lyt_Lys). In the other stable state, the concentration of Lyt is almost

Figure 5: Equilibrium values correspond to those parameter sets which gave maximum stochastic success rate for their respective models (see Table 4). (**A**) Deterministic simulations. (**B**) Stochastic simulations. Note how the values of Lys at MoI of 1 and Lyt at MoI of 2 for 2_Lyt_Lys and 4_Lyt_Lys are much higher than those of the any other model.

zero and that of Lys is about half of its concentration at MoI of 2. Arguably, in lambda's system,

the level of Lys in the second stable state would be high enough to maintain lysogeny.

For 4_Lyt_Lys, none of the parameter sets produced bistability at MoI of 1. For 2_Lyt_Lys,

for Hill coefficients set of a=2, b=2 one parameter set generated bistability at MoI of 1, but its

stochastic success rate was just 7.6% (Bistability exists for two more parameter sets, but their

second stable states are at very high values of Lyt (>50) and very low values of Lys (<2); hence,

inconsequential for lysogeny maintenance, and in any case, never reached by the phase point).

For Hill coefficients set of a=2, b=4, the only parameter set which did not exhibit bistability

at MoI of 1 had stochastic success rate of 97%, while maximum stochastic success rate among

other parameter sets was 50% (as aforementioned in the section for stochastic simulations). All

of the three-protein models exhibited bistability at MoI of 1. The Lyt and Lys values of second

stable states at MoI of 1 in three-protein models are about same as those of second stable states

in two-protein models at the said MoI.

DNA between OL and OR sites forms a loop that has been shown to be important for the

stable maintenance of lysogeny [12]. The loop forms due to interaction between CI dimers

bound at OL1 and OL2 with those bound at OR1 and OR2 [13]. Therefore, the contribution of

OL-CI-OR complex to production of CI would be represented by adding a term proportional to

[CI], raised to the power 8, to numerator and denominator. Since bistability at MoI of 1 in the

two-protein models is the consequence of *lys*' transcription getting activated by its own product

in a cooperative manner (i.e., by the binding of Lys dimer), in lambda's GRN, activation of *cI*'s

promoter present in a looped DNA, stabilized by CI octamer, would either generate bistability

or contribute to already existing bistability due to two CI dimers activating the transcription of

cI. Thus, it is reasonable to propose that the role of OL_CI_OR loop formation is to produce or

strengthen bistability at MoI of 1. This argument becomes stronger in the light of the finding

that looping also activates transcription from *pRM* by allowing the α-CTD of RNAP bound at

pRM to contact UP element at OL [16]. In the stochastic simulations, however, none of the

two-protein and three-protein models produced bistability at MoI of 1.

At MoI of 2, only two models, viz. 2_Lyt_Lys and 6_Lyt_Lys, show bistability for about

80% and 60%, respectively, of the parameter sets. Notably, only these two models have Lyt

repressing the transcription of *lys*. Since at the second stable state the concentration of Lyt is

very high and that of Lys is very low, a parameter set would not, if at all, generate switch with

high DSQ if its phase point reached this stable fixed point. Hence, bistability at MoI of 2 is

inconsequential.

Figure 6: (**A-B**)Phase diagram of 1A_Lyt_Lys corrresponding to the parameter set that gave maximum stochastic success rate, at Mo1 of 1 and 2. Green and red full circles are stable fixed points, whereas empty black circle is unstable fixed point. Green stable point is where system reaches when a single phage infects a bacterium. Red stable point is where system reaches when lysogeny is established by two phages, but only one of them gets integrated into the host's genome.

Why positive feedback?

There can be two reasons why lysis/lysogeny switch is based upon the positive feedback: 1) biological properties of the switch, viz. a) highest switch quotient and presence of bistability at MoI of 1, and 2) quickest evolution of such a model. It should be noted, however, that speed of evolution would not matter if evolution is path-independent. That is, it's possible that nature initially evolves a sub-optimal design but which, given enough time, gets superseded by an optimal one.

Switch quotient: As mentioned in the previous sections, SQs generated in the deterministic and the stochastic simulations, respectively, for models possessing the positive feedback are much greater than those of the models lacking positive feedback.

Bistability at MoI of 1: As stated in the last section, for models not possessing the positive feedback loop, no parameter set, if at all, having sufficiently good stochastic success rate generated bistability. If one ignores the possibility of any other mechanism generating bistability, such as the formation of OL-CI-OR complex, this reason alone is sufficient for nature to choose models which possess the positive feedback loop over those which do not.

Speed of evolution: Even though the maximum stochastic success rate is very low for 3_Lyt_Lys and (especially) 6_Lyt_Lys, they are still compared with 4_Lyt_Lys and 2_Lyt_Lys, respectively, as these two are the only pairs within which mathematical comparison with regard to the positive feedback loop is possible. 2_Lyt_Lys and 4_Lyt_Lys differ from 6_Lyt_Lys and 3_Lyt_Lys, respectively, only in not having the positive feedback loop. Thus, model equations of former two models differ from those of latter two only in the dynamics of Lys. In models with the positive feedback loop, the term representing binding of Lys to the intergenic region (i.e., Y^b/K_{D2}) is multiplied by rate constant for transcriptional activation of *lys* by Lys, k_4. On the other hand, in models without the positive feedback loop Y^b/K_{D2} is multiplied by k_3, the basal expression rate of *lys*. Thus, 2_Lyt_Lys and 4_Lyt_Lys can be thought of as being equivalent to 6_Lyt_Lys and 3_Lyt_Lys, respectively, whose k_4 is equal to k_3. That is, the former two models are those latter two models, respectively, whose rate constant for transcriptional activation of

lys by Lys is equal to the basal expression rate of *lys*. This constrain of having $k_3 = k_4$ reduces

the potential parameter space for 2_Lyt_Lys and 4_Lyt_Lys by one dimension. Hence, the two

parameters being independent in 3_Lyt_Lys and 6_Lyt_Lys makes nature more likely to discover

them. This explains why 2_Lyt_Lys (11, 11) and 4_Lyt_Lys (2, 2) produced fewer parameter sets

than 6_Lyt_Lys (16, 16) and 3_Lyt_Lys (11, 13), respectively, for both sets of Hill coefficients

during the order search (as shown in the parenthesis).

Now, qualitative equivalence of 3_Lyt_Lys and 6_Lyt_Lys with 1B_Lyt_Lys, which is equivalent

to 1A_Lyt_Lys, is shown. 1B_Lyt_Lys is qualitatively equivalent to 3_Lyt_Lys for the reason that

in the former, transcriptional activation of *lys'* is achieved by binding of Lyt to its promoter;

whereas, in the latter, *lys* possesses basal expression. 6_Lyt_Lys differs from 3_Lyt_Lys in

having Lyt as a repressor of *lys*. This interaction is expendable, as at MoI of 2, concentration of

Lyt is anyway very low, and qualitatively speaking, at MoI of 1 repression of *lys* by Lyt can be

compensated by reducing basal expression of *lys*. For a given set of Hill coefficients, average k_3

is at least a few times higher for 6_Lyt_Lys as compared to that for 3_Lyt_Lys (data not shown).

Methods

Derivation of model equations

The model, using the fact that binding of protein to itself or DNA is a much quicker process than transcription and translation, assumes quick equilibration for the processes of protein binding to itself or DNA, in order to calculate the "combined" dissociation constants of proteins. In the expressions below, P, X, and Y are promoter, Lys, and Lyt, respectively.

$$X + X \overset{K_{DX}}{\rightleftharpoons} X_2 \qquad\qquad \frac{[X]^2}{[X_2]} = K_{DX}$$

$$Y + Y \overset{K_{DY1}}{\rightleftharpoons} Y_2 \qquad\qquad \frac{[Y]^2}{[Y_2]} = K_{DY1}$$

$$Y_2 + Y_2 \overset{K_{DY2}}{\rightleftharpoons} Y_4 \qquad\qquad \frac{[Y_2]^2}{[Y_4]} = K_{DY2}$$

$$P + X \overset{K'_{D0}}{\rightleftharpoons} PX \qquad\qquad \frac{[P][X]}{[PX]} = K'_{D0} = K_{D0}$$

$$P + X_2 \overset{K'_{D1}}{\rightleftharpoons} PX_2 \qquad\qquad \frac{[P][X^2]}{[PX_2]} = K'_{D1} K_{DX} = K_{D1}$$

$$P + Y_4 \overset{K'_{D2}}{\rightleftharpoons} PY_4 \qquad\qquad \frac{[P][Y^4]}{[PY_4]} = K'_{D2} K^2_{DY1} K_{DY2} = K_{D2}$$

Above expressions for concentrations of promoter-protein complexes are for cases where a) Lyt binds as monomer, b) Lyt binds as dimer, and c) Lys binds as tetramer. They exhaust all other cases, viz. monomeric and dimeric Lys, and monomeric and dimeric CII.

Processes of transcription and translation are not considered explicity except for *lyt-CII*

424 genes in the three-protein models. Hence, the model equations describe concentrations of

425 proteins only. With expressions for concentrations of promoter-protein complexes, one can

426 write generalized form of term representing protein production.

428
$$\frac{b + \sum\limits_{i} k_i.[DNA - Prot_i]}{[Unbound\ DNA] + \sum\limits_{i} k_i.[DNA - Prot_i]}$$

where b is, in case present, basal expression and k_i is rate constant for transcriptional activation

by i$_{th}$ protein.

Parameter sets, viz. rate constants and dissociation constants, of model equations were

searched deterministically in two stages, viz. order search and linear search (as they are named

here). In the order search, rate constants and dissociation constants were searched as 3's

exponent, which was varied between -5 and 5 with the difference of 1, in a nested fashion.

Thus, the number of parameter sets searched was equal to the number of parameters raised

to the power 11. Notably, switch quotients generated by this approach are unrefined because

rate constants and dissociation constants were increased geometrically, thereby causing a lot

of intervening values to remain unsampled. Therefore, parameter sets generated from order

search were further refined by linear search, which searches the neighbourhood of parameter

set arithmetically. It was noted that those parameter sets generated in the order search whose

SQs were too close to each other were either rescaled form of each other, or differed in

those parameters to which SQ was resilient up to a certain range. Thus, in order to remove

redundancy and in the interest of time, for linear search, the parameter sets were taken in such

a way that the difference between consecutive SQs is at least 0.01.

Parameter sets, and thus accompanied SQs, generated through order search were refined

by linear search in the following way. The value of each parameter (say, V) of a set was

varied between -3*V/5 and 3*V/5 with the increment of V/5, in a nested fashion. Thus, the

number of parameter sets searched was equal to the number of parameters raised to the power

7. However, for three-protein model, which had eight parameters, in the interest of saving time,

each parameter was varied between -2*V/5 and 2*V/5 with the increment of V/5, in a nested

fashion. Search was ended if the latest SQ was either lower than the previous one (which never

happened) or if ((latest SQ - previous SQ)/previous SQ) was less than 0.01. Again, in the

interest of saving time, for three-protein model, the search was ended if the SQ at the end of

the last iteration was more than or equal to 0.95. It should be noted that linear search is path

dependent: it may happen that a path which initially yields lower SQs leads to higher SQ in

the end than a path which initially yields higher SQs, and thus, treaded by the search. For both

order and linear search and for all of the models, in order to expedite search, those parameter

sets were rejected whose accompanying SQ was lower than the SQ of the previous parameter

set. The values of the parameters were normalized such that the Lyt's equilibrium concentration

was 10 arb. units. This was done for two purposes: a) to ensure that lowest values of Lyt at

MoI of 1 and Lys at MoI of 2 never drop to zero in the stochastic simulations; b) in order to

make comparison of parameter sets and equilibrium values of proteins visually easier. For both

order and linear search, simulations were carried for time 100 arb. units. Thus, there was a

possibility of a system of equations, defining a particular model, not reaching equilibrium in

100 arb. units for a given parameter set. In order to eliminate such parameter sets, simulations

were done for 10^5 arb. units. Only few parameter sets had not reached equilibrium, and all

of such parameter sets produced negative SQ. In order to calculate stochastic switch quotie nt,

levels of proteins were averaged between 100 and 200 arb. units. The transient kinetics, viz.

inital rise and plateauing at MoI of 1 and bell-shaped trajectory MoI 2, were completed at most

by 50 arb. units.

Acknowledgement

472 The author thanks Dr. Supreet Saini for hosting him in his lab in the Department of Chemical

473 Engineering at IIT Bombay.

References

[1] Kourilsky P (1973) Lysogenization by bacteriophage lambda. I. Multiple infection and the lysogenic response. Mol. Gen. Genet. 122:183195.

[2] Kobiler O, Rokney A, Friedman N, Court DL, Stavans J, Oppenheim AB (2005) Quantitative kinetic analysis of the bacteriophage lambda genetic network. Proc Natl Acad Sci U S A 102:4470-4475.

[3] Avlund M, Dodd IB, Semsey S, Sneppen K, Krishna S (2009) Why do phage play dice? J Virol. 83:11416-11420.

[4] Kourilsky P (1974) Lysogenization by bacteriophage lambda. II. Identification of genes involved in the multiplicity dependent processes. Biochimie 56:15171523.

[5] Banuett F, Hoyt MA, McFarlane L, Echols H, Herskowitz I (1986) hflB, a new Escherichia coli locus regulating lysogeny and the level of bacteriophage lambda cII protein. J Mol Biol 187:213-224.

[6] Kobiler O, Rokney A, Oppenheim AB (2007) Phage lambda CIII: a protease inhibitor regulating the lysis-lysogeny decision. PLoS One 2:e363.

[7] Weitz JS, Mileyko Y, Joh RI, Voit EO (2008) Collective decision making in bacterial viruses. Biophys J 95:2673-2680.

[8] Court DL, Oppenheim AB, Adhya SL (2007) A new look at bacteriophage lambda genetic networks. J Bacteriol 189:298-304.

[9] Avlund M, Krishna S, Semsey S, Dodd IB, Sneppen K. (2010) Minimal gene regulatory circuits for a lysis-lysogeny choice in the presence of noise. PLoS One 5:e15037.

[10] Hoopes BC, McClure WR (1985) A cII-dependent promoter is located within the Q gene of bacteriophage lambda. Proc Natl Acad Sci U S A 82:3134-3138.

[11] Ho YS, Rosenberg M (1985) Characterization of a third, cII-dependent, coordinately

activated promoter on phage lambda involved in lysogenic development. J Biol Chem

260:11838-11844.

[12] Svenningsen SL, Costantino N, Court DL, Adhya S (2005) On the role of Cro in lambda

prophage induction. Proc Natl Acad Sci U S A 102: 4465-4469.

[13] Révet B, von Wilcken-Bergmann B, Bessert H, Barker A, Mller-Hill B (1999) Four

dimers of lambda repressor bound to two suitably spaced pairs of lambda operators

[14] Ho Y, Lewis M, Rosenberg M (1982) Purification and properties of a transcriptional

activator. The cII protein of phage lambda. J Biol Chem 257: 9128-9134.

[15] Jain D, Kim Y, Maxwell KL, Beasley S, Zhang R, Gussin GN, Edwards AM, Darst SA

(2005) Crystal structure of bacteriophage lambda cII and its DNA complex. Molecular

Cell 19: 259-269.

[16] Cui L, Murchland I, Shearwin KE, Dodd IB (2013) Enhancer-like long-range

transcriptional activation by CI-mediated DNA looping. PNAS 110: 29222927.

[17] Elowitz MB, Levine AJ, Siggia ED, Swain PS (2002) Stochastic gene expression in

single cell. Science 297: 11831186.

[18] Golding I, Paulsson J, Zawilski SM, Cox EC (2005) Real-time kinetics of gene activity

in individual bacteria. Cell 123: 10251036.

[19] Gillespie DT (1977) Exact stochastic simulation of coupled chemical reactions. J Phys

Chem 81: 23402361.