A majority of HIV persistence during antiretroviral therapy is due to infected cell proliferation

Daniel B. Reeves[1], Elizabeth R. Duke[1,2], Thor A. Wagner[3,4], Sarah E. Palmer[5], Adam M. Spivak[6], Joshua T. Schiffer[1,2,7*]

[1]Vaccine and Infectious Diseases Division, Fred Hutchinson Cancer Research Center, 1100 Fairview Ave. Seattle, WA 98122, USA
[2]Department of Medicine, University of Washington, 1959 NE Pacific St. Seattle, WA 98195, USA
[3]Department of Pediatrics, University of Washington, 1959 NE Pacific St. Seattle, WA 98195, USA
[4]Center for Global Infectious Disease Research, Seattle Children's Research Institute, 1900 9th Ave, Seattle, WA 98101, USA
[5]Centre for Virus Research, The Westmead Institute for Medical Research, 176 Hawkesbury Rd, Westmead NSW 2145, Australia
[6]Department of Medicine, University of Utah, 30 N 1900 E, Salt Lake City, UT 84132, USA
[7]Clinical Research Division, Fred Hutchinson Cancer Research Center, 1100 Fairview Ave. Seattle, WA 98122, USA
*Corresponding author, email: jschiffe@fhcrc.org (JTS)

Abstract

Antiretroviral therapy (ART) suppresses viral replication in people living with HIV. Yet, infected cells persist for decades on ART and viremia returns if ART is stopped. Persistence has been attributed to viral replication in an ART sanctuary and long-lived and/or proliferating latently infected cells. Using ecological methods and existing data, we infer that >99% of infected cells are members of clonal populations after one year of ART. We reconcile our results with observations from the first months of ART, demonstrating mathematically how a "fossil record" of historic HIV replication permits observed viral evolution even while most new infected cells arise from proliferation. Together, our results imply cellular proliferation generates a majority of infected cells during ART. Therefore, reducing proliferation could decrease the size of the HIV reservoir and help achieve a functional cure.

Introduction

Antiretroviral therapy (ART) limits HIV replication in previously uninfected cells leading to elimination of most infected CD4+ T cells.[1] Yet, some infected cells persist and are cleared from the body at an extremely slow rate despite decades of treatment.[2,3] There is debate whether infection remains due to HIV replication within a small population of cells[4,5] or due to persistence of memory CD4+ T cells with HIV integrated into human chromosomal DNA.[3,6,7] If the latter mechanism predominates, prolonged cellular lifespan and/or frequent cellular proliferation may sustain stable numbers of infected cells.

To optimize HIV cure strategies, mechanisms sustaining infection must be understood. Persistent viral replication in a "sanctuary" where ART levels are inadequate implies a need to improve ART delivery.[8] If HIV persists without replication as a *latent reservoir* of memory CD4+ T cells, then the survival mechanisms of these cells are ideal therapeutic targets. Infected cell longevity might be addressed by reactivating the lytic HIV replication cycle[9] and strengthening the anti-HIV cytolytic immune response,

42 leading to premature cellular demise. Anti-proliferative therapies could limit homeostatic or antigen
43 driven proliferation.[10-12]
44
These competing hypotheses have been studied by analyzing HIV evolutionary dynamics. Due to the
high mutation rate of HIV reverse transcriptase and the large viral population size,[13] HIV replication in
the absence of ART produces large viral diversity.[13-15] Over time, new strains become dominant due to
continuous positive immunologic selection pressure against the virus. Repeated "selective sweeps"
cause genetic divergence, or a positive molecular evolution rate,[16] often measured by continual growth
in genetic distance between the consensus strain and the founder virus.[17-19]
51
A recent study documented new HIV mutants during months 0-6 of ART in three participants at a rate
equivalent to pre-ART time points. New mutations were noted across multiple anatomic compartments,
implying widespread circulation of evolving strains.[4] One possible explanation for this data is the
presence of a drug sanctuary in which ART levels are insufficient to stop new infection events.
Alternative proposed interpretations are experimental error related to PCR resampling, or variable
cellular age structure within the phylogenetic trees.[20,21]
58
In other studies of participants on more prolonged ART (at least one year), viral evolution was not
observed despite sampling of multiple anatomic compartments.[22-25] Identical HIV DNA sequences were
noted in samples obtained years apart,[14,26,27] suggesting long-lived latently infected cells as a possible
mechanism of HIV persistence.[3,6,7,24,25] Clonal expansions of identical HIV DNA sequences were also
observed, demonstrating that cellular proliferation generates new infected cells.[4,12,24,28-30] Multiple,
equivalent sequences were noted in blood, gut-associated lymphoid tissue (GALT), and lymph nodes,
even during the first month of ART.[24,29,30]
66
The majority of these studies relied on sequencing single genes including *env*, *gag* and *pol*: this approach
may overestimate HIV clonality because mutations in other genome segments could go unobserved.[17,31]
In addition, these studies also measured total HIV DNA. However, a majority of HIV DNA sequences have
incurred deleterious mutations and do not constitute the true replication competent HIV reservoir.[32,33]
To address these issues, a more recent study utilized a comprehensive, whole-genome sequencing
approach to confirm the presence of abundant replication competent sequence clones.[34] In a separate
cohort of patients, rebounding HIV sequences arose from replication competent clonal populations.[35]
74
Another approach to define HIV clonality involves sequencing of the HIV integration site within human
chromosomal DNA.[36-40] While HIV tends to integrate into the same genes,[39,41] it is extremely unlikely that
two cellular infection events would result in HIV integration within precisely the same human
chromosomal locus by chance alone.[37] Thus, integration site analyses abrogate the challenge of
overestimating clonality due to incomplete sequencing and provide an elegant surrogate for whole
genome sequencing. Previous studies of integration sites found significant numbers of repeated
integration sites, providing strong evidence that these infected cells arose from cellular proliferation.[42,43]
These studies are not absolutely conclusive for HIV persistence because integration site sequencing
cannot confirm or deny replication competency of the integrated virus.[39]
84
While HIV sequence clonality has been widely observed, existing studies observed equivalent sequences
in a minority (<50%) of observed sequences. Here, we demonstrate that this finding can be explained by
incomplete sampling. Using tools adapted from ecology and data from two integration site studies[36,37]
and a replication competent HIV DNA study,[34] we show that nearly all observed unique sequences are

likely to be members of clonal populations which derived from cellular proliferation. We predict that the
HIV reservoir consists of a small number of massive clones, and a massive number of small clones.
91
Based on these results, we used a mechanistic mathematical model to reconcile apparent evolution
during the early months of ART with apparent clonality after a year or more of ART. The model includes
the major proposed mechanisms for HIV persistence: a drug sanctuary, long-lived infected cells, and
proliferating infected cells. The model highlights that observed HIV evolution during the first 6 months
of ART can be caused by serial observations of long-lived (or proliferated) cells that were once generated
by viral replication. We suggest sampling sequences during early ART may result in detection of a
positive molecular evolution rate due to the "fossil record" of past infections rather than current viral
replication in a drug sanctuary. Based on observed cellular rates, model output after one week of ART
shows that a majority of new infected cells are generated by proliferation.
101
102 While it remains impossible to rule out a completely unobserved drug sanctuary, our combined
103 approaches suggest that cellular proliferation predominantly drives observed HIV persistence on ART.
104 Consequently, anti-proliferative therapies embody a meaningful therapeutic approach for HIV cure.

Results

Defining genetic markers of HIV persistence. During untreated infection, HIV integrates its DNA copy
into human chromosomal DNA in each infected CD4+ T cell.[44] A majority of new infections are marked
by novel mutations due to the high error rate of HIV reverse transcriptase and integration into a unique
chromosomal location (**Fig 1**). Therefore, continual accrual of new mutations during ART would suggest
that ongoing viral replication, perhaps due to inadequate drug delivery to certain micro-anatomic
regions, allows HIV to persist during ART.
112
113 In a subset of infected CD4+ T cells, HIV replication does not progress beyond chromosomal integration
114 and the virus enters latency.[44] If the same HIV sequences (or integration sites) are found over long time
115 intervals, either cellular longevity or proliferation of latently infected cells allowed HIV to persist. If
116 equivalent HIV sequences with identical chromosomal integration sites are identified in multiple cells,
117 then these viruses were generated via cellular proliferation, rather than HIV replication (**Fig 1**).

118
119 *Figure 1. Possible mechanisms for HIV reservoir persistence and their genetic signatures. Viral*
120 *replication despite ART would lead to accrual of new mutations (color change) and novel chromosomal*
121 *integration sites in newly infected cells. Alternatively, longevity of latently infected cells maintains*
122 *sequences and integration sites. Finally, cellular proliferation of latently infected cells produces clonal*
123 *populations of equivalent HIV sequences and integration sites.*

124
125 Throughout the paper, we contrast the impact of HIV replication and cellular proliferation on HIV
126 persistence during ART by quantifying the numbers or fractions of *unique* sequences and *equivalent*
127 sequences. Human DNA polymerase has much higher copying fidelity than HIV's reverse transcriptase.
128 Thus, we assume cells whose origin is viral replication will contain unique sequences while cells whose
129 origin is cellular proliferation will contain equivalent sequences and be members of clonal populations.

130
Fractions of equivalent total HIV DNA sequences may be extrapolated to replication competent
sequences. Most integrated HIV DNA has accrued mutations that render the virus replication
incompetent. Quantification of total HIV DNA copies therefore overestimates the size of the replication
competent reservoir by 2-3 orders of magnitude relative to viral outgrowth assays.[32] Replication
incompetent, equivalent HIV sequences are commonly present in multiple cells[24,29]. Precisely because
these sequences are terminally mutated, they are concrete evidence that some other mechanism
(cellular proliferation) copies HIV DNA. The proportion of clonal sequences is similar when analysis
includes only replication competent sequences, or all HIV DNA.[34] As a result, while total HIV DNA may
not predict quantity of replication competent viruses, estimates of clonal frequency using total HIV DNA

might be extrapolated to the smaller replication competent reservoir.[33] We use total HIV DNA as it
allows a greater sample size for analysis.
142
Clonal HIV DNA sequences and clonal replication competent sequences are detectable at various time
points during ART. To examine the structure of clonal total and replication competent HIV DNA, we
ranked observed sequences from several studies according to their abundance: rank-abundance curves
are ordered histograms denoted $a(r)$ such that $a(1)$ is the abundance of the largest clone. These curves
facilitate identification of quantities of interest like the richness $R = \max(r)$, sample size $N = \sum_r a(r)$,
and the number of singletons $N_1 = \sum_r I[a(r) = 1]$. Here $I[\cdot]$ is the indicator function equal to 1 when
its argument is true and 0 otherwise.
150

151
Figure 2. Evidence for clonal HIV sequences. *Raw data rearranged as rank abundance curves.* ***A.*** *Total*
HIV DNA from integration site data (Wagner et al., and Maldarelli et al.)[36,37]. Each panel represents a
participant, and each marker a duration of ART (indicated in years in the panel legend). W and M in the
panel headings distinguish the study. ***B.*** *Replication competent HIV DNA (Hosmane et al.)[34]. Each panel*
represents a participant. Participants used for analyses below have more than 20 sequences observed
(noted by asterisks in panel headings). ***C & D.*** *Sample size of HIV DNA* ***(C)*** *and replication competent HIV*
DNA ***(D).*** *Measuring total HIV DNA increases the number of observed unique sequences (observed*
sequence richness). The number of total sequences at each time point is plotted against the observed
sequence richness. For all HIV DNA samples and when $N > 20$ for replication competent HIV DNA, the
observed richness is always less than the sample size (to the right of the dotted line y=x), owing to the
presence of sequence clones. ***E.*** *Sample rarefaction curves for all 17 time points from the 8 study*
participants in ***A*** *demonstrate the observed number of distinct integration sites as a function of HIV DNA*
sequence experimental sample size. ***F.*** *Sample rarefaction curves for all 5 study participants in* ***B***
demonstrate the observed number of distinct replication competent HIV DNA sequences as a function of
sequence sample size. In both cases, at low sample size, distinct sequences are commonly observed with
each new sample. As sample size increases, distinct sequences are increasingly less likely to be detected
owing to the presence of repeatedly detected sequence clones. As more and more unique sequences are
detected, the curves would flatten until all unique sequences are detected and the curve is completely
flat.

Wagner *et al.* sampled HIV DNA in three participants at three time points 1.1-12.3 years following ART
initiation.[37] Maldarelli *et al.* sampled HIV DNA from five participants at one to three time points 0.2-14.5
years following ART initiation.[36] In these studies, 1-16% (mean: 7%) of sequences were members of
observed sequence clones (**Fig 2A**),[36,37] meaning that HIV DNA was identified in the same chromosomal
integration site in at least two cells. The absolute number of observed sequence clones $N_{i>1}$ in the 17
samples ranged from 1-150 (mean: 15). The remaining sequences were identified in a specific
chromosomal integration site in only one cell (*observed singletons*).[37] For total HIV DNA, at each
participant time point, certain sequences predominated: the largest observed sequence clone contained
2-62 sequences (mean: 11), accounting for 3-26% (mean= 9%) of total observed sequences.

Hosmane *et al.* sequenced replication competent HIV isolates from 12 study participants on ART: 0-28%
(mean: 11%) of sequences were members of *observed sequence clones* (**Fig 2B**).[34] The lack of detected
clones in 3 participants may reflect their low sequence sample size. Participants with fewer than 20
total sequences were therefore excluded from individual analyses described below but were included
for population level evaluations. For replication competent HIV DNA in the 5 persons having sequence
sample-size $N > 20$, certain sequences dominated: the largest observed sequence clone contained 3-9
sequences (mean: 6.8), accounting for 11-42% (mean= 28%) of total observed sequences. The number
of non-singleton sequence clones $N_{i>1}$ in the 5 samples ranged from 1-7 (mean: 3.8).

Sequence sampling depth is low relative to total population size. There was a higher number of
experimentally detected sequences (N) for total HIV DNA (**Fig 2C**) than for replication competent HIV
(**Fig 2D**). For total HIV DNA, the number of observed *unique* sequences (R^{obs} or the *observed sequence
richness*) was always less than N (**Fig 2C**) due to clonal populations. Where $N > 20$ for replication
competent viruses, R^{obs} was always less than N, again due to the presence of clones **(Fig 2D)**. There was
a higher R^{obs} as the sequence sample size increased (**Fig 2C&D**), suggesting that detection of unique
clones increases with deeper sampling.

Thus, we can infer that further sampling would likely uncover new unique sequences. To quantify the
relationship between sample size and discovery, we generated sample rarefaction curves (see **Methods**
and **Supplementary Methods**) using the rank-abundance distributions (**Fig 2E&F**). These curves
interpolate the data to demonstrate the likely discovery of new sequences as sampling increases up to
the sample size of the original experiment. At low sample size, a new sequence is likely to be found with
each additional sample. As sampling increases, the chance of sampling a previously documented
sequence increases, and the slope of the rarefaction curve begins to flatten. As sample size approaches
the true richness of the population, the curve plateaus and few new unique sequences remain to be
sampled. Current sampling depth remains on the steep, initial portion of the curve.

Ecological estimates of lower bounds on true HIV sequence richness from limited samples. To estimate
a lower bound for true sequence richness, we used the Chao1 estimator, a nonparametric ecologic tool
that uses frequency ratios of observed singletons N_1 and doubletons N_2 (see **Methods** and
Supplementary Methods).[45,46] For the HIV reservoir, theoretical values for true richness range from one
(if all sequences were identical and originated from a single proliferative cell) to the total population size
(if all sequences were distinct and originated from error-prone viral replication). We found estimated
lower bounds for true sequence richness exceeded observed richness, typically by an order of
magnitude in both total HIV DNA and replication competent HIV (**Fig 3**). These initial lower bound
estimates for sequence richness are far lower than previously estimated population sizes for HIV DNA
and replication competent HIV DNA sequences,[2,3,6] suggesting that clones may predominate.

219
220

221
Figure 3. The actual total number of distinct HIV sequences far exceeds the observed total number of
distinct HIV sequences during ART. *Observed sequence richness underestimates the true HIV sequence*
richness. For both data sources, Chao1 provides an estimate of the lower bound (min) of true sequence
*richness (error bars are asymmetric confidence intervals, see **Supplementary Methods**). In all cases,*
Chao1 estimates are above observed values. Our modeling technique estimates a much higher upper
bound (max) for true sequence richness. Nevertheless, the total HIV sequence population size (dashed
lines: 10^9 for total HIV DNA and 10^7 for replication competent HIV) is 1-2 orders of magnitude above the
upper bound estimates for sequence richness, suggesting substantial clonality of HIV sequences.
230

231 **A majority of observed HIV sequences are members of large proliferative clones.** The Chao1 estimator
232 does not include information about the total population size. However, estimates for the total number
233 of total DNA and replication competent sequences in the entire reservoir exist.[33] Using that additional
234 information, we developed an ecologic model to extrapolate the true rank-abundance of HIV sequences
235 for each participant time point.
236

Based on the observation that observed data was roughly log-log-linear **(Figure 2A)**, we chose a power-
law model for rank-abundance: $a(r) \propto r^{-\alpha}$. Other functional forms were explored (exponential, linear,
and biphasic power law) but were worse or equivalent for data fitting (not shown). Our model requires 3
parameters, the power law exponent (α), the sequence population size (L), and the sequence richness
(R). Model fitting is described in the **Methods** with additional detail in the **Supplementary Methods**.
Briefly, we generated 2,500 possible models for each data set, choosing a plausible fixed population size
from available data ($L = 10^9$ for HIV DNA and $L = 10^7$ for intact, replication competent HIV
DNA).[2,3,6,33,47] We then recapitulated the experiment by taking N random samples from each model
distribution and comparing sampled data to experimental data to find optimal model parameters. This
resampling method correctly inferred the power law exponent from simulated power law data
(Supplementary Fig 1).
248

However, for experimental data we could not precisely identify R. Recognizing this uncertainty, we
developed an integral approximation to estimate the largest possible richness (least clonality) given L
and the best-fit α (derivation in **Supplementary Methods** and illustration in **Supplementary Fig 2**). Then,
using the lower bound estimate from the Chao1 estimator, we were able to fully constrain the estimate
of true HIV sequence richness in the reservoir. Our maximal estimates for sequence richness were

notably several orders of magnitudes higher than Chao1 estimates (**Fig 3**) but lower than the total
sequence population size (L).

Our method demonstrated excellent fit to cumulative proportional abundances of observed clones for
total HIV DNA (**Fig 4A**) and replication competent HIV DNA (**Fig 5A**). For total HIV DNA (**Fig 4B**) and
replication competent HIV DNA (**Fig 5B**), optimal fit was noted within narrow ranges for the power law
slope parameter but across a wide possible range of true sequence richness. Using the top 5 best fit
models, we generated extrapolated distributions of the entire HIV sequence rank-abundance for each
participant time point. We observed similar estimates for the population size of the largest clones,
which account for approximately 50% of the reservoir (200-2,000 clones for HIV DNA in **Fig 4C** and 2-7
clones for replication competent HIV DNA in **Fig 5C**). However, the tail of the reservoir, which consists of
thousands of smaller clones, varied considerably across the parameter sets with 900-100,000 possible
clones accounting for 90% of the HIV DNA and 100-2,000 possible clones accounting for 90% of
replication competent HIV. This variability reflects the fact that true sequence richness is only partially
identifiable using our procedure.

270
271 **Figure 4. Ecologic modeling suggests a majority of HIV DNA sequences are members of sequence**
272 **clones.** *To model the true rank abundance distribution of the HIV reservoir, we used a power law model*

*and recapitulated experimental sampling (sample size equal to the experimental sample size) from 2,500 theoretical power law distributions to fit the best model to participant data in **Fig 2A**. Theoretical distributions varied according to the slope of the power law and the true sequence richness but were fixed at 10^9 total HIV DNA sequences. **A.** Five best model fits to cumulative proportional abundance curves from a single representative participant (WR, 12 years on ART). Black circles represent the experimental data; the 5 colored model lines are superimposed based on virtually equivalent fit to the data. **B.** Heat diagram representing model fit according to power law exponent α and true sequence richness R with best fit noted by minimum error score (blue color, see details of calculation in results above); black shaded areas represent parameter sets excluded based on the Chao1 estimator (lower bound on sequence richness) and mathematical constraints of the power law (upper bound for sequence richness). A wide range of values for sequence richness allow excellent model fit while the power law exponent is well defined. **C.** Extrapolations of the best-fit cumulative distribution function to the entire pool of 10^9 infected cells; under the most conservative estimates, the top 200,000 ranked clones constitute the entire reservoir. **D.** Extrapolations of the best fit power law to the entire pool of 10^9 infected cells; the top 1000 clones consist of >10^4 cells each. **E.** Extrapolations of the best fit cumulative distribution function to the entire pool of 10^9 infected cells for all participant time points in **Fig 2A**; we assume the maximum possible sequence richness in each case and still note a predominance of sequence clones. **F.** Extrapolations of the best-fit power law to the entire pool of 10^9 infected cell for all participants in **Fig 2A**; the top 1,000 clones each consist of >10^4 cells each. A large number of clones (~10^6) contain many fewer cells (<100).*

Even under the most conservative assumptions (maximum possible true sequence richness in **Fig 3**), the vast majority of sequences were predicted to be members of true sequence clones. For the participant in **Fig 4C**, a maximum of 200,000 clones were needed to reach 100% cumulative abundance for HIV DNA. The ratio of estimated true sequence richness to the total number of infected cells R/L with HIV DNA (~10^5: 10^9) represents an upper limit on the fraction of sequences that are true singletons: we estimate that greater than 99.9% of infected cells contain true clonal sequences (**Fig 3**).

Similarly, the ratio of estimated true sequence richness to the total number of infected cells with replication competent HIV for the participant in **Fig 5C** was 10^5:10^7. Hence, at least 99% of cells contain true clonal sequences (**Fig 3**). Of note, this ratio is stable regardless of assumed reservoir size. For instance, if we assume a true reservoir size of 10^6, then our estimate of true sequence richness is ~10^4.

The model fitting procedure was used on all data in **Fig 2**. We biased against a clonally dominated reservoir to the greatest extent possible by selecting the best fitting power law exponent and then calculating the maximum possible sequence richness (**Fig 3**). The power law slope parameter was on average lower across participants for HIV DNA ($\alpha = 0.9 \pm 0.1$) than for replication competent HIV DNA ($\alpha = 1.4 \pm 0.2$). As a result, the predicted cumulative distribution of HIV DNA (**Fig 4E**) was often concave-up with log rank as compared to concave-down with log rank noted for replication competent HIV DNA (**Fig 5E**), suggesting that a smaller number of extremely large clones might make up a higher proportion of the replication competent HIV reservoir.

For both HIV DNA (**Fig 4F**) and replication competent virus (**Fig 5F**), the top 100 clones in all participants are estimated to be massive (>10^5 and >10^4 cells respectively). However, there are also large numbers of much smaller clones with fewer than 1,000 cells (>10^6 and >10^4 clones respectively). In contrast to observed data, a majority of sequences are clonal, suggesting that proliferation is the major generative mechanism of persistent HIV-infected cells.

320
321 ***Figure 5. Ecologic modeling suggests a majority of replication competent HIV sequences are members***
322 ***of sequence clones.*** *To recapitulate experimental conditions in **Fig 2B**, we performed in silico sampling*
323 *(sample size equal to the experimental sample size) from 2,500 theoretical power law distributions of*
324 *replication competent HIV clone size distributions sorted by rank. Theoretical distributions varied*
325 *according to the exponent of the power law model and the true sequence richness and were fixed at a*
326 *reservoir size of 10^7 replication competent HIV DNA sequences. **A.** Five best model fits to cumulative*
327 *proportional abundance curves from a single representative participant (S10). Black circles represent the*
328 *experimental data; the 5 colored model lines are from five separate parameter sets. **B.** Heat map*
329 *representing model fit according to power law slope α and true sequence richness R with best fit noted*
330 *by lowest error (blue color); the black shaded area represents parameter sets excluded based on*
331 *mathematical constraints of the power law (upper bound on sequence richness). A wide range of values*
332 *for sequence richness (<10^5 sequences) allow excellent model fit while power law slope falls within a*
333 *narrow range. **C.** Extrapolations of the best-fit cumulative distribution function to the entire pool of 10^7*

infected cells; under the most conservative estimates, the top 10^5 ranked clones constitute the entire
reservoir. ***D.*** *Extrapolations of the best fit power law to the entire pool of 10^7 infected cells; the top 100*
clones consist of >10^4 cells each. ***E.*** *Extrapolations of the best fit cumulative distribution function to the*
entire pool of 10^7 infected cells for all participants and time points (see original data in ***Fig 2B****); we*
assume the largest possible observed sequence richness in each case and still note a predominance of
sequence clones. ***F.*** *Extrapolations of the best-fit power law to the entire pool of 10^7 infected cell for all*
participants in ***Fig 2B****; the top 100 clones again consist of >10^4 cells each. A large number of clones*
(~10^4) contain many fewer cells (<100).

Modeling combined population data gives similar results as individual fitting. To increase sample size
and eliminate bias related to excluding participants with low sample sizes, we combined results from all
participant time points for HIV DNA (17 time points) and replication competent HIV (12 time points) into
single rank order distribution curves. We then fit the power law models to both sets of data
(**Supplementary Fig 3A&B, E&F**). We again noted a narrow range of possible values for the power law
exponent and a large range of possible values for true sequence richness. The exponent was again $\alpha <$
1 for total HIV DNA and $\alpha \approx 1$ for replication competent virus (**Supplementary Fig 3A&E**), leading to
concave-up and linear relationships between cumulative proportional abundance and log rank,
respectively (**Supplementary Fig 3C&G**). We estimated that at least 99.9% of cells with HIV DNA
(**Supplementary Fig 3C**) and 99.8% of cells with replication competent HIV (**Supplementary Fig 3G**)
contain true clonal sequences. The top 100 HIV DNA clones (**Supplementary Fig 3D**) and replication
competent clones (**Supplementary Fig 3H**) contained >10^6 and >10^4 cells respectively.

Using the population level data, we generated sample rarefaction curves from the extrapolated rank-
abundance curves. These curves show that after 10,000 sequences were sampled, the observed
sequence richness would continue to increase with more sampling (**Supplementary Fig 4**). Even if
experimental sample sizes could be increased 100-fold from the present data, sequences would
continue to be dominated by those from large clones. Our statistical inference approach is therefore
necessary to provide a more realistic estimate of the clonal distribution of the HIV reservoir.

A mechanistic model that includes both an ART sanctuary and cellular proliferation can reconcile
observations from early and late ART. Our analyses above identify the critical role of cellular
proliferation in generating infected cells after a year of ART but do not capture the dynamic mechanisms
underlying this observation or explain possible evidence of viral evolution during months 0-6 of ART.[4]
We therefore developed a viral dynamic mathematical model. Our model (**Fig 6A**) consists of differential
equations, described in detail in the **Methods**. Most model parameter values are obtained from the
literature (**Table 1**).

Figure 6. A mechanistic model recapitulates HIV RNA decay and predicts rough equivalence of virus
produced by the sanctuary and virus produced by a reactivating reservoir up until months 4-6 of ART.
A*. Model schematic: I_1 cells produce virus, pre-integration latent cells I_2 are longer lived and eventually*
transition to I_1, and long-lived latently infected cells $I_{3(j)}$ proliferate and die at measured rates
depending on cell phenotype j (e.g. effector memory, central memory, naïve. Sanctuary cells I_S allow
ongoing HIV replication despite ART. Parameters and their values are discussed in the Methods and listed
*in Sup Table 1. **B**. The mathematical model recapitulates observed HIV RNA data (Palmer et al.[51]) over*
*weeks and years of ART. V_1 is virus derived from I_1 while V_S is derived from I_S. **C**. I_2 and I_3 become the*
predominant cell types early during ART. I_S remains very low throughout the duration of ART which is
necessary to explain the lack of detectable viremia on fully suppressive ART.

Briefly, we classify rapid death δ_1 and viral production within actively infected cells I_1. Cells with longer
half-life I_2 are activated to I_1 at rate ξ_2. I_2 may represent CD4+ T cells with a prolonged pre-integration
phase, but their precise biology does not affect model outcomes.[48] The state $I_{3(j)}$ represents latently
infected reservoir cells of phenotype j, which contain a single chromosomally integrated HIV DNA
provirus.[44] I_3 reactivates to I_1 at rate ξ_3.[49] The probabilities of a newly infected cell entering
$I_1, I_2, I_{3(j)}$, are $\tau_1, \tau_2, \tau_{3(j)}$. Because we are focused on the role of proliferation, we assume sub-
populations of I_3,[12] including effector memory (T$_{em}$), central memory (T$_{cm}$), and naïve (T$_n$) CD4+ T cells,
which have been experimentally proven to turn over at different rates $\alpha_{3(j)}, \delta_{3(j)}$.[12,42,43]

ART potency $\epsilon \in [0,1]$ characterizes decrease in viral infectivity due to ART.[50] Other dynamic features of
infection such as death rate of infected cells, latent cell proliferation rate and reactivation rates of latent
cells, are unchanged on ART. In our simulations, the basic reproductive number becomes $R_0(1 - \epsilon)$ on
ART and is <1 when $\epsilon > 0.95$, meaning that each cell infects fewer than one other cell and viral load
declines from its previous steady state until becoming undetectable. Only short stochastic chains of new
infection can occur.

To make a model inclusive of viral evolution despite ART, we allow for the possibility of a drug sanctuary
state (I_S) that reproduces with reproductive number $R_0(1 - \epsilon_S) \sim 8$. In the drug sanctuary, ART potency
is assumed to be negligible ($\epsilon_S = 0$) such that the sanctuary reproductive number is equivalent to the
value from a model without ART. Target cell limitation or a local immune response must result in a
sanctuary viral set point to prevent infected cells and viral load from growing exponentially. The
sanctuary size must also be limited (0.001-0.01% of the original burden of replicating HIV) to achieve
realistic viral decay kinetics.[51] In the absence of contradictory information, we assumed homogeneous
mixing of V_1 and V_S in blood and lymph nodes.[4]

Based on the observation that activated, uninfected CD4+ T cells (*S*), the targets for replicating HIV,
decrease in numbers after initiation of ART we also simulate the model with and without the possibility
of slow target cell decline within the HIV drug sanctuary. We approximate this process with an
exponential decay of target cells with rate ζ (per day).[52,53] The decay rate is lower than concurrent decay
rates measured from HIV RNA[50,51,54] because abnormal T cell activation persists for more than a year
after ART.[53]

The model accurately simulates viral dynamics during ART. We fit the model to ultra-sensitive viral load
measurements collected from multiple participants in Palmer *et al*.[51] We included experimentally
derived values for most parameter values (**Table 1**), solving only for activation rates ξ_2 and ξ_3 by fitting
to viral load. Simulations reproduce three phases of viral clearance (**Fig 6B**) and predict trajectories of
infected cell compartments (**Fig 6C**). Of note, the model is able to achieve fit to the data with different
assumptions of starting values of the three infected cell compartments (the relative proportion of which
are unknown pre-ART): in this circumstance, we arrive at different values of ξ_2 and ξ_3 without impacting
overall model conclusions regarding the HIV reservoir. The size of the sanctuary (expressed as the
fraction of infected cells φ_S) is only constrained to be below a value $<10^{-5}$ to ensure accurate model fit
for a static sanctuary model.

Cellular proliferation sustains HIV infection during ART whether or not a small drug sanctuary exists.
We next used the model to estimate the fraction of cells generated by cellular proliferation versus viral
replication. We conservatively assumed that prior to ART all infected cells were generated by viral
replication. Then, we tracked the number of cells whose origin was replication and the number whose
origin was cellular proliferation. Without directly simulating a phylogeny, the fraction of all cells that
derive from replication provides a surrogate for the expected fraction of cells that would give a signal of
evolution. We also distinguish the *current replication percentage*, the fraction of infected cells currently
being generated from viral replication, from the *net replication percentage*, the fraction of total infected
CD4+ T cells at a given time whose origin was HIV replication. This distinction allows us to contrast the
net number of surviving, historically-infected cells with the number of cells that are presently being
generated via HIV infection. Because many long-lived cells were once generated by HIV infection, the
net replication percentage may exceed the current replication percentage.

We then simulated the model under several plausible sanctuary and reservoir conditions to assess the
relative contributions of infection and cellular proliferation in sustaining infected cells. We considered
different reservoir compositions based on evidence that effector memory (T_{em}), central memory (T_{cm})
and naïve (T_n) cells proliferate at different rates and that distributions of infection in these cells differ
among infected patients.[12,42,43] Further, because a drug sanctuary has not been observed, its true
volume is unknown and may vary across persons. We therefore conducted simulations with a static
sanctuary, a slowly diminishing sanctuary, and no drug sanctuary **(Fig 7A)**.

446
Figure 7. The vast majority of infected cells are generated via proliferation within 6 months of ART
initiation. *Model simulations contrast the number of cells generated by viral replication with those*
generated by cellular proliferation. The fraction of cells generated by replication at any time point is
referred to as the current replication percentage. The fraction of cells that remain alive whose ultimate
origin was viral replication is referred to as the net replication percentage. Different assumptions
*regarding sanctuary (I_S) and latent cell populations (I_3) were simulated corresponding to columns. **A.***
Moving left to right, we assume a static drug sanctuary, a slowly declining drug sanctuary and no drug
sanctuary. Pie charts on the right indicate the reservoir composition by T cell phenotypes and correspond
*with colored lines in **B-D. B.** Under all assumptions, once ART is initiated, most new infected cells arise*
*due to cellular proliferation as opposed to HIV replication after 12 months of ART. **C.** New latently*
infected reservoir cells (I_3) are generated almost entirely by proliferation soon after ART is initiated
*under all conditions. **D.** The observed proportion of infected cells originally generated by HIV infection*
rather than cellular proliferation will overestimate the actual ongoing proportion during the first 6
months of ART assuming a small or large sanctuary volume. This trend is more notable when the
reservoir contains a higher proportion of slowly proliferating naïve T cells.

Regardless of assumed pre-treatment reservoir composition and sanctuary size, the contribution of
replication to generation of new infected cells is negligible after one year of ART. The contribution of
new replication diminishes rapidly with time on ART regardless of whether a sanctuary is assumed (**Fig
7B**). The fraction of long lived latently infected cells (I_3) generated by viral replication (**Fig 7C**, note log
scale) is negligible within days of ART initiation. This finding captures the extent of the impact of
proliferation even when a sanctuary is assumed.

**Observable HIV DNA sequence evolution during early ART can represent a fossil record of prior
replication events.** In all simulations, the net fraction of cells generated from viral replication rather
than cellular proliferation at 6 months of ART (5-25% in **Fig 7D**) is higher than the current percentage
generated by replication (**Fig 7B**). A higher fraction of slowly proliferating T_n cells exacerbates the
difference between historical and contemporaneous generation of infected cells (**Fig 7D**, green line).
Because the net fraction is what will be observed experimentally, the model reveals why ongoing
evolution might be observed even while the dominant mechanism sustaining the reservoir is cellular
proliferation. In keeping with the first section of our paper, after 12 months of ART, the net and current
percentage of infected cells generated by HIV replication become negligible for all simulated parameter
sets. Importantly, the lag between net and current viral replication generation emerges whether or not
a small drug sanctuary is included in the model.

We refer to the phenomenon that long-lived cells may contain signatures of past viral replication as the
"fossil record". To emphasize the concept, the fossil record finding is qualitatively illustrated in **Fig 8**
using a population of 30 infected cells. At 3 time points following the initiation of ART, we compare the
net and current percentage of cells generated by viral replication. At day 60, 30% of cells remain that
were originally generated by viral replication. This means 30% of observed sequences might produce a
signal of evolution. However, at that time an overwhelming majority of new infected cells are being
generated by proliferation.

490
Figure 8. Qualitative illustration of the fossil record phenomenon. In an example population of 30
infected cells, the proportion of infected cells that were once generated by HIV replication (the net
replication percentage, or "fossil record" of HIV replication) remains >30% for the first 2 months of ART.
However, in this time, the proportion of cells newly generated by HIV replication (shaded box) becomes
negligible. The net fraction is observed experimentally, so our simulations indicate a contemporaneous
representation of the HIV reservoir cannot be observed until the "fossil record" is completely washed out,
sometime between 6 months and a year of ART.
498
Different factors drive net (observed) and current replication percentage during early ART. We next
performed sensitivity analyses to identify parameters that impact the timing of transition from HIV
replication to cellular proliferation as a source for new and observed infected cells. Under all parameter
assumptions, the majority of new infected cells arose from proliferation after a year of chronic ART (**Fig**
9A). Only the sanctuary decay rate (ζ) had an important impact on generation of new infected cells. Our
analysis included a sanctuary in which target cell availability did not decay at all. In that scenario, 5-10%
of new infected cells were generated by HIV replication after a year of ART (**Fig 9A**), which is not
consistent with lack of viral evolution observed at this timepoint. Rapid disappearance of HIV replication
as a source of new infected cells was identified regardless of initial reservoir volume, drug sanctuary
volume, ART efficacy, and reservoir composition (fraction of T_{em}, T_{cm}, and T_n).
509
510 The net replication percentage was completely unaffected by the decay rate of target cells within the
511 drug sanctuary. Only an increase in the percentage of slowly proliferating reservoir cells (T_n) predicted
512 an increase in the net replication percentage (**Fig 9A**). The drivers of current infected cell and net
513 infected cell origin therefore differed completely, highlighting the major differences between observed
514 sequence data and contemporaneous mechanisms generating new infected cells.
515

Figure 9. Transition from replication to proliferation as the dominant mechanism of HIV persistence during ART occurs under a wide range of parameter assumptions. A-C. *See Methods for complete simulated parameter ranges.* **A.** *Local sensitivity analysis (green: current infection, red: net infection) revealed no meaningful difference in percentage of new infected cells generated by viral replication after a year of ART despite variability in initial reservoir volume $I_3(0)$, sanctuary fraction φ_S, and ART effectiveness in and out of the sanctuary (ϵ_S and ϵ). Only an extremely low, or zero, sanctuary decay rate ζ predicted that a meaningful percentage (25%) of infected cells would be newly generated by HIV replication at one year, despite the fact that signals of evolution are not typically observed at this timepoint. Including a high percentage of slowly proliferating naïve CD4+ T cells (T_n) in the reservoir alters the percentage of net, but not current, replication percentage.* **B.** *50 examples from 1,000 global sensitivity analysis simulations. HIV replication accounted for fewer than 25% of current and net infected cells after a year of ART in a majority of simulations.* **C.** *The parameters most correlated with current and net replication percentage at 1 year of ART are different. Current replication percentage inversely correlates with sanctuary decay rate while net (observed) replication percentage positively correlates with reservoir composition (the fraction of naïve latently infected cells). Correlations are measured with a Spearman correlation coefficient.*

To confirm these results, we simulated 10^4 possible patients in a global sensitivity analysis in which all parameter values were simultaneously varied. A rapid transition to proliferation as the source of new infected cells occurred during year one of ART in a majority of simulated patients, and the same variables correlated significantly with net and current replication percentage, respectively (**Fig 9B&C**). Overall, this analysis does not rule out the possibility of a drug sanctuary but does confirm that its relative impact compared to cellular proliferation is likely to be minimal.

Discussion

To eliminate HIV infected cells during prolonged ART, it is necessary to understand the mechanisms by which they persist. In this paper, we used existing data and two methods – inference of HIV clone distributions and mechanistic mathematical modeling – to determine that a majority of infected cell persistence is due to cellular proliferation rather than HIV replication. These conclusions suggest strategies that enhance ART delivery to anatomic drug sanctuaries are less likely to be effective at

546 reducing infected cell burden relative to reservoir reduction strategies. In particular, antiproliferative
547 therapies provide an ideal response to the observed dominance of proliferation.
548
In the first part of the paper, we used existing data to infer the true clonal distributions within the entire
reservoir of HIV sequences in infected participants on long term ART. While the raw data indicate
substantial fractions of *observed* singleton sequences, when the total reservoir size is considered, these
observed singletons are revealed to be predominately members of clonal populations. In fact, the HIV
reservoir appears to be defined by a rank-abundance distribution of clone sizes that can be roughly
approximated as a power-law relationship. This distribution implies that a small number of massive
clones, and a massive number of small clones, comprise a large percentage of sequences.
556
A power-law distribution can be created when a heterogeneous population grows multiplicatively with a
widely variable growth rate.[55] This suggests that the distribution of clone sizes in the reservoir is likely to
have a mechanistic basis. It is plausible, though unproven, that such variable growth arises from rapid
bursts of CD4+ T cell proliferation due to cognate antigen recognition. HIV integration into tumor
suppression genes could also account for some observed clonal dominance.[36,37] Smaller clones may arise
from homeostatic proliferation, or less frequent exposure to smaller amounts of cognate antigen.
563
Another consequence of our inference is that we can more precisely define the mechanism sustaining
equivalent sequences observed in longitudinal samples separated by many years. While we cannot rule
out cellular longevity as a cause of HIV persistence in certain cells, the observation of multiple clonal
sequences could not arise from purely long-lived latently infected cells. In fact, our analysis suggests that
most observed singlet sequences arise from resampling clonal populations that have undergone many
rounds of proliferation.
570
571 The first analysis does not include time-dynamics in the reservoir. Consequently, in the second part of
572 the paper we develop a mechanistic model to reconcile observations from early and late ART. This
573 model is the first to include the three main mechanistic hypotheses for reservoir persistence: an ART
574 sanctuary, long-lived latent cells, and proliferation of latent cells. The model recapitulates known HIV
575 RNA decay kinetics while tracking cells that originate from ongoing replication and cellular proliferation.
576
The model helps to explain how a "fossil record" of evolution would be observed early during ART,
whether or not a small drug sanctuary exists. The model tracks both the fraction of cells that were
generated by viral replication at a given time (current replication percentage) and the fraction that were
generated by viral replication at any time point but are "fossilized" in a long-lived latently infected state
(net, or observed, replication percentage). The net replication percentage remains non-negligible in the
first months of ART even while the current replication percentage drops rapidly. Thus, an observed
sequence that was once created by viral replication (and thus might give a signal of divergence from the
founder virus) can represent a historic replication event rather than current replication. Because time of
detection does not correlate linearly with sequence age, inference of evolution early during ART is
problematic.[20,21] However, the fossil record is transient: within a year of effective ART, observed
phylogenetic data is more likely to represent true reservoir dynamics. Our model agrees with
observations reflecting a lack of contemporaneous HIV evolution after this time.[14,22-27,29,30,36,37]
589
Our sensitivity analysis shows that the major variable correlating with higher observed replication
percentages (a larger proportion of slowly proliferating CD4+ T cells in the reservoir) is not the same
variable that correlates with higher new replication percentages (a slower decrease in sanctuary size).
Replication percentage correlates with the amount of ongoing evolution in viral populations. Without

requiring any phylogenetic simulation, this simple model provides an explanation for evolution during
the first months of ART and no observed HIV evolution in participants with a year of ART.[14,22-27,29,30,36,37]
If we assume a large drug sanctuary and do not allow it to contract as a result of target cell decline, a
persistent low-level sanctuary would emerge that stabilizes at 6 months and generates ongoing
evolution at later ART timepoints. Notably, this has not been observed in clinical studies.
599
Our modeling results inform experiments in two ways. Using rarefaction, we suggest reasonable sample
sizes to verify our hypotheses experimentally (see **Supplementary Fig 4**). We demonstrate that observed
values of sequence richness and clone size, are substantial underestimates. Current studies only sample
the "tip of the iceberg" of the HIV reservoir. Hundreds of thousands of infected cells from a single time
point would be required to capture true reservoir diversity. This sampling depth could only be feasibly
achieved as part of an autopsy study.
606
607 By using dynamical modeling, we also demonstrate that the wash-out period for the fossil record of HIV
608 replication may be up to a year post ART. Thus, we suggest that future reservoir studies are conducted
609 after this time point to avoid observation of historic evolution rather than contemporaneous dynamics.
610
The work presented here carries several important caveats. Current integration site data is still
uncommon and, while robust, is limited to a handful of participants in only a few studies. Modeling rank
abundance curves makes a large assumption about the continuity of the data. The power law model
represents but one approach, and future work should attempt to uncover why that distribution appears
to provide good fit to the data. Extrapolating abundance curves has been criticized: we note that our
attempt to design a simple parametric model was based on the additional information of reservoir size
and our goal to define an upper limit on reservoir richness;[56] we also emphasize that the tail of our
distributions is impossible to precisely characterize with our methods. Our approach is calibrated against
sequence data from blood. However, the dynamics of HIV within lymph tissue may have different
distributions. While historically, blood samples have been taken as a surrogate for HIV infected cells, we
cannot rule out the possibility that the drug sanctuary that does not exchange virus or infected cells with
blood. This sanctuary would be unobservable until probed anatomically. It seems unlikely that such a
sanctuary could be sustained because some trafficking of CD4+ T cells from other compartments seems
necessary to avoid terminal target cell limitation. However, future studies should address possible one-
way trafficking or local proliferation of target cells.
626
627 In conclusion, we demonstrate that the majority of HIV infected cells arise from proliferation after the
628 first year of ART. We have also provided an explanation for incongruent observations of evolution
629 before and after a year of ART. Because proliferation appears to be the dominant force sustaining the
630 HIV reservoir,[34] we suggest limiting proliferation as a prime therapeutic target.[10,11,57]

Methods

631
Rank abundance of HIV integration sites. We used an ecological framework to study the abundance of
clonal HIV. To do so, we applied methods to integration site and replication competent HIV sequence
data. Cellular DNA found with HIV integrated into different integration sites in the human genome were
defined as distinct "clones". The number of times a cell was found with the same integration site added
to the "abundance" of that clone. By ordering (ranking) the clones from largest to smallest by
abundance, we developed a rank abundance curve, $a(r)$, for each participant time point. No
assumptions were made about the stability or dynamics of the reservoir rank abundance over time.
639

In our analysis of data from Wagner *et al.*,[37] we combine measurements taken closely in time and use
the median time point as done in that published paper. In our analysis of Maldarelli *et al.*,[36] we grouped
by integration site, or nearest measured integration site when integration site was not noted. It is
important to note that the methods used by Wagner *et al.* and Maldarelli *et al.* are slightly different. The
ISLA method used by Wagner *et al.* is lower throughput than the next generation shotgun sequencing
method used by Maldarelli *et al.* The absolute number of viruses identified by each group therefore
differs. However, the percentage of observed singletons is similar between the two studies.

We manually counted the abundance of replication competent HIV sequences using phylogenetic trees
in Hosmane *et al.*[34]

Calculation of rarefaction curves. We used rarefaction curves to estimate the expected number of
distinct sequences that would still be present in a subsample of k sequences from the observed data
with sample size of N:

$$\langle n_k \rangle = R^{obs} - \binom{N}{k}^{-1} \sum_{r=1}^{R^{obs}} \binom{N-a(r)}{k},$$ (1)

where the parentheses indicate binomial coefficients, e.g. $\binom{N}{k} = \frac{N!}{k!(N-k)!}$. Later, we extrapolated
rarefaction curves using the modeled distributions for the total reservoir size L. Because the number of
samples we allowed was orders of magnitude smaller than the number of cells in the reservoir, $k \ll L$,
we used Stirling's approximation to simplify the binomial coefficients. The expected number of
sequences after k samples is then

$$\langle \tilde{n}_k \rangle = R - \sum_{r=1}^{R} \left[1 - \frac{a(r)}{L} \right]^k,$$ (2)

an expression which avoids computation of large factorials (derivation in the **Supplementary Methods**).

Nonparametric estimation of species richness. We employed the Chao1 estimator to set a lower bound
on the sequence or integration site richness.[58] A derivation of the estimator is included in the
Supplementary Methods. Chao1 is not a mechanistic model and requires no free parameters. Inference
relies on only the number of observed singleton (N_1) and observed doubleton (N_2) sequences such that

$$R^{Chao1} = R^{obs} + \frac{N_1(N_1-1)}{2(N_2+1)}.$$ (3)

We display an asymmetric confidence interval in **Fig 3** (see Chao *et al.*[58] or **Supplementary Methods** for
the calculation). We also note it is possible the data are undersampled to the extent that a one-sided
confidence interval may be more appropriate. Thus, for our biological conclusions we take the Chao1
point estimate as a lower bound, and constrain the upper bound using the parametric model (**Eq 4**).
Other richness estimators (jackknife 1 and 2) were tested but provided similar and consistently lower
estimates of richness than the Chao1 estimator. These were not included in our results because the
Chao1 was interpreted as a lower bound on true sequence richness.

Parametric models to extrapolate sequence abundance curves. Estimates of the size of the HIV
reservoir (both replication competent and total) were gathered from the published literature.[33] We then
developed a parametric model to quantify the true rank abundance distribution of the complete HIV
reservoir. Examination of the data indicated a possible log-log-linear relationship, so we chose a discrete

integer power law model so that the probability of a rank is described by $p(r) = \psi(R)r^{-\alpha}$ where the
coefficient $\psi(R) = \sum_{r=1}^{R} r^{-\alpha}$ is the normalization constant for the power law. Then, to describe the true
rank abundance $a(r)$ we chose the reservoir size depending on the model context (replication
competent $L = 10^7$ or total HIV DNA $L = 10^9$). To ensure integer number of cells, we rounded this
distribution, and forced the total number of cells to equal the reservoir size. That is,

$a(r; \alpha, R, L) = |[L\psi(R)r^{-\alpha}]|$ (4)

where $|[]|$ indicates rounding to the nearest integer. Thus, our model depended on two free
parameters, a power law exponent α, and the reservoir richness R. Other functional forms were
explored but simplicity and accurate reproduction of the data were optimal with the power law.

Fitting the rank-abundance model to experimental data. Using the experimental data we found the
best-fit model using the following procedure. We fixed the reservoir size L depending on the model
context (replication competent or total HIV DNA). We chose a value for R and α from ranges $R \in$
$[10^3, 10^7]$ and $\alpha \in [0,2]$ to specify the model. Then, we sampled the extrapolated distribution 10 times
using multinomial sampling with the same number of samples as the experimental data being fit,
$\mathcal{M}(N^{obs}, p(r))$. This procedure assumes that sampling cells does not change the distribution of the
reservoir, which is reasonable given the reservoir size. Each sampled data set was compared to the
experimental data by computing the residual sum of squares (rss) error of the cumulative proportional
abundance (cpa) curves. For each model then, the reported error is the average rss over the 10
resamplings. Because the rss error is not symmetric across the domain of the cpa, this approach
becomes similar to minimizing the Kolmogorov-Smirnov (KS) statistic: the maximum deviation between
two cumulative distributions. For each experimental data set, 2500 model parameter sets were
generated, and fitting results are visualized as heat maps (see **Figs 4A, 5A** for example). Because the
procedure becomes computationally expensive as $R > 10^7$, we did not explore values above this
threshold. In theory, it is possible to have a distribution with all clones having a single member $R =$
$L, \alpha = 0$. For the total DNA reservoir, this value would result in $R = 10^9$. However, this model was never
optimal. In fact, as richness increased beyond $R \approx 10^6$, the model was no longer sensitive to R. Thus, it
appeared that finding the best fit α was sufficient to specify the model if proper bounds on richness
were included.

We excluded models where $R < R^{Chao1}$, but we also sought to identify an upper bound for R. Indeed,
certain model parameter combinations are mathematically impossible. For example, for a given power
law exponent, the richness is constrained below a certain value for a given reservoir size. This
observation has been considered previously in ecology under the terminology of 'feasible sets'.[59] To
determine the largest possible richness that still has the best fit, we chose the roughly constant value of
α that emerged when R was large enough to be unidentifiable. Then, we noted that for large R it is a
reasonable approximation to allow $\sum_{r=1}^{R} a(r) = \int_{1}^{R} a(r) \, dr$. R is thus approximately bounded, and we
solved for the maximal value or the upper bound on the richness given the best fit α and the chosen L. A
discussion and numerical validation of this approximation is presented in the **Supplementary Methods**
and **Supplementary Fig 2**. The upper bound provides the sequence abundance most permissive of true
singleton sequences – the reservoir with the most evidence of HIV replication as opposed to
proliferation. In extrapolated reservoirs, we used the maximum richness model to ensure we were
biasing the results as strongly as possible against our own hypothesis.

Model fitting validation with simulated data. A discussion and demonstration of model validation is included in the **Supplementary Methods** and **Supplementary Fig 1**. The exercise shows that simply fitting a power law to the experimental data (using log-log-linear regression) without the extra sampling step necessarily underestimates the power-law exponent, demonstrating the utility of our approach. Moreover, it shows that a published maximum likelihood approach[60] is not as accurate for these data as our resampling approach (code hosted at http://tuvalu.santafe.edu/~aaronc/powerlaws/ last accessed July 2018) . We simulated a reservoir with known power law exponent **Supplementary Fig 1A** and tested for recovery of this known value. The fitting validation proceeded identically to the data fitting, 2500 distributions were generated (225 examples are shown in **Supplementary Fig 1D**), the simulated data was sampled **Supplementary Fig 1B**, and reranked **Supplementary Fig 1C**. Fitting results **Supplementary Fig 1E&F** are shown analogous to **Figs 4&5,A&B**. Finally, the most correct parameter estimation of three methods tried came from our modeling approach **Supplementary Fig 1G**.

Mechanistic model for the persistence of the HIV reservoir. The canonical model for HIV dynamics describes the time-evolution of the concentrations of susceptible S and infected I CD4+ T cells and HIV virus V.[50,54,61] Our model grows from the canonical model, simplifying with several approximations and extending the biological detail to simulate HIV dynamics on ART, including a long-lived latent reservoir and a potential drug sanctuary. Perelson *et al.* first noticed and quantified a 'biphasic' clearance of HIV virus upon initiation of ART and showed that viral half-lives of 1.5 and 14 days correspond with the half-lives of two infected cell compartments.[50,54] With longer observation times and single-copy viral assays, Palmer *et al.* found four-phases of viral clearance after initiation of ART.[51] Because of uncertainty in distinguishing the third and fourth phase in that study, we focus on the first three decay rates and corresponding cellular compartments, attributing a mixture of the third and fourth phase decay to the clearance of the productively infectious latent reservoir (half-life 44 months) as measured by Siliciano *et al.* and recently corroborated by Crooks *et al*.[2,3] and the clearance of HIV DNA.[47] We developed a mechanistic mathematical model that has three types of infected cells I_1, I_2, I_3 that are meant to simulate productively infected cells, pre-integration infected cells, and latently infected cells, respectively. We classify rapid death δ_1 and viral production within actively infected cells I_1. Cells with longer half-life that may represent pre-integration infected cells I_2 are activated to I_1 at rate ξ_2. I_2 may represent CD4+ T cells with a prolonged pre-integration phase, but their precise biology does not affect model outcomes.[48]

The state $I_{3(j)}$ represents latently infected reservoir cells of phenotype j, which contain a single chromosomally integrated HIV DNA provirus.[44] I_3 reactivates to I_1 at rate ξ_3 which at present is assumed to be constant across cell phenotypes.[49] The probabilities of a newly infected cell entering $I_1, I_2, I_{3(j)}$, are $\tau_1, \tau_2, \tau_{3(j)}$. Because we are focused on the role of proliferation, we assume sub-populations of I_3,[12] including effector memory (T$_{em}$), central memory (T$_{cm}$), and naïve (T$_n$) CD4+ T cells, which proliferate and die at different rates $\alpha_{3(j)}, \delta_{3(j)}$.[12,42,43] Parameter values and initial conditions for the model are collected in **Table 1**.

Including a decreasing sanctuary in the model. A recent hypothesis about reservoir persistence suggests there may be a small, anatomic sanctuary (1 in 10^5 infected cells) in which ART is not therapeutic.[4] Thus, we included the state variable I_S that is maintained at a constant set-point level prior to ART, where all new infected cells arise from ongoing replication. We opted for this simplification because it biased against our conclusions. The amount of virus produced by the sanctuary V_S is extremely low relative to non-sanctuary regions because ART results in levels undetectable by sensitive assays.[51]

Many studies have demonstrated that HIV accelerates immunosenescene through abnormal activation
of CD4+ T cells.[62-64] ART results in a marked reduction of T cell activation and apoptosis, a potential
signature of HIV susceptible cells.[65] By examining the decline of activation markers for CD4+ T cells, we
approximated the decay kinetics of activated T cells upon ART, inferring approximate decay kinetics of
the target cells in our model.[52,53,66] A range of initial values exists (from ~5–20% activation) depending
on stage of HIV infection, yet after a year of ART, a large percentage of patients return to almost normal,
or slightly elevated CD4+ T cell activation levels (2-3%).[52] Because we assume that target cell depletion is
minimal at viral load set-point, we can approximate that the susceptible cell concentration decreases
over time as the immune activation decreases, i.e., $S = S(0)e^{-\zeta t}$. This single exponential decay is
simplified (it may be biphasic but the data are not granular enough to discriminate this dynamic
subtlety). From existing data, the decay constant should be in the range $\zeta \sim [0.002, 0.01]$ day^{-1}.[52,66] We
extend this decay into the sanctuary, allowing the number of susceptible cells over the whole body to
decrease so that we have $I_S = I_1(0)\varphi_S e^{-\zeta t}$ where φ_S is the fraction of infected cells that are in a
sanctuary. Model simulations are also performed without this assumption of target cell contraction.

Last, we use the quasi-static approximation that virus is proportional to the number of actively infected
cells in all compartments $V = n(I_1 + I_S)$ where $n = \pi/\gamma$, the ratio of the viral production rate to the
viral clearance rate (**Table 1**). The model is thus

$\dot{I}_1 = \tau_1 \beta_\epsilon SV - \delta_1 I_1 + \xi_2 I_2 + \sum_j \xi_3 I_{3(j)}$
$\dot{I}_2 = \tau_2 \beta_\epsilon SV + (\alpha_2 - \delta_2 - \xi_2)I_2$ (5)
$\dot{I}_{3(j)} = \tau_{3(j)} \beta_\epsilon SV + (\alpha_{3(j)} - \delta_{3(j)} - \xi_3)I_{3(j)}$,

where we use the over-dot to denote the time derivative.

Comparing proliferation and viral replication: 'net' and 'current' percentages. By solving the ODE
model (**Eq 6**), we have the time solution for each infected cell state. From these, we can compute the
total number of newly infected cells generated in a given time interval Δt by ongoing replication. That
value is $I^{rep}(t) = (\beta_\epsilon SV + \phi_S \beta SV_S)\Delta t$. The total number of newly infected cells generated by
proliferation of a previously infected cell can be computed similarly in a time interval as $I^{pro}(t) =$
$\sum_{i(j)} \alpha_{i(j)} I_{i(j)} \Delta t$. Therefore, the percentage of infected cells generated by current replication is written

$\Phi^{current}(t) = 100 \cdot \dfrac{I^{rep}(t)}{I^{rep}(t) + I^{pro}(t)}.$ (6)

We can further subset this newly generated fraction by examining the percentage of newly infected cells
that enter the long-lived latent state I_3 by defining $I^{rep(3)}(t) = \tau_3(\beta_\epsilon SV + \phi_S \beta SV_S)\Delta t$ and $I^{pro(3)}(t) =$
$\sum_j \alpha_{3(j)} I_{3(j)} \Delta t$ so that

$\Phi^{current(3)}(t) = 100 \cdot \dfrac{I^{rep(3)}(t)}{I^{rep(3)}(t) + I^{pro(3)}(t)}.$ (7)

The net (or observed) replication percentage, is the fraction of cells that remain that were once
generated by viral replication. To compute this quantity, we use an additional set of ODEs that we refer
to as "tracking equations" because they do not change the dynamics of the system, and only are used to
track specific variables. To denote the net value as opposed to new value we use a subscript Σ. The net
cells generated by viral replication in state i of phenotype j is governed by the differential equation
825

$$\dot{I}_{i(j)}{}^{(\Sigma)rep} = \tau_{i(j)}\beta_\epsilon SV - (\delta_{i(j)} - \xi_{i(j)})I_{i(j)}{}^{(\Sigma)rep}.$$ (8)

Likewise, the net cells generated by proliferation in state i of phenotype j is governed by the differential
equation

$$\dot{I}_{i(j)}{}^{(\Sigma)pro} = \alpha_{i(j)}I_{i(j)} - (\delta_{i(j)} - \xi_{i(j)})I_{i(j)}{}^{(\Sigma)pro}.$$ (9)

We note that because we only allow these two mechanisms, $\dot{I}_{i(j)} = \dot{I}_{i(j)}{}^{(\Sigma)rep} + \dot{I}_{i(j)}{}^{(\Sigma)pro}$ and
$I_{i(j)}(t) = I_{i(j)}^{(\Sigma)rep}(t) + I_{i(j)}^{(\Sigma)pro}$. By solving the tracking equations separately, we can then find the net
replication percentage by summing over cell types and phenotypes to ultimately write

$$\Phi^\Sigma(t) = 100 \cdot \frac{\sum_{i(j)} I_{i(j)}^{(\Sigma)rep}(t)}{\sum_{i(j)} I_{i(j)}^{(\Sigma)rep}(t) + I_{i(j)}^{(\Sigma)pro}(t)}.$$ (10)

In all simulations, we assumed that 100% of infected cells at the initiation of ART were generated by
viral replication, that is $\Phi^\Sigma(0) = 100$. This assumption biases results in favor of replication. However,
we choose it because, to the best of our knowledge, studies of proliferation during chronic untreated
HIV have not been performed.

Sensitivity analysis. Using estimated parameter bounds [lower, upper], we completed a local and global
sensitivity analysis. These ranges were chosen to cover a wide range of possible assumptions. We
allowed $I_3(0) = [0.02,2]$ cells μL^{-1}, $\varphi_S = [10^{-6}, 10^{-4}]$ unitless, $\zeta = [0,0.2]$ day^{-1}, $\epsilon = [0.9,0.99]$
unitless, $\epsilon_S = [0,0.9]$ unitless, $I_{3(n)}(0) = [0,0.5] \times I_3(0)$ cells μL^{-1}. For the local analysis, we used all
values as in **Table 1** and modified one parameter at a time over each listed range above. The global
analysis was performed by using 10^4 Latin Hypercube samplings of the complete 6-dimensional
parameter space.[67] The key outcome, the replication percentage (net and current) at 1 year of ART, was
correlated to each parameter using the Spearman correlation coefficient—defined by the ratio of the
covariance between the outcome and the variable divided by the standard deviations of each when the
variables were rank-ordered by value.

Data and code availability. Computational code for all calculations and simulations was performed in
Python and Matlab and can be found at **https://github.com/dbrvs/reservoir_persistence**. Sequence
data was obtained from the Retrovirus Integration Database (RID).[68]

References

1. Volberding, P.A. & Deeks, S.G. Antiretroviral therapy and management of HIV infection.
Lancet **376**, 49-62 (2010).
2. Crooks, A.M.*, et al.* Precise Quantitation of the Latent HIV-1 Reservoir: Implications for
Eradication Strategies. *J Infect Dis* **212**, 1361-1365 (2015).
3. Siliciano, J.D.*, et al.* Long-term follow-up studies confirm the stability of the latent
reservoir for HIV-1 in resting CD4+ T cells. *Nat Med* **9**, 727-728 (2003).
4. Lorenzo-Redondo, R.*, et al.* Persistent HIV-1 replication maintains the tissue reservoir
during therapy. *Nature* **530**, 51-56 (2016).

5.   Gunthard, H.F.*, et al.* Evolution of envelope sequences of human immunodeficiency
virus type 1 in cellular reservoirs in the setting of potent antiviral therapy. *J Virol* **73**,
9404-9412 (1999).
6.   Finzi, D.*, et al.* Identification of a reservoir for HIV-1 in patients on highly active
antiretroviral therapy. *Science* **278**, 1295-1300 (1997).
7.   Finzi, D.*, et al.* Latent infection of CD4+ T cells provides a mechanism for lifelong
persistence of HIV-1, even in patients on effective combination therapy. *Nat Med* **5**,
512-517 (1999).
8.   Fletcher, C.V.*, et al.* Persistent HIV-1 replication is associated with lower antiretroviral
drug concentrations in lymphatic tissues. *Proc Natl Acad Sci U S A* **111**, 2307-2312
(2014).
9.   Archin, N.M.*, et al.* Administration of vorinostat disrupts HIV-1 latency in patients on
antiretroviral therapy. *Nature* **487**, 482-485 (2012).
10.   Chapuis, A.G.*, et al.* Effects of mycophenolic acid on human immunodeficiency virus
infection in vitro and in vivo. *Nat Med* **6**, 762-768 (2000).
11.   Garcia, F.*, et al.* Effect of mycophenolate mofetil on immune response and plasma and
lymphatic tissue viral load during and after interruption of highly active antiretroviral
therapy for patients with chronic HIV infection: a randomized pilot study. *J Acquir*
Immune Defic Syndr **36**, 823-830 (2004).
12.   Chomont, N.*, et al.* HIV reservoir size and persistence are driven by T cell survival and
homeostatic proliferation. *Nat Med* **15**, 893-900 (2009).
13.   Maldarelli, F.*, et al.* HIV populations are large and accumulate high genetic diversity in a
nonlinear fashion. *J Virol* **87**, 10313-10323 (2013).
14.   Nickle, D.C.*, et al.* Evolutionary indicators of human immunodeficiency virus type 1
reservoirs and compartments. *J Virol* **77**, 5540-5546 (2003).
15.   Sanjuan, R. & Domingo-Calap, P. Mechanisms of viral mutation. *Cell Mol Life Sci* **73**,
4433-4448 (2016).
16.   Poon, A.F.*, et al.* Reconstructing the dynamics of HIV evolution within hosts from serial
deep sequence data. *PLoS Comput Biol* **8**, e1002753 (2012).
17.   Zanini, F.*, et al.* Population genomics of intrapatient HIV-1 evolution. *Elife* **4**(2015).
18.   Shankarappa, R.*, et al.* Consistent viral evolutionary changes associated with the
progression of human immunodeficiency virus type 1 infection. *J Virol* **73**, 10489-10502
(1999).
19.   Lemey, P.*, et al.* Synonymous substitution rates predict HIV disease progression as a
result of underlying replication dynamics. *PLoS Comput Biol* **3**, e29 (2007).
20.   Kearney, M.F.W., A.; Shao, W.; McManus W.R.; Bale, M.J.; Luke, B.; Maldarelli, F.;
Mellors, J.M.; Coffin, J.M. Ongoing HIV Replication During ART Reconsidered. *Open*
Forum Infectious Diseases **4**, ofx173 (2017).
21.   Rosenbloom, D.I.S., Hill, A.L., Laskey, S.B. & Siliciano, R.F. Re-evaluating evolution in the
HIV reservoir. *Nature* **551**, E6-E9 (2017).
22.   Evering, T.H.*, et al.* Absence of HIV-1 evolution in the gut-associated lymphoid tissue
from patients on combination antiviral therapy initiated during primary infection. *PLoS*
Pathog **8**, e1002506 (2012).

23.   Frenkel, L.M.*, et al.* Multiple viral genetic analyses detect low-level human
immunodeficiency virus type 1 replication during effective highly active antiretroviral
therapy. *J Virol* **77**, 5721-5730 (2003).
24.   Josefsson, L.*, et al.* The HIV-1 reservoir in eight patients on long-term suppressive
antiretroviral therapy is stable with few genetic changes over time. *Proc Natl Acad Sci U*
S A **110**, E4987-4996 (2013).
25.   Kearney, M.F.*, et al.* Lack of detectable HIV-1 molecular evolution during suppressive
antiretroviral therapy. *PLoS Pathog* **10**, e1004010 (2014).
26.   Rothenberger, M.K.*, et al.* Large number of rebounding/founder HIV variants emerge
from multifocal infection in lymphatic tissues after treatment interruption. *Proc Natl*
Acad Sci U S A **112**, E1126-1134 (2015).
27.   Brodin, J.*, et al.* Establishment and stability of the latent HIV-1 DNA reservoir. *Elife*
5(2016).
28.   Bull, M.E.*, et al.* Monotypic human immunodeficiency virus type 1 genotypes across the
uterine cervix and in blood suggest proliferation of cells with provirus. *J Virol* **83**, 6020-
6028 (2009).
29.   von Stockenstrom, S.*, et al.* Longitudinal Genetic Characterization Reveals That Cell
Proliferation Maintains a Persistent HIV Type 1 DNA Pool During Effective HIV Therapy. *J*
Infect Dis **212**, 596-607 (2015).
30.   Wagner, T.A.*, et al.* An increasing proportion of monotypic HIV-1 DNA sequences during
antiretroviral treatment suggests proliferation of HIV-infected cells. *J Virol* **87**, 1770-
1778 (2013).
31.   Alizon, S. & Fraser, C. Within-host and between-host evolutionary rates across the HIV-1
genome. *Retrovirology* **10**, 49 (2013).
32.   Bruner, K.M.*, et al.* Defective proviruses rapidly accumulate during acute HIV-1
infection. *Nat Med* **22**, 1043-1049 (2016).
33.   Ho, Y.C.*, et al.* Replication-competent noninduced proviruses in the latent reservoir
increase barrier to HIV-1 cure. *Cell* **155**, 540-551 (2013).
34.   Hosmane, N.N.*, et al.* Proliferation of latently infected CD4+ T cells carrying replication-
competent HIV-1: Potential role in latent reservoir dynamics. *J Exp Med* **214**, 959-972
(2017).
35.   Joos, B.*, et al.* HIV rebounds from latently infected cells, rather than from continuing
low-level replication. *Proc Natl Acad Sci U S A* **105**, 16725-16730 (2008).
36.   Maldarelli, F.*, et al.* HIV latency. Specific HIV integration sites are linked to clonal
expansion and persistence of infected cells. *Science* **345**, 179-183 (2014).
37.   Wagner, T.A.*, et al.* HIV latency. Proliferation of cells with HIV integrated into cancer
genes contributes to persistent infection. *Science* **345**, 570-573 (2014).
38.   Boritz, E.A.*, et al.* Multiple Origins of Virus Persistence during Natural Control of HIV
Infection. *Cell* **166**, 1004-1015 (2016).
39.   Cohn, L.B.*, et al.* HIV-1 integration landscape during latent and active infection. *Cell* **160**,
420-432 (2015).
40.   Simonetti, F.R.*, et al.* Clonally expanded CD4+ T cells can produce infectious HIV-1 in
vivo. *Proc Natl Acad Sci U S A* **113**, 1883-1888 (2016).

41.   Schroder, A.R.*, et al.* HIV-1 integration in the human genome favors active genes and
954   local hotspots. *Cell* **110**, 521-529 (2002).

955   42.   Macallan, D.C.*, et al.* Rapid turnover of effector-memory CD4(+) T cells in healthy
956   humans. *J Exp Med* **200**, 255-260 (2004).

957   43.   McCune, J.M.*, et al.* Factors influencing T-cell turnover in HIV-1-seropositive patients. *J*
958   *Clin Invest* **105**, R1-8 (2000).

959   44.   Josefsson, L.*, et al.* Single cell analysis of lymph node tissue from HIV-1 infected patients
960   reveals that the majority of CD4+ T-cells contain one HIV-1 DNA molecule. *PLoS Pathog*
961   **9**, e1003432 (2013).

962   45.   Eren, M.I., Chao, A., Hwang, W.H. & Colwell, R.K. Estimating the richness of a population
963   when the maximum number of classes is fixed: a nonparametric solution to an
964   archaeological problem. *PLoS One* **7**, e34179 (2012).

965   46.   Seymour, A.M. Imaging cardiac metabolism in heart failure: the potential of NMR
966   spectroscopy in the era of metabolism revisited. *Heart Lung Circ* **12**, 25-30 (2003).

967   47.   Besson, G.J.*, et al.* HIV-1 DNA decay dynamics in blood during more than a decade of
968   suppressive antiretroviral therapy. *Clin Infect Dis* **59**, 1312-1321 (2014).

969   48.   Cardozo, E.F.*, et al.* Treatment with integrase inhibitor suggests a new interpretation of
970   HIV RNA decay curves that reveals a subset of cells with slow integration. *PLoS Pathog*
971   **13**, e1006478 (2017).

972   49.   Hill, A.L., Rosenbloom, D.I., Fu, F., Nowak, M.A. & Siliciano, R.F. Predicting the outcomes
973   of treatment to eradicate the latent reservoir for HIV-1. *Proc Natl Acad Sci U S A* **111**,
974   13475-13480 (2014).

975   50.   Perelson, A.S.*, et al.* Decay characteristics of HIV-1-infected compartments during
976   combination therapy. *Nature* **387**, 188-191 (1997).

977   51.   Palmer, S.*, et al.* Low-level viremia persists for at least 7 years in patients on suppressive
978   antiretroviral therapy. *Proc Natl Acad Sci U S A* **105**, 3879-3884 (2008).

979   52.   Hunt, P.W.*, et al.* T cell activation is associated with lower CD4+ T cell gains in human
980   immunodeficiency virus-infected patients with sustained viral suppression during
981   antiretroviral therapy. *J Infect Dis* **187**, 1534-1543 (2003).

982   53.   Kaufmann, G.R.*, et al.* Rapid restoration of CD4 T cell subsets in subjects receiving
983   antiretroviral therapy during primary HIV-1 infection. *AIDS* **14**, 2643-2651 (2000).

984   54.   Perelson, A.S., Neumann, A.U., Markowitz, M., Leonard, J.M. & Ho, D.D. HIV-1 dynamics
985   in vivo: virion clearance rate, infected cell life-span, and viral generation time. *Science*
986   **271**, 1582-1586 (1996).

987   55.   M., M. A brief history of generative models for power law and lognormal distributions.
988   *Internet Mathematics* **1(2)**, 226-251. (2004).

989   56.   Willis, A. Extrapolating abundance curves has no predictive power for estimating
990   microbial biodiversity. *Proc Natl Acad Sci U S A* **113**, E5096 (2016).

991   57.   Reeves, D.B.*, et al.* Anti-proliferative therapy for HIV cure: a compound interest
992   approach. *Sci Rep* **7**, 4011 (2017).

993   58.   Chao, A. Estimating the population size for capture-recapture data with unequal
994   catchability. *Biometrics* **43**, 783-791 (1987).

995   59.   Locey, K.J. & White, E.P. How species richness and total abundance constrain the
996   distribution of abundance. *Ecol Lett* **16**, 1177-1185 (2013).

60. Clauset, A.S., C. R.; Newman, M. E. . Power-law distributions in empirical data. *SIAM*
Review **Nov 6;51(4)**, 661-703. (2009).

61. Perelson, A.S., Kirschner, D.E. & De Boer, R. Dynamics of HIV infection of CD4+ T cells.
Math Biosci **114**, 81-125 (1993).

62. Rutishauser, R.L*., et al.* Early and Delayed Antiretroviral Therapy Results in Comparable
Reductions in CD8+ T Cell Exhaustion Marker Expression. *AIDS Res Hum Retroviruses*
(2017).

63. Serrano-Villar, S*., et al.* HIV-infected individuals with low CD4/CD8 ratio despite effective
antiretroviral therapy exhibit altered T cell subsets, heightened CD8+ T cell activation,
and increased risk of non-AIDS morbidity and mortality. *PLoS Pathog* **10**, e1004078
(2014).

64. Cockerham, L.R*., et al.* Programmed death-1 expression on CD4(+) and CD8(+) T cells in
treated and untreated HIV disease. *AIDS* **28**, 1749-1758 (2014).

65. Appay, V. & Sauce, D. Immune activation and inflammation in HIV-1 infection: causes
and consequences. *J Pathol* **214**, 231-241 (2008).

66. Autran, B*., et al.* Positive effects of combined antiretroviral therapy on CD4+ T cell
homeostasis and function in advanced HIV disease. *Science* **277**, 112-116 (1997).

67. Sanchez, M.A. & Blower, S.M. Uncertainty and sensitivity analysis of the basic
reproductive rate. Tuberculosis as an example. *Am J Epidemiol* **145**, 1127-1137 (1997).

68. Shao, W*., et al.* Retrovirus Integration Database (RID): a public database for retroviral
insertion sites into host genomes. *Retrovirology* **13**, 47 (2016).

69. Ribeiro, R.M*., et al.* Estimation of the initial viral growth rate and basic reproductive
number during acute HIV-1 infection. *J Virol* **84**, 6096-6102 (2010).

70. Huang, Y., Liu, D. & Wu, H. Hierarchical Bayesian methods for estimation of parameters
in a longitudinal HIV dynamic system. *Biometrics* **62**, 413-423 (2006).

71. Luo, R., Piovoso, M.J., Martinez-Picado, J. & Zurakowski, R. HIV model parameter
estimates from interruption trial data including drug efficacy and reservoir dynamics.
PLoS One **7**, e40198 (2012).

72. Conway, J.M. & Perelson, A.S. Residual Viremia in Treated HIV+ Individuals. *PLoS*
Comput Biol **12**, e1004677 (2016).

73. Ramratnam, B*., et al.* Rapid production and clearance of HIV-1 and hepatitis C virus
assessed by large volume plasma apheresis. *Lancet* **354**, 1782-1785 (1999).

74. Markowitz, M*., et al.* A novel antiviral intervention results in more accurate assessment
of human immunodeficiency virus type 1 replication dynamics and T-cell decay in vivo. *J*
Virol **77**, 5037-5038 (2003).

75. Blankson, J.N*., et al.* Biphasic decay of latently infected CD4+ T cells in acute human
immunodeficiency virus type 1 infection. *J Infect Dis* **182**, 1636-1642 (2000).

1034

1035 ## End Notes

1036 The authors declare no competing interests.

1037

1038 **Acknowledgments**. DBR thanks F Boshier and O Hyrien for many illuminating conversations and JM
1039 Drake for originally suggesting the Chao1 estimator. DBR is supported by a Washington Research

1040 Foundation (WRF) Postdoctoral Fellowship. ERD is supported by the National Center for Advancing
1041 Translational Sciences of the NIH under Award Number KL2 TR002317. JTS is supported by NIH grants:
1042 P01 AI030371-24, U19 AI113173-02, R01 AI121129-01, UM1 AI126623, UM1 AI068635.

1043

Author contributions statement. DBR, ERD, and JTS conceived the study. TAW, SEP, and AMS
contributed ideas and data sources for the project. DBR assembled data, wrote all code, performed all
calculations, ran the models, and analyzed output data. JTS and DBR wrote the manuscript with
contributions from all other authors.

Tables

Table 1: Model parameters

Parameter	Value	Meaning	Units	Source
R_0	8	Basic reproductive number of HIV	[]	[69]
β_0	2×10^{-4}	Viral infectivity, used in $\beta_\epsilon = \beta_0(1-\epsilon)$	[μL copy^{-1} day^{-1}]	[61,70,71]
ϵ	0.95	ART efficacy outside the sanctuary	[]	[71,72]
π	10^3	Viral production rate, used in $n = \pi/\gamma$	[μL copy^{-1} day^{-1}]	[61,70,71]
γ	23	Viral clearance rate, used in $n = \pi/\gamma$	[day^{-1}]	[73]
α_S	150	Susceptible cell production rate	[μL copy^{-1} day^{-1}]	[54,70,71]
δ_S	0.2	Susceptible cell death rate	[day^{-1}]	[61,70,71]
δ_1	0.8	Productively infected cell (I_1) clearance rate	[day^{-1}]	[71,74]
δ_2	0.02	Pre-integration cell (I_2) death rate	[day^{-1}]	[48,50]
α_2	0.047	Pre-integration cell proliferation rate	[day^{-1}]	[42]
ξ_2	0.08	Pre-integration cell activation rate	[day^{-1}]	Fit
$\alpha_{3(j)}$	[0.047,0.015, 0.002]	Proliferation rate of latently infected cells $j \in$ [T$_{em}$, T$_{cm}$, T$_n$] phenotypes, respectively	[day^{-1}]	[42]
ξ_3	0.0003	Latent cell activation rate (for all j)	[day^{-1}]	Fit
$\delta_{3(j)}$		Calculated from latent clearance rate as $\theta_L = \alpha_{3(j)} - \delta_{3(j)} - \xi_3$ where $\theta_L = -5.2 \times 10^{-4}$	[day^{-1}]	[2,3]
$\tau_{i(j)}$	[1, 10^{-2}, 10$^{-4}\varrho_j$]	Probability of infection of each compartment, taken from y-intercepts in Ref. 50	[]	[51]
ϱ_j	[0.2,0.75,0.05]	Fraction of latent infected cells of each phenotype (e.g. from patient #5 in Ref. 12)	[cells μL^{-1}]	[12]
$V(0)$	10^2	Initial viral load (from typical set-point value 10^5 copies/mL)	[copy μL^{-1}]	[69]
$I_1(0)$	2	Initial concentration of productively infected cells, calculated from $I_1(0) = V(0)/n$	[cells μL^{-1}]	[75]
$I_2(0)$	0.2	Initial concentration of pre-integration infected cells	[cells μL^{-1}]	[75]
$I_{3(j)}(0)$	$0.2\varrho_j$	Initial concentration of each latent phenotype, calculated from ~10^6 latently infected cells in ~5L of blood	[cells μL^{-1}]	[2,12]
$I_S(0)$	180	Initial concentration of sanctuary cells, calculated from equilibrium model $I_S(0) = \frac{\alpha_S}{\delta_1} - \frac{\delta_S}{n\beta_0(1-\epsilon_S)}$, e.g. Ref. 56 SI	[cells μL^{-1}]	Calc
ζ	0.007	Decay rate of T cell activation	[day^{-1}]	[52]
ϵ_S	0	ART efficacy in the sanctuary, minimum value	[]	Min
φ_S	10^{-5}	Fraction of cells in sanctuary	[]	[4]

1050