PathCORE: visualizing globally co-occurring pathways in large transcriptomic compendia

Kathleen M. Chen[1], Jie Tan[2], Gregory P. Way[1], Georgia Doing[3], Deborah A. Hogan[3], Casey S. Greene[1,*]

[1] Department of Systems Pharmacology and Translational Therapeutics. Perelman School of Medicine. University of Pennsylvania. Philadelphia PA. 19104
[2] Department of Molecular and Systems Biology. Geisel School of Medicine at Dartmouth. Hanover NH. 03755
[3] Department of Microbiology and Immunology. Geisel School of Medicine at Dartmouth. Hanover NH. 03755

[*] To whom correspondence should be addressed.
Casey Greene
University of Pennsylvania
3400 Civic Center Blvd.
Philadelphia PA 10104
Phone: 215-573-2991
Email: csgreene@upenn.edu

Abstract

Background: Investigators often interpret genome-wide data by analyzing the expression levels of genes within pathways. While this within-pathway analysis is routine, the products of any one pathway can affect the activity of other pathways. Past efforts to identify relationships between biological processes have evaluated overlap in knowledge bases or evaluated changes that occur after specific treatments. Individual experiments can highlight condition-specific pathway-pathway interactions; however, constructing a complete network of such relationships across many conditions requires analyzing results from many studies.
Results: We developed the PathCORE software to predict global pathway-pathway interactions, i.e. those evident across a broad data compendium. PathCORE starts with the results of robust feature construction algorithms, which are now being developed and applied to transcriptomic data. PathCORE identifies pathways grouped together in features more than expected by chance as functionally *co-occurring*. We performed example analyses using PathCORE for a microbial compendium for which eADAGE features were already available and a TCGA dataset of 33 cancer types that we analyzed via NMF. PathCORE recapitulated previously described pathway-pathway interactions and suggested additional edges with biological plausibility that still remain to be explored. The software also identifies genes associated with each relationship and includes a user-installable web interface where users can (1) visualize the resulting network and (2) review the expression levels of associated genes in the original data, which helps biologists using the PathCORE software design experiments to test the relationships that were identified.
Conclusions: PathCORE is a hypothesis generation tool that identifies co-occurring pathways from the results of unsupervised analysis of the growing body of gene expression data. Software that steps beyond within-pathway relationships to between-pathway relationships can reveal levels of organization that have been less frequently considered.
Keywords: gene expression; unsupervised feature construction; crosstalk; unsupervised, pathway interactions

Background

The number of publicly available genome-wide datasets is growing rapidly [1]. High-throughput sequencing technologies that measure gene expression quickly with high accuracy and low cost continue to enable this growth [2]. Expanding public data repositories have laid the foundation for computational methods that consider entire compendia of gene expression data to extract biological patterns [3]. These patterns may be difficult to detect in measurements from a single experiment. Unsupervised approaches, which identify important signals in the data without relying on prior knowledge, may discover new expression modules [4, 5].

Feature extraction methods are a class of unsupervised algorithms that can reveal unannotated
biological processes from genomic data [5]. Features can be constructed as representative "meta-genes":
each feature has a set of influential genes, and these genes suggest the biological or technical pattern
captured by the feature. However, these features are often designed to be independent or may be
considered in isolation [6, 7]. When examined in the context of knowledgebases such as the Kyoto
Encyclopedia of Genes and Genomes (KEGG) [8], most features are significantly enriched for more than
one pathway [5]; it follows then that such features can be described by a set of functionally related
pathways. We introduce the PathCORE (**path**way **co**-occurrence **re**lationships) software, an approach for
connecting features learned from the data to known biological gene sets, e.g. pathways from KEGG or
other databases.
66
PathCORE offers a data-driven approach for predicting and visualizing global pathway-pathway
interactions. Interactions are drawn based on the sets of pathways, annotated in a resource of gene sets,
occurring within constructed features. To avoid simply discovering relationships between gene sets that
share many genes, PathCORE incorporates an optional pre-processing step that corrects for a situation
Donato et al. refer to as pathway crosstalk [9]. Donato et al. recognized that pathways with shared genes
were often discovered together due to overlapping genes in gene sets. We implement Donato et al.'s
maximum impact estimation in a Python package separate from, but used in, PathCORE (PyPI: crosstalk-
correction). With this correction, the PathCORE software allows a user to examine how pathways
influence each other in a biological system based on how genes are expressed as opposed to which
genes are shared.
77
We demonstrate PathCORE by applying the software to both a microbial and a cancer
expression dataset. Briefly, for the microbial analysis we created a network of KEGG pathways from
recently described ensemble Analysis using Denoising Autoencoders for Gene Expression (eADAGE)
models trained on a compendium of *Pseudomonas aeruginosa* (*P. aeruginosa*) gene expression data [5].
We provide a live demo of the PathCORE web application for the *P. aeruginosa* KEGG network at
pathcore-demo.herokuapp.com/PAO1. PathCORE can be used with other feature construction
approaches as well. For example, we perform PathCORE analysis of the same *P. aeruginosa*
compendium using non-negative matrix factorization (NMF) on expression datasets [10, 11]. We also
demonstrate PathCORE's use in large cancer genomics data by creating a Pathway Interaction Database
(PID) [12] pathway-pathway network of NMF features extracted from a The Cancer Genome Atlas
(TCGA) pan-cancer dataset of 33 different tumor types [13]. In these applications, PathCORE
successfully discerns biologically important pathway-pathway interactions from the constructed features.
90
Related work
Most published approaches that capture pathway-pathway interactions from gene expression
experiments were designed for disease-specific, case-control studies [14, 15]. Pham et al. developed
Latent Pathway Identification Analysis to find pathways that exert latent influences on transcriptionally
altered genes [16]. Under this approach, the transcriptional response profiles for a binary condition
(disease/normal), in conjunction with the pathway specified in the KEGG and functions in Gene Ontology
(GO), are used to construct a pathway-pathway network where key pathways are identified by their
network centrality scores [16, 17]. Similarly, Pan et al. measured the betweenness centrality of pathways
in disease-specific genetic interaction and coexpression networks to identify those most likely to be
associated with bladder cancer risk [18]. These methods captured pathway relationships associated with
a particular disease state. Our approach diverges from such studies in its intent: PathCORE finds
pathway relationships within a biological system that are discernable in features constructed from diverse
transcriptomic data--not necessarily specific to any one condition or disease.
104
Fewer publications to date have focused on the construction of a general pathway-pathway
interaction network. Those that did determined the absence or presence of a pathway-pathway interaction
based on shared genes between gene sets, protein-protein interactions or other curated knowledgebases
[19-22]. A function-based method of constructing a global network, detailed by Li et al., relied on publicly
available protein interaction data to determine pathway-pathway interactions [21]. Two pathways were
connected in the network if the number of protein interactions between the pair was significant with
respect to the computed background distribution. Networks of this kind rely on databases of interactions,

though they can be subsequently used for pathway-centric analyses of transcriptomic data [21, 23]. Glass
and Girvan described another network structure that relates functional terms in GO based on shared
gene annotations [24]. In contrast with this approach, PathCORE specifically removes gene overlap in
pathway definitions before they are used to build a network. Our software reports pathway-pathway
connections from global gene expression patterns, as opposed to protein-protein interactions, while
controlling for the fact that some pathways share genes.
118

119
Figure 1 The approach implemented in PathCORE to construct a pathway co-occurrence network from
an expression compendium.
(**a**) A user-selected feature extraction method is applied to expression data. Such methods assign each
gene a weight, according to some distribution, that represents the gene's contribution to the feature. The
set of genes that are considered highly representative of a feature's function is referred to as a feature's
gene signature. The gene signature is user-defined and should be based on the weight distribution
produced by the unsupervised method of choice. In the event that the weight distribution contains both
positive and negative values, a user can specify criteria for both a positive and negative gene signature. A
test of pathway enrichment is applied to identify corresponding sets of pathways from the gene
signature(s) in a feature. We consider pathways significantly overrepresented in the same feature to *co-*
occur. Pairwise co-occurrence relationships are used to build a network. Each edge is weighted by the
number of features containing both pathways.
(**b**) *N* permuted networks are generated to assess the statistical significance of a co-occurrence relation
in the network. Two invariants are maintained during a permutation: (1) pathway side-specificity (positive
and negative, when applicable) and (2) the number of distinct pathways in a feature's (side-specific) gene
signature.
(**c**) For each edge observed in the co-occurrence network, we compare its weight against the weight
distribution generated from *N* (default: 10,000) permutations of the network. Edges with a q-value below
alpha (default: 0.05) are kept in the final co-occurrence network.
139

Implementation
PathCORE identifies functional links between known pathways from the output of feature
construction methods applied to gene expression data. The result is a network of pathway co-occurrence
relationships that represents the grouping of biological processes or pathways within those features. We
correct for gene overlap in the pathway annotations to avoid identifying co-occurrence relationships
driven by shared genes. Additionally, PathCORE implements a permutation test for evaluating and
removing edges—pathway relationships—in the resulting network that cannot be distinguished from a null
model of random associations. Our software is written in Python and pip-installable (PyPI package name:
pathcore). Each of the functions that we describe here can be used independently; however, we expect

most users to employ the complete approach for interpreting pathways shared in extracted features (Fig.
4).
151
152   <u>Data organization</u>
153   PathCORE requires the following inputs:
154   (1) A **weight matrix** that connects each gene to each feature. We expect that this will be generated
155   by applying a feature construction algorithm to a compendium of gene expression data. In
156   principal component analysis (PCA), this is the loadings matrix [25]; in independent component
157   analysis (ICA), the unmixing matrix [26]; in ADAGE or eADAGE it is termed the weight matrix [5,
158   7]; in NMF it is the matrix W, where the NMF approximation of the input dataset A is $A \sim WH$ [10].
159   The primary requirements are that features must contain the full set of genes in the compendium
160   and genes must have been assigned weights that quantify their contribution to a given feature.
161   Accordingly, a weight matrix will have the dimensions n x k, where n is the number of genes in
162   the compendium and k is the number of features constructed.
163   (2) A **gene signature definition**. To construct a pathway co-occurrence network, the weight matrix
164   must be processed into gene signatures by applying a threshold to weights. Subsequent pathway
165   overrepresentation will be determined by the set(s) of genes within these signatures. These are
166   often the weights at the extremes of the distribution. How gene weights are distributed will
167   depend on the user's selected feature construction algorithm; because of this, a user must
168   specify the criterion for including a gene in a gene signature. PathCORE permits rules for a single
169   gene signature or both a positive and a negative gene signature. The use of 2 signatures may be
170   appropriate when the feature construction algorithm produces positive and negative weights, the
171   extremes of which both characterize a feature (e.g. PCA, ICA, ADAGE or eADAGE).
172   (3) A list of **pathway definitions**, where each pathway contains a set of genes (e.g. KEGG
173   pathways, PID pathways, GO biological processes).
174
175   <u>Weight matrix construction and signature definition</u>
176   In practice, users can obtain a weight matrix from many different methods. For the purposes of
177   this paper, we demonstrate generality by constructing weight matrices via eADAGE and NMF.
178
179   *eADAGE*
180   eADAGE is an unsupervised feature construction algorithm developed by Tan et al. [5] that uses
an ensemble of neural networks (an ensemble of ADAGE models) to capture biological signatures
embedded in the expression compendium. By initializing eADAGE with different random seeds, Tan et al.
produced 10 eADAGE models that each extracted k=300 features from the compendium of genome-scale
P. aeruginosa data. Because PathCORE supports the aggregation of co-occurrence networks created
from different models on the same input data, we use all 10 of these models in the PathCORE analysis of
eADAGE models (doi:10.5281/zenodo.583172).
187
188   Tan et al. refers to the features constructed by eADAGE as nodes. They are represented as a
weight matrix of size n x k, where n genes in the compendium are assigned positive or negative gene
weights, according to a standard normal distribution, for each feature k. Tan et al. determined that the
gene sets contributing the highest positive or highest negative weights (+/- 2.5 standard deviations) to a
feature described gene expression patterns across the compendium, and thus referred to the gene sets
as signatures. Because a feature's positive and negative gene signatures did not necessarily correspond
to the same biological process or function, Tan et al. analyzed each of these sets separately [5]. Tan et
al.'s gene signature rules are specified as an input to the PathCORE analysis as well.
196
NMF
We also constructed NMF models for the *P. aeruginosa* dataset and the TCGA pan-cancer dataset.
Given an NMF approximation of $A \sim WH$ *[10]*, where A is the input expression dataset of size n x s (n
genes by s samples), NMF aims to find the optimal reconstruction of A by WH such that W clusters on
samples (size n x k) and H clusters on genes (size k x s). We set k to the desired number of features,
k=300, and use W as the input weight matrix for the PathCORE software. We found that the gene weight
distribution of an NMF feature is right-skewed and (as the name suggests) non-negative (Fig. S1). In this

case, we defined the gene signature to be the set of genes with weights 2.0 standard deviations above
the mean weight of each feature.
Construction of a pathway co-occurrence network
We employ a Fisher's exact test to determine the pathways significantly associated with each
gene signature. When considering significance of a particular pathway, the two categories of gene
classification are as follows: (1) presence or absence of the gene in the gene signature and (2) presence
or absence of the gene in the pathway definition. We specify a contingency table for each pathway and
calculate the p-value. After false discovery rate (FDR) correction, pathways with a q-value of less than
alpha (default: 0.05) are considered significantly enriched. Two pathways co-occur, or share an edge in
the pathway co-occurrence network, if they are both overrepresented in a gene signature. The number of
times such a pathway pair is present over all features corresponds to its edge weight in the pathway-
pathway network (Fig. 1a).
216
Permutation test
The network that results from the preceding method is densely connected, and many edges may
be spurious. To remove correlations that cannot be distinguished from random pathway associations, we
define a statistical test that determines whether a pathway-pathway relationship appearing x times in a k--
feature model is unexpected under the null hypothesis—the null hypothesis being that the relationship
does not appear more often than it would in a random network. We create N weighted null networks by
permuting overrepresented pathways across the model's features while preserving the number of
pathways for which each feature is enriched. In the case where we have positive and negative gene
signatures, overrepresentation can be positive or negative. Because certain pathways may display bias
toward one side—for example, a pathway may be overrepresented more often in features' positive gene
signatures—we perform the permutation separately for each side. The N random networks produce the
background weight distribution for every observed edge; significance can then be assessed by comparing
the true (observed) edge weight against the null (Fig. 1b). Pathway-pathway relationships with a q-value
above alpha (default: 0.05) are considered insignificant by this statistical test and are removed from the
network of co-occurring pathways (Fig. 1c).
232
233   Because we can derive the expected weight of every edge from the N random networks, we can
234 divide the observed edge weights by their respective expected weights (divide by 1 if the edge is not
235 present in any of the N permutations). Edges in the final network are weighted by their odds ratios.
236
Gene overlap correction
Pathways can co-occur because of shared genes (Fig. 2a, b, d). Though some approaches use
the overlap of genes to identify connected pathways, we sought to capture pairs of pathways that
persisted even when this overlap was removed. The phenomenon of observing enrichment of multiple
pathways due to gene overlap has been previously termed as "crosstalk," and Donato et al. have
developed a method to correct for it [9]. Due to confusion around the term, we refer to this as *overlapping
genes* in this work, except where specifically referencing Donato et al. Their approach, called maximum
impact estimation, begins with a membership matrix indicating the original assignment of multiple genes
to multiple pathways. It uses expectation maximization to estimate the pathway in which a gene
contributes its greatest predicted impact (its maximum impact) and assigns the gene only to this pathway
[9]. This provides a set of new pathway definitions that no longer share genes (Fig. 2c, e).
248
We provide an implementation of Donato et al.'s maximum impact estimation as a Python
package separate from PathCORE so that it is available for any pathway analyses (PyPI package name:
crosstalk-correction). The procedure is written using NumPy functions and data structures, which allows
for efficient implementation of array and matrix operations in Python [28]. In PathCORE, overlapping
genes are addressed before pathway overrepresentation analysis so that the resulting pathway co-
occurrence network identifies interactions that are not driven by gene overlap. We incorporate this
correction into the PathCORE workflow by default; however, users can choose to disable it as well.

Figure 2 Correcting for gene overlap results in a sparser pathway co-occurrence network.
(**a**) The KEGG pathway annotations for the sulfonate transport system are a subset of those for sulfur metabolism. 12 genes annotated to the sulfonate transport system are also annotated to sulfur metabolism. (**b**) Without applying the overlap correction procedure, 25 of the genes in the positive and negative gene signatures of the eADAGE feature "Node 11" are annotated to sulfur metabolism--of those, 8 genes are annotated to the sulfonate transport system as well. (**c**) All 8 of the overlapping genes are mapped to the sulfur metabolism pathway after overlap correction.
(**d**) A co-occurrence network built without applying the overlap correction procedure will report co-occurrence between the sulfonate transport system and sulfur metabolism, whereas (**e**) no such relation is identified after overlap correction.

PathCORE network visualization and support for experimental follow-up

As an optional step, a Flask application can be set up for each PathCORE network. Metadata gathered from the analysis are saved to TSV files, and we use a script to populate collections in a MongoDB database with this information. The co-occurrence network is rendered using the D3.js force-directed graph layout [29]. Users can select a pathway-pathway relationship in the network to view a new page containing details about the genes annotated to one or both pathways (Fig. 3a).

We created a web interface for deeper examination of interactions present in the pathway co-occurrence network. When presented with a visualization of the PathCORE network, our collaborators suggested that additional support for determining potential gene targets and experimental conditions would help them design experiments to validate novel relationships. The details we included in an edge-specific page address their suggestion by (1) highlighting up to twenty genes--annotated to either of the two pathways in the edge--contained in features that also contain this edge, after controlling for the total number of features that contain each gene, and (2) displaying the expression levels of these genes in each of the fifteen samples where they were most and least expressed. The quantity of information (twenty genes, thirty samples total) we choose to include in an edge page is intentionally limited so that users can review it in a reasonable amount of time.

To implement the functionality in (1), we computed an odds ratio for every gene annotated to one or both pathways in the edge. The odds ratio measures how often we observe a feature enriched for both the given gene and the edge of interest relative to how often we would expect to see this occurrence. We

289 calculate the proportion of observed cases and divide by the expected proportion--equivalent to the
290 frequency of the edge appearing in the model's features.
291
292   Let K be the number of features from which the PathCORE network was built. K_G is the number of
293 features that contain gene G (i.e. G is in feature K's gene signature), K_E the number of features that
294 contain edge E (i.e. the two pathways connected by E are overrepresented in feature K), and $K_{G \& E}$ the
295 number of features that contain both gene G and edge E. The odds ratio is computed as follows:

$$Observed = K_{G \& E} / K_G$$
$$Expected = K_E / K$$
$$Odds\ ratio = Observed / Expected$$

299 An odds ratio above 1 suggests that the gene is more likely to appear in features enriched for this pair of
300 pathways: we rank the genes by their odds ratio to highlight genes most observed with the co-occurrence
301 relationship.
302
303   The information specified in (2) requires an "expression score" for every sample. A sample
304 expression score is calculated using the twenty genes we selected in goal (1): it is the average of the
305 normalized gene expression values weighted by the normalized gene odds ratio. Selection of the most
306 and least expressed samples is based on these scores. We use two heatmaps to show the twenty genes'
307 expression values in each of the fifteen most and least expressed samples (Fig. 3b).
308
309   For each sample in an edge page, a user can examine how the expression values of the edge's
310 twenty genes in that sample compare to those recorded for all other samples in the dataset that are from
311 the same experiment (Fig. 3c). Genes that display distinct expression patterns under a specific setting
312 may be good candidates for follow-up studies.
313

314
Figure 3 A web application used to analyze pathway-pathway interactions in the eADAGE-based, *P.*
aeruginosa KEGG network.
(**a**) A user clicks on an edge (a pathway-pathway interaction) in the network visualization and (**b**) is
directed to a page that displays expression data from the original transcriptomic dataset specific to the
selected edge (goo.gl/Hs5A3e). The expression data is visualized as two heatmaps that indicate the
fifteen most and fifteen least expressed samples corresponding to the edge. To select the "most" and
"least" expressed samples, we assign each sample a summary "expression score." The expression score

is based on the expression values of the genes (limited to the top twenty genes with an odds ratio above
1) annotated to one or both of the pathways. Here, we show the heatmap of least expressed samples
specific to the [Phosphate transport - Type II general secretion] relationship. (**c**) Clicking on a square in
the heatmap directs a user to an experiment page (goo.gl/KYNhwB) based on the sample corresponding
to that square. A user can use the experiment page to identify whether the expression values of genes
specific to an edge and a selected sample differ from those recorded in other samples of the experiment.
In this experiment page, the first three samples (labeled in black) are *P. aeruginosa* "baseline" replicates
grown for 72 h in drop-flow biofilm reactors. The following three samples (labeled in blue) are *P.
aeruginosa* grown for an additional 12 h (84 h total). Labels in blue indicate that the three 84 h replicates
are in the heatmap of least expressed samples displayed on the [Phosphate transport – Type II general
secretion] edge page.
333
Results
Interpreting features extracted by unsupervised clustering algorithms with PathCORE.
Networks modeling the relationships between curated processes in a biological system offer a
means for developing new hypotheses about which pathways influence each other and when. PathCORE
creates a network of globally co-occurring pathways based on features observed in a compendium of
gene expression data. Biological patterns in the data are extracted by a feature construction algorithm
such as PCA [30], ICA [6], NMF [10], GWCoGAPS [4], or eADAGE [5]. These algorithms capture sources
of variability in the data that induce coordinated changes in gene expression as features. The genes that
contribute the most to these features covary. This provides a data-driven categorization of the biological
system that can then be analyzed at the pathway-level by identifying annotated pathways
overrepresented in each feature.
345

346
Figure 4 The PathCORE software analysis workflow.
(**a**) A user applies a feature construction algorithm to a transcriptomic dataset of genes-by-samples. The
features constructed must preserve the genes in the dataset and assign weights to each of these genes
according to some distribution. (**b**) Inputs required to run the complete PathCORE analysis workflow. The
features constructed are stored in a weight matrix and the user-defined gene signature rules--up to 2 for
both a positive and negative gene signature--should be based on the algorithm's specified feature weight
distribution. A list of pathway definitions will be used to interpret the features constructed and build a
pathway-pathway co-occurrence network. (**c**) Methods in the PathCORE analysis workflow (capitalized
and in purple), can be employed independently of each other so long as the necessary input(s) are
provided.
357

The methods we implement in PathCORE can be used independently of each other (Fig. 4).
Here, we present analyses that can be produced by applying the full PathCORE pipeline to models
created from a transcriptomic compendium by an unsupervised feature construction algorithm. Input
pathway definitions are "overlap-corrected" (correcting for gene overlap between definitions) for each
feature before enrichment analysis. An overlap-corrected, weighted pathway co-occurrence network is
built by connecting the pairs of pathways that are overrepresented in features of the model. Finally, we
remove edges that cannot be distinguished from a null model of random associations based on the
results of a permutation test.
366
PathCORE also offers support for users interested in experimentally verifying a pathway-pathway
relationship (Fig. 3). We provide the code for setting up a web application where the network can be
visualized and its edges analyzed using the original input information. A pathway-pathway edge page
contains 2 heatmaps that display the samples in the compendium where the underlying genes are most
and least expressed. When available, information about each sample can be included on the page so that
users can refer to the conditions in which the expression patterns occurred (Fig. 3c).
373

374
Figure 5 eADAGE features constructed from publicly available *P. aeruginosa* expression data describe
known KEGG pathway relationships.
(**a**) The glycolysis/gluconeogenesis, pentose phosphate, and Entner-Doudoroff pathways share common
functions related to glucose catabolism .
(**b**) Organophosphate and inorganic phosphate transport- and metabolism-related processes frequently
co-occur with bacterial secretion systems; in particular, we highlight the pairwise relationships between
type II secretion and the phosphate-related processes.
(**c**) Pathways involved in the catabolism of sulfur-containing molecules, taurine (NitT/TauT family
transport) and methionine (D-Methionine transport), and the general sulfur metabolism process are
functionally linked.
(**d**) We observe pairwise relationships between zinc transport, iron transport, and the MacAB-TolC
transporter.
387
<u>PathCORE identifies interactions between KEGG pathways in *P. aeruginosa* using features extracted</u>
<u>from publicly available *P. aeruginosa* gene expression experiments</u>
We used PathCORE to create a network of co-occurring pathways out of the expression
signatures extracted from a *P. aeruginosa* compendium. For every feature, overlap correction was
applied to the *P. aeruginosa* KEGG pathway annotations and overlap-corrected annotations were used in
the overrepresentation analysis. PathCORE aggregates multiple networks by taking the union of the

edges across all networks and summing the weights of common pathway-pathway connections. We do
this to emphasize the pathway-pathway co-occurrence relationships that are more stable [31] —that is,
the relationships that appear across multiple models. Finally, we removed edges in the aggregate
network that were not significant after FDR correction when compared to the background distributions
generated from 10,000 permutations of the network. We applied PathCORE to features built by both NMF
(Fig. S2) and eADAGE (discussed below). The pathway-pathway network generated by NMF [27] is
smaller than that generated by eADAGE; that is, there are fewer pathway-pathway connections in the
network. It is possible that this is due to a difference in the stability of the results from the two algorithms
or the comprehensiveness of the features extracted by each approach. eADAGE includes an ensemble
step that improves model consistency. The eADAGE authors also observed that models constructed by
this ensemble procedure also more comprehensively captured pathways than non-ensemble models [5].
The PathCORE analysis of a 300 feature NMF decomposition of the *P. aeruginosa* compendium
produced a KEGG network that is similar in size to the PID network (Fig. 6, S2).

The eADAGE co-occurrence network identifies a number of pathway-pathway interactions that
have been previously characterized (Fig. 5). This suggests that PathCORE can capture functional links
between biological pathways. Three glucose catabolism processes co-occur in the network: glycolysis,
pentose phosphate, and the Entner-Doudoroff pathway (Fig. 5a). We also found a cluster relating
organophosphate and inorganic phosphate transport- and metabolism-related processes (Fig. 5b).
Notably, phosphate uptake and acquisition genes are directly connected to the *hxc* genes that encode a
type II secretion system. This Hxc secretion system is responsible for the secretion of alkaline
phosphatases, which are phosphate scavenging enzymes [32, 33] and the phosphate binding DING
protein [34]. Furthermore, alkaline phosphatases, DING and the *hxc* genes are regulated by the
transcription factor PhoB which is most active in response to phosphate limitation. As shown in Fig. 5c,
we also identified linkages between two pathways involved in the catabolism of sulfur-containing
molecules, taurine and methionine, and the general sulfur metabolism process. Other connections
between pathways involved in the transport of iron (ferrienterobactin binding) [35] and zinc (the *znu*
uptake system [36]) were identified (Fig. 5d). Interestingly, genes identified in the edge between the zinc
transport and MacAB-TolC pathways include the *pvd* genes involved in pyoverdine biosynthesis and
regulation, a putative periplasmic metal binding protein, as well as other components of an ABC
transporter (genes PA2407, PA2408, and PA2409 at goo.gl/bfqOk8) [37].

We used the PathCORE web application for the eADAGE KEGG *P. aeruginosa* network
(pathcore-demo.herokuapp.com/PAO1) to analyze the connection between the phosphate transport
system and a type II general secretion system pathway (Fig. 3a, b; edge page at goo.gl/Hs5A3e). The
sixteen genes reported on the edge page all have odds ratios above 34; these genes are at least 34
times more likely to appear in the gene signatures in which both of these pathways are overrepresented.
Such genes may help to reveal the biological basis of the co-occurrence relationship. In this case, the
results suggest that there may be some overlap between machinery for transporting phosphate into the
cell and secreting substances out of the cell via type II secretion. Alternatively, the two processes may be
mechanistically separate but coregulated such that phosphate scavenging molecules may be secreted by
type II secretion coincidentally to aid in phosphate acquisition. The heatmap of the fifteen least expressed
samples shows that the *pstB* and *pstS* genes, annotated to the phosphate transport system, are
consistently expressed higher relative to the other genes in the edge for these samples. The *pstS, pstC,*
pstA, and *pstB* genes are proximal of the phosphate-specific transport (Pst) operon that encodes a high-
affinity orthophosphate transport system [38]. Future studies will examine whether deletion of the Pst
phosphate system impairs secretion by the type II secretion system.

To assess whether the relationships identified in the pathway analysis paralleled gene expression
patterns in the context of a published experiment, we looked across experiments to determine if the
genes contained in the edge were co-regulated. As an example of the types of relationships that we
observed, we present a single experiment, E-GEOD-22164 from Folsom et al. (Fig. 3c; experiment page
at goo.gl/KYNhwB), that contains data from two sample types with three replicates each [39]. One set of
samples, referred to as the baseline, is of *P. aeruginosa* grown for 72 h in drop-flow biofilm reactors. The
other is of *P. aeruginosa* grown for an additional 12 h (84 h total). We found that three of the samples with
the lowest expression of the genes within the shared edge were samples from the later timepoint. All six

of these samples (three replicates each) are displayed in the experiment page. The sixteen edge genes
showed differential expression between the 72 h and 84 h timepoints. Particularly in the case of *pstA* and
pstC, the baseline replicates had normalized expression values near the center of the range from the
compendium whereas the 84 h samples had expression levels at the low end of the range. This suggests
that genes involved in both phosphate transport and type II secretion are less expressed at 84 h
compared to 72 h. Future studies will determine if this is due to a physiological change in the biofilm cells
at the late time point such that phosphate demands were lower or different sources of phosphate become
available.
458

459
Figure 6 PID pathway-pathway interactions discovered in NMF features constructed from the TCGA pan-
cancer gene expression dataset.
(**a**) Pathways in this module are responsible for cell cycle progression.
(**b**) Wnt signaling interactions with nuclear Beta-catenin signaling, FGF signaling, and BMP signaling
have all been linked to cancer progression.
(**c**) Here, we observe functional links between pathways responsible for angiogenesis and those
responsible for cell proliferation.
(**d**) The VEGF-VEGFR pathway interacts with the S1P3 pathway through Beta3 integrins.
(**e**) This module contains many interactions related to immune system processes. The interaction cycle
formed by T-Cell Receptor (TCR) signaling in naïve CD4+ T cells and IL-12/IL-4 mediated signaling
events, outlined in yellow, is one well-known example. The cycle in blue is formed by the ATF2, NFAT,
and AP1 pathways; pairwise co-occurrence of these three transcription factor networks may suggest that
dysregulation of any one of these pathways can trigger variable oncogenic processes in the immune
system.
474
PathCORE identifies interactions among PID pathways from the TCGA pan-cancer gene expression
dataset.
PathCORE is not specific to a certain dataset or organism. We also constructed a 300-feature
NMF model of TCGA pan-cancer gene expression, which is comprised of 33 different cancer-types from
various organ sites, and applied the PathCORE software to those features. We chose NMF because it
has been used in previous studies to identify biologically relevant patterns in transcriptomic data [10] and

by many studies to derive molecular subtypes [40-42]. The 300 NMF features were analyzed using
overlap-corrected PID pathways, a collection of 196 human cell signaling pathways with a particular focus
on processes relevant to cancer [12].
484
485 We found that PathCORE detects modules of co-occurring pathways consistent with our current
486 understanding of cancer-related interactions (Fig. 6). Importantly, because the connections were
487 constructed from many different cancer-types, these modules may represent pathway-pathway
488 interactions present in a large proportion of all tumors and may be good candidates for targeted
489 treatments.
490
For example, a module composed of a FoxM1 transcription factor network, an E2F transcription
factor network, Aurora B kinase signaling, ATR signaling, PLK1 signaling, and members of the Fanconi
anemia DNA damage response pathway are densely connected (Fig 6a). When two pathways share an
edge in the co-occurrence network, they are overrepresented together in one or more features. The
connections in this module recapitulate well known cancer hallmarks including cellular proliferation
pathways and markers of genome instability, such as the activation of DNA damage response pathways
[43]. We found that several pairwise pathway co-occurrences correspond with previously reported
pathway-pathway interactions [44-46]. We also observed a hub of pathways interacting with Wnt signaling
(Fig. 6b). In our network, pathways that co-occur with Wnt signaling include the regulation of nuclear
Beta-catenin signaling, FGF signaling, and BMP signaling. The Wnt and BMP pathways are functionally
integrated in biological processes that contribute to cancer progression [47]. Additionally, Wnt/Beta-
catenin signaling is a well-studied regulatory system, and the effects of mutations in Wnt pathway
components on this system have been linked to tumorigenesis [48]. Wnt/Beta-catenin and FGF together
influence the directional migration of cancer cell clusters [49].
505
Two modules in the network relate to angiogenesis, or the formation of new blood vessels (Fig.
6c, d). Tumors transmit signals that stimulate angiogenesis because a blood supply provides the
necessary oxygen and nutrients for their growth and proliferation. One module relates angiogenesis
factors to cell proliferation. This module connects the following pathways: PDGFR-beta signaling, FAK-
mediated signaling events, VEGFR1 and VEGFR2-mediated signaling events, nuclear SMAD2/3
signaling regulation, and RB1 regulation (Fig. 6c). These functional connections are known to be involved
in tumor proliferation [50-52]. The other module indicates a direct relationship by which the VEGF
pathway interacts with the S1P3 pathway through Beta3 integrins (Fig. 6d). *S1P3* is a known regulator of
angiogenesis [53], and has been demonstrated to be associated with treatment-resistant breast cancer
and poor survival [54]. Moreover, this interaction module has been observed to promote endothelial cell
migration in human umbilical veins [55]. Taken together, this independent module may suggest a distinct
angiogenesis process activated in more aggressive and metastatic tumors that is disrupted and regulated
by alternative mechanisms [56].
519
Finally, PathCORE revealed a large, densely connected module of immune related pathways
(Fig. 6e). We found that this module contains many interactions that align with immune system
processes. One such example is the well characterized interaction cycle formed by T-Cell Receptor
(TCR) signaling in naïve CD4+ T cells and IL-12/IL-4 mediated signaling events [57-59]. At the same
time, PathCORE predicts additional immune-related interactions. We observed a cycle between the three
transcription factor networks: ATF-2, AP-1, and CaN-regulated NFAT-dependent transcription. These
pathways can take on different, often opposing, functions depending on the tissue and subcellular
context. For example, ATF-2 can be an oncogene in one context (e.g. melanoma) and a tumor
suppressor in another (e.g. breast cancer) [60]. AP-1, comprised of Jun/Fos proteins, is associated with
both tumorigenesis and tumor suppression due to its roles in cell survival, proliferation, and cell death
[61]. Moreover, NFAT in complex with AP-1 regulates immune cell differentiation, but dysregulation of
NFAT signaling can lead to malignant growth and tumor metastasis [62]. The functional association
observed between the ATF-2, AP-1, and NFAT cycle together within the immunity module might suggest
that dysregulation within this cycle has profound consequences for immune cell processes and may
trigger variable oncogenic processes.
535
536 **Conclusions**

537   Unsupervised methods can identify previously undiscovered patterns in large collections of data.
538 PathCORE overlays curated knowledge after feature construction to help researchers interpret
539 constructed features in the context of existing knowledgebases. Specifically, PathCORE aims to clarify
540 how expert-annotated gene sets work together from a gene expression perspective.
541
Gene set analyses can be heavily confounded by shared genes. Some pathways may be
observed together because they depend on each other, while others may simply contain some of the
same genes. In PathCORE, pathway annotations undergo a procedure called maximum impact
estimation, described in a publication by Donato et al., that maps each gene in each feature to the one
pathway in which it has the greatest estimated impact [9]. We provide this overlap correction algorithm as
a Python package (PyPI package name: crosstalk-correction) available under the BSD 3-Clause license.
Though the algorithm had been described, no publicly available implementation existed.
549
PathCORE includes software for analysis and visualization and can be broadly applied to
constructed features. We demonstrate PathCORE in two different contexts, analyses of the bacterium *P.*
aeruginosa and human pan-cancer datasets, using two different feature construction methods (eADAGE
and NMF). We provide a demonstration application containing the results of the eADAGE *P. aeruginosa*
analysis for researchers to explore. For each edge, users can explore heatmaps displaying the
expression levels of predicted driver genes in the original samples. This provides support for assessing
computationally-derived relationships in experimental follow-ups.
557
Unsupervised analyses of genome-scale datasets that summarize key patterns in the data have
the potential to improve our understanding of how a biological system operates via complex interactions
between molecular processes. However, interpreting the features generated by unsupervised approaches
is still challenging. PathCORE is a component of a software ecosystem that connects the features
extracted from data to curated resources. The specific niche that PathCORE aims to fill is in revealing to
researchers which gene sets most are most closely related to each other in machine learning-based
models of gene expression, which genes play a role in this co-occurrence, and which conditions drive this
interaction will help researchers most effectively employ these algorithms.
566
Project name: PathCORE
Project home page: https://pathcore-demo.herokuapp.com
Archived version: https://github.com/greenelab/PathCORE-analysis/releases/tag/v1.0 (links to
download .zip and .tar.gz files are provided here)
Operating system: Platform independent
Programming language: Python
Other requirements: Python 3 or higher
License: BSD 3-clause
575
576 <u>**Declarations**</u>
577
578 **Ethics approval and consent to participate**
579 Not applicable
580
581 **Consent for publication**
582 Not applicable
583
584 **Availability of data and materials**
585 Files for each of the PathCORE networks described in the results are provided in Supplementary
586 material.
587
588 *Data sets*
589 *P. aeruginosa* eADAGE models: doi:10.5281/zenodo.583172
590 TCGA pan-cancer dataset: doi:10.5281/zenodo.56735
591
592 *Source code* (all links are from https://github.com/)

593 PathCORE analysis: (greenelab/PathCORE-analysis/tree/v1.0) This repository contains all the scripts to
594 reproduce the analyses described in this paper. The Python scripts here should be used as a starting
595 point for new PathCORE analyses. Instructions for setting up a web application for a user's specific
596 PathCORE analysis are provided in this repository's README.
597
598 Overlap correction: (kathyxchen/crosstalk-correction/tree/v1.0.4) Donato et. al's procedure for overlap
599 correction [9] is a pip-installable Python package 'crosstalk-correction' that is separate from, but listed as
600 a dependency in, PathCORE. It is implemented using NumPy [28].
601
602 PathCORE methods: (greenelab/PathCORE/tree/v1.0) The methods included in the PathCORE analysis
603 workflow (Fig. 4c) are provided as a pip-installable Python package 'pathcore'. It is implemented using
604 Pandas [62], SciPy (specifically, scipy.stats) [63], StatsModels [64], and the crosstalk-correction package.
605
PathCORE demo application: (kathyxchen/PathCORE-demo/tree/v1.0) The project home page, pathcore-
demo.herokuapp.com provides links to
(1) The **web application** for the eADAGE-based KEGG *P. aeruginosa* described in the first case
study.
(2) A **view** of the NMF-based PID pathway co-occurrence network described in the second case
study.
(3) A **quick view page** where users can temporarily load and visualize their own network file
(generated from the PathCORE analysis).
614
615 **Competing interests**
616 The authors declare that they have no competing interests
617

618 **Funding**
619 KMC was funded by an undergraduate research grant from the Penn Institute for Biomedical Informatics.
620 CSG was funded in part by a grant from the Gordon and Betty Moore Foundation (GBMF 4552). JT, GD,
621 DAH, and CSG were funded in part by a pilot grant from the Cystic Fibrosis Foundation (STANTO15R0).
622 DAH was funded in part by R01-AI091702.
623
624 **Authors' contributions**
625 KMC implemented the software, performed the analyses, and drafted the manuscript. JT and GPW
626 contributed computational reagents. KMC, DAH, and CSG designed the project. KMC, JT, GPW, GD,
627 DAH, and CSG interpreted the results. JT, GPW, GD, DAH, and CSG provided critical feedback and
628 revisions on the manuscript. JT and GPW reviewed source code.
629
630 **Acknowledgements**
631 The authors would also like to thank Daniel Himmelstein, Kurt Wheeler, and René Zelaya for helping to
632 review the source code.
633

References
1. Greene CS, Foster JA, Stanton BA, Hogan DA, Bromberg Y: **Computational Approaches to Study**
Microbes and Microbiomes. Pacific Symposium on Biocomputing 2016, **21**:557.
2. Tatlow PJ, Piccolo SR: **A cloud-based workflow to quantify transcript-expression levels in public**
cancer compendia. Scientific Reports 2016, **6**.
3. Kolesnikov N, Hastings E, Keays M, Melnichuk O, Tang YA, Williams E, Dylag M, Kurbatova N,
Brandizi M, Burdett T: **ArrayExpress update—simplifying data submissions.** Nucleic Acids Res 2014,
:gku1057.
4. Stein-O'Brien G, Carey J, Lee W, Considine M, Favorov A, Flam E, Guo T, Li L, Marchionni L,
Sherman T: **PatternMarkers and Genome-Wide CoGAPS Analysis in Parallel Sets (GWCoGAPS) for**
data-driven detection of novel biomarkers via whole transcriptome Non-negative matrix
factorization (NMF). bioRxiv 2016, :083717.
5. Tan J, Doing G, Lewis KA, Price CE, Chen KM, Cady KC, Perchuk B, Laub MT, Hogan DA, Greene
CS: **System-wide automatic extraction of functional signatures in *Pseudomonas aeruginosa* with**
eADAGE. Cell Systems 2016, In Press; doi:10.1101/078659.

6. Engreitz JM, Daigle BJ, Marshall JJ, Altman RB: **Independent component analysis: mining**
microarray data for fundamental human gene expression modules. J Biomed Inform 2010,
43(6):932-944.
7. Tan J, Hammond JH, Hogan DA, Greene CS: **Adage-based integration of publicly available**
Pseudomonas aeruginosa **gene expression data with denoising autoencoders illuminates**
microbe-host interactions. mSystems 2016, **1**(1):25.
8. Kanehisa M, Goto S: **KEGG: Kyoto Encyclopedia of Genes and Genomes.** Nucleic Acids Res 2000,
28(1):27-30.
9. Donato M, Xu Z, Tomoiaga A, Granneman JG, MacKenzie RG, Bao R, Than NG, Westfall PH, Romero
R, Draghici S: **Analysis and correction of crosstalk effects in pathway analysis.** Genome Res 2013,
23(11):1885-1893.
10. Brunet J, Tamayo P, Golub TR, Mesirov JP: **Metagenes and molecular pattern discovery using**
matrix factorization. Proceedings of the national academy of sciences 2004, **101**(12):4164-4169.
11. Kim PM, Tidor B: **Subsystem identification through dimensionality reduction of large-scale**
gene expression data. Genome Res 2003, **13**(7):1706-1718.
12. Schaefer CF, Anthony K, Krupa S, Buchoff J, Day M, Hannay T, Buetow KH: **PID: the Pathway**
Interaction Database. Nucleic Acids Res 2009, **37**(suppl 1):D679.
13. Weinstein JN, Collisson EA, Mills GB, Shaw KRM, Ozenberger BA, Ellrott K, Shmulevich I, Sander C,
Stuart JM, Cancer Genome Atlas Research Network: **The Cancer Genome Atlas pan-cancer analysis**
project. Nat Genet 2013, **45**(10):1113-1120.
14. Visakh R, Nazeer KA: **Identifying epigenetically dysregulated pathways from pathway–pathway**
interaction networks. Comput Biol Med 2016, **76**:160-167.
15. Yang J, Luo R, Yan Y, Chen Y: **Differential pathway network analysis used to identify key**
pathways associated with pediatric pneumonia. Microb Pathog 2016, **101**:50-55.
16. Pham L, Christadore L, Schaus S, Kolaczyk ED: **Network-based prediction for sources of**
transcriptional dysregulation using latent pathway identification analysis. Proceedings of the
National Academy of Sciences 2011, **108**(32):13347-13352.
17. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS,
Eppig JT: **Gene Ontology: tool for the unification of biology.** Nat Genet 2000, **25**(1):25-29.
18. Anonymous *Proceedings of the Ecal:* Citeseer; 2013.
19. Bell L, Chowdhary R, Liu JS, Niu X, Zhang J: **Integrated bio-entity network: a system for**
biological knowledge discovery. PloS one 2011, **6**(6):e21474.
20. Chowbina SR, Wu X, Zhang F, Li PM, Pandey R, Kasamsetty HN, Chen JY: **HPD: an online**
integrated human pathway database enabling systems biology studies. BMC Bioinformatics 2009,
10(11):S5.
21. Li Y, Agarwal P, Rajagopalan D: **A global pathway crosstalk network.** Bioinformatics 2008,
24(12):1442-1447.
22. Wu X, Chen JY: **Molecular interaction networks: topological and functional characterizations.**
Automation in Proteomics and Genomics: An Engineering Case-Based Approach 2009, **145**.
23. de Anda-Juregui G, Meja-Pedroza RA, Espinal-Enrquez J, Hernndez-Lemus E: **Crosstalk events in**
the estrogen signaling pathway may affect tamoxifen efficacy in breast cancer molecular
subtypes. Computational biology and chemistry 2015, **59**:42-54.
24. Glass K, Girvan M: **Finding new order in biological functions from the network structure of gene**
annotations. PLoS Comput Biol 2015, **11**(11):e1004565.
25. Abdi H, Williams LJ: **Principal component analysis.** Wiley interdisciplinary reviews: computational
statistics 2010, **2**(4):433-459.
26. Stone JV: *Independent component analysis:* Wiley Online Library; 2004.
27. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer P,
Weiss R, Dubourg V: **Scikit-learn: Machine learning in Python.** Journal of Machine Learning Research
2011, **12**(Oct):2825-2830.
28. Walt Svd, Colbert SC, Varoquaux G: **The NumPy array: a structure for efficient numerical**
computation. Computing in Science & Engineering 2011, **13**(2):22-30.
29. Bostock M: **D3. js.** Data Driven Documents 2012, **492**.
30. Yeung KY, Ruzzo WL: **Principal component analysis for clustering gene expression data.**
Bioinformatics 2001, **17**(9):763-774.
31. Yu B: **Stability.** Bernoulli 2013, **19**(4):1484-1500.

32. Liu X, Long D, You H, Yang D, Zhou S, Zhang S, Li M, He M, Xiong M, Wang X:
Phosphatidylcholine affects the secretion of the alkaline phosphatase PhoA in *Pseudomonas*
strains. Microbiol Res 2016, **192**:21-29.
33. Ball G, Durand R, Lazdunski A, Filloux A: **A novel type II secretion system in *Pseudomonas***
aeruginosa. Mol Microbiol 2002, **43**(2):475-485.
34. Ball G, Viarre V, Garvis S, Voulhoux R, Filloux A: **Type II-dependent secretion of a *Pseudomonas***
***aeruginosa* DING protein.** Res Microbiol 2012, **163**(6):457-469.
35. Stephens DL, Choe MD, Earhart CF: ***Escherichia coli* periplasmic protein FepB binds**
ferrienterobactin. Microbiology 1995, **141**(7):1647-1654.
36. Ellison ML, IIII JMF, Parrish W, Danell AS, Pesci EC: **The transcriptional regulator Np20 is the zinc**
uptake regulator in *Pseudomonas aeruginosa*. PloS one 2013, **8**(9):e75389.
37. Winsor GL, Lam DK, Fleming L, Lo R, Whiteside MD, Nancy YY, Hancock RE, Brinkman FS:
***Pseudomonas* Genome Database: improved comparative analysis and population genomics**
capability for *Pseudomonas* genomes. Nucleic Acids Res 2010, :gkq869.
38. Rico-Jimnez M, Reyes-Darias JA, Ortega I, Pea AID, Morel B, Krell T: **Two different mechanisms**
mediate chemotaxis to inorganic phosphate in *Pseudomonas aeruginosa*. Scientific Reports 2016,
6.
39. Folsom JP, Richards L, Pitts B, Roe F, Ehrlich GD, Parker A, Mazurie A, Stewart PS: **Physiology of**
***Pseudomonas aeruginosa* in biofilms as revealed by transcriptome analysis.** BMC microbiology
2010, **10**(1):294.
40. Bailey P, Chang DK, Nones K, Johns AL, Patch A, Gingras M, Miller DK, Christ AN, Bruxner TJ,
Quinn MC: **Genomic analyses identify molecular subtypes of pancreatic cancer.** Nature 2016, .
41. Cancer Genome Atlas Research Network: **Integrated genomic analyses of ovarian carcinoma.**
Nature 2011, **474**(7353):609-615.
42. Yuan Y, Van Allen EM, Omberg L, Wagle N, Amin-Mansour A, Sokolov A, Byers LA, Xu Y, Hess KR,
Diao L: **Assessing the clinical utility of cancer genomic and proteomic data across tumor types.**
Nat Biotechnol 2014, **32**(7):644-652.
43. Hanahan D, Weinberg RA: **Hallmarks of cancer: the next generation.** Cell 2011, **144**(5):646-674.
44. Moldovan G, D'Andrea AD: **How the fanconi anemia pathway guards the genome.** Annu Rev
Genet 2009, **43**:223-249.
45. Sadasivam S, DeCaprio JA: **The DREAM complex: master coordinator of cell cycle-dependent**
gene expression. Nature Reviews Cancer 2013, **13**(8):585-595.
46. Tomida J, Itaya A, Shigechi T, Unno J, Uchida E, Ikura M, Masuda Y, Matsuda S, Adachi J,
Kobayashi M: **A novel interplay between the Fanconi anemia core complex and ATR-ATRIP kinase**
during DNA cross-link repair. Nucleic Acids Res 2013, **41**(14):6930-6941.
47. Itasaki N, Hoppler S: **Crosstalk between Wnt and bone morphogenic protein signaling: a**
turbulent relationship. Developmental Dynamics 2010, **239**(1):16-33.
48. MacDonald BT, Tamai K, He X: **Wnt/β-catenin signaling: components, mechanisms, and**
diseases. Developmental cell 2009, **17**(1):9-26.
49. Aman A, Piotrowski T: **Wnt/β-catenin and Fgf signaling control collective cell migration by**
restricting chemokine receptor expression. Developmental cell 2008, **15**(5):749-761.
50. Petersen M, Pardali E, Van Der Horst G, Cheung H, Van Den Hoogen C, Van Der Pluijm G, Ten Dijke
P: **Smad2 and Smad3 have opposing roles in breast cancer bone metastasis by differentially**
affecting tumor angiogenesis. Oncogene 2010, **29**(9):1351-1361.
51. Yoon H, Dehart JP, Murphy JM, Lim SS: **Understanding the roles of FAK in cancer: inhibitors,**
genetic models, and new insights. Journal of Histochemistry & Cytochemistry 2015, **63**(2):114-128.
52. Chinnam M, Goodrich DW: **RB1, development, and cancer.** Curr Top Dev Biol 2011, **94**:129.
53. Takuwa Y, Du W, Qi X, Okamoto Y, Takuwa N, Yoshioka K: **Roles of sphingosine-1-phosphate**
signaling in angiogenesis. World journal of biological chemistry 2010, **1**(10):298-306.
54. Watson C, Long JS, Orange C, Tannahill CL, Mallon E, McGlynn LM, Pyne S, Pyne NJ, Edwards J:
High expression of sphingosine 1-phosphate receptors, S1P 1 and S1P 3, sphingosine kinase 1,
and extracellular signal-regulated kinase-1/2 is associated with development of tamoxifen
resistance in estrogen receptor-positive breast cancer patients. The American journal of pathology
2010, **177**(5):2205-2215.
55. Paik JH, Chae S, Lee M, Thangada S, Hla T: **Sphingosine 1-phosphate-induced endothelial cell**
migration requires the expression of EDG-1 and EDG-3 receptors and Rho-dependent activation of

αvβ3-and β1-containing integrins. J Biol Chem 2001, **276**(15):11830-11837.
56. Serini G, Valdembri D, Bussolino F: **Integrins and angiogenesis: a sticky business.** Exp Cell Res
2006, **312**(5):651-658.
57. Brogdon JL, Leitenberg D, Bottomly K: **The potency of TCR signaling differentially regulates**
NFATc/p activity and early IL-4 transcription in naive CD4 T cells. The Journal of Immunology 2002,
168(8):3825-3832.
58. Hsieh C, Macatonia SE, Tripp CS, Wolf SF, O'Garra A, Murphy KM: **Development of TH1 CD4 T**
Cells Through IL-12. Science 1993, **260**:547.
59. Vacaflores A, Chapman NM, Harty JT, Richer MJ, Houtman JC: **Exposure of Human CD4 T Cells to**
IL-12 Results in Enhanced TCR-Induced Cytokine Production, Altered TCR Signaling, and
Increased Oxidative Metabolism. PloS one 2016, **11**(6):e0157175.
60. Lau E, Ze'ev AR: **ATF2–at the crossroad of nuclear and cytosolic functions.** J Cell Sci 2012,
125(12):2815-2824.
61. Shaulian E, Karin M: **AP-1 as a regulator of cell life and death.** Nat Cell Biol 2002, **4**(5):E136.
62. Mller MR, Rao A: **NFAT, immunity and cancer: a transcription factor comes of age.** Nature
Reviews Immunology 2010, **10**(9):645-656.