Single-cell RNA-seq of dopaminergic neurons informs candidate gene selection for sporadic

Parkinson's disease

Paul W. Hook[1], Sarah A. McClymont[1], Gabrielle H. Cannon[1], William D. Law[1], A. Jennifer

Morton[2], Loyal A. Goff[1,3*], Andrew S. McCallion[1,4,5*]

[1]McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of

Medicine, Baltimore, Maryland, United States of America

[2]Department of Physiology Development and Neuroscience, University of Cambridge,

Cambridge, United Kingdom

[3]Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore,

Maryland, United States of America

[4]Department of Comparative and Molecular Pathobiology, Johns Hopkins University School of

Medicine, Baltimore, Maryland, United States of America

[5]Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland,

United States of America

*, To whom correspondence should be addressed: andy@jhmi.edu and loyalgoff@jhmi.edu

ABSTRACT

Parkinson's disease (PD) is caused by the collapse of *substantia nigra* (SN) dopaminergic (DA)

neurons of the midbrain (MB), while other DA populations remain relatively intact. Common

variation influencing susceptibility to sporadic PD has been primarily identified through genome

wide association studies (GWAS). However, like many other common genetic diseases, the

genes impacted by common PD-associated variation remain to be elucidated. Here, we used

single-cell RNA-seq to characterize DA neuron populations in the mouse brain at embryonic and

early postnatal timepoints. These data allow for the unbiased identification of DA neuron

subpopulations, including a novel postnatal neuroblast population and SN DA neurons.

Comparison of SN DA neurons with other DA neurons populations in the brain reveals a unique

transcriptional profile, novel marker genes, and specific gene regulatory networks. By integrating

these cell population specific data with published GWAS, we develop a scoring system for

prioritizing candidate genes in PD-associated loci. With this, we prioritize candidate genes in all

32 GWAS intervals implicated in sporadic PD risk, the first such systematically generated list.

From this we confirm that the prioritized candidate gene *CPLX1* disrupts the nigrostriatal

pathway when knocked out in mice. Ultimately, this systematic rationale leads to the

identification of biologically pertinent candidates and testable hypotheses for sporadic PD that

will inform a new era of PD genetic research.

The most commonly used genetic tool today for studying complex disease is the genome wide

association study (GWAS). As a strategy, GWAS was initially hailed for the insight it might

provide into the genetic architecture of common human disease risk. Indeed, the collective data

from GWAS since 2005 has revealed a trove of variants and genomic intervals associated with

an array of phenotypes[1]. The majority of variants identified in GWAS are located in non-coding

DNA[2] and are enriched for characteristics denoting regulatory DNA[2,3]. This regulatory variation

is expected to impact expression of a nearby gene, leading to disease susceptibility.

Traditionally, the gene closest to the lead SNP has been prioritized as the affected gene.

However, recent studies show that disease-associated variants can act on more distally located

genes, invalidating genes that were previously extensively studied[4,5]. The inability to

systematically connect common variation with the genes impacted limits our capacity to

elucidate potential therapeutic targets and can waste valuable research efforts.

Although GWAS is inherently agnostic to the context in which disease-risk variation acts, the

biological impact of common functional variation has been shown to be cell context

dependent[2,6]. Extending these observations, Pritchard and colleagues recently demonstrated that

although genes need only to be expressed in disease-relevant cell types to contribute to risk,

those expressed preferentially or exclusively therein contribute more per SNP[7]. Thus, accounting

for the cellular and gene regulatory network (GRN) contexts within which variation act may

better inform the identification of impacted genes. These principles have not yet been applied

systematically to many of the traits for which GWAS data exists. We have chosen Parkinson's

disease (PD) as a model complex disorder for which a significant body of GWAS data remains to

be explored biologically in a context dependent manner.

PD is the most common progressive neurodegenerative movement disorder. Incidence of PD

increases with age, affecting an estimated 1% worldwide beyond 70 years of age[8–10]. The genetic

underpinnings of non-familial or sporadic PD have been studied through the use of GWAS with

a recent meta-analysis highlighting 32 loci associated with sporadic PD susceptibility[11]. While a

small fraction of PD GWAS loci contain genes known to be mutated in familial PD (*SNCA* and

LRRK2)[12,13], most indicted intervals do not contain a known causal gene or genes. Although PD

ultimately affects multiple neuronal centers, preferential degeneration of DA neurons in the SN

leads to functional collapse of the nigrostriatal pathway and loss of fine motor control. The

preferential degeneration of SN DA neurons in relation to other mesencephalic DA neurons has

driven research interest in the genetic basis of selective SN vulnerability in PD. Consequently,

one can reasonably assert that a significant fraction of PD-associated variation likely mediates its

influence specifically within the SN.

In an effort to illuminate a biological context in which PD GWAS results could be better

interpreted, we undertook single-cell RNA-seq (scRNA-seq) analyses of multiple DA neuronal

populations in the brain, including ventral midbrain DA neurons. This analysis defined the

heterogeneity of DA populations over developmental time in the brain, revealing gene

expression profiles specific to discrete DA neuron subtypes. These data further facilitated the

definition of GRNs active in DA neuron populations including the SN. With these data, we

establish a framework to systematically prioritize candidate genes in all 32 PD GWAS loci and

begin exploring their pathological significance.

RESULTS

scRNA-seq characterization defines DA neuronal subpopulation heterogeneity

In order to characterize DA neuron molecular phenotypes, we undertook scRNA-seq on cells

isolated from distinct anatomical locations of the mouse brain over developmental time. We used

fluorescence activated cell sorting (FACS) to retrieve single DA neurons from the Tg(Th-

EGFP)DJ76Gsat BAC transgenic mouse line, which expresses eGFP under the control of the

tyrosine hydroxylase (*Th*) locus[14]. We microdissected both MB and FB from E15.5 mice,

extending our analyses to MB, FB, and OB in P7 mice (Figure 1a). E15.5 and P7 time points

were chosen based on their representation of stable MB DA populations, either after neuron birth

(E15.5) or between periods of programmed cell death (P7) (Figure 1a)[15].

Quality control and outlier analysis identify 396 high quality cell transcriptomes to be used in

our analyses. We initially sequenced RNA from 473 single cells to an average depth of ~8 x 10^5

50 bp paired-end fragments per cell. Using Monocle 2, we converted normalized expression

estimates into estimates of RNA copies per cell[16]. Cells were filtered based on the distributions

of total mass, total number of mRNAs, and total number of expressed genes per cell

(Supplementary Figure 1a, 1b, 1c; detailed in Methods). After QC, 410 out of 473 cells were

retained. Using principal component analysis (PCA) as part of the iterative analysis described

below, we identified and removed 14 outliers determined to be astrocytes, microglia, or

oligodendrocytes (Supplementary Figure 1e; Supplementary Table 1), leaving 396 cells (~79

cells/timepoint-region; Supplementary Figure 1d).

To confirm that our methods can discriminate between different populations of neurons, we first

explored differences between timepoints. Following a workflow similar to the recently described

"dpFeature" procedure[17], we identified genes with highly variable transcriptional profiles and

performed PCA. As anticipated, we observed that the greatest source of variation was between

developmental ages (Figure 1b). Genes associated with negative PC1 loadings (E15.5 cells) were

enriched for gene sets consistent with mitotically active neuronal, undifferentiated precursors

(Figure 1c). In contrast, genes associated with positive PC1 loadings (P7 cells) were enriched for

ontology terms associated with mature, post-mitotic neurons (Figure 1c). This initial analysis

establishes our capacity to discriminate among biological classes present in our data using PCA

as a foundation.

Further, we attempted to identify clusters of single cells between and within timepoints and

anatomical regions. In order to do this, we selected the PCs that described the most variance in

the data and used t-Stochastic Neighbor Embedding (t-SNE)[18] to further cluster cells in an

unsupervised manner (see Methods). Analysis of all cells revealed that the E15.5 cells from both

MB and FB cluster together (Figure 1d), supporting the notion that they are less differentiated.

By contrast, cells isolated at P7 mostly cluster by anatomical region, suggesting progressive

functional divergence with time (Figure 1d). We next applied this same scRNA-seq analysis

workflow (See Methods) in a recursive manner individually in all regions at both timepoints to

further explore heterogeneity. This revealed a total of 13 clusters (E15.5 FB.1-2, MB.1-2; P7

OB.1-3, FB.1-2, MB.1-4; Figure 1e), demonstrating the diversity of DA neuron subtypes and

providing a framework upon which to evaluate the biological context of genetic association

signals across closely-related cell types. Using known markers, we confirmed that all clusters

expressed high levels of pan-neuronal markers (*Snap25*, *Eno2*, and *Syt1*) (Supplementary Figure

2a). In contrast, we observed scant evidence of astrocyte (*Aldh1l1*, *Slc1a3*, *Aqp4*, and *Gfap*;

Supplementary Figure 2a) or oligodendrocyte markers (*Mag*, *Mog*, and *Mbp*; Supplementary

Figure 2a), thus confirming we successfully isolated our intended substrate, *Th+* neurons.

scRNA-seq revealed biologically and temporally discriminating transcriptional signatures

With subpopulations of DA neurons defined in our data, we set out to assign a biological identity

to each cluster. Among the four clusters identified at E15.5, two were represented in t-SNE space

as a single large group that included cells from both MB and FB (E15.MB.1, E15.FB.1), leaving

two smaller clusters that were comprised solely of MB or FB cells (Supplementary Figure 3a).

The latter MB cluster (E15.MB.2; Supplementary Figure 3a, Supplementary Figure 3b)

specifically expressed *Foxa1*, *Lmx1a*, *Pitx3*, and *Nr4a2* and thus likely represents a post-mitotic

DA neuron population[19] (Supplementary Table 2; Supplementary Table 3). Similarly, the

discrete E15.FB.2 cluster expressed markers of post-mitotic FB/hypothalamic neurons

(Supplementary Figure 3b), including *Six3, Six3os1*, *Sst,* and *Npy* {Supplementary Table 2;

Supplementary Table 3}. These embryonic data did not discriminate between cells populating

known domains of DA neurons, such as the SN.

By contrast, P7 cells mostly cluster by anatomical region and each region has defined subsets

(Figure 1d, 1e, 2a). Analysis of P7 FB revealed two distinct cell clusters (Figure 2b). Expression

of the neuropeptides *Gal* and *Ghrh* and the *Gsx1* transcription factor place P7.FB.1 cells in the

arcuate nucleus (Supplementary Table 2; Supplementary Table 3)[20–23]. The identity of P7.FB.2,

however, was less clear, although subsets of cells therein did express other arcuate nucleus

markers for $Th^+/Ghrh^-$ neuronal populations e.g. *Onecut2*, *Arx*, *Prlr*, *Slc6a3*, and *Sst*

(Supplementary Figure 3c; Supplementary Table 3)[23]. All three identified OB clusters (Figure

2c) express marker genes of OB DA neuronal development or survival (Supplementary Table 2,

Supplementary Table 3; Supplementary Figure 3d)[24]. It has previously been reported that *Dcx*

expression diminishes with neuronal maturation[25] and *Snap25* marks mature neurons[26]. We

observe that these OB clusters seem to reflect this continuum of maturation wherein expression

of *Dcx* diminishes and *Snap25* increases with progression from P7.OB1 to OB3 (Supplementary

Figure 3d). This pattern is mirrored by a concomitant increase in OB DA neuron fate

specification genes (Supplementary Figure 3d)[24,27]. In addition, we identified four P7 MB DA

subset clusters (Figure 2d). Marker gene analysis confirmed that three of the clusters correspond

to DA neurons from the VTA (*Otx2* and *Neurod6*; P7.MB.1)[28,29], the PAG (*Vip* and *Pnoc*;

P7.MB.3)[30,31], and the SN (*Sox6*, *Aldh1a7*, *Ndnf*, *Serpine2*, *Rbp4*, and *Fgf20*; P7.MB.4)[28,32–34]

(Supplementary Table 2; Supplementary Table 3). These data are consistent with recent scRNA-

seq studies of similar populations[33,35]. Through this marker gene analysis, we successfully

assigned a biological identity to 12/13 clusters.

The only cluster without a readily assigned identity was P7.MB.2. This population of P7 MB DA

neurons, P7.MB.2 (Figure 2d), is likely a progenitor-like population. Like the overlapping

E15.MB.1 and E15.FB.1 clusters (Supplementary Figure 3a), this cluster preferentially expresses

markers of neuronal precursors/differentiation/maturation {Supplementary Table 2,

Supplementary Table 3}. In addition to sharing markers with the progenitor-like E15.MB.1

cluster, P7.MB.2 exhibits gene expression consistent with embryonic mouse neuroblast

populations[33], cell division, and neuron development[36–40] (Supplementary Table 2,

Supplementary Table 3). Consistent with the hypothesis, this population displayed lower levels

of both *Th* and *Slc6a3,* markers of mature DA neurons, than the terminally differentiated and

phenotypically discrete P7 MB DA neuron populations of the VTA, SN and PAG (Figure 2e).

With this hypothesis in mind, we sought to ascertain the spatial distribution of P7.MB.2 DA

neurons through multiplex, single molecule fluorescence *in situ* hybridization (smFISH) for *Th*

(pan-P7 MB DA neurons), *Slc6a3* (P7.MB.1, P7.MB.3, P7.MB.4), and one of the neuroblast

marker genes identified through our analysis, either *Lhx9* or *Ldb2* (P7.MB.2) (Figure 2e). In each

experiment, we scanned the ventral midbrain for cells that were Th+/Slc6a3- and positive for the

third gene. *Th+/Slc6a3-/Lhx9+* cells were found scattered in the dorsal SN *pars compacta*

(SNpc) along with cells expressing *Lhx9* alone (Figure 2f, 2h). Expression of *Ldb2* was found to

have a similar pattern to *Lhx9*, with *Th+/Slc6a3-/Ldb2+* cells found in the dorsal SNpc (Figure

2f, 2h). Expression of *Lhx9* and *Ldb2* was low or non-existent in *Th+/Slc6a3+* cells in the SNpc

(Figure 2e, 2f). Importantly, cells expressing these markers express *Th* at lower levels than

Th+/Slc6a3+ neurons (Figure 2f, 2g), consistent with our scRNA-seq data (Figure 2e). Thus,

with the resolution of the spatial distribution of this novel neuroblast-like P7 MB DA population,

we assign biological identity to each defined brain DA subpopulation.

Novel SN-specific transcriptional profiles and GRNs highlight its association with PD

Overall our analyses above allowed us to successfully separate and identify 13 brain DA

neuronal populations present at E15.5 and P7, including SN DA neurons. Motivated by the

clinical relevance of SN DA neurons to PD, we set out to understand what makes them

transcriptionally distinct from the other MB DA neuron populations.

In order to look broadly at neuronal subtypes, we evaluated expression of canonical markers of

other neuronal subtypes in our *Th+* neuron subpopulations. Interestingly, we observed

inconsistent detection of *Th* and eGFP in some E15.5 clusters (Supplementary Figure 2b). This

likely reflects lower *Th* transcript abundance at this developmental state, but sufficient

expression of the eGFP reporter to permit FACS collection (Supplementary Figure 2c). The

expression of other DA markers, *Ddc* and *Slc18a2*, mirror *Th* expression, while *Slc6a3*

expression is more spatially and temporally restricted (Supplementary Figure 2b). The SN cluster

displays robust expression of all canonical DA markers (Supplementary Figure 2b). Multiple

studies have demonstrated that *Th+* neurons may also express markers characteristic of other

major neuronal subtypes[41–43]. We found that only the SN and PAG showed no expression of

either GABAergic (*Gad1*/*Gad2*/*Slc32a1*) or glutamatergic (*Slc17a6*) markers (Supplementary

Figure 2b). This neurotransmitter specificity is a potential avenue for exploring the preferential

vulnerability of the SN in PD.

Next, we postulated that genes whose expression defined the P7 SN DA neuron cluster might

illuminate their preferential vulnerability in PD. We identified 110 SN-specific genes, by first

finding all differentially expressed genes between P7 subset clusters and then using the Jensen-

Shannon distance to identify cluster specific genes (See Methods). Prior reports confirm the

expression of 49 of the 110 SN-specific genes (~45%) in postnatal SN (Supplementary Table 4).

We then sought evidence to confirm or exclude SN expression for the remaining, novel 61 genes

(55%). Of these, 25/61 (~41%) were detected in adult SN neurons by *in situ* hybridization (ISH)

of coronal sections in adult (P56) mice (Allen Brain Atlas, ABA; http://developingmouse.brain-

map.org), including *Col25a1*, *Fam184a*, *Ankrd34b*, *Nwd2*, and *Cadps2* (Figure 3, Supplementary

Table 5). Only 4/61 genes, for which ISH data existed in the ABA, lacked clear evidence of

expression in the adult SN (Supplementary Table 5). The ABA lacked coronal ISH data on 32/61

genes, thus we were unable to confirm their presence in the SN. Collectively, we identify 110

postnatal SN DA marker genes and confirm the expression of those genes in the adult mouse SN

for 74 (67%) of them, including 25 novel markers of this clinically relevant cell population that

we confirmed using the ABA image catalog.

We next asked whether we could identify significant relationships between cells defined as being

P7 SN DA neurons and distinctive transcriptional signatures in our data. We identify 16 co-

expressed gene modules by performing weighted gene co-expression network analysis

(WGCNA)[44,45] on all expressed genes of the P7 subset (Supplementary Figure 4; Supplementary

Table 6). By calculating pairwise correlations between modules and P7 subset clusters, we reveal

that 7/16 modules are significantly and positively correlated (Bonferroni corrected p < 3.5e-04)

with at least one subset cluster (Figure 3c). We graphically represent the eigenvalues for each

module in each cell in P7 t-SNE space, confirming that a majority of these significant modules

(6/7) displayed robust spatial, isotype enrichment (Figure 3d).

In order identify the biological relevance of these modules, each module was tested for

enrichment for Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways, Gene Ontology

(GO) gene sets, and Reactome gene sets. Two modules, the "brown" and "green" modules, were

significantly associated with the Parkinson's Disease KEGG pathway gene set (Figure 3c;

Supplementary Table 7). Interestingly, the "brown" module was also significantly correlated

with the P7 VTA population (P7.MB.1) and enriched for addiction gene sets (Supplementary

Table 7) highlighting the link between VTA DA neurons and addiction[46]. Strikingly, only the P7

SN cluster was significantly correlated with both PD-enriched modules (Figure 3c). This specific

correlation suggests these gene modules may play a role in the preferential susceptibility of the

SN in PD.

Integrating SN DA neuron specific data enables prioritization of genes within PD-associated

intervals

With these context-specific data in hand, we posited that SN DA neuron-specific genes and the

broader gene co-expression networks that correlate with SN DA neurons might be used to

prioritize genes within loci identified in PD GWAS. Such a strategy would be agnostic to prior

biological evidence and independent of genic position relative to the lead SNP, the traditional

method used to prioritize causative genes.

To investigate pertinent genes within PD GWAS loci, we identified all human genes within

topologically associated domains (TADs) and a two megabase interval encompassing each PD-

associated lead SNP. TADs were chosen because regulatory DNA impacted by GWAS variation

is more likely to act on genes within their own TAD[47]. While topological data does not exist for

261 SN DA neurons, we use TAD boundaries from hESCs as a proxy, as TADs are generally

262 conserved across cell types[48]. To improve our analyses, we also selected +/- 1 megabase interval

263 around each lead SNP thus including the upper bounds of reported enhancer-promoter

264 interactions[49,50]. All PD GWAS SNPs interrogated were identified by the most recent meta-

265 analysis (32 SNPs in total)[11], implicating a total of 1132 unique genes. We then identified

266 corresponding one-to-one mouse to human homologs (670/1132; ~59%), primarily through the

267 Mouse Genome Informatics homology database (Methods).

269 To prioritize these genes in GWAS loci, we developed a gene-centric score that integrates our

270 data as well as data in the public domain. We began by intersecting the PD loci genes with our

271 scRNA-seq data as well as previously published SN DA expression data[33], identifying 285 genes

272 (285/670; ~43%) with direct evidence of expression in SN DA neurons in at least one dataset.

273 Each PD-associated interval contained ≥1 SN-expressed gene (Table 1; Supplementary Table 8).

274 Emphasizing the need for a novel, systematic strategy, in 13/32 GWA intervals (~41%), the most

275 proximal gene to the lead SNP was not detectably expressed in mouse SN DA neuron

276 populations (Table 1; Supplementary Table 8). Surprisingly, two loci contained only one SN

277 DA-expressed gene (Table 1): *Mmp16* (*MMP16* locus) and *Tsnax* (*SIPA1L2* locus) (Figure 4a).

278 The relevance of these candidate genes to neuronal function/dysfunction is well supported[51,52].

279 This establishes gene expression in a relevant tissue as a powerful tool in the identification of

280 causal genes.

282 In order to prioritize likely diseases-associated genes in the remaining 30 loci, we scored genes

283 on three criteria: whether genes were identified as specific markers for the P7.MB.4 (SN) cluster

(Supplementary Table 2), whether the genes were differentially expressed between all P7 DA

neuron populations, and whether the genes were included in PD gene set enriched and SN

correlated gene modules uncovered in WGCNA (Supplementary Table 6). This strategy

facilitated further prioritization of a single gene in 14 additional loci including *SNCA*, *LRRK2*,

and *GCH1* loci (Figure 4a; Table 1). Importantly, using this approach we indict the familial PD

gene encoding alpha-synuclein (SNCA)*,* as responsible for the observed PD association within

4q22.1 (Figure 4a, Table 1). Thus, by using context-specific data alone, we were able to

prioritize a single candidate gene in exactly half of the PD-GWAS associated loci.

Furthermore, at loci in which a single gene did not emerge, we identified dosage sensitive genes

by considering the probability of being loss-of-function (LoF) intolerant (pLI) metric from the

ExAC database[53,54]. Since most GWAS variation is predicted to impact regulatory DNA and in

turn impact gene expression, it follows that genes in GWAS loci that are more sensitive to

dosage levels may be more likely to be candidate genes. With that in mind, the pLI for each gene

was used to further "rank" the genes within loci where a single gene was not prioritized. For

those loci, including *MAPT* and *DDRGK1* loci (Figure 4a), we report a group of top scoring

candidate genes (≤ 5) (Table 1). Expression of prioritized genes in the adult SN adds to the

validity of the genes identified as possible candidates (Figure 4b).

Two interesting examples that emerge from this scoring are found at the *MAPT* and *TMEM175-*

GAK-DGKQ loci. Although *MAPT* has previously been implicated in multiple neurodegenerative

phenotypes, including PD (OMIM: 168600), we instead prioritize two genes before it (*CRHR1*

and *NSF*; Table 1). We detect *Mapt* and *Nsf* expression consistently across all assayed DA

neurons (Figure 4c). By contrast, expression of *Crhr1,* encoding the corticotropin releasing

hormone receptor 1*,* is restricted to P7 DA neurons in the SN and the more mature OB neuronal

populations (Figure 4c). Similarly, at the *TMEM175-GAK-DGKQ* locus, our data shows that

although all three proximal genes are expressed in the SN, the adjacent *CPLX1* was one of the

prioritized gene (Table 1). There are multiple lines of evidence that strengthen *CPLX1* as a

candidate gene. Expression of CPLX1 is elevated both in the brains of PD patients and the brains

of mice overexpressing the *SNCA* A53T PD mutation[55,56]. Additionally, mice deficient in *CPLX1*

display an early-onset, cerebellar ataxia along with prolonged motor and behavioral

phenotypes[57,58]. However, the impact of *Cplx1* deficiency on the integrity of the nigrostriatal

pathway, to date, has not been explored. In order to confirm *CPLX1* as a candidate gene, we

performed immunohistochemistry (IHC) for *Th* in the *Cplx1* knockout mouse model

(Supplementary Table 9)[57–59]. We measured the density of *Th+* innervation in the striatum of

Cplx1 -/- mice and controls (Figure 4d, Supplementary Table 10) and found that *Cplx1* -/- mice

had significantly lower *Th+* staining in the striatum (p-value = 3.385e-08; Figure 4e). This

indicates that *Cplx1* KO mice have less *Th+* fiber innervation and a compromised nigrostriatal

pathway, supporting its biological significance in MB DA populations and to PD.

The systematic identification of causal genes underlying GWAS signals is essential in order for

the scientific and medical communities to take full advantage of all the GWAS data published

over the last decade. Taken collectively, we demonstrate how scRNA-seq data from disease-

relevant populations can be leveraged to illuminate GWAS results, facilitate systematic

prioritization of GWAS loci implicated in PD, and can leads to the functional characterization of

previously underexplored candidate genes.

DISCUSSION

Midbrain DA neurons in the SN have been the subject of intense research since being

definitively linked to PD nearly 100 years ago[60]. While degeneration of SN DA neurons in PD is

well established, they represent only a subset of brain DA populations. It remains unknown why

nigral DA neurons are particularly vulnerable. We set out to explore this question using scRNA-

seq. Recently, others have used scRNA-seq to characterize the mouse MB, including DA

neurons[33]. Here, we extend these data significantly, extensively characterizing the transcriptomes

of multiple brain DA populations longitudinally and discovering GRNs associated with specific

populations.

Most importantly, our data facilitate the iterative and biologically informed prioritization of gene

candidates for all PD-associated genomic intervals, the first such systematically generated

dataset. In practice, the gene closest to the lead SNP identified within a GWAS locus is

frequently treated as the prime candidate gene, often without considering tissue-dependent

context. Our study overcomes this by integrating genomic data derived from specific cell

contexts with analyses that are agnostic to one another. We posit that genes pertinent to PD are

likely expressed within SN DA neurons. This hypothesis is consistent with the recent description

of the "omnigenic" nature of common disease, wherein variation impacting genes expressed in a

disease tissue explain the vast majority of risk[7].

First, we identify intervals that reveal one primary candidate, i.e. those that harbor only one SN-

expressed gene. Next, we examine those intervals with many candidates, and prioritize based on

a cumulative body of biological evidence. In total, we prioritize 5 or fewer candidates in all 32

PD GWAS loci studied, identifying a single gene in sixteen loci (16/32; 50%) and three or fewer

genes in ~84% of loci (27/32). Ultimately this prioritization reduces the candidate gene list for

PD GWAS loci dramatically from 1132 genes to 65 genes.

The top genes we identify in three PD loci (*SNCA*, *FGF20*, *GCH1*) have been directly associated

with PD, MB DA development, and MB DA function[34] (OMIM: 163890, 128230). Furthermore,

our prioritization of *CPLX1* over other candidates in the *TMEM175-GAK-DGKQ* locus is

supported by multiple lines of evidence. Additionally, we demonstrate that the integrity of the

nigrostriatal pathway is disrupted in *Cplx1* knockout mice. Dysregulation of *CPLX1* RNA is also

a biomarker in individuals with pre-PD prodromal phenotypes harboring the *PARK4* mutation

(*SNCA* gene duplication)[61]. These results validate our approach and strengthen the argument for

the use of context specific data in pinpointing candidate genes in GWAS loci.

Many of the genes prioritized (Table 1) have been shown to have various mitochondrial

functions[62–68]. The identification of genes associated with mitochondrial functions is especially

interesting in light of the "omnigenic" hypothesis of complex traits[7]. Since mitochondrial

dysfunction has been extensively implicated in PD[69], the prioritized genes may represent "core"

genes that in turn can affect the larger mitochondrial-associated regulatory networks active in the

disease relevant cell-type (SN DA neurons). It is notable that one of these genes is the presenilin

associated rhomboid like gene or *PARL*. *PARL* cleaves *PINK1*, a gene extensively implicated in

PD pathology and recently a variant in *PARL* has been associated with early-onset PD (OMIM:

607858)[70–72].

While our method successfully prioritized one familial PD gene (*SNCA*), we do not prioritize

LRRK2, another familial PD gene harbored within a PD GWAS locus. *Lrrk2* is not prioritized

simply because it is not detectably expressed in our SN DA neuronal population. This is

expected as numerous studies have reported little to no *Lrrk2* expression in *Th+* MB DA neurons

both in mice and humans[73,74]. Instead, our method prioritizes *PDZRN4*. This result does not

necessarily argue against the potential relevance of *LRRK2* but instead provides an additional

candidate that may contribute to PD susceptibility. The same logic should be noted for two other

PD-associated loci, wherein our scoring prioritizes different genes (*KCNN3* and *CRHR1*/*NSF*,

respectively) than one previously implicated in PD (*GBA* and *MAPT*) (OMIM: 168600). Notably,

KCNN3, *CRHR1*, and *NSF*, all have previous biological evidence making them plausible

candidates[75–77].

Despite this success, we acknowledge several notable caveats. First, not all genes in PD-

associated human loci have identified mouse homologs. Thus, it remains possible that we may

have overlooked the contribution of some genes whose biology is not comprehensively queried

in this study. Secondly, we assume that identified genetic variation acts in a manner that is at

least preferential, if not exclusive, to SN DA neurons. Lastly, by prioritizing SN-expressed

genes, we assume that PD variation affects genes whose expression in the SN does not require

insult/stress. These caveats notwithstanding, our strategy sets the stage for a new generation of

independent and combinatorial functional evaluation of gene candidates for PD-associated

genomic intervals.

METHODS

Data availability

Raw data will be made available on Sequence Read Archive (SRA) and Gene Expression

Omnibus (GEO) prior to publication. Summary data is available where code is available below

(https://github.com/pwh124/DA_scRNA-seq).

Code Availability

Code for analysis, for the production of figures, and summary data is deposited at

https://github.com/pwh124/DA_scRNA-seq

Animals.

The Th:EGFP BAC transgenic mice (Tg(Th-EGFP)DJ76Gsat/Mmnc) used in this study were

generated by the GENSAT Project and were purchased through the Mutant Mouse Resource &

Research Centers (MMRRC) Repository (https://www.mmrrc.org/). Mice were maintained on a

Swiss Webster (SW) background with female SW mice obtained from Charles River

Laboratories (http://www.criver.com/). The Tg(Th-EGFP)DJ76Gsat/Mmnc line was primarily

maintained through matings between Th:EGFP positive, hemizygous male mice and wild-type

SW females (dams). Timed matings for cell isolation were similarly established between

hemizygous male mice and wild-type SW females. The observation of a vaginal plug was

defined as embryonic day 0.5 (E0.5). All work involving mice (husbandry, colony maintenance

and euthanasia) were reviewed and pre-approved by the institutional care and use committee.

Cplx1 knockout mice and wild type littermates used for immunocytochemistry were taken from a

colony established in Cambridge using founder mice that were a kind gift of Drs K. Reim and N.

Brose (Gottingen, Germany). *Cplx1* mice in this colony have been backcrossed onto a C57/Bl6J

inbred background for at least 10 generations. All experimental procedures were licensed and

undertaken in accordance with the regulations of the UK Animals (Scientific Procedures) Act

1986. Housing, rearing and genotyping of mice has been described in detail previously[57,58].

Mice were housed in hard-bottomed polypropylene experimental cages in groups of 5-10 mice in

a housing facility was maintained at $21 - 23^{o}C$ with relative humidity of $55 \pm 10\%$. Mice had *ad*

libitum access to water and standard dry chow. Because homozygous knockout *Cplx1* mice have

ataxia, they have difficulty in reaching the hard pellets in the food hopper and drinking from the

water bottles. Lowered waterspouts were provided and access to normal laboratory chow was

improved by providing mash (made by soaking 100 g of chow pellets in 230 ml water for 60 min

until the pellets were soft and fully expanded) on the floor of the cage twice daily. *Cplx1*

genotyping to identify mice with a homozygous or heterozygous deletion of the *Cplx1* gene was

conducted as previously described[57], using DNA prepared from tail biopsies.

Dissection of E15.5 brains.

At 15.5 days after the timed mating, pregnant dams were euthanized and the entire litter of

embryonic day 15.5 (E15.5) embryos were dissected out of the mother and immediately placed

in chilled Eagle's Minimum Essential Media (EMEM). Individual embryos were then

decapitated and heads were placed in fresh EMEM on ice. Embryonic brains were then removed

and placed in Hank's Balanced Salt Solution (HBSS) without Mg^{2+} and Ca^{2+} and manipulated

while on ice. The brains were immediately observed under a fluorescent stereomicroscope and

$EGFP^{+}$ brains were selected. $EGFP^{+}$ regions of interest in the forebrain (hypothalamus) and the

midbrain were then dissected and placed in HBSS on ice. This process was repeated for each

EGFP[+] brain. Four EGFP[+] brain regions for each region studied were pooled together for

dissociation.

Dissection of P7 brains.

After matings, pregnant females were sorted into their own cages and checked daily for newly

born pups. The morning the pups were born was considered day P0. Once the mice were aged to

P7, all the mice from the litter were euthanized and the brains were then quickly dissected out of

the mice and placed in HBSS without Mg^{2+} and Ca^{2+} on ice. As before, the brains were then

observed under a fluorescent microscope, EGFP[+] status for P7 mice was determined, and EGFP[+]

brains were retained. For each EGFP[+] brain, the entire olfactory bulb was first resected and

placed in HBSS on ice. Immediately thereafter, the EGFP[+] forebrain and midbrain regions for

each brain were resected and also placed in distinct containers of HBSS on ice. Five EGFP[+]

brain regions for each region were pooled together for dissociation.

Generation of single cell suspensions from brain tissue.

Resected brain tissues were dissociated using papain (Papain Dissociation System, Worthington

Biochemical Corporation; Cat#: LK003150) following the trehalose-enhanced protocol reported

by Saxena, et. al, 2012[78] with the following modifications: The dissociation was carried out at

37°C in a sterile tissue culture cabinet. During dissociation, all tissues at all time points were

triturated every 10 minutes using a sterile Pasteur pipette. For E15.5 tissues, this was continued

for no more than 40 minutes. For P7, this was continued for up to 1.5 hours or until the tissue

appeared to be completely dissociated.

Additionally, for P7 tissues, after dissociation but before cell sorting, the cell pellets were passed

through a discontinuous density gradient in order to remove cell debris that could impede cell

sorting. This gradient was adapted from the Worthington Papain Dissociation System kit.

Briefly, after completion of dissociation according to the Saxena protocol[78], the final cell pellet

was resuspended in DNase dilute albumin-inhibitor solution, layered on top of 5 mL of albumin-

inhibitor solution, and centrifuged at 70g for 6 minutes. The supernatant was then removed.

FACS and single-cell collection.

For each timepoint-region condition, pellets were resuspended in 200 μL of media without serum

comprised of DMEM/F12 without phenol red, 5% trehalose (w/v), 25 μM AP-V, 100 μM

kynurenic acid, and 10 μL of 40 U/μl RNase inhibitor (RNasin® Plus RNase Inhibitor, Promega)

at room temperature. The resuspended cells were then passed through a 40 uM filter and

introduced into a Fluorescence Assisted Cell Sorting (FACS) machine (Beckman Coulter MoFlo

Cell Sorter or Becton Dickinson FACSJazz). Viable cells were identified via propidium iodide

staining, and individual neurons were sorted based on their fluorescence (EGFP+ intensity, See

Supplementary Figure 2c) directly into lysis buffer in individual wells of 96-well plates for

single-cell sequencing (2 μL Smart-Seq2 lysis buffer + RNAase inhibitor, 1 μL oligo-dT primer,

and 1 μL dNTPs according to Picelli et al., 2014[79]. Blank wells were used as negative controls

for each plate collected. Upon completion of a sort, the plates were briefly spun in a tabletop

microcentrifuge and snap-frozen on dry ice. Single cell lysates were subsequently kept at -80°C

until cDNA conversion.

Single-cell RT, library prep, and sequencing.

Library preparation and amplification of single-cell samples were performed using a modified

version of the Smart-Seq2 protocol[79]. Briefly, 96-well plates of single cell lysates were thawed to

4°C, heated to 72°C for 3 minutes, then immediately placed on ice. Template switching first-

strand cDNA synthesis was performed as described above using a 5'-biotinylated TSO oligo.

cDNAs were amplified using 20 cycles of KAPA HiFi PCR and 5'-biotinylated ISPCR primer.

Amplified cDNA was cleaned with a 1:1 ratio of Ampure XP beads and approximately 200 pg

was used for a one-quarter standard sized Nextera XT tagmentation reaction. Tagmented

fragments were amplified for 14 cycles and dual indexes were added to each well to uniquely

label each library. Concentrations were assessed with Quant-iT PicoGreen dsDNA Reagent

(Invitrogen) and samples were diluted to ~2 nM and pooled. Pooled libraries were sequenced on

the Illumina HiSeq 2500 platform to a target mean depth of ~8.0 x 10^5 50bp paired-end

fragments per cell at the Hopkins Genetics Research Core Facility.

RNA sequencing and alignment.

For all libraries, paired-end reads were aligned to the mouse reference genome (mm10)

supplemented with the Th-EGFP$^+$ transgene contig, using HISAT2[80] with default parameters

except: -p 8. Aligned reads from individual samples were quantified against a reference

transcriptome (GENCODE vM8)[81] supplemented with the addition of the eGFP transcript.

Quantification was performed using cuffquant with default parameters and the following

additional arguments: --no-update-check –p 8. Normalized expression estimates across all

samples were obtained using cuffnorm[82] with default parameters.

Single-cell RNA data analysis.

Expression estimates.

Gene-level and isoform-level FPKM (Fragments Per Kilobase of transcript per Million) values

produced by cuffquant[82] and the normalized FPKM matrix from cuffnorm was used as input for

the Monocle 2 single cell RNA-seq framework[83] in R/Bioconductor[84]. Genes were annotated

using the Gencode vM8 release[81]. A CellDataSet was then created using Monocle (v2.2.0)[83]

containing the gene FPKM table, gene annotations, and all available metadata for the sorted

cells. All cells labeled as negative controls and empty wells were removed from the data.

Relative FPKM values for each cell were converted to estimates of absolute mRNA counts per

cell (RPC) using the Monocle 2 Census algorithm[16] using the Monocle function "relative2abs."

After RPCs were inferred, a new cds was created using the estimated RNA copy numbers with

the expression Family set to "negbinomial.size()" and a lower detection limit of 0.1 RPC.

QC Filtering.

After expression estimates were inferred, the cds containing a total of 473 cells was run through

Monocle's "detectGenes" function with the minimum expression level set at 0.1 transcripts. The

following filtering criteria were then imposed on the entire data set:

i. Number of expressed genes - The number of expressed genes detected in each cell in the

dataset was plotted and the high and low expressed gene thresholds were set based on

observations of each distribution. Only those cells that expressed between 2,000 and 10,000

genes were retained.

ii. Cell Mass - Cells were then filtered based on the total mass of RNA in the cells calculated by

Monocle. Again, the total mass of the cell was plotted and mass thresholds were set based on

observations from each distribution. Only those cells with a total cell mass between 100,000 and

1,300,000 fragments mapped were retained.

iii. Total RNA copies per cell - Cells were then filtered based on the total number of RNA

transcripts estimated for each cell. Again, the total RNA copies per cell was plotted and RNA

transcript thresholds were set based on observations from each distribution. Only those cells with

a total mRNA count between 1,000 and 40,000 RPCs were retained.

A total of 410 individual cells passed these initial filters. Outliers found in subsequent, reiterative

analyses described below were analyzed and removed resulting a final cell number of 396. The

distributions for total mRNAs, total mass, and number of expressed, can be found in

Supplementary Figure 1.

Log distribution QC.

Analysis using Monocle relies on the assumption that the expression data being analyzed follows

a log-normal distribution. Comparison to this distribution was performed after initial filtering

prior to continuing with analysis and was observed to be well fit.

Reiterative single-cell RNA data analysis.

After initial filtering described above, the entire cds as well as subsets of the cds based on "age"

and "region" of cells were created for recursive analysis. Regardless of how the data was

subdivided, all data followed a similar downstream analysis workflow.

Determining number of cells expressing each gene.

The genes to be analyzed for each iteration were filtered based on the number of cells that

expressed each gene. Genes were retained if they were expressed in > 5% of the cells in the

dataset being analyzed. These are termed "expressed_genes." For example, when analyzing all

cells collected together (n = 410), a gene had to be expressed in 20.5 cells (410 x 0.05 = 20.5) to

be included in the analysis. Whereas when analyzing P7 MB cells (n = 80), a gene had to be

expressed in just 4 cells (80 x 0.05 = 4). This was done to include genes that may define rare

populations of cells that could be present in any given population.

Monocle model preparation.

The data was prepared for Monocle analysis by retaining only the expressed genes that passed

the filtering described above. Size factors were estimated using Monocle's

"estimateSizeFactors()" function. Dispersions were estimated using the "estimateDispersions()"

function.

High variance gene selection.

Genes that have a high biological coefficient of variation (BCV) were identified by first

calculating the BCV by dividing the standard deviation of expression for each expressed gene by

the mean expression of each expressed gene. A dispersion table was then extracted using the

dispersionTable() function from Monocle. Genes with a mean expression > 0.5 transcripts and a

"dispersion_empirical" >= 1.5*dispersion_fit or 2.0*dispersion_fit were identified as "high

variance genes."

Principal component analysis (PCA).

PCA was then run using the R "prcomp" function on the centered and scaled log2 expression

values of the "high variance genes." PC1 and PC2 were then visualized to scan the data for

obvious outliers as well as bias in the PCs for age, region, or plates on which the cells were

sequenced. If any visual outliers in the data was observed, those cells were removed from the

original subsetted cds and all filtering steps above were repeated. Once there were no obvious

visual outliers in PC1 or PC2, a screeplot was used plot the PCA results in order to determine the

number of PCs that contributed most significantly to the variation in the data. This was manually

determined by inspecting the screeplot and including only those PCs that occur before the

leveling-off of the plot.

t-SNE and clustering.

Once the number of significant PCs was determined, t-Distributed Stochastic Neighbor

Embedding (t-SNE)[18] was used to embed the significant PC dimensions in a 2-D space for

visualization. This was done using the "tsne" package available through R with "whiten =

FALSE." The parameters "perplexity" and "max_iter" were tested with various values and set

according what was deemed to give the cleanest clustering of the data.

After dimensionality reduction via t-SNE, the number of clusters was determined in an unbiased

manner by fitting multiple Gaussian distributions over the 2D t-SNE projection coordinates using

the R package ADPclust[85] and the t-SNE plots were visualized using a custom R script. The

number of genes expressed and the total mRNAs in each cluster were then compared.

Differential expression Analyses.

Since the greatest source of variation in the data was between ages (Figure 1), differential

expression analyses and downstream analyses were performed separately for each age.

In order to find differentially expressed genes between brain DA populations at each age, the

E15.5 and P7 datasets were annotated with regional cluster identity ("subset cluster").

Differential expression analysis was performed using the "differentialGeneTest" function from

Monocle that uses a likelihood ratio test to compare a vector generalized additive model

(VGAM) using a negative binomial family function to a reduced model in which one parameter

of interest has been removed. In practice, the following models were fit:

"~subset.cluster" for E15.5 or P7 dataset

Genes were called as significantly differentially expressed if they had a q-value (Benjamini-

Hochberg corrected p-value) < 0.05.

Cluster specific marker genes.

In order to identify differentially expressed genes that were "specifically" expressed in a

particular subset cluster, R code calculating the Jensen-Shannon based specificity score from the

R package cummeRbund[86] was used similar to what was described in Burns *et al*[87].

Briefly, the mean RPC within each cluster for each expressed gene as well as the percentage of

cells within each cluster that express each gene at a level > 1 transcript were calculated. The

".specificity" function from the cummRbund package was then used to calculate and identify the

cluster with maximum specificity of each gene's expression. Details of this specificity metric can

be found in Molyneaux, *et al*[88].

To identify subset cluster specific genes, the distribution of specificity scores for each subset

cluster was plotted and a specificity cutoff was chosen so that only the "long right tail" of each

distribution was included (i.e. genes with a specificity score above the cutoff chosen). For each

iterative analysis, the same cutoff was used for each cluster or region (specificity $\geq$0.4). Once the

specificity cutoff was chosen, genes were further filtered by only retaining genes that were

expressed in >= 40% of cells within the subset cluster that the gene was determined to be

specific for.

Gene Set Enrichment Analyses.

Gene set enrichment analyses were performed in two separate ways depending upon the

situation. A Gene Set Enrichment Analysis (GSEA) PreRanked analysis was performed when a

ranked list (e.g. genes ranked by PC1 loadings) using GSEA software available from the Broad

Institute (v2.2.4)[89,90]. Ranked gene lists were uploaded to the GSEA software and a

"GSEAPreRanked" analysis was performed with the following settings: 'Number of

Permutations' = 1000, 'Collapse dataset to gene symbols' = true, 'Chip platform(s)' =

GENE_SYMBOL.chip, and 'Enrichment statistic' = weighted. Analysis was performed against

Gene Ontology (GO) collections from MSigDB, including c2.all.v5.2.symbols and

c5.all.v5.2.symbols. Top ten gene sets were reported for each analysis (Supplementary Table 1).

Figures and tables displaying the results were produced using custom R scripts.

Unranked GSEA analyses for lists of genes was performed using hypergeometric tests from the

R package clusterProfiler implemented through the functions 'enrichGO', 'enrichKEGG', and

'enrichPathway' with 'pvalueCutoff' set at 0.01, 0.1, 0.1, respectively with default settings[91].

These functions were implemented through the 'compareCluster' function when analyzing

WGCNA data.

Weighted Gene Co-Expression Network Analysis (WGCNA).

WGCNA was performed in R using the WGCNA package (v1.51)[44,45] following established

pipelines laid out by the packages authors (see

https://labs.genetics.ucla.edu/horvath/CoexpressionNetwork/Rpackages/WGCNA/ for more

detail). Briefly, an expression matrix for all P7 neurons containing all genes expressed in >= 20

cells (n = 12628) was used with expression counts in log2(Transcripts + 1). The data were

initially clustered in order to identify and remove outliers (n = 1) to leave 223 total cells

(Supplementary Figure 4a). The soft threshold (power) for WGCNA was then determined by

calculating the scale free topology model fit for a range of powers (1:10, 12, 14, 16, 18, 20)

using the WGCNA function "pickSoftThreshold()" setting the networkType = "signed". A power

of 10 was then chosen based on the leveling-off of the resulting scale independence plot above

0.8 (Supplementary Figure 4b). Network adjacency was then calculated using the WGCNA

function "adjacency()" with the following settings: power = 10 and type = "signed." Adjacency

calculations were used to then calculate topological overlap using the WGCNA function

"TOMsimilarity()" with the following settings: TOMtype = "signed." Distance was then

calculated by subtracting the topological overlap from 1. Hierarchical clustering was then

performed on the distance matrix and modules were identified using the "cuttreeDynamic"

function from the dynamicTreeCut package[92] with the following settings: deepSplit = T;

pamRespectsDendro = FALSE, and minClusterSize = 20. This analysis initially identified 18

modules. Eigengenes for each module were then calculated using the "moduleEigengenes()"

function and each module was assigned a color. Two modules ("grey" and "turquoise") were

removed at this point. Turquoise was removed because it contained 11567 genes or all the genes

that could not be grouped with another module. Grey was removed because it only contained 4

genes, falling below the minimum set module size of 20. The remaining 16 modules were

clustered (Supplementary Figure 4c) and the correlation between module eigengenes and subset

cluster identity was calculated using custom R scripts. Significance of correlation was

determined by calculated the Student asymptotic p-value for correlations by using the WGCNA

"corPvalueStudent()" function. Gene set enrichments for modules were determined by using the

clusterProfiler R package[91]. The correlation between the t-SNE position of a cell and the module

eigengenes was calculated using custom R scripts.

Prioritizing Genes in PD GWAS Loci.

Topologically Associated Domain (TAD) and Megabase Gene Data.

The data for human TAD boundaries were obtained from human embryonic stem cell (hESC)

Hi-C data[48] and converted from human genome hg18 to hg38 using the liftOver tool from UCSC

Genome Browser (http://genome.ucsc.edu/). PD GWAS SNP locations in hg38 were intersected

with the TAD information to identify TADs containing a PD GWAS SNP. The data for +/- 1

megabase regions surrounding PD GWAS SNPs was obtained by taking PD GWAS SNP

locations in hg38 and adding or subtracting 1e+06 from each location. All hg38 Ensembl

(version 87) genes that fell within the TADs or megabase regions were then identified by using

the biomaRt R package[93,94]. All genes were then annotated with PD locus and SNP information.

Mouse homologs for all genes were identified using human to mouse homology data from

Mouse Genome Informatics (MGI)

(http://www.informatics.jax.org/downloads/reports/HOM_MouseHumanSequence.rpt; Date

accessed: 07/07/2017). Homologs of protein coding genes that did not have a mouse homolog in

the data above were manually curated by searching the human gene name in the MGI database

(http://www.informatics.jax.org/). Of the 462 genes with no mouse homologs, only 60 (60/462,

~13%) were annotated as protein coding genes (Supplementary Figure 5). 17 loci include at least

one protein coding gene with no identified, one-to-one mouse homolog (Supplementary Figure

5). All 670 genes with mouse homologs are annotated as "protein_coding." Genes homologs

were manually annotated if a homolog was found to exist. The TAD and megabase tables were

then combined to create a final PD GWAS locus-gene table.

PD GWAS Loci Gene Scoring.

Genes within PD GWAS loci were initially scored using two gene lists: Genes with an average

expression ≥0.5 transcripts in the SN cluster in our data (points = 1) and genes with an average

expression ≥0.5 transcripts in the SN population in La Manno, *et al*[33] (points = 1). Further

prioritization was accomplished by using three gene lists: genes that were differentially

expressed between subset clusters (points = 1); Genes found to be "specifically" expressed in the

P7 MB SN cluster (points = 1); Genes found in the WGCNA modules that are enriched for PD

(points = 1). Expression in the SN cluster was considered the most important feature and was

weighted as such through the use of two complementary datasets with genes found to be

expressed in both receiving priority. Furthermore, a piece of external data, pLI scores for each

gene from the ExAC database[54], was added to the scores in order to rank loci that were left with

≥2 genes in the loci after the initial scoring. pLI scores

(fordist_cleaned_exac_r03_march16_z_pli_rec_null_data.txt) were obtained from

http://exac.broadinstitute.org/ (Date dowloaded: March 30, 2017).

In situ hybridization.

In situ hybridization data was downloaded from publically available data from the Allen Institute

through the Allen Brain Atlas (http://www.brain-map.org/). The image used in Figure 3a was

obtained from the Reference Atlas at the Allen Brain Atlas (http://mouse.brain-

map.org/static/atlas). URLs for all Allen Brain Atlas in situ data analyzed and downloaded for

SN marker genes (Figure 3b) are available in Supplementary Table 6. Data for SN expression *in*

situ data for PD GWAS genes (Figure 4b) were obtained from the following experiments: 1056

(*Th*), 79908848 (*Snca*), 297 (*Crhr1*), 74047915 (*Atp6v1d*), 72129224 (*Mmp16*), and 414 (*Cntn1*).

Data accessed on 03/02/17.

Single molecule in situ hybridization (smFISH).

740 For *in situ* hybridization experiments, untimed pregnant Swiss Webster mice were ordered from

741 Charles River Laboratories (Crl:CFW(SW); http://www.criver.com/). Mice were maintained as

742 previously described. Pups were considered P0 on the day of birth. At P7, the pups were

743 decapitated, the brain was quickly removed, and the brain was then washed in 1x PBS. The intact

744 brain was then transferred to a vial containing freshly prepared 4% PFA in 1x PBS and incubated

745 at 4^{o}C for 24 hours. After 24 hours, brains were removed from PFA and washed three times in 1x

746 PBS. The brains were then placed in a vial with 10% sucrose at 4^{o}C until the brains sunk to the

747 bottom of the vial (usually ~1 hour). After sinking, brains were immediately placed in a vial

748 containing 30% sucrose at 4^{o}C until once again sinking to the bottom of the vial (usually

749 overnight). After cryoprotection, the brains were quickly frozen in optimal cutting temperature

750 (O.C.T.) compound (Tissue-Tek) on dry ice and stored at -80^{o}C until use. Brains were sectioned

751 at a thickness of 14 micrometers and mounted on Superfrost Plus microscope slides

752 (Fisherbrand, Cat. # 12-550-15) with two sections per slide. Sections were then dried at room

753 temperature for at least 30 minutes and then stored at -80^{o}C until use.

755 RNAscope *in situ* hybridization (https://acdbio.com/) was used to detect single RNA transcripts.

756 RNAscope probes were used to detect *Th* (C1; Cat No. 317621, Lot: 17073A), *Slc6a3* (C2; Cat

757 No. 315441-C2, Lot: 17044A), *Lhx9* (C3; Cat No. 495431-C3, Lot: 17044A), and *Ldb2* (C3; Cat

758 No. 466061-C3, Lot: 17044A). The RNAscope Fluorescent Multiplex Detection kit (Cat No.

759 320851) and the associated protocol provided by the manufacturer were used. Briefly, frozen

760 tissues were removed from -80^{o}C and equilibrated at room temperature for 5 minutes. Slides

761 were then washed at room temperature in 1x PBS for 3 minutes with agitation. Slides were then

762 immediately washed in 100% ethanol by moving the slides up and down 5-10 times. The slides

were then allowed to dry at room temperature and hydrophobic barriers were drawn using a

hydrophobic pen (ImmEdge Hydrophobic Barrier PAP Pen, Vector Laboratories, Cat. # H-4000)

around the tissue sections. The hydrophobic barrier was allowed to dry overnight. After drying,

the tissue sections were treated with RNAscope Protease IV at room temperature for 30 minutes

and then slides were washed in 1x PBS. Approximately 100 uL of multiplex probe mixtures (C1

- *Th*, C2 - *Slc6a3*, and C3 - one of *Lhx9* or *Ldb2*) containing either approximately 96 uL C1: 2 uL

C2: 2 uL C3 (*Th*:*Slc6a3*:*Lhx9*) or 96 uL C1: 0.6 uL C2: 2 uL C3 (*Th*:*Slc6a3*:*Ldb2*) were applied

to appropriate sections. Both mixtures provided adequate *in situ* signals. Sections were then

incubated at 40°C for 2 hours in the ACD HybEZ oven. Sections were then sequentially treated

with the RNAscope Multiplex Fluorescent Detection Reagents kit solutions AMP 1-FL, AMP 2-

FL, AMP 3-FL, and AMP 4 Alt B-FL, with washing in between each incubation, according to

manufacturer's recommendations. Sections were then treated with DAPI provided with the

RNAscope Multiplex Fluorescent Detection Reagents kit. One drop of Prolong Gold Antifade

Mountant (Invitrogen, Cat # P36930) was then applied to each section and a coverslip was then

placed on the slide. The slides were then stored in the dark at 4°C overnight before imaging.

Slides were further stored at 4°C throughout imaging. Manufacturer provided positive and

negative controls were also performed alongside experimental probe mixtures according to

manufacturer's protocols. Four sections that encompassed relevant populations in the P7 ventral

MB (SN, VTA, etc.) were chosen for each combination of RNAscope smFISH probes and

subsequent analyses.

smFISH Confocal Microscopy.

RNAscope fluorescent *in situ* experiments were analyzed using the Nikon A1 confocal system

equipped with a Nikon Eclipse Ti inverted microscope running Nikon NIS-Elements AR 4.10.01

64-bit software. Images were captured using a Nikon Plan Apo λ 60x/1.40 oil immersion lens

with a common pinhole size of 19.2 μM, a pixel dwell of 28.8 μs, and a pixel resolution of 1024

x 1024. DAPI, FITC, Cy3, and Cy5 channels were used to acquire RNAscope fluorescence.

Positive and negative control slides using probe sets provided by the manufacturer were used in

order to calibrate laser power, offset, and detector sensitivity, for all channels in all experiments

performed.

smFISH image analysis and processing.

Confocal images were saved as .nd2 files. Images were then processed in ImageJ as follows.

First, the .nd2 files were imported into ImageJ and images were rotated in order to reflect a

ventral MB orientation with the ventral side of the tissue at the bottom edge. Next the LUT

ranges were adjusted for the FITC (range: 0-2500), Cy3 (range: 0-2500), and Cy5 (range: 0-

1500) channels. All analyzed images were set to the same LUT ranges. Next, the channels were

split and merged back together to produce a "composite" image seen in Figure 2. Scale bars were

then added. Cells of interest were then demarcated, duplicated, and the channels were split.

These cells of interest were then displayed as the insets seen in Figure 2.

Immunohistochemistry and quantification of *Th* striatum staining in *Cplx1* mice.

Mice (N=8 Cplx 1$^{-/-}$; N=3 WT littermates; ages between 4-7.5 weeks) were euthanized and their

brains fresh-frozen on powdered dry ice. Brains were sectioned at 35 mm and sections and

mounted onto Superfrost-plus glass slides (VWR International, Poole, UK). Sections were

peroxidase inactivated, and one in every 10 sections was processed immunohistochemically for

tyrosine hydroxylase. Sections were incubated in primary anti-tyrosine hydroxylase antibody

(AB152, Millipore) used at 1/2000 dilution in 1% normal goat serum in phosphate-buffered

saline and 0.2% Triton X-100 overnight at 4^oC. Antigens were visualised using a horseradish

peroxidase-conjugated anti-rabbit second antibody (Vector, PI-1000, 1/2000 dilution) and

visualized using diaminobenzidine (DAB; Sigma). The slides were stored in the dark (in black

slide boxes) at room temperature (21 C).

Images of stained striatum were taken using a Nikon AZ100 microscope equipped with a 2x lens

(Nikon AZ Plan Fluor, NA 0.2, WD45), a Nikon DS-Fi2 camera, and NIS-Elements AR 4.5

software. Appropriate zoom and light exposure were determined before imaging and kept

constant for all slides and sections. Density of Th+ DAB staining was measured using ImageJ

software. Briefly, images were imported into ImageJ and the background was subtracted (default

50 pixels with "light background" selected). Next, images were converted to 8-bit and the image

was inverted. Five measurements of density were taken for each side of a striatum in a section

along with a density measurement from adjacent, unstained cortex. Striosomes were avoided

during measuring when possible. Striatal measurements had background (defined as staining in

the adjacent cortex in a section) subtracted. The mean section measurements (intensity/pixels

squared) for each brain were calculated and represented independent measurements of the same

brain. Variances were compared between the WT and KO populations. A two sample t-test was

then used to compare WT vs. *Cplx1 -/-* section densities with the following parameters in R

using the "t.test" function: alternative = "two-sided", var.equal = "T".

ACKNOWLEDGEMENTS

The authors wish to thank Stephen M. Brown for implementation and optimization of smFISH.

Dr. Zhiguang Zheng and Mrs. Wendy Leavens for excellent technical support with the *Cplx1*

knockout mice and immunohistochemistry and Drs Kerstin Reim and Niels Brose for the gift of

the founder mice for the Cambridge Cplx1 knockout mice colony. This research was supported

in part by US National Institutes of Health grants R01 NS62972 and MH106522 to ASM and a

grant from CHDI *Inc*. to AJM.

838 AUTHOR CONTRIBUTIONS

839 PWH, ASM, and LAG designed the study and wrote the paper. PWH, SAM, WDL, GAC, and

840 AJM performed the experiments. PWH and LAG implemented the computational algorithms to

841 process the raw data and conduct analyses thereof. PWH, LAG, and ASM analyzed and

842 interpreted the resulting data. LAG contributed novel computational pipeline development.

843 Correspondence to ASM (andy@jhmi.edu) and LAG (loyalgoff@jhmi.edu).

845 FINANCIAL INTERESTS STATEMENT

846 The authors declare no competing financial interests.

REFERENCES

1. Visscher, P. M., Brown, M. A., McCarthy, M. I. & Yang, J. Five Years of GWAS Discovery. *Am. J. Hum. Genet.* **90,** 7–24 (2012).

2. Maurano, M. T. *et al.* Systematic Localization of Common Disease-Associated Variation in Regulatory DNA. *Science (80-.).* **337,** 1190–1195 (2012).

3. Farh, K. K. *et al.* Genetic and epigenetic fine mapping of causal autoimmune disease variants. *Nature* **518,** 337–343 (2015).

4. Smemo, S. *et al.* Obesity-associated variants within FTO form long-range functional connections with IRX3. (2014). doi:10.1038/nature13138

5. Gupta, R. M. *et al.* A Genetic Variant Associated with Five Vascular Diseases Is a Distal Regulator of Endothelin-1 Gene Expression In Brief A common sequence variant that perturbs long-range enhancer interactions mediates risk for different vascular diseases. A Genetic Variant Associated with Five Vascular Diseases Is a Distal Regulator of Endothelin-1 Gene Expression. *Cell* **170,** 522–533 (2017).

6. Lee, D. *et al.* A method to predict the impact of regulatory variants from DNA sequence. *Nat. Genet.* **47,** 955–61 (2015).

7. Boyle, E. A., Li, Y. I. & Pritchard, J. K. An Expanded View of Complex Traits: From Polygenic to Omnigenic. *Cell* **169,** 1177–1186 (2017).

8. de Rijk, M. C. *et al.* Prevalence of parkinsonism and Parkinson's disease in Europe: the EUROPARKINSON Collaborative Study. European Community Concerted Action on the Epidemiology of Parkinson's disease. *J Neurol Neurosurg Psychiatry* **62,** 10–15 (1997).

9. Pringsheim, T., Jette, N., Frolkis, A. & Steeves, T. D. The prevalence of Parkinson's disease: a systematic review and meta-analysis. *Mov Disord* **29,** 1583–1590 (2014).

10.   Savitt, J. M., Dawson, V. L. & Dawson, T. M. Diagnosis and treatment of Parkinson

disease: molecules to medicine. *J Clin Invest* **116,** 1744–1754 (2006).

11.   Nalls, M. a *et al.* Large-scale meta-analysis of genome-wide association data identifies six

new risk loci for Parkinson's disease. *Nat. Genet.* **56,** 1–7 (2014).

12.   Puschmann, A. Monogenic Parkinson's disease and parkinsonism: clinical phenotypes and

frequencies of known mutations. *Park. Relat Disord* **19,** 407–415 (2013).

13.   Klein, C. & Westenberger, A. Genetics of Parkinson's disease. *Cold Spring Harb*

Perspect Med **2,** a008888 (2012).

14.   Heintz, N. Gene Expression Nervous System Atlas (GENSAT). *Nat. Neurosci.* **7,** 483–483

(2004).

15.   Barallobre, M. J. *et al.* DYRK1A promotes dopaminergic neuron survival in the

developing brain and in a mouse model of Parkinson's disease. *Cell Death Dis.* **5,** e1289

(2014).

16.   Qiu, X. *et al.* Single-cell mRNA quantification and differential analysis with Census. *Nat*

Methods **14,** 309–315 (2017).

17.   Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories. *Nat.*

Methods (2017). doi:10.1038/nmeth.4402

18.   Van Der Maaten, L. & Hinton, G. Visualizing Data using t-SNE. *J. Mach. Learn. Res.* **9,**

2579–2605 (2008).

19.   Arenas, E., Denham, M. & Villaescusa, J. C. How to make a midbrain dopaminergic

neuron. *Development* **142,** 1918–36 (2015).

20.   Björklund, A. & Dunnett, S. B. Dopamine neuron systems in the brain: an update. *Trends*

Neurosci **30,** 194–202 (2007).

21.   Li, H., Zeitler, P. S., Valerius, M. T., Small, K. & Potter, S. S. Gsh-1, an orphan Hox

895   gene, is required for normal pituitary development. *EMBO J* **15,** 714–724 (1996).

896   22.   McNay, D. E., Pelling, M., Claxton, S., Guillemot, F. & Ang, S. L. Mash1 is required for

897   generic and subtype differentiation of hypothalamic neuroendocrine cells. *Mol Endocrinol*

898   **20,** 1623–1632 (2006).

899   23.   Campbell, J. N. *et al.* A molecular census of arcuate hypothalamus and median eminence

900   cell types. *Nat Neurosci* **20,** 484–496 (2017).

901   24.   Agoston, Z. *et al.* Meis2 is a Pax6 co-factor in neurogenesis and dopaminergic

902   periglomerular fate specification in the adult olfactory bulb. *Development* **141,** 28–38

903   (2014).

904   25.   Francis, F. *et al.* Doublecortin is a developmentally regulated, microtubule-associated

905   protein expressed in migrating and differentiating neurons. *Neuron* **23,** 247–256 (1999).

906   26.   Gokce, O. *et al.* Cellular Taxonomy of the Mouse Striatum as Revealed by Single-Cell

907   RNA-Seq. *Cell Rep.* **16,** 1126–1137 (2016).

908   27.   Vergaño-Vera, E. *et al.* Nurr1 blocks the mitogenic effect of FGF-2 and EGF, inducing

909   olfactory bulb neural stem cells to adopt dopaminergic and dopaminergic-GABAergic

910   neuronal phenotypes. *Dev Neurobiol* **75,** 823–841 (2015).

911   28.   Panman, L. *et al.* Sox6 and Otx2 control the specification of substantia nigra and ventral

912   tegmental area dopamine neurons. *Cell Rep.* **8,** 1018–1025 (2014).

913   29.   Viereckel, T. *et al.* Midbrain Gene Screening Identifies a New Mesoaccumbal

914   Glutamatergic Pathway and a Marker for Dopamine Cells Neuroprotected in Parkinson's

915   Disease. *Sci Rep* **6,** 35203 (2016).

30.   Kozicz, T., Vigh, S. & Arimura, A. The source of origin of PACAP- and VIP-

immunoreactive fibers in the laterodorsal division of the bed nucleus of the stria terminalis

in the rat. *Brain Res.* **810,** 211–219 (1998).

31.   Darland, T., Heinricher, M. M. & Grandy, D. K. Orphanin FQ/nociceptin: A role in pain

and analgesia, but so much more. *Trends in Neurosciences* **21,** 215–221 (1998).

32.   Cai, H., Liu, G., Sun, L. & Ding, J. Aldehyde Dehydrogenase 1 making molecular inroads

into the differential vulnerability of nigrostriatal dopaminergic neuron subtypes in

Parkinson's disease. *Transl. Neurodegener.* **3,** 27 (2014).

33.   La Manno, G. *et al.* Molecular Diversity of Midbrain Development in Mouse, Human, and

Stem Cells. *Cell* **167,** 566–580.e19 (2016).

34.   Itoh, N. & Ohta, H. Roles of FGF20 in dopaminergic neurons and Parkinson's disease.

Front Mol Neurosci **6,** 15 (2013).

35.   Poulin, J. F. *et al.* Defining midbrain dopaminergic neuron diversity by single-cell gene

expression profiling. *Cell Rep.* **9,** 930–943 (2014).

36.   Uhde, C. W., Vives, J., Jaeger, I. & Li, M. Rmst is a novel marker for the mouse ventral

mesencephalic floor plate and the anterior dorsal midline cells. *PLoS One* **5,** (2010).

37.   Ng, S. Y., Bogu, G. K., Soh, B. & Stanton, L. W. The long noncoding RNA RMST

interacts with SOX2 to regulate neurogenesis. *Mol. Cell* **51,** 349–359 (2013).

38.   Ellis, B. C., Molloy, P. L. & Graham, L. D. CRNDE: A long non-coding RNA involved in

CanceR Neurobiology, and DEvelopment. *Frontiers in Genetics* **3,** 1–15 (2012).

39.   Lin, M. *et al.* RNA-Seq of human neurons derived from iPS cells reveals candidate long

non-coding RNAs involved in neurogenesis and neuropsychiatric disorders. *PLoS One* **6,**

(2011).

40.   Guttman, M. *et al.* lincRNAs act in the circuitry controlling pluripotency and

differentiation. (2011). doi:10.1038/nature10398

41.   Morales, M. & Margolis, E. B. Ventral tegmental area: cellular heterogeneity,

connectivity and behaviour. *Nat Rev Neurosci* **18,** 73–85 (2017).

42.   Everitt, B. J., Hökfelt, T., Wu, J. Y. & Goldstein, M. Coexistence of tyrosine hydroxylase-

like and gamma-aminobutyric acid-like immunoreactivities in neurons of the arcuate

nucleus. *Neuroendocrinology* **39,** 189–191 (1984).

43.   Asmus, S. E. *et al.* Increasing proportions of tyrosine hydroxylase-immunoreactive

interneurons colocalize with choline acetyltransferase or vasoactive intestinal peptide in

the developing rat cerebral cortex. *Brain Res* **1383,** 108–119 (2011).

44.   Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9,** 559 (2008).

45.   Langfelder, P. & Horvath, S. Fast R Functions for Robust Correlations and Hierarchical

Clustering. *J Stat Softw* **46,** (2012).

46.   Pascoli, V., Terrier, J., Hiver, A. & Lüscher, C. Sufficiency of Mesolimbic Dopamine

Neuron Stimulation for the Progression to Addiction. *Neuron* **88,** 1054–1066 (2015).

47.   Dekker, J., Marti-Renom, M. A. & Mirny, L. A. Exploring the three-dimensional

organization of genomes: interpreting chromatin interaction data. *Nat. Rev. Genet.* **14,**

390–403 (2013).

48.   Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of

chromatin interactions. *Nature* **485,** 376–380 (2012).

49.   Lettice, L. A. *et al.* A long-range Shh enhancer regulates expression in the developing

limb and fin and is associated with preaxial polydactyly. *Hum. Mol. Genet.* **12,** 1725–1735

(2003).

50.   Benko, S. *et al.* Highly conserved non-coding elements on either side of SOX9 associated

964   with Pierre Robin sequence. *Nat. Genet.* **41,** 359–364 (2009).

965   51.   Yong, V. W., Power, C., Forsyth, P. & Edwards, D. R. Metalloproteinases in biology and

966   pathology of the nervous system. *Nat Rev Neurosci* **2,** 502–511 (2001).

967   52.   Li, Z., Wu, Y. & Baraban, J. M. The Translin/Trax RNA binding complex: clues to

968   function in the nervous system. *Biochim Biophys Acta* **1779,** 479–485 (2008).

969   53.   Doan, R. N. *et al.* Mutations in Human Accelerated Regions Disrupt Cognition and Social

970   Behavior. *Cell* **167,** 341–354.e12 (2016).

971   54.   Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536,**

972   285–291 (2016).

973   55.   Basso, M. *et al.* Proteome analysis of human substantia nigra in Parkinson's disease.

974   *Proteomics* **4,** 3943–3952 (2004).

975   56.   Gispert, S. *et al.* Complexin-1 and Foxp1 Expression Changes Are Novel Brain Effects of

976   Alpha-Synuclein Pathology. *Mol. Neurobiol.* **52,** 57–63 (2015).

977   57.   Glynn, D., Drew, C. J., Reim, K., Brose, N. & Morton, A. J. Profound ataxia in complexin

978   I knockout mice masks a complex phenotype that includes exploratory and habituation

979   deficits. *Hum. Mol. Genet.* **14,** 2369–2385 (2005).

980   58.   Glynn, D., Sizemore, R. J. & Morton, A. J. Early motor development is abnormal in

981   complexin 1 knockout mice. *Neurobiol. Dis.* **25,** 483–495 (2007).

982   59.   Kielar, C., Sawiak, S. J., Negredo, P. N., Tse, D. H. Y. & Morton, A. J. Tensor-based

983   morphometry and stereology reveal brain pathology in the complexin1 knockout mouse.

984   *PLoS One* **7,** (2012).

60.   Parent, M. & Parent, A. Substantia nigra and Parkinson's disease: a brief history of their

long and intimate relationship. *Can J Neurol Sci* **37,** 313–319 (2010).

61.   Lahut, S. *et al.* Blood RNA biomarkers in prodromal PARK4 and REM sleep behavior

disorder show role of complexin-1 loss for risk of Parkinson's disease. *Dis. Model. Mech.*

dmm.028035 (2017). doi:10.1242/dmm.028035

62.   Hildick-Smith, G. J. *et al.* Macrocytic anemia and mitochondriopathy resulting from a

defect in sideroflexin 4. *Am. J. Hum. Genet.* **93,** 906–914 (2013).

63.   Islam, M. M., Suzuki, H., Makoto, Y. & Tanaka, M. Primary structure of the smallest

(6.4-kDa) subunit of human and bovine ubiquinol-cytochrome c reductase deduced from

cDNA sequences. *Biochem Mol Biol Int.* **41,** 1109–1116 (1997).

64.   Swartz, D. A., Park, E. I., Visek, W. J. & Kaput, J. The e subunit gene of murine F1F0-

ATP synthase. Genomic sequence, chromosomal mapping, and diet regulation. *J. Biol.*

Chem. **271,** 20942–20948 (1996).

65.   Tomar, D. *et al.* MCUR1 Is a Scaffold Factor for the MCU Complex Function and

Promotes Mitochondrial Bioenergetics. *Cell Rep.* **15,** 1673–1685 (2016).

66.   Plovanich, M. *et al.* MICU2, a Paralog of MICU1, Resides within the Mitochondrial

Uniporter Complex to Regulate Calcium Handling. *PLoS One* **8,** (2013).

67.   Wonsey, D. R., Zeller, K. I. & Dang, C. V. The c-Myc target gene PRDX3 is required for

mitochondrial homeostasis and neoplastic transformation. *Proc. Natl. Acad. Sci. U. S. A.*

99, 6649–54 (2002).

68.   Curran, J. E. *et al.* Genetic variation in PARL influences mitochondrial content. *Hum.*

Genet. **127,** 183–190 (2010).

69.   Winklhofer, K. F. & Haass, C. Mitochondrial dysfunction in Parkinson's disease. *Biochim*

Biophys Acta **1802,** 29–44 (2010).

70.   Shi, G. *et al.* Functional alteration of PARL contributes to mitochondrial dysregulation in

Parkinson's disease. *Hum. Mol. Genet.* **20,** 1966–1974 (2011).

71.   Shi, G. & McQuibban, G. A. The Mitochondrial Rhomboid Protease PARL Is Regulated

by PDK2 to Integrate Mitochondrial Quality Control and Metabolism. *Cell Rep.* **18,** 1458–

1472 (2017).

72.   Jin, S. M. *et al.* Mitochondrial membrane potential regulates PINK1 import and

proteolytic destabilization by PARL. *J. Cell Biol.* **191,** 933–942 (2010).

73.   Galter, D. *et al.* LRRK2 expression linked to dopamine-innervated areas. *Ann Neurol* **59,**

714–719 (2006).

74.   Higashi, S. *et al.* Expression and localization of Parkinson's disease-associated leucine-

rich repeat kinase 2 in the mouse brain. *J Neurochem* **100,** 368–381 (2007).

75.   Soden, M. E. *et al.* Disruption of Dopamine Neuron Activity Pattern Regulation through

Selective Expression of a Human KCNN3 Mutation. *Neuron* **80,** 997–1009 (2013).

76.   Abuirmeileh, A., Harkavyi, A., Kingsbury, A., Lever, R. & Whitton, P. S. The CRF-like

peptide urocortin greatly attenuates loss of extracellular striatal dopamine in rat models of

Parkinson's disease by activating CRF1 receptors. *Eur. J. Pharmacol.* **604,** 45–50 (2009).

77.   Simunovic, F. *et al.* Gene expression profiling of substantia nigra dopamine neurons:

further insights into Parkinson's disease pathology. *Brain* **132,** 1795–1809 (2009).

78.   Saxena, A. *et al.* Trehalose-enhanced isolation of neuronal sub-types from adult mouse

brain. *Biotechniques* **52,** 381–385 (2012).

79.   Picelli, S. *et al.* Full-length RNA-seq from single cells using Smart-seq2. *Nat. Protoc.* **9,**

171–181 (2014).

80.   Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory

requirements. *Nat. Methods* **12,** 357–60 (2015).

81.   Mudge, J. M. & Harrow, J. Creating reference gene annotation for the mouse C57BL6/J

genome assembly. *Mamm Genome* **26,** 366–378 (2015).

82.   Trapnell, C. *et al.* Differential gene and transcript expression analysis of RNA-seq

experiments with TopHat and Cufflinks. *Nat. Protoc.* **7,** 562–78 (2012).

83.   Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by

pseudotemporal ordering of single cells. *Nat. Biotechnol.* **32,** 381–6 (2014).

84.   Huber, W. *et al.* Orchestrating high-throughput genomic analysis with Bioconductor. *Nat*

Methods **12,** 115–121 (2015).

85.   Wang, X.-F. & Xu, Y. Fast clustering using adaptive density peak detection. *Stat. Methods*

Med. Res. 1–14 (2015). doi:10.1177/0962280215609948

86.   Trapnell, C. *et al.* Differential analysis of gene regulation at transcript resolution with

RNA-seq. *Nat. Biotechnol.* **31,** 46–53 (2013).

87.   Burns, J. C., Kelly, M. C., Hoa, M., Morell, R. J. & Kelley, M. W. Single-cell RNA-Seq

resolves cellular complexity in sensory organs from the neonatal inner ear. *Nat Commun*

6, 8557 (2015).

88.   Molyneaux, B. J. *et al.* DeCoN: Genome-wide analysis of invivo transcriptional dynamics

during pyramidal neuron fate selection in neocortex. *Neuron* **85,** 275–288 (2015).

89.   Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for

interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102,** 15545–

15550 (2005).

90.   Mootha, V. K. *et al.* PGC-1alpha-responsive genes involved in oxidative phosphorylation

are coordinately downregulated in human diabetes. *Nat Genet* **34,** 267–273 (2003).

91.   Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R package for comparing

biological themes among gene clusters. *OMICS* **16,** 284–7 (2012).

92.   Langfelder, P., Zhang, B. & Horvath, S. Defining clusters from a hierarchical cluster tree:

the Dynamic Tree Cut package for R. *Bioinformatics* **24,** 719–720 (2008).

93.   Durinck, S. *et al.* BioMart and Bioconductor: A powerful link between biological

databases and microarray data analysis. *Bioinformatics* **21,** 3439–3440 (2005).

94.   Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the

integration of genomic datasets with the R/Bioconductor package biomaRt. (2009).

doi:10.1038/nprot.2009.97

FIGURES

Figure 1

1066

Figure 1. scRNA-seq analysis of isolated cells allows their separation by developmental time. a) Diagram of

scRNA-seq experimental procedures for isolating and sequencing EGFP+ cells. b) Principal component analysis

(PCA) on all cells collected using genes with highly variant transcriptional profiles. The greatest source of variation

(PC1) is explained by the time point at which the cells were collected, not the region from which the cells were

collected. c) The top ten Gene Ontology (GO) gene sets enriched in genes with positive (red) and negative (green)

PC1 loadings. Genes with negative PC1 loadings and negative normalized enrichment scores (NES) were enriched

for terms indicative of mitotically active cells. Genes with positive PC1 loadings and NES scores were enriched for

terms expected of more mature neurons. d) A t-distributed Stochastic Neighbor Embedding (t-SNE) plot of all

collected cells colored by regional identity. E15.5 cells cluster together while P7 cells cluster primarily by regional

identity. e) A t-SNE plot of all collected cells colored by subset cluster identity. Through iterative analysis,

timepoint-regions collected can be separated into multiple subpopulations (13 in total). Midbrain, Mb; Forebrain,

FB; Olfactory bulb; OB; Fluorescence activated cell sorting; FACS.

1079

1080   **Figure 2**

1081

Figure 2. Subclusters of P7 *Th+* neurons are identified based on marker gene analyses. a) A t-SNE plot of all P7

neurons collected using colored by subset cluster identity. The neurons mostly cluster by regional identity. b) t-SNE

plot of P7 FB neurons. P7 FB neurons cluster into two distinct populations. c) t-SNE plot of P7 OB neurons. P7 OB

neurons cluster into three populations. These populations represent a trajectory of Th+ OB maturation (Table S3) as

indicated by the red arrow. d) A t-SNE plot of P7 MB neurons. P7 MB neurons cluster into four clusters: the

substantia nigra (SN), the ventral tegmental area (VTA), the periaqueductal grey area (PAG), and a novel

progenitor-like population. e) Boxplots displaying the expression of four genes (*Th*, *Slc6a3*, *Lhx9*, and *Ldb2*) across

all subclusters identified. The novel P7 MB progenitor-like cluster (P7.MB.2) has a similar expression profile to

E15.5 MB neuroblast population (E15.MB.1) (Table S2). +/- 1.5x interquartile range is represented by the whiskers

on the boxplots. Data points beyond 1.5x interquartile range are considered as outliers and plotted as black points. f)

Representative image of multiplex single molecule fluorescent *in situ* hybridization (smFISH) for *Th*, *Slc6a3*, and

Lhx9, in the mouse ventral midbrain. Zoomed-in panels represent cell populations observed. Scale bar, 50 uM. g)

Representative image of multiplex smFISH for *Th*, *Slc6a3*, and *Ldb2*, in the mouse ventral midbrain. Zoomed-in

panels represent cell populations observed. h) Diagram of ventral midbrain summarizing the results of smFISH.

Th+/Slc6a3-/Lhx9+ and Th+/Slc6a3-/Ldb2+ cells are both found in the dorsal SN. Scale bar, 50 uM. NB,

neuroblast; SN, substantia nigra; VTA, ventral tegmental area; IPN, interpeduncular nucleus.

1098

Figure 3

Figure 3. Novel markers and gene modules reveal context specific SN DA biology. a) Reference atlas diagram from

the Allen Brain Atlas (ABA; http://www.brain-map.org/) of the P56 mouse ventral midbrain. b) Confirmation of

novel SN DA neuron marker genes through the use of ABA *in situ* hybridization data (http://www.brain-map.org/).

Coronal, P56 mouse *in situ* data was explored in order to confirm the expression of 25 novel SN markers.

Th expression in P56 mice was used as an anatomical reference during analysis. c) Correlation heatmap of the

Pearson correlation between module eigengenes and P7 Th+ subset cluster identity. Modules are represented by

their assigned colors at the bottom of the matrix. Modules that had a positive correlation with a subset cluster and

had a correlation P-value less than the Bonferroni corrected significance level (P-value < 3.5e-04) contain an

asterisk. SN cluster (P7.MB.4) identity, denoted by a black rectangle, was found to be highly correlated with two

modules ("green" and "brown") that were enriched for the "Parkinson's Disease" KEGG gene set (labeled with

"PD"). d) The eigengene value for each P7 neuron in the seven WGCNA modules shown to be significantly

positively associated with a subset cluster overlaid on the t-SNE plot of all P7 neurons (Figure 2a). Plotting of

eigengenes confirms strict spatial restriction of module association. Only the "lightcyan" module does not seem to

show robust spatial restriction.

1116

1117 **Figure 4**

1118

Figure 4. Context specific SN DA data allows for the prioritization of candidate genes in PD GWAS loci. a) A locus

plot displaying four megabase regions in the human genome (hg38) centered on PD GWAS SNPs in six loci. Genes

are displayed as boxes on their appropriate strand. Genes are shaded by their prioritization score and gene names are

displayed for genes with a score of 3 or higher in each locus. b) *In situ* hybridization from the ABA

([http://www.brain-map.org/)](http://www.brain-map.org/) of five prioritized genes along with *Th* for an anatomical reference. Coronal, P56

mouse *in situ* data was used. c) Boxplots displaying expression of prioritized genes from the *MAPT* locus (Figure 4a;

Table 1). +/- 1.5x interquartile range is represented by the whiskers on the boxplots. Data points beyond 1.5x

interquartile range are considered as outliers and plotted as black points. d) Representative light microscopy images

of *Th+* innervation density in the striatum of WT and *Cplx1* knockout (KO) mice. Scale bar, 1 mm. e) Boxplots

comparing the level of Th+ striatum innervation between WT and *Cplx1* KO mice. DAB staining density was

measured in 35 uM, horizontal sections in WT mice (mice = 3, sections = 16) and *Cplx1* KO mice (mice = 8,

sections = 40). Each point in the boxplot represents a stained, 35 uM section. Statistical analyses were performed

between conditions with section averages in order to preserve observed variability (WT n = 16, Cplx1 KO n = 40).

A two sample t-test revealed that Th+ innervation density was significantly lower in *Cplx1* KO mice (t = 6.4395, df

= 54, p = 3.386e-08). Data points outside of 1.5x interquartile range, represented by the whiskers on the boxplots,

are considered as outliers and plotted as black points.

1135
1136

1137
1138

1139

1140

1141

1142

1143

1144

1145

 TABLES

 Table 1. Summary of the systematic scoring of genes in 32 GWAS loci associated with PD

Locus	Genes	Mouse Homologs	Expressed Genes	Top Candidate Genes	Prioritized by	Closest Gene Expressed
ACMSD-TMEM163	17	9	5	*UBXN4; CCNT2; R3HDM1; RAB3GAP1*	SN expression; pLI	No, neither
BCKDK-STX1B	90	54	19	*MAPK3; VKORC1; BOLA2B*	SN expression; Differential expression; pLI	Yes, both
BST1	19	10	3	*CPEB2*	SN expression; Differential expression	No
CCDC62	50	41	17	*ARL6IP4*	SN expression; Differential expression	No
DDRGK1	54	37	13	*ATRN; NOP56; MRPS26; C20orf27; IDH3B*	SN expression; Differential expression; pLI	Yes
DLG2	15	7	4	*DLG2; CCDC90B*	SN expression; Differential expression; pLI	Yes
FAM47E-SCARB2	33	22	10	*G3BP2; CCNI; CDKL2*	SN expression; Differential expression; pLI	Yes - *SCARB2*, No - *FAM47E*
FGF20	18	12	8	*FGF20; ZDHHC2; TUSC3; MICU3; MTMR7*	SN expression; Differential expression; SN specific; pLI	Yes
GBA-SYT11	84	56	36	*KCNN3*	SN expression; Differential expression; SN specific; WGCNA module	Yes, both
GCH1	29	15	7	*GCH1*	SN expression; Differential expression; SN specific; WGCNA module	Yes
GPNMB	28	13	5	*RAPGEF5*	SN expression; Differential expression	No
HLA-DBQ1	164	99	31	*ATP6V1G2*	SN expression; Differential expression; WGCNA module	No
INPP5F	29	13	7	*PRDX3; NANOS1; INPP5F; SFXN4*	SN expression; Differential expression; pLI	Yes
ITGA8	27	15	5	*FAM171A1*	SN expression; Differential expression	No
KRT8P25-APOOP2	17	7	2	*CHMP2B*	SN expression; Differential expression	No, neither are in mouse
LRRK2	10	7	4	*PDZRN4*	SN expression; Differential expression; WGCNA module	No
MAPT	40	20	8	*CRHR1; NSF; MAPT*	SN expression; Differential expression; pLI	Yes
MCCC1	25	11	5	*DCUN1D1; ABCC5; PARL*	SN expression; Differential expression; pLI	No
MIR4697	14	11	6	*OPCML*	SN expression; Differential expression	Not in mouse
MMP16	9	2	1	*MMP16*	SN expression	Yes
NMD3	20	10	3	*B3GALNT1*	SN expression; Differential expression	Yes
RAB7L1-NUCKS1	42	31	11	*LRRN2; KLHDC8A; SRGAP2*	SN expression; Differential expression; pLI	Yes - *NUCKS1*, No - *RAB7L1* (*Rab29*)
RIT2	6	3	3	*RIT2; SYT4*	SN expression; Differential expression; pLI	Yes
SIPA1L2	15	6	1	*TSNAX*	SN expression	No
SNCA	11	7	4	*SNCA*	SN expression; Differential expression; WGCNA module	Yes
SPPL2B	80	65	29	*UQCR11*	SN expression; Differential expression; WGCNA module	Yes
SREBF1-RAI1	67	26	12	*COPS3; NT5M*	SN expression; Differential expression; pLI	No, neither

STK39	17	10	4	*STK39; B3GALT1*	SN expression; Differential expression; pLI	Yes
TMEM175-GAK-DGKQ	40	25	10	*MAEA; CPLX1; ATP5I; TMEM175*	SN expression; Differential expression; WGCNA module; pLI	Yes, all three
TMEM229B	24	15	8	*VTI1B; ATP6V1D*	SN expression; Differential expression; pLI	Yes
USP25	27	6	2	*HSPA13*	SN expression	Yes
VPS13C	11	5	2	*TLN2; RORA*	SN expression; pLI	No

1148   Scoring was carried out at described in the Results and Methods. Candidate genes are presented for each of 32 PD

1149   GWAS loci (identified by Nalls, *et al*[11]. Additional information for each PD GWAS locus is presented including the

1150   number of unique genes scored, the number genes with a mouse homolog, the number of genes expressed in mouse

1151   SN DA neurons, which data prioritized the top genes, and whether the closest gene to the lead SNP is expressed.

1152   Detailed scoring for each gene can be found in Supplementary Table 8.

1153

1154

1155

1156

1157

1158

1159