INTERCELLULAR COMMUNICATION VIA THE *COMX*-INDUCING PEPTIDE (XIP) OF *STREPTOCOCCUS MUTANS*

Justin Kaspar[1], Simon A. M. Underhill[2], Robert C. Shields[1], Adrian Reyes[1], Suzanne Rosenzweig[2], Stephen J. Hagen[2], and Robert A. Burne[1#]

[1]Department of Oral Biology, University of Florida, Gainesville, Florida 32610

[2]Department of Physics, University of Florida, Gainesville, Florida 32611-8440

Running title: Competence signaling in biofilms

Keywords: dental caries, genetic competence, biofilms, ComX/SigX, stress response

[#] Corresponding author.

Mailing address: Department of Oral Biology, University of Florida, College of Dentistry,

P.O. Box 100424, Gainesville, FL 32610.

Phone: (352) 273-8850

Fax: (352) 273-8829

E-mail: rburne@dental.ufl.edu

ABSTRACT

Gram-positive bacteria utilize externalized peptides to coordinate genetic and

physiological processes required for biofilm formation, stress responses and ecological

competitiveness. One example is activation of natural genetic competence by ComR and the

comX-inducing peptide (XIP) in *Streptococcus mutans*. Although the competence pathway can

be activated by addition of synthetic XIP in defined medium, the hypothesis that XIP is able to

function as an intercellular signal molecule has not been rigorously tested. Co-culture model

systems were developed that included a "sender" strain that overexpressed the XIP precursor

(ComS) and a "responder" strain harboring a GFP reporter fusion to a ComR-activated gene

(*comX*) promoter. The ability of the sender strain to provide a signal to activate GFP expression

was monitored at the individual cell and population levels using i) planktonic culture systems, ii)

cells suspended in an agarose matrix or iii) cells growing in biofilms. XIP was shown to be freely

diffusible and XIP signaling between the *S. mutans* sender and responder strains did not require

cell-to-cell contact. The presence of a sucrose-derived exopolysaccharide matrix diminished the

efficiency of XIP signaling in biofilms, possibly by affecting spatial distribution of XIP senders

and potential responders. Intercellular signaling was greatly impaired in a strain lacking the

primary autolysin, AtlA, and was substantially greater when the sender strain underwent lysis.

Collectively, these data provide evidence that *S. mutans* XIP can indeed function as a peptide

signal between cells and highlight the importance of studying signaling with endogenously-

produced peptide(s) in populations in various environments and physiologic states.

IMPORTANCE

The *comX*-inducing peptide (XIP) of *Streptococcus mutans* is a key regulatory element

in the activation of genetic competence, which allows cells to take up extracellular DNA. XIP has

been found in cell culture fluids and addition of synthetic XIP to physiologically receptive cells

can robustly induce competence gene expression. However, there is controversy as to whether

XIP can function as an intercellular communication signal. Here, we show that XIP indeed

signals between cells in *S. mutans*, but that cell lysis may be a critical factor, as opposed to a

dedicated secretion/processing system, in allowing for release of XIP into the environment. The

results have important implications in the context of the ecology, virulence and evolution of a

ubiquitous human pathogen and related organisms.

INTRODUCTION

Evidence for extensive complexity and diversity in bacterial intercellular communication strategies has rapidly accumulated since quorum sensing was first described by Greenberg and colleagues (1). The study of the genetics, biochemistry and ecology of interbacterial communication has yielded valuable insights into the molecular basis for the behaviors of bacterial communities and has revealed novel pathways that can be targeted to disrupt the formation, persistence and pathogenic potential of biofilms (2). Because bacterial biofilm communities exist at relatively high cell densities and are typically rich in exopolymeric materials, mass transport limitations contribute to the development of considerable spatial heterogeneity. Despite this heterogeneity, biofilm communities coordinate population-wide responses when challenged with exogenous and endogenous stressors by employing intercellular communication pathways. In the well-characterized Gram-negative quorum sensing systems, cell-cell communication occurs primarily via N-acyl homoserine lactones (3), whereas Gram-positive bacteria communicate mainly via oligopeptides, often through two-component signal transduction systems (TCS) consisting minimally of a membrane-associated histidine kinase and a cytosolic response regulator (4). Alternatively, signaling may occur through a pathway that requires active internalization of a peptide, followed by specific binding of the peptide by a cytosolic transcriptional regulator (5). Genetic competence in Gram-positive bacteria is usually controlled by one of two peptide-based systems. The most thoroughly studied are similar to the ComCDE pathway of *Streptococcus pneumoniae.* ComC is the pro-peptide of competence stimulating peptide (CSP), which is processed and externalized by a specialized secretion apparatus, then signals for the activation of competence genes via the ComDE TCS. The second is the more recently characterized ComRS pathway, which consists of the ComS precursor for *comX*-inducing peptide (XIP) and the cytosolic regulator ComR (6).

Streptococcus mutans, which colonizes the human oral cavity, is unusual among streptococci in that both CSP and XIP can activate transcription of the master regulator of

competence development, an alternative sigma factor encoded by the *sigX* (σ^X) or *comX* (7–9)

gene; SigX and ComX are homologues in different organisms. Thus, *S. mutans* has become an

intriguing model for the study of peptide-based communication strategies (10). CSP in *S.*

mutans is generated by secretion and processing of ComC through the ComAB exporter, with

further processing by the SepM protease to yield the most active form (18-aa) of CSP (11). In a

chemically-complex medium that is rich in peptides, such as brain heart infusion (BHI), CSP

signaling occurs via ComDE and directly activates bacteriocin production. However, ComCDE

of *S. mutans* are not true homologues of *S. pneumoniae* ComCDE, rather they are most similar

in sequence and function to the BlpCRH system of the pneumococcus, with which *S. mutans*

ComCDE share ancestral origin (6, 9). Provision of CSP to early exponential phase cells results

in up-regulation of *comX* transcription through an as-yet-undefined indirect mechanism, and

leads to a dramatic increase (~10^3-fold) in efficiency of transformation. The second route for

induction of competence in *S. mutans* involves the ComRS pathway, consisting of the Rgg-like

transcriptional regulator ComR and XIP, the latter being a 7-aa peptide derived from the C-

terminus of the 17-aa precursor, ComS. The current view of ComRS-dependent activation of

comX in *S. mutans* is that the ribosomally translated ComS peptide is exported into the

extracellular space and cleaved by a protease(s) to yield XIP (7, 9, 12), although the

mechanisms for secretion or processing have not been identified. Notably, three independent

research groups have detected XIP in supernatant fluids of *S. mutans* (13–15). In *Streptococcus*

thermophilus (16), the Eep protease processes ComS to XIP, but an equivalent function for

proteases in *S. mutans* with characteristics similar to Eep has not been demonstrated (13).

Additionally, in *S. thermophilus*, the exported XIP appears to remain associated with the cell

surface, as intercellular signaling by XIP in a co-culture system required cell-cell contact (16).

The current model for how the ComRS-XIP system functions is that, after XIP is produced and

externalized, it is transported back into the cell by the oligopeptide ABC transporter Opp (9).

The re-imported XIP can then be specifically bound by ComR to form a dimeric ComR-XIP

complex that functions as a transcriptional activator of the promoters of *comX* and *comS* (17).

The activation of *comS* creates a positive feedback loop that amplifies ComS and possibly XIP

production (9, 17). The ComR-XIP complex recognizes a ComR-box consisting of a 20-bp

palindromic motif with a conserved central GACA/TGTC inverted repeat (7, 18). The degree of

conservation of the ComR-box with the consensus sequence has been correlated with levels of

ComR-regulon expression (17).

Several different orthologs of ComR and ComS are present within streptococci (19).

ComR, along with apparent orthologous proteins that include PlcR of *Bacillus thuringiensis* and

PrgX of *Enterococcus faecalis,* are part of the RNPP superfamily (Rap/Npr/PlcR/PrgX) of

transcriptional regulators that interact with pheromones in cell-cell signaling pathways. Rgg

regulators typically contain an N-terminal helix-turn-helix (HTH) DNA-binding element and an

approximately 220-aa C-terminal alpha-helical domain thought to be involved in binding of

cognate small hydrophobic peptides (SHPs) (20). ComS peptides show substantial size and

primary sequence variation between species of streptococci (21, 22). The so-called type I ComS

peptides of *Streptococcus salivarius* and *S. thermophilus* harbor a P(F/Y)F motif and lack

charged residues. The type II peptides, like ComS of *S. mutans*, along with the Pyogenes and

Bovis groups of streptococci, contain a WW motif and basic and/or acidic residues (7, 17). In all

streptococci that possess ComRS, *comS* is located immediately downstream of *comR.* The

ComR-box upstream of *comS* is immediately followed by a "T-tract" that, in conjunction with the

ComR-box palindrome, may function as a Rho-independent transcriptional terminator for *comR*

(7, 17). All type II ComRS systems, including that of *S. mutans*, are located about 50 kbp from

the origin of replication and are associated with clusters of genes involved in purine biosynthesis

and the RuvB Holliday junction DNA helicase (7).

In the period since the ComRS system was first described by Gardan (23), Mashburn-

Warren (7) and their co-workers, there has been minimal investigation into how XIP is

externalized and whether it can actually function as an intercellular communication molecule for

S. mutans and related organisms. In this report, we designed and constructed an *S. mutans* co-

culture system that allowed us to investigate intercellular ComRS signaling through observation

of individual cells in planktonic cultures, in an agarose gel matrix and within single-species

biofilm communities.

RESULTS AND DISCUSSION

Development of in vitro co-cultivation models to monitor XIP signaling.

While numerous studies have used synthetic XIP peptide (sXIP) to explore the activation of *the* ComRS signaling pathway of *S. mutans* in planktonic cultures, little attention has been devoted to self-activation of the system, and only recently has sXIP-dependent gene activation been examined in *S. mutans* growing in biofilms (24), which is the normal growth mode of this organism in the oral cavity. Here, we developed a system to test whether the ComRS system can self-activate (intracellular) or cross-activate (intercellular) *comX* in the absence of exogenously supplied sXIP. Using previously described strains for our studies of genetic competence in *S. mutans* (25–27), we constructed a co-cultivation model consisting of two genetically modified *S. mutans*: a "sender" strain harboring a *comS* overexpressing plasmid (pIB184comS) along with a plasmid carrying the dsRed fluorescent protein (RFP) under the control of the *comX* promoter (P*comX*), and a "responder" strain expressing GFP also under the control of P*comX* (Figure 1A). The responder also carried the empty pIB184 vector so that the two strains were as genetically similar as possible. In certain cases, *comS* was deleted from the responder to remove any confounding effects of XIP production and auto-feedback by the responder strain. As structured, then, the sender constitutively expresses *comS* from the P_{23} promoter on pIB184, overproducing the pro-peptide ComS and hence XIP. Intracellular signaling or self-activation could be monitored in the sender strain through RFP fluorescence; the sender strain was easily distinguished from the co-cultivated responder in microscopy images by its lack of green fluorescence in combination with its strong red fluorescence. If XIP can serve as an intercellular signal, the responder strain should then import the XIP from the sender strain using Opp, where it can complex with the ComR regulator and activate the P*comX* promoter, resulting in production of GFP. Intercellular signaling can be visualized by fluorescence microscopy, by measuring total fluorescence in a plate reader or by flow cytometry of both planktonic cultures and/or dispersed biofilms.

The ability of the sender strain to activate *comX* in the responder strain, as measured by

GFP production, was first examined with a plate reader-based assay. Mid-exponential phase

planktonic cultures of both sender and responder strains were diluted 1:100 into fresh FMC

medium, such that the sender strain was present in a ratio of 2.3:1 with the responder strain

(Supplemental Figure 1), resulting in a final overall dilution in terms of absolute cell numbers of

approximately 1:50. After 18 hours of growth, no detectable fluorescence was observed when

wild-type (WT) *S. mutans* strain UA159 was co-cultivated with the P*comX*::*gfp* responder strain

(Figure 1B). In contrast, the *comS*-overexpressing strain, i.e. the sender, was able to elicit high

expression of P*comX*::*gfp* by the responder under similar growth conditions. To verify that

P*comX*::*gfp* expression was derived from imported extracellular XIP provided by the sender

strain, we evaluated GFP fluorescence with two mutant backgrounds of the responder strain,

Δ*comS* and Δ*opp*. GFP fluorescence was still detectable in the responder strain lacking *comS*,

albeit at a level 4-fold lower than in the strain with an intact *comS* gene. However, no GFP

production was observed in the *opp* mutant of the responder strain. Both results were as

expected: a lower level of fluorescence in a responder lacking *comS* would be unable to amplify

comX expression through the internal positive feedback arising from the ComR-XIP complex

activating the *comS* promoter, and XIP provided by the sender must be imported via Opp to

activate *comX*. Collectively, the results indicated that the sender was able provide a XIP signal

to the responder and that *comX* was activated by this XIP signal, consistent with the current

model for XIP-dependent activation of *comX* via active internalization by Opp, complexation with

ComR, and signal amplification through activation of *comS* transcription.

To provide a rough estimate of the amount of XIP generated by the sender strain, we

compared the fluorescence of the pIB184ComS/UA159;P*comX*::*gfp*/UA159 co-cultivation results

to the fluorescence of cultures obtained by addition of various concentrations of sXIP to a mixed

culture of UA159 and P*comX*::*gfp*/UA159 at the same 2.3:1 ratio used in the previous

experiment. From the measured fluorescence, it could be estimated that the amount of XIP

provided by the *comS* over-expresser was between 37.5 and 50 nM (Figure 1C), which is

similar to the concentration of XIP required to achieve ComRS-dependent activation of *comX* in

other *S. mutans* UA159 derivatives in microfluidic and planktonic experiments (9). These data

collectively show not only that intercellular signaling can occur via XIP and the ComRS pathway

in *S. mutans*, but also that signaling can occur in the co-culture model at physiologically relevant

levels of the peptide signal.

XIP signaling does not require cell-cell contact.

Early studies concluded that *S. mutans* XIP was a secreted, diffusible signal molecule

(14). However, studies with *S. thermophilus* demonstrated that cell-cell contact was required for

intercellular signaling by XIP (16). Since *S. mutans* and *S. thermophilus* are not closely related

within the genus *Streptococcus* and evolved in very different environments, we tested whether

the signaling observed in Figure 1 required cell-cell contact. Our initial test involved determining

whether exposure of the responder strain to cell-free supernatant fluids derived from the sender

strain could induce *comX* expression. Overnight cultures of either UA159 or the *comS*-

overexpressing strain were grown in FMC medium. After pelleting the cells, the supernates were

filter sterilized, the pH was adjusted to 7.0, and a concentrated solution of sterile glucose was

added to provide an additional 20 mM glucose. Supernates from *S. mutans* UA159

supplemented with 50 nM sXIP served as a positive control. The P*comX*::*gfp*/UA159 responder

was then suspended in the supernatant fluids and fluorescence was measured during growth of

the cells. No fluorescence was observed from the responder strain grown in the supernates

from UA159, unless sXIP was added (Figure 2A). However, when the P*comX*::*gfp* responder

strain was grown in supernates from the *comS*-overexpressing strain, robust fluorescence was

evident. These data clearly show that cell-free supernates of *S. mutans* are sufficient for

intercellular signaling, albeit overexpression of *comS* was required to yield sufficient signal

peptide in the supernates under the conditions tested.

Definitive evidence that the sender could produce a cell-free signal that could activate

comX expression in the responder strain was obtained when the *comS*-overexpressing sender

and PcomX::*gfp*/UA159 responder strains were cultured in separate chambers of a transwell

apparatus (Figure 2B), with the strains separated by a 10 μm thick, 0.4 μm pore-size

polycarbonate filter membrane. The membrane allows passage of small molecules and

peptides, but prevents physical contact between cells in the different compartments. Addition of

50 nM of sXIP to medium alone in the upper compartment activated P*comX*::*gfp* expression in

the lower compartment (Figure 2C). Consistent with the supernatant transfer experiments in

Figure 2A, UA159 was unable to stimulate *comX* expression in the P*comX*::*gfp* responder over

the course of a 24-h incubation. In contrast, the *comS*-overexpressing strain in the top

compartment of the transwell apparatus readily stimulated *comX* expression in the responder

strain in the lower compartment. These experiments show that signaling occurs through the

polycarbonate filter membrane and does not absolutely require contact, which suggests that XIP

is released by the sender strain, is diffusible and that signaling can occur without direct contact

between sender and responder strains of *S. mutans*.

In previous reports, the XIP peptide could be detected in supernatant fractions of the

wild-type strain UA159 at a high cell densit*y* (OD_{600} = 1.0) and cell supernates could also

stimulate a P*comX* reporter strain (14). In this study, we were not able to observe activation of

the P*comX*::*gfp* reporter by co-cultivation of strain UA159 or by using supernates from overnight

cultures of *S. mutans* UA159. The discrepancy between these studies and ours could be due to

the previously documented differences in XIP signaling seen between the chemically-defined

media CDM (14) and FMC (28). In particular, XIP signaling is exquisitely sensitive to low pH

(28, 29) and the drop of pH in FMC due to carbohydrate metabolism may be too rapid when

compared to that in CDM, since the latter is formulated with substantially greater (phosphate)

buffer capacity. While levels of XIP in supernates from high-density, overnight cultures have

been reported to be as high as 1 μM (14), such levels are not required to activate *comX*

transcription or induce genetic competence in early exponential phase cells when the pH is near

neutrality (28, 29) (Figure 1C). In fact, later in the growth phase, when XIP concentrations in the

medium may approach μM levels, the ComRS system may be inactive due to acidification of the

environment. Instead, optimal signaling occurs at the threshold for P*comX* activation and

theoretically at lower cell densities, when the inhibitory effects of low pH generated by

carbohydrate fermentation are minimalized. Thus, the co-culture model that is presented here

permits the study of competence signaling under conditions that may be more physiologically

relevant.

XIP diffuses in an agarose matrix

Oral biofilms, the natural habitat of *S. mutans,* are rich in exopolymeric material of

bacterial and host origin. To verify that the XIP is able to diffuse freely in an aqueous

environment containing *S. mutans*, we conducted a diffusion experiment using the

P*comX*::*gfp*/Δ*comS* reporter strain grown in FMC medium and embedded in low-melting-point

agarose/FMC medium. Cells were loaded into an IBIDI microslide channel (IBIDI GmbH) and

allowed to attach to the channel surface. A 2% low-melting point agarose/FMC mixture was then

injected into the channel to prevent advective transport of XIP and to immobilize the cells in the

channel. sXIP, diluted in FMC to a final concentration of 1 μM, was then deposited at one end of

the channel with an equal volume of water deposited at the other end to balance the hydrostatic

pressure along the length of the channel. Cells at different locations in the channel were then

imaged by fluorescence microscopy at regular time intervals as the XIP diffused through the

channel (Figure 3B-D). The diffusion coefficient of XIP was estimated by modeling the channel

as one-dimensional with a concentrated XIP source at one end from which the peptide spreads

diffusively (that is, the cells produce no XIP and there is no hydrodynamic flow). GFP

fluorescence versus time and spatial position was fit to a 1D diffusion equation (Figure 3A),

leading to an estimate for the diffusion coefficient of XIP through FMC/agarose. The resulting

estimate of $1.8 \pm 0.3 \times 10^{-6}$ cm^2/s was of the expected magnitude, based on the molecular mass

of XIP and diffusion in an aqueous medium. These data, along with the transwell experiment

presented in Figure 2, verify that XIP can signal by diffusion through aqueous medium,

independently of cell-cell contact.

Study of ComRS signaling between individual cells in an agarose matrix.

272   To begin visualizing ComRS signaling at the single-cell level using a co-culture

approach, we first mixed the *comS*-overexpressing sender with the responder, which carried

P*comX*::*gfp* in either the UA159 or Δ*comS* genetic background, in liquid FMC when the cells

reached an OD$_{600}$ = 0.1. The mixture was loaded into a microfluidic channel, followed by

injection of the 2% low-melting-point agarose/FMC mixture described above and in Figure 3.

Both the sender and the responder were imaged at 1-h intervals for up to 5 h to monitor

intercellular activation of P*comX* via diffusion of XIP. Figure 4 plots the measured fluorescence

of both RFP and GFP at the 2 h and 3 h time points. The scatter plots of individual cell

fluorescence in the culture are shown for both a UA159-based responder (Figures 4A and 4E)

and for a Δ*comS* responder (Figures 4C and 4G). In both co-cultures, P*comX*::*gfp* activation in

the responders was not significantly different from baseline fluorescence of the wild-type strain

UA159, which lacks a *gfp* reporter (data not shown). The experiment was repeated with a

positive control in which 1 μM sXIP was added into co-cultures containing responders with a

UA159 (Figures 4B and 4F) or Δ*comS* genetic background (Figures 4D and 4H). In this case,

most cells produced either red fluorescence (senders) or green fluorescence (responders), but

not both. This was observed for responders in the wild-type (Figure 4C) or Δ*comS* (Figure 4E)

genetic background. These data show that, in this experimental system, ComRS signaling from

the sender strain elicits a much weaker response from the responder than does the addition of

synthetic XIP or was seen in the microtiter-based experiments detailed above. Since diffusion

limitation cannot explain the results, we posited that the differences between this experiment

and those described above could arise from a number of different factors that include a shorter

experimental duration, a greater average distance between sender and responder cells in the

agarose matrix than in the co-culture studies, more limited lysis of the senders (see below), or

insufficient externalization of XIP by the sender under these particular conditions.

One interesting finding from this experiment is that self-activation from the sender strain

(intracellular signaling) was more apparent and of a greater magnitude than observed for the

responder strain (intercellular signaling). As seen in Figure 4A and 4C, the senders are self-

activating and the responders remain largely inactive, in terms of *comX* promoter activity, to the

signal in comparison to when sXIP was added (Figs. 4B and 4D). Such behavior may indicate,

at least within the confines of this experimental design, that the ComRS pathway is more

efficient for intracellular signaling than for intercellular communication. The *S. mutans*

competence pathway is unique among streptococci in that a ComCDE-like system, the

activation of which is dependent on the quorum sensing molecule CSP, is linked to ComRS and

that addition of sCSP to a peptide-rich medium results in a ~10^3 increase in transformation

efficiency. One must therefore consider, based on these findings, whether the ComRS pathway

in *S. mutans* functions primarily as an intracellular signal driven by its positive feedback loop.

Such a model for the importance of internal auto-activation by ComRS in complex medium,

such as BHI, has been previously presented (9). Importantly, an ability for XIP to function in self-

activation of cells and also as an intercellular signaling molecule are not mutually exclusive.

There is a high degree of specificity of the *S. mutans* ComR protein for XIP from *S. mutans*, but

XIP variants from other streptococcal species do not appear to interact with ComR of *S. mutans*;

whereas certain other ComR proteins are able to interact with XIP from non-cognate species

(21). Also of interest is the fact the ComS is extremely highly conserved between isolates of *S.*

mutans (32)*. Perhaps the stringency of the *S. mutans* ComR-XIP interaction and high degree of

conservation of XIP sequence in this organism reflects evolutionary pressures and niche

adaptations that favored activation of competence only with a signal input from other strains of

S. mutans in complex populations in humans.

ComRS signaling in biofilm populations analyzed by microscopy and flow cytometry.

As noted above, the natural environment for *S. mutans* is in biofilms, enmeshed in

exopolymeric material of bacterial and host origin, so we next evaluated ComRS signaling from

sender to responder in an *in vitro* biofilm model system where the exopolysaccharides were

generated during the experiment by enzymatic activities of *S. mutans*. To achieve an

approximately equal number of sender and responder cells in the model biofilm system for the

duration of the experiments, an optimal ratio of sender:responder inoculum was first determined.

Beginning with a 1:1 ratio of sender:responder grown in FMC medium and incubating the

biofilms for 18 hours, it was determined after biofilm dispersal and plating that the *comS*-

overexpressing strain was underrepresented with respect to the P*comX*::*gfp* responder strain,

with the responder constituting 91 ± 3% of the viable colonies recovered (Supplemental Figure

2). We attributed this observation to the somewhat slower growth of the *comS* overexpressing

strain (Supplemental Figure 3), and possibly to an enhancement in lytic behaviors associated

with *comS* overexpression. However, since ComRS signaling is impaired in acidic conditions

(28, 29), we reasoned that different inoculum ratios of sender:responder coupled with

replacement of the supernates with fresh medium after 6 h might enhance the representation of

the *comS*-overexpressing (sender) strain in the biofilm. Allowing for sufficient sender

representation was considered essential for the study of signaling. We tested three different

ratios of sender:responder biofilm inoculum: 4:1, 1:1, and 1: 4. The 4:1 sender:responder

inoculum resulted in roughly equivalent representation of the two strains after 18 h of incubation;

48 ± 10% of the viable colonies recovered were senders. The 4:1 ratio of sender:responder also

yielded the greatest amount of biomass, as measured by crystal violet staining at the end of the

18 h incubation period. Hence, all biofilm experiments were conducted with the 4:1 biofilm

inoculum of sender:responder, unless otherwise noted.

ComRS signaling within co-culture biofilm populations of *S. mutans* was visualized after

18 h of incubation in FMC medium, with replacement of spent medium with fresh medium at 6 h.

As a positive control, sXIP was added with the fresh medium at the 6-h time point in a final

concentration of 50 nM. Fluorescence images from both the sXIP-treated control and co-culture

biofilms that received no sXIP treatment were obtained for both the RFP-marked sender strain

and the GFP responder strain using confocal microscopy (Figure 5A). To quantify the proportion

of GFP-positive responders, biofilm populations grown under the same conditions were

harvested, sonicated to isolate individual cells, and analyzed via flow cytometry. A robust

response to sXIP was seen in the control biofilm population, with 65 ± 3% of single cells

expressing GFP (Figure 5B). In comparison, the co-culture biofilm population had measurable

GFP expression, but in a diminished proportion of the cells (12 ± 2%). In addition, the sender

strain constituted a greater proportion of the co-culture population (45 ± 4%) than in biofilms

treated with sXIP (18 ± 3%), likely due to enhanced sensitivity of the ComS-overproducing strain

to growth inhibition and/or induction of lysis by sXIP.

As the *comS*-overexpressing (sender) strain has apparently decreased fitness in biofilm

populations compared to a similar strain with an otherwise wild-type genetic background

(responder), we next grew biofilms with an original inoculum of either sender or responder, and

added the other strain at 6 h, along with fresh medium, to establish the co-cultures. Under

growth conditions in which the *comS*-overexpressing strain was allowed to establish first in the

biofilm, an increased proportion (61 ± 3%) of cells displayed *dsRed* fluorescence by flow

cytometry, compared to when the strains were added together in the initial inoculum (45 ± 4%).

However, the proportion of the GFP-positive PcomX::*gfp* strain remained unchanged when

contrasting co-inoculation with inoculation with the responder strain at 6 h (11 ± 4% to 12 ± 2%,

respectively). However, when the responder strain was established first, very little GFP or RFP

fluorescence activity was observed, either by confocal microscopy or flow cytometry. This is

most likely due to the *comS*-overexpressing strain not being able to establish and/or persist

after the strain with the wild-type genetic background had already established a biofilm. Overall,

ComRS signaling in co-culture biofilms was strongly dependent on the timing of introduction of

the second strain and the proportions of the sender; a finding that is not surprising in light of the

established impacts of low pH, growth phase and other factors on the efficiency of XIP-

dependent activation of *comX*.

Sucrose reduces intercellular XIP signaling in biofilms.

S. mutans strains are genomically and phenotypically diverse members of the oral

microbiome that predominantly colonize hard surfaces. A substantial body of evidence

implicates these organisms as primary etiological agents of human dental caries (31). Among

the many attributes that enable *S. mutans* to be an effective caries pathogen are its potent

acidogenic and aciduric properties, coupled with its capacity to utilize sucrose to form copious

quantities of extracellular polysaccharide (EPS) via three glucosyl- and one fructosyl-transferase

enzymes (Gtfs and Ftfs) (32). Gtfs, through *in situ* synthesis on the acquired enamel pellicle,

provide initial colonization sites for *S. mutans*, and produce the insoluble EPS matrix that

encases the bacteria, leading to the establishment of complex 3-dimensional biofilm structures

(33). Within these structures are highly diverse microenvironments that may influence diffusion

of chemical compounds and peptides that function in interbacterial communication. All

experiments herein were, to this point, conducted using FMC containing 20 mM glucose as the

sole carbohydrate source. To explore how addition of sucrose and production of an EPS matrix

would impact XIP signaling, we grew co-culture biofilms with low (2.5 mM to 15 mM), medium (5

mM to 10 mM) and high (9 mM to 2 mM) ratios of sucrose to glucose; sucrose is a dissacharide,

so the weight-to-volume concentrations of fermentable carbohydrates were constant across all

experiments. In all conditions in which sucrose was present, XIP activation within the co-culture

biofilm was evident (Figure 6). Interestingly, though, when sucrose was provided, a spatial

organization pattern between the *comS*-overexpressing sender and PcomX::*gfp*/UA159

responder became apparent, with the two strains segregating into different regions of the

biofilms (Figure 6A). In terms of both proportion of cells responding and overall *gfp* intensity

from the PcomX::*gfp* strain, measureable *gfp* fluorescence was reduced under conditions in

which sucrose was added to the growth medium (Figures 6B). In 20 mM glucose alone, the

proportion of *gfp* expressers was 19 ± 2% and median GFP intensity was 4.3 ± 0.5 au (arbitrary

fluorescent units), whereas 5 ± 1% of the population expressed *gfp* and intensity was recorded

as 3.1 ± 0.1 au under all sucrose conditions tested. Similar to the co-culture experiments within

the FMC/agarose gel, an increase in the strength of intracellular signaling was observed as RFP

intensity measured in the sender strain increased from 17.9 ± 0.9 au in biofilms formed in 20

mM glucose to 25.3 ± 0.9 au when the highest sucrose concentration was present.

406   To further evaluate the basis for the negative impact of sucrose and the resultant

polysaccharide matrix on intercellular signaling by XIP within biofilm populations, biofilms of

P*comX*::*gfp*/UA159 were grown with low, medium, and high ratios of sucrose to glucose for 5 h

before sXIP was added. GFP production was monitored at selected time intervals, both by

relative fluorescence in a plate reader and by confocal microscopy (Supplemental Figure 4).

Ample *comX* activation was noted in biofilms cultured in low-sucrose conditions when a final

concentration of 200 nM or 2 µM sXIP was provided. However, at the intermediate sucrose

concentration, activation was only seen with 2 µM sXIP and yielded a relative fluorescence that

was 3.5-fold lower compared to the low-sucrose condition. The most substantial effects were

seen in the high-sucrose condition, where 2 µM sXIP was unable to activate the P*comX*

responder strain. In fact, 10 µM sXIP was needed to measure GFP production within the high-

sucrose biofilms. Collectively, these data highlight that provision of sucrose, which dramatically

alters the biofilm EPS matrix and influences the physiology and transformability of *S. mutans*

(34, 35), has an overall negative impact on the response of cells to XIP.

One potential explanation for the reduced response by the responder strain when

various amounts of sucrose were present in the growth medium is slower diffusion of the

secreted XIP peptide within the biofilms due to increased EPS formation. The reduced diffusion

of XIP could lead to changes in the spatial distribution and temporal dynamics of ComRS

signaling within biofilm populations, leading to increased phenotypic heterogeneity within the

biofilm. Such observations were recently noted in the study of quorum sensing systems within

biofilms under flowing conditions (36). The images in Figure 6A show that some spatial

correlation between sender and responder is evident, such that clusters of senders and

responders appear segregated, even though the biofilm inoculum consists of uniform

suspension of senders and responders. A comparison of merged and brightfield images also

shows that certain cells within the biofilm population remain unresponsive to the sender's signal

(Supplemental Figure 5). Similar phenotypic heterogeneity was recently noted in monoculture

biofilms that had been treated with sXIP peptide, and notably, there was little evidence of cell

death in biofilms cultured under similar conditions (24). Thus, it will be interesting to unravel why

these cells within the biofilm are unresponsive, although it can be hypothesized that some are

unresponsive due to slow growth or to being confined to microenvironments that inhibit uptake

or responses to signal(s) inputs. Further study of the spatial organization of various modified

sender and responder strains should yield a more complete understanding the role of biofilm

architecture on signaling, and vice versa.

Lysis of the sender strain contributes significantly to XIP signaling.

A critical gap in our current understanding of the ComS/XIP system of *S. mutans* is that

the export apparatus and protease that generate externalized XIP from ComS have not

identified. Several attempts have been made to identify both the presumptive exporter and the

protease in *S. mutans* (13, 20, 37). The ABC transporter PptAB was identified recently in *S.*

pyogenes as an SHP exporter, but XIP secretion in *S. mutans* was only partially reduced in a

pptAB deletion (38). These results indicate that other mechanisms exist by which XIP appears

in the supernatant fraction of *S. mutans* cultures (13, 15). Since the studies detailed above

provide ample evidence for intercellular activation of ComRS under conditions in which

substantial amounts of cell lysis occur, such as overnight planktonic growth (Figure 2) and

growth within biofilms (Figures 5 and 6), but a lack of intercellular activation when early

exponential phase cells were spatially separated and fixed within an agarose suspension

(Figure 4), we examined whether active cell lysis was important for the release of the ComS/XIP

signal to potential responders. To accomplish this, we monitored XIP signaling activity in

supernates of *comS*-overexpressing cells in a wild-type genetic background and in a strain

lacking the major autolysin, AtlA (Δ*atlA*) (Figure 7) (39). Whereas supernates from UA159 or the

Δ*atlA* mutant failed to activate the P*comX*::*gfp*/UA159 reporter, the supernates from the *comS*-

overexpressing strain in an AtlA-positive strain activated P*comX* expression in the responder,

similar to the results reported above (Figure 2A). However, no detectable GFP fluorescence

was observed when the supernates from the *comS*-overexpressing strain carrying the *atlA*

deletion was supplied to the responder strain (Figure 7A). These data are corroborated by

growth curves of the P*comX*::*gfp*/UA159 responder strain in the selected supernates, as growth

in the supernates from the 184*comS*/UA159 strain was slower than in supernates from UA159,

Δ*atlA* and 184*comS*/Δ*atlA,* attributable to diminished XIP-dependent growth inhibition and/or

XIP-induced lysis of the responder. Experiments using a P*comX*::*gfp*/Δ*comS* responder strain,

which eliminates the auto-feedback loop and self-activation through activation of *comS* by

ComR-XIP, also showed no activation in response to the *atlA* mutant supernates (Figures 7B).

These data provide intriguing new evidence that externalization of ComS/XIP in *S. mutans* may

depend, entirely or in part, on cell lysis or loss of membrane integrity, rather than on a dedicated

secretion apparatus, as is the case for a number of other streptococci.

A number of properties of the ComRS system of *S. mutans* in particular may provide an

explanation for why cell lysis could be the primary pathway for XIP externalization. First,

ComRS is extremely highly conserved among *S. mutans* isolates, with almost no variation in

ComS sequence and none in the sequence of XIP (32). Similarly, ComR of *S. mutans* very

specifically recognizes *S. mutans* XIP, but not XIP from other sources. It is also notable that *S.*

mutans has an obligatory biofilm lifestyle, where regulated cell death and lysis, along with eDNA

release (42), are critically important for biofilm maturation and stability (43). *S. mutans* may thus

have evolved in a way that a dedicated ComS/XIP exporter was dispensable, as autolysis of

bacterial cells during biofilm growth could be sufficient to release XIP at levels that are effective

for its function(s) (42). The release by lysis also would serve as an altruistic signal to viable *S.*

mutans that *S. mutans* DNA has been released into the environment, which may have aided in

the diversification of the species by tuning competence to the preferential assimilation of DNA

that could be sufficiently homologous to recombine with the competent population. If confirmed,

these findings may also explain why *S. mutans* bacteriocin production and autolysis are more

intimately intertwined with competence development than in some other streptococcal species

(44, 45). CSP-induced activation of bacteriocins, some of which can kill sensitive strains of *S.*

mutans, could trigger lysis in the sensitive sub-population, which would then release XIP and

DNA for the now-competent members of the biofilm. The hypothesis also takes into

consideration Wenderska *et. al.*'s observation that deletion of *comX* reduces the abundance of

XIP in culture supernates (37), as this deletion would remove competence-driven lysis as a

release pathway. It is also interesting to posit that maturation of ComS to XIP may actually be a

consequence of induced cell lysis and activation of proteolytic pathways, such that when a cell

lyses, XIP would be released.

SUMMARY

Recent analysis of bacterial communication at the single cell level and *in situ* in biofilm

systems has led to revision of numerous views of intercellular signaling in "natural" populations.

While many of these advances have been made with well-established model organisms (e.g.

Bacillus subtilis, Staphylococcus aureus, Escherichia coli, Pseudomonas aeruginosa), the oral

pathogen *S. mutans* regulates multiple biological processes, including genetic competence,

bacteriocin production, biofilm maturation and tolerance of environmental insults, through a suite

of small hydrophobic peptides (26, 27) and dedicated transcriptional regulators (44, 45). Many

of these are unique to this organism, function differently than in paradigm organisms, and are

critical to the ability of *S. mutans* to cause disease.

Overall, the work presented here highlights a new model system for studying Gram-

positive bacterial communication based on naturally-derived signal oligopeptides from an

overexpressing population, as opposed to the addition of synthetic peptide(s). The model shows

that XIP can, in fact, act as an intercellular signal that does not require cell-cell contact. The

model also allows investigation of responses to XIP in a biological setting that more closely

mimics the natural growth environment of the organism. This work also emphasizes the

differences in signaling behaviors that occur between planktonically grown cultures and those

that are grown within biofilms, where spatial and temporal distribution patterns can have a more

significant impact on interpretation of signal inputs. Development of this system facilitates the

modelling of ComRS behaviors within an environment such as dental plaque, where competition

of *S. mutans* with commensal streptococci strongly influences the pathogenic potential of the

biofilms (46). Importantly, this work also highlights for the first time the connection between lysis

and XIP release, offering an alternative hypothesis for why XIP has been detected mainly in

culture supernates. Future investigations can provide insight into how signaling impacts genetic

and physiological control of early biofilm growth, responses to environmental stresses and

competition between competing oral species, ultimately yielding information that can be used to

disrupt key components of the signaling circuit to decrease the proportions of pathogens in

dental biofilms.

EXPERIMENTAL PROCEDURES

Bacterial strains and growth conditions. S. mutans wild-type strain UA159 and its derivatives

(Table 1) were grown in either brain heart infusion (BHI) (Difco) or FMC medium (47) that was

supplemented with 10 µg ml^{-1} erythromycin and 1 mg ml^{-1} of kanamycin or spectinomycin, as

needed. Unless otherwise noted, cultures were grown overnight in BHI medium with the

indicated antibiotics, if needed, at 37°C in a 5% CO_2, aerobic atmosphere. The next day,

cultures were harvested by centrifugation, washed twice in 1 mL of phosphate-buffered saline

(PBS), and resuspended in PBS to remove all traces of BHI. PBS cell suspensions were then

transferred to 5 mL polystyrene round-bottom tubes (Corning Incorporated). The samples were

then sonicated using a Fisher Scientific Model 120 Sonic Dismembrator in the water bath mode

at 100% amplitude for three intervals of 30 s each, with placement on ice for the intervals.

Sonicated cell suspensions were subjected to a final centrifugation to remove any cellular debris

and resuspended in the desired medium before diluting to begin each experiment. Synthetic XIP

(sXIP, aa sequence = GLDWWSL), corresponding to residues 11-17 of ComS, was synthesized

and purified to 96% homogeneity by NeoBioSci (Cambridge, MA). The lyophilized sXIP was

reconstituted with 99.7% dimethyl sulfoxide (DMSO) to a final concentration of 2 mM and stored

in 100 µL aliquots at -20°C

Construction of bacterial strains. Mutant strains of *S. mutans* were created using a PCR ligation

mutagenesis approach, as previously described (48). Overexpression of genes was achieved

by amplifying the structural genes of interest from *S. mutans* UA159 and cloning into the

expression plasmid pIB184 (25). Transformants were confirmed by PCR and sequencing after

selection on BHI agar with appropriate antibiotics. Plasmid DNA was isolated from *E. coli* using

QIAGEN (Chatsworth, Calif.) columns, and restriction and DNA-modifying enzymes were

obtained from Invitrogen (Gaithersburg, Md.) or New England Biolabs (Beverly, Mass.). PCRs

were carried out with 100 ng of chromosomal DNA by using *Taq* DNA polymerase, and PCR

products were purified with the QIAquick kit (QIAGEN).

Measurements of GFP fluorescence with plate reader. For measurements of GFP fluorescence,

co-cultures were inoculated from washed and sonicated overnight cultures in FMC medium at a

1:1 ratio, unless noted otherwise. Inoculated medium (175 µL) was added to each well along

with a 50 µL mineral oil overlay in a Costar™ 96 well assay plate (black plate with clear bottom;

Corning Incorporated) and incubated at 37°C. At intervals of 30 minutes for a total of 18 hours,

absorbance at 600 nm along with GFP fluorescence (excitation 485/20 nm, emission 528/20

nm) was measured with a Synergy 2 multimode microplate reader (BioTek). Relative

expression was calculated by subtracting the background fluorescence of UA159 (mean from

six replicates) from raw fluorescence units of the reporter strains and then dividing by OD_{600}.

Measurement of diffusive XIP spreading. To estimate a diffusion constant for sXIP and the

ComRS system, P*comX*::*gf*p/Δ*comS* cells were injected into an IBIDI microslide (ibidi GmbH, µ-

slide VI), a slide having 6 parallel, narrow channels, each with a loading port at both ends. The

cells were allowed to settle to the surface of the channels before a 2% low-melting-point

agarose/FMC gel was pushed through each channel to immobilize individual cells on the

channel window. Injection of the agarose/FMC mixture removed any cells that were not stuck to

the slide. sXIP that was reconstituted in DMSO and diluted into FMC to a final 1 µM

concentration was then deposited in one port of each channel, with an equal volume of DI water

in the opposite port to balance the hydrostatic pressure in the channel. The resulting culture was

then incubated at 37°C with mineral oil sealing the loading ports to minimize drying of the cooled

agarose. Green fluorescence of individual P*comX*::*gf*p/Δ*comS* reporter cells was measured as

a function of position and time, relative to the XIP loading, in order to estimate the diffusion

constant for sXIP in the medium. Cells were imaged at 20X (CFI Plan Fluor DLL, NA 0.5, Nikon)

onto a cooled CCD camera (CoolSNAP HQ2, Photometrics). Images were collected in phase

contrast, GFP fluorescence (using a Nikon C-FL GFP HC HISN zero shift filter cube), or red

fluorescence (C-FL Y-2E/C Texas Red filter cube, Nikon). Three images of each channel were

collected at intervals of one hour, for periods up to five hours, after introduction of sXIP. The

expression of GFP in individual cells was quantified by a previously described method (9, 49).

Fitting of the diffusion coefficient was performed in Matlab® (The Mathworks, Inc.) and was

done by minimizing the difference between measured GFP response and one calculated from a

1D diffusion equation model with a delta function of XIP concentration at the origin as an initial

condition. The error on this fit was checked by a 1000-iteration bootstrap method, with the

standard deviation of calculated diffusion coefficient values used as the fit error.

Co-culture within agarose/FMC mixture. P*comX*::*dsRed*/pIB184*comS* (sender strain) and

P*comX*::*gfp* reporter in either a UA159 or Δ*comS* background (responder strain) were mixed

together at OD_{600} of 0.1 in liquid FMC at a 2.3::1 ratio before injection into an IBIDI microslide

and immobilized with a 2% low-melting temperature agarose/FMC mixture. The resulting culture

was grown in an incubated chamber with mineral oil on top of the injection holes to minimize

drying. Imaging was performed at intervals to examine reporter fluorescence, as previously

described (9, 29). sXIP (1 μM) was added as a positive control, and channels containing solely

GFP reporter cells were also imaged as a negative control.

Fitness assessment. Fitness between the *comS*-overexpressing strain and the PcomX::*gfp*

responder strain carrying a kanamycin resistance marker (pPMZ) was determined from 18 h

biofilms inoculated with either a 4::1, 1::1, or 1::4 ratio of sender::responder and grown in 6 well

polystyrene flat bottom plates (Corning Incorporated). At the end of the 18 h incubation, biofilms

were washed twice in PBS to remove unattached cells, scraped from the 6 well plates using a

cell scraper, and sonicated in a water bath sonicator for 3 intervals of 30 seconds in 5 mL

polystyrene round-bottom tubes. Single cells resuspened in PBS were then serially diluted and

plated on either BHI, erythromycin-BHI, or kanamycin-BHI plates for selective plating. After 48

hours of incubation at 37°C in 5% CO_2, plates were removed, imaged, and viable colonies

enumerated from each plate. Fitness of each strain was then determined by taking the sum of

viable colonies from the erythromycin-BHI and kanamycin-BHI plates, dividing by the

erythromycin-BHI count and multiplying by 100 to determine a viable count percentage returned

of each strain from the grown biofilm.

Confocal laser scanning microscopy. Sonicated overnight cultures were washed and re-

suspended in FMC before being diluted 1:50 overall into fresh medium with the desired

sender:responder co-culture ratio. Diluted cell suspensions (350 µL) were inoculated into each

well of an 8-well µ-Slide (ibidi USA) chambered coverslip as well as a 6-well polystyrene flat

bottom plate (2.5 mL) to be used for flow cytometry analysis. Plates were incubated at 37°C in a

5% CO_2, aerobic atmosphere for a total of 18 h. At the 6-h time point, spent medium was

removed from the biofilms and fresh medium applied along with either sXIP or another strain, if

desired. Prior to analysis by microscopy, wells were washed 3 times with PBS and were kept

hydrated with 100 µL of PBS. Biofilm images were acquired using a spinning disk confocal

system connected to a Leica DMIRB inverted fluorescence microscope equipped with a

Photometrics cascade-cooled EMCCD camera. GFP fluorescence was detected by excitation at

488 nm and emission was collected using a 525 nm (±25 nm) bandpass filter. Detection of

dsRed fluorescence (RFP) was performed using a 642-nm excitation laser and a 695-nm (±53-

nm) bandpass filter. All z-sections were collected at 1 µm intervals using an 63X/1.40 oil

objective lens. Image acquisition and processing was performed using VoxCell (VisiTech

International).

Flow cytometry. Biofilms grown in 6-well flat bottom plates were scraped from the surface of

wells before being run through a FACSCalibur™ (BD Biosciences) flow cytometer. At the end of

the 18-h incubation, biofilms were washed twice in PBS to remove unattached cells, scraped

from the 6-well plates using a cell scraper, and sonicated in a water bath sonicator for 3

intervals of 30 seconds in 5 mL polystyrene round-bottom tubes to achieve primarily single cells

for analysis. Forward and side scatter signals were set stringently to allow sorting of single cells.

In total, 5×10^4 cells were counted from each event, at a maximum rate of 2×10^3 cells per

second, and each experiment was performed in triplicate. Detection of GFP fluorescence was

through a 530 nm (± 30 nm) bandpass filter, and dsRed was detected using a 670-nm long pass

filter. Data were acquired for unstained cells and single-color positive controls so that data

collection parameters could be properly set. The data were collected using Cell Quest Pro (BD

Biosciences) and analyzed with FCS Express 4 (De Novo Software). Gating for quadrant

analysis was selected by using a dot density plot with forward and side scatter, with gates set to

capture the densest section of the plot. Graphing and statistical analyses were performed using

Prism (GraphPad Software). *x-* and *y*-axis data represent logarithmic scales of fluorescent

intensity (arbitrary units).

Monitoring XIP-signaling using filtered supernates. Filtered culture supernates from selected

strains of *S. mutans* were produced by centrifugation of cultures diluted 20-fold from overnight

and grown to OD = 0.85-1.0. The resulting supernatant fluid was adjusted to pH 7.0 using 6.25

N NaOH and concentrated glucose was added to increase the amount of glucose in the

supernates by 20 mM. The resulting solution was then filtered using a 0.22 µm PVDF syringe

filter. P*comX*::*gfp* reporter cells with either UA159 or Δ*comS* background were diluted 20-fold

from an overnight culture, grown to OD_{600} of 0.1 and collected via centrifugation. The supernates

of the reporter cells were removed and replaced with the same volume of the pH- and glucose-

corrected supernates of the strains of interest. For sXIP controls, supernates were replaced with

fresh FMC, after which sXIP was added in the indicated concentrations. Two mL of the resulting

reporter cell-supernate combinations were then placed in a Falcon 24 well plate (Corning Inc.),

covered with mineral oil and the OD_{600} and green fluorescence (excitation 485nm, emission

528nm filter set) were measured in a BioTek Synergy 2[®] plate reader, shaking gently before

each reading to prevent cell settling.

ACKNOWLEDGEMENTS

Research reported in this publication was supported by the National Institute of Dental and

Craniofacial Research of the National Institutes of Health under Award Numbers R01

DE023339, R01 DE13239, T90 DE21990 and F31 DE024416.

REFERENCES

1. **Fuqua WC**, **Winans SC**, **Greenberg EP**. 1994. Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. J Bacteriol **176**:269–75.

2. **Kostakioti M**, **Hadjifrangiskou M**, **Hultgren SJ**. 2013. Bacterial biofilms: development, dispersal, and therapeutic strategies in the dawn of the postantibiotic era. Cold Spring Harb Perspect Med **3**:a010306.

3. **Boyer M**, **Wisniewski-Dyé F**. 2009. Cell-cell signalling in bacteria: not simply a matter of quorum. FEMS Microbiol Ecol **70**:1–19.

4. **Lyon GJ**, **Novick RP**. 2004. Peptide signaling in *Staphylococcus aureus* and other Gram-positive bacteria. Peptides **25**:1389–1403.

5. **Monnet V**, **Juillard V**, **Gardan R**. 2014. Peptide conversations in Gram-positive bacteria. Crit Rev Microbiol 1–13.

6. **Johnston C**, **Martin B**, **Fichant G**, **Polard P**, **Claverys J-P**. 2014. Bacterial transformation: distribution, shared mechanisms and divergent control. Nat Rev Microbiol **12**:181–96.

7. **Mashburn-Warren L**, **Morrison DA**, **Federle MJ**. 2010. A novel double-tryptophan peptide pheromone controls competence in *Streptococcus* spp. via an Rgg regulator. Mol Microbiol **78**:589–606.

8. **Håvarstein LS**. 2010. Increasing competence in the genus *Streptococcus*. Mol Microbiol **78**:541–4.

9. **Son M**, **Ahn S-J**, **Guo Q**, **Burne RA**, **Hagen SJ**. 2012. Microfluidic study of competence regulation in *Streptococcus mutans*: environmental inputs modulate bimodal and unimodal expression of *comX*. Mol Microbiol **86**:258–72.

10. **Lemos JA**, **Quivey RG**, **Koo H**, **Abranches J**. 2013. *Streptococcus mutans*: a new Gram-positive paradigm? Microbiology **159**:436–45.

11. **Hossain MS**, **Biswas I**. 2012. An extracelluar protease, SepM, generates functional competence-stimulating peptide in *Streptococcus mutans* UA159. J Bacteriol **194**:5886–96.

12. **Shanker E**, **Federle M**. 2017. Quorum sensing regulation of competence and bacteriocins in *Streptococcus pneumoniae* and *mutans*. Genes (Basel) **8**:15.

13. **Khan R**, **Rukke H V**, **Ricomini Filho AP**, **Fimland G**, **Arntzen MØ**, **Thiede B**, **Petersen FC**. 2012. Extracellular identification of a processed type II ComR/ComS pheromone of *Streptococcus mutans*. J Bacteriol **194**:3781–8.

14. **Desai K**, **Mashburn-Warren L**, **Federle MJ**, **Morrison DA**. 2012. Development of competence for genetic transformation of *Streptococcus mutans* in a chemically defined medium. J Bacteriol **194**:3774–80.

714   15.   **Wenderska IB**, **Lukenda N**, **Cordova M**, **Magarvey N**, **Cvitkovitch DG**, **Senadheera**
715   **DB**. 2012. A novel function for the competence inducing peptide, XIP, as a cell death
716   effector of *Streptococcus mutans*. FEMS Microbiol Lett **336**:104–12.
717
718   16.   **Gardan R**, **Besset C**, **Gitton C**, **Guillot A**, **Fontaine L**, **Hols P**, **Monnet V**. 2013.
719   Extracellular life cycle of ComS, the competence-stimulating peptide of *Streptococcus*
720   *thermophilus*. J Bacteriol **195**:1845–55.
721
722   17.   **Fontaine L**, **Goffin P**, **Dubout H**, **Delplace B**, **Baulard A**, **Lecat-Guillet N**, **Chambellon**
723   **E**, **Gardan R**, **Hols P**. 2013. Mechanism of competence activation by the ComRS
724   signalling system in streptococci. Mol Microbiol **87**:1113–32.
725
726   18.   **Fontaine L**, **Boutry C**, **de Frahan MH**, **Delplace B**, **Fremaux C**, **Horvath P**, **Boyaval P**,
727   **Hols P**. 2009. A novel pheromone quorum-sensing system controls the development of
728   natural competence in *Streptococcus thermophilus* and *Streptococcus salivarius*. J
729   Bacteriol **192**:1444–1454.
730
731   19.   **Cook LC**, **Federle MJ**. 2014. Peptide pheromone signaling in *Streptococcus* and
732   *Enterococcus*. FEMS Microbiol Rev **38**:473–92.
733
734   20.   **Fleuchot B**, **Gitton C**, **Guillot A**, **Vidic J**, **Nicolas P**, **Besset C**, **Fontaine L**, **Hols P**,
735   **Leblond-Bourget N**, **Monnet V**, **Gardan R**. 2011. Rgg proteins associated with
736   internalized small hydrophobic peptides: a new quorum-sensing mechanism in
737   streptococci. Mol Microbiol **80**:1102–19.
738
739   21.   **Shanker E**, **Morrison DA**, **Talagas A**, **Nessler S**, **Federle MJ**, **Prehna G**. 2016.
740   Pheromone recognition and selectivity by ComR proteins among *Streptococcus* species.
741   PLOS Pathog **12**:e1005979.
742
743   22.   **Talagas A**, **Fontaine L**, **Ledesma-Garca L**, **Mignolet J**, **Li de la Sierra-Gallay I**, **Lazar**
744   **N**, **Aumont-Nicaise M**, **Federle MJ**, **Prehna G**, **Hols P**, **Nessler S**. 2016. Structural
745   insights into streptococcal competence regulation by the cell-to-cell communication
746   system ComRS. PLOS Pathog **12**:e1005980.
747
748   23.   **Gardan R**, **Besset C**, **Guillot A**, **Gitton C**, **Monnet V**. 2009. The oligopeptide transport
749   system is essential for the development of natural competence in *Streptococcus*
750   *thermophilus* strain LMD-9. J Bacteriol **191**:4647–55.
751
752   24.   **Shields RC**, **Burne RA**. 2016. Growth of *Streptococcus mutans* in biofilms alters peptide
753   signaling at the sub-population Level. Front Microbiol **7**:1075.
754
755   25.   **Biswas I**, **Jha JK**, **Fromm N**. 2008. Shuttle expression plasmids for genetic studies in
756   *Streptococcus mutans*. Microbiology **154**:2275–82.
757
758   26.   **Ahn S-J**, **Kaspar J**, **Kim JN**, **Seaton K**, **Burne RA**. 2014. Discovery of novel peptides
759   regulating competence development in *Streptococcus mutans*. J Bacteriol **196**:3735–
760   3745.
761
762   27.   **Kaspar J**, **Ahn S-J**, **Palmer SR**, **Choi SC**, **Stanhope MJ**, **Burne RA**. 2015. A unique
763   ORF within the *comX* gene of *Streptococcus mutans* regulates genetic competence and
764   oxidative stress tolerance. Mol Microbiol **96**:463–482.

765   28.   **Guo Q**, **Ahn S-J**, **Kaspar J**, **Zhou X**, **Burne RA**. 2014. Growth phase and pH influence
766   peptide signaling for competence development in *Streptococcus mutans*. J Bacteriol
767   **196**:227–36.
768
769   29.   **Son M**, **Ghoreishi D**, **Ahn S-J**, **Burne RA**, **Hagen SJ**. 2015. Sharply tuned pH response
770   of genetic competence regulation in *Streptococcus mutans*: a microfluidic study of the
771   environmental sensitivity of *comX*. Appl Environ Microbiol **81**:5622–31.
772
773   30.   **Palmer SR**, **Miller JH**, **Abranches J**, **Zeng L**, **Lefebure T**, **Richards VP**, **Lemos JA**,
774   **Stanhope MJ**, **Burne RA**. 2013. Phenotypic heterogeneity of genomically-diverse
775   isolates of *Streptococcus mutans*. PLoS One **8**:e61358.
776
777   31.   **Takahashi N**, **Nyvad B**. 2011. The role of bacteria in the caries process: ecological
778   perspectives. J Dent Res **90**:294–303.
779
780   32.   **Bowen WH**, **Koo H**. 2011. Biology of *Streptococcus mutans* derived
781   glucosyltransferases: Role in extracellular matrix formation of cariogenic biofilms. Caries
782   Res **45**:69–86.
783
784   33.   **Koo H**, **Falsetta ML**, **Klein MI**. 2013. The exopolysaccharide matrix: a virulence
785   determinant of cariogenic biofilm. J Dent Res **92**:1065–73.
786
787   34.   **Zeng L**, **Burne RA**. 2015. Sucrose- and fructose-specific effects on the transcriptome of
788   *Streptococcus mutans* probed by RNA-Seq. Appl Environ Microbiol **82**:AEM.02681-15.
789
790   35.   **Moye ZD**, **Son M**, **Rosa-Alberty AE**, **Zeng L**, **Ahn S-J**, **Hagen SJ**, **Burne RA**. 2016.
791   Effects of carbohydrate source on genetic competence in *Streptococcus mutans*. Appl
792   Environ Microbiol AEM.01205-16.
793
794   36.   **Kim MK**, **Ingremeau F**, **Zhao A**, **Bassler BL**, **Stone HA**. 2016. Local and global
795   consequences of flow on bacterial quorum sensing. Nat Microbiol **1**:15005.
796
797   37.   **Chang JC**, **LaSarre B**, **Jimenez JC**, **Aggarwal C**, **Federle MJ**. 2011. Two group A
798   streptococcal peptide pheromones act through opposing Rgg regulators to control biofilm
799   development. PLoS Pathog **7**:e1002190.
800
801   38.   **Chang JC**, **Federle MJ**. 2016. PptAB exports Rgg quorum-sensing peptides in
802   *Streptococcus*. PLoS One **11**:e0168461.
803
804   39.   **Ahn S-J**, **Burne RA**. 2006. The *atlA* operon of *Streptococcus mutans*: role in autolysin
805   maturation and cell surface biogenesis. J Bacteriol **188**:6877–88.
806
807   40.   **Liao S**, **Klein MI**, **Heim KP**, **Fan Y**, **Bitoun JP**, **Ahn S-J**, **Burne RA**, **Koo H**, **Brady LJ**,
808   **Wen ZT**. 2014. *Streptococcus mutans* extracellular DNA is upregulated during growth in
809   biofilms, actively released via membrane vesicles, and influenced by components of the
810   protein secretion machinery. J Bacteriol **196**:2355–66.
811
812   41.   **Bayles KW**. 2007. The biological role of death and lysis in biofilm development. Nat Rev
813   Microbiol **5**:721–726.
814
815

816   42.   **Son M**, **Shields RC**, **Ahn S-J**, **Burne RA**, **Hagen SJ**. 2015. Bidirectional signaling in the
817   competence regulatory pathway of *Streptococcus mutans*. FEMS Microbiol Lett
818   **362**:fnv159.
819

820   43.   **Dufour D**, **Cordova M**, **Cvitkovitch DG**, **Lévesque CM**. 2011. Regulation of the
821   competence pathway as a novel role associated with a streptococcal bacteriocin. J
822   Bacteriol **193**:6552–9.
823

824   44.   **Kaspar J**, **Kim JN**, **Ahn S-J**, **Burne RA**. 2016. An essential role for (p)ppGpp in the
825   integration of stress tolerance, peptide signaling, and competence development in
826   *Streptococcus mutans*. Front Microbiol **7**:1162.
827

828   45.   **Seaton K**, **Ahn S-J**, **Sagstetter AM**, **Burne RA**. 2011. A transcriptional regulator and
829   ABC transporters link stress tolerance, (p)ppGpp, and genetic competence in
830   *Streptococcus mutans*. J Bacteriol **193**:862–74.
831

832   46.   **Marsh PD**. 2003. Are dental diseases examples of ecological catastrophes? Microbiology
833   **149**:279–294.
834

835   47.   **Terleckyj B**, **Shockman GD**. 1975. Amino acid requirements of *Streptococcus mutans*
836   and other oral streptococci. Infect Immun **11**:656–664.
837

838   48.   **Lau PC.**, **Sung CK**, **Lee JH**, **Morrison DA**, **Cvitkovitch DG**. 2002. PCR ligation
839   mutagenesis in transformable streptococci: application and efficiency. J Microbiol
840   Methods **49**:193–205.
841

842   49.   **Kwak IH**, **Son M**, **Hagen SJ**. 2012. Analysis of gene expression levels in individual
843   bacterial cells without image segmentation. Biochem Biophys Res Commun **421**:425–30.
844

845   50.   **LeBlanc DJ**, **Lee LN**, **Abu-Al-Jaibat A**. 1992. Molecular, genetic, and functional analysis
846   of the basic replicon of pVA380-1, a plasmid of oral streptococcal origin. Plasmid **28**:130–
847   145.
848

849   51.   **Liu Y**, **Zeng L**, **Burne RA**. 2009. AguR is required for induction of the *Streptococcus*
850   *mutans* agmatine deiminase system by low pH and agmatine. Appl Environ Microbiol
851   **75**:2629–37.
852

853 **TABLES**

854 TABLE 1. List of strains

Strain or plasmid	Relevant Characteristics*	Source or Reference
S. mutans Strains		
UA159	Wild-type	ATCC 700610
P*comX*::*gfp*/ UA159	UA159 harboring P*comX*::*gfp*	(9)
P*comX*::*gfp*/Δ*comS*	Δ*comS* harboring P*comX*::*gfp*	(9)
P*comX*::*gfp*/Δ*oppA*	Δ*oppA* harboring P*comX*::*gfp*	(9)
P*comX*::*gfp*/pIB184	UA159 harboring P*comX*::*gfp* and pIB184	(28)
P*comX*::*gfp*/pPMZ	UA159 harboring P*comX*::*gfp* and pPMZ	This study
pIB184comS/ UA159	UA159 harboring pIB184comS	(28)
P*comS*::*dsRed/*pIB184comS	UA159 harboring P*comS*::*dsRed* and pIB184comS	This study
P*comX*::*dsRed/*pIB184comS	UA159 harboring P*comX*::*dsRed* and pIB184comS	This study
Δ*atlA*	*atlA* (SMU.704c) :: NPKm[R]	(39)
pIB184comS / Δ*atlA*	Δ*atlA* harboring pIB184comS	This study
Plasmids		
pDL278	*E. coli - Streptococcus* shuttle vector, Sp[R]	(50)
pIB184	Shuttle expression plasmid with the constitutive P23 promoter, Em[R]	(25)
pPMZ	LacZ fusion integration vector based on pMC195 and pMC340B; Km[R]	(51)

855   * Em, erythromycin; Km, kanamycin; Sp, spectinomycin.

 FIGURE LEGENDS

 Figure 1. Model of co-culture system. (A) To monitor natural ComRS signaling, two strains of *S.*

 mutans are grown together in a co-culture system. The first strain, the "sender", contains a

 plasmid that allows for the overexpression of the XIP peptide precursor *comS* under the control

 of the constitutive P_{23} promoter. The sender strain also contains pDL278 carrying the gene for

 the dsRed fluorescent protein under the control of the *comX* promoter. The second strain, the

 "responder", harbors the P*comX*::*gfp* reporter plasmid on pDL278 which becomes activated

 when external XIP is imported into the responder via the oligopeptide permease, Opp. The

 empty pIB184 vector is also harbored in the responder strain to keep sender and responder

 strains as genetically similar as possible. (B) Relative GFP expression and OD_{600}

 measurements during co-culture growth of UA159 and P*comX*::*gfp*/UA159 (green; circles),

 pIB184comS/UA159 and P*comX*::*gfp*/UA159 (blue; triangles), pIB184comS/UA159 and

 P*comX*::*gfp*/Δ*comS* (orange; diamonds), and pIB184comS/UA159 and P*comX*::*gfp*/Δ*oppA* (red;

 squares). (C) Relative GFP expression measurements during co-culture of UA159 (sender) and

 P*comX*::*gfp*/UA159 (responder) using different concentrations of sXIP (legend) in comparison to

 growth between pIB184comS/UA159 and P*comX*::*gfp*/UA159 (black squares; co-culture). Each

 assay was performed with biological triplicates.

 Figure 2. XIP signaling is cell-cell contact independent. (A) Relative GFP expression and OD_{600}

 measurements of P*comX*::*gfp*/UA159 grown either in supernatants of either UA159 (green,

 circles), UA159 supplemented with 50 nM sXIP (blue, squares), or pIB184comS/UA159

 (orange, triangles). Supernatants were taken from overnight cultures of respective strains and

 filter sterilized, its pH was adjusted to 7.0, and glucose was added to the spent medium in an

 amount equivalent to a concentration of 20 mM. (B) Model of transwell assay using co-cultures

 of pIB184comS/UA159 and P*comX*::*gfp*/UA159. P*comX*::*gfp*/UA159 (green) strain is inoculated

 first in the bottom well, followed by placement of the 0.4 µM polycarbonate membrane insert.

The pIB184comS/UA159 (red) strain is then inoculated in the top well above the membrane. (C)

Relative GFP expression measurements from transwell experiments. Labeling of x-axis denotes

strain inoculated in the upper well first, followed up the strain inoculated in the lower well. Each

assay was performed with biological triplicates. Statistical analysis was performed by Student's

t-test. N.S. = not significant.

Figure 3. Measurements of XIP spatial diffusion. P*comX*::*gfp*/Δ*comS* cells were injected into an

IBIDI microslide and allowed to settle before an agarose/FMC mixture was pushed through the

slide to immobilize the cells. sXIP (1 µM) was injected into one end of the channel and green

fluorescence of individual cells was monitored at different distances from the site of XIP

injection. (A) 3D plot of average GFP fluorescence of cells in the channel (solid) overlaid with a

fit to a diffusive spreading model (mesh). (B) Phase/fluorescence overlaid images of

P*comX*::*gfp*/Δ*comS* at distances of 3, 4, 5 and 6 mm from the XIP injection site at 1 h after

injection; (C) 3 h after injection; (D) 5 h after injection.

Figure 4. Single-cell observations of ComRS signaling in co-cultures. Co-cultures of *comS*-

overexpressing sender (P*comX*:*dsRed*/pIB184*comS*) and either P*comX*::gfp/UA159 or

P*comX*::gfp/Δ*comS* responder in a low-melting-point agarose/FMC gel. After loading the gel into

a microfluidic channel, cells were imaged at hourly intervals by phase contrast and fluorescence

microscopy. (A) Fluorescence of P*comX*::*dsRed*/pIB184*comS* and P*comX*::gfp/UA159 co-

culture at 2 h (blue; triangles) and 3 h (red; circle) time points as a single cell scatter plot. (B)

P*comX*::*dsRed*/pIB184*comS* and P*comX*::gfp/UA159 co-culture at 2 h and 3 h with 1 µM sXIP

injected as control. (C) P*comX*::*dsRed*/pIB184*comS* and P*comX*::gfp/Δ*comS* co-culture at 2 h

and 3 h time points. (D) P*comS*::*dsRed*/pIB184*comS* and P*comX*::gfp/Δ*comS* co-culture at 2 h

and 3 h time points with 1 µM sXIP injected as control. (E-H) Representative images of (A-D) in

the same order.

Figure 5. ComRS signaling in biofilms. Observation of ComRS signaling in 18 h biofilms. (A)

Selected maximum intensity z-section confocal microcopy images of co-culture biofilms. Images

are of a 10 µm section of fluorescent range within the biofilm, collected at 1 µm intervals using a

63X/1.40 oil objective lens. Y- axis labeling of the panel denotes the order of biofilm inoculation

at time 0 h. (B) Quadrant analysis of collected flow cytometry data from similarly-grown co-

culture biofilms shown in A. Y- axis shows dsRed intensity and X- axis shows GFP intensity.

Quadrants are set to UA159 control. Flow cytometry data was collected from three independent

experiments with triplicate samples.

*Figure 6.Impact of sucrose on ComRS signaling in biofilms.*Observation of ComRS signaling in

18-h biofilms grown in different concentrations of sucrose. (A) Selected maximum intensity z-

section confocal microcopy images of co-culture biofilms. Images are a 10 µm section of

fluorescent range within the biofilm, collected at 1 µm intervals using an 63X/1.40 oil objective

lens. X-axis labeling of the panel denotes amount of carbohydrate source used in biofilm growth

medium (No sucrose = 20 mM glucose; low sucrose = 15 mM glucose and 2.5 mM sucrose;

medium sucrose = 10 mM glucose and 5 mM sucrose; high sucrose = 2 mM glucose and 9 mM

sucrose). As sucrose is a disaccharide, carbohydrate concentration (w/v) was the same in each

condition. (B) Quadrant analysis of collected flow cytometry data from similarly-grown co-

culture biofilms as shown in A. Y-axis shows dsRed intensity and X- axis shows GFP intensity.

Quadrants are set to UA159 control. Flow cytometry data was collected from three independent

experiments with triplicate samples.

Figure 7. Impact of cell lysis on ComRS Signaling. Fluorescence reporter activity (colored lines)

and growth curves (black lines) of responder strains grown in supernates of sender strains. (A)

GFP fluorescence of P*comX*::*gfp*/UA159 reporter cells grown in overnight supernates (filtrates)

of either pIB184/UA159 (green; squares), pIB184/Δ*atlA* (blue; circles), pIB184*comS*/UA159

(orange; upward triangles), or pIB184*comS*/Δ*atlA* (red; downward triangles). Overnight

supernates were pH-corrected to 7.0 using 6.25N NaOH and glucose was replenished to a

concentration equivalent to an additional 20 mM before filtering the supernatant fluids through a

0.22 μM PVDF syringe. (B) GFP fluorescence of P*comX*::*gfp*/Δ*comS* reporter cells grown in

overnight supernates (filtrates) as in A. Each assay was performed with biological triplicates.

A

Senders

P$_{23}$

comS

pIB184

P*comX*

dsRed

pDL278

XIP — Opp

Opp

P$_{23}$

--

pIB184

P*comX*

gfp

pDL278

Responders

B

Relative Expression (RFU/OD$_{600}$) vs Time (Hours), with Optical Density 600 nm on right axis

- UA159;P*comX*::*gfp*
- 184*comS*;P*comX*::*gfp*
- 184*comS*;P*comX*::*gfp*Δ*comS*
- 184*comS*;P*comX*::*gfp*Δ*oppA*

C

Relative Expression (RFU/OD$_{600}$) vs Time (Hours)

- 184*comS*;P*comX*::*gfp*
- 50 nM sXIP
- 37.5 nM sXIP
- 25 nM sXIP
- 12.5 nM sXIP
- 5 nM sXIP
- 1 nM sXIP

A

| sXIP Control | Same Time |
| Sender First | Responder First |

B

UA159 Control | sXIP Control | Same Time

Sender First | Responder First

Senders (P*comX::dsRed*)

Responders (P*comX::gfp*)

UA159 Control: 0.3 ± 0, 0 ± 0, 99 ± 0, 0.6 ± 0

sXIP Control: 18 ± 3, 2 ± 0.6, 15 ± 1, 65 ± 3

Same Time: 45 ± 4, 1 ± 0.4, 42 ± 3, 12 ± 2

Sender First: 61 ± 6, 9 ± 1, 19 ± 2, 11 ± 4

Responder First: 3 ± 1, 0 ± 0.6, 94 ± 3, 3 ± 1