Some reasons why latent period should not be considered constant over the course of a plant disease epidemic

Frédéric Suffert

UMR BIOGER, INRA, AgroParisTech, Université Paris-Saclay, 78850 Thiverval-Grignon, France

✉ frederic.suffert@inra.fr

Abstract – The latent period is a crucial life history trait, particularly for polycyclic plant diseases, because it determines how many complete monocycles could theoretically occur during an epidemic. Many studies have focused on the variation of latent period with pathogen or host genotype, or its plasticity in response to environmental factors. The focus on these aspects is unsurprising, as these factors classically form the apices of the epidemiological triangle. Experiments in controlled conditions are generally used to assess pathogenicity and host susceptibility. Most epidemiological studies consider implicitly that the latent period of a plant pathogen population is constant in time. Once estimated for one or several pairs of host-pathogen genotypes, the value of the parameter 'latent period' is usually considered fixed and is often used "as is" in epidemiological models. My thesis here is that the latent period can display non-negligible variability over the course of a disease epidemic, and that this variability has multiple sources, some of which have complex, antagonistic impacts. I develop arguments for four sources of variation challenging the implicit assumption that the latent period remains constant: daily fluctuations in leaf temperature, nature of inoculum, host stage or age of host tissues, intra-population competition and selection for aggressiveness traits. I focus on the wheat fungal pathogen Septoria tritici blotch (*Zymoseptoria tritici*), making use of empirical datasets collected during my own research projects and a targeted literature review. Finally I discuss in which cases certain sources of these variations should be accounted for into epidemiological models.

Keywords – latent period, plant disease epidemiology, *Zymoseptoria tritici*

INTRODUCTION

In plant pathogens latent period is defined as "the length of time between the start of the infection process by a unit of inoculum and the start of production of infectious units" (Madden *et al.*, 2007). It determines the interval between successive generations of pathogens, analogously to age of reproductive maturity in nonparasitic organism, even if this view is an over-simplification in the case of pathogens for which disease cycles overlap. The importance of the latent period for the understanding and prediction of pathogen development has long been recognized in plant disease epidemiology (Vanderplank, 1963; Zadoks, 1972). It is a crucial life history trait and component of aggressiveness (Lannou, 2012), especially for polycyclic diseases, because it determines how many complete monocycles could theoretically occur during an epidemic and the final intensity of the epidemic. Of course several other monocyclic parameters (infection efficiency, infectious period, sporulation intensity) are of great importance, but my objective is not here to assess and discuss their relative impact on epidemics.

Many plant pathology studies have investigated variations of the latent period with pathogen or host genotype, or its plasticity in response to environmental factors, such as temperature and humidity (Davis & Fitt, 1994; Shaw, 1990; Tomerlin & Jones, 1983; Webb & Nutter, 1997). Such approaches are relevant, because these factors lie at the apices of the epidemiological triangle (host, pathogen, environment; Zadoks, 1972). Most of empirical studies focused on the mean value of the latent period with statistical features such as standard error (related to the definition of latent period that is used; see below), in order to reduce the uncertainty of the measure, but rarely on the intrinsic variance (i.e. not due to a bias of measurement). There is very little information about the expected magnitude and between-individual temporal variability of the latent period in a fungal plant pathogen population. Such a variability was detected over a longer period in rare cases, for example in poplar rust: the latent period of five *Melampsora larici-populina* pathotypes collected in 1998 on Beaupré, a largely cultivated poplar cultivar, was 7 days, compared to 8 days required by the older isolates of the same pathotype (Pinon & Frey, 2005). Focusing on soilborne plant pathogens, Leclerc *et al.* (2014) similarly noticed that there is little information about how the incubation period (the time between host infection and the expression of first disease symptom, so shorter than latent period) varies within host populations. Finally, no empirical

study has ever considered the possibility that the latent period of a pathogen population may display variability over the course of a single annual epidemic.

In compartmental models SIR and SEIR (Kermack & McKendrick, 1927; van der Plank, 1963; Madden *et al.*, 2007) used for simulating polycyclic plant disease epidemics, the latent period is usually considered to be constant (e.g. *Plasmopara viticola* [Rossi *et al.*, 2009], *Phytophthora infestans* [Andrade-Piedra *et al.*, 2005], *Puccinia arachidis* [Savary *et al.*, 1990]); sometimes it is considered to be specific to an interaction between a pathogen genotype and a host genotype (e.g. Viljanen-Rollinson *et al.* 2005). The simplest case is the SIR model, in which the compartments represent susceptible, infectious and removed individuals, while the SEIR model also includes an exposed compartment, containing hosts who are in a latent stage (infected but not yet infectious). Several standard SEIR formulations in botanical, but also veterinary and medical epidemiology, assume that the stage durations (including infectious and latent periods) are exponentially or gamma distributed. However, real distributions are narrower and centred around the mean, and one can imagine that these distributions change over time, for instance over the course of a single annual epidemic. Effects of the infectious and/or latent period distribution on prediction concerning epidemiological transitions were rarely investigated: for example in childhood disease dynamics by Krylova & Earn (2013), in plant disease dynamics by Parry *et al.* (2014) who includes seasonal and host age effects in their model of the latent period of Huanglongbing, and by Cunniffe *et al.* (2012) who considers flexible distributions of latent periods in generic SEIR models and demonstrates how mis-specified or time-varying latent periods can lead to incorrect assessment of final intensity of an epidemic.

Several experiments are performed in controlled conditions to assess the pathogenicity of the pathogen (separated into a qualitative term, virulence, and a quantitative term, aggressiveness; Lannou, 2012) or the susceptibility of the host. Once estimated for one or several pairs of host-pathogen genotypes, one can be tempted to fix the value of the parameter 'latent period' and to use it into the model "as is" (except for temperature, which is often taken into account - imperfectly, as explained below - through the use of a thermal time scale). This approach is entirely appropriate if there is no great heterogeneity in the pathogen population, the host population or the environmental conditions during an epidemic. However, in practice, the pathogenicity of the active part of a pathogen population (e.g. the strains responsible for secondary infections in the case of a polycyclic epidemic), the susceptibility of a host population (e.g. landraces or varietal mixtures), and the environmental conditions (e.g. diurnal or seasonal temperature variations) may vary in time in significant proportions. The values taken by the parameters 'pathogenicity', 'susceptibility', and 'temperature' may therefore be heterogeneous locally at a given time step. In such conditions, one may wonder if the variability of the latent period, as important as its mean value, should not be better estimated and taken into account in models for which it is important. Here I would like to convince the reader that the short-term variability of the latent period has numerous sources, some of which are antagonistic, and is sometimes not negligible. To this end I have identified four sources of variability that challenge the implicit assumption that the mean latent period of a local pathogen population remains constant over the course of an epidemic. To present this view, I focus on the wheat pathogen Septoria tritici blotch (*Zymoseptoria tritici*), making use of empirical datasets collected during my own research projects and a targeted literature review. This fungal disease is particularly suitable for this analysis because the effects of several factors have been highlighted and are now well documented. *Z. tritici* is a cyclic heterothallic pathogen reproducing both sexually and asexually, resulting in infections initiated by two types of spores (ascospores and pycnidiospores), with relative contributions to the epidemic that change over the course of the year (Suffert *et al.*, 2011). The pathogen population displays a high degree of genetic diversity (Linde *et al.*, 2002) and there may be considerable phenotypic variability in latent period between strains (Morais *et al.*, 2015; 2016). Wheat has a long growth cycle and infections occur from early fall to late spring, under the influence of heterogeneous environmental selective pressures driven by abiotic conditions such as temperature (Lovell *et al.*, 2004a), but also biotic conditions such as the physiological stage of wheat or its fertilization regime (Robert *et al.*, 2005; Précigout *et al.*, 2017). As Septoria tritici blotch epidemics are polycyclic and results from an integration of many overlapping infection cycles, the latent period is a crucial fitness trait. It is quite long, facilitating the detection and quantification of any differences by *in planta* experiments, and it may display signs of local adaptation to climatic conditions (Suffert *et al.*, 2015).

Several operational definitions of latent period can be used. For example, the latent period for Septoria

tritici blotch is usually estimated at the scale of a lesion, as the time between inoculation and the appearance of the first pycnidium (Shearer & Zadoks, 1972; Armour *et al.*, 2004) or, for the sake of convenience, 5% of the final number of pycnidia or 5% of the maximum percentage of area covered by pycnidia (Suffert *et al.*, 2013). When several lesions rather than a single lesion are considered, particularly if methodological constraints make it necessary (e.g. impossibility of replicating individual inoculation with a given *Z. tritici* genotype using the ascosporic form, contrary to the conidial form; Morais *et al.*, 2015), the latent period is estimated at the scale of a leaf, as the time between inoculation and the appearance of half the final sporulating lesions (Shaw, 1990; Lovell *et al.*, 2004a).

THE LATENT PERIOD CAN VARY WITH FLUCTUATIONS IN LEAF TEMPERATURE DESPITE AN IDENTICAL DAILY MEAN AIR TEMPERATURE

The development of plant pathogens responds strongly to the temperature of the surrounding environment. The effects of temperature are so well recognized in plant epidemiology that linear thermal time (referring to the accumulation of degrees above a given base temperature over a specified period of time; Lovell *et al.*, 2004b) is widely preferred over physical time for assessments and the modeling of disease development, particularly for Septoria tritici blotch. Consequently, the latent period is usually expressed in degree days rather than as a number of physical days. This accounts, for example, for the decrease in the latent period of *Z. tritici* estimated as a number of days over the spring epidemic period: a 350 degree-day latent period (with a base temperature of -2.4°C; Lovell *et al.*, 2004a) typically corresponds in average to 33 days in early spring (April) but only 22 days in late spring (June) in France (average monthly temperature in Poissy, Yvelines; https://en.climate-data.org). Taking into account the impact of temperature by this way is however not completely adequate because relationships between temperature and the efficiency or duration of a given epidemiological process are actually not linear and not monotonous. Consequently the latent period, while assessed using thermal time, should not be considered constant in time, particularly if the time step used for the calculation is large (e.g. daily), for at least two reasons.

Firstly, thermal time is usually calculated from air temperature, whereas the development of foliar fungal pathogens, including *Z. tritici*, reacts more directly to leaf temperature (the temperature actually perceived by the fungus), which can be very different from air temperature (Bernard *et al.*, 2013). Leaf temperature is harder to measure than air temperature, but it could be estimated indirectly estimated from soil-vegetation-atmosphere transfer (SVAT) models including data recorded at standard weather stations (Xiao *et al.*, 2006).

Secondly, the latent period is usually assessed under fluctuating temperature regimes, with a thermal scale based on the accumulation of daily mean temperatures. The effects of diurnal fluctuations are, therefore, not taken into account. Bernard (2012) established the impact of two patterns of leaf temperature variation, in which mean temperature was equal (18°C) but daily temperature range differed (±2°C and ±5°C), on the latent period of *Z. tritici*: the larger temperature range increased the latent period by 1.3 days on average. Similar results have been obtained for other plant pathogens (Scherm *et al.*, 1994; Xu, 1999). The differences in pathogen development between constant and fluctuating environments are partly due to 'rate summation' or the Kaufmann effect (Ruel & Ayres, 1999), a mathematical consequence of the nonlinear shape of thermal performance curves (TPCs). The length of the latent period under a fluctuating temperature regime can be predicted with the rate summation scheme (Hau *et al.*, 1985):

$$S = \sum_{i=1,Y} [R(T)\Delta t]$$

where Δt is the integration step (e.g. 1 hour); S, the accumulated development, is dimensionless and defined as zero initially and one at the completion of a process (i.e. appearance of the first pycnidia), and $R(T)$ is the rate function of this process.

Finally, degree-hours should be preferred over degree-days once the TPC of the latent period is available.

The mean TPC of the latent period for *Z. tritici* was established empirically, with a limited number of fungal isolates, in natural (Shaw, 1990) or controlled conditions (Bernard *et al.*, 2012). The variability of the latent period between pathogen populations of different geographic origins has never before been characterized in detail. The latent period TPCs presented in Fig. 1a were obtained from two groups of nine *Z. tritici* samples collected from two regions of France with different climates (Brittany and Burgundy). The thermal optimum differed between the two populations: <20°C for the isolates from Brittany, >21°C for the isolates from Burgundy. The

Figure 1 – Illustration of four sources of variability of the latent period (expressed in days post-inoculation [dpi] or in degree-days post-inoculation [ddpi] with a base temperature of 0°C) in the wheat pathogen *Zymoseptoria tritici* over the course of an annual epidemic. Latent period was calculated in cases 1a, 1c and 1d as the time between inoculation and the appearance of 5% of the maximum percentage of area covered by pycnidia, calculated by fitting a Gompertz growth curve to experimental data as described in Suffert *et al.* (2013), and in case 1b as the time between inoculation and the appearance of 5% of the maximum number of pycnidia in each individual lesion as described in Morais *et al.* (2015).

1a. Effect of daily mean wheat leaf temperature on the latent period of two *Z. tritici* populations (2 × 9 isolates collected from cv. Apache in two French regions; black diamond = Dijon in Burgundy; white diamond = Ploudaniel in Brittany) assessed after pycnidiospore inoculation on wheat adult plants cv. Apache. The thermal performance curve (order 2 polynomial) was adjusted with six replicates per temperature.

1b. Relationship between the latent period of 12 *Z. tritici* isolates assessed after ascospore and pycnidiospore inoculation on wheat adult plants cv. Apache (from Morais *et al.*, 2015). Each point corresponds to the mean of several values for pycnidiospore inoculation (vertical bars represent the standard deviation) and a single value for ascospore inoculation.

1c. Relationship between the latent periods of *Z. tritici* populations (2 × 9 isolates collected from cv. Apache in two French regions; colored diamond = Dijon in Burgundy; white diamond = Ploudaniel in Brittany) assessed after pycnidiospore inoculation on wheat seedlings and wheat adult plants cv. Apache for five different wheat cultivars: Apache (green), common to both Brittany and Burgundy; Altamira (red) and Paledor (yellow), mostly cultivated in Brittany; Arezzo (blue) and Altigo (brown), mostly cultivated in Burgundy. Each point is the mean value for assessments on six replicates.

1d. Mean latent period of two *Z. tritici* subpopulations (2 × 15 isolates collected on seedlings cv. Soissons very early [Pi] in the epidemic and the upper leaf layers at the end of the same epidemic [Pf]), assessed under winter (on wheat seedlings cv. Soissons at 8.9°C) and spring (on adult plants cv. Soissons at 18.1°C) conditions (from Suffert *et al.*, 2015). Asterisks indicate that the mean differs significantly (P < 0.1) between Pi and Pf.

effect of temperature on latent period can differ between pathogen populations expressing local patterns of climatic adaptation.

Several published models of Septoria tritici blotch development consider the latent period to be proportional to temperature: 330 dd in Robert *et al.* (2008); 250, 350, 450, and 550 dd in Eriksen *et al.* (2001); 256 dd in Lovell *et al.* (2004a); 310 dd in Baccar *et al.* (2011). The mean TPC of the latent period is thus considered to be a decreasing linear function, with base temperature estimated at -2.4°C (Lovell *et al.*, 2014a) or 0°C (Armour *et al.*, 2004). The assumption of linearity for the suboptimal part of the TPC (from 10°C to 18°C) is acceptable, but not taking the supraoptimal part of the temporal TPC correctly into account can have deleterious consequences, particularly in fluctuating thermal regimes

THE LATENT PERIOD IS AFFECTED BY THE FORM OF INOCULUM: ASCOSPORES VS. PYCNIDIOSPORES

Models of Septoria tritici blotch development considering both ascospores and pycnidiospores in an explicit manner, either throughout the cropping season (Eriksen *et al.*, 2001) or solely at the onset of the epidemic (Robert *et al.*, 2008), assume that the infection process after spore deposition is the same for both types of spore. Morais *et al.* (2015) showed however that the latent period of *Z. tritici* was significantly longer (about 60 degree-days, i.e. 3-4 days in late spring) after infection with ascospores than after infection with pycnidiospores (Fig. 1b). This empirical result is consistent with results obtained for other plant pathogens in studies considering the efficiency of different types of spore without specifically focusing on the latent period (Gilles *et al.*, 2001; Karolewski *et al.*, 2002; Li *et al.*, 2004). In *Z. tritici*, one concrete consequence of this difference is that the mean latent period early in the epidemic, when first lesions are caused by wind-dispersed ascospores, is intrinsically longer than that during the spring epidemic stage, when infections are caused mostly by splash-dispersed pycnidiospores (Suffert *et al.*, 2011). A hypothetical, theoretical distribution of the number of new lesions induced by a pathogen population consisting of different *Z. tritici* strains according to their latent periods results in the superimposition of two unimodal distributions centered around the mean latent period value of each type of spore (Fig. 2). The resulting distribution may or may not be bimodal, depending on the relative contributions of the two types of spore to the infection. This example reveals another reason for which the latent period of

a plant pathogen with both sexual and asexual reproduction modes should not be considered constant over the course of an epidemic.

THE LATENT PERIOD DEPENDS ON HOST STAGE AND HOST TISSUE AGE

An increase in the latent period with increasing host development is classically observed for several plant pathogens, such as *Puccinia hordei* (Parlevliet, 1975) and *Puccinia striiformis* (Tomerlin *et al.*, 1983). This finding is consistent with the lack of univocal relationship between seedling and adult plant resistance, in wheat rusts for example, because many resistance genes are expressed in adult plants but not in seedlings (McIntoch *et al.*, 1995). I assessed the latent periods of two groups of *Z. tritici* isolates collected in two climatically different regions of France (Brittany and Burgundy), on both seedlings and adult plants. I found a large difference between plants of different ages, with a mean latent period of 301 dd for seedlings and 534 dd for adult plants (Fig. 1c). Moreover, other experimental studies have suggested that the susceptibility of wheat tissues varies with leaf layer for synchronous measurements (i.e. on the same date) on adult plants, probably due to differences in leaf age (interactions between the susceptibility of host tissues, natural senescence and nitrogen status; Ben Slimane *et al.*, 2012; Bernard *et al.*, 2013; Suffert *et al.*, 2015). The increase in latent period with developmental stage (seedling vs. adult plant), and, more generally, with leaf age (time between leaf emergence and leaf infection), has been investigated in detail for *Puccinia arachidis* (Savary, 1987). These findings provide further support for the contention that the latent period of a plant pathogen should not be considered constant over the course of an epidemic.

THE LATENT PERIOD IS STRAIN-DEPENDENT AND, THEREFORE, AFFECTED BY COMPETITION WITHIN A LOCAL PATHOGEN POPULATION

As mentioned above, latent period depends on pathogen genotype. Variability within a local pathogen population may be high or low, according to the intrinsic structure of the population (sexual reproduction leading to high levels of variability or clonality). Locally, at the scale of a single annual epidemic, some authors consider average aggressiveness, and, thus, latent period, to be stable (for a given type of spore). This assumption is usually implicit, but might be erroneous. I showed that the mean latent period of *Z. tritici* pycnidiospores can vary significantly during a single annual epidemic (Suffert *et al.*, 2015): isolates

Figure 2 – Hypothetical, theoretical distribution of the number of new lesions (on wheat plants, per m^2 and per week) induced by a pathogen population consisting of different *Z. tritici* strains according to their latent period, taking into account the nature of the spores and the epidemic stage (a = early stage of the epidemic in December; b = intermediate stage of the epidemic in April, c = late stage of the epidemic in June).

Red dotted lines correspond to ascospore-initiated lesions; blue dotted lines correspond to pycnidiospore-initiated lesions; black lines are the cumulative curves. Curves were built with the hypothesis of the mean latent period is 505 ddpi for pycnidiospore infection and 557 ddpi for ascospore infection, based on the results obtained by Morais *et al.* (2015) and re-used in Fig.1b. Both latent periods have here a gamma distribution, with a similar range of variance for ascospores and pycnidiospores. The relative height of the curves does not rely on actual experimental values, which do not exist, and is to be considered as an order of magnitude. This order of magnitude is inspired by the relative importance of the two types of inoculum from early (a) to intermediate epidemic stage (b) estimated using data of inoculum availability (Suffert & Sache, 2011; Morais *et al.*, 2015) and data of simulated origin of inoculum causing lesions from intermediate (b) to late (c) epidemic stage (Duvivier, 2015). The effect of host stage (Fig. 1c), which would likely shorten latent period at the early stage of the epidemic (a), was here not taken into account.

collected on the upper leaf layers of wheat at the end of epidemic have a shorter latent period than those collected from seedlings very early in the same epidemic. This difference in latent period between disease expressed under spring conditions (adult plants, warm temperature) and under winter conditions (seedlings, cold temperature) and suggested that strains with a shorter latent period are selected during the second part of the epidemic (spring), when the disease is propagated by the upward splash dispersal of spores (Fig. 1d). During this period, a short latent period is a key fitness trait conferring a real competitive advantage. These conclusions were corroborated by the observed decrease in between-genotype variance for the latent period. The decrease in the mean latent period of a pathogen population over the course of the epidemic is consistent with the increase in other aggressiveness traits recorded in various fungal pathogens after a few cycles of asexual reproduction (Newton & McGurk, 1991; Villaréal & Lannou, 2000; Andrivon *et al.*, 2007; Le May *et al.*, 2012). Once again, these findings indicate that the latent period of a plant pathogen should not be considered constant over the course of an epidemic.

DISCUSSION

This dissertation asserts that the mean latent period of the active part of a local pathogen population should not be implicitly considered constant over the course of the annual plant disease epidemic. Mean latent period can vary in time, and the sources of variation are numerous: daily fluctuations in leaf temperature, nature of the inoculum, host stage or age of host tissues, intra-population competition and selection for aggressiveness traits. Some of these sources of variation may have complex, antagonistic impacts.

On the one hand, mean latent period may decrease during the epidemic period because of intra-population competition driven by external factors, for instance temperature in the case of *Z. tritici*. On the other hand, it may decrease during the same period with changes in the ratio of the two spore types (pycnidiospores > ascospores), and finally increase as the ratio inverses (ascospores > pycnidiospores) during the late epidemic stage due to the occurrence of sexual reproduction before the end of the growing season. Shaw (1990) suggested that the increase in latent period he observed at high mean temperatures reflects the adaptation of *Z.*

tritici to local climatic conditions, such as the cool summers in the UK, and a physiological trade-off between an ability to grow rapidly at high temperatures and an ability to grow rapidly at low temperatures. This hypothesis is consistent with the conclusion of Suffert *et al.* (2015) that seasonal changes can drive short-term selection for fitness traits, recently confirmed by Boixel *et al.* (unpubl. data). However, Shaw's results were obtained in field conditions, and might therefore also be due to an artifact related to host stage effects (latent period shorter on seedlings than on adult plants), the use of air temperature rather than leaf temperature (Bernard *et al.*, 2013), and a greater amplitude of daily fluctuations during spring than during winter (Bernard, 2012). I suspect there are several causal factors involved, as is often the case in plant disease epidemiology.

The key message is that the mean latent period of a plant pathogen population can vary locally, in the short term. The direction (decrease or increase) and causes of this variation are difficult to determine, but plant pathologists need to be aware that the latent period is not necessarily a fixed parameter and that they may have an interest in incorporating, when necessary, a certain variability into epidemiological models. This variability should not be only described by a statistical distribution (e.g. exponential or gamma distribution) but also be structured by epidemiological processes as illustrated by the four sources of variation described above. The main difficulty is to know exactly in which case these sources of variation should be accounted for in a modelling work. Modelling is about simplification made according to the objective of the model and of hypotheses made. The detail of the parameterization of a model cannot take into account too many unconstrained parameters. Sources of short term variability in latent period should be analyzed and potentially incorporated into three type of epidemiological models at least: in the case of *Z. tritici*, forecasting models used by growers to determine the proper timing for effective fungicide sprays (e.g. Audsley *et al.*, 2005; El-Jarroudi *et al.*, 2009), mechanistic models used as research tools for testing the impact of different epidemiological parameters on disease development (e.g. Eriksen *et al.*, 2017; Baccar *et al.*, 2011), and evolutionary models in which latent period is under selective pressure (Suffert *et al.*, 2015), is involved in a trade-off (e.g. Heraudet *et al.*, 2008; Pariaud *et al.*, 2012; Suffert *et al.*, 2017), or is expected to tend towards an evolutionary optimum (e.g. Précigout *et al.*, 2017). To justify this, I have found only two examples in botanical epidemiology. In the first

example, Cunniffe *et al.* (2012) proposed an extension to the generic SEIR model, splitting the latent and infection compartments and thereby allowing time-varying infection rates and more realistic distributions of latent and infectious periods to be represented. Their results demonstrated that extending a model that has such a simplistic representation of the infection dynamics may not, in fact, lead to more accurate results, and they suggested that modelers should carefully consider the underlying assumptions of the simplest compartmental models. In the second example, Leclerc *et al.* (2014) conducted experiments on the soilborne pathogenic fungus *Rhizoctonia solani* in sugar beet and used spatially-explicit models to estimate the incubation period distribution. They demonstrate the impact of differing distributional estimations about the incubation period (i.e. exponential vs. gamma) on the epidemiological understanding of the disease development. These two examples show that including additional biological realism in a model about the distribution or the variation in latent period can be more or less critical in assessing the disease development at the scale of a single annual epidemic.

The aim of this article was to highlight a point that had not been discussed until now: the latent period is not necessarily constant in the short term. I hope this consideration will allow to justify the addition of relevant empirical knowledge into epidemiological models, since their precision or their robustness could be improved.

REFERENCES

Andrade-Piedra, J., Hijmans, R.J., Forbes, G.A., Fry, W.E., and Nelson, R.J. 2005. Simulation of potato late blight in the Andes: I - Modification and parameterization of the LATEBLIGHT model. Phytopathology 95:1191-1199.

Andrivon, D., Pilet, F., Montarry, J., Hafidi, M., Corbière, R., Achbani, E., Pelle, R., and Ellissèche, D. 2007. Adaptation of *Phytophthora infestans* to partial resistance in potato: evidence from French and Moroccan populations. Phytopathology 97:338-343.

Armour, T., Viljanen-Rollinson, S.L.H., Chng, S., Butler, R.C., Jamieson, P.D., and Zyskowski, R.F. 2004. Examining the latent period of septoria tritici blotch in a field trial of winter wheat. New Zeal. Plant Protect. 57:116-120.

Audsley, E., Milne, A., and Paveley, N. 2005. A foliar disease model for use in wheat disease management decision support systems. Ann. Appl. Biol. 147:161-172.

Baccar, R., Fournier, C., Dornbusch, T., Andrieu, B., Gouache, D., and Robert, C. 2011. Modelling the

effect of wheat canopy architecture as affected by sowing density on *Septoria tritici* epidemics using a coupled epidemic-virtual plant model. Ann. Bot. 108:1179-1194.

Ben Slimane, R., Bancal, P., Suffert, F., and Bancal, M.O. 2011. Localized septoria leaf blotch lesions in winter wheat flag leaf do not accelerate apical senescence during necrotrophic stage. J. Plant Pathol. 94:543-553.

Bernard, F. 2012. The development of a foliar fungal pathogen does react to temperature, but to which temperature? PhD thesis, AgroParisTech, France.

Bernard, F., Sache, I., Suffert, F., and Chelle, M. 2013. The development of a foliar fungal pathogen does react to leaf temperature! New Phytol. 198:232-240.

Cunniffe, N.J., Stutt, R.O.J.H., van den Bosch, F., and Gilligan, C.A. 2012. Time-dependent infectivity and flexible latent and infectious periods in compartmental models of plant disease. Phytopathology 102:365-380.

Davis, H., and Fitt, B.D.L. 1994. Effects of temperature and leaf wetness on the latent period of *Rhynchosporium secalis* (leaf blotch) on leaves of winter barley. J. Phytopath. 140:269-279.

Duvivier, M. 2015. Distribution of the airborne inoculum of wheat leaf rust and septoria tritici blotch. PhD thesis, Université Catholique de Louvain, Belgique.

El Jarroudi, M., Delfosse, P., Maraite, H., Hoffmann, L., and Tychon, B. 2009. Assessing the accuracy of simulation model for Septoria leaf blotch disease progress on winter wheat. Plant Dis. 93:983-992.

Eriksen, L.M.W., Shaw, M.W., and Ostergard, H. 2001. A model of the effect of pseudothecia on genetic recombination and epidemic development in populations of *Mycosphaerella graminicola*. Phytopathology 91:240-248.

Gilles, T., Fitt, B.D.L., McCartney, H.A., Papastamati, K., and Steed, J.M. 2001. The roles of ascospores and conidia of *Pyrenopeziza brassicae* in light leaf spot epidemics on winter oilseed rape (*Brassica napus*) in the UK. Ann. Appl. Biol. 138:141-152.

Hau, B., Eisensmith, S.P., and Kranz, J. 1985. Construction of temporal models II: Simulation of aerial epidemics. In: Gilligan, C.A. Advances in Plant Pathology, Vol. 3: Mathematical Modelling of Crop Disease. Academic Press, London.

Heraudet, V, Salvaudon, L, Shykoff, J.A. 2008. Trade-off between latent period and transmission success of a plant pathogen revealed by phenotypic correlations. Evol. Ecol. Res. 10:913-924.

Karolewski, Z., Evans, N., Fitt, B.D.L., Todd, A.D., and Baierl, A. 2002. Sporulation of *Pyrenopeziza brassicae* (light leaf spot) on oilseed rape (*Brassica napus*) leaves inoculated with ascospores or conidia at different temperatures and wetness durations. Plant Pathol. 51:654-665.

Kermack, W.O., and McKendrick, A.G. 1927. A contribution to the mathematical theory of epidemics. Proc. R. Soc. London A 115:700-721.

Krylova, O., and Earn, D.J.D. 2013. Effects of the infectious period distribution on predicted transitions in childhood disease dynamics. J. R. Soc. Interface 10:20130098.

Lannou, C. 2012. Variation and selection of quantitative traits in plant pathogens. Ann. Rev. Phytopath. 50:319-338.

Le May, C., Guibert M., Leclerc, A., Andrivon, and D., Tivoli, B. 2012. A single, plastic population of *Mycosphaerella pinodes* causes ascochyta blight on winter and spring peas (*Pisum sativum*) in France. Appl. Environ. Microb. 78:8431-8440.

Leclerc, M., Doré, T., Gilligan, C.A., Lucas, P., and Filipe, J.A.N. 2014. Estimating the delay between host infection and disease (incubation period) and assessing its significance to the epidemiology of plant diseases. PLoS One 9:e86568.

Li, H., Sivasithamparam, K., Barbetti, M.J., and Kuo, J. 2004. Germination and invasion by ascospores and pycnidiospores of *Leptosphaeria maculans* on spring-type *Brassica napus* canola varieties with varying susceptibility to blackleg. J. Gen. Plant Pathol. 70:261-269.

Linde, C.C., Zhan, J., and McDonald, B.A. 2002. Population structure of *Mycosphaerella graminicola*: from lesions to continents. Phytopathology 92:946-955.

Lovell, D.J., Hunter, T., Powers, S.J., Parker, S.R., and van den Bosch, F. 2004a. Effect of temperature on latent period of septoria leaf blotch on winter wheat under outdoor conditions. Plant Pathol. 53:170-181.

Lovell, D.J., Powers, S.J., Welham, S.J., and Parker S.R. 2004b. A perspective on the measurement of time in plant disease epidemiology. Plant. Pathol. 53:705-712.

Madden, L.V., Hughes, G, van den Bosch, F. 2007. The study of plant disease epidemics. APS Press, St Paul, MN.

McIntosh, R.A., Wellings C.R., and Park, R.F. 1995. Wheat rusts: An atlas of resistance genes. Kluwer Academic Press, London, UK.

Morais, D., Laval, V., Sache, I., and Suffert, F. 2015. Comparative pathogenicity of sexual and asexual spores of *Zymoseptoria tritici* (Septoria tritici blotch) on wheat leaves. Plant Pathol. 64:1429-1439.

Morais, D., Laval, V., Sache, I., and Suffert, F. 2016. Inferring the origin of primary inoculum of *Zymoseptoria tritici* from differential adaptation of resident and immigrant populations to wheat cultivars. Eur. J. Plant Pathol. 145:393-404.

Newton, A.C., and McGurk, L. 1991. Recurrent selection for adaptation of *Erysiphe graminis* f.sp. *hordei* to partial resistance of barley. J. Phytopathol. 132:328-338.

Pariaud, B., van den Berg, F., van den Bosch, F., Powers, S.J., Kaltz, O., and Lannou, C. 2012. Shared influence of pathogen and host genetics on a trade-off between latent period and spore production capacity in the wheat pathogen, *Puccinia triticina*. Evol. Appl. 6:303-312.

Parlevliet, J.E. 1975. Partial resistance of barley to leaf rust, *Puccinia hordei*. I. Effect of cultivar and development stage on latent period. Euphytica 24:21-27.

Parry, M., Gibson, G. J., Parnell, S., Gottwald, T.R., Irey, M.S., Gast, T.C., and Gilligan, C.A. 2014. Bayesian inference for an emerging arboreal epidemic in the presence of control. Proc. Nat. Acad. Sci. 111:6258-6262.

Pinon, J., and Frey, P. 2005. Interactions between poplar clones and *Melampsora* populations and their implications for breeding for durable resistance. *In*: Pei, M.H., McCracken, A.R. (eds.). Rust diseases of willow and poplar, CABI International, p. 139-154.

Précigout, P-A., Claessen, D., and Robert, C. 2017. Crop fertilisation impacts epidemics and optimal latent period of biotrophic fungal pathogens. Phytopathology, in press.

Robert, C., Bancal, M.-O., Ney, B., and Lannou, C. 2005. Wheat leaf photosynthesis loss due to leaf rust, with respect to lesion development and leaf nitrogen status. New phytol. 165:227-241.

Robert, C., Fournier, C., Andrieu, B., and Ney, B. 2008. Coupling a 3D virtual wheat (*Triticum aestivum*) plant model with a Septoria tritici epidemic model (Septo3D): A new approach to investigate plant-pathogen interactions linked to canopy architecture. Funct. Plant Biol. 35:997-1013.

Rossi, V., Giosuè, S., and Caffi, T. 2009. Modelling the dynamics of infections caused by sexual and asexual spores during *Plasmopara viticola* epidemics. J. Plant. Pathol. 91:615-627.

Ruel, J.J., and Ayres, M.P. 1999. Jensen's inequality predicts effects of environmental variation. Trends Ecol. Evol. 14:361-366.

Savary, S. 1987. Decrease by plant development and leaf age of susceptibility of groundnut to rust (*Puccinia arachidis*) in a susceptible cultivar. Neth. J. Plant. Pathol. 93:25-31.

Savary, S., De Jong, P.D., Rabbinge, R., and Zadoks, J.C. 1990. Dynamic simulation of groundnut rust: a preliminary model. Agric. Syst. 32:113-141.

Scherm, H., and van Bruggen, A.H.C. 1994. Effects of fluctuating temperatures on the latent period of lettuce downy mildew (*Bremia lactucae*). Phytopathology 84:853-859.

Shaw, M.W. 1990. Effects of temperature, leaf wetness and cultivar on the latent period of *Mycosphaerella graminicola* on winter wheat. Plant Pathol. 39:255-268.

Shearer, B.L., and Zadoks, J.C. 1972. The latent period of *Septoria nodorum* in wheat. 1. The effect of temperature and moisture treatments under controlled conditions. Neth. J. Plant Pathol. 78:237-247.

Suffert, F., Goyeau, H., Sache, I., Carpentier, F., Gélisse, S., Morais, D., and Delestre, G. 2017. Trade-off between intra- and interannual scales in the evolution of aggressiveness in a local plant pathogen population. bioRxiv, https://doi.org/10.1101/151068.

Suffert, F., Ravigné, V., and Sache, I. 2015. Seasonal changes drive short-term selection for fitness traits in the wheat pathogen *Zymoseptoria tritici*. Appl. Env. Microb. 81:6367-6379.

Suffert, F., and Sache, I. 2011. Relative importance of different types of inoculum to the establishment of *Mycosphaerella graminicola* in wheat crops in north-west Europe. Plant Pathol. 60:878-889.

Suffert, F., Sache, I., and Lannou, C. 2011. Early stages of septoria tritici blotch epidemics of winter wheat: Build-up, overseasoning, and release of primary inoculum. Plant Pathol. 60:166-177.

Suffert, F., Sache, I., and Lannou, C. 2013. Assessment of quantitative traits of aggressiveness in *Mycosphaerella graminicola* on adult wheat plants. Plant Pathol. 62:1330-1341.

Tomerlin, J.R., and Jones, A.L. 1983. Effect of temperature and relative humidity on the latent period of *Venturia inaequalis* in apple leaves. Phytopathology 73:51-54.

Tomerlin, J.R., Eversmeyer, M.G., Kramer, C.L., and Browder, L.E. 1983. Temperature and host effects on latent and infectious periods and on urediniospore production of *Puccinia recondita* f. sp. *tritici*. Phytopathology 73:414-419.

van der Plank, J.E. 1963. Plant diseases: epidemics and control. Academic Press, London.

Villaréal, L.M.M.A., and Lannou C. 2000. Selection for increased spore efficacy by host genetic background in a wheat powdery mildew population. Phytopathology 90:1300-1306.

Viljanen-Rollinson, S.L.H., Marroni M.V., Butler, R.C., Deng Y., and Armour, T. 2005. Latent periods of Septoria tritici blotch on ten cultivars of wheat. New Z. Plant Prot. 58:256-260.

Webb, D.H., and Nutter, F.W.J. 1997. Effects of leaf wetness duration and temperature on infection efficiency, latent period, and rate of pustule appearance of rust in alfalfa. Phytopathology 87:946-950.

Xiao, W., Yu, Q., Flerchinger, G.N, and Zheng, Y.F. 2006. Evaluation of SHAW model in simulating energy balance, leaf temperature, and micrometeorological variables within a maize canopy. Agron. J. 98:722-729.

Xu, X.M. 1999. Effects of temperature on the latent period of the rose powdery mildew pathogen, *Sphaerotheca pannosa*. Plant Pathol. 48:662-667.

Zadoks, J.C. 1972. Methodology of epidemiological research. Ann. Rev. Phytopath. 10:253-276.