Sperm chemotaxis is driven by the slope of the chemoattractant

concentration field

Ramírez-Gómez H.V.[1], Jiménez-Sabinina V.[2], Tuval I.[3], Velázquez-Pérez M.[1], Beltrán

C.[1], Carneiro J.[4], Wood C.D.[1, 5], Darszon A.[1, *], Guerrero A.[1, 5, *].

[1] Departamento de Genética del Desarrollo y Fisiología Molecular, Instituto de

Biotecnología, Universidad Nacional Autónoma de México (UNAM), Cuernavaca,

Morelos, 62210, México.

[2] Cell Biology and Biophysics Unit, European Molecular Biology Laboratory (EMBL),

Heidelberg, Germany.

[3] Mediterranean Institute for Advanced Studies, IMEDEA (CSIC-UIB), Esporles,

Balearic Islands, Spain.

[4] Instituto Gulbenkian de Ciência (IGC), Rua da Quinta Grande, 6 2780-156, Oeiras,

Portugal.

[5] Laboratorio Nacional de Microscopía Avanzada, Instituto de Biotecnología,

Universidad Nacional Autónoma de México (UNAM), Cuernavaca, Morelos, 62210,

México.

[*] Correspondence and requests for materials should be addressed to A.G. or A.D.

(email: adanog@ibt.unam.mx, darszon@ibt.unam.mx).

Abstract

Spermatozoa are attracted to their conspecific female gamete by diffusive molecules

released from the egg investments, a process called chemotaxis. The decapeptide

speract induces metabolic and permeability changes in *Strongylocentrotus purpuratus*

sea urchin sperm. In spite of decades since speract purification from *S. purpuratus* egg

investments, sperm chemotaxis has not been demonstrated in this species. By studying

how the stimulus function, which spermatozoa experience during the accumulation of

bound chemoattractants throughout their trajectory, influences both their motility

response and their internal Ca^{2+} oscillations, we were able to show, for the first time,

that *S. purpuratus* spermatozoa exhibit chemotaxis under sufficiently steep speract

concentration gradients. We demonstrate that this process arises through frequency

entrainment of the coupled metabolic oscillators.

Introduction

Broadcast spawning organisms, such as marine invertebrates, release their gametes into open sea, where they are often subject to extensive dilution that reduces the probability of gamete encounter (Lotterhos, 2010). In many marine organisms, female gametes release diffusible molecules that attract homologous spermatozoa (Lillie, 1913, Miller, 1985, Suzuki, 1995). Propelled by their beating flagella, spermatozoa detect and respond to chemoattractant concentration gradients by steering their swimming trajectory toward the gradient source: the egg. Though it was in bracken ferns where sperm chemotaxis was first identified (Pfeffer, 1884), sea urchins are currently the best-characterized model system for studying sperm chemotaxis at a molecular level (Alvarez et al., 2012, Cook et al., 1994, Darszon et al., 2008, Strunker et al., 2015, Wood et al., 2015).

The sea urchin egg is surrounded by an extracellular matrix which contains short sperm-activating peptides (SAPs), that modulate sperm motility through altering intracellular Ca^{2+} concentration ($[Ca^{2+}]_i$) and other signaling intermediates (Darszon et al., 2008, Suzuki, 1995). The probability of sperm-egg encounter is enhanced by the prompt transduction of the biochemical signals triggered by SAPs into the appropriate reorientation of the sperm trajectory.

The decapeptide speract is one of best characterized members of the SAP family due to its powerful stimulating effect on metabolism, permeability and motility in *S. purpuratus* and *L. pictus* spermatozoa. In 2003, it was shown that the binding of speract to its receptor located in the flagellar plasma membrane, triggers a train of $[Ca^{2+}]_i$ increases in immobilized *S. purpuratus* spermatozoa (Wood et al., 2003). This calcium signal was thought to regulate the activity of dynein motor proteins in the flagellum, thus might modulate the swimming path of sperm (Brokaw, 1979).

A direct link between $[Ca^{2+}]_i$ signaling and sperm motility was established through

the use of optochemical techniques to rapidly, and non-turbulently, expose swimming

sea urchin spermatozoa to their conspecific attractant in a well-controlled experimental

regime (Wood et al., 2005, Bohmer et al., 2005). Currently, it is well established that

the transient $[Ca^{2+}]_i$ increases triggered by chemoattractants produce a sequence of turns

and straight swimming episodes (turn-and-run), where each turning event results from

the rapid increase in the $[Ca^{2+}]_i$ (Wood et al., 2005, Bohmer et al., 2005, Shiba et al.,

2008, Alvarez et al., 2012). The turn-and-run seems to be a general requirement for

sperm chemotaxis, however it is not sufficient on its own to produce a chemotactic

response (Guerrero et al., 2010a, Strunker et al., 2015, Wood et al., 2007, Wood et al.,

2005).

Our current understanding of chemotaxis, suggests that sperm first sample the

chemoattractant concentration gradient by swimming in periodic paths (either circular

2D or helical 3D). During the sampling phase, the accumulation of bound

chemoattractants triggers $[Ca^{2+}]_i$ transients that control the waveform of the flagellar

beat. In this way, the alternate periods of asymmetrical (turn) and symmetrical (run)

flagellar beating give rise to a looping swimming pattern that guides up to the source of

the chemoattractant gradient.

Friedrich and Jülicher proposed a generic theory that captures the essence of sperm

navigation following periodic paths in a non-homogeneous chemoattractant field, where

the sampling of a periodic concentration stimulus $s(t)$ is translated by intracellular

signaling $i(t)$ into the periodic modulation of the swimming path curvature $k(t)$

(Friedrich and Jülicher, 2007, Friedrich and Jülicher, 2008). As result, the periodic

swimming path drifts in a direction that depends on the internal dynamics of the

signaling system. In this theory, the latency of the intracellular signaling (the $[Ca^{2+}]_i$

signal), expressed as the phase shift between *s(t)* and *k(t)*, is a crucial determinant of the

directed looping of the swimming trajectory up the chemical concentration field. This

theory also predicts that chemotaxis is a robust property of the system that does not

require fine-tuning of parameters (if the signaling system is adaptive) (Friedrich and

Jülicher, 2008, Friedrich and Jülicher, 2009). In other words, there is a large range of

parameters for which sperm chemotaxis is a robust outcome, providing an effective way

for sampling the local chemoattractant concentration field and detecting the direction of

the concentration gradient (Kaupp et al., 2003, Kashikar et al., 2012, Pichlo et al., 2014,

Friedrich and Jülicher, 2008, Friedrich and Jülicher, 2007).

Even though the conceptual framework of Friedrich & Jülicher provides insights into

the mechanism governing sperm chemotaxis, it does not explore the scenario whereby

chemoattractants trigger an autonomous $[Ca^{2+}]_i$ oscillator operating in the absence of a

periodic stimulus. The existence of an autonomous $[Ca^{2+}]_i$ oscillator triggered by

chemoattractants (Wood et al., 2003, Espinal et al., 2011, Aguilera et al., 2012) suggests

that sperm chemotaxis might operate in a dynamical space where two autonomous

oscillators, namely the stimulus function and the internal Ca^{2+} oscillator, reach

frequency entrainment (Pikovsky et al., 2001).

In spite of 30 years of research since speract's isolation from *S. purpuratus* oocytes

(Hansbrough and Garbers, 1981, Suzuki, 1995), chemotaxis of *S. purpuratus* sperm

towards this peptide has not yet been demonstrated (Cook et al., 1994, Darszon et al.,

2008, Guerrero et al., 2010b, Kaupp, 2012, Miller, 1985, Wood et al., 2015). A

comparison between individual *L. pictus* and *S. purpuratus* sperm responses to a

specific chemoattractant concentration gradient generated by photoactivating caged

speract (CS) revealed that only *L. pictus* spermatozoa exhibit chemotaxis under these

conditions (Guerrero et al., 2010a). In that study, *L. pictus* spermatozoa experience $[Ca^{2+}]_i$ fluctuations and pronounced turns while swimming in descending speract gradients, that result in spermatozoa reorienting their swimming behavior along the positive chemoattractant concentration gradient. In contrast, *S. purpuratus* spermatozoa experience similar trains of $[Ca^{2+}]_i$ fluctuations that in turn drive them to relocate, but with no preference towards the center of the chemoattractant gradient (Guerrero et al., 2010a).

The precise triggering of $[Ca^{2+}]_i$ fluctuations associated to sperm turning events towards the chemoattractant gradient, as well as the turn-and-run behavior seem to be general requirements for sperm chemotaxis in marine invertebrates (Bohmer et al., 2005, Guerrero et al., 2010a, Jikeli et al., 2015, Kashikar et al., 2012, Shiba et al., 2008, Strunker et al., 2015, Wood et al., 2015), and could be important features of sperm chemotaxis in general.

In the present work, we investigate whether *S. purpuratus* spermatozoa can undergo chemotaxis. Particularly, we examined whether there is a physical limit to the sampling of the chemoattractant concentration gradient needed for detection that has, to date, prevented observation and characterization of their chemotactic response. We report that *S. purpuratus* spermatozoa are chemotactic only when exposed to much steeper speract concentration gradients than those previously employed. Furthermore, we explored the coupling between the recruitment of speract molecules during the sperm voyage, the triggered signaling cascade and the internal Ca^{2+} oscillator; and demonstrate that sperm chemotaxis arises through coupled metabolic oscillators.

Results

Chemotaxis refers to the directed movement of an organism or a cell in a chemical

gradient. The first step in a chemotactic response is the sampling of a chemoattractant

field, from where in a given time T, statistical fluctuations limit the precision with

which the searcher can determine the concentration of the chemoattractant (Berg and

Purcell, 1977, Vergassola et al., 2007, Dusenbery, 2011).

Assuming that the number of ligands that collide with a single receptor per unit of

time is proportional to radius of the receptor s, the concentration of the attractant c and

the diffusion constant of the attractant D, the total number of collisions between N_R

receptors in a time interval T is $x = N_R s D c T$. The statistical fluctuation on the number

of ligand-receptor collisions x follows a Poisson distribution, and therefore the expected

value $E[x]$ and the variance $Var[x]$ are the same. In this scenario, the coefficient of

variation is $\frac{SD[x]}{E[x]} = \frac{\sqrt{Var[x]}}{E[x]} = \frac{\sqrt{E[x]}}{E[x]}$. This coefficient can be interpreted as the smallest

fractional error attainable in the determination of the concentration of the attractant (see

Theory section) (Berg and Purcell, 1977, Dusenbery, 2011, Vergassola et al., 2007).

$$u_1 = (N_R s D c T)^{-1/2} \qquad (1)$$

Uncertainty in the determination of the presence of chemoattractant molecules

The flagella of sea urchin spermatozoa possess a large number of high affinity SAP

receptors (10^4-10^6 depending on species and author) (Nishigaki and Darszon, 2000,

Shimomura and Garbers, 1986, Smith and Garbers, 1982, Kaupp et al., 2008). This

great number of SAP receptors provides spermatozoa with a fine-tuned sensory system

able to respond to a wide range of SAP concentrations (10^{-12}-10^{-6} M). That these

receptors are distributed along the entire flagellum increases the probability of capturing

even a few arriving molecules (Kaupp et al., 2003, Kashikar et al., 2012).

To understand the theoretical sampling capabilities of sea urchin spermatozoa, given

a specific speract concentration (in a homogeneous concentration field), and flagellar

receptor availability, we first computed the smallest fractional error attainable in the

detection of a chemoattractant u_1 at different concentrations, considering distinct

receptor numbers. A theoretical *S. purpuratus* spermatozoon possessing $N_R = 2 \times 10^4$

receptors, will sample a medium containing 10 nM of speract in time $T = 0.5$ s with an

uncertainty of $u_1 = 0.006$ (**Figure 1 – figure supplement 1a** and **Figure 1 – table**

supplement 1). Here, the time scale T was considered as the time needed for a *S.*

purpuratus spermatozoon to traverse a distance equivalent to approximately half a

circumference of its circular trajectory when confined to the water-glass boundary, in a

homogeneous field of speract. The least fractional error attainable in the determination

of the concentration of the attractant has no upper boundary, however u_1 values greater

than 1 indicates that stochastic fluctuations dominate over the signal (Berg and Purcell,

1977). For simplicity, u_1 was considered to operate in the range [0, 1]. In other words,

our model spermatozoon, when exposed to a chemoattractant concentration of 10 nM,

will be able to determine the presence of speract molecules with 0.6% of uncertainty in

0.5 s.

If the spermatozoon finds itself in a sampling space of 1 pM of speract, this level of

uncertainty increases to 60% over the same interval. It is likely that at speract

concentrations below picomolar, an *S. purpuratus* spermatozoon will require relatively

extended sampling times to determine the presence of speract molecules. For example,

for a spermatozoon sampling a 1 pM concentration field of speract for 60 seconds, the

uncertainty of receptor occupancy drops to 6% (**Figure 1 – figure supplement 1b**).

An important difference between *S. purpuratus* and *L. pictus* spermatozoa is their

respective receptor density; the former possesses 3.15x fewer receptors than the latter (2

x 10^4 versus 6.3 x 10^4 receptors per cell, respectively) (**Figure 1 – table supplement 1**)

(Nishigaki and Darszon, 2000, Nishigaki et al., 2001). Note that if *S. purpuratus*

spermatozoa were to possess a receptor density equivalent to *L. pictus*, they would be

able to sample a speract concentration field of 1 pM within 0.5 seconds with an

uncertainty of $u_1 = 0.4$ (40%) (**Figure 1 – figure supplement 1a**). Given the differences

in sensitivities between these two species, the lack of chemotactic responses in *S.*

purpuratus spermatozoa could possibly be explained, at least partially, by their lesser

density of speract receptors, compared to *L. pictus*.

Uncertainty in the determination of sperm position in a non-homogeneous

chemoattractant concentration field

To detect a chemoattractant concentration gradient, the signal at two given points,

sampled throughout a time interval must be greater than the noise, which in this case

would arise from spontaneous stochastic activation of one or more components of the

chemoattractant signaling pathway (**Figure 1a**) (Berg and Purcell, 1977, Vergassola et

al., 2007, Wood et al., 2015, Dusenbery, 2011). Information regarding the direction of

the chemoattractant gradient results from the magnitude difference of the signal at two

sampled positions (for further explanation see **Theory** section). The smallest fractional

error in determination of the chemoattractant gradient direction due to the occupation

state of receptors is:

$$u_2 = v^{-1}T^{-3/2}(N_R sD)^{-1/2}c^{1/2}\varepsilon^{-1} \qquad (2)$$

where *v* is the swimming velocity and $\varepsilon = dc/dr$ is the chemoattractant concentration

gradient (**Figure 1a**) (Berg and Purcell, 1977, Dusenbery, 2011, Vergassola et al.,

2007). A locality requirement must also be met: the change in concentration across the

distance spanned during *T* must be small, compared to the concentration itself

(Vergassola et al., 2007).

The slope of the chemoattractant concentration gradient ε directly impinges on the

ability to reliably determine the source of the attractant (**Figure 1b**). At high

concentrations of speract (10^{-8} to 10^{-6} M) the change in receptor occupancy experienced

by *S. purpuratus* spermatozoa, at two given distinct locations, allows the reliable

assessment of the chemical gradient, when $u_2 < 0.05$ (**Figure 1b**) (As for u_1, u_2 was

considered to operate in the range [0, 1]) (Vergassola et al., 2007, Berg and Purcell,

1977). However, at low concentrations of speract (below of 10^{-9} M), where other

parameters are identical, stochastic fluctuations tend to dominate over the signal

(**Figure 1b**). Under these low-concentration regimes, the slope of the chemoattractant

gradient is determinant; shallow gradients ($\varepsilon < 10^{-12}$ M μm^{-1}) increase the uncertainty

for detection of gradient polarity. In contrast, steeper chemoattractant gradients, *i.e.* $\varepsilon \geq$

10^{-11} M μm^{-1}, would allow *S. purpuratus* spermatozoa to determine the orientation of

the chemical gradient dependably (**Figure 1b**).

Previously it was shown that *L. pictus* spermatozoa experience chemotaxis in

response to exposure to a gradient of speract (Guerrero et al., 2010a). Notably, under

the same experimental regime *S. purpuratus* spermatozoa undergo a motility response

to the presence of speract, yet do not demonstrate chemotaxis. It seems reasonable to

speculate that the lack of chemotactic responses in *S. purpuratus* spermatozoa is due to

their reduced receptor density relative to *L. pictus* spermatozoa. The chemoreception

model predicts that, in contrast to *L. pictus*, *S. purpuratus* spermatozoa might determine

the direction of the attractant gradient in the range of 10^{-10} to 10^{-9} M (**Figure 1b**, orange

and yellow lines with $u_2 < 5\%$), and thus can only determine the orientation of the

chemical gradient in steep, but not shallow concentration gradients (**Figure 1b**, yellow

and cyan lines with $u_2 < 5\%$). If the latter holds to be true, then the model predicts that

S. purpuratus spermatozoa should be able to experience chemotaxis when exposed to

relatively steep speract gradients, which have not been tested experimentally to date.

The ratio of uncertainties for reliable assessment of the direction of a speract gradient

between *S. purpuratus* and *L. pictus* spermatozoa can be expressed as:

$$\frac{u_{2\,purpuratus}}{u_{2\,pictus}} = S\frac{\varepsilon_{pictus}}{\varepsilon_{purpuratus}},$$

$$S = \left(\frac{v_{purpuratus}}{v_{pictus}}\right)^{-1}\left(\frac{T_{purpuratus}}{T_{pictus}}\right)^{-3/2}\left(\frac{N_{R\,purpuratus}}{N_{R\,pictus}}\right)^{-1/2} \quad\quad\quad (3, 4)$$

with *S* being a factor that scales the slope of the speract gradient to a regime that

allows *S. purpuratus* spermatozoa to detect the local direction of the chemical gradient.

In other words, the model predicts that *S. purpuratus* spermatozoa should undergo

chemotaxis in a speract gradient approximately three times steeper than the gradient that

drives chemotaxis in *L. pictus* spermatozoa, with $\varepsilon_{purpuratus} = 3.25\ \varepsilon_{pictus}$.

In summary, the chemoreception model suggests that *S. purpuratus* spermatozoa

detect chemoattractant gradients with less sensitivity than those of *L. pictus*. It also

predicts that *S. purpuratus* spermatozoa may detect chemoattractant gradients in the 10^{-9}

M regime with sufficient certainty only if the difference in sampling concentration is

greater than 10^{-11} M μm^{-1} (steep concentration gradients) (**Figure 1b**). Given this

prediction, we designed and implemented an experimental condition where *S.*

purpuratus spermatozoa should experience chemotaxis.

***S. purpuratus* spermatozoa accumulate at steep speract concentration gradients**

Our experimental setup is designed to generate specific concentration gradients by

focusing a brief (200 ms) flash UV light along an optical fiber, through the objective,

and into a field of swimming *S. purpuratus* spermatozoa containing CS at 10 nM in

artificial sea water (Tatsu et al., 2002, Guerrero et al., 2010a). To test experimentally

whether *S. purpuratus* undergo chemotaxis, as predicted from the chemoreception

model, we varied the slope of the chemoattractant gradient by separately employing

four optical fibers of distinct diameters that could be arranged into five different

configurations (*f1, f2, f3, f4, f5*).

Each configuration produces a characteristic pattern of UV illumination within the

imaging field (**Figure 2**). The UV intensity was measured at the back focal plane of the

objective for each fiber configuration (**Figure 2a** and **Figure 2 – table supplement 1**).

The spatial derivative of the imaged UV light profile was computed for use as a proxy

for the slope of the speract concentration gradient (**Figure 2b**). By examining these UV

irradiation patterns, we calculated that, at the gradient peak, the highest concentration of

speract released through photo-liberation from CS is generated by the *f5* fiber, followed

by *f4 > f3 > f2 > f1*. The steepest UV irradiation gradients are those generated by the *f2,*

f3 and *f5* fibers (**Figure 2b**).

Irrespective of the optical fiber used, the photo-activation of caged speract triggers

the stereotypical Ca^{2+}-dependent motility responses of *S. purpuratus* spermatozoa

(**Figure 2c, Movies 1, 2, 5, 6** and **7**). To determine whether these changes lead to sperm

accumulation, we developed an algorithm which automatically scores the number of

spermatozoa at any of four defined concentric regions (R1, R2, R3, and R4) relative to

the center of the speract concentration gradient (**Figure 2 – figure supplement 1**).

Photo-liberation of speract through either *f2* or *f3* fiber, but not through *f1, f4* or *f5*

fibers, lead to the accumulation of *S. purpuratus* spermatozoa towards the center of the

speract gradient (zones R1 and R2) within the first 5-10 seconds after UV irradiation

(**Figure 3**, **Figure 3 - figure supplement 1, Figure 3 - figure supplement 2**, **Movies 1**,

and **2**).

Interestingly, for *f2*, the number of spermatozoa increases in R1 and R2 and

decreases in R3 and R4, indicating that cells from the R3 and R4 regions most probably

relocate towards R1 and R2 (**Figure 3a**, **Figure 3 - figure supplent 2** and **Movie 1**). In

the case of the *f3* gradient, the number of spermatozoa increases in the R1, R2 and R3

regions and decreases in the R4 region, suggesting that spermatozoa in R4 and possibly

outside of the imaged field are entering the other regions (**Figure 3a** and **Movie 2**). The

maximum sperm accumulation (about two-fold) occurs in R1 for the *f3* gradient (**Figure**

3a and **Movie 2**).

In the case of exposure to the *f5* speract concentration gradient the number of

spermatozoa showed a tendency to increase in R2, R3 and R4, although it was only

statistically significant in R4 (**Figure 3a**, **Figure 3 - figure supplement 2** and **Movie**

6). Gradients *f1, f4* and negative controls (Low $[Ca^{2+}]_i$ or High $[K^+]_e$) did not show

increased sperm number in any region (**Figure 3a**, **Figure 3 - figure supplement 1**,

Figure 3 - figure supplement 2, **Movies 3**, **4**, **5** and **6**).

We also evaluated the corresponding $[Ca^{2+}]_i$ changes across the imaging field for

each imposed speract concentration gradient. Gradients *f2* to *f5* increase $[Ca^{2+}]_i$ in

spermatozoa at least two-fold, while the increase for the *f1* gradient was modest (**Figure**

3b). Interestingly, $[Ca^{2+}]_i$ levels rose highest upon exposure to the *f4* gradient, even

though the number of spermatozoa did not increase significantly (**Figure 3a**), which

underlines the notion that elevated $[Ca^{2+}]_i$ levels are necessary, but not sufficient to

drive the accumulation of spermatozoa (**Figure 3**), as previously suggested (Kaupp et

al., 2003, Wood et al., 2005, Bohmer et al., 2005, Guerrero et al., 2010a, Alvarez et al.,

2012).

In summary, *S. purpuratus* spermatozoa accumulate towards the center of the speract

gradients generated by the *f2-* and *f3*-fibers, which are the two optical fibers that

generate UV light profiles with steeper slopes compared to the *f1* and *f4* fibers (**Figure**

2b). Notably, use of fibers *f4* and *f5* photo-releases higher concentrations of speract (by

providing higher UV energies than other fibers) (**Figure 2a, Figure 2 table**

supplement 1), yet they trigger neither the maximum accumulation of *S. purpuratus*

spermatozoa at the center of the chemoattractant field, nor the most elevated $[Ca^{2+}]_i$

response.

S. purpuratus spermatozoa undergo chemotaxis upon exposure to steep speract

gradients

The spatial derivative of the UV profiles shown in **Figure 2b** indicates that the

steeper light gradients generated from UV irradiation are those of *f2, f3* and *f5*, which

are assumed to generate the most pronounced speract gradients of similar form. This

assumption is strictly only valid at the instant of UV exposure, as subsequently the

speract gradient dissipates over time with a diffusion constant of $D \approx 240 \ \mu m^2 \ s^{-1}$.

We further sought to understand how the stimulus function, which *S. purpuratus*

spermatozoa experience during the accumulation of bound speract throughout their

trajectory, influences their motility response. For this purpose, we computed the spatio-

temporal dynamics of the speract gradient for *f1, f2, f3, f4* and *f5* fibers (**Figure 4a, b**

and **Figure 4 - figure supplement 1**) and analyzed the trajectories of spermatozoa

swimming in these five distinct speract gradient configurations (**Figure 4c** and **Figure 4**

- figure supplement 2). From these trajectories, we derived the chemotaxis indices

(LECI, (Yoshida et al., 2002)) (**Figure 4d** and **Figure 5**), and computed the stimulus

function of individual spermatozoa in response to each of the five speract gradient

forms (**Figure 4e** and **Figure 4 - figure supplement 2**).

Fibers *f2*, *f3* and *f5* triggered chemotaxis in *S. purpuratus* spermatozoa (**Figure 4**,

Figure 4 - figure supplement 2 and **Figure 5**). Under such conditions, only those

spermatozoa located in the regions R3 and R4 at the moment of speract uncaging

underwent chemotaxis (**Figure 5**; see **Figure 4** and **Figure 4 - supplement figure 2** for

single sperm reorienting towards the center of speract gradients generated by the *f2* and

f3 fibers, respectively). For both, *f2* and *f3* conditions, sperm chemotaxis initiated within

the first 3 s after speract exposure, and lasted less than 10 s (**Figure 5**, **Figure 3 - figure**

supplement 1, **Movies 1** and **2**).

Figure 4b shows that, for the *f2* condition, the slope of the speract gradient within

R2 and regions is barely altered during the initial 3 seconds of speract exposure (gray

shading). Spermatozoa in that region sample a pseudo-static speract gradient over a

short (2-3 s) period, which is apparently sufficient to permit detection of the direction of

the gradient. In contrast, spermatozoa located at other regions (R1, R3, R4) experience a

speract gradient with a slope that changes during the sampling time (**Figure 4a**, **4b**).

In summary, *f2*, *f3* and *f5* optical fibers generate speract concentration gradients that

drive the accumulation of *S. purpuratus* spermatozoa. Of note, the *f4* fiber generates a

speract gradient that activates motility responses of *S. purpuratus* spermatozoa but not

chemotaxis (**Movie 6**), although it was previously shown that it triggers the

accumulation of *L. pictus* spermatozoa (Guerrero et al., 2010a).

The model of chemoreception presented in the previous section (**equations (3, 4)**)

predicts a scaling rule for chemotactic responses between *S. purpuratus* and *L. pictus*

spermatozoa of *S > 3.25*. Moreover, the UV-irradiation profiles shown in **figure 2b**

indicate that *f2*, *f3*, and *f5* fibers generate steeper speract gradients than *f4* fiber.

To reliably determine the direction of the chemoattractant concentration gradient, the

signal difference *dc* between two sampled positions *dr* must be greater than the noise

(**Figure 1a**). In order to test the prediction of the chemoreception model, we computed

the local relative slope ξ detected by single spermatozoa exposed to a given speract

concentration gradient, with $\xi = c^{-1/2}dc/dr$ (**Figure 4e**).

We found that, in agreement with the chemoreception model, the maximum relative

slope $\xi_{max} = Max(\xi_1, \xi_2, \xi_3, ..., \xi_n)$ experienced by *S. purpuratus* spermatozoa, when

exposed to *f2* and *f3* speract gradients, were up to 2-3 times greater than that

experienced when exposed to the *f4*-generated speract gradient (**Figure 6a**). In addition,

they were also up to 2-3 times greater than the relative slope experienced by *L. pictus*

spermatozoa when exposed to *f4* speract gradient (**Figure 6a**), hence supporting the

predicted scaling rule for the detection of the speract concentration gradient between *L.*

pictus and *S. purpuratus* spermatozoa.

The slope of the speract gradient is the critical determinant for the strength of

coupling between the stimulus function and the internal Ca^{2+} oscillator

To test the idea that the slope of the speract gradient regulates the coupling between

the stimulus function and the internal Ca^{2+} oscillator triggered by speract, we made use

of a generic model for coupled phase oscillators (Pikovsky et al., 2001). In its simplest

form, the model describes two phase oscillators of intrinsic frequencies ω_1 and ω_2

coupled with a strength γ through the antisymmetric function of their phase difference ϕ

$= \varphi_1 - \varphi_2$. The time evolution of ϕ then follows an Adler equation $d\phi/dt = \Delta\omega - 2\gamma \sin(\phi)$,

which is the leading order description for weakly-coupled non-linear oscillators. In the

present case, the two coupled oscillators are the internal Ca^{2+} oscillator and the

oscillations in the stimulus function induced in spermatozoa swimming across a speract

gradient. The former occurs even for immotile cells, for which there are no stimulus

oscillations under a spatially uniform speract field (**Figure 6 - supplement 1,** and

Movie 8); while the later exists under two tested negative controls: cells swimming in

Low Ca^{2+} and in High K$^+$ artificial sea water, both of which inhibit Ca^{2+} oscillations (see

Figure 3, Movie 3 and **4).**

There are two immediate predictions from the Adler model: first, there is a minimum

coupling strength necessary for the two oscillators to synchronize ($\gamma_{min} = \Delta\omega/2$). For

weaker coupling (*i.e.* $\gamma < \gamma_{min}$), the two oscillators run with independent frequencies and,

hence, the phase difference increases monotonically with time; second, and within the

synchronous region (*i.e.* $\gamma > \gamma_{min}$), the phase difference between the oscillators is

constant and it does not take any arbitrary value, but rather follows a simple relation to

the coupling strength ($\phi_{sync} = arcsin(\Delta\omega/2\gamma)$). **Figure 6b** shows the two regions in the

parameter space given by $\Delta\omega$ and γ. The boundary between these two regions

corresponds to the condition $\gamma = \gamma_{min}$ and it delimits what is known as an Arnold's

tongue.

We measure the difference in intrinsic frequency by looking at the instantaneous

frequency of the internal Ca^{2+} oscillator just before and after the speract gradient is

established. The range of measured $\Delta\omega$ is shown in the same figure as a band of

accessible conditions in our experiments (mean of $\Delta\omega$, black line; mean $\pm$ standard

deviation, green dashed lines). If the driving coupling force between the oscillators is

the maximum slope of the speract gradient, *i.e.* $\gamma = \xi_{max}$, we would expect to find a

minimum slope ($\overline{\xi^{*}_{max}}$) below which no synchrony is observed.

This is indeed the case as clearly shown in **Figure 6a**, **6c** and **6d** (magenta line).

Moreover, and for cells for which synchronization occurs, the measured phase

difference is constrained by the predicted functional form of $\phi_{sync} = \phi_{sync}(\Delta\omega, \gamma)$ as can

be verified from the collapsed data shown in **Figure 6c**, and **6d** within the theoretical

estimates.

Altogether, the excellent agreement of this simple model of coupled phase oscillators

with our data, points to the slope of the speract gradient as the driving force behind the

observed synchronous oscillations and, as a result, for the chemotactic ability of sea

urchin spermatozoa.

Discussion

Marine spermatozoa, together with many motile microorganisms, explore their

environment via helical swimming paths, whereupon encountering a surface these

helices collapse to circular trajectories. The intrinsic periodicity of either swimming

behavior commonly results in the periodic sampling of the cells chemical environment

with direct implications for their ability to accurately perform chemotaxis.

A strict requirement for sperm chemotaxis is the presence of extracellular Ca^{2+}. For

chemotaxis to occur, the timing of the Ca^{2+} transients (*i.e.* the intracellular Ca^{2+}

oscillations) triggered by the chemoattractants must also be kept in phase with the

polarity of the chemoattractant concentration field, which in this, and other studies, is

referred as the stimulus function (Bohmer et al., 2005, Guerrero et al., 2010a, Kaupp et

al., 2008, Friedrich and Jülicher, 2008). This requisite coupling ensures that the turning

events start at the descending phase of the chemoattractant concentration field;

otherwise spermatozoa are driven away by Ca^{2+}-dependent motility adjustments. The

periodic sampling of chemoattractants by the sperm flagellum continuously feeds back

to the signaling pathway governing the intracellular Ca^{2+} oscillator, hence providing a

potential coupling mechanism for sperm chemotaxis. Indirect evidence for the existence

of a feedback loop operating between the stimulus function and the Ca^{2+} oscillator

triggered by chemoattractants has been found in *L. pictus*, *A. punctulata* and *Ciona*

intestinalis species, whose spermatozoa show robust chemotactic responses towards

their conspecific chemoattractants (Guerrero et al., 2010a, Shiba et al., 2008, Jikeli et

al., 2015, Bohmer et al., 2005).

For almost three decades, chemotaxis had not been observed for the widely studied

S. purpuratus species under diverse experimental conditions, raising doubts about their

capabilities to 'sense' and respond to the spatial cues provided by the speract

concentration gradients. To tackle whether *S. purpuratus* spermatozoa are able to

'sense' a chemoattractant concentration gradient, we use a model of chemoreception

developed by Berg and Purcell, which considers the minimal requirements needed for a

single searcher (*i.e.* a sperm cell) to gather sufficient information to determine the

orientation of a non-uniform concentration field. By considering the difference between

L. pictus and *S. purpuratus* spermatozoa in terms of the number of chemoattractant

receptors, receptor pocket size, sampling time, swimming velocity, sampling distance,

and the local mean and slope of the chemoattractant concentration field, we predicted

that *S. purpuratus* should be able to detect the polarity of a speract concentration field.

The model predicts that speract gradient necessary to guide *S. purpuratus* spermatozoa

would be up to three times steeper than the gradient that drives chemotactic responses

on *L. pictus* spermatozoa. We tested this prediction experimentally by exposing *S.*

purpuratus spermatozoa to various defined speract concentration gradients. We show

that *S. purpuratus* spermatozoa exhibit chemotactic responses but, as predicted by the

chemoreception model, only if the speract concentration gradients are sufficiently steep

(*i.e.* speract gradients that are at least three times steeper than the speract concentration

gradient that drives chemotaxis in *L. pictus* spermatozoa). The shallower speract

gradients previously tested are therefore unable to generate any chemotactic response in

S. purpuratus spermatozoa.

To investigate further the molecular mechanism involved in sperm chemotaxis, we

measured both the stimulus function and the triggered-internal Ca^{2+} oscillations for up

to five hundred *S. purpuratus* spermatozoa exposed to five distinct speract

concentration gradients. We demonstrate that the slope of the chemoattractant

concentration field is a major determinant for sperm chemotaxis in *S. purpuratus*, and

might be an uncovered feature of sperm chemotaxis in general. A steep slope of the

speract gradient entrains the frequencies of the stimulus function and the internal Ca^{2+}

oscillator triggered by the periodic sampling of a non-uniform speract concentration

field. We assessed the transition boundary of the coupling term (the slope of the speract

concentration field) for the two oscillators to synchronize, and found it to be very close

to the boundary where *S. purpuratus* start to experience chemotaxis. The agreement of

our data with a model of weakly coupled phase oscillators, points to the slope of the

speract gradient as the driving force behind the observed synchronous oscillations and,

as a result, for the chemotactic ability of sea urchin spermatozoa.

One can further hypothesize about the evolutionary origin of the described

differences in sensitivity to chemoattractant concentration gradients between *S.*

purpuratus and *L. pictus* spermatozoa given the significant differences between their

ecological reproduction niches. The turbulent environment where sea urchin reproduce

directly impinges on the dispersion rates of small molecules such as speract, hence,

imposing ecological pressure against sperm chemotaxis. For instance, the reproduction

success of *L. pictus*, *S. purpuratus* and *Abalone* species has been shown to peak at

particular hydrodynamic shearing values (Zimmer and Riffell, 2011, Riffell and

Zimmer, 2007, Mead and Denny, 1995, Hussain et al., 2017). We might ask what are

the typical values of the chemoattractant gradients encountered by the different species

in their natural habitats. The correct scale to consider when discussing the small-scale

distribution of chemicals in the ocean is the Batchelor scale, $l_B = (\eta D^2/\zeta)^{1/4}$, where η is

kinematic viscosity, D the molecular diffusivity and ζ is the turbulent dissipation rate

(Batchelor, 2006, Aref et al., 2014). Turbulence stirs dissolved chemicals in the ocean,

stretching and folding them into sheets and filaments at length scales down to the

Batchelor scale: below l_B molecular diffusion dominates and chemical gradients are

smoothened out.

S. purpuratus is primarily found in the low intertidal zone. The purple sea urchin

thrives amid strong wave action and areas with churning aerated water. These more

energetic zones, including tidal channels and breaking waves, generate relatively high

levels of turbulence ($\zeta \sim 10^{-4}$ m^{2}s^{-3}) which lead to relatively small values of l_B and,

hence, to steep gradients (*i.e.* l/l_B). *L. pictus*, on the contrary, is mostly found at the edge

of or inside kelp beds, well below the low tide mark where the levels of turbulence are

much more moderate ($\zeta \sim 10^{-6}$ m^{2}s^{-3}) (Jiménez, 1997, Thorpe, 2007). This difference in

turbulent kinetic energy dissipation rate has a significant effect on the largest chemical

gradients available in a particular habitat for sperm chemotaxis. The ratio of l_B for the

different habitats scales as $l_{Bpurpuratus}/l_{Bpictus} \sim (\zeta_{pictus}/\zeta_{purpuratus})^{1/4} \sim 3$, which fits

considerably well with the relative sensitivity to speract of the two species.

Furthermore, we have shown that *S. purpuratus* spermatozoa experience chemotaxis

toward steeper speract gradients than those that guide *L. pictus* spermatozoa, which is

also compatible with the distinct chemoattractant gradients they might naturally

encounter during their voyage searching for the egg.

The chemoattractant concentration gradients generated in the present study were

near-instantaneously set up by the photo-release of speract in still water. Further

experimental studies are needed to assess the chemotactic ability of sea urchin

spermatozoa to more realistic chemoattractant gradients (as those shaped, for instance,

by hydrodynamic forces in their natural environment) and to shed light into the

mechanisms governing chemotaxis and their ecological implications.

Materials and Methods

Materials

Undiluted *S. purpuratus* or *L. pictus* spermatozoa (Pamanes S. A. de C. V.,

Ensenada, Mexico and Marinus Scientific, LLC. Newport Beach, CA, USA

respectively) were obtained by intracoelomic injection of 0.5 M KCl and stored on ice

until used within a day. Artificial seawater (ASW) was 950 to 1050 mOsm and

contained (in mM): 486 NaCl, 10 KCl, 10 $CaCl_2$, 26 $MgCl_2$, 30 $MgSO_4$, 2.5 $NaHCO_3$,

10 HEPES and 1 EDTA (pH 7.8). For experiments with *L. pictus* spermatozoa, slightly

acidified ASW (pH 7.4) was used to reduce the number of spermatozoa experiencing

spontaneous acrosome reaction. Low Ca^{2+} ASW was ASW but pH 7.0 and with 1 mM

$CaCl_2$, and Ca^{2+}-free ASW was ASW with no added $CaCl_2$. [Ser5; nitrobenzyl-

Gly6]speract, referred to throughout the text as caged speract (CS), was prepared as

previously described (Tatsu et al., 2002). Fluo-4-AM and pluronic F-127 were from

Molecular   Probes,   Inc.   (Eugene,   OR,   USA).   PolyHEME   [poly(2-

hydroxyethylmethacrylate)]   was   from   Sigma-Aldrich   (Toluca,   Edo de   Mexico,

Mexico).

Loading of Ca^{2+}-fluorescent indicator into spermatozoa

Undiluted spermatozoa were suspended in 10 volumes of low Ca^{2+} ASW containing

0.2% pluronic F-127 plus 20 μM of fluo-4-AM and incubated for 2.5 h at 14 ℃.

Spermatozoa were stored in the dark and on ice until use.

Imaging of fluorescent swimming spermatozoa

The cover slips were briefly immersed into a 0.1% wt/vol solution of poly-HEME in

ethanol, hot-air blow-dried to rapidly evaporate the solvent and mounted on reusable

chambers fitting a TC-202 Bipolar temperature controller (Medical Systems Corp.). The

temperature plate was mounted on a microscope stage (Eclipse TE-300; Nikon) and

maintained at a constant 15 ºC. Aliquots of labeled sperm were diluted in ASW and

transferred to an imaging chamber (final concentration ~2×10^5 cells ml^{-1}).

Epifluorescence images were collected with a Nikon Plan Fluor 40x 1.3 NA oil-

immersion objective using the chroma filter set (ex HQ470/40x; DC, 505DCXRU; em,

HQ510LP) and recorded on a DV887 iXon EMCCD Andor camera (Andor Bioimaging,

NC). Stroboscopic fluorescence illumination was supplied by a Cyan LED no. LXHL-

LE5C (Lumileds Lighting LLC, San Jose, USA) synchronized to the exposure output

signal of the iXon camera (2 ms illumination per individual exposure). Images were

collected with AndoriQ 1.8 software (Andor Bioimaging, NC) at 30.80 fps in full-chip

mode (observation field of 200 x 200 µm).

Image processing

The background fluorescence was removed by generating an average pixel intensity

time-projection image from the first 94 frames before uncaging, which was then

subtracted from each frame of the image stack by using the Image calculator tool of

ImageJ 1.49u (Schneider et al., 2012). For **Figure 2c**, the maximum pixel intensity time

projections were created every 3 s from background-subtracted images before and after

the UV flash.

Quantitation of global changes of spermatozoa number and $[Ca^{2+}]_i$

To study the dynamics of overall sperm motility and $[Ca^{2+}]_i$ signals trigger by the

distinct speract gradients, we developed an algorithm that provides an efficient

approach to automatically detect the head of every spermatozoa in every frame of a

given video-microscopy (C/C++, OpenCV 2.4, Qt-creator 2.4.2). Fluorescence

microscopy images generated as described previously were used. The following steps

summarize the work-flow of the algorithm (**Figure 2 - figure supplement 1**):

1. Segment regions of interest from background: This step consists of thresholding

each image (frame) of the video (x, y, t) to segment the zones of interest (remove

noise and atypical values). Our strategy includes performing an automatic

selection of threshold value for each Gaussian blurred image (I_G) (σ = 3.5 µm)

considering the mean value (M_I) and the standard deviation (S_I) of the image I_G.

The threshold value is defined by: $T_I = M_I + 6S_I$.

2. Compute the connected components: The connected components labeling is used

to detect connected regions in the image (a digital continuous path exists between

all pairs of points in the same component ‑ the sperm heads). This heuristic

consists of visiting each pixel of the image and creating exterior boundaries using

pixel neighbors, accordingly to a specific type of connectivity.

3. Measure sperm head fluorescence. For each region of interest, identify the

centroid in the fluorescence channel (sperm head) and measure the mean value.

4. Compute the relative positions of the sperm heads within the imaging field, and

assign them to either R1, R2, R3 or R4 concentric regions around the centroid of

the UV flash intensity distribution. The radii of R1, R2, R3 or R4, were 25, 50,

75 and 100 µm, respectively.

5. Repeat steps 1 to 4 in a frame-wise basis.

Step 1 of the algorithm filters out shot noise and atypical values; step 2 divides the

images in N connected components on the location of the sperm heads; step 3

quantitates sperm head fluorescence, and finally step 4 computes the relative sperm

position on the imaging field. A similar approach has been recently used to identify

replication centers of adenoviruses in fluorescence microscopy images (Garces et al.,

2016).

We automatically analyzed up to 267 videos of *S. purpuratus* spermatozoa, each

containing tens of swimming cells, exposed to five distinct speract concentration

gradients.

Computing the dynamics of speract concentration gradients

The dynamics of the chemoattractant gradient was computed using the Green's

function of the diffusion equation:

$$c(r,t) = \frac{C_0}{\sqrt{2\pi\sigma}} e^{-r^2/2\sigma^2} + C_b \qquad\qquad (5)$$

This equation for the concentration tells us that the profile has the form of a

Gaussian. The width of the Gaussian is $\sigma = \sqrt{4D(t + t_0)}$, and hence it increases as the

square root of the time. C_b is the basal concentration of the chemoattractant, D is the

molecular diffusivity.

The speract concentration gradients were generated via the photolysis of 10 nM

caged speract (CS) with a 200 ms UV pulse delivered through each of four different

optical fibers with internal diameters of 200 μm, 600 μm, 2 mm and 4 mm (at two

different positions). Light intensity was normalized dividing each point by the sum of

all points of light intensity for each fiber and multiplying it by the fiber potency

(measured at the back focal plane of the objective) in miliwatts (mW) (**Figure 2 - table**

supplement 1). Each spatial distribution of instantaneously-generated speract

concentration gradient was computed by fitting their corresponding normalized spatial

distribution of UV light (Residual standard error: 2.738 x 10^{-5} on 97 degrees of

freedom), considering an uncaging efficiency of 5%, as reported (Tatsu et al., 2002).

The diffusion coefficient of a molecule is related to its size by the Stokes-Einstein

equation:

$$D = \frac{kT}{6\pi\eta R_h} \qquad\qquad (6)$$

where k is Boltzmann's constant, T is the temperature, η is the viscosity of the

solvent, and R_h is the hydrodynamic radius (Lakowicz, 2006). The hydrodynamic radius

R_h of speract was calculated by modeling the molecules in terms of equivalent

hydrodynamic spheres.

$$R_h = \left(\frac{3M\bar{v}}{4\pi}\right)^{1/3}$$ (7)

where M is the molecular weight, and $\bar{v}$ is the specific gravity (Lakowicz, 2006). The

volume of an equivalent spherical particle is $V_e = 4/3\,\pi R_h^3$. Equations **6** and **7** show

that the radius and diffusion coefficient are weakly dependent on the molecular weight.

The diffusion coefficient of speract has not been measured experimental, nonetheless

it can be estimated following equations 6 and 7. The diffusion coefficient of a similar

chemoattractant molecule, resact (with fourteen amino acids), has been reported, D_{resact}

$= 239 \pm 7\ \mu m^2\ s^{-1}$ (Kashikar et al., 2012). If we consider that speract is a decapeptide,

the 1.4 fold difference in molecular weight between speract and resact would imply a

$(1.4)^{1/3}$ fold difference in their diffusion coefficients, which is close to the experimental

error reported (Kashikar et al., 2012). For the sake of simplicity, the spatio-temporal

dynamics of the distinct instantaneously generated speract gradients was modeled

considering a speract diffusion coefficient of $D_{speract} = 240\ \mu m^2\ s^{-1}$.

The hydrodynamic radius of speract ($R_h = 8.8$ Å) was computed with equation (**6**),

with $D_{speract} = 240\ \mu m^2\ s^{-1}$, $k = 1.38 \times 10^{-23}$ J K^{-1}, $T = 288.15$ K and $\eta = 0.001$ N s /m^2.

Computing $[Ca^{2+}]_i$ dynamics, chemotactic behavior, and the stimulus function of

single spermatozoa

Spermatozoa were tracked semi-automatically by following the head centroid with the

MtrackJ plugin (Meijering et al., 2012) of ImageJ 1.49u. Single cell $[Ca^{2+}]_i$ signals were

computed from the mean value of a 5 x 5 pixels region, centered at each sperm head

along the time.

Chemotactic behavior was quantified using the linear equation chemotaxis index

(*LECI*) employing different temporal windows before (control) and after uncaging

speract. *LECI* is defined as the negative value of the slope (*LECI* = *-l*) of a least square

linear regression *r(t)* = *lt* + *r₀* (Yoshida et al., 2002) where *r(t)* is the sperm head

distance to the center of the speract concentration gradient, *t* is the time after UV pulse

measured in seconds, with *t ∈ [0 - 3; 3.2 - 6.2; 6.2 - 9.2]* seconds. Positive *LECI* indicate

movement towards the chemoattractant source. The center of the speract gradient was

identified as the centroid of the UV flash intensity distribution.

The head position of each spermatozoa was used to compute the local concentration

of speract at $r(x, y)$ over each frame. The stimulus function of single spermatozoa

$s = c(r, t)$ was computed by solving equation (**5**), considering both their swimming

trajectories, and the spatio-temporal evolution of a given speract concentration gradient.

The profiles of UV light were used to compute the initial conditions at $c(r, t_o)$.

The phase- and temporal-shifts between time derivative of the stimulus function

ds/dt and the internal Ca^{2+} oscillator triggered by speract, were computed from their

normalized cross-correlation function.

Analysis of speract induced Ca^{2+} transients with immobilized spermatozoa.

Imaging chambers were prepared by coating cover slips with 50 μg/ml poly-D-

lysine, shaking off excess, and allowing to air-dry. Coated cover slips were then

attached to imaging chambers. Fluo-4 labeled spermatozoa were diluted 1:40 in ASW,

immediately placed into the chambers, and left for 2 min, after which unattached sperm

were removed by washing with ASW. The chambers were then filled with 0.5 ml of

ASW containing 500 nM of caged speract, and mounted in a TC-202 Bipolar

temperature controller (Medical Systems Corp.). Images were collected with Andor iQ

1.7 software (Andor Bioimaging, NC) at 90 fps in full-chip mode, binning 4x4

(observation field of 200 μm x 200 μm). The imaging setup was the same as that used

for swimming spermatozoa. The caged speract was photo-released with a 200 ms UV

pulse delivered through an optical fiber (4 mm internal diameter) coupled to a Xenon

UV lamp (UVICO, Rapp Opto Electronic). The optical fiber was mounted on a

"defocused" configuration to minimized the generation of UV light heterogeneities.

Images were processed off-line using ImageJ 1.4.5s. Overlapping spermatozoa and

any incompletely adhered cells, which moved during the experiment, were ignored.

Fluorescence measurements in individual sperm were made by manually drawing a

region of interest around the flagella with the line tool of ImageJ.

Programs were written in R statistical software.

Theory

Constraints over the detection of chemoattractant concentration fields.

Strategies for chemoattractant sampling based on local concentration gradients require

the concentration to be high enough to ensure that the measured average difference at

two nearby locations is larger than typical fluctuations (Wood et al., 2015). As

stochastic fluctuations limits the precision with which a microorganism can, in a given

time T, determine the concentration of a surrounding chemoattractant (Berg and Purcell,

1977, Vergassola et al., 2007), the rate of encounter between a chemoattractant c

diffusing with effective diffusivity D and its receptor, with a binding site of effective

radius s, is given by:

$$J(\boldsymbol{r}) = 4\pi s D c(\boldsymbol{r}) \tag{8}$$

For reliable assessment of the local concentration $c(r)$, a spermatozoon provided with

N_R number of chemoattractant receptors, collects detection events over time T_{int}. The

average number of detection events will then be $J(r)N_R T_{int}$. Typical stochastic

fluctuations are of the order of the square root of the mean (Berg and Purcell, 1977).

Then the condition for the signal to emerge from the noise reads:

$$\sqrt{N_R s D c T_{int}} \ll 1 \qquad (9)$$

Then, it follows that the spermatozoon will experience an uncertainty $u_1 =$

$(N_R s D c T_{int})^{-1/2}$ (**eq. (1)**), for measuring a variation on the local concentration of

chemoattractant (**Figure 1 - figure supplement 1**).

To reliably measure a concentration gradient of chemoattractant, the difference in

counts collected by the spermatozoon in the interval measurement must be above the

noise level (**Figure 1a**). The corresponding conditions can be stated as (Berg and

Purcell, 1977, Vergassola et al., 2007):

$$\left(v T_{int} \frac{dc}{dr} \right) N_R s D T_{int} \gg \sqrt{N_R s D c T_{int}} \qquad (10)$$

$$v T_{int} \frac{d \log c}{dr} \ll 1 \qquad (11)$$

Here, v is the swimming velocity and dc/dr is the concentration gradient of the

chemoattractant. The first inequality indicates that the signal-to-noise ratio for the

difference in the number of hits experienced by the swimming spermatozoon across the

integration time T_{int} be larger than unity. The second inequality is the requirement of

locality, *i.e.* the change in concentration across the distance spanned during T_{int} needs

to be small compared to the local concentration itself.

The information to determine the direction of the chemoattractant gradient results

from the magnitude difference of the signal at two sampled positions. Then the

condition for the signal to emerge from the noise on a chemoattractant field reads:

$$v^{-1} T^{-3/2} (N_R s D)^{-1/2} c^{1/2} \varepsilon^{-1} \ll 1 \qquad (12)$$

where $\varepsilon = dc/dr$ is the chemoattractant concentration gradient. Then, it follows that

the least fractional error of the chemoattractant gradient direction due to the state of

occupation of receptors is $u_2 = v^{-1} T^{-3/2} (N_R s D)^{-1/2} c^{1/2} \varepsilon^{-1}$ (**Figure 1, eq. (2)**).

699 Assuming an exponentially decaying concentration, reliable integration time T_{int}

700 scales as $exp(r/3\lambda)$ (Vergassola et al., 2007).

701 *Statistical analyses*

702 Data are presented for individual spermatozoa (n) collected from up to three sea

703 urchins. All statistical tests were performed using R software (R Core Team, 2016). The

704 significance level was set at 95%.

706 **Acknowledgements**

707 The authors thank Dr. Tatsu Yoshiro for providing the caged speract, and Drs.

708 Hermes Gadêlha, David Smith and Nina Pastor for stimulating discussions and a critical

709 reading of the manuscript. AG thanks Dr. Manabu Yoshida and Dr. Kaoru Yoshida for

710 feedback about sperm chemotaxis in marine invertebrates, and to the Japan Society for

711 the Promotion of Science (JSPS invitation fellowship for research in Japan to A.G.,

712 short term JSPS/236, ID no. S16172).

714 **Author contributions**

715 A.G., A.D. and I.T. conceived the project; A.G. and V.J.S. performed the

716 experiments; H.R.G., A.G., I.T., V.J.S. and M.V. analyzed the data; A.G., I.T. and

717 H.R.G performed the mathematical model calculations and wrote the corresponding

718 section; A.D., A.G., H.R.G. and I.T. participated in the design and drafting of the

719 manuscript J.C., C.D.W. and C.B. provide feedback for conceptualization and drafting

720 of the manuscript. All authors approved the final version of the article.

722 **Competing interests**

723 The authors declared that no competing interests exist.

Funding

A.D. and A.G. acknowledges grants from the Consejo Nacional de Ciencia y

Tecnología (CONACyT Fronteras 71, CONACyT Ciencia Básica 252213), A.G., A.D.

and C.B to Programa de Apoyo a Proyectos de Investigación e Innovación Tecnológica

UNAM (PAPIIT/DGAPA) (IA202417, IN205516, IN206016 and IN112514

respectively). We acknowledge CONACYT and PAPIIT for fellowships to M.V.P.; the

Spanish Ministry of Economy and Competitiveness Grants No. FIS2013-48444-C2-1-P,

FIS2016-77692-C2-1- P, the Subprogram Ramón-y-Cajal and the Ibero-America

Program-Santander Universities 2015 (I.T.).

Figure 1. Physics of chemoreception. a. Determining the direction of the chemoattractant gradient requires that the signal difference *dc* between two sampled positions *dr* must be greater than the noise. **b-d.** The uncertainty in the determination of the chemoattractant gradient direction, u_2, plotted against the slope of the gradient, *dc/dr*, in log-log scale, for different chemoattractant concentrations, **(b)** *S. purpuratus*, **(c)** *L. pictus*, and **(d)** *A. punctulata* spermatozoa (See **Figure 1 - table supplement 1** for the list of parameter values taken in consideration for panels **b-d**).

Figure 1 - figure supplement 1. Uncertainty for the detection of a homogeneous chemoattractant concentration field. The uncertainty for determining the chemoattractant concentration u_1 versus the receptor number N_R for different concentrations (10^{-6} to 10^{-12} M), for a sampling interval of $T = 0.5$ s **(a)**, or a $T = 60$ s **(b)**. Vertical lines indicate the number of receptors of Sp = *S. purpuratus* (dark green), Lp = *L. pictus* (dark blue) and Ap = *A. punctulata* (black) spermatozoa. Parameters: sampling interval $T = 0.5$ s **(a)** and 60 s **(b)**, speract diffusion coefficient $D = 239$ $\mu m^2 s^{-1}$, hydrodynamic radius of speract (as proxy of receptor pocket size) $R_h = 0.79$ nm (see **figure 1 - figure supplement 2 and figure 1 - table supplement 1**).

Spermatozoan species	S. purpuratus	L. pictus	A. punctulata
N_R [per cell]	2 x 10[4b]	6.3 x 10[4b]	3 x 10[5]
D [cm^{2}s^{-1}]	2.4 x 10[-6c]	2.4 x 10[-6c]	2.4 x 10[-6c]
s [cm]	1.65 x 10[-7c]	1.65 x 10[-7c]	1.65 x 10[-7c]
T [s]	0.39 ± 0.08[a]	0.52 ± 0.22[a]	0.6
v [cms^{-1}]	121 ± 13 x 10[-4a]	144 ± 36 x 10[-4a]	200 x 10[-4]
dr [cm]	28 ± 6 x 10[-4a]	46 ± 14 x 10[-4a]	60 x 10[-4]

Figure 1 - table supplement 1. Parameters of the chemoattractant sampling model for each species. Note that the main differences between species are the number of receptors N_R and the velocity v of the spermatozoon. D diffusion coefficient; s hydrodynamic radius of speract (as proxy of chemoattractant receptor's pocket radius); T sampling time (time to swim half circumference in the boundary close to the water-glass interface); dr = sampling interval (circumference diameter). [a]Measured in this study (mean ± sd; N = 3 sea urchins; n = 495 (*S. purpuratus*), 56 (*L. pictus*) spermatozoa. [b](Nishigaki et al., 2001, Nishigaki and Darszon, 2000). [c]Calculated in this study (see section: ***Computing the dynamics of speract concentration gradients*** in **Materials and Methods**, and **Figure 1 - figure supplement 2**).

Figure 1 - figure supplement 2. Speract and resact modeling. Five speract and resact

structures, respectively, were modeled using PEP-FOLD (Maupetit et al., 2009,

Thevenet et al., 2012, Maupetit et al., 2010), and used to estimate an equivalent

hydrodynamic radius by computing the length of the mayor axis of each model: **a.**

Speract (16.18 ± 0.41 Å). **b.** Resact (18.97 ± 1.79 Å). The estimates of the mayor axis

of the speract model were similar to their corresponding hydrodynamic diameter

estimates ($d = 2R_h = 2(8)$ Å = 1.6 nm) computed with the Stokes-Einstein equation (R_h

= 0.88 nm) (see section: *Computing the dynamics of speract concentration gradients* in

Materials and Methods). Models were visualized with chimera 1.11.2 (Pettersen et al.,

2004).

Figure 2. Screening of speract concentration gradients. a. Radial profile of the UV

light scattered at the glass-liquid interface for each optical fiber (*f1-f5*). **b**. Derivatives of

radial distribution for each optical fiber. **c**. Spatial distribution of the UV flash energy

(left), and typical motility and $[Ca^{2+}]_i$ responses of spermatozoa exposed to different

concentration gradients of speract (right). F-F$_0$ time projections, showing spermatozoa

head fluorescence each 3 s before and after 200 ms UV photoactivation of 10 nM caged

speract in artificial sea water. The pseudo-color scale represents the relative

fluorescence of fluo-4, a Ca^{2+} indicator, showing maximum (red) and minimum (blue)

relative $[Ca^{2+}]_i$. Scale bars of 50 μm.

Figure 2 - figure supplement 1. Automatic segmentation of swimming spermatozoa. a. Working-flow of the algorithm: Video microscopy images were background subtracted by removing the temporal average intensity projection (static fraction) of the first un-stimulated frames (93 frames = 3 seconds) from the whole video (25 seconds). The resulting images were convolved with a low-pass spatial frequency filter to eliminate typical detector shot-noise. The resulting images were thresholded to generate arrays of regions of interest (ROIs), a heuristic search for connected components is then applied to label single ROIs and to assign the corresponding pixels to unique spermatozoa. Scale bar of 15 μm. **b.** The positions of the sperm heads within the imaging field are assigned to either R1, R2, R3 or R4 concentric regions around the centroid of the UV flash intensity distribution. Each ROI was also used to compute the sperm head fluorescence from the raw video microscopy images (as the mean value of the ROI). Scale bar of 50 μm.

	Physical diameter (mm)	UV power at the back focal plane of the objective (mW)*
f1	0.2	0.07
f2	0.6	1.25
f3	2	4.7
f4	4	7.8
f5	4	9.46

Figure 2 - table supplement 1. Physical diameter of the optical fibers, and UV light

power measured at the back focal plane of a 40×/1.3NA oil-immersion objective.

*Typically, there is a 20% loss of light power due to scattering within the optics.

Figure 3. Motility and $[Ca^{2+}]_i$ responses of *S. purpuratus* spermatozoa exposed to

specific concentration gradients of speract. a. Fold change in sperm number, defined

as the number of spermatozoa at the peak of the response (6 s) relative to the mean

number before speract stimulation (0-3 s) (see **figure 3 - figure supplement 1**). **b**.

Relative changes in $[Ca^{2+}]_i$ experienced by spermatozoa at the peak response (6 s) after

speract stimulation. Negative controls for spermatozoa chemotaxis are artificial sea

water with nominal Ca^{2+} (Low Ca^{2+}); and artificial sea water with 40 mM of K^+ (High

K^+). Both experimental conditions prevent chemotactic responses by inhibiting the Ca^{2+}

membrane permeability alterations triggered by speract; the former disrupts the Ca^{2+}

electrochemical gradient, and the later disrupt the K^+ electrochemical gradient required

as electromotive force needed for the opening of Ca^{2+} channels. Number of experiments

(n) on the top of each experimental condition. Each box contains 50% of single events,

the inner lines indicate the median and the error bars delimit the 95% outliers. The

notch display a confidence interval around the median (median +/- 1.57 x $IQRn^{-1/2}$). If

two boxes notches do not overlap, there is evidence that their medians differ (95%

confidence). *Statistical significance, $p < 0.05$; Multiple comparison test after Kruskal-

Wallis. Studies at the population level are presented in **figure 3 - figure supplement 1**,

while studies considering the absolute number of spermatozoa at two periods: 3s and 6s

(3 seconds before and after speract exposure, respectively) are presented in **figure 3 -**

figure supplement 2.

Figure 3 - figure supplement 2. Absolute number of sperm cells, three seconds before and three seconds after speract photo-release through either *f1*, *f2*, *f3*, *f4*, *f5* optical fibers. Absolute number of spermatozoa scored at either R1, R2, R3 or R4 concentric regions around the centroid of the UV flash intensity distribution. Each box contains 50% of single events, the inner lines indicate the median and the error bars delimit the 95% outliers. *Statistical significance, $p < 0.05$; Mann-Whitney test.

$$\xi_{max}= Max(\xi_1, \xi_2, \xi_3, ..., \xi_n)$$

Figure 4. Steep speract gradients attract *S. purpuratus* spermatozoa. a. Dynamics

of the *f2* speract gradient. The blue dashed line ($t_0 = 0$ s) corresponds to a Gaussian

distribution fitted to the UV light profile, and illustrates the putative shape of the

instantaneously-generated speract concentration gradient. Solid black lines illustrate the

temporal evolution of the speract concentration field after t = 1, 2, 3, …, 20 seconds. **b.**

Temporal changes in *f2* speract field computed radially (each 10 µm) from the center of

the gradient. A region where the concentration of speract barely changes is shown in

gray. **c.** Characteristic motility changes of a *S. purpuratus* spermatozoon exposed to the

f2 speract gradient. Solid lines illustrate its swimming trajectory 3 s before (gray) and 6

s after (black) speract exposure. **d**. Spermatozoon head distance to the source of the

speract gradient versus time, calculated from black trajectory in **c**. LECI index is

computed from the least squared linear regression of the head distance to the source, a

negative slope (as shown) indicates chemotaxis (positive LECI). **e**. Stimulus function

computed from the swimming behavior of the spermatozoon in **c**, considering the

dynamics of **a**.

Figure 4 - figure supplement 1. Modeling of the dynamics of speract gradient based on the UV light profile of distinct optical fibers. The radial profiles of the UV light scattered at the glass-liquid interface of *f1*, *f3*, *f4*, *f5* optical fibers are shown in gray. The speract gradient was generated as in **figure 4**, but with the corresponding *f1* (**a**), *f3* (**c**), *f4* (**e**) and *f5* (**g**) optical fibers. *Left panels* - The dynamics of the speract gradient computed as is in **figure 4**. The blue dashed line ($t_0 = 0$) corresponds to a

867   Gaussian distribution fitted to the UV light profile, and illustrates the putative shape of

868   the instantaneously generated speract gradient. Solid black lines illustrate the shape of

869   the speract gradient after t = 1, 2, 3, …, 20 seconds. *Right panels* - Simulated temporal

870   changes in speract concentration gradients of *f1* (**a**), *f3* (**c**), *f4* (**e**) and *f5* (**g**) at each 10

871   radial μm point from the center of the concentration gradient.

Figure 4 - figure supplement 2. Characteristic motility changes of a *S. purpuratus* spermatozoon exposed to *f3 and f4* speract gradients (chemotactic vs non-chemotactic response). Panels **a**, **b** and **c** show single cell responses to the *f3* speract gradient (chemotactic); and panels **d**, **e** and **f** to the *f4* speract gradient (non-chemotactic). **a, d.** Solid lines illustrate the spermatozoon swimming trajectory 3 s before (gray) and 6 s after (black) speract gradient exposure. **b, e**. Spermatozoon head distance to the source of the speract gradient versus time calculated from **a** and **d**, respectively. **b, e**. Stimulus function computed from **a** and **d**, considering the spatio-temporal dynamics of speract computed for the *f3* and *f4* gradients, respectively.

Figure 5. *S. purpuratus* **spermatozoa selectively experience chemotaxis towards speract gradients.** Chemotactic behavior quantified using the linear equation chemotaxis index (*LECI*) before (**a**), as control experiment, and after exposure to either *f1*, *f2*, *f3*, *f4* or *f5* speract gradients (**b**, **c**). The color scale indicates the mean *LECI* value calculated from tens of *S. purpuratus* spermatozoa located in either R1, R2, R3, or R4 regions prior speract photo-release (3 s after UV light irradiation (**b**), or 6 s after UV light exposure (**c**)). Positive LECIs (shown in red) indicate movement towards the chemoattractant source. Note that spermatozoa that were in R3 and R4 distal regions prior speract gradient exposure experience chemotaxis only in response to *f2*, *f3*, and *f5* speract gradients (**b**, **c**). Numerals indicate the number of spermatozoa analyzed in each experimental condition. *Statistical significance, $p < 0.05$; Student's t-test.

Figure 6. The slope of the speract gradient generates a frequency-locking phenomenon between the stimulus function and the internal Ca^{2+} oscillator triaggered by speract. a. Maximum relative slopes (ξ_{max}) experienced by *S. purpuratus* (Sp) when exposed to *f1, f2, f3, f4, f5* speract gradients. The maximum relative slopes experienced by *L. pictus* spermatozoa (Lp) towards *f4* experimental regime are also shown. Note that ξ_{max} for *f2, f3*, and *f5*, are up to 2-3 times greater than in *f4*, regardless of the species. **b.** Arnold's tongue indicating the difference in intrinsic frequency of the internal Ca^{2+} oscillator of *S. purpuratus* spermatozoa, just before and after the speract gradient exposure. **c**. Phase difference between the time derivative of the stimulus function and the internal Ca^{2+} oscillator of *S. purpuratus* spermatozoa, obtained by computing the cross-correlation function between both time series (**Figure 6 - figure**

supplement 2). **d**. Phase difference between the time derivative of the stimulus function

and the internal Ca^{2+} oscillator of *S. purpuratus* spermatozoa expressed in temporal

delays. **b-d**. Gray points represent the collapsed data of all *f1, f2, f3, f4, f5* experimental

regimes. Red, black and blue points indicate chemotactic spermatozoa (LECI > 1),

located in R3, and R4 regions just before the speract gradient is established under *f2, f3*

and *f5* experimental regimes, respectively. Magenta lines represent the transition

boundary ($\gamma_{min} = \overline{\xi^*_{max}} \sim 2.7 \times 10^{-4}$) below which synchrony is not observed, obtained

from the theoretical estimates (black curves) of panels **c** and **d**. Green lines indicate

confidence intervals.

Figure 6 - figure supplement 1. Speract induces Ca^{2+} oscillations in immobilized *S.*

***purpuratus* spermatozoa.** Spermatozoa were immobilized on cover slips coated with

poly-D-lysine (see **Materials and Methods**), and ASW containing 500 nM caged

speract added. Recordings were performed 3s before and during 6s after 0.2s of UV

irradiation. *f4* optical fiber was used for the UV light path, to generate the speract

concentration gradient. The optical fiber was mounted on a "defocused" configuration

to minimize UV light heterogeneities producing photoactivated speract gradients. Time

traces indicate the [Ca^{2+}]$_i$ of selected spermatozoa of **Movie 8**. Note that the photo-

release of speract induces a train of [Ca^{2+}]$_i$ episodes in immobilized spermatozoa, hence

provides evidences for the presence of an internal Ca^{2+} oscillator triggered by speract.

Figure 6 - figure supplement 2. Cross-correlation analysis of $[Ca^{2+}]_i$ and stimulus

function derivative (ds) signals. Representative examples of $[Ca^{2+}]_i$ and the derivate of

the stimulus function (ds) were plotted and then analyzed by cross-correlation analysis

(CCF). Examples of a pair of spermatozoa for the two principal chemotactic gradients

(*f2* and *f3*) are shown. **a, b.** Representative examples of two spermatozoa in *f2* gradient.

c, d. Representative examples of two spermatozoa in *f3* gradient.

Movies

Movie 1. Typical motility and Ca^{2+} responses of *S. purpuratus* spermatozoa towards *f2*-generated speract concentration gradient. Spermatozoa swimming in artificial sea water containing 10 nM caged speract, 3 s before and 5 s after 200 ms UV irradiation. An optical fiber of 0.6 mm internal diameter was used for the UV light path to generate the speract concentration gradient. Real time: 31 frames s^{-1}, 40x /1.3NA oil-immersion objective. Note that spermatozoa located at R2, R3 and R4 regions prior speract exposure swim up the speract concentration gradient, towards the center of the imaging field. The pseudo-color scale represents the relative fluorescence of fluo-4, a Ca^{2+} indicator, showing maximum (red) and minimum (blue) relative [Ca^{2+}]$_i$. Six *S. purpuratus* spermatozoa were manually tracked for visualization purposes. Scale bar of 50 µm.

https://www.dropbox.com/s/v1h4kx6oqxc0f5l/Movie%202.avi?dl=0

Movie 2. Typical motility and Ca^{2+} responses of *S. purpuratus* spermatozoa

towards *f3*-generated speract concentration gradient. An optical fiber of 2 mm

internal diameter was used for the UV light path to generate the speract concentration

gradient. Other imaging conditions were set up as for Movie 1. Note that spermatozoa

located at R2, R3 and R4 regions prior to speract exposure swim up the speract

concentration gradient, towards the center of the imaging field. The pseudo-color scale

represents the relative fluorescence of fluo-4, a Ca^{2+} indicator, showing maximum (red)

and minimum (blue) relative [Ca^{2+}]$_i$. Six *S. purpuratus* spermatozoa were manually

tracked for visualization purposes. Scale bar of 50 µm.

Movie 3. Chemotaxis of *S. purpuratus* spermatozoa requires extracellular calcium.

Spermatozoa swimming in artificial sea water with nominal calcium containing 10 nM

caged speract 3 s before and 5 s after exposure to 200 ms UV light. Nominal calcium

disrupts the electrochemical gradient required for Ca^{2+} influx, hence blocking the

triggering of the internal Ca^{2+} oscillator by speract. The *f2* fiber (0.6 mm diameter) was

used to uncage speract in this control. Other imaging conditions were set up as for

Movie 1. Note that spermatozoa re-located after speract uncaging but they fail to

experience the Ca^{2+}-driven motility alteration triggered by speract. As a consequence

they fail to experience chemotaxis (compare with Movie 1). The pseudo-color scale

represents the relative fluorescence of fluo-4, a Ca^{2+} indicator, showing maximum (red)

and minimum (blue) relative $[Ca^{2+}]_i$. Six *S. purpuratus* spermatozoa were manually

tracked for visualization purposes. Scale bar of 50 µm.

Movie 4. Disrupting the K$^+$ electrochemical gradient blocks chemotaxis of *S.*

***purpuratus* spermatozoa.** Cells were swimming in artificial sea water containing 40

mM of KCl, and 10 nM caged speract 3 s before and 5 s after exposure to 200 ms UV

light. High K$^+$ in the ASW blocks the hyperpolarization required for opening Ca^{2+}

channels, and hence prevents the triggering of the internal Ca^{2+} oscillator by speract

exposure. The *f2* fiber (0.6 mm diameter) was used to uncage speract in this control.

Other imaging conditions were set up as for Movie 1. Note that spermatozoa re-located

after speract uncaging but they fail to experience the Ca^{2+}-driven motility alteration

triggered by speract, and thus they fail to experience chemotaxis (compare with Movie

1). The pseudo-color scale represents the relative fluorescence of fluo-4, a Ca^{2+}

indicator, showing maximum (red) and minimum (blue) relative [Ca^{2+}]$_i$. Six *S.*

purpuratus spermatozoa were manually tracked for visualization purposes. Scale bar of

50 µm.

https://www.dropbox.com/s/l3l7nmh1yw9juov/Movie%205.avi?dl=0

Movie 5. Typical motility and Ca^{2+} responses of *S. purpuratus* spermatozoa

towards *f1*-generated speract concentration gradient. An optical fiber of 0.2 mm

internal diameter was used for the UV light path to generate the speract concentration

gradient. Other imaging conditions were set up as for Movie 1. Note that some

spermatozoa re-located after speract uncaging but they fail to experience chemotaxis

(compare with Movie 1). The pseudo-color scale represents the relative fluorescence of

fluo-4, a Ca^{2+} indicator, showing maximum (red) and minimum (blue) relative $[Ca^{2+}]_i$.

Six *S. purpuratus* spermatozoa were manually tracked for visualization purposes. Scale

bar of 50 µm.

Movie 6. Typical motility and Ca^{2+} responses of *S. purpuratus* spermatozoa

towards *ƒ4*-generated speract concentration gradient. An optical fiber of 4 mm

internal diameter was used for the UV light path to generate the speract concentration

gradient. Other imaging conditions were set up as for Movie 1. Note that spermatozoa

re-located after speract uncaging but they fail to experience chemotaxis (compare with

Movie 1). The pseudo-color scale represents the relative fluorescence of fluo-4, a Ca^{2+}

indicator, showing maximum (red) and minimum (blue) relative $[Ca^{2+}]_i$. Six *S.*

purpuratus spermatozoa were manually tracked for visualization purposes. Scale bar of

50 µm.

Movie 7. Typical motility and Ca^{2+} responses of *S. purpuratus* spermatozoa

towards *f*5-generated speract concentration gradient. An optical fiber of 4 mm

internal diameter was used for the UV light path to generate the speract concentration

gradient. Other imaging conditions were set up as for Movie 1. Note that spermatozoa

located at R2, R3 and R4 regions prior to speract exposure swim up the speract

concentration gradient, towards the center of the imaging field. The pseudo-color scale

represents the relative fluorescence of fluo-4, a Ca^{2+} indicator, showing maximum (red)

and minimum (blue) relative $[Ca^{2+}]_i$. Six *S. purpuratus* spermatozoa were manually

tracked for visualization purposes. Scale bar of 50 µm.

1015

1016

Movie 8. Photo-release of caged speract induces Ca^{2+} oscillations in immobilized *S.*

***purpuratus* spermatozoa.** Spermatozoa were immobilized, by coating the cover slip

with a thin layer of poly-D-lysine, in artificial sea water containing 500 nM caged

speract, 3 s before and during 6 s after 0.2 s of UV irradiation. The *f4* optical fiber was

used for the UV light path to generate the speract concentration gradient. The optical

fiber was mounted on a "defocused" configuration to minimize the generation of UV

light heterogeneities. 93 frames s^{-1}, 40x/1.3NA oil-immersion objective, 4x4 binning.

The pseudo-color scale represents the relative fluorescence of fluo-4, a Ca^{2+} indicator,

showing maximum (red) and minimum (blue) relative $[Ca^{2+}]_i$. The brightness and

contrast scale was adjusted for better visualization of $[Ca^{2+}]_i$ transients in the sperm

flagella (as a consequence some heads look artificially oversaturated, however no

fluorescence saturation was observed in the raw data).

1029

1030

References

2011. R: A Language and Environment for Statistical Computing. Vienna, Austria : the R Foundation for Statistical Computing: R Development Core Team.

AGUILERA, L. U., GALINDO, B. E., SANCHEZ, D. & SANTILLAN, M. 2012. What is the core oscillator in the speract-activated pathway of the Strongylocentrotus purpuratus sperm flagellum? *Biophys J,* 102**,** 2481-8.

ALVAREZ, L., DAI, L., FRIEDRICH, B. M., KASHIKAR, N. D., GREGOR, I., PASCAL, R. & KAUPP, U. B. 2012. The rate of change in Ca(2+) concentration controls sperm chemotaxis. *J Cell Biol,* 196**,** 653-63.

AREF, H., BLAKE, J. R., BUDIŠIĆ, M., CARTWRIGHT, J. H. E., CLERCX, H. J. H., FEUDEL, U., GOLESTANIAN, R., GOUILLART, E., LE GUER, Y., VAN HEIJST, G. F., KRASNOPOLSKAYA, T. S., MACKAY, R. S., MELESHKO, V. V., METCALFE, G., MEZIĆ, I., DE MOURA, A. P. S., EL OMARI, K., PIRO, O., SPEETJENS, M. F. M., STURMAN, R., THIFFEAULT, J. L. & TUVAL, I. 2014. Frontiers of chaotic advection. *Reviews of Modern Physics,* 1403**,** 61.

BATCHELOR, G. K. 2006. Small-scale variation of convected quantities like temperature in turbulent fluid Part 1. General discussion and the case of small conductivity. *Journal of Fluid Mechanics,* 5**,** 113.

BERG, H. C. & PURCELL, E. M. 1977. Physics of chemoreception. *Biophys J,* 20**,** 193-219.

BOHMER, M., VAN, Q., WEYAND, I., HAGEN, V., BEYERMANN, M., MATSUMOTO, M., HOSHI, M., HILDEBRAND, E. & KAUPP, U. B. 2005. Ca2+ spikes in the flagellum control chemotactic behavior of sperm. *EMBO J,* 24**,** 2741-52.

BROKAW, C. J. 1979. Calcium-induced asymmetrical beating of triton-demembranated sea urchin sperm flagella. *J Cell Biol,* 82**,** 401-11.

COOK, S. P., BROKAW, C. J., MULLER, C. H. & BABCOCK, D. F. 1994. Sperm chemotaxis: egg peptides control cytosolic calcium to regulate flagellar responses. *Dev Biol,* 165**,** 10-9.

DARSZON, A., GUERRERO, A., GALINDO, B. E., NISHIGAKI, T. & WOOD, C. D. 2008. Sperm-activating peptides in the regulation of ion fluxes, signal transduction and motility. *Int J Dev Biol,* 52**,** 595-606.

DUSENBERY, D. B. 2011. *Living at micro scale: The unexpected physics of being small*, Harvard University Press.

ESPINAL, J., ALDANA, M., GUERRERO, A., WOOD, C., DARSZON, A. & MARTINEZ-MEKLER, G. 2011. Discrete dynamics model for the speract-activated Ca2+ signaling network relevant to sperm motility. *PLoS One,* 6**,** e22619.

FRIEDRICH, B. M. & JÜLICHER, F. 2007. Chemotaxis of sperm cells. *Proc Natl Acad Sci U S A,* 104**,** 13256-61.

FRIEDRICH, B. M. & JÜLICHER, F. 2008. The stochastic dance of circling sperm cells: sperm chemotaxis in the plane. *New Journal of Physics,* 10**,** 123025.

FRIEDRICH, B. M. & JÜLICHER, F. 2009. Steering chiral swimmers along noisy helical paths. *Phys Rev Lett,* 103**,** 068102.

GARCES, Y., GUERRERO, A., HIDALGO, P., LOPEZ, R. E., WOOD, C. D., GONZALEZ, R. A. & RENDON-MANCHA, J. M. 2016. Automatic detection and measurement of viral replication compartments by ellipse adjustment. *Sci Rep,* 6**,** 36505.

GUERRERO, A., NISHIGAKI, T., CARNEIRO, J., YOSHIRO, T., WOOD, C. D. &
DARSZON, A. 2010a. Tuning sperm chemotaxis by calcium burst timing. *Dev*
Biol, 344**,** 52-65.
GUERRERO, A., WOOD, C. D., NISHIGAKI, T., CARNEIRO, J. & DARSZON, A.
2010b. Tuning sperm chemotaxis. *Biochem Soc Trans,* 38**,** 1270-4.
HANSBROUGH, J. R. & GARBERS, D. L. 1981. Speract. Purification and
characterization of a peptide associated with eggs that activates spermatozoa. *J*
Biol Chem, 256**,** 1447-52.
HUSSAIN, Y. H., SADILEK, M., SALAD, S., ZIMMER, R. K. & RIFFELL, J. A.
2017. Individual female differences in chemoattractant production change the
scale of sea urchin gamete interactions. *Dev Biol,* 422**,** 186-197.
JIKELI, J. F., ALVAREZ, L., FRIEDRICH, B. M., WILSON, L. G., PASCAL, R.,
COLIN, R., PICHLO, M., RENNHACK, A., BRENKER, C. & KAUPP, U. B.
2015. Sperm navigation along helical paths in 3D chemoattractant landscapes.
Nat Commun, 6**,** 7985.
JIMÉNEZ, J. 1997. Oceanic turbulence at millimeter scales. *Scientia Marina,* 61**,** 47-56.
KASHIKAR, N. D., ALVAREZ, L., SEIFERT, R., GREGOR, I., JACKLE, O.,
BEYERMANN, M., KRAUSE, E. & KAUPP, U. B. 2012. Temporal sampling,
resetting, and adaptation orchestrate gradient sensing in sperm. *J Cell Biol,* 198**,**
1075-91.
KAUPP, U. B. 2012. 100 years of sperm chemotaxis. *J Gen Physiol,* 140**,** 583-6.
KAUPP, U. B., KASHIKAR, N. D. & WEYAND, I. 2008. Mechanisms of sperm
chemotaxis. *Annu Rev Physiol,* 70**,** 93-117.
KAUPP, U. B., SOLZIN, J., HILDEBRAND, E., BROWN, J. E., HELBIG, A.,
HAGEN, V., BEYERMANN, M., PAMPALONI, F. & WEYAND, I. 2003. The
signal flow and motor response controling chemotaxis of sea urchin sperm. *Nat*
Cell Biol, 5**,** 109-17.
LAKOWICZ, J. R. 2006. *Principles of Fluorescence Spectroscopy*, Springer US.
LILLIE, F. R. 1913. The mechanism of fertilization. *Science,* 38**,** 524-528.
LOTTERHOS, K. E. L., D. R 2010. *Gamete release and spawning behavior in*
broadcast spawning marine invertebrates, New York, NY, Oxford University
Press.
MAUPETIT, J., DERREUMAUX, P. & TUFFERY, P. 2009. PEP-FOLD: an online
resource for de novo peptide structure prediction. *Nucleic Acids Res,* 37**,** W498-
503.
MAUPETIT, J., DERREUMAUX, P. & TUFFERY, P. 2010. A fast method for large-
scale de novo peptide and miniprotein structure prediction. *J Comput Chem,* 31**,**
726-38.
MEAD, K. S. & DENNY, M. W. 1995. The effects of hydrodynamic shear stress on
fertilization and early development of the purple sea urchin Strongylocentrotus
purpuratus. *Biol Bull,* 188**,** 46-56.
MEIJERING, E., DZYUBACHYK, O. & SMAL, I. 2012. Methods for cell and particle
tracking. *Methods Enzymol,* 504**,** 183-200.
MILLER, R. L. 1985. Sperm Chemo-Orientation in the Metazoa. *In:* METZ, C. B. &
MONROY, A. (eds.) *Biology of Fertilization.* New York: Academic Press.
NISHIGAKI, T. & DARSZON, A. 2000. Real-time measurements of the interactions
between fluorescent speract and its sperm receptor. *Dev Biol,* 223**,** 17-26.
NISHIGAKI, T., ZAMUDIO, F. Z., POSSANI, L. D. & DARSZON, A. 2001. Time-
resolved sperm responses to an egg peptide measured by stopped-flow
fluorometry. *Biochem Biophys Res Commun,* 284**,** 531-5.

PETTERSEN, E. F., GODDARD, T. D., HUANG, C. C., COUCH, G. S., GREENBLATT, D. M., MENG, E. C. & FERRIN, T. E. 2004. UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem,* 25**,** 1605-12.

PFEFFER, W. 1884. Locomotorische Richtungsbewegungen durch chemische Reize. *In Untersuchungen aus dem botanischen Institut zu Tübingen,* 1**,** 363-482.

PICHLO, M., BUNGERT-PLUMKE, S., WEYAND, I., SEIFERT, R., BONIGK, W., STRUNKER, T., KASHIKAR, N. D., GOODWIN, N., MULLER, A., PELZER, P., VAN, Q., ENDERLEIN, J., KLEMM, C., KRAUSE, E., TROTSCHEL, C., POETSCH, A., KREMMER, E., KAUPP, U. B., KORSCHEN, H. G. & COLLIENNE, U. 2014. High density and ligand affinity confer ultrasensitive signal detection by a guanylyl cyclase chemoreceptor. *J Cell Biol,* 206**,** 541-57.

PIKOVSKY, A., ROSENBLUM, M. & KURTHS, J. 2001. *Synchronization. A Universal Concept in Nonlinear Sciences*, Cambridge University Press; 1 edition.

RIFFELL, J. A. & ZIMMER, R. K. 2007. Sex and flow: the consequences of fluid shear for sperm-egg interactions. *J Exp Biol,* 210**,** 3644-60.

SCHNEIDER, C. A., RASBAND, W. S. & ELICEIRI, K. W. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods,* 9**,** 671-5.

SHIBA, K., BABA, S. A., INOUE, T. & YOSHIDA, M. 2008. Ca2+ bursts occur around a local minimal concentration of attractant and trigger sperm chemotactic response. *Proc Natl Acad Sci U S A,* 105**,** 19312-7.

SHIMOMURA, H. & GARBERS, D. L. 1986. Differential effects of resact analogues on sperm respiration rates and cyclic nucleotide concentrations. *Biochemistry,* 25**,** 3405-10.

SMITH, A. C. & GARBERS, D. L. 1983. The binding of an 125I-speract analogue to spermatozoa. *In:* LENNON, D. L., STRATMAN, F. W. & ZAHLTEN, R. N., eds. Biochemistry of Metabolic Processes, 1982 Madison, Wisconsin, U.S.A. Amsterdam, New York, U.S.A.: Elsevier, 496.

STRUNKER, T., ALVAREZ, L. & KAUPP, U. B. 2015. At the physical limit - chemosensation in sperm. *Curr Opin Neurobiol,* 34**,** 110-6.

SUZUKI, N. 1995. Structure, function and biosynthesis of sperm-activating peptides and fucose sulfate glycoconjugate in the extracellular coat of sea urchin eggs. *Zoolog Sci,* 12**,** 13-27.

TATSU, Y., NISHIGAKI, T., DARSZON, A. & YUMOTO, N. 2002. A caged sperm-activating peptide that has a photocleavable protecting group on the backbone amide. *FEBS Lett,* 525**,** 20-4.

THEVENET, P., SHEN, Y., MAUPETIT, J., GUYON, F., DERREUMAUX, P. & TUFFERY, P. 2012. PEP-FOLD: an updated de novo structure prediction server for both linear and disulfide bonded cyclic peptides. *Nucleic Acids Res,* 40**,** W288-93.

THORPE, S. A. 2007. *An Introduction to Ocean Turbulence*, Cambridge University Press.

VERGASSOLA, M., VILLERMAUX, E. & SHRAIMAN, B. I. 2007. 'Infotaxis' as a strategy for searching without gradients. *Nature,* 445**,** 406-9.

WOOD, C. D., DARSZON, A. & WHITAKER, M. 2003. Speract induces calcium oscillations in the sperm tail. *J Cell Biol,* 161**,** 89-101.

WOOD, C. D., GUERRERO, A., PRIEGO-ESPINOSA, D. A., MARTÍNEZ-MEKLER, G., CARNEIRO, J. & DARSZON, A. 2015. Sea Urchin Sperm Chemotaxis. *In:*

COSSON, J. J. (ed.) *Flagellar Mechanics and Sperm Guidance.* Czech
Republic: Bentham Science Publishers.
WOOD, C. D., NISHIGAKI, T., FURUTA, T., BABA, S. A. & DARSZON, A. 2005.
Real-time analysis of the role of Ca(2+) in flagellar movement and motility in
single sea urchin sperm. *J Cell Biol,* 169**,** 725-31.
WOOD, C. D., NISHIGAKI, T., TATSU, Y., YUMOTO, N., BABA, S. A.,
WHITAKER, M. & DARSZON, A. 2007. Altering the speract-induced ion
permeability changes that generate flagellar Ca2+ spikes regulates their kinetics
and sea urchin sperm motility. *Dev Biol,* 306**,** 525-37.
YOSHIDA, M., MURATA, M., INABA, K. & MORISAWA, M. 2002. A
chemoattractant for ascidian spermatozoa is a sulfated steroid. *Proc Natl Acad
Sci U S A,* 99**,** 14831-6.
ZIMMER, R. K. & RIFFELL, J. A. 2011. Sperm chemotaxis, fluid shear, and the
evolution of sexual reproduction. *Proc Natl Acad Sci U S A,* 108**,** 13200-5.
1193