Sperm chemotaxis is driven by the slope of the chemoattractant

concentration field

Ramírez-Gómez H.V.[1], Jimenez Sabinina V.[2], Velázquez-Pérez M.[1], Beltrán C.[1],

Carneiro J.[3], Wood C.D.[4], Tuval I.[5], Darszon A.[1, *], Guerrero A.[4, *].

[1] Departamento de Genética del Desarrollo y Fisiología Molecular, Instituto de

Biotecnología, Universidad Nacional Autónoma de México (UNAM), Avenida

Universidad 2001, Colonia Chamilpa, Cuernavaca, Morelos, 62210, México.

[2] Cell Biology and Biophysics Unit, European Molecular Biology Laboratory (EMBL),

Heidelberg, Germany.

[3] Instituto Gulbenkian de Ciência (IGC), Rua da Quinta Grande, 6 2780-156, Oeiras,

Portugal.

[4] Laboratorio Nacional de Microscopía Avanzada, Instituto de Biotecnología,

Universidad Nacional Autónoma de México (UNAM), Cuernavaca, Morelos, 62210,

México.

[5] Mediterranean Institute for Advanced Studies, IMEDEA (CSIC-UIB), Esporles,

Balearic Islands, Spain.

[*] Correspondence and requests should be addressed to A.G. or A.D. (email:

adanog@ibt.unam.mx, darszon@ibt.unam.mx).

Abstract

Spermatozoa of marine invertebrates are attracted to their conspecific female

gamete by diffusive molecules, called chemoattractants, released from the egg invest-

ments in a process known as chemotaxis. The information from the egg chemoattractant

concentration field is decoded into intracellular Ca^{2+} concentration ($[Ca^{2+}]_i$) changes

that regulate the internal motors that shape the flagellum as it beats. By modeling sea

urchin species-specific differences in sperm chemoattractant-receptor characteristics

we conclude that receptor density constrains the steepness of the chemoattractant con-

centration gradient detectable by spermatozoa. Through analyzing different chemoat-

tractant gradient forms, we demonstrate for the first time that *Strongylocentrotus pur-*

puratus sperm are chemotactic and this response is consistent with frequency entrain-

ment of two coupled physiological oscillators: i) the stimulus function and ii) the $[Ca^{2+}]_i$

changes. We demonstrate that the slope of the chemoattractant gradients provides the

coupling force between both oscillators, arising as a fundamental requirement for sperm

chemotaxis.

Introduction

Broadcast spawning organisms, such as marine invertebrates, release their gametes into open water, where they are often subject to extensive dilution that reduces the probability of gamete encounter (Lotterhos et al., 2010). In many marine organisms, female gametes release diffusible molecules that attract homologous spermatozoa (Lillie, 1913; Miller, 1985; Suzuki, 1995), which detect and respond to chemoattractant concentration gradients by swimming toward the gradient source: the egg. Although it was in bracken ferns where sperm chemotaxis was first identified (Pfeffer, 1884), sea urchins are currently the best-characterized model system for studying sperm chemotaxis at a molecular level (Alvarez et al., 2012; Cook et al., 1994; Darszon et al., 2008; Strünker et al., 2015; Wood et al., 2015).

The sea urchin egg is surrounded by an extracellular matrix which contains sperm-activating peptides (SAPs) that modulate sperm motility through altering intracellular Ca^{2+} concentration ($[Ca^{2+}]_i$) and other signaling intermediates (Darszon et al., 2008; Suzuki, 1995). The biochemical signals triggered by SAPs guide the sperm trajectory towards the egg.

The decapeptide speract is one of best characterized members of the SAP family due to its powerful stimulating effect on metabolism, permeability and motility in *Strongylocentrotus purpuratus* and *Lytechinus pictus* spermatozoa. The binding of speract to its receptor, located in the flagellar plasma membrane, triggers a train of $[Ca^{2+}]_i$ increases in immobilized spermatozoa of both species (Wood et al., 2003). This calcium signal was proposed to regulate the activity of dynein motor proteins in the flagellum, and thus potentially modulate the trajectory of free-swimming spermatozoa (Brokaw, 1979; Mizuno et al., 2017).

A direct link between $[Ca^{2+}]_i$ signaling and sperm motility was established through

the use of optochemical techniques to rapidly, and non-turbulently, expose swimming

sea urchin spermatozoa to their conspecific attractant in a well-controlled experimental

regime (Böhmer et al., 2005; Wood et al., 2005). Currently, it is well established that

the transient $[Ca^{2+}]_i$ increases triggered by chemoattractants produce a sequence of

turns and straight swimming episodes (the "turn-and-run" response), where each

turning event results from a rapid increase in the $[Ca^{2+}]_i$ (Alvarez et al., 2012; Böhmer

et al., 2005; Shiba et al., 2008; Wood et al., 2005). The turn-and-run response seems to

be a general requirement for sperm chemotaxis in sea urchins, however it is not

sufficient on its own to produce a chemotactic response (Guerrero et al., 2010; Strünker

et al., 2015; Wood et al., 2007, 2005).

In spite of 30 years of research since speract's isolation from *S. purpuratus* oocytes

(Hansbrough et al., 1981; Suzuki, 1995), chemotaxis of *S. purpuratus* sperm in the

presence of this peptide has not yet been demonstrated (Cook et al., 1994; Darszon et

al., 2008; Guerrero et al., 2010; Kaupp, 2012; Miller, 1985; Wood et al., 2015). A

comparison between individual *L. pictus* and *S. purpuratus* sperm responses to a

specific chemoattractant concentration gradient generated by photoactivating caged

speract (CS) revealed that only *L. pictus* spermatozoa exhibit chemotaxis under these

conditions (Guerrero et al., 2010). In that study, *L. pictus* spermatozoa experience

$[Ca^{2+}]_i$ fluctuations and pronounced turns while swimming in descending speract

gradients, that result in spermatozoa reorienting their swimming behavior along the

positive chemoattractant concentration gradient. In contrast, *S. purpuratus* spermatozoa

experience similar trains of $[Ca^{2+}]_i$ fluctuations that in turn drive them to relocate, but

with no preference towards the center of the chemoattractant gradient (Guerrero et al.,

2010).

In the present work, we investigate boundaries that limit sperm chemotaxis of marine

invertebrates. Particularly, we examined whether the chemoattractant concentration

gradient must have a minimum steepness to provoke an adequate, chemotactic sperm

motility response. Previous studies of chemotactic amoebas crawling up a gradient of

cAMP, have shown that the slope of the chemical concentration gradient works as a

determinant factor in chemotaxis of this species, where the signal-to-noise relationship

of stimulus to the gradient detection mechanism imposes a limit for chemotaxis

(Amselem et al., 2012). In addition, recent theoretical studies by Kromer and colleagues

have shown that, in marine invertebrates, sperm chemotaxis operates efficiently within

a boundary defined by the signal-to-noise ratio of detecting ligands within a

chemoattractant concentration gradient (Kromer et al., 2018).

If certain, this detection limit may have prevented the observation and

characterization of chemotactic responses on *S. purpuratus* spermatozoa to date. In this

study, we identify the boundaries for detecting chemotactic signals of *S. purpuratus*

spermatozoa, and show that sperm chemotaxis arises only when sperm are exposed to

much steeper speract concentration gradients than those previously employed by

Guerrero et al. (2010). Furthermore, we examined the coupling between the recruitment

of speract molecules during sperm swimming (i.e. stimulus function) and the internal

Ca^{2+} oscillator, and demonstrate that sperm chemotaxis arises through coupling of these

physiological oscillators.

Results

Species-specific differences in chemoattractant-receptor binding rates:

chemoattractant sensing is limited by receptor density in S. purpuratus spermatozoa

Spermatozoa measure the concentration and the changes in concentration of egg-released chemoattractant during their journey. Cells detect chemoattractant molecules in the extracellular media by integrating chemoattractant-receptor binding events. A spermatozoon moving in a medium where the chemoattractant concentration is isotropic will collect stochastic chemoattractant-receptor binding events with a rate J, according to equation **(1)** (**Figure 1a**).

$$J = 4\pi D a \bar{c} \frac{N}{N + \pi a/s} = J_{max} \frac{N}{N + \pi a/s} \tag{1}$$

Where D is the diffusion coefficient of the chemoattractant, a is the radius of the cell, $\bar{c}$ is the mean chemoattractant concentration, N is the number of receptor molecules on the cell surface, s is the effective radius of the chemoattractant molecule, J_{max} is the maximal flux that the cell can experience, and $\frac{N}{N + \pi a/s}$ is the probability that a molecule that has collided with the cell will find a receptor (Berg and Purcell, 1977). The quantity $\pi a/s$ is the number of receptors that allows half maximal binding rate for any concentration of chemoattractant, which is hereafter denoted as $N_{1/2}$ (Berg and Purcell, 1977) (see ***1.1. On the estimate of maximal chemoattractant absorption*** in **supplementary material)**.

The expression above was used by Berg and Purcell (1977) to conclude that the chemoattractant binding and adsorption rate saturate as a function of the density of receptors, becoming diffusion limited, i.e. when $N \gg N_{1/2} = \pi a/s$ the chemoattractant adsorption flux becomes $J \cong J_{max}$ (see ***1.1. On the estimate of maximal chemoattractant absorption*** in **supplementary material**). If the density of chemoattractant receptor is such that spermatozoa of the different species operate under

this saturated or perfect absorber regime, then any postulated species-specific

differences would have to be downstream.

In **Table SI** we list the biophysical parameters considered for calculating the

species-specific rate of binding as a function of the chemoattractant concentration. The

different functions of the receptor density and the species receptor density are depicted

in **figure S1**. Our calculations (see **supplementary material**, section ***1.1. On the***

estimate of maximal chemoattractant absorption) indicate that only *S. purpuratus*

spermatozoa operate in a regime for which the rate of chemoattractant uptake is limited

by receptor density, therefore it cannot be considered as a perfect absorber. The actual

number of speract receptors for this species is approximately $2x10^4$ per sperm cell

which is fewer than the estimate of $N_{1/2} \sim 3x10^4$ (**Table SI**). In contrast, *L. pictus* and

A. punctulata spermatozoa seem to approximate towards operating as perfect absorbers

(**Figure S1** and **Table SI**). Both observations hold when considering the cylindrical

geometry of the sperm flagellum (see section ***1.1. On the estimate of maximal***

chemoattractant absorption in **supplementary material**). Therefore, there are

meaningful species-specific differences in chemoattractant receptor density which

could by themselves explain differences in chemotactic behavior.

Receptor density constrains the chemoattractant concentration gradient detectable

by spermatozoa

A functional chemotactic signaling system must remain unresponsive while the

cell swims through an isotropic chemoattractant concentration field and must trigger a

directional motility response if the cell moves across a concentration gradient (**Figure**

1a-c). This absolute prerequisite of the signaling system defines the minimal

quantitative constraints for reliable detection of a gradient and therefore for chemotaxis.

A cell moving along a circular trajectory in an isotropic chemoattractant field

(**Figure 1a**) will collect a random number of chemoattractant-receptor binding events

during the half revolution time Δt, that has a Poisson distribution with mean $J\Delta t$ and

standard deviation $\sqrt{J\Delta t}$. Because under these conditions there is no spatial positional

information to guide the cell, the chemotactic signaling system must be unresponsive

to the fluctuations in the number of binding events expected from the Poisson noise.

The chemotactic response should only be triggered when the cell moves into a

concentration gradient (**Figures 1b** and **1c**) sufficiently large to drive binding event

fluctuations over the interval Δt with an amplitude that supersedes that of the

background noise. As derived in the **supplementary material**, section *1.2. A condition*

for detecting a change in the chemoattractant concentration, the reliable detection of

a chemoattractant gradient requires the following condition dependent on the maximal

concentration difference experienced during half a revolution and on the mean

chemoattractant concentration $\bar{c}$:

167
$$\left(4\pi Da\bar{c}\,\frac{N}{N+\pi a/s}\Delta t \right) v\Delta t\frac{\partial c}{\partial r}\bar{c}^{-1} \;>\; \sqrt{4\pi Da\bar{c}\,\frac{N}{N+\pi a/s}\Delta t} \tag{2}$$

168   Noting that the left-hand side of the condition represents the chemotactic signal and

169   the right-hand side is a measurement of the background noise, equation (**2**) can be

170   rewritten in terms of signal-to-noise ratio:

171
$$SNR = v\Delta t^{3/2}\left(4\pi Da\bar{c}\,\frac{N}{N+\pi a/s}\right)^{1/2}\xi > 1 \tag{3}$$

Where v is the mean linear velocity, and $\xi = \bar{c}^{-1}\frac{\partial c}{\partial r}$ is the relative slope of the

chemoattractant concentration gradient (see section *1.2. A condition for detecting a*

change in the chemoattractant **concentration** in **supplementary material** for the

derivation). This quantity ξ measures the strength of the stimulus received when

sampling a particular position *r* relative to the mean concentration $\bar{c}$ (**Figure 1c**). As ξ

increases, the strength of the chemotactic signal increases.

Equation (**3**) means that the ability to reliably determine the source of the attractant

depends critically on the relative slope of the chemoattractant concentration gradient ξ,

which must be steep enough to be distinguishable from noise (**Figures 1b** and **1c**, and

Table SI; for further explanation see *1.2. A condition for detecting a change in the*

chemoattractant concentration in **supplementary material**).

We modeled the *SNR* corresponding to different gradients, and within a range of mean

concentrations of chemoattractant between 10^{-11} to 10^{-6} M for three sea urchin species:

S. purpuratus, *L. pictus* and *A. punctulata* (**Figures 1d-f**). For all species studied, at

high mean concentrations of chemoattractant (10^{-8} to 10^{-6} M), the change in

chemoattractant receptor occupancy experienced at two given distinct positions allows

reliable assessment of relatively shallow chemical gradients ($\xi \sim [10^{-3}, 10^{-4}]$ µm^{-1}), with

SNR > 1 for a wide range of ξ (**Figures 1d-f**). However, at low concentrations of

chemoattractant (below 10^{-8} M), keeping all other parameters equal, stochastic

fluctuations begin to mask the signal. In this low-concentration regime, the steepness

of the chemoattractant concentration gradient is determinant for chemoattractant

detection. Shallow gradients result in insufficient *SNR*, while steeper chemoattractant

gradients ($\xi > 10^{-3}$ µm^{-1}) are dependably detected by spermatozoa, i.e. *SNR* > 1 (**Figures**

1d-f).

Previous reports show that *A. punctulata* spermatozoa are very sensitive to resact

(presumably reacting to single molecules) due the high density of resact receptors

($\sim 3 \times 10^{5}$ per cell) which allows them to sense this chemoattractant at low picomolar

concentrations (Kashikar et al., 2012). In contrast, *L. pictus* and *S. purpuratus*

spermatozoa bear lower densities of chemoattractants receptors, approximately 6.3×10^{4}

and 2×10^4 receptors/cell, respectively (Nishigaki et al., 2001; Nishigaki and Darszon,

2000). According to these species-specific differences in chemoattractant receptor

densities, **figures 1d-f** suggest that the spermatozoa of *A. punctulata* are likely more

sensitive to resact, than those of either *L. pictus* or *S. purpuratus* species to the same

mean concentration gradients of speract; with the spermatozoa of *S. purpuratus* being

less sensitive than those of *L. pictus* species to equivalent speract gradients and mean

concentrations. Moreover, the constraints on *SNR* imply that *S. purpuratus* spermatozoa

should only respond to the chemoattractants at higher mean speract concentrations and

at steeper gradients than those that elicit chemotaxis in *L. pictus* spermatozoa (compare

Figures 1d and **1e**).

To understand the differential sensitivity between the spermatozoa of *S. purpuratus*

and *L. pictus* we analyzed the scenario in which the capacity to detect the gradient for

both spermatozoa species were equal, i.e. they would have the same signal-to-noise

ratios, $SNR_{purpuratus} = SNR_{pictus}$. We compute the ratio of the slopes of the speract

concentration gradient experienced by either *S. purpuratus* or *L. pictus* spermatozoa,

which represents a scaling factor (*SF*) in the gradient slope, expressed as:

$$SF = \frac{\xi_{purpuratus}}{\xi_{pictus}} = \left(\frac{v_{pictus}}{v_{purpuratus}} \right) \left(\frac{\Delta t_{pictus}}{\Delta t_{purpuratus}} \right)^{3/2} \left(\frac{Z_{pictus}}{Z_{purpuratus}} \right)^{1/2} \sim 3 \qquad \textbf{(4)}$$

where $Z = \left(\frac{Na}{N + \pi a/s} \right)$ is the probability that a speract molecule that has collided with

the cell will bind to a receptor (Berg and Purcell, 1977), multiplied by the radius a of

the cell.

The estimation of the scaling factor *SF* predicts that *S. purpuratus* spermatozoa

should undergo chemotaxis in a speract gradient three times steeper than the gradient

that elicits chemotaxis in *L. pictus* spermatozoa, with $\xi_{purpuratus} \sim 3\xi_{pictus}$.

In summary, the chemoreception model suggests that *S. purpuratus* spermatozoa

detect chemoattractant gradients with lower sensitivity than those of *L. pictus*. It also

predicts that *S. purpuratus* spermatozoa may detect chemoattractant gradients in the 10^-

^{9} M regime with sufficient certainty only if the slope of the chemoattractant

concentration gradient is greater than 3×10^{-3} μm^{-1} (i.e. steep concentration gradients)

(**Figure 1d**).

If the latter holds true, then *S. purpuratus* spermatozoa should be able to experience

chemotaxis when exposed to steeper speract gradients than those tested experimentally

so far. Given this prediction, we designed and implemented an experimental condition

for which we expect *S. purpuratus* spermatozoa to experience chemotaxis. In general

terms, this scaling rule for sensing chemoattractant gradients might also apply for other

species of marine invertebrates.

S. purpuratus spermatozoa accumulate at steep speract concentration gradients

Our experimental setup is designed to generate determined speract concentration

gradients by focusing a brief (200 ms) flash of UV light along an optical fiber, through

the objective, and into a field of swimming *S. purpuratus* spermatozoa containing

caged-speract (CS) at 10 nM in artificial sea water (Guerrero et al., 2010; Tatsu et al.,

2002). To test experimentally whether *S. purpuratus* spermatozoa undergo chemotaxis,

as predicted from the chemoreception model, we varied the slope of the speract

concentration gradient by separately employing four optical fibers of distinct diameters,

arranged into five different configurations (*f1, f2, f3, f4, f5*) (**Figure 2c**).

Each configuration produces a characteristic pattern of UV illumination within the

imaging field (**Figure 2c**). The UV intensity, measured at the back focal plane of the

objective for each fiber configuration, is shown in **Table SII**. The spatial derivative of

the imaged UV light profile was computed as a proxy for the slope of the speract

concentration gradient (**Figure 2b**). By examining these UV irradiation patterns, the

highest concentration of speract released through photo-liberation from CS is generated

by the *f5* fiber, followed by *f4 > f3 > f2 > f1* (**Figure 2a**). The steepest UV irradiation

gradients are those generated by the *f2, f3* and *f5* fibers (**Figure 2b**).

Irrespective of the optical fiber used, the photo-activation of caged speract triggers

the stereotypical Ca^{2+}-dependent motility responses of *S. purpuratus* spermatozoa

(**Figures 2d, Movies 1, S1, S4-S6**). To determine whether these changes lead to sperm

accumulation, we developed an algorithm, which automatically scores the number of

spermatozoa at any of the four defined concentric regions (R1, R2, R3, and R4) relative

to the center of the speract concentration gradient (**Figures 3a** and **S2**).

As you can see in **Table I**, the photo-liberation of speract through the different fibers

used here triggered various response types (**Figures 3b**, **3c** and **S3**). Negative controls

(Low $[Ca^{2+}]_i$ or High extracellular K^+ ($[K^+]_e$) for *f2* gradient) did not show increased

sperm numbers in any region (**Figures 3b** and **S3**; **Movies S2** and **S3**, respectively).

In summary, *S. purpuratus* spermatozoa accumulate significantly towards the center

of the speract gradients generated by the *f2*- and *f3*-fibers, which generate UV light

profiles with steeper slopes compared to the *f1* and *f4* fibers (**Figure 2b**). These

observations agree with the chemoreception model, in that spermatozoa exposed to

steeper gradients experience lower uncertainty (i.e. higher *SNR*) to determine the

direction of the source of the chemoattractant.

Notably, the use of fibers *f4* and *f5* uncages higher concentrations of speract (by

providing higher UV energies than other fibers) (**Figure 2a** and **Table SII**), yet they do

not trigger the maximum accumulation of *S. purpuratus* spermatozoa at the center of

the chemoattractant field.

S. purpuratus spermatozoa undergo chemotaxis upon exposure to steep speract gradients

The sperm accumulation responses observed in any of *f2* and *f3* conditions suggest that the slope of the chemoattractant concentration gradient might indeed function as a driving force for sperm chemotaxis. However, the accumulation of spermatozoa at the center of the field might also imply other factors, such as cell trapping, or cell death (Yoshida and Yoshida, 2011).

To more reliably scrutinize the trajectories described by *S. purpuratus* spermatozoa in response to speract gradients, chemotactic behavior was quantified using a chemotactic index (CI) that considers the sperm speed and direction both before and after the chemotactic stimulus (see **Figures 4a** and **4b)**. This CI takes values from -1 (negative chemotaxis) to 1 (positive chemotaxis), with 0 being no chemotaxis at all (**Movie 3**). The temporal evolution of the CI, for each of *f1, f2, f3, f4, f5* speract concentration fields, was computed (**Figure 4c)**, and their distributions across time were analyzed by a binomial test (**Figure 4d,** and **Movie S7)** (for further explanation, see *Chemotactic index* section in **Materials and methods**).

The speract fields created by fibers *f2, f3* and *f5* produce significantly positive CI values compared to other conditions (*f1*, *f4* and negative controls), confirming that steeper speract concentration gradients trigger chemotactic responses in *S. purpuratus* spermatozoa. Again, the lack of chemotactic responses in *S. purpuratus* spermatozoa observed by Guerrero et al., 2010, was reproduced through stimulation with *f4*, zero Ca^{2+}, or High K^+ experimental regimes (a scrutiny of non-chemotactic cells is presented in **Figure S8** and section *2.7. Sperm swimming behavior in different chemoattractant gradients* in **supplementary material**).

302  Chemotactic efficiency, which in our work is reported by CI, contains information

303 regarding the capability of single cells to detect and undergo a direct response towards

304 a chemotactic stimulus. It also provides information about the percent of responsive

305 cells that, after detecting a stimulus, can experience chemotaxis. As sperm chemotaxis,

306 and chemotaxis in general, has evolved to operate optimally in the presence of noise

307 (Amselem et al., 2012; Kromer et al., 2018; Lazova et al., 2011), we examined the

308 boundary of *SNR* where sperm chemotaxis operates efficiently for *S. purpuratus* sper-

309 matozoa (**Figure 4e**). Note that, in the regime of *SNR* < 0.7, chemotactic efficiency

310 scales monotonically; for *SNR* > 1, saturation or adaptation mechanisms might impinge

311 on the chemotactic efficiency, as reported in other chemotactic signaling systems

312 (Amselem et al., 2012; Kromer et al., 2018; Lazova et al., 2011). In agreement with

313 these results, we found that the percentage of *S. purpuratus* spermatozoa experiencing

314 relocation increases monotonically with the *SNR* (**Figure 4f**), within the noise limits of

315 0.3 < *SNR* < 1.4, which is also in agreement with the findings of sperm chemotaxis

316 operating optimally in the presence of noise (Amselem et al., 2012; Kromer et al., 2018;

317 Lazova et al., 2011).

319 ***The magnitude of slope of the gradient is a major determinant of sperm chemotaxis***

320  The spatial derivative of the UV profiles shown in **figure 2b** indicates that the

321 steeper light gradients generated from UV irradiation are those of *f2, f3* and *f5*, which

322 are assumed to generate the steepest speract gradients of similar form. This assumption

323 is strictly valid at the instant of UV exposure; subsequently the speract gradient

324 dissipates over time with a diffusion coefficient of $D \approx 240 \ \mu m^2 s^{-1}$. However, the

325 gradient steepness that each spermatozoon experiences during swimming is determined

by the combination of UV flash duration, the speract diffusion time, and the sperm

motility response by itself.

We sought to understand how the stimulus function, which *S. purpuratus*

spermatozoa experience during the accumulation of bound speract throughout their

trajectory, influences their motility response. For this purpose, we computed the spatio-

temporal dynamics of the speract gradient for *f1, f2, f3, f4* and *f5* fibers (**Figure 5a, 5b**

and **S4**) and analyzed the trajectories of spermatozoa swimming in these five distinct

speract gradient configurations (**Figure 5c**, **S5a** and **S5c**). Moreover, we examined the

stimulus function of individual spermatozoa in response to each of the five speract

gradient forms (**Figure 5e**, **S5b, S5d** and **Movie 2**).

The model of chemoreception presented in the previous sections (see equations (**2**)

and (**3**)) predicts a scaling rule for chemotactic responses between *S. purpuratus* and *L.*

pictus spermatozoa of *SF ~ 3* (equation (**4**)). Moreover, the derivatives of the UV-

irradiation profiles shown in **figure 2b** indicate that the *f2*, *f3*, and *f5* fibers generate

steeper speract gradients than the f1 and *f4* fibers.

To determine the direction of the chemoattractant concentration gradient, the signal

difference ∂c between two sampled positions ∂r must be greater than the noise (**Figure**

1c). To test the prediction of the chemoreception model, we computed the local relative

slope of the chemoattractant concentration gradient ξ detected by single spermatozoa

exposed to a given speract concentration gradient, with $\xi = \bar{c}^{-1}\frac{\partial c}{\partial r}$ (**Figure 5e**).

We found that, in agreement with the chemoreception model, the maximum relative

slope of the chemoattractant concentration gradient $\xi_{max} = max(\xi_1, \xi_2, \xi_3, ..., \xi_n)$

required by *S. purpuratus* spermatozoa to undergo chemotaxis is created when the *f2*

and *f3* fibers are employed to generate speract gradients. This relative slope of the

chemoattractant concentration gradients is at least 3 times greater than that experienced

when exposed to the *f4*-generated speract gradient (**Figure 6b**). In addition, *L. pictus*

spermatozoa undergo chemotaxis when exposed to the *f4* speract gradient, which is 2-

3 times smaller than that required by *S. purpuratus* (**Figure 6b**). These findings support

the predicted scaling rule for the detection of the speract concentration gradient between

L. pictus and *S. purpuratus* spermatozoa (**Figures 6b** and **6c**).

The slope of the speract concentration gradient is the critical determinant for the

strength of coupling between the stimulus function and the internal Ca^{2+} oscillator

Friedrich and Jülicher proposed a general theory that captures the essence of sperm

navigation traversing periodic paths in a non-homogeneous chemoattractant field, in

which the sampling of a stimulus function *S(t)* is translated by intracellular signaling

into the periodic modulation of the swimming path curvature *k(t)* (Friedrich and

Jülicher, 2008, 2007). As a result, the periodic swimming path drifts in a direction that

depends on the internal dynamics of the signaling system. In this theory, the latency of

the intracellular signaling (i.e. the $[Ca^{2+}]_i$ signal), expressed as the phase shift between

S(t) and *k(t)*, is a crucial determinant of the directed looping of the swimming trajectory

up the chemical concentration field (Friedrich and Jülicher, 2009, 2008).

Even though this conceptual framework provides insights into the mechanism gov-

erning sperm chemotaxis, it does not explore the scenario where chemoattractants trig-

ger an autonomous $[Ca^{2+}]_i$ oscillator (Aguilera et al., 2012; Espinal et al., 2011; Wood

et al., 2003), which suggests that sperm chemotaxis might operate in a dynamical space

where two autonomous oscillators, namely the stimulus function and the internal Ca^{2+}

oscillator, reach frequency entrainment (**Figure 6a**).

To test the hypothesis that the slope of the speract concentration gradient regulates

the coupling between the stimulus function and the internal Ca^{2+} oscillator triggered by

speract, we made use of a generic model for coupled phase oscillators (Pikovsky et al.,

2003). In its simplest form, the model describes two phase oscillators of intrinsic fre-

quencies ω_1 and ω_2 coupled with a strength γ through the antisymmetric function of

their phase difference $\phi = \varphi_1 - \varphi_2$. The time evolution of ϕ then follows an Adler equa-

tion $d\phi/dt = \Delta\omega - 2\gamma \sin(\phi)$, which is the leading order description for weakly-coupled

non-linear oscillators. In the present case, the two coupled oscillators are the internal

Ca^{2+} oscillator and the oscillations in the stimulus function induced in spermatozoa

swimming across a speract gradient (**Figure 6a**). The former occurs even for immotile

cells, for which there are no stimulus oscillations under a spatially uniform speract field

(**Figure S6,** and **Movie S8**); while the latter exists under two tested negative controls:

cells swimming in Low Ca^{2+} and in High K^+ artificial sea water, both of which inhibit

Ca^{2+} oscillations (see **Figure 3c, S3** and **Movies S2** and **S3**, respectively)**.**

Wood et al., showed that immobilized *S. purpuratus* spermatozoa might experience

spontaneous Ca^{2+} transients (Wood et al., 2003). To provide insight into the mechanism

of sperm chemotaxis we characterized and compared the spontaneous vs the speract-

induced $[Ca^{2+}]_i$ oscillations, and conclude that they are of different oscillatory nature,

hence the spontaneous oscillations do not have a role in sperm chemotaxis (see **Figure**

S9 and section *2.8. Spontaneous vs speract-induced $[Ca^{2+}]_i$ oscillations* in **supplemen-**

tary material).

There are two immediate predictions from the Adler model: first, there is a minimum

coupling strength necessary for the two oscillators to synchronize ($\gamma_{min} = \Delta\omega/2$). For

weaker coupling (i.e. $\gamma < \gamma_{min}$), the two oscillators run with independent frequencies

and, hence, the phase difference increases monotonically with time; second, and within

the synchronous region (i.e. $\gamma > \gamma_{min}$), the phase difference between the oscillators is

constant and does not take any arbitrary value, but rather follows a simple relation to

the coupling strength ($\phi_{sync} = arcsin(\Delta\omega/2\gamma)$). **Figure 6d** shows the two regions in the

parameter space given by $\Delta\omega$ and γ. The boundary between these two regions

corresponds to the condition $\gamma = \gamma_{min}$ and it delimits what is known as an Arnold's

tongue.

We measured the difference in intrinsic frequency by looking at the instantaneous

frequency of the internal Ca^{2+} oscillator just before and after the speract gradient is

established. The range of measured $\Delta\omega$ is shown in **Figure 6d** as a band of accessible

conditions in our experiments (mean of $\Delta\omega$, black line; mean $\pm$ standard deviation,

green dashed lines). If the driving coupling force between the oscillators is the

maximum slope of the speract concentration gradient, i.e. $\gamma = \xi_{max}$, we would expect to

find a minimum slope ($\overline{\xi^*_{max}}$) below which no synchrony is observed. This is indeed

the case as clearly shown in **Figure 6b**, **6e** and **6f** (magenta line). Moreover, and for

cells for which synchronization occurs, the measured phase difference is constrained

by the predicted functional form of $\phi_{sync} = \phi_{sync}(\Delta\omega, \gamma)$ as can be verified from the

collated data shown in **Figure 6e**, and **6f** within the theoretical estimates (see also

Figure S7). Altogether, the excellent agreement of this simple model of coupled phase

oscillators with our data, points to the slope of the speract concentration gradient as the

driving force behind the observed synchronous oscillations and, as a result, for the

chemotactic ability of sea urchin spermatozoa.

Discussion

What are the boundary conditions that limit a sperm's capacity to determine the source of guiding molecules?

During their journey, spermatozoa must measure both the concentration and change on concentration of chemoattractants. Diffusing molecules bind to receptors as discrete packets arriving randomly over time with statistical fluctuations, imposing a limit on detection. By following the differences in the mean concentration of chemoattractants, sampled at a particular time, spermatozoa gather sufficient information to assess the source of the gradient. However, there is a lower detection limit to determine the direction of the chemical gradient, which depends on the swimming speed of the sperm, the sampling time, and as shown in this work, on the steepness of the slope of the chemoattractant concentration gradient.

For almost three decades, chemotaxis had not been observed for the widely-studied *S. purpuratus* species under diverse experimental conditions, raising doubts about their chemotactic capabilities in response to the speract concentration gradients. To examine whether *S. purpuratus* spermatozoa are able to detect spatial information from specific chemoattractant concentration gradient, we use a model of chemoreception developed by Berg and Purcell, 1977, which considers the minimal requirements needed for a single searcher (i.e. a sperm cell) to gather sufficient information to determine the orientation of a non-uniform concentration field. By considering the difference between *L. pictus* and *S. purpuratus* spermatozoa in terms of the number of chemoattractant receptors, receptor pocket effective size, cell size, sampling time, mean linear velocity, sampling distance, and the local mean and slope of the chemoattractant concentration gradient, our model predicts that *S. purpuratus* spermatozoa would need a speract gradient three times steeper than the gradient that drives chemotactic responses for *L.*

pictus spermatozoa. We tested this experimentally by exposing *S. purpuratus* sperma-

tozoa to various defined speract concentration gradients.

We showed that *S. purpuratus* spermatozoa can undergo chemotaxis, but only if the

speract concentration gradients are sufficiently steep, as predicted by the chemorecep-

tion model (i.e. speract gradients that are at in the region of three times steeper than the

speract concentration gradient that drives chemotaxis in *L. pictus* spermatozoa). This

confirms and explains why the shallower speract gradients previously tested are unable

to generate any chemotactic response in *S. purpuratus* spermatozoa (Guerrero et al.,

2010a), despite inducing characteristic "turn and run" motility responses.

These findings indicate that the guiding chemical gradient must have a minimum

steepness to elicit sperm chemotaxis, where the signal-to-noise relationship (*SNR*) of

stimulus to the gradient detection mechanism imposes a limit for the chemotactic effi-

ciency. Our results are in agreement with recent theoretical studies by Kromer and col-

leagues, indicating that sperm chemotaxis of marine invertebrates operates optimally

within a boundary defined by the *SNR* of collecting ligands within a chemoattractant

concentration gradient (Kromer et al., 2018). We showed that *SNR* can be tuned by the

steepness of the chemical gradient, where higher *SNR*'s are reached at steeper gradients,

hence increasing the probabilities of locating the source of the gradient.

The large majority of marine spermatozoa characterized to date, together with many

motile microorganisms, explore their environment via helical swimming paths, where-

upon encountering a surface these helices collapse to circular trajectories. The intrinsic

periodicity of either swimming behavior commonly results in the periodic sampling of

the cell chemical environment with direct implications for their ability to accurately

perform chemotaxis.

The periodic sampling of chemoattractants by the sperm flagellum continuously

feeds back to the signaling pathway governing the intracellular Ca^{2+} oscillator, hence

providing a potential coupling mechanism for sperm chemotaxis. Indirect evidence for

the existence of a feedback loop operating between the stimulus function and the Ca^{2+}

oscillator triggered by chemoattractants has been found in *L. pictus*, *A. punctulata* and

Ciona intestinalis (ascidian) species, whose spermatozoa show robust chemotactic re-

sponses towards their conspecific chemoattractants (Böhmer et al., 2005; Guerrero et

al., 2010a; Jikeli et al., 2015; Shiba et al., 2008).

To investigate further the molecular mechanism involved in sperm chemotaxis, we

measured both the stimulus function and the triggered $[Ca^{2+}]_i$ oscillations for up to one

thousand *S. purpuratus* spermatozoa exposed to five distinctly-shaped speract concen-

tration gradients. We demonstrate that the steepness of the slope of the chemoattractant

concentration gradient is a major determinant for sperm chemotaxis in *S. purpuratus*

and might be an uncovered feature of sperm chemotaxis in general. A steep slope of

the speract concentration gradient entrains the frequencies of the stimulus function and

the internal Ca^{2+} oscillator triggered by the periodic sampling of a non-uniform speract

concentration field. We assessed the transition boundary of the coupling term (the slope

of the speract concentration gradient) for the two oscillators to synchronize and found

it to be very close to the boundary where *S. purpuratus* starts to experience chemotaxis.

The agreement of our data with a model of weakly-coupled phase oscillators suggests

that the slope of the speract concentration gradient is the driving force behind the ob-

served synchronous oscillations and, as a result, for the chemotactic ability of sea urchin

spermatozoa.

Caution needs to be taken with the interpretations of the agreement of our data with

such a generic model for coupled phase oscillators, particularly when considering only

a few steps of the oscillatory cycles. The latter is relevant for assessing frequency

entrainment, which in some cases demands a certain delay before reaching the synchro-

nized state, i.e. when the natural frequencies of the connected oscillators are very dis-

tinct. The chemotactic responses scored in the present study encompass a few steps

(<10) of both the stimulus function and the internal Ca^{2+} oscillator triggered by speract

(**Figures S5-S7**). Our data indicate that within the chemotactic regime, frequency en-

trainment of the stimulus function and the internal Ca^{2+} oscillator of *S. purpuratus* sper-

matozoa seems to occur almost instantaneously, within the first three oscillatory steps

(**Figure S7**). Such interesting findings can be explained by the proximity of the natural

frequencies of both oscillators (**Figure 6d**), which may relieve the need of longer delay

for reaching frequency entrainment. Whether the proximity of the frequencies of both

oscillators is sculped by the ecological niche where sperm chemotaxis occurs is an open

question, however, a near-instantaneous entrainment would confer obvious evolution-

ary advantage under the reproductively competitive conditions of synchronized spawn-

ing as undertaken by sea urchins.

One can further hypothesize about the evolutionary origin of the described differ-

ences in sensitivity to chemoattractant concentration gradients between *S. purpuratus*

and *L. pictus* spermatozoa if we consider their respective ecological reproductive

niches. The turbulent environment where sea urchins reproduce directly impinges on

the dispersion rates of small molecules such as speract, hence, imposing ecological lim-

its that constrain permissive chemoattractant gradient topologies within different hy-

drodynamic regimes. For instance, the reproductive success of *L. pictus*, *S. purpuratus*

and abalone species has been shown to peak at defined hydrodynamic shearing values

(Hussain et al., 2017; Mead and Denny, 1995; Riffell and Zimmer, 2007; Zimmer and

Riffell, 2011). What are the typical values of the chemoattractant gradients encountered

by the different species in their natural habitat? The correct scale to consider when

discussing the small-scale distribution of chemicals in the ocean is the Batchelor scale,

$l_B = (\eta D^2/\zeta)^{1/4}$, where η is kinematic viscosity, D the diffusion coefficient and ζ is the

turbulent dissipation rate (Aref et al., 2017; Batchelor et al., 1959). Turbulence stirs

dissolved chemicals in the ocean, stretching and folding them into sheets and filaments

at spatial dimensions down to the Batchelor scale: below l_B molecular diffusion domi-

nates and chemical gradients are smoothened out.

S. purpuratus is primarily found in the low intertidal zone. The purple sea urchin

lives in a habitat with strong wave action and areas with shaking aerated water. These

more energetic zones, including tidal channels and breaking waves, generate relatively

high levels of turbulence ($\zeta \sim 10^{-4}$ m^{2}s^{-3}) which lead to somewhat small values of l_B

and, hence, to steep gradients (i.e. $1/l_B$). *L. pictus*, on the contrary, is mostly found at

the edge of or inside kelp beds, well below the low tide mark where the levels of tur-

bulence are much more moderate ($\zeta \sim 10^{-6}$ m^{2}s^{-3}) (Jimenez, 1997; Thorpe, 2007). This

difference in the turbulent kinetic energy dissipation rate has a significant effect on the

spatial dimensions of chemical gradients for sperm chemotaxis present in a particular

habitat. The ratio of l_B for the different habitats scales as $l_{Bpurpuratus}/l_{Bpictus} \sim (\zeta_{pictus}/\zeta_{pur-}$

$_{puratus})^{1/4} \sim 3$, which fits considerably well with the relative sensitivity to speract of the

two species. Furthermore, we have shown that *S. purpuratus* spermatozoa experience

chemotaxis toward steeper speract gradients than those that guide *L. pictus* spermato-

zoa, which is also compatible with the distinct chemoattractant gradients they might

naturally encounter during their journey in search of an egg.

Materials and Methods

Materials

Artificial seawater (ASW), and Low Ca^{2+} ASW were prepared as in Guerrero et al., 2010, their detailed composition, together with an extended list of other materials is presented in the **supplementary material**. Caged speract (CS), was prepared as described previously (Tatsu et al., 2002).

Loading of Ca^{2+}-fluorescent indicator into spermatozoa and microscopy imaging

S. purpuratus or *L. pictus* spermatozoa were labeled with fluo-4-AM (as described in section **2.2. Loading of Ca^{2+}-fluorescent indicator into spermatozoa** in **supplementary material**), and their swimming behavior was studied at the water-glass interface on an epifluorescence microscope stage (Eclipse TE-300; Nikon). The cover slips were covered with poly-HEME to prevent the attachment of the cells to the glass. Images were collected with a Nikon Plan Fluor 40x 1.3 NA oil-immersion objective. Temperature was controlled directly on the imaging chamber at a constant 15 ℃. Stroboscopic fluorescence excitation was provided by a Cyan LED synchronized to the exposure output signal of the iXon camera (2 ms illumination per individual exposure, observation field of 200 x 200 µm), the fluorescence cube was set up accordingly (see **supplementary material**).

Image processing and quantification of global changes of spermatozoa number and $[Ca^{2+}]_i$

To study the dynamics of overall sperm motility and $[Ca^{2+}]_i$ signals triggered by the distinct speract gradients, we developed an algorithm that provides an efficient approach to automatically detect the head of every spermatozoa in every frame of a given video-microscopy file. A detailed description of the algorithm is provided in the **supplementary material**.

Computing the dynamics of speract concentration gradients

The dynamics of the chemoattractant gradient was computed using Green's function of the diffusion equation:

$$c = f(r, t) = \frac{c_0}{\sqrt{4\pi D t}} e^{-r^2/\sigma^2} + c_b \tag{5}$$

Equation **(5)** for the concentration tells us that the profile has a Gaussian form, where D is the diffusion coefficient of the chemoattractant, c_b is the basal concentration of the chemoattractant, t is the time interval, r is the distance to the center of the gradient and c_0 is the initial concentration. The width of the Gaussian is $\sigma = \sqrt{4D(t + t_0)}$, and hence it increases as the square root of time.

The speract concentration gradients were generated via the photolysis of 10 nM caged speract (CS) with a 200 ms UV pulse delivered through each of four different optical fibers with internal diameters of 0.2, 0.6, 2, and 4 mm (at two different positions). Light intensity was normalized dividing each point by the sum of all points of light intensity for each fiber and multiplying it by the fiber potency (measured at the back focal plane of the objective) in milliwatts (mW) (**Table SII**). Each spatial distribution of instantaneously-generated speract concentration gradient was computed by fitting their corresponding normalized spatial distribution of UV light (Residual standard error: 2.7×10^{-5} on 97 degrees of freedom), considering an uncaging efficiency of 5-10%, as reported (Tatsu et al., 2002).

The diffusion coefficient of speract has not been measured experimentally. However, the diffusion coefficient of a similar chemoattractant molecule, resact (with fourteen amino acids), has been reported, $D_{resact} = 239 \pm 7$ μm^2 s^{-1} (Kashikar et al., 2012). If we consider that speract is a decapeptide, the 1.4 fold difference in molecular weight between speract and resact would imply a $(1.4)^{1/3}$ fold difference in their diffusion coefficients, which is close to the experimental error reported (Kashikar et al.,

2012). For the sake of simplicity, the spatio-temporal dynamics of the distinct

instantaneously generated speract gradients was modeled considering a speract

diffusion coefficient of $D_{speract} = 240\ \mu m^2\ s^{-1}$.

Computing $[Ca^{2+}]_i$ dynamics and the stimulus function of single spermatozoa

Spermatozoa were tracked semi-automatically by following the head centroid with

the MtrackJ plugin (Meijering et al., 2012) of ImageJ 1.49u. Single cell $[Ca^{2+}]_i$ signals

were computed from the mean value of a 5 x 5 pixel region, centered at each sperm

head along the time. The head position of each spermatozoa x was used to compute the

mean concentration of speract at r over each frame. The stimulus function of single

spermatozoa $S = f(c)$ was computed by solving equation (**5**), considering both their

swimming trajectories, and the spatio-temporal evolution of a given speract

concentration gradient. The profiles of UV light were used to compute the initial

conditions at $c(r, t_o)$.

The phase- and temporal-shifts between the time derivative of the stimulus function

dS/dt and the internal Ca^{2+} oscillator triggered by speract, were computed from their

normalized cross-correlation function.

Programs were written in R statistical software (R Development Core Team, 2016).

Chemotactic index (CI)

Each sperm trajectory was smoothened using a moving average filter, with a window

of 60 frames (two seconds approximately) (**Figure 4b and Movie 3**). A linear model

was then fitted to the smoothed trajectory; the corresponding line was forced to go

through the mean point of the smoothed trajectory (orange point in **Figure 4b and**

Movie 3). The θ angle between red and black vectors was calculated in each frame from

the second 4.5 to 10.

The chemotactic index was defined based on the progressive displacement of the

sperm trajectory as $CI = \frac{|u|cos\theta - |v|cos\varphi}{|u|+|v|}$, being ϕ and θ the angles between gray and ma-

genta, and red and black vectors, respectively; and $|v|$ and $|u|$ the magnitude of the sperm

progressive speed before and after speract uncaging, respectively (**Figure 4b and**

Movie 3). The CI considers the sperm displacement before speract uncaging (i.e. un-

stimulated drift movement at 0-3 seconds), and then subtracts it from the speract in-

duced effect (at 3-10 seconds). The CI takes values from -1 (negative chemotaxis) to 1

(positive chemotaxis), being 0 no chemotaxis at all.

Statistical analyses

The normality of the CI distributions, each obtained from *f1* to *f5* speract gradient stim-

uli, was first assessed using the Shapiro-Wilk test; none of them were normal (Gauss-

ian), so each CI distribution was analyzed using non-parametric statistics (**Figure 4d**

and **Movie S7**). The curves obtained from medians of each CI distribution were

smoothed using a moving average filter, with a window of 20 frames (0.6 seconds)

(**Figure 4c**).

Data are presented for individual spermatozoa (n) collected from up to three sea

urchins. All statistical tests were performed using R software (R Development Core

Team, 2016). The significance level was set at 95% or 99%.

Acknowledgements

The authors thank Dr. Tatsu Yoshiro for providing the caged speract, and Drs. Hermes

Gadêlha, David Smith and Nina Pastor for stimulating discussions and a critical reading

of the manuscript. AG thanks Dr. Manabu Yoshida and Dr. Kaoru Yoshida for feedback

regarding sperm chemotaxis in marine invertebrates, and to the Japan Society for the

Promotion of Science (JSPS invitation fellowship for research in Japan to A.G., short

term JSPS/236, ID no. S16172). AD performed part of this work while carrying out a

Sabbatical at the Instituto Gulbenkian de Ciencia (IGC) supported by UNAM/DGAPA

and IGC.

Author contributions

A.G., A.D. and I.T. conceived the project; A.G. and V.J.S. performed the

experiments; H.R., A.G., I.T., V.J.S and M.V. analyzed the data; H.R., A.G., I.T. and

J.C. carried out the mathematical model calculations and wrote the corresponding

section; H.R. and I.T. developed the chemotactic index; A.D., A.G., H.R. and I.T.

participated in the design and drafting of the manuscript; J.C., C.D.W. and C.B.

provided feedback for conceptualization and drafting of the manuscript. All authors

approved the final version of the article.

Competing interests

The authors declared that no competing interests exist.

Funding

A.D. and A.G. acknowledge grants from the Consejo Nacional de Ciencia y

Tecnología (CONACyT Fronteras 71, Ciencia Básica 252213 and 255914), A.G., A.D.

and C.B to Programa de Apoyo a Proyectos de Investigación e Innovación Tecnológica

UNAM (PAPIIT/DGAPA) (IA202417, IN205516, IN206016, IN215519 and

IN112514). We acknowledge CONACYT and PAPIIT for fellowships to M.V.P. and

H.R.; the Spanish Ministry of Economy and Competitiveness Grants No. FIS2013-

48444-C2-1-P, FIS2016-77692-C2-1- P, the Subprogram Ramón-y-Cajal and the

Ibero-America Program-Santander Universities 2015 (I.T.). M.V.P. had a scholarship

from PAPIIT/DGAPA IN206016.

Figure 1. Physics of chemoreception. a. A spermatozoon swimming in an isotropic chemoattractant concentration field, where the number of molecules detected (n) is within the noise of detection. **b.** A spermatozoon swimming near to an egg, while chemoattractant molecules are diffusing from its surrounding jelly layer creating a chemoattractant gradient. Note that the signal detected in this case is larger than the detection noise. **c.** The assessment of a chemoattractant concentration gradient requires that the signal difference ∂c between two sampled positions ∂r must be greater than the noise. **d-f.** The signal-to-noise ratio in the determination of the chemoattractant gradient *SNR* plotted against the relative slope of the chemoattractant concentration gradient in log scale, $\xi = \bar{c}^{-1}\frac{\partial c}{\partial r}$, for different chemoattractant concentrations of speract for either *S. purpuratus* (**d**), or *L. pictus* (**e**) spermatozoa, and of resact for *A. punctulata* (**f**) spermatozoa (see **Table SI** for the list of parameter values taken in consideration for panels **d-f**). *S. purpuratus* spermatozoa have lower capacity of

680   detection for the same chemoattractant concentrations at a given ξ than *L. pictus* and *A.*

681   *punctulata*. The tone of the shaded areas indicates shallow or steep gradient conditions.

682   The horizontal dotted line represents *SNR* = 1; the vertical magenta line represents ξ =

683   2.8 x 10^{-3} μm^{-1}.

Figure 2. Screening of speract concentration gradients. a. Radial profile and its

derivative (**b**) of the UV light scattered at the glass-liquid interface for each optical fiber

(*f1-f5*). **c**. Spatial distribution of the UV flash energy for each fiber. **d.** Representative

motility and $[Ca^{2+}]_i$ responses of *S. purpuratus* spermatozoa exposed to different

concentration gradients of speract. $F-F_0$ time projections, showing spermatozoa head

fluorescence at 3 s intervals before and after photoactivation of 10 nM caged speract in

691   artificial seawater with 200 ms UV flash. The pseudo-color scale represents the relative

692   fluorescence of fluo-4, an $[Ca^{2+}]_i$ indicator, showing maximum (red) and minimum

693   (blue) relative $[Ca^{2+}]_i$. Scale bars of 50 µm.

Figure 3. Motility and $[Ca^{2+}]_i$ responses of *S. purpuratus* spermatozoa exposed to specific concentration gradients of speract. a. Automated segmentation algorithm of swimming sperm. The positions of the sperm heads within the imaging field are assigned to either R1, R2, R3 or R4 concentric regions around the centroid of the UV flash intensity distribution. Each ROI was also used to compute the sperm head fluorescence from the raw video microscopy images (as the mean value of the ROI) (see **Figure S2**). Scale bar of 50 µm. **b**. Fold change in sperm number, defined as the number of spermatozoa at the peak of the response (6 s) relative to the mean number before speract stimulation (0-3 s) (see **Figure S3**). **c**. Relative changes in $[Ca^{2+}]_i$ experienced by spermatozoa at the peak response (6 s) after speract stimulation. Negative controls for spermatozoa chemotaxis are artificial seawater with nominal Ca^{2+} (Low Ca^{2+}); and artificial seawater with 40 mM of K^+ (High K^+). Both experimental conditions prevent chemotactic responses by inhibiting the Ca^{2+} membrane permeability alterations triggered by speract; the former disrupts the Ca^{2+} electrochemical gradient, and the later disrupt the K^+ electrochemical gradient required as electromotive force needed to elevate pH_i and open Ca^{2+} channels. The central line in each box plot represents the median value, the box denotes the data spread from 25% to 75%, and the whiskers reflect 10–90%. The number of experiments is indicated on

713   the bottom of each experimental condition. We used the same number of experiments

714   for the relative change in $[Ca^{2+}]_i$ (right panel). *Statistical significance, $p < 0.05$;

715   multiple comparison test after Kruskal-Wallis.

Figure 4. *S. purpuratus* **spermatozoa selectively experience chemotaxis towards specific speract gradients. a.** Sperm trajectory before (gray) and after (black) the UV irradiation (purple). **b.** Definition of a chemotactic index to score chemotactic responses. Dots represent sperm trajectory before (gray) and after (black) UV irradiation. Green and blue empty spirals indicate the smoothed trajectory before and after UV irradiation. Grey and black vectors are the progressive sperm displacement before and after stimulation, respectively; and the **v** and **u** vectors are the linear speed before and after stimulation; and ϕ and θ are the angles to their corresponding reference vectors to the center of the imaging field – the highest UV irradiated area, (magenta and red, respectively). Chemotactic index (CI) is defined as in the inset (see also **Movie 3**). **c.**

Temporal evolution of the chemotactic index. Functions were calculated from the me-

dian obtained from sperm trajectories of each of *f1, f2, f3, f4, f5, f2-ZeroCa^{2+},* and *f2-*

HighK$^+$ experimental conditions (**Movie S7**). **d.** Radial histograms at second 9 (vertical

dotted line at panel **c**). Significant differences (*Binomial test, p-value < 0.05*) were ob-

served only for *f2*, *f3* and *f5* fibers, compared to controls. n represents the number of

individual sperm trajectories analyzed. **e.** Chemotaxis index (CI) as a function of the

signal-to-noise ratio (*SNR*). Each parameter was calculated for single cells. Large filled

points represent the median for each gradient condition distribution **f.** Fraction of re-

sponding cells as a function of the *SNR*. Each parameter was calculated per gradient

condition.

Figure 5. Steep speract gradients provoke chemotaxis in *S. purpuratus* spermato-

zoa. a. Dynamics of the *f2* speract gradient. The blue dashed line ($t_0 = 0$ s) corresponds

to a Gaussian distribution fitted to the UV light profile and illustrates the putative shape

of the instantaneously-generated speract concentration gradient. Solid black lines illus-

trate the temporal evolution of the speract concentration field after t = 1, 2, 3, …, 20

seconds. **b.** Temporal changes in the *f2* speract field computed radially (each 10 µm)

from the center of the gradient. **c.** Characteristic motility changes of a *S. purpuratus*

spermatozoon exposed to the *f2* speract gradient. Solid lines illustrate its swimming

trajectory 3 s before (gray), during UV flash (purple) and 6 s after (black) speract ex-

posure. **d.** Spermatozoa head distance to the source of the speract gradient versus time,

calculated from sperm trajectory in **c**. **e.** Stimulus function computed from the swim-

ming behavior of the spermatozoon in **c**, considering the dynamics of **a** and **b**.

Figure 6. The slope of the speract concentration gradient generates a frequency-locking phenomenon between the stimulus function and the internal Ca²⁺ oscillator triggered by speract. a. Coupled oscillator model. Each sperm has two independent oscillators: i) stimulus function and ii) $[Ca^{2+}]_i$, which can be coupled through a forcing term that connects them, in our case the slope of the chemoattractant concentration gradient (ξ_0). **b.** Maximum relative slopes (ξ_{max}) of the chemoattractant concentration gradient experienced by *S. purpuratus* (*Sp*) spermatozoa when exposed to *f1, f2, f3, f4, f5* speract gradients. The maximum relative slopes of the chemoattractant concentration gradient experienced by *L. pictus* spermatozoa (*Lp*) towards *f4* experimental regime are also shown. Note that ξ_{max} for *f2, f3*, and *f5*, are up to 2-3 times greater than in *f4*, regardless of the species. **c.** Experimental signal-to-noise ratios (*SNR*) regimes experienced by spermatozoa swimming in different gradient conditions. Note that only *f2, f3* and *f5* have higher *SNR*, compared to other gradient conditions, for which stochastic fluctuations mask the signal. This *SNR* calculation assumes a 10% of speract uncaging.

d. Arnold's tongue indicating the difference in intrinsic frequency of the internal Ca^{2+}

oscillator of *S. purpuratus* spermatozoa, just before and after the speract gradient ex-

posure. **e**. Phase difference between the time derivative of the stimulus function and the

internal Ca^{2+} oscillator of *S. purpuratus* spermatozoa, obtained by computing the cross-

correlation function between both time series (**Figure S7**). **f**. Phase difference between

the time derivative of the stimulus function and the internal Ca^{2+} oscillator of *S. purpu-*

ratus spermatozoa expressed in temporal delays. **d-f**. Gray points represent the collated

data of all *f1, f2, f3, f4, f5* experimental regimes. Red, black and blue points indicate

chemotactic spermatozoa (CI > 0 at second 3 after UV flash), located in R3, and R4

regions just before the speract gradient is established under *f2, f3* and *f5* experimental

regimes, respectively. Magenta lines represent the transition boundary ($\gamma_{min} = \overline{\zeta^*_{max}}$ ~

2.8 x 10^{-3} µm^{-1}, see also **Figure 1d-f**) below which no synchrony is observed, obtained

from the theoretical estimates (black curves, mean of $\Delta\omega$) of panels **e** and **f**. Green

dashed lines indicate confidence intervals (mean ± standard deviation).

Table I. Accumulation responses triggered by different speract gradients.

Gradient	Sperm accumulation	Sperm depleted	$[Ca^{2+}]_i$ rise (fold)
f1	No	No	< 2
f2	R1 and R2	R3 and R4	> 2
f3	R1, R2 and R3	R4	> 2
f4	No	No	> 2
f5	No	No	~ 2

Accumulation responses were evaluated at second 6, i.e. 3 seconds after photo-libera-

tion of speract by a 200 ms UV flash.

https://www.dropbox.com/s/oe0mnc8j5r65l8s/Movie%201.avi?dl=0

Movie 1. Typical motility and Ca^{2+} responses of *S. purpuratus* spermatozoa to-

wards an *f2*-generated speract concentration gradient. Spermatozoa swimming in

artificial sea water containing 10 nM caged speract, 3 s before and 5 s after 200 ms UV

irradiation. An optical fiber of 0.6 mm internal diameter (*f2*) was used for the UV light

path to generate the speract concentration gradient. Real time: 30.8 frames s^{-1},

40x/1.3NA oil-immersion objective. Note that spermatozoa located at R2, R3 and R4

regions prior to speract exposure swim up the speract concentration gradient, towards

the center of the imaging field. The pseudo-color scale represents the relative fluores-

cence of fluo-4, a Ca^{2+} indicator, showing maximum (red) and minimum (blue) relative

[Ca^{2+}]$_i$. Six *S. purpuratus* spermatozoa were manually tracked for visualization pur-

poses. Scale bar of 50 µm.

https://www.dropbox.com/s/gtvau67qlaoriwj/Movie%202.1.avi?dl=0

Movie 2. Sperm trajectory analysis and stimulus function. Single-cell analysis was

performed for approximately 1000 sperm trajectories for the different speract gradients

(*f1-f5* and negative controls). The sperm trajectory shown here is representative of a

chemotactic sperm. This analysis was implemented after speract uncaging at 3 seconds

(from 3.2 - 10 seconds). Trajectory before, after and during the 200 ms UV flash is

shown in gray, black and purple, respectively.

Movie 3. Sperm trajectory analysis and chemotactic index (CI). Single-cell analysis

was performed for approximately 1000 sperm trajectories from the different speract

gradients (*f1-f5* and negative controls). Angle ϕ is calculated just once and is always

the same for each sperm trajectory. Angle θ is calculated per frame of the video for

each sperm trajectory, resulting in the chemotactic index kinetics for each sperm tra-

jectory (right panel). The sperm trajectory shown here represents a chemotactic sperm.

This analysis was implemented from 4.5 seconds to 10 seconds. Speract uncaging was

induced at 3 seconds. Trajectory before and after speract release is shown in gray and

black dots, respectively.

References

Aguilera LU, Galindo BE, Sánchez D, Santillán M. 2012. What Is the Core Oscillator
in the Speract-Activated Pathway of the Strongylocentrotus purpuratus Sperm
Flagellum? *Biophys J* **102**:2481–2488. doi:10.1016/j.bpj.2012.03.075
Alvarez L, Dai L, Friedrich BM, Kashikar ND, Gregor I, Pascal R, Kaupp UB. 2012.
The rate of change in Ca 2+ concentration controls sperm chemotaxis. *J Cell
Biol* **196**:653–663. doi:10.1083/jcb.201106096
Amselem G, Theves M, Bae A, Beta C, Bodenschatz E. 2012. Control Parameter
Description of Eukaryotic Chemotaxis. *Phys Rev Lett* **109**:108103.
doi:10.1103/PhysRevLett.109.108103
Aref H, Blake JR, Budišić M, Cardoso SSS, Cartwright JHE, Clercx HJH, El Omari
K, Feudel U, Golestanian R, Gouillart E, van Heijst GF, Krasnopolskaya TS, Le
Guer Y, MacKay RS, Meleshko V V., Metcalfe G, Mezić I, de Moura APS, Piro
O, Speetjens MFM, Sturman R, Thiffeault J-L, Tuval I. 2017. Frontiers of
chaotic advection. *Rev Mod Phys* **89**:025007.
doi:10.1103/RevModPhys.89.025007
Batchelor GK, Howells ID, Townsend AA. 1959. Small-scale variation of convected
quantities like temperature in turbulent fluid Part 2. The case of large
conductivity. *J Fluid Mech* **5**:134. doi:10.1017/S0022112059000106
Berg HC, Purcell EM. 1977. Physics of chemoreception. *Biophys J* **20**:193–219.
doi:10.1016/S0006-3495(77)85544-6
Böhmer M, Van Q, Weyand I, Hagen V, Beyermann M, Matsumoto M, Hoshi M,
Hildebrand E, Kaupp UB, Bohmer M, Van Q, Weyand I, Hagen V, Beyermann
M, Matsumoto M, Hoshi M, Hildebrand E, Kaupp UB. 2005. Ca2+ spikes in the
flagellum control chemotactic behavior of sperm. *EMBO J* **24**:2741–52.
doi:10.1038/sj.emboj.7600744
Brokaw CJ. 1979. Calcium-induced asymmetrical beating of triton-demembranated
sea urchin sperm flagella. *J Cell Biol* **82**:401–411. doi:10.1083/jcb.82.2.401
Cook SP, Brokaw CJ, Muller CH, Babcock DF. 1994. Sperm Chemotaxis: Egg
Peptides Control Cytosolic Calcium to Regulate Flagellar Responses. *Dev Biol*
165:10–19. doi:10.1006/dbio.1994.1229
Darszon A, Guerrero A, Galindo BE, Nishigaki T, Wood CD. 2008. Sperm-activating
peptides in the regulation of ion fluxes, signal transduction and motility. *Int J
Dev Biol* **52**:595–606. doi:10.1387/ijdb.072550ad
Espinal J, Aldana M, Guerrero A, Wood C, Darszon A, Martínez-Mekler G. 2011.
Discrete Dynamics Model for the Speract-Activated Ca2+ Signaling Network
Relevant to Sperm Motility. *PLoS One* **6**:e22619.
doi:10.1371/journal.pone.0022619
Friedrich BM, Jülicher F. 2009. Steering Chiral Swimmers along Noisy Helical Paths.
Phys Rev Lett **103**:068102. doi:10.1103/PhysRevLett.103.068102
Friedrich BM, Jülicher F. 2008. The stochastic dance of circling sperm cells: sperm
chemotaxis in the plane. *New J Phys* **10**:123025. doi:10.1088/1367-
2630/10/12/123025
Friedrich BM, Jülicher F. 2007. Chemotaxis of sperm cells. *Proc Natl Acad Sci*
104:13256–13261. doi:10.1073/pnas.0703530104
Guerrero A, Nishigaki T, Carneiro J, Tatsu Y, Wood CD, Darszon A, Yoshiro T,
Wood CD, Darszon A. 2010a. Tuning sperm chemotaxis by calcium burst
timing. *Dev Biol* **344**:52–65. doi:10.1016/j.ydbio.2010.04.013
Guerrero A, Wood CD, Nishigaki T, Carneiro J, Darszon A. 2010b. Tuning sperm

chemotaxis. *Biochem Soc Trans* **38**:1270–1274. doi:10.1042/BST0381270

Hansbrough JR, Garbers DL, Garbers L. 1981. Speract. Purification and characterization of a peptide associated with eggs that activates spermatozoa. *J Biol Chem* **256**:1447–52.

Hussain YH, Sadilek M, Salad S, Zimmer RK, Riffell JA. 2017. Individual female differences in chemoattractant production change the scale of sea urchin gamete interactions. *Dev Biol* **422**:186–197. doi:10.1016/j.ydbio.2017.01.006

Jikeli JF, Alvarez L, Friedrich BM, Wilson LG, Pascal R, Colin R, Pichlo M, Rennhack A, Brenker C, Kaupp UB. 2015. Sperm navigation along helical paths in 3D chemoattractant landscapes. *Nat Commun* **6**:7985. doi:10.1038/ncomms8985

Jimenez J. 1997. Oceanic turbulence at millimeter scales. *Sci Mar* **61**:47–56.

Kashikar ND, Alvarez L, Seifert R, Gregor I, Jäckle O, Beyermann M, Krause E, Kaupp UB, Benjamin Kaupp U. 2012. Temporal sampling, resetting, and adaptation orchestrate gradient sensing in sperm. *J Cell Biol* **198**:1075–1091. doi:10.1083/jcb.201204024

Kaupp UBB. 2012. 100 years of sperm chemotaxis. *J Gen Physiol* **140**:583–586. doi:10.1085/jgp.201210902

Kromer JA, Märcker S, Lange S, Baier C, Friedrich BM. 2018. Decision making improves sperm chemotaxis in the presence of noise. *PLOS Comput Biol* **14**:e1006109. doi:10.1371/journal.pcbi.1006109

Lazova MD, Ahmed T, Bellomo D, Stocker R, Shimizu TS. 2011. Response rescaling in bacterial chemotaxis. *Proc Natl Acad Sci* **108**:13870–13875. doi:10.1073/pnas.1108608108

Lillie FR. 1913. The mechanism of fertilization. *Science (80-)* **38**:524–528. doi:10.1126/science.38.980.524

Lotterhos K, Levitan D, Traits G. 2010. Gamete release and spawning behavior in broadcast spawning marine invertebrates. *Evol Prim Sex Characters Anim* 99–120.

Mead KS, Denny MW. 1995. The effects of hydrodynamic shear stress on fertilization and early development of the purple sea urchin Strongylocentrotus purpuratus. *Biol Bull* **188**:46–56. doi:10.2307/1542066

Meijering E, Dzyubachyk O, Smal I. 2012. Methods for Cell and Particle Tracking. pp. 183–200. doi:10.1016/B978-0-12-391857-4.00009-4

Miller RL. 1985. Sperm Chemo-Orientation in the Metazoa In: Metz CB, Monroy A, editors. Biology of Fertilization. New York: Academic Press. pp. 275–337.

Mizuno K, Shiba K, Yaguchi J, Shibata D, Yaguchi S, Prulière G, Chenevert J, Inaba K. 2017. Calaxin establishes basal body orientation and coordinates movement of monocilia in sea urchin embryos. *Sci Rep* **7**:1–10. doi:10.1038/s41598-017-10822-z

Nishigaki T, Darszon A. 2000. Real-Time Measurements of the Interactions between Fluorescent Speract and Its Sperm Receptor. *Dev Biol* **223**:17–26. doi:10.1006/dbio.2000.9734

Nishigaki T, Zamudio FZ, Possani LD, Darszon A. 2001. Time-resolved sperm responses to an egg peptide measured by stopped-flow fluorometry. *Biochem Biophys Res Commun* **284**:531–5. doi:10.1006/bbrc.2001.5000

Pfeffer W. 1884. Locomotorische Richtungsbewegungen durch chemische Reize. *Untersuchungen aus dem Bot Inst zu Tübingen.*

Pikovsky A, Rosenblum M, Kurths J. 2003. Synchronization: A Universal Concept in Nonlinear Sciences. *Cambridge Nonlinear Sci Ser 12* 432.

doi:10.1063/1.1554136

R Development Core Team. 2016. R: A language and environment for statistical
computing. Vienna, Austria: R Foundation for Statistical Computing.

Riffell J a, Zimmer RK. 2007. Sex and flow: the consequences of fluid shear for
sperm egg interactions. *J Exp Biol* **210**:3644–3660. doi:10.1242/jeb.008516

Shiba K, Baba SA, Inoue T, Yoshida M. 2008. Ca2+ bursts occur around a local
minimal concentration of attractant and trigger sperm chemotactic response.
Proc Natl Acad Sci U S A **105**:19312–7. doi:10.1073/pnas.0808580105

Strünker T, Alvarez L, Kaupp UB. 2015. At the physical limit - chemosensation in
sperm. *Curr Opin Neurobiol* **34**:110–116. doi:10.1016/j.conb.2015.02.007

Suzuki N. 1995. Structure, function and biosynthesis of sperm-activating peptides and
fucose sulfate glycoconjugate in the extracellular coat of sea urchin eggs.
ZoologSci **12**:13–27.

Tatsu Y, Nishigaki T, Darszon A, Yumoto N. 2002. A caged sperm-activating peptide
that has a photocleavable protecting group on the backbone amide. *FEBS Lett*
525:20–24. doi:10.1016/S0014-5793(02)03000-4

Thorpe S. 2007. An introduction to ocean turbulence. Cambridge University Press.

Wood CD, Darszon A, Whitaker M. 2003. Speract induces calcium oscillations in the
sperm tail. *J Cell Biol* **161**:89–101. doi:10.1083/jcb.200212053

Wood CD, Guerrero A, Priego-Espinosa DA, Martínez-Mekler G, Carneiro J,
Darszon A. 2015. Sea urchin sperm chemotaxis, Flagellar Mechanics and Sperm
Guidance. Bentham Science Publishers. doi:10.2174/9781681081281150101

Wood CD, Nishigaki T, Furuta T, Baba SA, Darszon A. 2005. Real-time analysis of
the role of Ca 2+ in flagellar movement and motility in single sea urchin sperm.
J Cell Biol **169**:725–731. doi:10.1083/jcb.200411001

Wood CD, Nishigaki T, Tatsu Y, Yumoto N, Baba SA, Whitaker M, Darszon A.
2007. Altering the speract-induced ion permeability changes that generate
flagellar Ca2+ spikes regulates their kinetics and sea urchin sperm motility. *Dev*
Biol **306**:525–537. doi:10.1016/j.ydbio.2007.03.036

Yoshida M, Yoshida K. 2011. Sperm chemotaxis and regulation of flagellar
movement by Ca2+. *Mol Hum Reprod* **17**:457–465. doi:10.1093/molehr/gar041

Zimmer RK, Riffell JA. 2011. Sperm chemotaxis, fluid shear, and the evolution of
sexual reproduction. *Proc Natl Acad Sci* **108**:13200–13205.
doi:10.1073/pnas.1018666108

Supplementary material. Sperm chemotaxis is driven by the slope of the

chemoattractant concentration field.

1. Theory

1.1. On the estimate of maximal chemoattractant absorption

Berg and Purcell (1977) derived a simple expression for the mean chemoattractant

binding and adsorption flux by a cell in the steady-state, denoted J:

$$J = 4\pi Da\bar{c}\frac{N}{N + \pi a/s} = J_{max}\frac{N}{N + \pi a/s} \tag{S1}$$

where D is the diffusion coefficient of the chemoattractant; a is the radius of the cell; $\bar{c}$ is

the mean concentration of the chemoattractant; N is the number of receptors on the

membrane of the cell; s is the effective radius of the receptor, assumed to be disk-like on the

cell surface and binding to chemoattractant molecules with high affinity; $J_{max} = 4\pi Da\bar{c}$,

which is the maximal flux of chemoattractant that a cell in the steady-state can experience;

and $\frac{N}{N + \pi a/s}$ is the probability that a molecule that has collided with the cell will find a

receptor (Berg and Purcell, 1977).

As pointed out by Berg and Purcell, the expression implies that for $N \gg \pi a/s =$

$N_{1/2}$ the flux of chemoattractant adsorption becomes $J \cong J_{max}$, which means that coverage

of only a small fraction of the cell surface by the receptors may lead to maximal flux. The

flux becomes practically independent of the number of receptors, proportional to the

concentration of the chemoattractant and limited by its diffusion, when receptor density is

sufficiently large. For a given chemoattractant concentration the half maximal flux ($J =$

$\frac{1}{2}J_{max}$) is reached when the number of receptors is $N = N_{1/2} = \pi a/s$.

It is worth getting a rough estimate of the number of receptors required for a maximal

influx of chemoattractant, in the specific case of spermatozoa by calculating $N_{1/2}$, assuming

a spherical cell with a surface area equivalent to that of the actual flagella. In the case of *S.*

purpuratus sperm, the flagellar width h $\cong$ 0.2 μm, and length L $\cong$ 40 μm that would give us

an approximate surface area A $\sim$ 25 μm^2 or an equivalent spherical radius $a_e \sim$ 1.4 μm and,

hence, a $J_{max} \cong 0.1 J_{max}^{sph}$, where J_{max}^{sph} represents the maximal influx of chemoattractants for

the spherical cell. The factor 0.1 relating the maximal flux J_{max} in a cylindrical flagellum

and J_{max}^{sph} in a spherical cell, arises by recalling that the expression for J_{max} in equation (**S1**).

The later stems from an analogy with electrostatics such that the total current depends on

the electrical capacitance C of the conducting material and, in particular, on the geometrical

arrangement. The capacitance of a simple spherical conductor equals the radius a of the

sphere but more generally we have $J_{max} = 4\pi C D \bar{c}$ (Berg and Purcell, 1977).

A more accurate computation can be obtained by taking into account the nearly cylindrical

shape of the flagellum. The capacitance of a finite cylinder can be obtained as a series

expansion in the logarithm of the cylinder aspect ratio $\Lambda = \ln(L/h)$ (Maxwell, 1877) and, to

a second order in 1/Λ, and is given by the following expression:

$$C = \frac{2\pi L}{\Lambda}\left[1 + \frac{1}{\Lambda}(1 - \ln 2) + \frac{1}{\Lambda^2}\left\{1 + (1 - \ln 2)^2 - \frac{\pi^2}{12}\right\} + O\left(\frac{1}{\Lambda^3}\right)\right] \qquad \textbf{(S2)}$$

For the case of the slender sperm flagellum with h/L << 1, $\Lambda \cong$ 5, this expression gives a

maximal influx for the cylinder that is, again, approximately one tenth that of the equivalent

sphere:

$$J_{max} \cong \frac{2\pi}{5} L D \bar{c} \cong 0.1 J_{max}^{sph} \qquad \textbf{(S3)}$$

The above description is valid only in the limit of instantaneous adsorption at the

receptors. For a finite rate of binding by the receptors we can simply modify the above

expressions to include an effective size for the binding sites $s_e = k_{on}/D$ (Phillips et al., 2012).

With $k_{on} = 24$ μM^{-1}s^{-1}, calculated for speract by Nishigaki in 2000 for *L. pictus* and in 2001

for *S. purpuratus* (Nishigaki et al., 2001; Nishigaki and Darszon, 2000), and $D = 240$ μm^{2}s$^-$

^{1}, $s_e = 1.9$ Å which is indeed much smaller than the physical size of the receptors (**Table SI**).

These equivalences were used to obtain the estimates in **Table SI**, which are discussed in

the main text. From these estimates, we can compute $N_{1/2} \sim 3\times10^4$ as the total number of SAP

receptors for the *S. purpuratus* sperm flagellum to act as a perfect absorber. As the actual

number of SAP receptors for this species is lower than that figure, *i.e.* $N < N_{1/2}$, we cannot

approximate the solution of equation **(S1)** to that of a perfect absorber. More specifically,

under these circumstances the absorption remains almost linearly dependent on the actual

number of receptors on the flagellum (**Figure S1**).

1.2. A condition for detecting a change in the chemoattractant concentration

A cell uses the chemoattractant it samples from the medium as a proxy of the

extracellular concentration of the chemoattractant at any given time. The flux of

chemoattractant J, calculated in the previous section, measures the sampling rate. Because

the number of chemoattractant molecules is finite and small, the actual number of molecules

sampled by the cell in an interval of time Δt is a random variable, denoted n, which is Poisson

distributed with expected value $E[n] = J\Delta$t and standard deviation $SD[n] = \sqrt{J\Delta t}$. The

chemotaxis signaling system of the spermatozoon should remain unresponsive while the cell

is swimming in an isotropic chemoattractant concentration field, as there are no spatial cues

for guidance (**Figure 1a**), although motility responses may still be triggered, as seen by the

stereotypical turn-and-run motility responses of *S. purpuratus* sperm in the presence of

speract (isotropic fields or weak gradients) – it has been previously reported that the turn-

and-run motility response is necessary, but not sufficient, for sea urchin sperm chemotaxis

(Wood et al., 2007). The Poisson fluctuations of sampled chemoattractant molecules,

measured by $\sqrt{J\Delta t}$ can be understood as background noise and hence should not elicit a

response. When the sperm is swimming confined to a plane in a chemoattractant gradient

produced by the egg (**Figures 1b** and **1c**), the chemotactic responses should be triggered only

when the amplitude of the sampling fluctuations are sufficiently large as to not be confused

with the background noise, i.e. when the difference in concentration at the two extremes of

the circular trajectory leads to fluctuations in chemoattractant sampling that are larger than

the background noise. These considerations lead to a minimal condition for reliable detection

of a chemotactic signal (Berg and Purcell, 1977; Vergassola et al., 2007), the corresponding

condition can be stated as:

82
$$\left(4\pi D a \bar{c} \, \frac{N}{N + \pi a/s} \Delta t \right) v \Delta t \frac{\partial c}{\partial r} \bar{c}^{-1} \; > \; \sqrt{4\pi D a \bar{c} \, \frac{N}{N + \pi a/s} \Delta t} \qquad \textbf{(S4)}$$

where $E[n] = J\Delta t = 4\pi D a \bar{c} \, \frac{N}{N + \pi a/s} \Delta t$; Δt is specifically the time the sperm takes to

make half a revolution in its circular trajectory; v is the mean linear velocity; and $\xi = \bar{c}^{-1} \frac{\partial c}{\partial r}$

is the relative slope of the chemoattractant concentration gradient.

As described in the main text, by interpreting the left-hand side of the equation **(S4)**

as the minimal chemotactic signal and the right-hand side as a measurement of the

background noise at a given mean concentration, one can obtain a minimal condition for the

smallest signal to noise ratio (*SNR*) necessary to elicit a chemotactic response. Equation **(S4)**

can be rewritten in terms of signal-to-noise ratio:

91
$$SNR = \ v\Delta t^{3/2} \left(4\pi Da\bar{c}\frac{N}{N+\pi a/s}\right)^{1/2} \xi > 1 \qquad \textbf{(S5)}$$

2. Extended Materials and Methods

2.1. Materials

Undiluted *S. purpuratus* or *L. pictus* spermatozoa (JAVIER GARCIA PAMANES,

Ensenada, Mexico and South Coast Bio-Marine San Pedro, CA 90731, USA respectively)

were obtained by intracoelomic injection of 0.5 M KCl and stored on ice until used within a

day. Artificial seawater (ASW) was 950 to 1050 mOsm and contained (in mM): 486 NaCl,

10 KCl, 10 CaCl$_2$, 26 MgCl$_2$, 30 MgSO$_4$, 2.5 NaHCO$_3$, 10 HEPES and 1 EDTA (pH 7.8).

For experiments with *L. pictus* spermatozoa, slightly acidified ASW (pH 7.4) was used to

reduce the number of spermatozoa experiencing spontaneous acrosome reaction. Low Ca^{2+}

ASW was ASW at pH 7.0 and with 1 mM CaCl$_2$, and Ca^{2+}-free ASW was ASW with no

added CaCl$_2$. [Ser5; nitrobenzyl‑Gly6]speract, referred to throughout the text as caged

speract (CS), was prepared as previously described (Tatsu et al., 2002). Fluo-4-AM and

pluronic F-127 were from Molecular Probes, Inc. (Eugene, OR, USA). PolyHEME [poly(2‑

hydroxyethylmethacrylate)] was from Sigma-Aldrich (Toluca, Edo de Mexico, Mexico).

2.2. Loading of Ca^{2+}-fluorescent indicator into spermatozoa

This was done as in Beltrán et al., 2014, as follows: undiluted spermatozoa were suspended

in 10 volumes of low Ca^{2+} ASW containing 0.2% pluronic F-127 plus 20 µM of fluo-4-AM

and incubated for 2.5 h at 14 ºC. Spermatozoa were stored in the dark and on ice until use.

2.3. Imaging of fluorescent swimming spermatozoa

The cover slips were briefly immersed into a 0.1% wt/vol solution of poly-HEME in ethanol,

hot-air blow-dried to rapidly evaporate the solvent, wash with distilled water twice followed

by ASW and mounted on reusable chambers fitting a TC-202 Bipolar temperature controller

(Medical Systems Corp.). The temperature plate was mounted on a microscope stage (Eclipse

TE-300; Nikon) and maintained at a constant 15 ℃. Aliquots of labeled sperm were diluted

in ASW and transferred to an imaging chamber (final concentration ~ $2x10^5$ cells ml^{-1}).

Epifluorescence images were collected with a Nikon Plan Fluor 40x 1.3 NA oil-immersion

objective using the Chroma filter set (ex HQ470/40x; DC 505DCXRU; em HQ510LP) and

recorded on a DV887 iXon EMCCD Andor camera (Andor Bioimaging, NC). Stroboscopic

fluorescence illumination was supplied by a Cyan LED no. LXHL-LE5C (Lumileds Lighting

LLC, San Jose, USA) synchronized to the exposure output signal of the iXon camera (2 ms

illumination per individual exposure). Images were collected with Andor iQ 1.8 software

(Andor Bioimaging, NC) at 30.80 fps in full-chip mode (observation field of ~200 x 200

µm).

2.4. Image processing

The background fluorescence was removed by generating an average pixel intensity time-

projection image from the first 94 frames (3 seconds) before uncaging, which was then

subtracted from each frame of the image stack by using the Image calculator tool of ImageJ

1.49u (Schneider et al., 2017). For **Figure 2d**, the maximum pixel intensity time projections

were created every 3 s from background-subtracted images before and after the UV flash.

2.5. Quantitation of global changes of spermatozoa number and $[Ca^{2+}]_i$

To study the dynamics of overall sperm motility and $[Ca^{2+}]_i$ signals triggered by the distinct

speract gradients we developed a segmentation algorithm that efficiently and automatically

detects the head of every spermatozoa in every frame of a given video-microscopy archive

(C/C++, OpenCV 2.4, Qt-creator 2.4.2). Fluorescence microscopy images generated as

described previously were used. The following steps summarize the work-flow of the

algorithm (**Figure S2**):

1. Segment regions of interest from background: This step consists of thresholding each

image (frame) of the video to segment the zones of interest (remove noise and atypical

values). Our strategy includes performing an automatic selection of a threshold value

for each Gaussian blurred image (I_G) ($\sigma = 3.5$ µm) considering the mean value (M_I)

and the standard deviation (SD_I) of the image I_G. The threshold value is defined by: T_I

$= M_I + 6SD_I$.

2. Compute the connected components: The connected components labeling is used to

detect connected regions in the image (a digital continuous path exists between all

pairs of points in the same component - the sperm heads). This heuristic consists of

visiting each pixel of the image and creating exterior boundaries using pixel neighbors,

accordingly to a specific type of connectivity.

3. Measure sperm head fluorescence. For each region of interest, identify the centroid in

the fluorescence channel (sperm head) and measure the mean value.

4. Compute the relative positions of the sperm heads within the imaging field, and assign

them to either R1, R2, R3 or R4 concentric regions around the centroid of the UV flash

intensity distribution. The radii of R1, R2, R3 or R4, were 25, 50, 75 and 100 µm,

respectively.

5. Repeat steps 1 to 4 in a frame-wise basis.

Step 1 of the algorithm filters out shot noise and atypical values; step 2 divides the images

into N connected components for the position of the sperm heads; step 3 quantitates sperm

head fluorescence, and finally step 4 computes the relative sperm position on the imaging

field. A similar approach has been recently used to identify replication centers of

adenoviruses in fluorescence microscopy images (Garcés et al., 2016).

We automatically analyzed 267 videos of *S. purpuratus* spermatozoa, each containing tens

of swimming cells, exposed to five distinct speract concentration gradients.

2.6. Analysis of speract induced Ca^{2+} transients with immobilized spermatozoa

Imaging chambers were prepared by coating cover slips with 50 µg/ml poly-D-lysine,

shaking off excess, and allowing to air-dry. Coated cover slips were then assembled into

imaging chambers. Fluo-4 labeled spermatozoa were diluted 1:40 in ASW, immediately

placed into the chambers, and left for 2 min, after which unattached sperm were removed by

washing with ASW. The chambers were then filled with 0.5 ml of ASW containing 500 nM

of caged speract and mounted in a TC-202 Bipolar temperature controller (Medical Systems

Corp.). Images were collected with Andor iQ 1.7 software (Andor Bioimaging, NC) at 90 fps

in full-chip mode, binning 4x4 (observation field of 200 µm x 200 µm). The imaging setup

was the same as that used for swimming spermatozoa. The caged speract was photo-released

with a 200 ms UV pulse delivered through an optical fiber (4 mm internal diameter) coupled

to a Xenon UV lamp (UVICO, Rapp Opto Electronic). The optical fiber was mounted on a

"defocused" configuration to minimize the generation of UV light heterogeneities.

Images were processed off-line using ImageJ 1.45s. Overlapping spermatozoa and any

incompletely adhered cells, which moved during the experiment, were ignored. Fluorescence

measurements in individual sperm were made by manually drawing a region of interest

around the flagella with the polygon selections tool of ImageJ.

2.7. Sperm swimming behavior in different chemoattractant gradients

The sperm swimming behavior in response to a chemoattractant concentration gradient can

be classified accordingly to their orientation angle (Θ), which is formed between their

reference and velocity vectors (**Figure 4b**). For the sake of simplicity, chemotactic drifts

(towards the source of the chemoattractant gradient) were considered to fall within the

category of ($\Theta < 60°$), the drift of swimming sperm in a direction perpendicular to the

gradient results from orientation angles falling within the range $60° \leq \Theta \leq 120°$. The instances

of negative chemotactic drifts (opposite to the source of the chemoattractant gradient) were

classified as those having higher orientation angles $\Theta > 120°$ (**Figure S8**).

The proportion of spermatozoa orientated with low Θ angles, i.e. towards the source of

the chemoattractant concentration gradient is enriched in those gradients that give

chemotactic responses: *f2 (p-value < 0.001)* and *f5 (p-value = 0.003); compare with f1 (p-*

value =0.2) and *f4 (p-value = 0.51)*. Statistical comparisons were performed with the

Pearson's Chi-squared test considering a probability of success of 1/3 for each type of

response (non-responding cells were not considered).

The two tested negative controls for chemotaxis (Low Ca^{2+} or High extracellular K^+

($[K^+]_e$) for *f2* gradient) showed a complete distinct distribution that the corresponding *f2*

gradient (*f2.0Ca p-value <0.001, f2.K p-value =0.01*, Fisher's exact test), i.e. as expected the

proportion of cells experiencing chemotactic drift was significantly reduced on the negative

controls.

Interestingly, the *f3* gradient provides the major stimulation of cell motility (the frequency of

non-responsive cells drops down to ~2%), however in this experimental condition the

proportion of cells responding towards the source of the chemoattractant gradient was not

significantly distinct from the other two types of responses *(p-value = 0.12*, Pearson's Chi-

squared test*)*.

In any tested gradient, the distributions of orientation angles have the same proportions

between perpendicular and opposite to the source responses: *f1 (p-value =0.63), f2 (p-value*

=1), f3 (p-value =0.4), f4 (p-value =0.84) and *f5 (p-value =0.15)*. Statistical comparisons

were performed with the exact binomial test considering a hypothesized probability of

success of 0.5.

2.8. Spontaneous vs speract-induced $[Ca^{2+}]_i$ oscillations

We characterized and compared the spontaneous vs the speract-induced Ca^{2+} oscillations

(**Figure S9**) and conclude that they are completely different phenomena. Spontaneous Ca^{2+}

oscillations are only observed in about 10% of the analyzed population of spermatozoa (see

Statistical analysis section in **Materials and Methods** on the manuscript). Most of the time

only one spontaneous oscillation is observed, and in the cases where more than one

spontaneous oscillation is present (which accounts for ~20% of the spontaneous oscillations,

i.e. only 2% of the total cells analyzed), they are significantly different in nature to the

speract-induced Ca^{2+} oscillations, judged as follows: they display a larger period and

amplitude (~ one order of magnitude) when compared to the speract induced oscillations

(**Figure S9c** and **S9d**). When these Ca^{2+} spontaneous oscillations occur, if not very large, the

cell will change direction randomly. If the oscillation is large enough, and is beyond a certain

$[Ca^{2+}]_i$ threshold, the cell stops swimming altogether (see for example: Wood et al., 2005;

Guerrero et al., 2013). After detection, we discarded cells undergoing spontaneous

oscillations in the present work.

Figure S1. Chemoattractant diffusive currents have a non-linear relationship to receptor coverage. For a spherical cell of radius a, with N disk-like receptors of effective radius s, the diffusive current saturates for $N \gg N_{1/2} = \pi a/s$ ($N_{1/2}$ is highlighted for *S. purpuratus*, *L. pictus* and *A. punctulata* with a green, blue and black dots, respectively). For *S. purpuratus*, however, the number of receptors is smaller than $N_{1/2}$ and the diffusive influx falls into an almost linear regime.

Figure S2. Automatic segmentation of swimming spermatozoa. Work-flow of the

segmentation algorithm: Video microscopy images were background subtracted by removing

the temporal average intensity projection (static fraction) of the first un-stimulated frames

(93 frames = 3 seconds) from the whole video (25 seconds). The resulting images were

convolved with a low-pass spatial frequency filter to reduce to a minimum typical noise

sources (detector, electronic, shot). The resulting images were thresholded to generate arrays

of regions of interest (ROIs), a heuristic search for connected components was then applied

to label single ROIs and to assign the corresponding pixels to unique spermatozoa. Scale bar

of 15 µm.

Figure S3. Sperm response to speract photo-release, collated data from individual experiments. Sperm motility responses to different speract concentration gradients (*f1, f2, f3, f4, f5*) at R1 (**a**), R2 (**b**), R3 (**c**) and R4 (**d**) concentric regions (see **Figure 3a**). Negative controls for sperm chemotaxis are artificial sea water with nominal 0 Ca^{2+} (Low Ca^{2+}); and artificial sea water with 40 mM of K^+ (High K^+). Each time trace represents the mean sperm density from up to 20 video microscopy experiments. Note that peak responses occurred around 6 seconds on the traces, some 3 seconds after speract exposure (indicated as vertical dotted lines). Purple vertical line indicates the UV flash (200 ms).

Figure S4. Modeling of the dynamics of speract gradient based on the UV light profile of distinct optical fibers. The radial profiles of the UV light scattered at the glass-liquid interface of *f1*, *f3*, *f4*, *f5* optical fibers are shown in gray. The speract gradient was generated

as in **figure 5**, but with the corresponding *f1* (**a** and **b**), *f3* (**c** and **d**), *f4* (**e** and **f**) and *f5* (**g** and

h) optical fibers. *Left panels* - The dynamics of the speract gradient computed as is in **figure**

5. The blue dashed line ($t_0 = 0$) corresponds to a Gaussian distribution fitted to the UV light

profile, and illustrates the putative shape of the instantaneously generated speract gradient.

Solid black lines illustrate the shape of the speract gradient after $t = 1, 2, 3, …, 20$ seconds.

Right panels - Simulated temporal changes in speract concentration gradients of *f1* (**a**), *f3* (**c**),

f4 (**e**) and *f5* (**g**) at each 10 µm radial point from the center of the concentration gradient.

Figure S5. Characteristic motility changes of a *S. purpuratus* spermatozoon exposed to *f3 and f4* speract gradients (chemotactic *vs* non-chemotactic response). Panels **a** and **b** show single cell responses to the *f3* speract gradient (chemotactic); and panels **c** and **d** to the *f4* speract gradient (non-chemotactic). **a, c.** Solid lines illustrate the spermatozoon swimming trajectory 3 s before (gray) and 6 s after (black) speract gradient exposure. **b, d.** Stimulus function computed from **a** and **c**, considering the spatio-temporal dynamics of speract computed for the *f3* and *f4* gradients, respectively.

Figure S6. Speract induces Ca^{2+} oscillations in immobilized *S. purpuratus* spermatozoa.
Spermatozoa were immobilized on cover slips coated with poly-D-lysine (see **Materials and Methods**), and ASW containing 500 nM caged speract added. Recordings were performed 3s before and during 6s after 200 ms of UV irradiation. *f4* optical fiber was used for the UV light path, to generate the speract concentration gradient. Time traces indicate the [Ca^{2+}]$_i$ of selected spermatozoa of **Movie S8**. Note that the photo-release of speract induces a train of [Ca^{2+}]$_i$ increases in immobilized spermatozoa, and hence provides evidence for the presence of an internal Ca^{2+} oscillator triggered by speract.

Figure S7. Cross-correlation analysis of $[Ca^{2+}]_i$ and stimulus function derivative (dS)

signals. Representative examples of $[Ca^{2+}]_i$ (red) and the derivative of the stimulus function

(dS) (black) were plotted and then analyzed by cross-correlation analysis (CCF). Red vertical

lines indicate the zero, blue vertical lines indicate the point of maximum correlation for each

case, which means that the phase shifting between $[Ca^{2+}]_i$ and dS is around 200 ms in these

cases. Examples of a pair of spermatozoa for the two principal chemotactic gradients (*f2* and

f3) are shown. **a, b.** Representative examples of two spermatozoa in an *f2* gradient. **c, d.**

Representative examples of two spermatozoa in an *f3* gradient.

Figure S8. Sperm swimming behavior in response to different chemoattractant gradients. *S. purpuratus* sperm behavior classified in four different classes: i) No response (purple), when spermatozoa keep swimming in concentric circles and do not move more than 15 µm of progressive swimming; ii) Towards the chemoattractant source (cyan), when spermatozoa respond drifting in swimming circles with an orientation angle (Θ) smaller than 60° ($\Theta < 60°$); iii) Opposite to the chemoattractant source (pink), when spermatozoa respond drifting in swimming circles with an orientation angle (Θ) bigger than 120° ($\Theta > 120°$); and iv) Perpendicular to the chemoattractant source (green), when spermatozoa respond drifting in swimming circles with an orientation angle (Θ) smaller than 120° but bigger than 60° ($60° \leq \Theta \leq 120°$). The orientation angle Θ is defined as the swimming angle of the drifting circles after speract uncaging (see **Figure 4b**). This analysis was made at second 6 after speract uncaging. Numbers in each bar represent the number of spermatozoa in each condition. Inset at the right panel shows the orientation angles (Θ). For further information see **Extended**

310   **Materials and Methods,** section *2.7. Sperm swimming behavior in different*

311   *chemoattractant gradients*.

Figure S9. Spontaneous *vs* speract-induced [Ca²⁺]ᵢ oscillations. a. Example of a

spontaneous Ca^{2+} oscillation (two oscillations). Only 20% of sperm experiencing

spontaneous oscillations suffer more than one oscillation, most of them experience only one

[Ca^{2+}]ᵢ increase. **b.** Representative trace of speract-induced oscillations. Caged-speract was

release after third second. **c.** Comparison of Ca^{2+} oscillation amplitude between spontaneous

(blue) *vs* speract-induced (red) oscillations. n = 75 for spontaneous and n = 56 for speract-

induced oscillations. **d.** Comparison of Ca^{2+} oscillation period between spontaneous (blue)

vs speract-induced (red) oscillations. n = 16 for spontaneous and n = 56 for speract-induced

oscillations. *Statistical significance, p < 0.01, Mann-Whitney *U* test. [Ca^{2+}]ᵢ oscillation

traces (panels **a** and **b**) were smoothed using an average filter with a 4 frame window in a

323   30.8 fps setting. For further information see **Extended Materials and Methods,** section *2.8.*

324   ***Spontaneous vs speract-induced $[Ca^{2+}]_i$ oscillations***.

325 **Table SI. Parameters of the chemoattractant sampling model for each species.**
326

	S. purpuratus	*L. pictus*	*A. punctulata*
N **[per cell]**	$2 \times 10^{4\,b}$	$6.3 \times 10^{4\,b}$	$3 \times 10^{5\,e}$
$N_{1/2}$	$2.9 \times 10^{4\,c}$	$3.6 \times 10^{4\,c}$	$1.8 \times 10^{4\,c}$
D **[cm^2 s^{-1}]**	$2.4 \times 10^{-6\,c}$	$2.4 \times 10^{-6\,c}$	$2.4 \times 10^{-6\,e}$
Kon **[M^{-1} s^{-1}]**	$2.7 \times 10^{-7\,b}$	$2.4 \times 10^{-7\,b}$	$5 \times 10^{-7\,f}$
s **[cm]**	$1.87 \times 10^{-8\,c}$	$1.66 \times 10^{-8\,c}$	$3.46 \times 10^{-8\,c}$
Δt [s]	0.39 ± 0.08 [a]	0.52 ± 0.22 [a]	0.6
v **[cm s^{-1}]**	$71.8 \times 10^{-4\,c}$	$88.5 \times 10^{-4\,c}$	$100 \times 10^{-4\,c}$
Δr [cm]	$28 \pm 6 \times 10^{-4\,a}$	$46 \pm 14 \times 10^{-4\,a}$	60×10^{-4}
L **[cm]**	$39.2 \pm 2.2 \times 10^{-4\,d}$	$48.7 \pm 2.1 \times 10^{-4\,d}$	$50 \times 10^{-4\,e}$
a **[cm]**	$1.4 \times 10^{-4\,c}$	$1.56 \times 10^{-4\,c}$	$1.58 \times 10^{-4\,c}$

327
Note that the main differences between species are the number of receptors N. $N_{1/2}$ number of

receptors that allows half maximal binding rate for any concentration of chemoattractant, i.e. $\pi a/s$. D

diffusion coefficient of the chemoattractant; Kon association rate constant; s effective radius of the

chemoattractant (as proxy of chemoattractant receptor's binding site radius); Δt sampling time (time

to swim half the circumference in the boundary close to the water-glass interface); v mean linear

speed of the spermatozoa, i.e. Δr/Δt; Δr sampling distance (circumference diameter); L length of

sperm flagellum; a spermatozoa radius, assuming that flagella are spheres. [a] Measured in this study

(mean ± sd); N = 3 sea urchins; n = 495 (*S. purpuratus*), n = 56 (*L. pictus*) spermatozoa. [b] Nishigaki

et al., 2001; Nishigaki and Darszon, 2000. [c] Calculated in this study (see section ***1.1. On the estimate***

of maximal chemoattractant absorption). [d] Measured in this study (mean ± sd); N = 1 sea urchin; n

= 26 (*S. purpuratus*), n = 39 (*L. pictus*) spermatozoa. [e] Kashikar et al., 2012. [f] Pichlo et al., 2014

reported 6.5×10^{-8} cm for the resact radius.

Table SII. Physical diameter of the optical fibers, and UV light power measured at the

back focal plane of the objective.

	Physical diameter (mm)	UV power at the back focal plane of the objective (mW)*
f1	0.2	0.07
f2	0.6	1.25
f3	2	4.7
f4	4	7.8
f5	4	9.46

*Typically, there is a 20% loss of light power between the back focal plane of the objective and the

sample, due to scattering within the optics.

Supplementary Movies

https://www.dropbox.com/s/v1h4kx6oqxc0f5l/Movie%202.avi?dl=0

Movie S1. Typical motility and Ca^{2+} responses of *S. purpuratus* spermatozoa towards

an *f3*-generated speract concentration gradient. An optical fiber of 2 mm internal diameter

(*f3*) was used for the UV light path to generate the speract concentration gradient. Other

imaging conditions were set up as for **Movie 1**. Note that spermatozoa located at R2, R3 and

R4 regions prior to speract exposure swim up the concentration field towards the center of

the gradient (R1). The pseudo-color scale represents the relative fluorescence of fluo-4, a

Ca^{2+} indicator, showing maximum (red) and minimum (blue) relative $[Ca^{2+}]_i$. Six *S.*

purpuratus spermatozoa were manually tracked for visualization purposes. Scale bar of 50

µm.

https://www.dropbox.com/s/27v2i3ofda57fqs/Movie%203.avi?dl=0

Movie S2. Chemotaxis of *S. purpuratus* spermatozoa requires extracellular calcium.

Spermatozoa swimming in artificial sea water with nominal calcium containing 10 nM caged

speract 3 s before and 5 s after exposure to 200 ms UV light. Nominal calcium disrupts the

electrochemical gradient required for Ca^{2+} influx, hence blocking the triggering of the

internal Ca^{2+} oscillation by speract. The *f2* fiber (0.6 mm diameter) was used to uncage

speract in this control. Other imaging conditions were set up as for **Movie 1**. Note that

spermatozoa re-located after speract uncaging but they failed to experience the Ca^{2+}-driven

motility alteration triggered by speract. As a consequence, they failed to experience

chemotaxis (compare with **Movie 1**). The pseudo-color scale represents the relative

fluorescence of fluo-4, a Ca^{2+} indicator, showing maximum (red) and minimum (blue)

relative $[Ca^{2+}]_i$. Six *S. purpuratus* spermatozoa were manually tracked for visualization

purposes. Scale bar of 50 µm.

https://www.dropbox.com/s/vft4aiw96fpf3sy/Movie%204.avi?dl=0

**Movie S3. Disrupting the K$^+$ electrochemical gradient blocks chemotaxis of *S.*

purpuratus spermatozoa.** Cells were swimming in artificial sea water containing 40 mM of

KCl, and 10 nM caged speract 3 s before and 5 s after exposure to 200 ms UV light. High K$^+$

in the ASW blocks the membrane potential hyperpolarization required for opening Ca^{2+}

channels, and hence prevents the triggering of the internal Ca^{2+} oscillator by speract

exposure. The *f2* fiber (0.6 mm diameter) was used to uncage speract in this control. Other

imaging conditions were set up as for **Movie 1**. Note that spermatozoa re-located after speract

uncaging but they failed to experience the Ca^{2+}-driven motility alteration triggered by

speract, and thus they failed to experience chemotaxis (compare with **Movie 1**). The pseudo-

color scale represents the relative fluorescence of fluo-4, a Ca^{2+} indicator, showing maximum

(red) and minimum (blue) relative [Ca^{2+}]$_i$. Six *S. purpuratus* spermatozoa were manually

tracked for visualization purposes. Scale bar of 50 µm.

https://www.dropbox.com/s/l3l7nmh1yw9juov/Movie%205.avi?dl=0

Movie S4. Typical motility and Ca^{2+} responses of _S. purpuratus_ spermatozoa towards

an _f1_-generated speract concentration gradient. An optical fiber of 0.2 mm internal

diameter (_f1_) was used for the UV light path to generate the speract concentration gradient.

Other imaging conditions were set up as for **Movie 1**. Note that some spermatozoa re-located

after speract uncaging but they failed to experience chemotaxis (compare with **Movie 1**). The

pseudo-color scale represents the relative fluorescence of fluo-4, a Ca^{2+} indicator, showing

maximum (red) and minimum (blue) relative $[Ca^{2+}]_i$. Six _S. purpuratus_ spermatozoa were

manually tracked for visualization purposes. Scale bar of 50 µm.

https://www.dropbox.com/s/qvwnyysbij6iz3b/Movie%206.avi?dl=0

Movie S5. Typical motility and Ca^{2+} responses of *S. purpuratus* spermatozoa towards

an *f4*-generated speract concentration gradient. An optical fiber of 4 mm internal diameter

(*f4*) was used for the UV light path to generate the speract concentration gradient. Other

imaging conditions were set up as for **Movie 1**. Note that spermatozoa re-located after speract

uncaging but they failed to experience chemotaxis (compare with **Movie 1**). The pseudo-

color scale represents the relative fluorescence of fluo-4, a Ca^{2+} indicator, showing maximum

(red) and minimum (blue) relative [Ca^{2+}]$_i$. Six *S. purpuratus* spermatozoa were manually

tracked for visualization purposes. Scale bar of 50 µm.

https://www.dropbox.com/s/o5o1e2jpmfitb1k/Movie%207.avi?dl=0

Movie S6. Typical motility and Ca^{2+} responses of *S. purpuratus* spermatozoa towards

an *f5*-generated speract concentration gradient. An optical fiber of 4 mm internal diameter

(*f5*) was used for the UV light path to generate the speract concentration gradient. Other

imaging conditions were set up as for **Movie 1**. Note that spermatozoa located at R2, R3 and

R4 regions prior to speract exposure swim up the speract concentration gradient, towards the

center of the imaging field (R1). The pseudo-color scale represents the relative fluorescence

of fluo-4, a Ca^{2+} indicator, showing maximum (red) and minimum (blue) relative [Ca^{2+}]$_i$. Six

S. purpuratus spermatozoa were manually tracked for visualization purposes. Scale bar of 50

µm.

https://www.dropbox.com/s/j4dlysh9k4r1lrp/Movie_10.avi?dl=0

Movie S7. Chemotactic index distributions. Radial histograms of chemotactic indices from

each different speract gradient. Black (*f2*) or red (rest) lines represent the median of each

distribution. This analysis was implemented from 4.5 seconds to 10 seconds. Speract

uncaging was induced at 3 seconds.

Movie S8. Photo-release of caged speract induces Ca^{2+} oscillations in immobilized *S.*

***purpuratus* spermatozoa.** Spermatozoa were immobilized, by coating the cover slip with

poly-D-lysine, in artificial sea water containing 500 nM caged speract, 3 s before and during

6 s after 200 ms of UV irradiation. The *f4* optical fiber was used for the UV light path to

generate the speract concentration gradient. The optical fiber was mounted in a "defocused"

configuration to minimize the generation of UV light heterogeneities. 93 frames s^{-1},

40x/1.3NA oil-immersion objective, 4x4 binning. The pseudo-color scale represents the

relative fluorescence of fluo-4, a Ca^{2+} indicator, showing maximum (red) and minimum

(blue) relative $[Ca^{2+}]_i$. The brightness and contrast scale was adjusted for better visualization

of $[Ca^{2+}]_i$ transients in the sperm flagella (as a consequence some heads look artificially

oversaturated, however no fluorescence saturation was observed in the raw data).

References

Beltrán C, Rodríguez-Miranda E, Granados-González G, García de De la Torre L, Nishigaki T, Darszon A. 2014. Zn2+ induces hyperpolarization by activation of a K+ channel and increases intracellular Ca2+ and pH in sea urchin spermatozoa. *Dev Biol* **394**:15–23. doi:10.1016/j.ydbio.2014.07.017

Berg HC, Purcell EM. 1977. Physics of chemoreception. *Biophys J* **20**:193–219. doi:10.1016/S0006-3495(77)85544-6

Garcés Y, Guerrero A, Hidalgo P, López RE, Wood CD, Gonzalez RA, Rendón-Mancha JM. 2016. Automatic detection and measurement of viral replication compartments by ellipse adjustment. *Sci Rep* **6**:1–14. doi:10.1038/srep36505

Guerrero A, Espinal J, Wood CD, Rendon JM, Carneiro J, Martinez-Mekler G, Darszon A, Rendón JM, Carneiro J, Martínez-Mekler G, Darszon A. 2013. Niflumic acid disrupts marine spermatozoan chemotaxis without impairing the spatiotemporal detection of chemoattractant gradients. *J Cell Sci* **126**:1477–87. doi:10.1242/jcs.121442

Kashikar ND, Alvarez L, Seifert R, Gregor I, Jäckle O, Beyermann M, Krause E, Kaupp UB, Benjamin Kaupp U. 2012. Temporal sampling, resetting, and adaptation orchestrate gradient sensing in sperm. *J Cell Biol* **198**:1075–1091. doi:10.1083/jcb.201204024

Maxwell C. 1877. On the Electrical Capacity of a long narrow Cylinder, and of a Disk of sensible Thickness. *Proc London Math Soc* **s1-9**:94–102. doi:10.1112/plms/s1-9.1.94

Nishigaki T, Darszon A. 2000. Real-Time Measurements of the Interactions between Fluorescent Speract and Its Sperm Receptor. *Dev Biol* **223**:17–26. doi:10.1006/dbio.2000.9734

Nishigaki T, Zamudio FZ, Possani LD, Darszon A. 2001. Time-resolved sperm responses to an egg peptide measured by stopped-flow fluorometry. *Biochem Biophys Res Commun* **284**:531–5. doi:10.1006/bbrc.2001.5000

Phillips R, Kondev J, Theriot J, Garcia H. 2012. Physical Biology of the Cell, 2nd ed. Garland Science.

Pichlo M, Bungert-Plümke S, Weyand I, Seifert R, Bönigk W, Strünker T, Kashikar ND, Goodwin N, Müller A, Körschen HG, Collienne U, Pelzer P, Van Q, Enderlein J, Klemm C, Krause E, Trötschel C, Poetsch A, Kremmer E, Kaupp UB. 2014. High density and ligand affinity confer ultrasensitive signal detection by a guanylyl cyclase chemoreceptor. *J Cell Biol* **206**:541–557. doi:10.1083/jcb.201402027

Schneider CA, Rasband WS, Eliceiri KW, Instrumentation C. 2017. NIH Image to ImageJ : 25 years of Image Analysis **9**:671–675.

Tatsu Y, Nishigaki T, Darszon A, Yumoto N. 2002. A caged sperm-activating peptide that has a photocleavable protecting group on the backbone amide. *FEBS Lett* **525**:20–24. doi:10.1016/S0014-5793(02)03000-4

Vergassola M, Villermaux E, Shraiman BI. 2007. 'Infotaxis' as a strategy for searching without gradients. *Nature* **445**:406–9. doi:10.1038/nature05464

Wood CD, Nishigaki T, Furuta T, Baba SA, Darszon A. 2005. Real-time analysis of the role of Ca 2+ in flagellar movement and motility in single sea urchin sperm. *J Cell Biol* **169**:725–731. doi:10.1083/jcb.200411001

Wood CD, Nishigaki T, Tatsu Y, Yumoto N, Baba SA, Whitaker M, Darszon A. 2007. Altering the speract-induced ion permeability changes that generate flagellar Ca2+ spikes regulates their kinetics and sea urchin sperm motility. *Dev Biol* **306**:525–537.