Title: **An optimized pipeline for parallel image-based quantification of gene expression and genotyping after *in situ* hybridization**

Running title: **Optimized ISH quantification**

Tomasz Dobrzycki[+,1], Monika Krecsmarik[+,1,2], Florian Bonkhofer[1], Roger Patient[1,2] and Rui Monteiro*[1,2]

1. MRC Molecular Haematology Unit, MRC Weatherall Institute of Molecular Medicine, John Radcliffe Hospital, University of Oxford, Oxford, OX3 9DS, United Kingdom
2. BHF Centre of Research Excellence, Oxford, United Kingdom

[+]These authors contributed equally.

*Contact details:

MRC Weatherall Institute of Molecular Medicine, John Radcliffe Hospital, University of Oxford, Oxford, OX3 9DS, United Kingdom

Email: rui.monteiro@imm.ox.ac.uk

Key words: zebrafish, mutant, bias, genotyping, image quantification

SUMMARY STATEMENT

Our optimized protocol to genotype zebrafish mutant embryos after *in situ* hybridization and digitally quantify the *in situ* signal will help to standardize existing experimental designs and methods of analysis.

ABSTRACT

Advances in genome engineering have resulted in the generation of numerous zebrafish mutant lines. A commonly used method to assess gene expression in the mutants is *in situ* hybridization (ISH). Because the embryos can be distinguished by genotype after ISH, comparing gene expression between wild type and mutant siblings can be done blinded and in parallel. Such experimental design reduces the technical variation between samples and minimises the risk of bias. This approach, however, requires an efficient method of genomic DNA extraction from post-ISH fixed zebrafish samples to ascribe phenotype to genotype. Here we describe a method to obtain PCR-quality DNA from 95-100% of zebrafish embryos, suitable for genotyping after ISH. In addition, we provide an image analysis protocol for quantifying gene expression of ISH-probed embryos, adaptable to analyse different expression patterns. Finally, we show that intensity-based image analysis enables accurate representation of the variability of gene expression detected by ISH and that it correlates well with quantitative methods like qRT-PCR. By combining genotyping after ISH and computer-based image analysis, we have established a high-confidence, unbiased methodology to assign gene expression levels to specific genotypes, and applied it to the analysis of molecular phenotypes of newly generated *lmo4a* mutants.

INTRODUCTION

The emergence of genome engineering technologies, including zinc-finger nucleases (ZFNs)(Doyon et al., 2008; Meng et al., 2008), transcription activator-like effector nucleases (TALENs)(Huang et al., 2011) and clustered regularly interspaced short palindromic repeats (CRISPR)/Cas9 (Hwang et al., 2013), has enabled zebrafish researchers to generate a wide range of mutant lines by precisely targeting genomic loci with high efficiency (Varshney et al., 2015). These methods provide alternatives to morpholino oligonucleotides (MOs), which have been used for gene knock down studies for over 15 years (Nasevicius and Ekker, 2000). In the light of recent concerns regarding the reliability of phenotypes induced by MOs (Kok et al., 2015), targeted gene knockouts have become a powerful tool to verify the transient MO phenotypes (Rossi et al., 2015) or to study genetic mutants, using MOs as a secondary tool (Gore et al., 2016). Low-cost protocols for DNA isolation (Meeker et

al., 2007) and for genotyping of zebrafish embryos (Wilkinson et al., 2013) have allowed faster generation and analysis of new mutant lines.

One of the most widely used methods to analyse molecular phenotypes during embryonic development is *in situ* hybridization (ISH). This technique is used to detect spatial expression patterns and tissue-specific changes in mRNA levels. Relevant protocols with several extensions have been standardised (Westerfield, 2007) and numerous validated probes are curated on the ZFIN database (Howe et al., 2013). Using ISH to analyse MO phenotypes requires processing of the MO-treated and control samples separately, which may result in an inherent bias and an increased risk of technical variation. Lack of reported measures to reduce the risk of bias has recently been exposed in a meta-analysis of *in vivo* animal studies (Macleod et al., 2015). Genomic mutants offer a way to overcome these issues. Wild type and mutant embryos can be processed in one sample, assessed phenotypically in a blinded manner and then distinguished based on their genotype. This way, the technical variation between samples is minimised and phenotypic assessment is largely bias-free. A few publications have reported approaches to genotype mutant lines after ISH, using either proteinase K treatment (Gore et al., 2016; Zhu et al., 2011) or commercial kits (Bresciani et al., 2014) for subsequent DNA extraction. However, the efficiency of these DNA extraction and genotyping methods has not been adequately demonstrated.

Reporting of phenotypes has usually been confined to one representative ISH image per condition, limiting the ability to quantitatively represent the variability of the phenotype. Approaches to score the expression levels as 'high', 'medium' or 'low' by eye (Blaser et al., 2017; Gao et al., 2016; Genthe and Clements, 2017; Peterkin et al., 2007; Place and Smith, 2017) are subjective and limited in how accurately they represent the effect of a knockout or a knockdown. Furthermore, visual scoring is inherently prone to poor reproducibility and low sensitivity. A more accurate way to quantitatively evaluate the change in gene expression levels involves counting the cells that contain the ISH signal (Espín-Palazón et al., 2014). However, this method is difficult to apply to compact anatomical structures and cell boundaries are hard to distinguish after ISH. Intensity-based image analysis using a selected region of interest (ROI) provides an objective alternative to visual scoring. Fan and colleagues have recently described a method using the ImageJ software to quantify ISH signal intensity in mouse embryos, where measurements were taken along a straight line drawn across the developing forelimb (Fan et al., 2015). Wen et al. have adapted this technique for use in the zebrafish to quantify gene expression levels in the mesoderm during early embryonic development (Wen et al., 2017). This approach can be further optimised to quantify gene expression after ISH in other regions and at other stages of the developing embryo.

Here we provide an optimised protocol for fast, inexpensive and highly efficient isolation of DNA from fixed zebrafish embryos for PCR and genotyping after ISH. This approach, based on the Hot Sodium Hydroxide and Tris (HotSHOT) method (Truett et al., 2000), is extremely reliable and allowed successful genotyping in 95-100% of the embryos. In addition, we propose a detailed step-by-step guide for gene expression intensity measurements based on modifications to the previously described quantitation methods (Fan et al., 2015; Wen et al., 2017). This pipeline (Fig. 1A) provides a tool for high-confidence, bias-free reporting of molecular phenotypes using standard ISH. We demonstrate its usefulness by assessing expression changes in *gpr65* morphants and *runx1* and *lmo4a* mutants.

RESULTS

HotSHOT genomic DNA extraction is suitable for PCR amplification in ISH-probed embryos

Our aim was to establish a fast and cost-effective way to extract DNA from individual embryos after ISH. First, we optimized the published HotSHOT protocol (Truett et al., 2000) for raw genomic DNA extraction, followed by JumpStart™ REDTaq® PCR. By testing amplicons designed for four different loci, we found that a 5-minute incubation time in the lysis buffer was sufficient to extract DNA to amplify fragments ranging from 144 to 600 base pairs (bp) by PCR (Fig. 1B). Increasing incubation time did not increase DNA yield (Fig. 1B). Next, we tested the 5-minute DNA extraction method on embryos at different stages by amplifying a 300bp fragment from the *runx1* locus. We successfully extracted DNA from fish ranging from 14hpf to 5dpf that had been stored for up to 2 years in glycerol at room temperature (Fig. 1C; Table 1). Incubation for less than 30 min was on average more efficient (allowing successful genotyping of more embryos) than incubation times longer than 65 min (Table 1). When comparing two different commercially available PCR master mixes, we found the JumpStart™ REDTaq® ReadyMix™ to be on average more efficient than Phire™ Green HotStart II PCR Master Mix (data not shown), likely due to differences in the optimal salt concentrations for the enzymes in each PCR mix.

Digital quantification of the ISH signal in the dorsal aorta reveals a significant decrease of dnmt3bb.1 mRNA levels in runx1$^{W84X/W84X}$ mutants and a significant increase of runx1 expression in gpr65 morphants

To demonstrate the usefulness of our genotyping protocol in a known mutant, we imaged 130 embryos from the incross of *runx1$^{+/W84X}$* heterozygotes (Jin et al., 2009), fixed at 33hpf and probed

for *dnmt3bb.1* mRNA, a known downstream target of *runx1* within the haemogenic endothelium (Gore et al., 2016) (Fig. 2A). Scoring of the images as 'high', 'medium' or 'low' showed a Mendelian 1:2:1 distribution of phenotypes (Fig. S1A). We then genotyped all the imaged embryos with 100% efficiency using the above protocol and RFLP (Fig. 2B). The observed Mendelian distribution of phenotypes, resulting from the first phenotypic assessment, did not entirely correspond to the respective genotypes. While the 'low' phenotype was significantly overrepresented in the homozygous mutant group (X^2=95.3, d.f.=4; p<0.001), there was no difference in the distribution of 'high'- and 'medium'-expressing embryos among wild type and heterozygous fish (X^2=1.35, d.f.=1; p>0.2) (Fig. S1B). To implement a more quantitative assessment, we used Fiji (Schindelin et al., 2012) to digitally quantify pixel intensities of each embryo in the dorsal aorta region along the yolk sac extension prior to genotyping. This approach relied on allocating two separate ROIs to each image: one containing the staining (yellow ROI, Fig. 1A) and another containing an equal area of the embryo without any staining (green ROI, Fig. 1A). By subtracting the background value from the staining value, a number was assigned to each embryo. After genotyping, we compared the signal intensity values in wild type, heterozygous and mutant embryos. As expected, the *runx1* mutant embryos showed a statistically significant reduction by approximately 50% of *dnmt3bb.1* signal compared to wild types or heterozygotes (μ_{wt}=54, μ_{het}=50.1, μ_{mut}=26.3; F=132.97, d.f.=2, 69.2; p<0.001). In contrast, there was no significant difference in signal intensity between wild types and heterozygotes (μ_{wt}=54, μ_{het}=50.1; p>0.3) (Fig. 2C). We found that the signal intensity values in all groups were very dispersed, with high coefficients of variation (24%, 22% and 21% for wild type, heterozygote and mutant groups, respectively).

We applied the same analysis method to the images of 16 *gpr65* morphants and 16 control (uninjected) siblings probed at 29hpf for *runx1* mRNA, previously shown to be downstream of *gpr65* (Gao et al., 2016). We found a significant increase of approximately 25% in pixel intensity levels of the *runx1* probe staining in *gpr65* morphants, compared to non-injected controls (μ_{wt}=29, μ_{gpr65}=37.2; t=2.38, d.f.=30; p<0.05) (Fig. 3). The values of the wild type embryos and the morphants showed dispersion with the coefficients of variation of 34% and 26%, respectively.

Digital quantification of runx1 ISH signal in the dorsal aorta reveals a discrepancy between the phenotype in lmo4a$^{\Delta25-29/\Delta25-29}$ mutants and previously described lmo4a morphants

We applied the ISH quantification and genotyping methods to analyse the phenotype of *lmo4a* mutants. We used TALENs to generate a 5bp deletion in the coding region of the *lmo4a* gene (*lmo4a$^{\Delta25-29}$*) (Fig. 4A). We then incrossed *lmo4a* heterozygous parents and quantified the expression of *runx1* in 28hpf embryos prior to genotyping them with 100% efficiency using RFLP (Fig. 4B).

MO-mediated knockdown of *lmo4a* was previously shown to result in reduced expression of *runx1* in the dorsal aorta at 30hpf (Meier et al., 2006). Surprisingly, we found no significant differences in *runx1* pixel intensity at 28hpf among wild type, heterozygous and homozygous siblings (μ_{wt}=30.3, μ_{het}=32.8, μ_{mut}=34.6; F=0.509, d.f.=2, 62; p>0.6) (Fig. 4C,D). The coefficients of variation of the values of the wild types, heterozygotes and homozygous mutants were 41%, 38% and 37%, respectively.

DISCUSSION

The HotSHOT method of genomic DNA isolation, originally designed for mouse ear notch samples (Truett et al., 2000), offers a fast and cost-effective way to genotype animals. While it has been used to isolate genomic DNA from PFA-fixed zebrafish samples before (Cooney et al., 2013; Meeker et al., 2007), its efficiency in this setting had not been reported. In consequence, the method has not been widely adopted in the community and many research groups rely on time-consuming and more expensive DNA extraction methods involving proteinase K treatment (Gore et al., 2016) or commercially available kits (Bresciani et al., 2014; Sood et al., 2013). Here we report that PCR-quality DNA can be rapidly extracted from 95-100% of fixed zebrafish embryos aged from 14hpf to 5dpf with an optimised HotSHOT protocol after ISH. This DNA can be subsequently used to genotype the samples with simple, inexpensive PCR followed by a digest to detect restriction enzyme sites disrupted by the mutation, as done previously for fresh tissue (Hruscha et al., 2013). To facilitate this, mutations can be designed to target restriction enzyme recognition sites. In fact, an online TALEN and CRISPR/Cas9 design tool Mojo Hand (Neff et al., 2012) readily provides restriction enzyme sites targeted by a desired mutation. While newly emerging alternatives to standard PCR and RFLP may provide a higher speed of genotyping (D'Agostino et al., 2016; Lee et al., 2016), the protocol described here for genotyping after ISH is attractive due to its high efficiency and demonstrated robustness in our hands.

Genotyping zebrafish embryos after ISH is important because it allows processing of mutant and wild type embryos in one batch, therefore limiting the technical variation between samples. In addition, expression levels of the target gene can be assessed in a non-biased way, because the embryos can be distinguished by their genotype only after phenotypic assessment. This is a powerful way to control for unconscious bias, a serious issue in *in vivo* animal research (Macleod et al., 2015). Assessing mRNA levels in post-ISH embryos has been performed visually, either by scoring the phenotypes into discrete groups (Blaser et al., 2017; Gao et al., 2016; Genthe and Clements, 2017) or by cell counting (Espín-Palazón et al., 2014). However, these approaches are prone to subjectivity and poor reproducibility. Furthermore, visual scoring can be difficult to carry out and interpret due to expression level differences between individuals of the same genotype. Indeed, we show here

that pixel intensities of the ISH signal in wild type embryos probed for the transcription factor *runx1* show high dispersion in wild type embryos with over 25% and up to 40% coefficient of variation (Figs 3B, 4D). These results indicate that the interpretation of phenotypes based purely on expected Mendelian distribution from heterozygous incrosses might be misleading. As we demonstrate, visual scoring of embryos from a heterozygous *runx1^(+/W84X)* incross into 'high', 'medium' and 'low' groups based on *dnmt3bb.1* expression levels gives a phenotypic Mendelian distribution of 1:2:1, which could suggest a haploinsufficiency effect on the regulation of *dnmt3bb.1* expression. Genotyping of these embryos revealed that the vast majority of 'low'-expressing ones were indeed genetically homozygous mutant. However, both 'high'- and 'medium'-expressing embryos were distributed similarly across wild type and heterozygous fish, disproving the haploinsufficiency hypothesis. As a possible explanation for this discrepancy, we found that the signal intensity values in all three genotypes were highly dispersed, with coefficients of variation over 20%. Therefore, each ISH experiment done on embryos from a heterozygous incross should be followed by genotyping to avoid misleading conclusions due to the variability of the ISH signal intensities in embryos of the same genotype.

We would argue that the use of digital image analysis on ISH-probed samples is critical for objective, statistical demonstration of changes in expression levels. Here we describe an imaging protocol based on previous studies (Fan et al., 2015; Wen et al., 2017) to measure gene expression intensity in the trunk region of 1-2 day old zebrafish embryos. We show that the average ISH staining intensity for *dnmt3bb.1* mRNA is significantly decreased in *runx1* mutants compared to wild type siblings, in agreement with previously reported qRT-PCR quantitation of *dnmt3bb.1* levels in whole embryos (Gore et al., 2016). Thus, our method is robust and should be adopted instead of less reliable visual scoring methods. It could also be used as an alternative to qRT-PCR experiments where these require larger numbers of animals and are prone to errors due to a limited number of highly reliable internal controls (Xu et al., 2016). Our quantification method addresses all of these limitations. Furthermore, it presents a way to measure changes in expression levels in a very tissue-specific manner, which is useful in the case of genes with multiple developmental roles. We believe it will be particularly helpful for studying other genes with expression patterns that are spatially restricted, such as *gata2b*, a haematopoietic gene expressed in the ventral wall of the dorsal aorta (Butko et al., 2015). An alternative modification to our method could involve subtracting the average intensity of another stained region from the intensity of the selected ROI – for instance, using the *runx1* or *dnmt3bb.1* signal in the head as an internal control. However, expression levels vary widely between different tissues and there is a risk of saturating the signal, reducing the dynamic range used for the

comparisons. Therefore, we believe that using an unstained region with the same area as a background measurement provides a more reliable way to quantify the ISH signal in each embryo.

We also propose a way to quantitatively represent variation in gene expression levels without relying on subjective and biased scoring. For instance, we could replicate the previously reported increase in *runx1* expression in *gpr65* morphants (Gao et al., 2016) with our method, but we represented it in a more objective quantitative way, also allowing powerful statistical analysis. In fact, we achieved 63% power to detect a 25% increase in pixel intensity at the *p*<0.05 level for sample sizes as small as 16 for each condition. This method of analysis also allows precise calculations of required sample sizes to achieve given power. In the presented example, 80% power would require 24 MO-injected embryos and 24 uninjected controls. Such calculations are essential in animal research (Dell et al., 2002), but they are notoriously unreported (Macleod et al., 2015). In fact, recently published updated guidelines for the use of MOs in zebrafish encourage statistical analysis of phenotypes and advocate the use of blinded assessment (Stainier et al., 2017) and both points can be addressed with our method. In addition, there is scope to automate the phenotypical analysis using this method. We have generated a batch conversion Fiji macro that converts all images in a given file directory (see Supplementary Detailed Protocol) ready for manual ROI selection. Future optimisations may involve automation of intensity measurements of the ISH images (Chen et al., 2011).

We have further applied the described method to analyse the molecular phenotype of an unpublished *lmo4a* mutant. In contrast to previous reports showing decreased *runx1* expression in *lmo4a* morphants (Meier et al., 2006), we found no difference in *runx1* pixel intensity levels between wild types, heterozygotes and *lmo4a* homozygous mutants. This discrepancy may result from compensatory mechanisms activated in mutants but not morphants, a phenomenon now frequently reported in zebrafish (El-Brolosy and Stainier, 2017). Potential compensation could come from the *lmo4b* paralogue or from *lmo2*, another gene from the *lmo* family that is expressed in the dorsal aorta and required for the specification of the haematopoietic programme (Patterson et al., 2007).

METHODS

Maintenance of zebrafish and morpholino oligonucleotide injections

All animal experiments were approved by the local ethics committee. Wild type and *runx1*[W84X] (Jin et al., 2009) and *lmo4a*[Δ25-29] mutant zebrafish (*Danio rerio* Hamilton) were maintained and bred according to standard procedures (Westerfield, 2007). Embryos were collected by natural mating and staged according to morphological features (Kimmel et al., 1995) corresponding to respective

age in hours or days post fertilisation (hpf or dpf, respectively). For *gpr65* knockdown, wild type one-cell stage embryos were injected with 4ng of GPR65_SP MO (Gao et al., 2016).

Generation of lmo4a mutants

For TALENs design, Mojo Hand software was used (http://www.talendesign.org)(Neff et al., 2012). The identified target site was GGAAAGCTCCGCGGTT. The RVD-containing repeats were assembled using the Golden Gate approach in pTAL3-DDD and pTAL3-RRR vectors (Cermak et al., 2011). The resulting DNA templates were verified by sequencing and linearised with NotI enzyme. The mRNAs were transcribed from 1µg linearised template with SP6 mMessage mMachine® kit (Ambion) and purified with RNeasy® Micro Kit (QIAGEN).

One-cell zebrafish embryos were injected with 100pg left-arm + 100pg right-arm TALEN mRNAs. Germline mutations in the founders (*lmo4a*$^{+/-}$) were identified using restriction fragment length polymorphism (RFLP)(Bedell et al., 2012) (see primer sequences below) with SacII enzyme on the genomic DNA extracted from their offspring. Mutations were identified by the presence of an undigested PCR product on a 2% agarose gel. The undigested mutant fragments were purified from the gel, cloned into pGEM®-T Easy vector (Promega) and sequenced. The *lmo4a*$^{+/-}$ founders were outcrossed to generate heterozygous carriers, identified with fin-clipping and genotyping at 3dpf as described previously (Wilkinson et al., 2013).

Whole-mount in situ hybridization

ISH was carried out according to the standard lab protocol (Jowett and Yan, 1996) using digoxygenin-labelled *dnmt3bb.1* (Gore et al., 2016) and *runx1* (Kalev-Zylinska et al., 2002) probes. Post hybridization, the embryos were bleached in 5% formamide/0.5% SSC/10% H_2O_2 (Monteiro et al., 2011) and imaged in 100% glycerol with QImaging MicroPublisher 5.0 RTV Camera and Q-Capture Pro 7™ software (version 7.0.3), using the same exposure, magnification and illumination settings for each embryo.

DNA extraction and genotyping

Genomic DNA was isolated from 4% paraformaldehyde(PFA)-fixed embryos using the original HotSHOT protocol (Truett et al., 2000) (Fig. 1a). Briefly, 40-75 µl of lysis buffer (25 mM NaOH, 0.2 mM EDTA) was added directly to a PCR tube with a freshly-imaged embryo in <5 µl 100% glycerol. To test the efficiency of the DNA extraction, embryos were suspended in the buffer and incubated at 95°C for 5-120 minutes, then cooled to 4°C, after which an equal volume of neutralisation buffer (40-75 µl 40 mM Tris-HCl) was added (see detailed Supplementary protocol). Genomic regions containing the mutated sites in the *runx1* locus were amplified with JumpStart™ REDTaq®

ReadyMix™ PCR or with Phire™ Green HotStart II PCR Master Mix according to manufacturer protocols, using 5 µl of DNA lysate in a 20 µl reaction volume and the following primer sequences: 5'-GCTCTGGTGGGCAAACTG-3' and 5'-CATGTGTTTGGACTGTGGGG-3' for *runx1*; and 5'-ACTTTGCCTCTGGATCTGCT-3' and 5'-TCTACATCAGTCCCGCCAAA-3' for *lmo4a*. The presence of *runx1* and *lmo4a* mutations were verified by restriction fragment length polymorphism (RFLP)(Hruscha et al., 2013) with HaeII or SacII enzymes respectively on a 2% agarose gel, using 100bp DNA Ladder as a reference. The sequences of additional primers used were: 5'-GCACCACAGTGGACATTGAT-3' and 5'-GTTGTAGAGGGCCAGCACTT-3' for the *tgfß3* locus; 5'-CATTAATGCGAGGGATACGG-3' and 5'-AAAAGAGCCACGGTAGGTGA-3' for the *tgfßr2* locus; and 5'-TGGCTAAGTGACCGTCAGAG-3' and 5'-TGAAACAAAACGCAGACGAC-3' for the *gata2a* locus.

Digital image analysis

Using Fiji software (Schindelin et al., 2012), the images were inverted to negative and converted to 8-bit grayscale. A Region of Interest (ROI) containing the ISH expression signal in the dorsal aorta along the yolk sac extension was drawn manually for each embryo. Then a second ROI with the same shape and area was created in a region of the embryo that has a uniform intensity and does not contain any ISH staining. In this particular instance, this region was placed just above the notochord (Fig. 1A). This area was used to define the background. A value for each region was then determined by measuring the average pixel intensity. After subtracting the value of the background region from the value of the stained region, the pixel intensity of the ISH signal was assigned to each embryo (Fig. 1A and detailed Supplementary protocol).

Statistical analysis

The numbers of embryos scored as 'high', 'medium' or 'low' were tested for equal distribution among wild type, heterozygous and mutant genotypes with contingency Chi-squared tests, applying Continuity Correction for 2x2 tables. For digitally analysed images, the pixel intensity values were assessed for normal distribution with Q-Q plots and transformed with sqrt function if necessary. Mean values (μ) of each experimental group were analysed with 2-tailed independent-samples *t*-test (for *gpr65* MO experiment) or with ANOVA (for mutant experiments) with 95% confidence levels, testing for the equality of variances with a Levene's test and applying the Welch correction when necessary. For ANOVA, differences between each two groups were assessed with either Tukey's post-hoc test (for equal variances) or with Games-Howell test (for unequal variances). For all these analyses the IBM® SPSS® Statistics (version 22) package was used. The degree of variability in each sample was assessed by calculating the coefficients of variation, defined as $\frac{s_{(x)}}{\mu}$, with $s_{(x)}$ being the standard deviation. The post-hoc power of the tests and required sample sizes were determined

with G*Power software (version 3.0.10)(Faul et al., 2007). The graphs presenting individual data points, means and ±s.d. were plotted using GraphPad Prism 7.

ACKNOWLEDGEMENTS

We thank Dr Dominic Waithe from the Wolfson Imaging Centre for advice on developing the image quantification method. We thank Rossella Rispoli for help with generating the batch conversion Fiji macro. We thank staff members of Biomedical Services at the John Radcliffe Hospital for monitoring and feeding zebrafish.

COMPETING INTERESTS

No competing interests declared.

FUNDING

T.D. was funded by a Wellcome Trust Chromosome and Developmental Biology PhD Scholarship (#WT102345/Z/13/Z). R.M. and M.K. were funded by the British Heart Foundation (BHF IBSR Fellowship FS/13/50/30436). R.K.P. and F.B. were funded by the Medical Research Council. R.M. acknowledges support from the BHF Centre of Research Excellence (RE/13/1/30181), Oxford.

DATA AVAILABILITY

All data generated or analysed during this study are included in this published article (and its Supplementary Information files).

REFERENCES

Bedell, V. M., Wang, Y., Campbell, J. M., Poshusta, T. L., Starker, C. G., Krug, R. G., 2nd, Tan, W., Penheiter, S. G., Ma, A. C., Leung, A. Y., et al. (2012). In vivo genome editing using a high-efficiency TALEN system. *Nature* **491**, 114-118.

Blaser, B. W., Moore, J. L., Hagedorn, E. J., Li, B., Riquelme, R., Lichtig, A., Yang, S., Zhou, Y., Tamplin, O. J., Binder, V., et al. (2017). CXCR1 remodels the vascular niche to promote hematopoietic stem and progenitor cell engraftment. *J. Exp. Med.* **214**, 1011-1027.

Bresciani, E., Carrington, B., Wincovitch, S., Jones, M., Gore, A. V., Weinstein, B. M., Sood, R. and Liu, P. P. (2014). CBFβ and RUNX1 are required at 2 different steps during the development of hematopoietic stem cells in zebrafish. *Blood* **124**, 70-78.

Butko, E., Distel, M., Pouget, C., Weijts, B., Kobayashi, I., Ng, K., Mosimann, C., Poulain, F. E., McPherson, A., Ni, C. W., et al. (2015). Gata2b is a restricted early regulator of hemogenic endothelium in the zebrafish embryo. *Development* **142**, 1050-1061.

Cermak, T., Doyle, E. L., Christian, M., Wang, L., Zhang, Y., Schmidt, C., Baller, J. A., Somia, N. V., Bogdanove, A. J. and Voytas, D. F. (2011). Efficient design and assembly of custom TALEN and other TAL effector-based constructs for DNA targeting. *Nucleic Acids Res.* **39**, e82.

Chen, S., Zhu, Y., Xia, W., Xia, S. and Xu, X. (2011). Automated analysis of zebrafish images for phenotypic changes in drug discovery. *J. Neurosci. Methods* **200**, 229-236.

Cooney, J. D., Hildick-Smith, G. J., Shafizadeh, E., McBride, P. F., Carroll, K. J., Anderson, H., Shaw, G. C., Tamplin, O. J., Branco, D. S., Dalton, A. J., et al. (2013). Teleost growth factor independence (gfi) genes differentially regulate successive waves of hematopoiesis. *Dev. Biol.* **373**, 431-441.

D'Agostino, Y., Locascio, A., Ristoratore, F., Sordino, P., Spagnuolo, A., Borra, M. and D'Aniello, S. (2016). A Rapid and Cheap Methodology for CRISPR/Cas9 Zebrafish Mutant Screening. *Mol. Biotechnol.* **58**, 73-78.

Dell, R. B., Holleran, S. and Ramakrishnan, R. (2002). Sample size determination. *Ilar Journal* **43**, 207-213.

Doyon, Y., McCammon, J. M., Miller, J. C., Faraji, F., Ngo, C., Katibah, G. E., Amora, R., Hocking, T. D., Zhang, L., Rebar, E. J., et al. (2008). Heritable targeted gene disruption in zebrafish using designed zinc-finger nucleases. *Nat. Biotechnol.* **26**, 702-708.

El-Brolosy, M. A. and Stainier, D. Y. (2017). Genetic compensation: A phenomenon in search of mechanisms. *PLoS genetics* **13**.

Espín-Palazón, R., Stachura, D. L., Campbell, C. A., García-Moreno, D., Del Cid, N., Kim, A. D., Candel, S., Meseguer, J., Mulero, V. and Traver, D. (2014). Proinflammatory signaling regulates hematopoietic stem cell emergence. *Cell* **159**, 1070-1085.

Fan, Y., Richelme, S., Avazeri, E., Audebert, S., Helmbacher, F., Dono, R. and Maina, F. (2015). Tissue-specific gain of RTK signalling uncovers selective cell vulnerability during embryogenesis. *PLoS Genet* **11**, e1005533.

Faul, F., Erdfelder, E., Lang, A.-G. and Buchner, A. (2007). G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav. Res. Methods* **39**, 175-191.

Gao, X., Wu, T., Johnson, K. D., Lahvic, J. L., Ranheim, E. A., Zon, L. I. and Bresnick, E. H. (2016). GATA Factor-G-Protein-Coupled Receptor Circuit Suppresses Hematopoiesis. *Stem cell reports* **6**, 368-382.

Genthe, J. R. and Clements, W. K. (2017). R-spondin 1 is required for specification of hematopoietic stem cells through Wnt16 and Vegfa signaling pathways. *Development* **144**, 590-600.

Gore, A. V., Athans, B., Iben, J. R., Johnson, K., Russanova, V., Castranova, D., Pham, V. N., Butler, M. G., Williams-Simons, L., Nichols, J. T., et al. (2016). Epigenetic regulation of hematopoiesis by DNA methylation. *eLife* **5**.

Howe, D. G., Bradford, Y. M., Conlin, T., Eagle, A. E., Fashena, D., Frazer, K., Knight, J., Mani, P., Martin, R., Moxon, S. A., et al. (2013). ZFIN, the Zebrafish Model Organism Database: increased support for mutants and transgenics. *Nucleic Acids Res.* **41**, 60.

Hruscha, A., Krawitz, P., Rechenberg, A., Heinrich, V., Hecht, J., Haass, C. and Schmid, B. (2013). Efficient CRISPR/Cas9 genome editing with low off-target effects in zebrafish. *Development* **140**, 4982-4987.

Huang, P., Xiao, A., Zhou, M., Zhu, Z., Lin, S. and Zhang, B. (2011). Heritable gene targeting in zebrafish using customized TALENs. *Nat. Biotechnol.* **29**, 699-700.

Hwang, W. Y., Fu, Y., Reyon, D., Maeder, M. L., Tsai, S. Q., Sander, J. D., Peterson, R. T., Yeh, J. R. and Joung, J. K. (2013). Efficient genome editing in zebrafish using a CRISPR-Cas system. *Nat. Biotechnol.* **31**, 227-229.

Jin, H., Sood, R., Xu, J., Zhen, F., English, M. A., Liu, P. P. and Wen, Z. (2009). Definitive hematopoietic stem/progenitor cells manifest distinct differentiation output in the zebrafish VDA and PBI. *Development* **136**, 647-654.

Jowett, T. and Yan, Y. L. (1996). Double fluorescent in situ hybridization to zebrafish embryos. *Trends Genet.* **12**, 387-389.

Kalev-Zylinska, M. L., Horsfield, J. A., Flores, M. V., Postlethwait, J. H., Vitas, M. R., Baas, A. M., Crosier, P. S. and Crosier, K. E. (2002). Runx1 is required for zebrafish blood and vessel development and expression of a human RUNX1-CBF2T1 transgene advances a model for studies of leukemogenesis. *Development* **129**, 2015-2030.

Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. and Schilling, T. F. (1995). Stages of embryonic development of the zebrafish. *Dev. Dyn.* **203**, 253-310.

Kok, F. O., Shin, M., Ni, C. W., Gupta, A., Grosse, A. S., van Impel, A., Kirchmaier, B. C., Peterson-Maduro, J., Kourkoulis, G., Male, I., et al. (2015). Reverse genetic screening reveals poor correlation between morpholino-induced and mutant phenotypes in zebrafish. *Dev. Cell* **32**, 97-108.

Lee, H. B., Schwab, T. L., Koleilat, A., Ata, H., Daby, C. L., Cervera, R., McNulty, M. S., Bostwick, H. S. and Clark, K. J. (2016). Allele-Specific Quantitative PCR for Accurate, Rapid, and Cost-Effective Genotyping. *Hum. Gene Ther.* **27**, 425-435.

Macleod, M. R., McLean, A., Kyriakopoulou, A., Serghiou, S., de Wilde, A., Sherratt, N., Hirst, T., Hemblade, R., Bahor, Z., Nunes-Fonseca, C., et al. (2015). Risk of Bias in Reports of In Vivo Research: A Focus for Improvement. *PLoS Biol.* **13**, e1002273.

Meeker, N., Hutchinson, S., Ho, L. and Trede, N. (2007). Method for isolation of PCR-ready genomic DNA from zebrafish tissues. *Biotechniques* **43**, 610-614.

Meier, N., Krpic, S., Rodriguez, P., Strouboulis, J., Monti, M., Krijgsveld, J., Gering, M., Patient, R., Hostert, A. and Grosveld, F. (2006). Novel binding partners of Ldb1 are required for haematopoietic development. *Development* **133**, 4913-4923.

Meng, X., Noyes, M. B., Zhu, L. J., Lawson, N. D. and Wolfe, S. A. (2008). Targeted gene inactivation in zebrafish using engineered zinc-finger nucleases. *Nat. Biotechnol.* **26**, 695-701.

Monteiro, R., Pouget, C. and Patient, R. (2011). The gata1/pu.1 lineage fate paradigm varies between blood populations and is modulated by tif1γ. *EMBO J.* **30**, 1093-1103.

Nasevicius, A. and Ekker, S. C. (2000). Effective targeted gene 'knockdown' in zebrafish. *Nat. Genet.* **26**, 216-220.

Neff, K. L., Argue, D. P., Ma, A. C., Lee, H. B., Clark, K. J. and Ekker, S. C. (2012). Mojo Hand, a TALEN design tool for genome editing applications. *BMC Bioinformatics* **14**, 1-7.

Patterson, L. J., Gering, M., Eckfeldt, C. E., Green, A. R., Verfaillie, C. M., Ekker, S. C. and Patient, R. (2007). The transcription factors Scl and Lmo2 act together during development of the hemangioblast in zebrafish. *Blood* **109**, 2389-2398.

Peterkin, T., Gibson, A. and Patient, R. (2007). Redundancy and evolution of GATA factor requirements in development of the myocardium. *Dev. Biol.* **311**, 623-635.

Place, E. S. and Smith, J. C. (2017). Zebrafish atoh8 mutants do not recapitulate morpholino phenotypes. *PLoS One* **12**, e0171143.

Rossi, A., Kontarakis, Z., Gerri, C., Nolte, H., Hölper, S., Krüger, M. and Stainier, D. Y. R. (2015). Genetic compensation induced by deleterious mutations but not gene knockdowns. *Nature* **524**, 230-233.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. *Nature methods* **9**, 676-682.

Sood, R., Carrington, B., Bishop, K., Jones, M., Rissone, A., Candotti, F., Chandrasekharappa, S. C. and Liu, P. (2013). Efficient methods for targeted mutagenesis in zebrafish using zinc-finger nucleases: data from targeting of nine genes using CompoZr or CoDA ZFNs. *PLoS One* **8**, e57239.

Stainier, D. Y. R., Raz, E., Lawson, N. D., Ekker, S. C., Burdine, R. D., Eisen, J. S., Ingham, P. W., Schulte-Merker, S., Yelon, D., Weinstein, B. M., et al. (2017). Guidelines for morpholino use in zebrafish. *PLOS Genetics* **13**.

Truett, G. E., Heeger, P., Mynatt, R. L., Truett, A. A., Walker, J. A. and Warman, M. L. (2000). Preparation of PCR-quality mouse genomic DNA with hot sodium hydroxide and tris (HotSHOT). *Biotechniques* **29**, 52-54.

Varshney, G. K., Sood, R. and Burgess, S. M. (2015). Understanding and Editing the Zebrafish Genome. *Adv. Genet.* **92**, 1-52.

Wen, B., Yuan, H., Liu, X., Wang, H., Chen, S., Chen, Z., de The, H., Zhou, J. and Zhu, J. (2017). GATA5 SUMOylation is indispensable for zebrafish cardiac development. *Biochim. Biophys. Acta*.

Westerfield, M. (2007). *The zebrafish book. A guide for the laboratory use of zebrafish (Danio rerio).* (5th edn). Eugene, OR: Univ of Oregon Press.

Wilkinson, R. N., Elworthy, S., Ingham, P. W. and van Eeden, F. J. M. (2013). A method for high-throughput PCR-based genotyping of larval zebrafish tail biopsies. *Biotechniques* **55**.

Xu, H., Li, C., Zeng, Q., Agrawal, I., Zhu, X. and Gong, Z. (2016). Genome-wide identification of suitable zebrafish Danio rerio reference genes for normalization of gene expression data by RT-qPCR. *J. Fish Biol.*

Zhu, C., Smith, T., McNulty, J., Rayla, A. L., Lakshmanan, A., Siekmann, A. F., Buffardi, M., Meng, X., Shin, J., Padmanabhan, A., et al. (2011). Evaluation and application of modularly assembled zinc-finger nucleases in zebrafish. *Development* **138**, 4555-4564.

Figure 1. Overview of the method to extract DNA for genotyping zebrafish mutants after ISH and measure the mRNA levels. A) Embryos collected from an incross of fish heterozygous for a mutant allele are probed for the measured gene with a standard ISH protocol. After imaging in 100% glycerol, genomic DNA is extracted using the HotSHOT protocol by adding the lysis buffer directly to the embryo in a 0.2ml PCR tube, followed by a ≥5 min incubation at 95°C. This DNA is used for genotyping of the embryos by PCR and restriction fragment length polymorphism (RFLP). In parallel, the images for each embryo are inverted and converted to 8-bit greyscale. ROIs of identical shape and size containing the ISH signal (yellow) and background (green) are manually selected and measured. The measurements, assigned to corresponding genotypes, are statistically analysed. B) DNA fragments amplified from single fixed and ISH-probed embryos after incubation in HotSHOT lysis buffer for 5, 15, 30, 45, 60 and 120 min followed by JumpStart™ REDTaq® ReadyMix™ PCR. Primer pairs designed for the *gata2a*, *runx1*, *tgfbr2b* and *tgfb3* produced 600, 300, 176 and 144 bp fragments, respectively. Three wild type embryos were used for each time point. First lane from the left: 100bp DNA ladder. C) DNA fragments amplified from single fixed and ISH-probed wild-type embryos after DNA extraction in HotSHOT lysis buffer for 5 min using a primer pair to amplify a 300bp fragment of the *runx1* locus with JumpStart™ REDTaq® ReadyMix™ PCR. First lane from the left: 100bp DNA ladder. The lanes after the ladder represent embryos that are 14h, 1, 2, 3 and 4 days old and have been stored in glycerol at room temperature for 2 years.

Figure 2. *Runx1* mutants have significantly reduced levels of *dnmt3bb.1* mRNA detected by ISH. A) A representative image of ISH for *dnmt3bb.1* in a 33hpf wild type embryo, showing the expression in the dorsal aorta. B) 2% agarose gel showing representative genotypes of wild type (WT), heterozygous (HET) and mutant (MUT) *runx1* embryos, distinguished by RFLP. Yellow: wild type 214bp + 124bp bands, pink: 338bp mutant band. First lane from the left: 100bp DNA ladder. C) Pixel intensity values of *dnmt3bb.1* mRNA in *runx1* mutant embryos (N=36) are significantly decreased compared to wild types (N=32) and heterozygotes (N=62). The coefficients of variation are 24%, 22% and 21% for wild type, heterozygote and mutant groups, respectively. The bars represent mean ± s.d. ***p<0.001 (Games-Howell post-hoc test).

Figure 3. *Gpr65* morphants have significantly increased levels of *runx1* mRNA detected by ISH. A) A representative image of ISH for *runx1* in a 29hpf wild type embryo, showing the expression in the dorsal aorta. B) Pixel intensity values of *runx1* mRNA in in *gpr65* MO-injected embryos (N=16) are significantly higher than in uninjected control siblings (N=16). The coefficients of variation are 34% and 26% for wild type and morphant groups, respectively. The bars represent mean ± s.d. *p<0.05 (*t* test). The power of the *t* test to detect the difference at 0.05 level was 63%.

Figure 4. *Runx1* levels detected by ISH are not affected in *lmo4a* mutants. A) TALENs were designed to a region (blue) ~20bp downstream of the *lmo4a* translation start site (green). Isolated mutant alleles carry 5bp deletions (Δ25-29) (red gaps) upstream of the conserved LIM domains (orange), resulting in a frameshift after S8. The resulting mutant protein is predicted to lack the LIM domains, including the crucial S39 (purple). B) 2% agarose gel showing representative genotypes of wild type (WT), heterozygous (HET) and mutant (MUT) *lmo4a* embryos, distinguished by RFLP. Yellow: wild type 258bp + 87bp bands, pink: 340bp mutant band. First lane from the left: 100bp DNA ladder. C) A representative image of ISH for *runx1* in a 28hpf embryo, showing the expression in the dorsal aorta. D) Quantification of the *runx1* mRNA signal, detected by ISH, from 28hpf wild type (N=15), heterozygous *lmo4a*$^{+/-}$ (het) (N=34) and *lmo4a*$^{Δ25-29/Δ25-29}$ mutant (N=18) embryos from one clutch shows no significant difference in *runx1* pixel intensity among the different genotypes. The coefficients of variation are 41%, 38% and 37% for wild type, heterozygote and mutant groups, respectively. The bars represent mean ± s.d.

Table 1. The efficiency of the HotSHOT protocol for post-ISH DNA extraction from zebrafish embryos at different stages depending on the lysis incubation time.

Incubation time / Age	≤30 min	≥65 min
22-26hpf	99.3% (N=9)	96.4% (N=1)
28-30hpf	99.3% (N=7)	94.2% (N=8)
33-48hpf	100% (N=2)	100% (N=2)
4-5dpf	96.8% (N=3)	88.7% (N=2)

N: number of independent experiments

Fig. 1

A

B Lysis incubation time

5' 15' 30' 45' 60' 120'

gata2a

runx1

tgfbr2b

tgfb3

C Embryonic stage

14hpf 1dpf 2dpf 3dpf 4dpf

Fig. 2

Fig. 3

A *runx1*

29hpf

B

uninjected   *gpr65* MO

*

runx1 signal [pixel intensity]

Fig. 4

A

DNA

wt
`ATG GTGAACAGTCGTGT GGAAAGCTCCGCGGTT GCGGTAACAGGAGGCAGTGGTGGGGCCGTA CGCTCCTGTGCTGGATGCGGTGGTCGGATATCGGAC`

lmo4a (Δ25-29)
`ATGGTGAACAGTCGTGTGGAAA----- GCGGTTGCGGTAACAGGAGGCAGTGGTGGGGCCGTACGCTCCTGTGCTGGATGCGGTGGTCGGATATCGGAC`

Protein

wt
`M VNSRVESSAVAVTGGSGGAV RSCAGCGGRISDRFLLFS MDRYWHTRCLKCSCCQAQLGEIGSTCFSKGGMILCRNDYIRLFGH SGA CSACGQSIPASEM`

Lmo4a (Δ25-29)
MVNSRVES GCGNRRQWWGRTLLCWMRWSDIGPLSALLYGSVLAHALPQVLLLPGSAGRDRQHLLQQRRDDPLQERLHQVVWSQWSLQCMWTVYSCQ----

wt
`VMRAQGNVYHLKCFTCATCRNRLVPGDRFHYVNGTIFCEHD RPGGALLSSHLTPLQSNTLLPDQKVC`

Lmo4a (Δ25-29)
`-----------------------------------------------------------------------`

Legend:
- Translation start site (green)
- TALENs target site (blue)
- LIM domains (orange)
- Crucial residue (dark red)

B

WT | MUT | HET

C

runx1

28hpf

D

28hpf

wild type | het | *lmo4a*^{Δ25-29/Δ25-29}

runx1 signal [pixel intensity]

SUPPLEMENTARY DETAILED PROTOCOL

DNA EXTRACTION AND IMAGE PROCESSING PROTOCOL

for

An optimized pipeline for parallel image-based quantification of gene expression and genotyping after *in situ* hybridization

Tomasz Dobrzycki, Monika Krecsmarik, Florian Bonkhofer, Roger Patient and Rui Monteiro

REAGENTS AND EQUIPMENT

- Stereomicroscope equipped with a digital camera and imaging software
- Fiji Software
- 0.2 ml PCR tubes
- glycerol
- forceps *or* P20 pipette and tips
- single- or multi-channel P200 pipette and tips
- BRAND® glass microscope slide with cavity for imaging embryos: L × W 76 mm × 26 mm, thickness 1.2-1.5 mm, with 1 concavity (BR475505, Sigma)
- HotSHOT alkaline lysis buffer (25 mM NaOH, 0.2 mM disodium EDTA, pH 12)
- HotSHOT neutralization buffer (Tris HCl 40 mM, pH 5)

PROCEDURES

Imaging

1. Prepare a glass slide with a drop of 100% glycerol.
2. Place the first ISH-probed embryo from 80% glycerol onto the slide and take a picture of the embryo with the appropriate orientation and magnification for the ISH staining.
3. Transfer the embryo either with forceps or a P20 pipette into a fresh 0.2 ml PCR tube with a minimum amount of glycerol.
4. Continue taking pictures of the remaining embryos using the same illumination, magnification and exposure settings.

DNA extraction

5. Add 40 µl of HotSHOT alkaline lysis buffer to the PCR tubes (from step **3**.) and make sure the embryos are in the solution. ** Using up to 75 µl does not affect the efficiency of the protocol.*
6. Incubate the tubes at 95°C for at least 5 minutes. ** Incubation longer than 65 min might reduce the success rate of the amplification of a PCR product.*
7. Cool the tubes to 4°C before adding an equal volume (40 µl) of HotSHOT neutralization buffer. ** Overnight incubation at 4°C after the neutralization improves PCR efficiency*. Spin down the samples before each use.
8. Use 5 µl of DNA lysate for PCR to genotype the embryos.

Image analysis

9. For image processing, open the image in Fiji and invert the image with "Edit▶Invert". Then change the image type to 8-bit ("Image▶Type▶8-bit"). * We have generated a Fiji macro that performs these actions simultaneously on all images in a directory.*

10. Using the polygon selection tool, draw your region of interest (ROI) manually on the image around the region containing the signal.

11. Press "t" to open the ROI manager. Use the ROI manager's Measure command to measure the intensity of the ROI. Copy the "mean" value from the Results window to Excel or equivalent calculation software.

12. Move the same ROI, ensuring the same size and shape as the original region, to a region of the zebrafish not containing any staining. Repeat step **11** to measure the background.

13. To obtain the mean pixel intensity of the ISH signal, subtract in Excel the mean intensity value of the background region from that of the stained region for each embryo.

Statistical analysis

14. Assign each intensity value to a genotype (from step **8.**). * If scoring phenotypes by eye (eg. 'high', 'medium' or 'low'), assign the phenotypes to genotypes and analyse with contingency chi-squared test.*

15. Plot all the values on one Q-Q plot to identify any deviations from normal distribution.
    * Normal distribution can also be verified with the Kolmogorov-Smirnov test or the Shapiro-Wilk test. However, for large sample sizes there is a high risk of false positives in these tests.*

16. If there are strong deviations from normal distribution, transform all the values (using e.g. ln or sqrt functions) to make sure they are normally distributed before proceeding.

17. Analyse the differences between the values (transformed if necessary) assigned to each genotype (wt vs heterozygote vs mutant) with 2-tailed ANOVA with 95% confidence levels, accounting for the equality of variances with a Levene's test and Welch correction. For pairwise comparisons between each pair of genotypes, use Tukey's (equal variances) or Games-Howell (unequal variances) post-hoc test.

18. Plot the untransformed values (from step **14**.) as dot plots for the best representation of the results.

Fig. S1

Figure S1. Phenotypic scoring of *dnmt3bb.1*-probed embryos does not fully correspond to their

***runx1* genotype.** A) The phenotypes of 130 33hpf *dnmt3bb.1*-probed embryos coming from the

incross of *runx1* heterozygotes scored as 'high' (N=34), 'medium' (N=64) and 'low' (N=32) follow a

Mendelian distribution. B) Distribution of the phenotypes according to the genotype. Images show a

representative phenotype scored as 'high', 'medium' or 'low'. Percentages show the occurrence of

the phenotypes for the given genotype.

```
// Recursively converts images to 8-bit/invert starting in a user-specified
directory.
macro "conv"{
dir1 = getDirectory("Choose Source Directory ");
list = getFileList(dir1);
dirD = getDirectory("Choose Destination Directory ");
counter = 1;
setBatchMode(true);
for (i=0; i<list.length; i++)
    {
      path = dir1+list[i];
      if (endsWith(path, "tif")){
            print(path);
            run("TIFF Virtual Stack...", "open=path");
            run("8-bit");
            run("Invert");
            counter_converted=IJ.pad(counter,2);
            print(counter_converted);
            saveAs("Tiff", dirD+"conv"+counter_converted);
      close();
      i = i + 1;
      call("java.lang.System.gc");
      }
   counter = counter+1;
   i = i-1;
  }
}
```