Pleistocene climate change and the formation of regional species pools

Joaquín Calatayud[1,2,3], Miguel Ángel Rodríguez[1], Rafael Molina-Vengas[4], María Leo[5],

Jose Luís Hórreo[6] and Joaquín Hortal[2].

1 Departamento de Ciencias de la Vida, Universidad de Alcalá, Edificio de Ciencias, Ctra.

Madrid-Barcelona km. 33,6, 28871 Alcalá de Henares, Madrid, Spain.

2 Departamento de Biogeografía y Cambio Global, Museo Nacional de Ciencias Naturales

(MNCN-CSIC), C/José Gutiérrez Abascal 2, 28006 Madrid, Spain.

3. Integrated Science Lab, Department of Physics, Umeå University, Naturvetarhuset, byggnad

G, NA plan 3, IceLab Umeå universitet, 901 87 Umeå, Sweden.

4. Institute of Plant Sciences, University of Bern, Altenbergrain 21, Bern 3013, Switzerland

5. Departamento de Biodiversidad y Conservación. Real Jardín Botánico de Madrid (CSIC).

28014 Madrid, España

6. Departamento de Biodiversidad y Biología Evolutiva, Museo Nacional de Ciencias Naturales

(MNCN-CSIC), C/José Gutiérrez Abascal 2, 28006 Madrid, Spain.

STRAPLINE: Pleistocene effects on species pools

Abstract

Despite the description of bioregions dates back from the origin of biogeography, the

processes originating their associated species pools have been seldom studied. Ancient

historical events are thought to play a fundamental role in configuring bioregions, but

the effects of more recent events on these regional biotas are largely unknown. We use a

network approach to identify regional and sub-regional faunas of European *Carabus*

beetles, and analyse the effects of dispersal barriers, niche similarities and phylogenetic

history on their configuration. We identify a transition zone matching the limit of the ice

sheets at Last Glacial Maximum. While southern species pools are mostly separated by

dispersal barriers, in the north species are mainly sorted by their environmental niches.

Strikingly, most phylogenetic structuration of *Carabus* faunas occurred during the

Pleistocene. Our results show how extreme recent historical events –such as Pleistocene

climate cooling, rather than just deep-time evolutionary processes, can profoundly

modify the composition and structure of geographic species pools.

Keywords: Pleistocene glaciations, regional community, species pool, regionalization,
occurrence networks, dispersal, niche tracking, *Carabus* ground beetles, Europe.

Introduction

Naturalists have long been captivated by the geographic distribution of world biotas. Rooted in the seminal ideas by Alexandre von Humbolt, this fascination has promoted a long-term research agenda aiming to delineate biogeographic regions according to their integrating faunas and floras (e.g. Wallace 1876, Holt et al. 2013, Rueda et al. 2013). Besides this, the large-scale eco-evolutionary processes that shape regional biotas are known to influence ecological and evolutionary dynamics at finer scales (Ricklefs 2008, 2015). For instance, regional species pools can modulate local diversity patterns (Ricklefs 2011, Medina et al. 2014, Ricklefs and He 2016), the structure and functioning of ecosystems (Naeslund and Norberg 2006), or co-evolutionary processes (Calatayud et al. 2016a). However, despite their fundamental importance, the processes that have configured regional biotas have been seldom studied (and particularly the historical ones), and most explanations on their origin and dynamics remain largely narrative (Crisp et al. 2011).

Perhaps the earliest speculations about the formation of regional species pools took place during the flourishment of bioregionalizations in the mid-19th century (reviewed by Ebach 2015). During that time, and beyond referring to geophysical factors (climate, soils, and physical barriers), some authors already started to emphasize historical influences as key elements determining the configuration of plant and animal regions. For instance, when Wallace (1876) proposed his ground-breaking zoogeographic regions, he argued that while the distribution of ancient linages such as genera and families would likely reflect major geological and climatic changes spanning the early and mid-Cenozoic, species distributions would be more influenced by recent events such as Pleistocene glaciations (see Rueda et al. 2013). These recent events could have fostered many additions and subtractions of species to regional faunas through dispersal and diversification processes. Indeed, increasing evidence suggests that Pleistocene glacial-interglacial dynamics may have driven population extinctions (e.g. Barnes et al. 2002), allopatric speciation in glacial refugia (e.g. Johnson et al. 2004) and post-glacial recolonization events (e.g. Hewitt 1999; Theissinger et al. 2013). Besides shaping phylogeographic patterns (Avise et al. 1998; Ursenbacher et al. 2006; Sommer and Nadachowski 2006; Provan and Bennett 2008), all these processes are likely underpinning diversity patterns for many taxa, particularly in the Holarctic (e.g. Svenning and Skov 2007; Hortal et al. 2011; Calatayud et al. 2016b). However, whether

the signature of Pleistocene glaciations scales up to the configuration of regional biotas

remains largely unknown.

Historical contingencies should act over the intricate interplay between

ecological (i.e. environmental tolerances and dispersal) and evolutionary (i.e.

diversification and adaptation to new habitats) processes underpinning the composition

and structure of regional species pools. On the one hand, niche-based processes may

determine the composition of regional species pools (Mittelbach and Schemske 2015),

mainly throughout their effects on species distribution ranges (Soberon 2007, Hortal et

al. 2010, 2012). These processes integrate responses to abiotic conditions along

geographical gradients and to local and regional biotic environments (Colwell et al.

2009), which may ultimately lead to the appearance of distinct regional communities in

areas of contrasted environmental conditions (Ricklefs 2015). Although species with

similar environmental tolerances/preferences can coexist in regions of similar climate,

their dispersal may be constrained by geographical barriers, which may lead to

divergent species pools under similar environmental conditions. Finally, evolutionary

processes also constrain all these mechanisms. For instance, environmental-driven

regions may be expected if occupancy of new areas is constrained by niche

conservatism (Hortal et al. 2011), which should also lead to species pools integrated of

evolutionary related species (i.e. niche conservatism generating phylogenetically

clustered species pools, Fig.1a). This, however, can be in turn filtered by

biogeographical processes (Gouveia et al. 2014). As such, diversification of lineages

within regions separated by strong dispersal barriers (e.g. mountain ranges) may also

lead to phylogenetically clustered pools of locally adapted species (i.e. geographically

driven niche conservatism; Fig.1a; Warren et al. 2014, Calatayud et al. 2016a).

Historical contingencies may contribute to the configuration of regional pools by

modifying the balance between these processes. For example, differential diversification

rates may be the predominant driver of regional species pools during climatically stable

periods (Cardillo 2011). Yet, regions with a greater influence of climatic fluctuations

such as Pleistocene glaciations may harbour pools of species mostly shaped by the joint

effects of current climate and post-glacial colonization dynamics (Svenning et al. 2015),

as well as by species' competition during these colonization processes (Horreo et al.

2018), thus eroding the signature of geographically-structured diversification processes.

In this study we aim to disentangle the relative importance of the processes

that may contribute to the formation of regional species pools, using European *Carabus*

(Coleoptera: Carabidae) as a model lineage. *Carabus* is a species-rich ground beetle

genus of great popularity due to the beautiful jewel-like appearance of some species

(Turin et al. 2003). In general, *Carabus* species are flightless nocturnal predators of

snails, earthworms and caterpillars. They hold hydrophilic adaptations and are typically

associated to deciduous forests (Deuve et al. 2012). Previous evidence suggests that the

richness of species from this genus in Europe is determined to a large extent by both

current environmental conditions (i.e. climate and habitat) and glacial-interglacial

dynamics (Calatayud et al. 2016b). This makes European *Carabus* an ideal case study to

evaluate the joint effects of evolutionary, ecological and historical contingency

processes as drivers of regional species pools.

Specifically, we use data on the distribution and evolutionary relationships of

Carabus species, along with network and phylogenetic analyses, to evaluate six

hypotheses: First, given the presumed low dispersal capacity of the species from this

genus (Turin et al. 2003), we hypothesize that (H1) European *Carabus* species pools are

mainly shaped by the main orographic barriers of the continent, but also, that (H2)

glacial-interglacial dynamics have led to strong differentiation between northern and

southern regional species pools. If this differentiation is true, northern European

Carabus faunas will be comprised of species that colonized newly vacant habitats after

the withdrawal of the ice sheet, and hence (H3) their regional distribution will be mostly

determined by current climate. In contrast, (H4) southern faunas will be mainly shaped

by the joint influence of diversification events and dispersal limitations, due to the

combined effect of higher climatic stability (e.g. climatic refugia) and a more complex

orography (Alps, Pyrenees, Carpathians). Therefore, (H5) species forming northern

regional pools will exhibit comparatively lower levels of regional endemicity, whereas

those forming southern regional pools will show comparatively higher levels of regional

affinity. Finally, according to Wallace (1876), the advance and retreat of the ice sheets

during the Pleistocene should have determined the spatial distribution of lineages,

eroding the effects of the former configuration of the distribution of the main *Carabus*

lineages. Therefore, (H6) we expect a temporal signal coincident with the Pleistocene in

the phylogenetic structure of *Carabus* faunas, and no effect of deep-time events on the

current geographical distribution of these lineages.

Material and methods

Rationale and structure of the analyses

Exploring the determinants of regional faunas requires jointly analysing ecological, evolutionary and historical factors. We did so through three consecutive steps (Fig.1b). First, we identified distinct regional species pools within Europe by using a network community detection algorithm. From this analysis we derived a species pairwise similarity matrix of occurrence into different modules, each one representing different regions. Second, we assessed the relative importance of environmental, spatial and evolutionary determinants of such similarity. To do so, we constructed four pairwise matrices to describe ecological, topographical and evolutionary relationships among species; namely, *i*) a matrix of climatic niche similarity, *ii*) a matrix of habitat similarity, *iii*) a matrix of spatial connectivity among distributional ranges, and *iv*) a phylogenetic distance matrix. Then, we used generalized partial matrix regressions to model the similarity in species occurrences as a function of these four matrices (Fig.1b). We used this workflow to explore the factors involved in the configuration of *Carabus* faunas both at regional (i.e. through analysing *Carabus* species co-occurrence across regions) and sub-regional scale (i.e. focusing on co-occurrence patterns within sub-regions). Finally, we also applied ancestral range estimation analysis in order to identify the time period from which ancestral areas are estimated with less uncertainty. By doing so, we aimed to detect important historical periods contributing to the regional organization of *Carabus* lineages.

The interpretation of the joint and independent effects of explanatory matrices can shed light on the different processes configuring regional faunas (see Fig.1a). Thus, if niche similarities (i.e. represented by the climatic and habitat similarity matrices) and phylogenetic distances altogether explained the regional co-occurrence of species, then this could be interpreted as indicative of constrained niche evolution (or a tendency to resemble ancestral niches) in shaping regional faunas (Fig.1a.i). However, if spatial connectivity also accounted for part of this co-occurrence, this would indicate that this niche conservatism pattern can be caused by geographical constrains (Fig.1a.ii). Further, the effects of niche similarities and spatial connectivity alone (i.e. without phylogenetic signal) can be most likely the consequence of a convergence of climatic niches due to geographic isolation (Fig.1a.iii), whereas the effects of connectivity and phylogeny would be indicative of a primacy of intra-regional speciation driven by geographical barriers. Niche similarities alone would point to an unconstrained niche evolution shaping regional faunas, while phylogeny alone would indicate a primacy of geographically unconstrained intra-regional speciation events. Finally, either a *cul-de-*

sac effect (i.e. the accumulation of species in past climatic refugia) or a primacy of

vicariant speciation events could lead to the existence of independent effects of

connectivity and regional co-occurrence (Fig.1a.iv).

Identification of regional species pools

We took advantage of community detection analysis —borrowed from network

theory— to identify *Carabus* regional species pools in Europe. We first generated a

bipartite network where species and grid cells constitute two disjoint sets of nodes that

are connected according to the presence of species in grid cells (e.g. Calatayud et al.

2016a). The species presence data comes from expert-based range maps of all *Carabus*

species inhabiting Europe (n = 131, Turin et al. 2003) overlaid into a 100-km equal-area

grid based on the LAEA pan-European grid system (available at

http://dataservice.eea.europa.euhttps://inspire.ec.europa.eu/, see Calatayud et al. 2016b

for details). Then, we conducted a modularity analysis using the index proposed by

Barber (2007) and the Louvain algorithm (Blondel et al. 2008) as implemented in the

Matlab function "Gen Louvain", (available at http://netwiki.amath.unc.edu; Mucha et al.

2010). This analysis is intended to find groups of nodes (i.e. species and grid cells) that

are more densely connected. Hence, in our case, the analysis identified groups of grid

cells, each group sharing *Carabus* species mainly distributed within its cells (i.e. regions

and their associated faunas). The Louvain algorithm was run 100 times, and the network

partition showing highest modularity value was retained. This optimal solution was

used to conduct all subsequent analyses, although all the solutions were qualitatively

similar. We evaluated the statistical significance of the modules by comparing their

associated modularity value to a null distribution of values (n = 100) where the original

presence-absence matrix was randomized using the independent swap algorithm (a

fixed-fixed null model implemented in the R package "picante", Kembel et al. 2010).

Finally, to detect potential sub-modules (i.e. sub-regions) nested within modules (i.e.

sub-regional species pools within regional species pools), we derived a new bipartite

network from each of the previously identified modules, and applied the procedure

described above in each case.

It is important to note that despite species and grid cells were assigned to just

one module, they could also occur in other modules with different degrees of

specificity. For example, despite most species in a grid cell will belong to the same

module the cell does, this cell could also hold species that are primarily associated to

other modules. Similarly, although a species will mostly be present in cells assigned to

its module, it may also occur in cells from other modules. Thus, we calculated the

degree of module specificity for each node (i.e. species and grid cells) as its number of

links with nodes of its module divided by its total number links (see Guimera and

Amaral's 2005 inter-modular participation index for a similar metric). Thus, higher

module specificity would correspond to species mainly distributed within its module

(highly endemic species), as well as to cells pertaining to well-defined regions; whereas

lower module specificity would indicate widespread species and cells located in

transition zones.

Assessing the determinants of regional species pools

To disentangle the determinants of the current configuration of *Carabus* faunas in

Europe, we first generated a species-per-module matrix, where each entry of the matrix

represents the percentage of the distributional range of a certain species that lies in a

given module. Then, we derived a co-occurrence pairwise similarity matrix from the

former matrix using the Schoener's index (Schoener 1970). This metric quantifies the

overlap between species pairs throughout the modules (see Krasnov et al. 2012 for a

previous application) and it ranges from 0 (no overlap) to 1 (identical distribution across

modules). Note that this similarity matrix reflects the co-occurrence similarities at

regional scale, thus ignoring distributional patterns at lower spatial scales (i.e. two

species may have identical regional distribution but appear as segregated at the local

scale). The resultant co-occurrence pairwise similarity matrix was used as dependent

variable. We generated four different pairwise dis/similarity matrices to be used as

explanatory variables. Two of them were used to account for environmental factors: (i)

a climatic-niche similarity matrix and (ii) a habitat similarity matrix. The remaining two

considered geographical and evolutionary factors: (iii) a spatial-connectivity matrix and

(iv) a phylogenetic distance matrix.

i) Climatic-niche similarity matrix. We characterized the climatic niche of each *Carabus*

species in the dataset following a similar approach as proposed by Broennimann et al.

(2012). We selected six bioclimatic variables to account for the main water and energy

aspects of climate –namely mean annual temperature, temperature of the warmest

quarter, temperature of the driest quarter, total annual precipitation, total precipitation of

the warmest quarter and total precipitation of the driest quarter– and altitudinal range to

account for the effects of mesoclimatic gradients within each grid cell. These variables

may be among the main determinants of the distribution of *Carabus* species diversity

within Europe (see Calatayud et al. 2016b). Bioclimatic variables were extracted from

Worldclim (v1.4 Hijmans et al. 2005; available at http://www.worldclim.org/), whereas

altitudinal data were derived from the 30-arcsecond digital elevation model GTOPO30

(available at https://lta.cr.usgs.gov/GTOPO30). We conducted a principal component

analysis on these variables to obtain a bidimensional climatic space defined by the two

main axes, that explained 81.4% of the variance (Fig. S1). Finally, we divided this

climatic space into a 100X100 grid and calculated species overlap in the gridded space

using Schoener's index (see above).

ii) Habitat similarity matrix. The distribution of *Carabus* species may also be shaped by

forest preferences (Turin et al. 2003). Accordingly, we used ten vegetation categories

derived from MODIS Land Cover at 5-minute resolution (Evergreen broadleaf forest,

deciduous needle-leaf forest, deciduous broadleaf forest, mixed forest, closed shrub

lands, open shrub lands, woody savannas, savannas and grasslands; Channan et al.

2014, available at http://glcf.umd.edu/data/lc/). For each species we computed the

proportion of each category overlaying its range. With this, we computed pairwise

similarities in the preference for different vegetation types using Schoener's index (see

above).

iii) Spatial connectivity matrix. To evaluate the potential influence of geophysical

barriers to dispersal on the current distribution of *Carabus* species, we first created a

dispersal-cost surface by dividing the study area in 1 Km^2 grid cells and weighting each

cell according to its topography (in this case, slope) and the presence of water bodies.

Slope values ranged from 0 to 100 at each pixel, being 0 the lowest dispersal cost and

100 the highest one, and were determined from GTOPO30 altitudinal data using the

GRASS tool r.slope (GRASS Development Team 2017). Grid cells including water

bodies as rasterized layers in the Nature Earth database (available at

http://www.naturalearthdata.com/) were further weighted by assigning arbitrary values

of friction to the dispersal of *Carabus* species, namely 30% for cells containing rivers

and lakes and 99% for cells that lay on sea water masses (note that *Carabus* species

show hydrophilic adaptations). Then, the connectivity between all pairs of cells was

calculated as the least-cost path over the dispersal-cost surface that connects both cells,

using the "gdistance" R package (van Etten 2015). Finally, the spatial connectivity

between each pair of species' distributional ranges in the dataset was estimated as the

average distance among all grid cells within the range of each species. Average

distances were preferred over absolute least-cost distances to avoid disproportionate

differences in spatial connectivity between overlapping and non-overlapping

distributional ranges.

iv) Phylogenetic distance matrix. To unravel the evolutionary history of the *Carabus*

lineage and assess the potential importance of evolutionary processes in determining the

formation of *Carabus* species pool, we reconstructed a species-level time-calibrated

molecular phylogeny including the 89 species for which we found available DNA

information on ten markers (eight mitochondrial regions: 12S rDNA,16S rDNA, 28S

rDNA, ND4, ND5, COI, Cytb and PEPCK; plus two nuclear ones: anonymous locus

and wingless; see Tables S1 and S2). We aligned each marker independently using

different algorithms: MAFFT (Katoh and Standley 2013), Clustal X (Thompson et al.

1994; Larkin et al. 2007), MUSCLE (Edgar 2004) and Kalign (Lassmann and

Sonnhammer 2005), and selected the most reliable alignment for each marker using the

multiple overlap score (MOS) provided by MUMSA (Lassmann and Sonnhammer

2006). We removed ambiguous or poorly aligned positions from the alignments with

trimAl (Capella-Gutiérrez et al. 2009). The final dataset was concatenated in 5603

basepairs following a total-evidence approach (Kluge 1998), and was used to conduct

Bayesian phylogenetic inference with BEAST v.2.4.6 software (Bouckaert et al. 2014).

We used a GTR model for sequence evolution, a Random Local Clock, a birth-death

model prior, and 100 million of MCMC chain length searching for convergence. There

is ongoing debate on the divergence time of the genus *Carabus* (Andujar et al. 2012,

Deuve et al. 2012), hence molecular dating was conducted under two different

scenarios. Firstly, according to Deuve et al.'s (2012) molecular dating, the crown age of

Carabus was set at 17.3 Mya. Secondly, and according to Andujar et al. (2012), the

origin of the group was set at 25.16 Mya. We used the software Tracer v 1.6 (available

at http://tree.bio.ed.ac.uk/software/tracer/) to check for MCMC chains convergence, and

FigTree v.1.4.2 (available at http://tree.bio.ed.ac.uk/software/figtree/) for viewing and

editing the phylogenetic trees (see Figures S2 and S3). To account for topological and

time-calibration uncertainties, we used 100 phylogenies sampled from the posterior

distribution for each molecular dating scenario. In addition, we used taxonomic

information and phylogenetic uncertainty methods (Rangel et al. 2015) to place species
lacking molecular information into the phylogeny (see Appendix S1). Thus, we derived
100 different phylogenetic hypotheses from each Bayesian posterior phylogeny by
randomly inserting missing species within their most derived consensus clade based on
taxonomic knowledge. In total, we generated 20,000 phylogenetic hypotheses (100
phylogenies per two molecular dating scenarios per 100 phylogenies accounting for
uncertainties associated with lack of molecular data) that were used in subsequent
analyses. Pairwise phylogenetic distances were calculated for each calibrated phylogeny
using the function *cophenetic* implemented in the APE R package (Paradis et al. 2004).
314

We used generalized multiple regression on distance matrices and deviance
partitioning to disentangle the relative importance of climatic niche, habitat preferences,
dispersal barriers and evolutionary history in determining *Carabus* species pools in
Europe. First, we conducted single regressions between the co-occurrence pairwise
similarity matrix and each of the four explanatory matrices described above to seek for
significant associations between the variables. We set a binomial family for error
distribution and "logit" as the link function (see Ferrier et al. 2007 and Calatayud et al.
2016a for a similar approach). To assess for significance, we randomized the observed
species per module matrix using the independent swap algorithm (see above) to derive
999 null occurrence similarity matrices. Then, we used simple regressions to relate each
null similarity matrix with each one of the explanatory matrices. The relationship
between an explanatory matrix and the observed species per module matrix was
considered to be significant when it explained a higher proportion of the deviance than
99% of the regressions performed on the null matrices. In the case of phylogenetic
pairwise distances we repeated this procedure for each phylogenetic hypothesis to
consider phylogenetic uncertainties, applying the same criterion for significance.
Finally, we retained those variables that showed significant relationships, and conducted
variance partitioning among explanatory matrices (Legendre and Legendre 2012) to
explore patterns of covariation among niche similarities (i.e. climatic and habitat
similarity matrices), dispersal barriers and phylogenetic history. We conducted the
analyses for the co-occurrence into modules (i.e. regions) and sub-modules (i.e. sub-
regions) both at a European and regional (i.e. co-occurrence into sub-modules of each
module) scales.
338

Ancestral range estimation

To assess whether deep historical signals were eroded by Pleistocene glaciations we used probabilistic models of geographic range evolution. These models estimate ancestral range state probabilities assuming different processes involved in range evolution. We used the Dispersal–Extinction–Cladogenesis model of range evolution (DEC; Ree and Smith 2008) implemented in the R package BioGeoBears (Matzke 2014) since this model has been shown to perform well even under complex scenarios (Beeravolu and Condamine 2018). Species ranges were coded as present/absent in each module detected in the former network clustering. We used this analysis for 1,000 randomly selected phylogenies of each dataset, since preliminary results showed little variations among phylogenies. The estimation of ancestral ranges usually tends to be more ambiguous in deeper nodes of the phylogeny, as the lability of geographical ranges would tend to blur deep-time signals (Losos and Glor 2003). Similarly, if the Pleistocene glacial periods had important effects on species distributions it could be expected that ancestral range estimations will increase in accuracy around the Pleistocene. That is, pre-Pleistocene signals on the evolution of species distribution ranges will be blurred. To explore this, we evaluated the existence of changes in the relationship between node age and the marginal probability of the single most-probable ancestral state at each internal node. Then, general additive mixed models (GAMMs) were fitted to the node marginal probability as a function of node age, including the phylogenetic hypothesis as a random factor. We also used generalized linear mixed models (GLMMs) combined with piecewise regression to detect potential major breakpoints (i.e. temporal shifts) in the relationship between marginal probability and node age for each of the two dating scenarios. Given the large amount of observations (130 nodes x 1,000 phylogenies) we firstly estimated the breakpoint independently for each phylogenetic hypothesis. To do so, we included the breakpoint as a new parameter in a generalized linear model of marginal probabilities as a function of node age, minimizing the deviance of the fitted model using the function "optimize" in the R package lme4 (Bates et al. 2014). Finally, to assess significance we used the averaged breakpoint value in a GLMM with the same fixed formulation but including the phylogenetic hypothesis as a random factor. Because node marginal probabilities ranged between 0 and 1, we used a binomial family and a loglink function to fit all models.

General assumptions of probabilistic ancestral range estimation models may

compromise subsequent interpretations (Ree and Sanmartin 2018). Moreover, our

dataset does not fulfil the assumption of ancestral range estimation models that

phylogenies should include all extant lineages, since some Carabus linages have

representatives from outside Europe (mainly Asiatic species) that were not included in

the analyses. Hence, we conducted additional analyses to provide further evidence on

the temporal signal in the phylogenetic structuration of *Carabus* faunas coinciding with

Pleistocene glaciations. To do so, we first generated a binomial variable based on the

module affiliation of the species descending from each internal phylogenetic node,

coding the phylogenetic nodes whose all descendant species were classified into the

same region as 1, and 0 otherwise. Then, we conducted the analytical approach

explained above, in this case fitting this binomial variable as a function of node age.

This analysis would further elucidate if there is a breakpoint in the relationship between

node age and the probability that all descendant species belong to the same region (see

Appendix S2 for details and another complementary approach). Finally, we also

explored whether there are differences between regions in the Pleistocene signal on the

phylogenetic structure of their faunas. For each region we calculated the probability that

a phylogenetic node has all its descendant species within the region, independently for

nodes ocurring either before and after the beginning of the Pleistocene (2.59 Mya;

herein pre-Pleistocene and post-Pleistocene nodes). This probability was calculated as

the number of pre- or post-Pleistocene nodes having all descendants associated to

region i divided by the total number of nodes in the phylogeny.

All analyses were carried out in R (R core team 2015), using the function *bam*

of the package mcvg for GAMM (Wood at al. 2015) and the package Lme4 for GLMM

analyses (Bates et al. 2014).

Results

Identification of regional faunas

The *Carabus* occurrence network was significantly modular (M=0.385, p=0.01),

dividing Europe in seven modules that group zoogeographically distinct regions with

their associated faunas (i.e., different regional species pools; Figs. 2a and S4).

Furthermore, all modules but module 2 showed significant sub-modular structure,

presenting a decrease in modularity with latitude (mean M=0.316, ranging from 0.154

to 0.468; all p-values < 0.05, see Table S4). Module 1 holds 21 species mainly living in

South-western Palearctic (Iberian Peninsula, North of Africa, Balearic Islands, Corsica,

Sardinia and the western half of Sicilia). This module was subdivided into four

submodules. Module 2 included only two species, both endemic of Crete. Module 3

identified an East Mediterranean region including the Italic Peninsula, part of Greece

and Turkey. This module holds 18 species and was subdivided into five submodules.

Module 4 depicted a Central European region embracing the Alps and the Carpathian

Mountains, as well as Central European plains. This module showed the highest species

richness, including 49 *Carabus* species, and was split into four submodules. Module 5

and module 6 comprised northern regions and showed the lowest species richness

values, holding 10 species each. The former comprised Iceland and the British Isles and

extended eastward up to the vicinity of the Ural Mountains. The latter included this

mountain range and expanded to the easternmost zone of the study area. Both modules

were divided into 3 submodules. Finally, module 7 included 21 species and embraced a

south-eastern central European region expanding from the Carpathian Mountains to the

south Ural Mountains. This module was split into three submodules.

Regarding transition zones between regions, and in agreement with our first

hypothesis, we found that they were clearly associated with geographical barriers such

as the Pyrenees, the Alps, the Carpathian and the Ural Mountains, as well as the Turkish

Straits System (that connects the Black Sea to the Aegean separating the Anatolian and

Greek peninsulas; Fig. 2b). Interestingly, we also identified a west-to-east transitional

belt between southern and northern regions that closely followed the southern limits of

the ice sheet at the Last Glacial Maximum (LGM). This transitional zone further

suggested a link between the configuration of *Carabus* regional faunas and Pleistocene

glacial conditions, supporting our hypothesis 2.

Correlates of regional co-occurrence

Matrix regressions showed that deviance of species co-occurrences across regions,

across sub-regions and within each region was significantly explained ($p < 0.01$),

primarily by environmental niche similarity, and secondarily by spatial connectivity,

except for northern regions (i.e. modules 5 and 6; Fig. 3 and Table S5). In contrast,

relationships with evolutionary relatedness were non significant in all instances and

regardless of the phylogenetic hypothesis used ($p > 0.01$ in all cases, see Table S5).

Comparing both types of subdivisions, environmental niche similarity explained more

deviance across sub-regions than across regions, whereas spatial connectivity did the

reverse (see Fig. 3). Comparing explained deviances between regions, the primacy of

environmental niche similarity (mostly climate, see Table S5) in the northern ones (5

and 6) is consistent with the notion that northern regional pools are geographically

sorted by current climate (our hypothesis 3), whereas the importance shown by spatial

connectivity in the remaining regions is consistent with the more complex orography of

central and southern Europe (consistent with hypothesis 4).

Ancestral range estimation

Both phylogenetic datasets (i.e. alternative calibration scenarios) yielded similar

qualitative and quantitative results (see Appendix S2). Thus, we only present here

ancestral range estimations based on Deuve's et al. (2012) calibration. GAMM results

showed that node marginal probability of the most probable state increased towards

younger nodes (P<0.01, explained deviance =7.49%, Fig. 4a). However, this increase

showed a steep increment coinciding with the Pleistocene. Indeed, piecewise regression

revealed that the relationship between marginal state probability and node age changed

at 1.51 Mya (median value; with 45^{th} and 55^{th} percentiles at 1.24 and 1.89 Mya.,

respectively; Fig. 4a and S5), suggesting that most of the phylogenetic structuration of

Carabus faunas began around the Pleistocene. Similarly, the probability for a

phylogenetic node to have all descendant associated to same region increased toward

younger nodes with a break point roughly associated with the Plio-Pleistocene transition

(3.70 Mya.; 45^{th} and 55^{th} percentiles at 2.47 and 4.48 Mya.; see Fig. S6). In both cases

the breakpoints associated to the Pleistocene were significant (P < 0.01).

In agreement with these results, we found that the probability of finding

phylogenetic nodes having all descendant belonging to the same region was higher for

post-Pleistocene nodes (median at 0.21; 25^{th} and 75^{th} percentiles at 0.20 and 0.23,

respectively; Fig. 4a) than for pre-Pleistocene ones (median at 0.10; 25^{th} and 75^{th}

percentiles at 0.08 and 0.11). The phylogenies based on Andujar's calibration yield

similar probabilities for both types of nodes (median at 0.15; 25^{th} and 75^{th} percentile at

0.14 and 0.17; and median at 0.16; 25^{th} and 75^{th} percentile at 0.14 and 0.18; respectively

for pre- and post-Pleistocene nodes, see Fig. S7). These low probabilities are,

nonetheless, congruent with the lack of phylogenetic signal in module co-occurrence

previously found. Interestingly, and regardless of calibration scheme, the probabilities

found for both type of nodes (i.e. pre- and post-Pleistocene) were higher in southern and

central regions than in northern ones (Fig. 4b). This suggests that regions not covered

by ice during the LGM can still reflect some old historical legacies (as shown by the

higher pre-Pleistocene node probability) while accumulating some related lineages that

diversify during and after the Pleistocene (as indicated by the higher post-Pleistocene

node probability).

Discussion

More than 140 years ago, Wallace (1876) foresaw that the influence of Pleistocene

glaciations on the distribution of diversity had been strong enough to erode the imprint

of previous events. Our results support Wallace's thoughts, showing a remarkable

coincidence between the distribution of the ice sheets at the Last Glacial Maximum and

the current configuration and evolutionary structure of European *Carabus* Faunas.

The first line of evidence supporting this idea comes from the close spatial

relationship between the southern limits of the ice sheet at LGM and the transition zone

separating the southern and northern regions. This border also coincides with the line –

identified by Calatayud et al. (2016b)– where the relationship between *Carabus* species

richness and current climate changes (Fig. 2). Thus, it seems that the climate changes

underwent during the Pleistocene not only shaped phylogeographic (Avise et al. 1998;

Hewitt 1999; Barnes et al. 2002; Johnson et al. 2004; Theissinger et al. 2013; Horreo et

al. 2016) and species richness patterns (e.g. Svenning and Skov 2007, Araújo et al.

2008, Hortal et al. 2011, Calatayud et al. 2016b), but that Ice ages have also left a strong

imprint on the geographical structure of species composition at a regional scale.

Accordingly, the species from the northernmost region (module 5) show the lowest

level of endemism (Fig. S8), as expected for regional faunas composed of species that

have recently colonized the north of Europe from southern glacial refugia (Araújo et al.

2008, Calatayud et al. 2016b, our hypothesis 5). In fact, although these species show

large distribution ranges in different parts of southern Europe, their ranges only overlap

near the northern Carpathian Mountains (Fig. S9). This area was a glacial refugia for a

large and taxonomically diverse array of northern European species (Ursenbacher et al.

2006; Sommer and Nadachowski, 2006, Provan and Bennett 2008), including *Carabus*

(Homburg et al. 2013). Additionally, the decrease in modularity values with latitude

also points to a lesser geographical structure of northern assemblages, which can be

interpreted as the result of a post-glacial colonization, together with less geographic

complexity in some areas.

Besides the Pleistocene effects in the definition and geographical structure of

regional species pools, we also found evidence of the imprint of this geological period

on the processes configuring the distribution of *Carabus* faunas. The general strong

relationship between regional patterns of co-occurrence and both niche similarities and

spatial connectivity shows that co-occurring species tend to have similar realized

environmental niches and that also tend to be geographically constrained by the same

dispersal barriers. This latter result was expected given the –presumed– low dispersal

capacity of *Carabus* species (see Turin et al. 2003), which is likely to be behind the

spatial coincidence of module transition zones and geographical barriers. Perhaps more

unexpected is the weak effect of phylogenetic distances despite the strong relationship

between regional co-occurrence and niche similarities. This implies that geographical

barriers rather than climatic-niche conservatism have restricted species distributions

even within regions of similar climate. These results also point to that *Carabus* niche

evolution is, to some extent, evolutionarily unconstrained, which is congruent with the

general high adaptation capacity of insects (e.g. Overgaard and Sørensen 2008).

Whatever the origin of the relationship between species occurrence and

environmental conditions, what is certainly true is that its strength changes between

regions. These changes follow a latitudinal gradient in the importance of environmental

niche similarities (Fig. 3). The occurrence into sub-regions is more strongly related to

the similarity in the realized niche in the north than in the south. This might be a direct

consequence of the effects of post-glacial colonization, where formerly glaciated areas

show a clear sorting of species due to its environmental preferences. On the contrary, in

southern regions, species are expected to had have more time to diversify and sort

geographically by other factors besides climate (Hortal et al. 2011). Our findings

corroborated this idea since we found strong effects of dispersal barriers in these areas.

Moreover, although we did not find a significant phylogenetic signal in the subregional

co-ocurrence over these regions, our analyses revealed that they hold a small but still

larger number of related species compared to northern ones, supporting that more stable

regions are more prone to accumulate related species.

Despite of these related species of southern regions, we found a generalized

lack of phylogenetic structuration of *Carabus* faunas. This can be the outcome of –

relatively– recent speciation events due to vicariance and/or a "cul-de-sac effect"

(O'Regan 2008). The former would imply the formation of dispersal barriers promoting

the geographical split of many lineages and subsequent allopatric speciation (Weeks et

al. 2016). Yet, the geophysical accidents that can be associated with the limits of the

Carabus regions we found here largely predate the origin of the genus (see Beccaluva et

al. 1998, Deuve et al. 2012). On the other hand, a generalized dispersion into climatic

refugia, together with a subsequent stagnancy within them (i.e. a "cul-de-sac" effect)

may also produce the observed mixing of unrelated lineages into regions. Although it is

difficult to distinguish between both processes, the latter seems more plausible, with

southern regions accumulating unrelated species while acting as glacial refugia, and

northern ones being recolonized by unrelated species with similar environmental niches

and/or simply higher dispersal capacity (Svenning and Skov 2007).

Supporting the Pleistocene signature, our results showed a temporal coincidence

between this geological period and the phylogenetic structuration of *Carabus* faunas.

This result was consistent regardless of the different approaches used and across the

different time calibration scenarios. This robust temporal coincidence supports that the

current regional organization of *Carabus* species and lineages is rooted at the

Pleistocene, which also explains the general lack of phylogenetic structure of regional

Carabus faunas. Our results partially contrast with ancestral range estimations for

clades inhabiting areas that were never glaciated, where more ancient signals were

found in the spatial sorting of lineages (Condamine et al. 2015, Economo et al. 2015,

Tänzler et al. 2016, Toussaint and Balke 2016). These previous findings are,

nonetheless, congruent with the higher probability of holding related *Carabus* species of

southern and more stable European regions. Interestingly, a visual inspection of 100

randomly chosen phylogenies (Appendix S5) reveals that several related lineages that

diversified before the Pleistocene and inhabit central regions (subgenera *Orinocarabus*

and *Platycarabus*; Fig. 4c and Appendix S5) mostly comprise montane and alpine

species (Turing et al. 2003). This suggests that older historical signals are mostly due to

presumably cold-adapted species that should be less affected by glacial conditions.

Nevertheless, this speculation should be taken with caution since the central-eastern

region also hosts related species (some *Morphocarabus* species, see Fig. 4) that are not

particularly adapted to cold environments (Turing et al. 2003). Further studies are

required to determine which characteristics allowed some *Carabus* to persist under

glacial conditions. In sum, these findings suggest that the repeated advances and retreats

of ice sheets and glacial conditions that characterize the European Pleistocene produced

repeated cycles of retreat to southern regions and advance towards the north of *Carabus*

species, a hustle-and-bustle process that ultimately led to the observed mixing of

unrelated lineages, with few related species inhabiting in less affected regions.

576    To summarize, our results provide solid arguments in favour of the importance

of Pleistocene glaciations along with geographical barriers and niche-based processes in

structuring the regional faunas of European *Carabus*. On the one hand, this group's

faunas are primarily delimited by the location of the southern limit of the ice sheet at

LGM, which separates two large regions that differ not only in species composition, but

also in the processes underlying the spatial organization of these species. On the other

hand, the phylogenetic structure of these faunas coincides with the beginning of the

Pleistocene. This implies that the geographical distribution of species and lineages is

profoundly shaped by past climates. Moreover, our results also suggest that ecological

(Naeslund and Norberg 2006, Madrigal et al. 2016) and evolutionary mechanisms

(Wüest et al. 2015, Calatayud et al. 2016a) that rely on procceses ocurring at regional

scales can be profoundly affected by the history of Earth's climates. Hence, the study of

these historical events may be essential to unravel both large and local scale diversity

patterns.

Acknowledgements

We are very grateful to Achille Casale for providing data on *Carabus* habitat

preferences and comments on an early version of the phylogeny, and the Scientific

Computation Centre of Andalusia (CICA) for the computing services they provided. We

acknowledge insightful discussions with Hortal lab members. JC was supported by a

FPU-fellowship of the Spanish Ministry of Education (FPU12/00575). This work is

partly supported by the Spanish Ministry of Economy, Industry and Competitiveness

(MINECO) project SCARPO (CGL2011-29317) to JC and JH. MAR and RMV were

supported through the grants CGL2017-86926-P and CGL2013-48768-P. ML and JLH

were supported by MINECO FPI and Juan de La Cierva grants, respectively.

Author contributions

603 JC and JH conceived research. JC designed the study with contributions of all authors.

604 JC, RMV, ML and JLH analysed the data. All authors discussed results. JC wrote the

605 paper with contributions of all authors.

References

1. Andújar, C., Serrano, J. and Gómez-Zurita, J. 2012. Winding up the molecular clock in the genus Carabus (Coleoptera: Carabidae): assessment of methodological decisions on rate and node age estimation. BMC evolutionary biology, **12**: 40.

2. Araújo, M. B., D. Nogués-Bravo, J. A. F. Diniz-Filho, A. M. Haywood, P. J. Valdes, and C. Rahbek. 2008. Quaternary climate changes explain diversity among reptiles and amphibians. Ecography **31**:8–15.

3. Avise, J. C., Walker, D., and Johns, G. C. 1998. Speciation durations and Pleistocene effects on vertebrate phylogeography. Proceedings of the Royal Society of London B: Biological Sciences **265**: 1707-1712.

4. Barber, M. J. 2007. Modularity and community detection in bipartite networks. Physical Review E **76**:066102.

5. Barnes, I., P. Matheus, B. Shapiro, D. Jensen, and A. Cooper. 2002. Dynamics of Pleistocene population extinctions in Beringian brown bears. Science **295**:2267–2270.

6. Bates, D., M. Maechler, B. Bolker, S. Walker, et al. 2014. lme4: Linear mixed-effects models using Eigen and S4. R package version **1**.

7. Beccaluva, L., M. Shallo, M. Coltorti, I. Premti, and F. Siena. 1998. Encyclopedia of European and Asian Regional Geology. Springer.

8. Beeravolu C.R., and F.L. Condamine. 2018. An Extended Maximum Likelihood Inference of Geographic Range Evolution by Dispersal, Local Extinction and Cladogenesis. bioRxiv. https://doi.org/10.1101/038695

9. Blondel, V. D., J.-L. Guillaume, R. Lambiotte, and E. Lefebvre. 2008. Fast unfolding of communities in large networks. Journal of Statistical Mechanics **2008**:P10008.

10. Broennimann, O., M. C. Fitzpatrick, P. B. Pearman, B. Petitpierre, L. Pellissier, N. G. Yoccoz, W. Thuiller, M.-J. Fortin, C. Randin, N. E. Zimmermann, et al. 2012. Measuring ecological niche overlap from occurrence and spatial environmental data. Global Ecology and Biogeography **21**:481–497.

11. Calatayud, J., J. L. Hórreo, J. Madrigal-González, A. Migeon, M. Á. Rodríguez, S. Magalhães, and J. Hortal. 2016*a*. Geography and major host evolutionary transitions shape the resource use of plant parasites. Proceedings of the National Academy of Sciences USA **113**:9840–9845.

12. Calatayud, J., J. Hortal, N. G. Medina, H. Turin, R. Bernard, A. Casale, V. M. Ortuño, L. Penev, and M. Á. Rodríguez. 2016*b*. Glaciations, deciduous forests, water availability and current geographical patterns in the diversity of European *Carabus* species. Journal of Biogeography **43**:2343–2353.

13. Capella-Gutiérrez, S., J.M. Silla-Martínez, and T. Gabaldón. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics **25**:1972–1973.

14. Cardillo, M. 2011. Phylogenetic structure of mammal assemblages at large geographical scales: linking phylogenetic community ecology with macroecology. Philosophical Transactions of the Royal Society of London B **366**:2545–2553.

15. Channan, S., K. Collins, and W. Emanuel. 2014. Global mosaics of the standard MODIS land cover type data. University of Maryland and the Pacific Northwest National Laboratory, College Park, Maryland, USA.

16. Colwell, R. K., and T. F. Rangel. 2009. Hutchinson's duality: the once and future
niche. Proceedings of the National Academy of Sciences USA **106**:19651–19658.

17. Condamine, F. L., E. F. Toussaint, A.-L. Clamens, G. Genson, F. A. Sperling, and G. J.
Kergoat. 2015. Deciphering the evolution of birdwing butterflies 150 years after
Alfred Russel Wallace. Scientific Reports **5**:11860.

18. Crisp, M. D., S. A. Trewick, and L. G. Cook. 2011. Hypothesis testing in
biogeography. Trends in Ecology and Evolution **26**:66–72.

19. Deuve, T., A. Cruaud, G. Genson, and J.-Y. Rasplus. 2012. Molecular systematics
and evolutionary history of the genus *Carabus* (Col. Carabidae). Molecular
Phylogenetics and Evolution **65**:259–275.

20. Ebach, M.C. 2015. Origins of Biogeography. Springer, New York.

21. Economo, E. P., E. M. Sarnat, M. Janda, R. Clouse, P. B. Klimov, G. Fischer, B. D.
Blanchard, L. N. Ramirez, A. N. Andersen, M. Berman, et al. 2015. Breaking out of
biogeographical modules: range expansion and taxon cycles in the hyperdiverse
ant genus Pheidole. Journal of Biogeography 42:2289–2301.

22. Edgar, R. C. 2004. MUSCLE: multiple sequence alignment with high accuracy and
high throughput. Nucleic Acids Research **32**:1792–1797.

23. Ehlers, J., and P. L. Gibbard. 2004. Quaternary glaciations-extent and chronology:
part I: Europe. Elsevier.

24. Ferrier, S., G. Manion, J. Elith, and K. Richardson. 2007. Using generalized
dissimilarity modelling to analyse and predict patterns of beta diversity in regional
biodiversity assessment. Diversity and Distributions **13**:252–264.

25. Gouveia, S. F., J. Hortal, M. Tejedo, H. Duarte, F. A. Cassemiro, C. A. Navas, and
J. A. F. Diniz-Filho. 2014. Climatic niche at physiological and macroecological
scales: the thermal tolerance–geographical range interface and niche
dimensionality. Global Ecology and Biogeography **23**:446–456.

26. GRASS Development Team, 2017. Geographic Resources Analysis Support System
(GRASS GIS) Software, Version 7.2. Open Source Geospatial Foundation.
http://grass.osgeo.org.

27. Guimerà, R., and L. A. N. Amaral. 2005. Functional cartography of complex
metabolic networks. Nature **433**:895–900.

28. Hewitt, G. M. 1999. Post-glacial re-colonization of European biota. Biological
Journal of the Linnean Society **68**:87–112.

29. Hijmans, R. J., S. E. Cameron, J. L. Parra, P. G. Jones, and A. Jarvis. 2005. Very high
resolution interpolated climate surfaces for global land areas. International
Journal of Climatology **25**:1965–1978.

30. Holt, B. G., J.-P. Lessard, M. K. Borregaard, S. A. Fritz, M. B. Araújo, D. Dimitrov, P.-
H. Fabre, C. H. Graham, G. R. Graves, K. A. Jønsson, et al. 2013. An update of
Wallace's zoogeographic regions of the world. Science **339**:74–78.

31. Homburg, K., C. Drees, M. M. Gossner, L. Rakosy, A. Vrezec, and T. Assmann. 2013.
Multiple glacial refugia of the low-dispersal ground beetle *Carabus irregularis*:
molecular data support predictions of species distribution models. PloS One
8:e61185.

32. Horreo, J.L., A. Jimenez-Valverde, and P.S. Fitze. 2016. Ecological change predicts
population dynamics and genetic diversity over 120 000 years. *Global Change*
Biology **22**:1737-1745.

33. Horreo J. L., M. L. Peláez, T. Suárez, M. C. Breedveld, B. Heulin, Y. Surget-Groba, T.
A. Oksanen, and P. S. Fitze. 2018. Phylogeography, evolutionary history, and effects
of glaciations in a species (Zootoca vivipara) inhabiting multiple biogeographic
regions*. Journal of Biogeography.*

34. Hortal, J., P. De Marco Jr, A. Santos, and J. A. F. Diniz-Filho. 2012. Integrating
703   biogeographical processes and local community assembly. Journal of
704   Biogeography **39**:627–628.

35. Hortal, J., J. A. F. Diniz-Filho, L. M. Bini, M. Á. Rodríguez, A. Baselga, D. Nogués-
706   Bravo, T. F. Rangel, B. A. Hawkins, and J. M. Lobo. 2011. Ice age climate,
707   evolutionary constraints and diversity patterns of European dung beetles. Ecology
708   Letters **14**:741–748.

36. Hortal, J., J. M. Lobo, and A. Jiménez-Valverde. 2012. Basic questions in
710   biogeography and the (lack of) simplicity of species distributions: putting species
711   distribution models in the right place. Natureza & Conservação **10**:108–118.

37. Hortal, J., N. Roura-Pascual, N. Sanders, and C. Rahbek. 2010. Understanding
713   (insect) species distributions across spatial scales. Ecography **33**:51.

38. Johnson, N. K., C. Cicero, and K. Shaw. 2004. New mitochondrial DNA data affirm
715   the importance of Pleistocene speciation in North American birds. Evolution
716   **58**:1122–1130.

39. Katoh, K., and D. M. Standley. 2013. MAFFT multiple sequence alignment software
718   version 7: improvements in performance and usability. Molecular Biology and
719   Evolution **30**:772–780.

40. Kembel, S. W., P. D. Cowan, M. R. Helmus, W. K. Cornwell, H. Morlon, D. D.
721   Ackerly, S. P. Blomberg, and C. O. Webb. 2010. Picante: R tools for integrating
722   phylogenies and ecology. Bioinformatics **26**:1463–1464.

41. Kluge, A.G. 1998. Total evidence or taxonomix congruence: cladistics or consensu
724   classification. *Cladistics* **14**:151-158.

42. Krasnov, B. R., M. A. Fortuna, D. Mouillot, I. S. Khokhlova, G. I. Shenbrot, and
726   R. Poulin. 2012. Phylogenetic signal in module composition and species
727   connectivity in compartmentalized host-parasite networks. The American
728   Naturalist **179**:501–511.

43. Lassmann, T. and E. L. Sonnhammer. 2005. Kalign - an accurate and fast multiple
730   sequence alignment algorithm. BMC bioinformatics **6**:1–9.

44. Lassmann, T. and E. L. Sonnhammer. 2006. Kalign, Kalignvu and Mumsa: web
732   servers for multiple sequence alignment. Nucleic Acids Research **34**: W596-W599

45. Legendre, P., and L. F. Legendre. 2012. Numerical ecology. Elsevier.

46. Losos, J. B., and R. E. Glor. 2003. Phylogenetic comparative methods and the
735   geography of speciation. Trends in Ecology and Evolution **18**:220–227.

47. Madrigal-González, J., P. Ruiz-Benito, S. Ratcliffe, J. Calatayud, G. Kändler,
737   A. Lehtonen, J. Dahlgren, C. Wirth, and M. A. Zavala. 2016. Complementarity
738   effects on tree growth are contingent on tree size and climatic conditions across
739   Europe. Scientific Reports **6**:32233.

48. Matzke, N. J., 2013. BioGeoBEARS: BioGeography with Bayesian (and Likelihood)
Evolutionary Analysis in R Scripts. University of California, Berkeley, Berkeley, CA.
http://CRAN.R-project.org/package=BioGeoBEARS.

49. Matzke, N. J. 2014. Model selection in historical biogeography reveals that
founder-event speciation is a crucial process in island clades. Systematic Biology
63:51–70.
50. Mazerolle, M. J. 2011. AICcmodavg: model selection and multimodel inference
based on (Q) AIC (c). R package version **1**.
51. Medina, N. G., B. Albertos, F. Lara, V. Mazimpaka, R. Garilleti, D. Draper, and
J. Hortal. 2014. Species richness of epiphytic bryophytes: drivers across scales on
the edge of the Mediterranean. Ecography **37**:80–93.
52. Mittelbach, G. G., and D. W. Schemske. 2015. Ecological and evolutionary
perspectives on community assembly. Trends in Ecology and Evolution **30**:241–
247.
53. Mucha, P. J., T. Richardson, K. Macon, M. A. Porter, and J.-P. Onnela. 2010.
Community structure in time-dependent, multiscale, and multiplex networks.
Science **328**:876–878.
54. Naeslund, B., and Norberg, J. 2006. Ecosystem consequences of the regional
species pool. *Oikos*, **115**:504-512.
55. O'Regan, H. J. 2008. The Iberian Peninsula–corridor or cul-de-sac? Mammalian
faunal change and possible routes of dispersal in the last 2 million years.
Quaternary Science Reviews **27**:2136–2144.
56. Overgaard, J., and J. G. Sørensen. 2008. Rapid thermal adaptation during field
temperature variations in *Drosophila melanogaster*. Cryobiology **56**:159–162.
57. Paradis, E., J. Claude and K. Strimmer. 2004. APE: analyses of phylogenetics and
evolution in R language. Bioinformatics **20**:289-290.
58. Provan, J., and Bennett, K. D. 2008. Phylogeographic insights into cryptic glacial
refugia. Trends in ecology & evolution **23**: 564-571.
59. Provine, W. B., 1989. Founder effects and genetic revolutions in microevolution
and speciation: an historical perspective. Pp. 43–76 *in* L. Val Giddings and K. Y.
Kaneshiro, editors. Genetics, speciation, and the founder principle. Oxford
University Press.
60. R Core Team, 2015. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/.
61. Rangel, T. F., R. K. Colwell, G. R. Graves, K. Fučíková, C. Rahbek, and J. A. F. Diniz-
Filho. 2015. Phylogenetic uncertainty revisited: Implications for ecological
analyses. Evolution **69**:1301–1312.
62. Ricklefs, R. E. 2008. Disintegration of the ecological community. The American
Naturalist **172**:741–750.
63. Ricklefs, R. E. 2011. Applying a regional community concept to forest birds of
eastern North America. Proceedings of the National Academy of Sciences USA
108:2300–2305.
64. Ricklefs, R. E. 2015. Intrinsic dynamics of the regional community. Ecology Letters
18:497–503.
65. Ricklefs, R. E., and F. He. 2016. Region effects influence local tree species diversity.
Proceedings of the National Academy of Sciences USA **113**:674–679.
66. Ree R.H., and I. Sanmartin. 2018. Conceptual and statistical problems with the
DEC+J model of founder-event speciation and its comparison with DEC via model
selection. Journal of Boigeography **45**:741-749

67. Ree, R. H., and S. A. Smith. (2008). Maximum likelihood inference of geographic
790   range evolution by dispersal, local extinction, and cladogenesis. Systematic biology
791   **57**:4-14.
792   68. Rueda, M., M. Á. Rodríguez, and B. A. Hawkins. 2013. Identifying global
793   zoogeographical regions: lessons from Wallace. Journal of Biogeography **40**:2215–
794   2225.
795   69. Schoener, T. W. 1970. Nonsynchronous spatial overlap of lizards in patchy
796   habitats. Ecology **51**:408–418.
797   70. Sommer, R. S., and Nadachowski, A. 2006. Glacial refugia of mammals in Europe:
798   evidence from fossil records. Mammal Review **36**: 251-265.
799   71. Svenning, J.-C., W. L. Eiserhardt, S. Normand, A. Ordonez, and B. Sandel. 2015. The
800   influence of paleoclimate on present-day patterns in biodiversity and ecosystems.
801   Annual Review of Ecology, Evolution and Systematics **46**:551–572.
802   72. Svenning, J.-C., and F. Skov. 2007. Could the tree diversity pattern in Europe be
803   generated by postglacial dispersal limitation? Ecology Letters **10**:453–460.
804   73. Tänzler, R., M. H. Van Dam, E. F. Toussaint, Y. R. Suhardjono, M. Balke, and
805   A. Riedel. 2016. Macroevolution of hyperdiverse flightless beetles reflects the
806   complex geological history of the Sunda Arc. Scientific Reports **6**:18793.
807   74. Theissinger, K., M. Bálint, K. A. Feldheim, P. Haase, J. Johannesen, I. Laube, and
808   S. U. Pauls. 2013. Glacial survival and post-glacial recolonization of an arctic–alpine
809   freshwater insect (*Arcynopteryx dichroa, Plecoptera, Perlodidae*) in Europe.
810   Journal of Biogeography **40**:236–248.
811   75. Thompson, J. D., D. G. Diggins, and T. J. Gibson. 1994. CLUSTAL W: improving the
812   sensitivity of progressive multiple sequence alignment through sequence
813   weighting, position-specific gap penalties and weight matrix choice. Nucleic Acids
814   Research **22**:4673–4680.
815   76. Toussaint, E. F., and M. Balke. 2016. Historical biogeography of Polyura butterflies
816   in the oriental Palaeotropics: trans-archipelagic routes and South Pacific island
817   hopping. Journal of Biogeography **43**:1560–1572.
818   77. Turin, H., L. Penev, A. Casale, E. Arndt, T. Assmann, K. Makarov, D. Mossakowski,
819   G. Szél, F. Weber, H. Turin, et al., 2003. Species accounts. Pp. 151–284 *in* H. Turin,
820   L. Penev, and A. Casale, editors. The Genus *Carabus* in Europe: A Synthesis.
821   Pensoft Publishers.
822   78. Ursenbacher, S., Carlsson, M., Helfer, V., Tegelström, H., and Fumagalli, L. 2006.
823   Phylogeography and Pleistocene refugia of the adder (Vipera berus) as inferred
824   from mitochondrial DNA sequence data. Molecular Ecology **15**: 3425-3437.
825   79. van Etten, J., 2015. gdistance: Distances and Routes on Geographical Grids.
826   http://CRAN.R-project.org/package=gdistance.
827   80. Wallace, A. R. 1876. The geographical distribution of animals: with a study of the
828   relations of living and extinct faunas as elucidating the past changes of the earth's
829   surface. Cambridge University Press.
830   81. Warren, D. L., M. Cardillo, D. F. Rosauer, and D. I. Bolnick. 2014. Mistaking
831   geography for biology: inferring processes from species distributions. Trends in
832   Ecology and Evolution **29**:572–580.
833   82. Weeks, B. C., S. Claramunt, and J. Cracraft. 2016. Integrating systematics and
834   biogeography to disentangle the roles of history and ecology in biotic assembly.
835   Journal of Biogeography **43**:1546–1559.

836   83. Wood, S.N., Goude, Y. and Shaw S. 2015. Generalized additive models for large
837   datasets. Journal of the Royal Statistical Society, Series C **64**: 139–155.
838   84. Wüest, R. O., A. Antonelli, N. E. Zimmermann, and H. P. Linder. 2015. Available
839   climate regimes drive niche diversification during range expansion. The American
840   Naturalist **185**:640–652.

 Figure 1.

Figure 1. Identifying the factors configuring regional faunas. a) Four (out of seven,

see b) hypothetical processes that may configure Regional faunas. Dotted lines depict

different regions while colours correspond with different climates. In each case, the tips

of the phylogeny point to regional distribution of the species. b) Workflow and potential

results: 1) Hypothetical results of modularity analysis over the occurrence network; 2)

similarity matrix of occurrence into modules; 3) pairwise matrix of environmental niche

similarities; phylogenetic distances and topographical connectivity; and 4) hypothetical

results and interpretations of a partial matrix regression on species occurrence

similarities as a function of niche similarities, phylogenetic distances and connectivity.

853 **Figure 2.**

a)

b)

Figure 2. Transition zones between regions were associated to geophysical

accidents and the border of the ice sheet at LGM. European *Carabus* regions found

by the network community detection analysis. a) Geographical location of modules (i.e.

regions) and submodules (i.e. sub-regions). b) Values of module affinity per grid cell;

green colours (i.e. cells with low affinity) identify transition zones. The dotted black

line corresponds with the southern limit of the ice sheet at LGM (extracted from Ehlers

and Gibbard 2004). The blue line depicts the breakpoint where the temperature-*Carabus*

richness relationship changes, as found in Calatayud et al. (2016b).

Figure 3

865
Figure 3. Regional co-occurrence was only explained by environmental niche

similarities and topographical connectivity. Results of the partial generalized matrix

regression of similarity in regional co-occurrence, as a function of environmental niche

similarity (climate and habitat), topographical connectivity and phylogenetic distances.

The first and second bars correspond with the models including co-occurrence

similarities among all modules and submodules, respectively. The remaining bars

correspond with the models where the similarities in submodule occurrence were

analysed independently for the species of each module.

Figure 4

875
Figure 4. Temporal coincidence between the Pleistocene and the phylogenetic

structuration of Carabus regions. a) GAMM predictions of the marginal probability

of the most probable state as a function of node age. The dashed red lines correspond

with the interval confidence at 95%. The dotted black line represents the median of the

breakpoint found by piecewise GLM regressions. The boxplot at the bottom represent

the 45th and 55th percentile breakpoint values, whereas the whiskers depict the 25th and

75th percentiles. b) Boxplot showing the probability of finding a phylogenetic node

having all descendant species grouped in the same region for pre-Pleistocene (pink) and

post- Pleistocene nodes (blue). This probability was calculated for all regions jointly

("All" in x axis) and independently (labelled according to Fig. 2a in x axis). c) An

example of a phylogenetic hypothesis where internal nodes are colored if all their

descendant species grouped in the same region. Node and tip colours correspond to the

regions where species were grouped, following Fig. 2a and the map at the bottom.

Lineages showing a high number of related species belonging to the same region are

higlighted as: "A" for *Platycarabus* subgenus; "B" for *Morphocarabus;* and "C" for

Orinocarabus . The dashed line corresponds with the average breakpoint (median)

where the probability of finding a node with all descendat species grouped in the same

region increases (see also Fig. S6). The shaded area depicts the 45^{th} and 55^{th} percentiles.