1

Drivers of genetic diversity in secondary metabolic gene clusters in a fungal population

Abigail L. Lind[1], Jennifer H. Wisecaver[2], Catarina Lameiras[3], Philipp Wiemann[4], Jonathan M.

Palmer[5], Nancy P. Keller[4], Fernando Rodrigues[6,7], Gustavo H. Goldman[8], Antonis Rokas[1,2]

1. Department of Biomedical Informatics, Vanderbilt University School of Medicine, Nashville,

Tennessee, USA.

2. Department of Biology, Vanderbilt University, Nashville, Tennessee, USA.

3. Department of Microbiology, Portuguese Oncology Institute of Porto, Porto, Portugal

4. Department of Medical Microbiology & Immunology, University of Wisconsin-Madison,

Madison, Wisconsin, USA

5. Center for Forest Mycology Research, Northern Research Station, US Forest Service, Madison,

Wisconsin, USA

6. Life and Health Sciences Research Institute (ICVS), School of Medicine, University of Minho,

Braga, Portugal

7. ICVS/3B's - PT Government Associate Laboratory, Braga/Guimarães, Portugal.

8. Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, São

Paulo, Brazil

†Corresponding author and lead contact: antonis.rokas@vanderbilt.edu

Summary

Filamentous fungi produce a diverse array of secondary metabolites (SMs) critical for defense,

virulence, and communication. The metabolic pathways that produce SMs are found in

contiguous gene clusters in fungal genomes, an atypical arrangement for metabolic pathways in

other eukaryotes. Comparative studies of filamentous fungal species have shown that SM gene

clusters are often either highly divergent or uniquely present in one or a handful of species,

hampering efforts to determine the genetic basis and evolutionary drivers of SM gene cluster

divergence. Here we examined SM variation in 66 cosmopolitan strains of a single species, the

opportunistic human pathogen *Aspergillus fumigatus.* Investigation of genome-wide

population-level variation showed that *A. fumigatus* strains contained five general types of

variation in SM gene clusters: non-functional gene polymorphisms, gene gain and loss

polymorphisms, whole cluster gain and loss polymorphisms, allelic polymorphisms where

different alleles corresponded to distinct, non-homologous clusters, and location

polymorphisms in which a cluster was found to differ in its genomic location across strains.

These polymorphisms affect the function of representative *A. fumigatus* SM gene clusters, such

as those involved in the production of gliotoxin, fumigaclavine, and helvolic acid, as well as the

function of clusters with undefined products. In addition to enabling the identification of

polymorphisms whose detection requires extensive genome-wide synteny conservation (e.g.,

mobile gene clusters and non-homologous cluster alleles), our population genomics approach

also implicated multiple underlying genetic drivers, including point mutations, recombination,

genomic deletion and insertion events, as well as horizontal gene transfer from distant fungi.

Finally, most of the population variants that we uncover have been previously hypothesized to

contribute to SM gene cluster diversity across entire fungal classes and phyla. We suggest that

the drivers of genetic diversity operating within a fungal population shown here are sufficient

to explain SM cluster macroevolutionary patterns.

Keywords: chemodiversity, specialized metabolism, genome evolution, genome architecture,

gene loss, genomic rearrangement

Introduction

Filamentous fungi produce a diverse array of small molecules that function as toxins, antibiotics, and pigments [1]. Though by definition these so-called specialized or secondary metabolites (SMs) are not strictly necessary for growth and development, they are critical to the lifestyle of filamentous fungi [2]. For example, antibiotic SMs gain give their fungal producers a competitive edge in environments crowded with other microbes [3]. SMs can additionally mediate communication between and within species, as well as contribute to virulence on animal and plant hosts in pathogenic fungi [4,5].

A genomic hallmark of SMs in filamentous fungi is that the biosynthetic pathways that produce them are typically organized into contiguous gene clusters in the genome [6]. These gene clusters contain the chemical backbone synthesis genes whose enzymatic products produce a core metabolite, such as non-ribosomal peptide synthases (NRPS) and polyketide synthases (PKS), tailoring enzymes that chemically modify the metabolite, transporters involved in product export, and transcription factors that control the expression of the clustered genes [6]. Filamentous fungal genomes, particularly those in the phylum Ascomycota [6], typically contain dozens of SM gene clusters. However, most individual SM gene clusters appear to be either species-specific or narrowly taxonomically distributed in only a handful of species [6,7]. SM gene clusters that are more broadly distributed show discontinuous taxonomic distributions and are often highly divergent between species. Consequently, the identity and total number of SM gene clusters can vary widely even between very closely related species whose genomes exhibit very high sequence and synteny conservation [8,9].

In the last decade, several comparative studies have described macroevolutionary patterns of SM gene cluster diversity. For example, studies centered on genomic comparisons of closely related species, such as members of the same genus, have identified several different types of inter-species divergence, from single nucleotide substitutions (e.g., differences in fumonisins produced by *Fusarium* species are caused by variants in one gene [10]), to gene gain / loss events (e.g., the trichothecene gene clusters in *Fusarium* species and the aflatoxin family SM gene clusters in *Aspergillus* species) [11–16], and genomic rearrangements (e.g., the

trichothecene gene clusters in *Fusarium*) [11]. Additionally, genetic and genomic comparisons

across fungal orders and classes have identified several instances of gene gain or loss [17–19]

and horizontal gene transfer [13,20–23] acting on individual genes or on entire gene clusters,

providing explanations for the diversity and discontinuity of the taxonomic distribution of

certain SM gene clusters across fungal species.

Although inter-species comparative studies have substantially contributed to our

understanding of SM diversity, the high levels of evolutionary divergence of SM clusters make

inference of the genetic drivers of SM gene cluster evolution challenging; simply put, it has

been difficult to "catch" the mechanisms that generate SM gene cluster variation "in the act".

Several previous studies have examined intra-species or population-level differences in

individual SM gene clusters, typically focusing on the presence and frequency of non-functional

alleles of clusters involved in production of mycotoxins. Examples of clusters exhibiting such

polymorphisms include the gibberellin gene cluster in *Fusarium oxysporum* [24], the fumonisin

gene cluster in *Fusarium fujikuroi* [25], the aflatoxin and cyclopiazonic gene clusters in

Aspergillus flavus [26], and the bikaverin gene cluster in *Botrytis cinerea* [27]. While these

studies have greatly advanced our understanding of SM gene cluster genetic variation and

highlighted the importance of population-level analyses, studies examining the entirety of SM

gene cluster polymorphisms in fungal populations are so far lacking. We currently do not

understand the types and frequency of SM gene cluster polymorphisms in populations,

whether these polymorphisms affect all types of SM gene clusters, as well as the genetic drivers

of SM gene cluster evolution.

To address these questions, we investigated the genetic diversity of all 36 known and

predicted SM gene clusters in whole genome sequence data from 66 strains, 8 of which were

sequenced in this study, of the opportunistic human pathogen *Aspergillus fumigatus*, a species

with cosmopolitan distribution and panmictic population structure [28]. We found that 13 SM

gene clusters were generally conserved and harbored low amounts of variation. In contrast, the

remaining 23 SM gene clusters were highly variable and contained one or more of five different

types of genetic variation: single-nucleotide polymorphisms including nonsense and frameshift

variants, individual gene gain and loss polymorphisms, entire cluster gain and loss

polymorphisms, polymorphisms associated with changes in cluster genomic location, and

clusters with non-homologous alleles resembling the idiomorphs of fungal mating loci. Many

clusters contained interesting combinations of these different polymorphisms, such as

pseudogenization in some strains and entire cluster loss in others. The types of variants we find

are likely generated by a combination of DNA replication and repair errors, recombination,

genomic insertions and deletions, and horizontal transfer. We additionally find an enrichment

for transposable elements (TEs) around horizontally transferred clusters, clusters that change in

genomic locations, and idiomorphic clusters. Taken together, our results provide a guide to

both the types of polymorphisms and the genetic drivers of SM gene cluster diversification in

filamentous fungi. As most of the genetic variants that we observe have been previously

associated with SM gene cluster diversity across much larger evolutionary distances and

timescales, we argue that population-level processes influencing SM gene cluster diversity are

sufficient to explain SM cluster macroevolutionary patterns.

Results

We analyzed the genomes of 66 globally distributed strains of *Aspergillus fumigatus* for polymorphisms in SM gene clusters. We performed whole-genome sequencing on 8 strains, and collected the remaining 58 strains from publicly available databases including NCBI Genome and the NCBI Short Read Archive (Figure 1, Table S1) [28–32]. We analyzed all strains for polymorphisms in 33 curated SM gene clusters present in the reference Af293 genome and additionally searched for novel SM gene clusters (see Methods). These examinations revealed five distinct types of polymorphisms which influence SM gene cluster variation (Table 1):

a) Single nucleotide and short indel polymorphisms. 33 / 33 SM gene clusters (present in the reference Af293 strain) contained multiple genes with missense SNPs and short indel variants in at least one strain in the population. 23 / 33 SM gene clusters contained one or more genes with frameshift or nonsense variants in the population.

b) Gene content polymorphisms involving loss or gain of one or more genes. 6 / 33 SM gene clusters contained a gene content polymorphism in the population.

c) Whole SM gene cluster gain and loss polymorphisms. 3 / 33 SM gene clusters present in the genome of the reference Af293 strain were absent in at least one strain in the population and an additional 3 previously unknown SM gene clusters were present in the population.

d) Idiomorphic polymorphisms. One locus contained multiple non-homologous SM gene cluster alleles in different strains of the population.

e) Genomic location polymorphisms. 2 / 33 SM gene clusters were found in different genomic locations (e.g., different chromosomes) between strains.

Single-nucleotide and indel polymorphisms

It is well established that single nucleotide polymorphisms (SNPs) and short indel polymorphisms are caused by errors in DNA replication and repair, and are a major source of genomic variation [33]. Non-synonymous SNPs and indels with missense, frameshift, and nonsense effects were widespread across the 33 SM reference gene clusters (Table S2). Every

strain contained numerous missense mutations and at least one nonsense or frameshift

mutation in its SM gene clusters. Although missense mutations are likely to influence SM

production, the functional effects of nonsense and frameshift mutations are comparatively

easier to infer from genomic sequence data because they often lead to truncated proteins

lacking a significant portion of their amino acid sequence. For example, a frameshift mutation in

the polyketide synthase (PKS) of the trypacidin gene cluster in the A1163 strain results in loss of

trypacidin production [34]. Interestingly, we identified a premature stop codon (Gln273*) in a

transcription factor required for trypacidin production, *tpcD,* in a strain sequenced in this study

(MO79587EXP). These data suggest that function of this SM gene cluster has been lost through

at least two independent genetic events in *A. fumigatus*.

Individual nonsense or frameshift variants ranged from very common in the population

to rarer variants present in one or a handful of strains. For example, the non-ribosomal peptide

synthase (NRPS) *pes3* gene (Afu5g12730) in SM gene cluster 21 harbors 16 nonsense or

frameshift polymorphisms in 55 strains. Seven of these polymorphisms are common and

present in 10 or more strains, while seven are rarer and found in 5 or fewer strains. Strains with

lab-mutated null alleles of the *pes3* gene are more virulent than strains with functional copies

[35], which may explain the widespread occurrence of null *pes3* alleles in the *Aspergillus*

population.

Gene content polymorphisms

We additionally identified several SM gene clusters that gained or lost genes in some

strains. These gene content polymorphisms were most likely generated through genomic

deletion or insertion events and were often present in high frequencies in the population (Table

1). In three cases, these polymorphisms impact backbone synthesis genes, rendering the SM

gene cluster non-functional. One example involves SM gene cluster 14, whose standard

composition includes a pyoverdine synthase gene, an NRPS-like gene, an NRPS backbone gene,

and several additional modification genes (Figure 2A). We discovered that 4 / 66 strains lack an

11-kb region on the 3' end of the cluster which normally contains an NRPS gene and two

additional cluster genes, and the first non-SM genes on the 3' end flanking the cluster. All *A.*

fumigatus strains contain a *copia* family transposable element [36,37] at the 3' end of the

cluster, suggesting that transposable elements may have been involved in the generation of this

polymorphism. Three of the strains lacking the 11-Kb region form a clade in the SNP-based

phylogeny (the fourth, RP-2014 was not included in this phylogeny as short read data were not

available). While this polymorphism could have arisen through a deletion event, a homologous

cluster lacking the 11-kb region is also present in the reference genomes of *Aspergillus lentulus*

and *Aspergillus fischerianus*, close relatives of *A. fumigatus* (Figure 2). The most parsimonious

explanation is that the genome of the *A. fumigatus* ancestor contained an SM gene cluster that

lacked the 11-kb region, and that this genomic region was subsequently gained and increased in

frequency in the *A. fumigatus* population.

Two additional gene content polymorphisms affecting SM backbone genes were

restricted to one strain each and appear to have arisen through genomic deletion events.

Specifically, strain IF1SWF4 lacks an 8-Kb region near the helvolic acid SM gene cluster,

resulting in the loss of the backbone oxidosqualene cyclase gene as well an upstream region

containing two non-SM genes (Figure S1). Strain LMB35Aa lacks a 54-kb region on the end of

chromosome 2, which includes five genes from the telomere-proximal fumigaclavine C cluster

(Figure S1).

In three other cases, gene content polymorphisms involved gene loss or truncation

events of non-backbone structural genes. We found that the second half of the ORF of the *gliM*

O-methyltransferase gene in the gliotoxin gene cluster has been lost in 2 / 66 strains (Figure

S1), and that the first half of the permease *fmqE* in the fumiquinazoline gene cluster has been

lost in 4 / 66 strains (Figure S1). Finally, an ABC transporter gene in SM cluster 21 has been

almost entirely lost in 21 / 66 strains (Figure S2). This deletion event is found in strains that are

related in the SNP phylogeny but does not perfectly mirror population structure (Figure S2).

Whole gene cluster gain and loss polymorphisms

Several SM gene clusters were gained or lost entirely in 13 / 66 strains. We observed

instances where a cluster present in the genome of the reference Af293 strain was absent or

pseudogenized in other strains as well as cases in which SM clusters present in other strains

were absent from the reference Af293 strain.

The most notable example of an SM gene cluster that was present in the Af293

reference genome but absent or pseudogenized in others was SM cluster 4. This cluster

contains 5 genes on the tip of the Af293 chromosome 1 and contains orthologs to five of the six

genes in the fusarielin gene cluster in *Fusarium graminearum* [38]. This cluster is also present in

several other *Aspergillus* species, including *A. clavatus* and *A. niger* [38]. Phylogenetic analysis

of the genes in this SM gene cluster is consistent with horizontal gene transfer between fungi in

the class Sordariomycetes and fungi in the class Eurotiomycetes, or alternatively with extensive

gene loss in both Sordariomycetes and Eurotiomycetes (Figure S3). This gene cluster is entirely

absent in 4 / 66 strains, and its genes are undergoing pseudogenization in an additional 43

strains via multiple independent mutational events (Figure 3A). These four strains form a single

clade on the SNP-based phylogeny, suggesting that this cluster was lost in a single deletion

event (Figure 3B). Further, 19 strains shared a single frameshift variant in the polyketide

synthase gene (4380_4381insAATGGGCT; frameshift at Glu1461 in Afu1g17740) and an

additional 13 strains shared a single frameshift variant (242delG; frameshift at Gly81) in an

aldose 1-epimerase gene (Afu1g17723). Eleven other strains each contained one to several

frameshift or nonsense polymorphisms involving nine unique mutational sites, suggesting that

this pathway is undergoing multiple independent pseudogenization events. Five of these strains

contained multiple distinct frameshifts and premature stop codons in more than one gene in

the cluster, indicating that the entire pathway is pseudogenized in these strains. The

distributions of pseudogenizing variants only in the aldose 1-epimerase gene and of

pseudogenizing variants in multiple genes in the cluster roughly correlate with population

structure, while the distribution of pseudogenizing variants in the PKS backbone gene does not

(Figure 3B).

By searching for novel SM gene clusters in the genomes of the other 65 *A. fumigatus*

strains, we found three SM gene clusters that were absent from the genome of the Af293

reference strain. As SM gene clusters are often present in repeat-rich and subtelomeric regions

that are challenging to assemble [39,40], these strains might harbor additional novel SM gene

clusters.

One of the novel SM gene clusters that we identified, cluster 34, was present in all but

two of the strains (Af293 and F7763). This cluster contains a PKS backbone gene, one PKS-like

gene with a single PKS associated domain, nine genes with putative biosynthetic functions

involved in secondary metabolism, and six hypothetical proteins (Figure 3C). The two strains

that lack this cluster contain a likely non-functional cluster fragment that includes the PKS-like

gene, two biosynthetic genes, and three hypothetical proteins. Interestingly, the 3' region

flanking this cluster is syntenic across all 66 strains but the 5' region is not, suggesting that a

recombination or deletion event may have resulted in the loss of this cluster in the Af293 and

F7763 strains.

The other SM gene clusters that were absent from the Af293 genome are present at

lower frequencies in the population; cluster 35 is present in 2 / 66 strains and cluster 36 in 4 /

66 strains. Cluster 35 is located in a region syntenic with an Af293 chromosome 4 region and is

flanked on both sides by transposable elements (Figure S4). The two strains that contain this

novel cluster are not sister to each other in the SNP-based phylogeny (Figure S4). Eight of the

14 genes in this SM gene cluster are homologous to genes in an SM gene cluster in the genome

of the insect pathogenic fungus *Metarhizium anisopliae* (Figure S4). Phylogenetic analysis of

these 8 genes is consistent with a horizontal transfer event (Figure S5). Cluster 36 is an NRPS

containing cluster located on genomic scaffolds that lack homology to either the Af293 or

A1163 genomes, making it impossible to determine on which chromosome this cluster is

located (Figure S3). Two of the strains containing this novel cluster are sister to each other in

the SNP-based phylogeny, while the third is distant to these two (Figure S4). The evolutionary

histories of the genes in the cluster are consistent with vertical inheritance and are present in

multiple *Aspergillus* species.

Idiomorph polymorphisms

One of the most peculiar types of polymorphisms that we identified is a locus containing

different unrelated alleles of SM gene clusters, reminiscent of the idiomorph alleles at the

fungal mating loci [41]. This locus, which resides on chromosome 3 and corresponds to cluster

10 in the Af293 genome (Figure 4), was previously described as being strain-specific in a

comparison between Af293 and A1163 [30] and is thought to reside in a recombination hot

spot [28]. Our analysis showed that there are at least 6 different alleles of this cluster in the

population, containing 4 different types of key enzymes involved in natural product

biosynthesis: a polyketide synthase (PKS) non-ribosomal peptide synthetase (NRPS) hybrid, a

highly reducing (HR) PKS, a non-reducing (NR) PKS and an NRPS-like enzyme (Figure 4). Two

additional alleles were present in only one strain each (Figure S6).

In the Af293 reference genome, the cluster present at this locus contains one NR-PKS

along with an NRPS-like gene (Allele B). In the A1163 reference genome and 17 other strains,

there is a PKS-NRPS and an HR-NRPS (Allele E). These alleles show an almost complete lack of

sequence similarity except for a conserved hypothetical protein and a fragment of the HR-PKS

in the Af293 allele; in contrast, the upstream and downstream flanking regions of the two

alleles, which do not contain any backbone genes, are syntenic. Remarkably, another allele,

present in 12 strains, contains all of the genes from both the Af293 and A1163 clusters (Allele

D). The remaining three alleles contain various combinations of these genes. One allele found in

22 strains contains some A1163-specific genes including the HR-PKS and no Af293-specific

genes (Allele F), while another allele found in 3 strains contains some Af293-specific genes

including the NRPS-like gene but no A1163 genes (Allele A). The final allele, present in 8 strains,

contains the entire Af293 allele as well as part of the A1163 allele containing the HR-PKS (Allele

C). Every allele is littered with long terminal repeat sequence fragments from *gypsy* and *copia*

TE families as well as with sequence fragments from DNA transposons from the *mariner* family

[36]. In some cases, these TEs correspond with breakpoints in synteny between alleles,

suggesting that the diverse alleles of this SM gene cluster may arise via TE-driven

recombination. Further, both of the alleles that are restricted to a single strain had an insertion

event of several genes near a TE, though the rest of the locus is highly similar to one of the

more common alleles (Figure S6).

Preliminary metabolite analysis comparing one of the strains (08-19-02-30) that

contains the allele harboring all four key enzymes (Allele D) to two strains (08-12-12-13 and 08-

19-02-10) that only codes for the putative cluster the HR-PKS backbone gene (Allele F) by

untargeted XCMS analysis [42] revealed the presence of 2 unique masses in 08-19-02-30 (Table

S4; Fig. 4B). Further analysis is underway to investigate whether any of these m/z can be

directly linked to the allele present in 08-19-02-30.

The evolutionary history of this highly diverse locus is unclear (Figure 4C). While it is

tempting to speculate that the largest allele containing all observed genes represents the

ancestral state, it does not explain the presence of a shared hypothetical protein and PKS gene

fragment between the Af293 locus (Allele C) and the A1163 locus (Allele B). The distributions of

Alleles A, B, C, and F roughly correlate with the population structure of the strains, while the

distribution of Alleles D and E do not (Figure 4C).

Genomic location polymorphisms

The final type of polymorphism that we observed is associated with SM gene clusters

that are located in different genomic locations in different strains, suggesting that these SM

gene clusters are behaving like mobile elements. This type of polymorphism was observed in

SM gene clusters 1 and 33, both of which produce as yet identified products, and are present at

low frequencies in the population.

SM gene cluster 1, which is present in six strains at three different genomic locations

(Figure 5A), consists of a PKS and four other modification genes that are always flanked by a 15

Kb region (upstream) and a 43 Kb region (downstream) containing TEs. In the reference Af293

strain and in strain F7763, this SM gene cluster and its flanking regions are located on

chromosome 1, while in strains dutch3, F13619, and Z5 they are located between Afu4g07320

and Afu4g07340 on chromosome 4. In contrast, in strain JCM_10253, the cluster and flanking

regions are located on chromosome 8 immediately adjacent to the 3' end of the intertwined

fumagillin and pseurotin SM gene supercluster [43]. The strains containing the allele on

Chromosome 1 are sister to each other in the SNP-based phylogeny, while the other strains are

scattered across the tree and do not reflect the underlying population structure (Figure 5C).

In 5 / 6 strains, the cluster appears to be functional and does not contain nonsense SNPs

or indels. However, the cluster found on chromosome 1 in strain F7763 contains two stop

codons in the oxidoreductase gene (Gln121* and Gln220*) and two premature stop codons in

the polyketide synthase (Gln1156* and Gln1542*), suggesting this strain contains a null allele.

This "jumping" gene cluster is not present in any other sequenced *Aspergillus* genus,

and phylogenetic analysis of its constituent genes is consistent with horizontal gene transfer

between fungi (Figure S7). Specifically, this gene cluster is also present in *Phaeosphaeria*

nodorum, a plant pathogen from the class Dothideomycetes, *Pseudogymnoascus pannorum*, a

fungus isolated from permafrost from the Leotiomycetes, *Escovopsis weberi*, and a fungal

parasite of fungus-growing ants from the Sordariomycetes (Figure 5B). One additional species,

the endophyte *Hypoxylon* sp. CI4A from the class Sordariomycetes, contains four of the five

cluster genes but is missing Afu1g00970, an MFS drug transporter. However, this species

contains an unrelated gene annotated as an MFS drug transporter immediately adjacent to this

cluster, so this species may be using a different transporter (Figure 5B). None of these fungi

contain the upstream or downstream TE-rich flanking regions present in *A. fumigatus*, and each

fungus contains additional unique genes with putative biosynthetic functions adjacent to the

transferred cluster. The most likely explanation for this change in flanking regions is that this

SM gene cluster was transferred into *A. fumigatus* once and has subsequently moved its

location in the genome.

The second SM gene cluster that shows variation in its genomic location across strains,

cluster 33, contains a terpene synthase. This cluster is present in only 5 strains at 3 distinct

locations (Figure S8). Similar to cluster 1, this cluster is also flanked by TEs, and in one strain the

clusters is located 58 Kb from another SM gene cluster. Two strains that contain the cluster in

the same genomic location are sister to each other in the SNP-based phylogeny, while the other

three strains containing the cluster do not reflect the underlying population structure (Figure

S8). In contrast to cluster 1, this cluster does not appear to have been horizontally transferred

between fungi and its genes are present in other sequenced *Aspergillus* species. As this mobile

cluster was not horizontally transferred, it is possible that the horizontal transfer of cluster 1

and its mobile nature throughout the *A. fumigatus* genome are driven by different mechanisms.

Discussion

Our examination of 66 genomes from strains of *Aspergillus fumigatus* revealed that five

general types of polymorphisms describe variation in SM gene clusters in a fungal population.

These polymorphisms include variation in single nucleotides, gene and gene cluster gains and

losses, non-homologous clusters at the same genomic position, and changes in genomic

locations of clusters (Figure 6). In several cases, the genetic mechanisms that gave rise to these

polymorphisms and their functional consequences are clear. Polymorphisms in single-

nucleotides most likely arose from errors in DNA replication or repair, while gene loss

polymorphisms likely arose from genomic deletion or recombination events. Others, such as SM

gene clusters that exist as non-homologous alleles or that entered the population through

horizontal transfer, could arise through multiple genetic mechanisms. Transposable elements

are common in polymorphic SM gene clusters and are found flanking mobile and horizontally

transferred clusters, as well as in regions adjacent to non-homologous alleles and where gene

gain has occurred, suggesting they contribute to SM gene cluster diversity. Using a population

genomics approach to identify SM gene cluster variants allowed us to also capture and describe

novel polymorphisms, including mobile gene clusters and idiomorphic clusters, which are

difficult to identify in comparative genomic studies between species whose conservation of

genome-wide synteny is low. Strikingly, the variants and genetic drivers we observe at the

population level are also implicated as driving SM gene cluster variation between fungal

species, suggesting that the observed microevolutionary processes are sufficient to explain

macroevolutionary patterns of SM gene cluster evolution.

The first novel type of polymorphism was observed in SM gene clusters 1 and 33, which

by occupying different genomic locations in different strains, appear to behave in a manner

similar to mobile genetic elements. Interestingly, both clusters are located near or immediately

adjacent to other SM gene clusters in some strains. For example, cluster 1 is located

immediately adjacent to the intertwined fumagillin and pseurotin supercluster [43] in one

strain. This supercluster is regulated by the transcriptional factor *fapR* and is located in a

chromosomal region controlled by the master SM regulators *laeA* and *veA* [43,44], raising the

hypothesis that these mobile gene clusters might be co-opting the regulatory machinery

already in place. Previous work has hypothesized that the fumagillin and pseurotin supercluster

formed through genomic rearrangement events placing the once-independent gene clusters in

close proximity to each other [43]. Our observation that this mobile gene cluster is located in

this same region not only supports this hypothesis but also implicates transposable elements as

one of the mechanisms by which such genomic rearrangements are formed. These

superclusters may also represent an intermediate stage in the formation of new SM gene

clusters. Supercluster formation, potentially mediated by mobile gene clusters, and followed

by gene loss, could explain macroevolutionary patterns of SM gene clusters have shown that

clustered genes in one species can be dispersed over multiple gene clusters in other species

[9,11].

The second novel type of polymorphism was the presence of multiple non-homologous

alleles at the cluster 10 locus, echoing the structure of the idiomorphic mating type locus [41].

This region has previously been reported to be a recombination hotspot in *A. fumigatus* [28]

and all alleles contain numerous transposable elements. Thus, it is possible that polymorphism

at this locus originated via SM gene cluster fusion or splitting events driven by transposable

elements. Interestingly, two other previously described instances of SM gene cluster variation

bear close resemblance to the *A. fumigatus* idiomorphic SM gene cluster 10 locus. The first is

the presence of two non-homologous *Aspergillus flavus* alleles, where some strains contain a 9-

gene sesquiterpene-like SM gene cluster and others contain a non-homologous 6-gene SM gene

cluster at the same genomic location [45]. The second is the presence of two non-homologous

SM gene clusters at the same, well-conserved, locus in a comparison of six species of

dermatophyte fungi [46]. Based on these results, we hypothesize that idiomorphic clusters may

be common in fungal populations and contribute to the broad diversity of SM gene clusters

across filamentous fungi.

The remaining types of SM polymorphism in the *A. fumigatus* population have

previously been described at the species level. For example, null alleles have been reported for

many individual SM gene clusters across diverse fungal species [10,17,26,27] and our results

show that it is a widespread phenomenon, affecting two-thirds of the *A. fumigatus* SM gene

clusters. Consistent with previous literature reporting the presence of additional SM gene

clusters not present in the reference strain [24,25], we also identify four low-frequency SM

gene clusters in the *A. fumigatus* population, two of which are horizontally transferred from

distantly related fungi and two that appear vertically inherited. We find numerous cases of

gene loss and relatively fewer cases of gene gain; previous studies have implicated these

processes across numerous types of SM gene clusters and reported their effects on metabolite

production [47]. Previous work has indicated that gene and gene cluster gain and loss tends to

occur near telomeres, suggesting that higher rates of genetic events like recombination lead to

loss [48]. While we do find several cases of gene and gene cluster loss in subtelomeric clusters,

including the fumigaclavine gene cluster and the helvolic acid gene cluster, we also find gene

loss in clusters that are not located near telomeres, such as the gliotoxin gene cluster and the

fumiquinazoline gene cluster. These findings suggest that while gene and gene cluster loss can

occur in telomeric regions, clusters in other genomic regions also experience high rates of loss.

Previous work has demonstrated that SM gene clusters, like the metabolites that they

produce, are highly divergent between fungal species. Our examination of genome-wide

variation shows that these SM gene clusters are correspondingly diverse within individual

strains of a single fungal species. Furthermore, the observed genetic changes in SM gene

clusters are widespread across different types of gene clusters and are caused by many

underlying genetic drivers, including gene and gene cluster gain, loss, non-homologous cluster

alleles, and mobile gene clusters. The net effect of these substitutions, gains, losses, and

rearrangements raises the hypothesis that fungal SM gene clusters are likely in a state of

evolutionary flux, constantly altering their SM gene cluster repertoire, and consequently

modifying and diversifying their chemodiversity.

Methods

Strains analyzed

Eight strains of *A. fumigatus* were isolated from four patients with recurrent cases of
aspergillosis in the Portuguese Oncology Institute in Porto, Portugal. Each strain was
determined to be *A. fumigatus* using macroscopic features of the culture and microscopic
morphology observed in the slide preparation from the colonies with lactophenol solution [49].
Based on the morphological characterization, all clinical strains were classified as *A. fumigatus*
complex-Fumigati. After whole genome sequencing, retrieval and examination of the beta
tubulin and calmodulin sequences of each strain confirmed that all strains belonged to *A.*
fumigatus (see Phylogenetic Analysis and Figure S9)*.* The genomes of all eight strains were
sequenced using 150bp Illumina paired-end sequence reads at the Genomic Services Lab of
Hudson Alpha (Huntsville, Alabama, USA). Genomic libraries were constructed with the Illumina
TruSeq library kit and sequenced on an Illumina HiSeq 2500 sequencer. Samples of all eight
strains were sequenced at greater than 180X coverage or depth (Table S1). Short read
sequences for these 8 strains are available in the Short Read Archive under accession
SRP109032.

In addition to the 8 strains sequenced in this study, we retrieved 58 *A. fumigatus* strains
with publicly available whole genome sequencing data, resulting in a population genomics
dataset of 66 strains (Table S1). The strains used included both environmental and clinical
strains and were isolated from multiple continents. Genome assemblies for 10 of these strains,
including the Af293 and A1163 reference strains, were available for download from GenBank
[28–32,50]. For 6 of these strains, short read sequences were also available from the NCBI Short
Read Archive (SRA), which were used for variant discovery only (see Single nucleotide variant
(SNV) and indel discovery) and not for genome assembly. Short read sequences were not
available for the remaining 4 strains. Short read sequences were downloaded for an additional
48 strains from the Short Read Archive if they were sequenced with paired-end reads and at
greater than 30x coverage.

Single nucleotide variant (SNV) and indel discovery

All strains with available short read data (62 of 66 strains) were aligned to both the Af293 and A1163 reference genomes using BWA mem version 0.7.12-r1044 [51]. Coverage of genes present in the reference genome was calculated using bedtools v2.25.0 [52]. SNV and indel discovery and genotyping was performed relative to the Af293 reference genome and was conducted across all samples simultaneously using the Genome Analysis Toolkit version 3.5-0-g36282e4 with recommended hard filtering parameters [53–55] and annotated using snpEff version 4.2 [56].

De novo **genome assembly and gene annotation**

All 56 strains without publicly available genome assemblies were *de novo* assembled using the iWGS pipeline [57]. Specifically, all strains were assembled using SPAdes v3.6.2 and MaSuRCA v3.1.3 and resulting assemblies were evaluated using QUAST v3.2 [58–60]. The average N50 of assemblies constructed with this strategy was 463 KB (Table S1). Genes were annotated in these assemblies as well as in five GenBank assemblies with no predicted genes using augustus v3.2.2 trained on *A. fumigatus* gene models [61]. Repetitive elements were annotated in all assemblies using RepeatMasker version open-4.0.6 [62].

Secondary metabolic gene cluster annotation and discovery

Secondary metabolic gene clusters in the Af293 reference genome were taken from two recent reviews, both of which considered computational and experimental data to delineate cluster boundaries [63,64] (Table S3). The genomes of the other 65 strains were scanned for novel SM gene clusters using identified using antiSMASH v3.0.5.1 [65]. To prevent potential assembly errors from confounding the analysis, any inference about changes in genomic locations of genes or gene clusters was additionally verified by manually inspecting alignments and ensuring that paired end reads supported an alternative genomic location (see SNV and indel discovery). Cases where paired end reads did not support the change in genomic location or where mapping was ambiguous or low quality were discarded.

Phylogenetic analysis

To confirm all strains in this analysis belonged to the species *Aspergillus fumigatus*, the

genomic sequences of the beta tubulin and calmodulin genes were extracted from the

assembled genomes of all strains. Gene phylogenies were constructed using *Aspergillus*

fischerianus as an outgroup using RAxML v8.0.25 with the GTRGAMMA substitution model [66].

The tree was midpoint rooted and all branches with bootstrap support less than 80% were

collapsed (Figure S9).

To construct a SNP-based strain phylogeny, biallelic SNPs with no missing data were

pruned using SNPRelate v1.8.0 with a linkage disequilibrium threshold of 0.8 [67]. A total of

15,274 SNVs were used to create a phylogeny using RAxML v8.0.25 with the ASC_BINGAMMA

substitution model [66]. The tree was midpoint rooted and all branches with bootstrap support

less than 80% were collapsed.

To understand the evolutionary histories of specific SM gene clusters showing unusual

taxonomic distributions, we reconstructed the phylogenetic trees of their SM genes.

Specifically, SM cluster protein sequences were queried against a local copy of the NCBI non-

redundant protein database (downloaded May 30, 2017) using phmmer, a member of the

HMMER3 software suite [68] using acceleration parameters --F1 1e-5 --F2 1e-7 --F3 1e-10. A

custom perl script sorted the phmmer results based on the normalized bitscore (nbs), where

nbs was calculated as the bitscore of the single best-scoring domain in the hit sequence divided

by the best bitscore possible for the query sequence (i.e., the bitscore of the query aligned to

itself). No more than five hits were retained for each unique NCBI Taxonomy ID. Full-length

proteins corresponding to the top 100 hits (E-value < 1 × 10-10) to each query sequence were

extracted from the local database using esl-sfetch [68]. Sequences were aligned with MAFFT

v7.310 using the E-INS-i strategy and the BLOSUM30 amino acid scoring matrix [69] and

trimmed with trimAL v1.4.rev15 using its gappyout strategy [70]. The topologies were inferred

using maximum likelihood as implemented in RAxML v8.2.9 [66] using empirically determined

substitution models and rapid bootstrapping (1000 replications). The phylogenies were

midpoint rooted and branches with less than 80% bootstrap support were collapsed using the

ape and phangorn R packages [71,72]. Phylogenies were visualized using ITOL version 3.0 [73].

Differential metabolite analysis

For natural product analysis, 5 x 106 spores/mL for the indicated strains were grown in

50 mL liquid GMM [74] for five days at 25 °C and 250 rpm in duplicates. Supernatants were

extracted with equal volumes of ethyl acetate, dried down and resuspened in 20% acetonitrile

(ACN). Each sample was analyzed by ultra-high performance liquid chromatography (UHPLC)

coupled with mass spectrometry (MS). The samples were separated on a ZORBAX Eclipse XDB-

C18 column (Agilent, 2.1 x 150 mm with a 1.8 µM particle size using a binary gradient of 0.5 %

(v/v) formic acid (FA) as solvent A and 0.5 % (v/v) FA in ACN as solvent B that was delivered by a

VanquishTM UHPLC system (Thermo Scientific) with a flow rate of 0.2 mL/min. The binary

gradient started with 20% B that was increased with a linear gradient to 100% B in 15 min

followed by an isocratic step at 100% B for 5 min. Before every run, the system was equilibrated

for 5 min at 20% B. The UHPLC system was coupled to a Q Exactive hybrid quadrupole

OritrapTM MS (Thermo Scientific). For electrospray ionization, the ion voltage was set at +/-3.5

kV in positive and negative mode, respectively. Nitrogen was used as sheath gas at a flow rate

of 45 and as sweep gas at a flow rate of 2. Data analysis was performed using XCMS [42] and

Maven [75] software.

Acknowledgements

A.L.L. was supported by the U.S. National Library of Medicine training grant 2T15LM007450.

This work was supported in part by the National Science Foundation (DEB-1442113 to A.R.) and

by the Northern Portugal Regional Operational Programme (NORTE 2020), under the Portugal

2020 Partnership Agreement, through the European Regional Development Fund (FEDER)

(NORTE-01-0145-FEDER-000013 to F.R.). G.H.G. thanks the Fundação de Amparo à Pesquisa do

Estado de São Paulo (FAPESP) and Conselho Nacional de Desenvolvimento Científico e

Tecnológico (CNPq), both from Brazil, for funding and support. This work was conducted in part

using the resources of the Advanced Computing Center for Research and Education at

Vanderbilt University (http://www.accre.vanderbilt.edu/).

Author contributions

Conceptualization, A.R., G.H.G., A.L.L.; Methodology, A.L.L., J.H.W, C.L, P.W., J.M.P.;

Investigation, A.L.L.; Visualization, A.L.L., J.H.W, P.W.; Resources, G.H.G., F.R., N.P.K., C.L.;

Writing, A.L.L, A.R.

583   **Table 1. Types and rates of SM gene cluster variants in *A. fumigatus* strains.**

Description	Phenotype	Drivers	Frequency at cluster level	Frequency at strain level	Previous reports
Single-nucleotide polymorphisms and indels	Potential for protein function change (missense); abrogation of protein function (nonsense and frameshift)	DNA replication errors; relaxation of purifying selection	100% (33/33 clusters; missense); 70% (23/33 clusters; nonsense and frameshift)	Every strain affected	Bikaverin in *Botrytis* [17,27], aflatoxin in *Aspergillus oryzae* and *Aspergillus flavus* [26], fumonisins in *Fusarium* [10], many others
Gene content polymorphisms	Loss of gene cluster function; structural changes in the metabolite; change in cluster expression or metabolite transport	Deletion and insertion events; recombination; transposable elements	6 clusters	27 / 66 strains	Trichothecene in *Fusarium*, aflatoxin and sterigmatocystin in *Aspergillus* [11–15]
Whole gene cluster polymorphisms	Loss or gain of novel metabolites	Deletion and insertion events; horizontal gene transfer; transposable elements	6 clusters	13 / 66 strains	Gibberellin and fumonisin in *Fusarium* [24,25]
fCluster idiomorphs	Changes in metabolites produced or structure of metabolites	Transposable elements; recombination; other mechanisms?	1 gene cluster	8 unique identified alleles	Putative SM gene clusters in dermatophytes; putative SM gene cluster in *Aspergillus flavus* and *Aspergillus oryzae* [45,46]
Mobile gene clusters	Potential for change in gene regulation	Transposable elements; horizontal gene transfer; other mechanisms?	2 gene clusters	8 / 66 strains	None

585
586

Table S1. Summary of strains, sequence data, and assemblies used.

Table S2. All nonsynonymous variants in SM gene cluster genes.

Table S3. Description, locus tags, and gene annotations of all reference Af293 SM gene

clusters.

Table S4. XCMS analysis of extracted supernatants from 08-19-02-30, 08-12-12-13, and 08-19-

02-10.

References

1. Vining LC. Functions of secondary metabolites. Annu Rev Microbiol. Annual Reviews 4139 El Camino Way, P.O. Box 10139, Palo Alto, CA 94303-0139, USA; 1990;44: 395–427. doi:10.1146/annurev.mi.44.100190.002143

2. Schimek C. Evolution of Special Metabolism in Fungi : Concepts , Mechanisms , and Pathways. In: Pöggler S, Wöstmeyer J, editors. Evolution of Fungi and Fungal-Like Organisms, The Mycota. XIV. Berlin, Heidelberg: Springer-Verlag; 2011. pp. 293–328.

3. Fox EM, Howlett BJ. Secondary metabolism: regulation and role in fungal biology. Curr Opin Microbiol. 2008;11: 481–7. doi:10.1016/j.mib.2008.10.007

4. Scharf DH, Heinekamp T, Brakhage AA. Human and Plant Fungal Pathogens: The Role of Secondary Metabolites. Heitman J, editor. PLoS Pathog. Public Library of Science; 2014;10: e1003859. doi:10.1371/journal.ppat.1003859

5. Yim G, Wang HH, Davies J. Antibiotics as signalling molecules. Philos Trans R Soc Lond B Biol Sci. 2007;362: 1195–200. doi:10.1098/rstb.2007.2044

6. Keller NP, Turner G, Bennett JW. Fungal secondary metabolism - from biochemistry to genomics. Nat Rev Microbiol. 2005;3: 937–47. doi:10.1038/nrmicro1286

7. Bennett J, Bentley R. What's in a name?—Microbial secondary metabolism. Adv Appl Microbiol. 1989;34.

8. Khaldi N, Seifuddin FT, Turner G, Haft D, Nierman WC, Wolfe KH, et al. SMURF: Genomic mapping of fungal secondary metabolite clusters. Fungal Genet Biol. 2010;47: 736–41. doi:10.1016/j.fgb.2010.06.003

9. Lind AL, Wisecaver JH, Smith TD, Feng X, Calvo AM, Rokas A. Examining the evolution of the regulatory circuit controlling secondary metabolism and development in the fungal genus Aspergillus. PLoS Genet. Public Library of Science; 2015;11: e1005096. doi:10.1371/journal.pgen.1005096

10. Proctor RH, Busman M, Seo J-A, Lee YW, Plattner RD. A fumonisin biosynthetic gene cluster in Fusarium oxysporum strain O-1890 and the genetic basis for B versus C fumonisin production. Fungal Genet Biol. 2008;45: 1016–1026. doi:10.1016/j.fgb.2008.02.004

11. Proctor RH, McCormick SP, Alexander NJ, Desjardins AE. Evidence that a secondary metabolic biosynthetic gene cluster has grown by gene relocation during evolution of the filamentous fungus Fusarium. Mol Microbiol. 2009;74: 1128–42. doi:10.1111/j.1365-2958.2009.06927.x

12. Berry DB, Guan Q, Hose J, Haroon S, Gebbia M, Heisler LE, et al. Multiple Means to the Same End: The Genetic Basis of Acquired Stress Resistance in Yeast. Copenhaver GP,

editor. PLoS Genet. Public Library of Science; 2011;7: e1002353.
doi:10.1371/journal.pgen.1002353

13.   Slot JC, Rokas A. Horizontal transfer of a large and highly toxic secondary metabolic gene
cluster between fungi. Curr Biol. Elsevier Ltd; 2011;21: 134–9.
doi:10.1016/j.cub.2010.12.020

14.   Ehrlich KC, Chang P-K, Yu J, Cotty PJ. Aflatoxin Biosynthesis Cluster Gene cypA Is Required
for G Aflatoxin Formation. Appl Environ Microbiol. 2004;70: 6518–6524.
doi:10.1128/AEM.70.11.6518-6524.2004

15.   Carbone I, Ramirez-Prado JH, Jakobek JL, Horn BW. Gene duplication, modularity and
adaptation in the evolution of the aflatoxin gene cluster. BMC Evol Biol. 2007;7: 111.
doi:10.1186/1471-2148-7-111

16.   Yu J, Chang P-K, Ehrlich KC, Cary JW, Bhatnagar D, Cleveland TE, et al. Clustered pathway
genes in aflatoxin biosynthesis. Appl Environ Microbiol. American Society for
Microbiology; 2004;70: 1253–62. doi:10.1128/AEM.70.3.1253-1262.2004

17.   Campbell M a, Rokas A, Slot JC. Horizontal transfer and death of a fungal secondary
metabolic gene cluster. Genome Biol Evol. 2012;4: 289–93. doi:10.1093/gbe/evs011

18.   Kroken S, Glass NL, Taylor JW, Yoder OC, Turgeon BG. Phylogenomic analysis of type I
polyketide synthase genes in pathogenic and saprobic ascomycetes. Proc Natl Acad Sci U
S A. 2003;100: 15670–5. doi:10.1073/pnas.2532165100

19.   Bushley KE, Turgeon BG. Phylogenomics reveals subfamilies of fungal nonribosomal
peptide synthetases and their evolutionary relationships. BMC Evol Biol. 2010;10: 26.
doi:10.1186/1471-2148-10-26

20.   Patron NJ, Waller RF, Cozijnsen AJ, Straney DC, Gardiner DM, Nierman WC, et al. Origin
and distribution of epipolythiodioxopiperazine (ETP) gene clusters in filamentous
ascomycetes. BMC Evol Biol. 2007;7: 174. doi:10.1186/1471-2148-7-174

21.   Khaldi N, Collemare J, Lebrun M-H, Wolfe KH. Evidence for horizontal transfer of a
secondary metabolite gene cluster between fungi. Genome Biol. BioMed Central; 2008;9:
R18. doi:10.1186/gb-2008-9-1-r18

22.   Khaldi N, Wolfe KH. Evolutionary Origins of the Fumonisin Secondary Metabolite Gene
Cluster in Fusarium verticillioides and Aspergillus niger. Int J Evol Biol. Hindawi Publishing
Corporation; 2011;2011: 423821. doi:10.4061/2011/423821

23.   Reynolds HT, Slot JC, Divon HH, Lysøe E, Proctor RH, Brown DW. Differential Retention of
Gene Functions in a Secondary Metabolite Cluster. Mol Biol Evol. Oxford University Press
(OUP); 10: e1004816. doi:10.1093/molbev/msx145

24.   Wiemann P, Sieber CMK, von Bargen KW, Studt L, Niehaus E-M, Espino JJ, et al.
Deciphering the Cryptic Genome: Genome-wide Analyses of the Rice Pathogen Fusarium

fujikuroi Reveal Complex Regulation of Secondary Metabolism and Novel Metabolites.
Mitchell AP, editor. PLoS Pathog. Public Library of Science; 2013;9: e1003475.
doi:10.1371/journal.ppat.1003475

25.   Chiara M, Fanelli F, Mulè G, Logrieco AF, Pesole G, Leslie JF, et al. Genome Sequencing of
675   Multiple Isolates Highlights Subtelomeric Genomic Diversity within *Fusarium fujikuroi*.
676   Genome Biol Evol. 2015;7: 3062–3069. doi:10.1093/gbe/evv198

677   26.   Chang P-K, Horn BW, Dorner JW. Sequence breakpoints in the aflatoxin biosynthesis gene
678   cluster and flanking regions in nonaflatoxigenic Aspergillus flavus isolates. Fungal Genet
679   Biol. 2005;42: 914–923. doi:10.1016/j.fgb.2005.07.004

680   27.   Schumacher J, Gautier A, Morgant G, Studt L, Ducrot P-H, Le Pêcheur P, et al. A
681   Functional Bikaverin Biosynthesis Gene Cluster in Rare Strains of Botrytis cinerea Is
682   Positively Controlled by VELVET. Yun S-H, editor. PLoS One. Public Library of Science;
683   2013;8: e53729. doi:10.1371/journal.pone.0053729

684   28.   Abdolrasouli A, Rhodes J, Beale MA, Hagen F, Rogers TR, Chowdhary A, et al. Genomic
685   Context of Azole Resistance Mutations in Aspergillus fumigatus Determined Using
686   Whole-Genome Sequencing. MBio. 2015;6. doi:10.1128/mBio.00536-15

687   29.   Nierman WC, Pain A, Anderson MJ, Wortman JR, Kim HS, Arroyo J, et al. Genomic
688   sequence of the pathogenic and allergenic filamentous fungus Aspergillus fumigatus.
689   Nature. 2005;438: 1151–6. doi:10.1038/nature04332

690   30.   Fedorova ND, Khaldi N, Joardar VS, Maiti R, Amedeo P, Anderson MJ, et al. Genomic
691   islands in the pathogenic filamentous fungus Aspergillus fumigatus. Richardson PM,
692   editor. PLoS Genet. Public Library of Science; 2008;4: e1000046.
693   doi:10.1371/journal.pgen.1000046

694   31.   Knox BP, Blachowicz A, Palmer JM, Romsdahl J, Huttenlocher A, Wang CCC, et al.
695   Characterization of Aspergillus fumigatus Isolates from Air and Surfaces of the
696   International Space Station. mSphere. 2016;1.

697   32.   Paul S, Zhang A, Ludeña Y, Villena GK, Yu F, Sherman DH, et al. Insights from the genome
698   of a high alkaline cellulase producing Aspergillus fumigatus strain obtained from Peruvian
699   Amazon rainforest. J Biotechnol. 2017;251: 53–58. doi:10.1016/j.jbiotec.2017.04.010

700   33.   Roberts JD, Kunkel TA. Fidelity of DNA replication. In: DePamphilis ML, editor. DNA
701   replication in eukaryotic cells. Cold Spring Harbor, New York: Cold Spring Harbor
702   Laboratory Press; 1996. pp. 217–247.

703   34.   Throckmorton K, Lim FY, Kontoyiannis DP, Zheng W, Keller NP. Redundant synthesis of a
704   conidial polyketide by two distinct secondary metabolite clusters in Aspergillus
705   fumigatus. Environ Microbiol. 2015; doi:10.1111/1462-2920.13007

35.   O'Hanlon KA, Cairns T, Stack D, Schrettl M, Bignell EM, Kavanagh K, et al. Targeted

Disruption of Nonribosomal Peptide Synthetase pes3 Augments the Virulence of
Aspergillus fumigatus. Infect Immun. 2011;79: 3978–3992. doi:10.1128/IAI.00192-11

36. Kapitonov V V., Jurka J. A universal classification of eukaryotic transposable elements
implemented in Repbase. Nat Rev Genet. Nature Publishing Group; 2008;9: 411–412.
doi:10.1038/nrg2165-c1

37. Mount SM, Rubin GM. Complete nucleotide sequence of the Drosophila transposable
element copia: homology between copia and retroviral proteins. Mol Cell Biol. 1985;5:
1630–8.

38. Sørensen JL, Hansen FT, Sondergaard TE, Staerk D, Lee TV, Wimmer R, et al. Production
of novel fusarielins by ectopic activation of the polyketide synthase 9 cluster in Fusarium
graminearum. Environ Microbiol. Blackwell Publishing Ltd; 2012;14: 1159–1170.
doi:10.1111/j.1462-2920.2011.02696.x

39. Treangen TJ, Salzberg SL. Repetitive DNA and next-generation sequencing: computational
challenges and solutions. Nat Rev Genet. NIH Public Access; 2011;13: 36–46.
doi:10.1038/nrg3117

40. Palmer JM, Keller NP. Secondary metabolism in fungi: does chromosomal location
matter? Curr Opin Microbiol. 2010;13: 431–6. doi:10.1016/j.mib.2010.04.008

41. Metzenberg RL, Glass NL. Mating type and mating strategies inNeurospora. BioEssays.
Wiley Subscription Services, Inc., A Wiley Company; 1990;12: 53–59.
doi:10.1002/bies.950120202

42. Smith CA, Want EJ, O'Maille G, Abagyan R, Siuzdak G. XCMS: Processing Mass
Spectrometry Data for Metabolite Profiling Using Nonlinear Peak Alignment, Matching,
and Identification. Anal Chem. 2006;78: 779–787. doi:10.1021/ac051437y

43. Wiemann P, Guo C-J, Palmer JM, Sekonyela R, Wang CCC, Keller NP. Prototype of an
intertwined secondary-metabolite supercluster. Proc Natl Acad Sci U S A. 2013;110:
17065–70. doi:10.1073/pnas.1313258110

44. Lin H-C, Chooi Y-H, Dhingra S, Xu W, Calvo AM, Tang Y. The Fumagillin Biosynthetic Gene
Cluster in Aspergillus fumigatus Encodes a Cryptic Terpene Cyclase Involved in the
Formation of β-trans-Bergamotene. J Am Chem Soc. 2013;135: 4616–9.
doi:10.1021/ja312503y

45. Gibbons JG, Salichos L, Slot JC, Rinker DC, McGary KL, King JG, et al. The evolutionary
imprint of domestication on genome variation and function of the filamentous fungus
Aspergillus oryzae. Curr Biol. 2012;22: 1403–9. doi:10.1016/j.cub.2012.05.033

46. Zhang H, Rokas A, Slot JCJ. Two different secondary metabolism gene clusters occupied
the same ancestral locus in fungal dermatophytes of the Arthrodermataceae. PLoS One.
2012;7: e41903. doi:10.1371/journal.pone.0041903

47. Schardl CL, Young CA, Hesse U, Amyotte SG, Andreeva K, Calie PJ, et al. Plant-Symbiotic
Fungi as Chemical Engineers: Multi-Genome Analysis of the Clavicipitaceae Reveals
Dynamics of Alkaloid Loci. Heitman J, editor. PLoS Genet. Public Library of Science;
2013;9: e1003323. doi:10.1371/journal.pgen.1003323

48. Young CA, Schardl CL, Panaccione DG, Florea S, Takach JE, Charlton ND, et al. Genetics,
genomics and evolution of ergot alkaloid diversity. Toxins. 2015. pp. 1273–1302.
doi:10.3390/toxins7041273

49. de Hoog G, Guarro J, Gené J, Figueras M. Atlas of Clinical Fungi. Washington, DC: ASM
Press; 2001.

50. Liu D, Zhang R, Yang X, Wu H, Xu D, Tang Z, et al. Thermostable cellulase production of
Aspergillus fumigatus Z5 under solid-state fermentation and its application in
degradation of agricultural wastes. Int Biodeterior Biodegradation. 2011;65: 717–725.
doi:10.1016/j.ibiod.2011.04.005

51. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 2009;25: 1754–1760. doi:10.1093/bioinformatics/btp324

52. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. Oxford University Press; 2010;26: 841–2.
doi:10.1093/bioinformatics/btq033

53. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The Genome
Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing
data. Genome Res. Cold Spring Harbor Laboratory Press; 2010;20: 1297–303.
doi:10.1101/gr.107524.110

54. Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, et
al. From FastQ Data to High-Confidence Variant Calls: The Genome Analysis Toolkit Best
Practices Pipeline. Current Protocols in Bioinformatics. Hoboken, NJ, USA: John Wiley &
Sons, Inc.; 2013. p. 11.10.1-11.10.33. doi:10.1002/0471250953.bi1110s43

55. DePristo MA, Banks E, Poplin R, Garimella K V, Maguire JR, Hartl C, et al. A framework for
variation discovery and genotyping using next-generation DNA sequencing data. Nat
Genet. 2011;43: 491–498. doi:10.1038/ng.806

56. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. A program for
annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in
the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly (Austin). Taylor &
Francis; 2012;6: 80–92. doi:10.4161/fly.19695

57. Zhou X, Peris D, Kominek J, Kurtzman CP, Hittinger CT, Rokas A. in silico Whole Genome
Sequencer & Analyzer (iWGS): A Computational Pipeline to Guide the Design and
Analysis of de novo Genome Sequencing Studies. G3:
Genes|Genomes|Genetics. 2016; doi:10.1534/g3.116.034249

58.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new
781   genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol.
782   Mary Ann Liebert, Inc.; 2012;19: 455–77. doi:10.1089/cmb.2012.0021

783   59.   Zimin A V., Marcais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. The MaSuRCA genome
784   assembler. Bioinformatics. 2013;29: 2669–2677. doi:10.1093/bioinformatics/btt476

785   60.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome
786   assemblies. Bioinformatics. 2013;29: 1072–1075. doi:10.1093/bioinformatics/btt086

787   61.   Stanke M, Morgenstern B. AUGUSTUS: a web server for gene prediction in eukaryotes
788   that allows user-defined constraints. Nucleic Acids Res. Oxford University Press; 2005;33:
789   W465-7. doi:10.1093/nar/gki458

790   62.   Smit A, Hubley R, Green P. Repeatmasker Open-4.0 [Internet]. Available:
791   http://www.repeatmasker.org

792   63.   Inglis DO, Binkley J, Skrzypek MS, Arnaud MB, Cerqueira GC, Shah P, et al.
793   Comprehensive annotation of secondary metabolite biosynthetic genes and gene
794   clusters of Aspergillus nidulans, A. fumigatus, A. niger and A. oryzae. BMC Microbiol.
795   2013;13: 91. doi:10.1186/1471-2180-13-91

796   64.   Bignell E, Cairns TC, Throckmorton K, Nierman WC, Keller NP. Secondary metabolite
797   arsenal of an opportunistic pathogenic fungus. Philos Trans R Soc B Biol Sci. 2016;371.

798   65.   Medema MH, Blin K, Cimermancic P, de Jager V, Zakrzewski P, Fischbach M a, et al.
799   antiSMASH: rapid identification, annotation and analysis of secondary metabolite
800   biosynthesis gene clusters in bacterial and fungal genome sequences. Nucleic Acids Res.
801   2011;39: W339-46. doi:10.1093/nar/gkr466

802   66.   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
803   phylogenies. Bioinformatics. 2014;30: 1312–1313. doi:10.1093/bioinformatics/btu033

804   67.   Zheng X, Levine D, Shen J, Gogarten SM, Laurie C, Weir BS. A high-performance
805   computing toolset for relatedness and principal component analysis of SNP data.
806   Bioinformatics. Oxford University Press; 2012;28: 3326–3328.
807   doi:10.1093/bioinformatics/bts606

808   68.   Eddy SR. A new generation of homology search tools based on probabilistic inference.
809   Genome Inform. 2009;23: 205–11.

810   69.   Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7:
811   Improvements in Performance and Usability. Mol Biol Evol. 2013;30: 772–780.
812   doi:10.1093/molbev/mst010

813   70.   Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. trimAl: a tool for automated
814   alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25: 1972–
815   1973. doi:10.1093/bioinformatics/btp348

816   71.   Paradis E, Claude J, Strimmer K. APE: Analyses of Phylogenetics and Evolution in R
817   language. Bioinformatics. Oxford University Press; 2004;20: 289–290.
818   doi:10.1093/bioinformatics/btg412

819   72.   Schliep KP. phangorn: phylogenetic analysis in R. Bioinformatics. Oxford University Press;
820   2011;27: 592–3. doi:10.1093/bioinformatics/btq706

821   73.   Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and
822   annotation of phylogenetic and other trees. Nucleic Acids Res. 2016;44: W242–W245.
823   doi:10.1093/nar/gkw290

824   74.   Shimizu K, Keller NP. Genetic involvement of a cAMP-dependent protein kinase in a G
825   protein signaling pathway regulating morphological and chemical transitions in
826   Aspergillus nidulans. Genetics. 2001;157: 591–600.

827   75.   Melamud E, Vastag L, Rabinowitz JD. Metabolomic Analysis and Visualization Engine for
828   LC−MS Data. Anal Chem. 2010;82: 9818–9826. doi:10.1021/ac1021166

Figure 1. SNP-based phylogeny of *A. fumigatus* strains. The phylogeny was constructed using 15,274 biallelic SNPs with no missing data. The tree is midpoint rooted and all branches with bootstrap support less than 80% are collapsed. This phylogeny does not include strains Af10, Af210, Z5, or RP-2014 as short read data were not available.

Figure 2. Differences in gene content in SM gene cluster 14 in A. fumigatus strains and closely related species.
Four *A. fumigatus* strains lack an 11-Kb region in this cluster, including an NRPS backbone gene. (A) Regions upstream and downstream of this cluster are syntenic. LMB35Aa also contains a large inversion that moves a transcription factor, oxidoreductase, and hypothetical protein 275 kb away from the cluster. *Aspergillus fischerianus* and *Aspergillus lentulus*, close relatives of *A. fumigatus*, contain a cluster lacking the 11-kb region. (B) Strains without 11-Kb region highlighted in red.

Figure 3. Pseudogenization and gene loss in SM gene clusters. (A) Positions of frameshift variants and nonsense variants in the fusarielin-like SM gene cluster 4. (B) Strains lacking the fusarielin-like cluster or containing frameshift and nonsense variants in the cluster. (C) SM gene cluster found in most *A. fumigatus* strains but absent from the Af293 reference and from the F7763 strain. EOS denotes end of scaffold.

Figure 4. Six alleles of an idiomorphic SM gene cluster. (A) Alleles of SM gene cluster 10 on chromosome 3. Red boxes denote transposable elements. Green arrows denote backbone genes (PKS or NRPS). (B) Extracted ion chromatograms for the two m/z identified in negative mode from XCMS analysis comparing extracts from strains with Alleles D and F. (C) Distribution of the six cluster alleles across a SNP-based phylogeny of strains. Bolded and italicized strains have unique alleles but appear derived from another allele (see also Figure S5).

Figure 5. Multiple genomic locations of a horizontally transferred SM gene cluster. (A) SM gene cluster 1 (Afu1g00970-01010) and flanking region is found in different genomic locations.The flanking regions contain transposon-derived open reading frames including two putative reverse transcriptases. (B) Synteny of A. fumigatus SM gene cluster 1 with clusters in *Phaeosphaeria nodorum*, *Pseudogymnoascus pannorum*, *Escovopsis weberi*, and *Hypoxylon sp. CI4A*. EOS denotes end of scaffold. All species contain non-syntenic genes predicted by antiSMASH to be part of a biosynthetic gene cluster. (C) Strains containing SM gene cluster 1 in SNP-based phylogeny.

Figure 6. Types and frequencies of all SM gene cluster variants in the *A. fumigatus* population.

Figure S1. Alignments showing deletion of genes in SM gene clusters. A) Deletion of helvolic acid genes in IF1SWF4.
B) Deletion of fumigaclavine genes in LMB35Aa. C) Partial deletion of gliM in the gliotoxin gene cluster in two strains.
D) Partial deletion of fmqE in the fumiquinazoline gene cluster in 3 strains.

Figure S1. Partial deletion of ABC transporter gene in SM gene cluster 21. (A,B) Coverage of 21 strains with partial deletion of SM gene cluster 21. (C) Position of strains with partial gene deletion in SNP based phylogeny.

Figure S2. Gene phylogenies of the fusarielin-like SM gene cluster 4. These phylogenies are consistent with horizontal transfer between Eurotiomycete and Sordariomycete fungi or with extensive gene loss.

S2A

Tree scale: 0.1

S2B

Tree scale: 0.1 ⊢———⊣

Taxonomy
- Leotiomycetes
- Sordariomycetes
- Eurotiomycetes
- Dothideomycetes
- other_Pezizomycotina
- other_Ascomycota
- other_Fungi
- other_Opisthokonta
- other_Eukaryota
- Bacteria
- Archaea
- Viruses

XP 018156681.1-759273-Colletotrichum higginsian-Sordariomycetes
ENH79013.1-1213857-Colletotrichum orbiculare-Sordariomycetes
CZT02535.1-914238-Rhynchosporium agropyri-Leotiomycetes
CZT43006.1-38038-Rhynchosporium secalis-Leotiomycetes
CZS89923.1-914237-Rhynchosporium commune-Leotiomycetes
XP 013277062.1-1442369-Rhinocladiella mackenziei-Eurotiomycetes
XP 007736464.1-1182542-Capronia epimyces CBS 606-Eurotiomycetes
XP 016240750.1-91928-Exophiala spinifera-Eurotiomycetes
XP 016258459.1-215243-Exophiala oligosperma-Eurotiomycetes
XP 007827070.1-1229662-Pestalotiopsis fici W106-Sordariomycetes
XP 007817533.1-655844-Metarhizium robertsii ARS-Sordariomycetes
KFG81511.1-5530-Metarhizium anisopliae-Sordariomycetes
KJK74649.1-1291518-Metarhizium anisopliae BR-Sordariomycetes
XP 014540250.1-1276141-Metarhizium brunneum ARSE-Sordariomycetes
KID84095.1-1276136-Metarhizium guizhouense A-Sordariomycetes
XP 014573814.1-1276143-Metarhizium majus ARSEF 2-Sordariomycetes
XP 013431313.1-1043004-Aureobasidium namibiae CB-Dothideomycetes
GAW21476.1-1813822-fungal sp No 14919-other Fungi
XP 018155051.1-759273-Colletotrichum higginsian-Sordariomycetes
EQL01261.1-911162-Ophiocordyceps sinensis C-Sordariomycetes
CZT13654.1-914237-Rhynchosporium commune-Leotiomycetes
KDN72350.1-1173701-Colletotrichum sublineola-Sordariomycetes
XP 008100387.1-645133-Colletotrichum graminicol-Sordariomycetes
XP 002544623.1-336963-Uncinocarpus reesii 1704-Eurotiomycetes
XP 013340600.1-1043005-Aureobasidium subglaciale-Dothideomycetes
GAM82793.1-1603295-fungal sp No 11243-other Fungi
G3XMB7.1-380704-Aspergillus niger ATCC 10-Eurotiomycetes
XP 001393494.1-425011-Aspergillus niger CBS 513-Eurotiomycetes
XP 001594902.1-665079-Sclerotinia sclerotiorum-Leotiomycetes
XP 001560980.1-332648-Botrytis cinerea B05 10-Leotiomycetes
ESZ98313.1-1432307-Sclerotinia borealis F 41-Leotiomycetes
XP 008086832.1-1116229-Glarea lozoyensis ATCC 20-Leotiomycetes
KIN01422.1-913774-Oidiodendron maius Zn-Leotiomycetes
XP 018075682.1-149040-Phialocephala scopiformis-Leotiomycetes
OCK95926.1-794803-Cenococcum geophilum 1 58-Dothideomycetes
OCL10303.1-574774-Glonium stellatum-Dothideomycetes
XP 007293085.1-1072389-Marssonina brunnea f sp-Leotiomycetes
XP 018077409.1-149040-Phialocephala scopiformis-Leotiomycetes
CZT52794.1-38038-Rhynchosporium secalis-Leotiomycetes
CZT04210.1-914237-Rhynchosporium commune-Leotiomycetes
CZT13906.1-914238-Rhynchosporium agropyri-Leotiomycetes
ORY67810.1-1141098-Pseudomassariella vexata-Sordariomycetes
OJJ34618.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
KGQ02700.1-1245745-Beauveria bassiana D1 5-Sordariomycetes
XP 011319442.1-229533-Fusarium graminearum PH 1-Sordariomycetes
XP 009262582.1-1028729-Fusarium pseudograminearu-Sordariomycetes
KZL72807.1-708197-Colletotrichum tofieldiae-Sordariomycetes
XP 007603123.1-1445577-Colletotrichum fioriniae-Sordariomycetes
KXH26574.1-703756-Colletotrichum simmondsii-Sordariomycetes
XP 001270372.1-344612-Aspergillus clavatus NRRL-Eurotiomycetes
KOS43413.1-229535-Penicillium nordicum-Eurotiomycetes
OQD93913.1-60172-Penicillium solitum-Eurotiomycetes
CRL30865.1-5075-Penicillium camemberti-Eurotiomycetes
GAT21367.1-1069201-Aspergillus luchuensis-Eurotiomycetes
GAA83286.1-1033177-Aspergillus kawachii IFO-Eurotiomycetes
QUERY-Afu1g17720-330879-Aspergillus fumigatus Af2-Eurotiomycetes
KEY83928.1-1476192-Aspergillus fumigatus var-Eurotiomycetes
XP 001275177.1-344612-Aspergillus clavatus NRRL-Eurotiomycetes
EYE92016.1-1388766-Aspergillus ruber CBS 135-Eurotiomycetes
OJJ83693.1-1160497-Aspergillus glaucus CBS 5-Eurotiomycetes
ODM24346.1-573508-Aspergillus cristatus-Eurotiomycetes
KOC18776.1-1392242-Aspergillus flavus AF70-Eurotiomycetes
XP 001827115.1-510516-Aspergillus oryzae RIB40-Eurotiomycetes
XP 002384353.1-332952-Aspergillus flavus NRRL33-Eurotiomycetes
KJK67560.1-1403190-Aspergillus parasiticus S-Eurotiomycetes
XP 015412269.1-1509407-Aspergillus nomius NRRL 1-Eurotiomycetes
XP 020054870.1-690307-Aspergillus aculeatus ATC-Eurotiomycetes
OJJ68416.1-767769-Aspergillus brasiliensis-Eurotiomycetes
GAA92532.1-1033177-Aspergillus kawachii IFO-Eurotiomycetes
OJZ90081.1-1137211-Aspergillus luchuensis CB-Eurotiomycetes
OJI84725.1-767770-Aspergillus tubingensis C-Eurotiomycetes
GAQ35625.1-5061-Aspergillus niger-Eurotiomycetes
EHA20992.1-380704-Aspergillus niger ATCC 10-Eurotiomycetes
XP 001390020.1-425011-Aspergillus niger CBS 513-Eurotiomycetes
GAW19657.1-1813822-fungal sp No 14919-other Fungi
EKG13971.1-1126212-Macrophomina phaseolina M-Dothideomycetes
XP 007587274.1-1287680-Neofusicoccum parvum UCRN-Dothideomycetes
XP 020131637.1-236234-Diplodia corticola-Dothideomycetes
KKY24657.1-420778-Diplodia seriata-Dothideomycetes
OMP82498.1-420778-Diplodia seriata-Dothideomycetes
XP 008097989.1-645133-Colletotrichum graminicol-Sordariomycetes
OLN87541.1-708187-Colletotrichum chlorophyt-Sordariomycetes
XP 018161333.1-759273-Colletotrichum higginsian-Sordariomycetes
CCF43873.1-80884-Colletotrichum higginsian-Sordariomycetes
KZL77770.1-708197-Colletotrichum tofieldiae-Sordariomycetes
KZL83287.1-1573173-Colletotrichum incanum-Sordariomycetes
EQB56236.1-1237896-Colletotrichum gloeospori-Sordariomycetes
ENH76248.1-1213857-Colletotrichum orbiculare-Sordariomycetes
OHF02959.1-1209926-Colletotrichum orchidophi-Sordariomycetes
XP 007598459.1-1445577-Colletotrichum fioriniae-Sordariomycetes
KXH51170.1-1460502-Colletotrichum nymphaeae-Sordariomycetes
KXH33030.1-703756-Colletotrichum simmondsii-Sordariomycetes
KXH47137.1-1209931-Colletotrichum salicis-Sordariomycetes
XP 018758495.1-334819-Fusarium verticillioides-Sordariomycetes
CZR48165.1-1227346-Fusarium proliferatum ET1-Sordariomycetes
S0EES1.1-1279085-Fusarium fujikuroi IMI 58-Sordariomycetes
OBS18895.1-36050-Fusarium poae-Sordariomycetes
XP 009259311.1-1028729-Fusarium pseudograminearu-Sordariomycetes
XP 011327621.1-229533-Fusarium graminearum PH 1-Sordariomycetes

S2C

Transcription factor Oxidoreductase aldose 1-epimerase family protein CYP450 zinc binding enoyl reductase PKS

Afu1g17710 720 723 725 730 740

Tree scale: 0.1

Taxonomy

- Leotiomycetes
- Sordariomycetes
- Eurotiomycetes
- Dothideomycetes
- other_Pezizomycotina
- other_Ascomycota
- other_Fungi
- other_Opisthokonta
- other_Eukaryota
- Bacteria
- Archaea
- Viruses

XP 008081690.1-1116229-Glarea lozoyensis ATCC 20-Leotiomycetes
XP 007827071.1-1229662-Pestalotiopsis fici W106-Sordariomycetes
KID84094.1-1276136-Metarhizium guizhouense A-Sordariomycetes
KJK74648.1-1291518-Metarhizium anisopliae BR-Sordariomycetes
EXU96140.1-568076-Metarhizium robertsii-Sordariomycetes
KFG81512.1-5530-Metarhizium anisopliae-Sordariomycetes
XP 009262581.1-1028729-Fusarium pseudograminearu-Sordariomycetes
EYB31406.1-5518-Fusarium graminearum-Sordariomycetes
XP 011319441.1-229533-Fusarium graminearum PH 1-Sordariomycetes
OJJ34619.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
ORY67811.1-1141098-Pseudomassariella vexata-Sordariomycetes
XP 007919597.1-1286976-Phaeoacremonium minimum U-Sordariomycetes
KZL72809.1-708197-Colletotrichum tofieldiae-Sordariomycetes
KXH26575.1-703756-Colletotrichum simmondsii-Sordariomycetes
XP 007603122.1-1445577-Colletotrichum fioriniae-Sordariomycetes
XP 001270373.1-344612-Aspergillus clavatus NRRL-Eurotiomycetes
KOS43410.1-229535-Penicillium nordicum-Eurotiomycetes
CRL30864.1-5075-Penicillium camemberti-Eurotiomycetes
OQD93854.1-60172-Penicillium solitum-Eurotiomycetes
KEY83927.1-1476192-Aspergillus fumigatus var-Eurotiomycetes
QUERY-Afu1g17723-330879-Aspergillus fumigatus Af2-Eurotiomycetes
KMK54953.1-1437362-Aspergillus fumigatus Z5-Eurotiomycetes
EHA18932.1-380704-Aspergillus niger ATCC 10-Eurotiomycetes
GAA83287.1-1033177-Aspergillus kawachii IFO-Eurotiomycetes
GAT21368.1-1069201-Aspergillus luchuensis-Eurotiomycetes
GAM89934.1-1603295-fungal sp No 11243-other Fungi
OCK79255.1-1314670-Lepidopterella palustris-Dothideomycetes
KDB17570.1-1159556-Ustilaginoidea virens-Sordariomycetes
OLN87181.1-708187-Colletotrichum chlorophyt-Sordariomycetes
EME45950.1-675120-Dothistroma septosporum N-Dothideomycetes
XP 007925836.1-383855-Pseudocercospora fijiensi-Dothideomycetes
XP 016217501.1-253628-Verruconis gallopava-Dothideomycetes
XP 016217502.1-253628-Verruconis gallopava-Dothideomycetes
KPM41344.1-78410-Neonectria ditissima-Sordariomycetes
XP 003041148.1-660122-Nectria haematococca mpVI-Sordariomycetes
XP 003004171.1-526221-Verticillium alfalfae VaM-Sordariomycetes
XP 009654134.1-498257-Verticillium dahliae VdLs-Sordariomycetes
OQO06661.1-1507870-Rachicladosporium antarct-Dothideomycetes
OQO18097.1-1974281-Rachicladosporium sp CCF-Dothideomycetes
OQO10055.1-1507870-Rachicladosporium antarct-Dothideomycetes
KND94354.1-1163406-Tolypocladium ophioglosso-Sordariomycetes
KYK61204.1-98403-Drechmeria coniospora-Sordariomycetes
ODA80971.1-98403-Drechmeria coniospora-Sordariomycetes
OAA79236.1-1081108-Cordyceps confragosa RCEF-Sordariomycetes
XP 006670394.1-983644-Cordyceps militaris CM01-Sordariomycetes
OAA52451.1-1081107-Cordyceps brongniartii RC-Sordariomycetes
XP 013425836.1-1043004-Aureobasidium namibiae CB-Dothideomycetes
KEQ67761.1-1043003-Aureobasidium melanogenum-Dothideomycetes
XP 013347900.1-1043005-Aureobasidium subglaciale-Dothideomycetes
KEQ89864.1-1043002-Aureobasidium pullulans E-Dothideomycetes
XP 020127555.1-236234-Diplodia corticola-Dothideomycetes
OMP87711.1-420778-Diplodia seriata-Dothideomycetes
KKY19282.1-420778-Diplodia seriata-Dothideomycetes
XP 007582513.1-1287680-Neofusicoccum parvum UCRN-Dothideomycetes
EKG15688.1-1126212-Macrophomina phaseolina M-Dothideomycetes
GAW20816.1-1813822-fungal sp No 14919-other Fungi
XP 007789192.1-1287681-Eutypa lata UCREL1-Sordariomycetes
OTB04225.1-1001833-Hypoxylon sp CI 4A-Sordariomycetes
OTA91720.1-1001938-Hypoxylon sp CO27 5-Sordariomycetes
OTA57144.1-1001937-Hypoxylon sp EC38-Sordariomycetes
XP 013950960.1-413071-Trichoderma virens Gv29 8-Sordariomycetes
OPB39282.1-1491466-Trichoderma guizhouense-Sordariomycetes
XP 018658687.1-398673-Trichoderma gamsii-Sordariomycetes
XP 013944581.1-452589-Trichoderma atroviride IM-Sordariomycetes
OTA03574.1-858221-Trichoderma parareesei-Sordariomycetes
ETS05692.1-1344414-Trichoderma reesei RUT C-Sordariomycetes
XP 006963461.1-431241-Trichoderma reesei QM6a-Sordariomycetes
OAL52377.1-765867-Pyrenochaeta sp DS3sAY3a-Dothideomycetes
XP 018386305.1-5599-Alternaria alternata-Dothideomycetes
XP 008020889.1-671987-Setosphaeria turcica Et28-Dothideomycetes
KNG44838.1-183478-Stemphylium lycopersici-Dothideomycetes
XP 007710543.1-930089-Bipolaris zeicola 26 R 13-Dothideomycetes
XP 014557659.1-930091-Bipolaris victoriae FI3-Dothideomycetes
XP 007695836.1-665912-Bipolaris sorokiniana ND9-Dothideomycetes
XP 007688430.1-930090-Bipolaris oryzae ATCC 445-Dothideomycetes
KHJ30190.1-52586-Erysiphe necator-Leotiomycetes
CCU74944.1-546991-Blumeria graminis f sp-Leotiomycetes
EPQ61501.1-1268274-Blumeria graminis f sp-Leotiomycetes
ESZ92078.1-1432307-Sclerotinia borealis F 41-Leotiomycetes
CCD49728.1-999810-Botrytis cinerea T4-Leotiomycetes
EMR86983.1-1290391-Botrytis cinerea BcDW1-Leotiomycetes
CZT04403.1-914238-Rhynchosporium agropyri-Leotiomycetes
CZT06225.1-914237-Rhynchosporium commune-Leotiomycetes
CZT44003.1-38038-Rhynchosporium secalis-Leotiomycetes
KFY73687.1-1420912-Pseudogymnoascus sp VKM-Leotiomycetes
XP 018131908.1-342668-Pseudogymnoascus verrucos-Leotiomycetes
OBT57271.1-1622150-Pseudogymnoascus sp 24MN-Leotiomycetes
KFZ08350.1-1420914-Pseudogymnoascus sp VKM-Leotiomycetes
OBT40144.1-1622147-Pseudogymnoascus sp WSF-Leotiomycetes
OBT85056.1-1622148-Pseudogymnoascus sp 03VT-Leotiomycetes
OBT79024.1-1622149-Pseudogymnoascus sp 05NY-Leotiomycetes
KFY20101.1-1420901-Pseudogymnoascus sp VKM-Leotiomycetes
KFY26564.1-1420906-Pseudogymnoascus sp VKM-Leotiomycetes
KFX89420.1-1437433-Pseudogymnoascus sp VKM-Leotiomycetes
KFY07860.1-1420902-Pseudogymnoascus sp VKM-Leotiomycetes
KFY29642.1-1420907-Pseudogymnoascus sp VKM-Leotiomycetes
OBT62886.1-1524831-Pseudogymnoascus sp 2334-Leotiomycetes
KFZ11761.1-1420915-Pseudogymnoascus sp VKM-Leotiomycetes
KFZ01402.1-1420913-Pseudogymnoascus sp VKM-Leotiomycetes

S2D

Tree scale: 0.1 ⊢——⊣

Taxonomy

- Leotiomycetes
- Sordariomycetes
- Eurotiomycetes
- Dothideomycetes
- other_Pezizomycotina
- other_Ascomycota
- other_Fungi
- other_Opisthokonta
- other_Eukaryota
- Bacteria
- Archaea
- Viruses

XP 011319440.1-229533-Fusarium graminearum PH 1-Sordariomycetes
OJJ34620.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
KZL72754.1-708197-Colletotrichum tofieldiae-Sordariomycetes
XP 007603120.1-1445577-Colletotrichum fioriniae-Sordariomycetes
KXH26577.1-703756-Colletotrichum simmondsii-Sordariomycetes
ORY67812.1-1141098-Pseudomassariella vexata-Sordariomycetes
XP 001270374.1-344612-Aspergillus clavatus NRRL-Eurotiomycetes
CRL30864.1-5075-Penicillium camemberti-Eurotiomycetes
OQD93951.1-60172-Penicillium solitum-Eurotiomycetes
KOS43411.1-229535-Penicillium nordicum-Eurotiomycetes
EHA18933.1-380704-Aspergillus niger ATCC 10-Eurotiomycetes
GAT21369.1-1069201-Aspergillus luchuensis-Eurotiomycetes
GAA83288.1-1033177-Aspergillus kawachii IFO-Eurotiomycetes
KEY83926.1-1476192-Aspergillus fumigatus var-Eurotiomycetes
QUERY-Afu1g17725-330879-Aspergillus fumigatus Af2-Eurotiomycetes
KMK54952.1-1437362-Aspergillus fumigatus Z5-Eurotiomycetes
GAD97588.1-1356009-Byssochlamys spectabilis-Eurotiomycetes
KIV78800.1-1016849-Exophiala sideris-Eurotiomycetes
XP 013324239.1-1408163-Rasamsonia emersonii CBS-Eurotiomycetes
XP 013327701.1-1408163-Rasamsonia emersonii CBS-Eurotiomycetes
XP 007787724.1-1263415-Endocarpon pusillum Z0702-Eurotiomycetes
XP 018188082.1-1328760-Xylona heveae TC161-other Pezizomycotina
SLM39940.1-136370-Umbilicaria pustulata-other Pezizomycotina
XP 006962774.1-431241-Trichoderma reesei QM6a-Sordariomycetes
EKG11783.1-1126212-Macrophomina phaseolina M-Dothideomycetes
OMP88640.1-420778-Diplodia seriata-Dothideomycetes
XP 013270916.1-1442369-Rhinocladiella mackenziei-Eurotiomycetes
XP 016227003.1-212818-Exophiala mesophila-Eurotiomycetes
XP 016632485.1-1442371-Fonsecaea multimorphosa C-Eurotiomycetes
XP 018695363.1-1367422-Fonsecaea erecta-Eurotiomycetes
XP 016250130.1-569365-Cladophialophora immunda-Eurotiomycetes
OQV00536.1-569365-Cladophialophora immunda-Eurotiomycetes
XP 007739659.1-1182543-Cladophialophora psammoph-Eurotiomycetes
XP 016625804.1-1442370-Cladophialophora bantiana-Eurotiomycetes
OAL38606.1-856822-Fonsecaea nubica-Eurotiomycetes
XP 013284061.1-1442368-Fonsecaea pedrosoi CBS 27-Eurotiomycetes
OAG44669.1-254056-Fonsecaea monophora-Eurotiomycetes
OJJ30508.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
ODM14997.1-573508-Aspergillus cristatus-Eurotiomycetes
OJJ85962.1-1160497-Aspergillus glaucus CBS 5-Eurotiomycetes
EYE95192.1-1388766-Aspergillus ruber CBS 135-Eurotiomycetes
KKK12315.1-308745-Aspergillus rambellii-Eurotiomycetes
CEL05234.1-454130-Aspergillus calidoustus-Eurotiomycetes
XP 659488.1-227321-Aspergillus nidulans FGSC-Eurotiomycetes
OJJ64297.1-1036612-Aspergillus sydowii CBS 5-Eurotiomycetes
OJI98390.1-1036611-Aspergillus versicolor CB-Eurotiomycetes
XP 001215107.1-341663-Aspergillus terreus NIH26-Eurotiomycetes
XP 015404014.1-1509407-Aspergillus nomius NRRL 1-Eurotiomycetes
OGM50557.1-109264-Aspergillus bombycis-Eurotiomycetes
KJK64619.1-1403190-Aspergillus parasiticus S-Eurotiomycetes
KOC13210.1-1392242-Aspergillus flavus AF70-Eurotiomycetes
BAE57429.1-510516-Aspergillus oryzae RIB40-Eurotiomycetes
XP 020058705.1-690307-Aspergillus aculeatus ATC-Eurotiomycetes
OOF99045.1-602072-Aspergillus carbonarius I-Eurotiomycetes
OJJ77694.1-767769-Aspergillus brasiliensis-Eurotiomycetes
EHA19494.1-380704-Aspergillus niger ATCC 10-Eurotiomycetes
XP 001394301.2-425011-Aspergillus niger CBS 513-Eurotiomycetes
GAA83936.1-1033177-Aspergillus kawachii IFO-Eurotiomycetes
OJZ90757.1-1137211-Aspergillus luchuensis CB-Eurotiomycetes
XP 001272758.1-344612-Aspergillus clavatus NRRL-Eurotiomycetes
GAQ07125.1-293939-Aspergillus lentulus-Eurotiomycetes
XP 001260060.1-331117-Aspergillus fischeri NRRL-Eurotiomycetes
GAO90844.1-91492-Aspergillus udagawae-Eurotiomycetes
KEY79284.1-1476192-Aspergillus fumigatus var-Eurotiomycetes
XP 749561.1-330879-Aspergillus fumigatus Af2-Eurotiomycetes
EDP54085.1-451804-Aspergillus fumigatus A11-Eurotiomycetes
OGE49342.1-1835702-Penicillium arizonense-Eurotiomycetes
OQD80997.1-416450-Penicillium antarcticum-Eurotiomycetes
OQE08078.1-29845-Penicillium vulpinum-Eurotiomycetes
OQE47134.1-36646-Penicillium coprophilum-Eurotiomycetes
CDM35225.1-1365484-Penicillium roqueforti FM-Eurotiomycetes
OQE29031.1-254877-Penicillium flavigenum-Eurotiomycetes
OQE88303.1-60175-Penicillium nalgiovense-Eurotiomycetes
XP 002557726.1-500485-Penicillium rubens Wiscon-Eurotiomycetes
KZN85110.1-5076-Penicillium chrysogenum-Eurotiomycetes
KGO69212.1-40296-Penicillium italicum-Eurotiomycetes
XP 016600192.1-27334-Penicillium expansum-Eurotiomycetes
KUM56968.1-48697-Penicillium freii-Eurotiomycetes
OQD61678.1-60169-Penicillium polonicum-Eurotiomycetes
OQD87913.1-60172-Penicillium solitum-Eurotiomycetes
KOS46330.1-229535-Penicillium nordicum-Eurotiomycetes
OQD59998.1-60169-Penicillium polonicum-Eurotiomycetes
OQE15639.1-303698-Penicillium steckii-Eurotiomycetes
OQD74193.1-69771-Penicillium decumbens-Eurotiomycetes
EPS33859.1-933388-Penicillium oxalicum 114-Eurotiomycetes
OKO98199.1-1316194-Penicillium subrubescens-Eurotiomycetes
CEO60574.1-104259-Penicillium brasilianum-Eurotiomycetes
OOQ85244.1-104259-Penicillium brasilianum-Eurotiomycetes
XP 003021658.1-663202-Trichophyton verrucosum H-Eurotiomycetes
XP 003175950.1-535722-Nannizzia gypsea CBS 1188-Eurotiomycetes
XP 001275058.1-344612-Aspergillus clavatus NRRL-Eurotiomycetes
XP 015406159.1-1509407-Aspergillus nomius NRRL 1-Eurotiomycetes
GAQ09758.1-293939-Aspergillus lentulus-Eurotiomycetes
GAO90499.1-91492-Aspergillus udagawae-Eurotiomycetes
XP 001262985.1-331117-Aspergillus fischeri NRRL-Eurotiomycetes
EDP52282.1-451804-Aspergillus fumigatus A11-Eurotiomycetes
XP 754149.1-330879-Aspergillus fumigatus Af2-Eurotiomycetes
KMK54265.1-1437362-Aspergillus fumigatus Z5-Eurotiomycetes
KEY83423.1-1476192-Aspergillus fumigatus var-Eurotiomycetes

S2E

S2F

Tree scale: 0.1 ⊢——⊣

GAA82870.1-1033177-Aspergillus kawachii IFO-Eurotiomycetes
OOF98050.1-602072-Aspergillus carbonarius I-Eurotiomycetes
KXX75968.1-100816-Madurella mycetomatis-Sordariomycetes
KFY16432.1-1420902-Pseudogymnoascus sp VKM-Leotiomycetes
OGM43967.1-109264-Aspergillus bombycis-Eurotiomycetes
KFZ13812.1-1420914-Pseudogymnoascus sp VKM-Leotiomycetes
AFP89391.1-1116209-Cladosporium phlei-Dothideomycetes
XP 001274946.1-344612-Aspergillus clavatus NRRL-Eurotiomycetes
OHE97136.1-1209926-Colletotrichum orchidophi-Sordariomycetes
XP 008100501.1-645133-Colletotrichum graminicol-Sordariomycetes
ENH82918.1-1213857-Colletotrichum orbiculare-Sordariomycetes
KXH63779.1-1209931-Colletotrichum salicis-Sordariomycetes
XP 001394625.2-425011-Aspergillus niger CBS 513-Eurotiomycetes
XP 015405462.1-1509407-Aspergillus nomius NRRL 1-Eurotiomycetes
XP 020054826.1-690307-Aspergillus aculeatus ATC-Eurotiomycetes
OPB36213.1-1491466-Trichoderma guizhouense-Sordariomycetes
KFY96473.1-1420913-Pseudogymnoascus sp VKM-Leotiomycetes
OTB16775.1-1001832-Daldinia sp EC12-Sordariomycetes
XP 001258701.1-331117-Aspergillus fischeri NRRL-Eurotiomycetes
EKG15173.1-1126212-Macrophomina phaseolina M-Dothideomycetes
OBS22513.1-36050-Fusarium poae-Sordariomycetes
ALQ32891.1-36050-Fusarium poae-Sordariomycetes
XP 003177417.1-535722-Nannizzia gypsea CBS 1188-Eurotiomycetes
AFN68293.1-5599-Alternaria alternata-Dothideomycetes
XP 018384350.1-5599-Alternaria alternata-Dothideomycetes
BAR40283.1-1608308-Fusarium sp FN080326-Sordariomycetes
AGO86662.1-42747-Fusarium heterosporum-Sordariomycetes
ALQ32974.1-70751-Fusarium sp NRRL 25184-Sordariomycetes
KLO99472.1-5127-Fusarium fujikuroi-Sordariomycetes
ALQ33011.1-42677-Fusarium subglutinans-Sordariomycetes
ALQ32847.1-246455-Fusarium foetens-Sordariomycetes
XP 018760578.1-334819-Fusarium verticillioides-Sordariomycetes
EWZ45929.1-660027-Fusarium oxysporum Fo47-Sordariomycetes
EWY99916.1-909455-Fusarium oxysporum FOSC 3-Sordariomycetes
EXA51210.1-1080344-Fusarium oxysporum f sp-Sordariomycetes
EXK85226.1-1089458-Fusarium oxysporum f sp-Sordariomycetes
EGU88865.1-660025-Fusarium oxysporum Fo5176-Sordariomycetes
KKY30404.1-1214573-Diaporthe ampelina-Sordariomycetes
KUI65083.1-105487-Valsa mali-Sordariomycetes
KUI61972.1-694573-Valsa mali var pyri-Sordariomycetes
KEQ86259.1-1043002-Aureobasidium pullulans E-Dothideomycetes
XP 013343243.1-1043005-Aureobasidium subglaciale-Dothideomycetes
ORY67416.1-1141098-Pseudomassariella vexata-Sordariomycetes
OTA59680.1-1001937-Hypoxylon sp EC38-Sordariomycetes
OTA90454.1-1001938-Hypoxylon sp CO27 5-Sordariomycetes
KFY84670.1-1420913-Pseudogymnoascus sp VKM-Leotiomycetes
KFZ03054.1-1420915-Pseudogymnoascus sp VKM-Leotiomycetes
XP 018134601.1-342668-Pseudogymnoascus verrucos-Leotiomycetes
KFY76989.1-1420912-Pseudogymnoascus sp VKM-Leotiomycetes
OBT61811.1-1524831-Pseudogymnoascus sp 2334-Leotiomycetes
KFY49789.1-1420909-Pseudogymnoascus sp VKM-Leotiomycetes
KFY92340.1-1420911-Pseudogymnoascus sp VKM-Leotiomycetes
KMM67662.1-454284-Coccidioides posadasii RM-Eurotiomycetes
XP 020057498.1-690307-Aspergillus aculeatus ATC-Eurotiomycetes
EPS29069.1-933388-Penicillium oxalicum 114-Eurotiomycetes
ARF05979.1-69781-Penicillium oxalicum-Eurotiomycetes
OTA98397.1-1001833-Hypoxylon sp CI 4A-Sordariomycetes
SLM36899.1-136370-Umbilicaria pustulata-other Pezizomycotina
ORY61265.1-1141098-Pseudomassariella vexata-Sordariomycetes
KXX75016.1-100816-Madurella mycetomatis-Sordariomycetes
XP 007911967.1-1286976-Phaeoacremonium minimum U-Sordariomycetes
AFI23577.1-80884-Colletotrichum higginsian-Sordariomycetes
KFY06571.1-1420902-Pseudogymnoascus sp VKM-Leotiomycetes
KFY34117.1-1420907-Pseudogymnoascus sp VKM-Leotiomycetes
KFX98753.1-1437433-Pseudogymnoascus sp VKM-Leotiomycetes
KFY37827.1-1420908-Pseudogymnoascus sp VKM-Leotiomycetes
KFY92181.1-1420913-Pseudogymnoascus sp VKM-Leotiomycetes
OBT43992.1-1622147-Pseudogymnoascus sp WSF-Leotiomycetes
AKL78828.1-101852-Glarea lozoyensis-Leotiomycetes
XP 018156684.1-759273-Colletotrichum higginsian-Sordariomycetes
XP 007827066.1-1229662-Pestalotiopsis fici W106-Sordariomycetes
KID84093.1-1276136-Metarhizium guizhouense A-Sordariomycetes
XP 014573816.1-1276143-Metarhizium majus ARSEF 2-Sordariomycetes
XP 014540252.1-1276141-Metarhizium brunneum ARSE-Sordariomycetes
EXU96139.1-568076-Metarhizium robertsii ARS-Sordariomycetes
XP 007817535.1-655844-Metarhizium robertsii ARS-Sordariomycetes
KFG81513.1-5530-Metarhizium anisopliae-Sordariomycetes
KJK74647.1-1291518-Metarhizium anisopliae BR-Sordariomycetes
KJK88403.1-1291519-Metarhizium anisopliae BR-Sordariomycetes
XP 009262583.1-1028729-Fusarium pseudograminearu-Sordariomycetes
XP 011319443.1-229533-Fusarium graminearum PH 1-Sordariomycetes
EYB31408.1-5518-Fusarium graminearum-Sordariomycetes
OJJ34622.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
XP 007919595.1-1286976-Phaeoacremonium minimum U-Sordariomycetes
ORY67814.1-1141098-Pseudomassariella vexata-Sordariomycetes
KZL72765.1-708197-Colletotrichum tofieldiae-Sordariomycetes
KXH26578.1-703756-Colletotrichum simmondsii-Sordariomycetes
XP 007603119.1-1445577-Colletotrichum fioriniae-Sordariomycetes
XP 001270376.1-344612-Aspergillus clavatus NRRL-Eurotiomycetes
KOS43405.1-229535-Penicillium nordicum-Eurotiomycetes
OQD93852.1-60172-Penicillium solitum-Eurotiomycetes
CRL31206.1-5075-Penicillium camemberti-Eurotiomycetes
GAA83290.1-1033177-Aspergillus kawachii IFO-Eurotiomycetes
EHA18935.1-380704-Aspergillus niger ATCC 10-Eurotiomycetes
GAA83291.1-1033177-Aspergillus kawachii IFO-Eurotiomycetes
KEY83630.1-1476192-Aspergillus fumigatus var-Eurotiomycetes
KEY83624.1-1476192-Aspergillus fumigatus var-Eurotiomycetes
QUERY-Afu1g17740-330879-Aspergillus fumigatus Af2-Eurotiomycetes
KMK54950.1-1437362-Aspergillus fumigatus Z5-Eurotiomycetes

Figure S3. Novel SM gene clusters in A. fumigatus strains. (A) Synteny between a novel PKS containing cluster in two strains with an SM gene cluster in *Metarhizium anisopliae*. This novel PKS cluster is located between transposable elements in a region syntenic with the reference Af293 chromosome 4. (B) Novel SM gene cluster in MO54056EXP and 3 additional strains. This cluster is only located on one scaffold in MO540556EXP and is fragmented across the other strains (ends of scaffolds are marked). (C) Coverage data from short read alignments for MO54056EXP, 12-7504462, and 08-19-02-30 relative to the MO54056EXP scaffold containing the novel SM gene cluster.(D) Strains with novel clusters on SNP phylogeny.

Figure S4. Gene phylogenies of SM gene cluster 24. The phylogenies of several genes in this cluster are consistent with horizontal transfer between *Aspergillus fumigatus* and *Metarhizium* fungi.

S4A

S4C

Chr 4 — MFS transporter — Gypsy transposon — CYP450 — PKS — Transporter — Protein kinase — O-methyltransferase — Aldehyde reductase — Amidohydrolase — CYP450 — Mariner transposon — Gypsy transposon — Chr 4

Afu4g01560   AFUB_102040

Tree scale: 0.1

Taxonomy

- Leotiomycetes
- Sordariomycetes
- Eurotiomycetes
- Dothideomycetes
- other_Pezizomycotina
- other_Ascomycota
- other_Fungi
- other_Opisthokonta
- other_Eukaryota
- Bacteria
- Archaea
- Viruses

EHA21874.1-380704-Aspergillus niger ATCC 10-Eurotiomycetes
ODM24152.1-573508-Aspergillus cristatus-Eurotiomycetes
XP 749839.2-330879-Aspergillus fumigatus Af2-Eurotiomycetes
EHA21875.1-380704-Aspergillus niger ATCC 10-Eurotiomycetes
GAQ02781.1-293939-Aspergillus lentulus-Eurotiomycetes
OJJ52319.1-1036612-Aspergillus sydowii CBS 5-Eurotiomycetes
KKK18129.1-308745-Aspergillus rambellii-Eurotiomycetes
KGO71756.1-40296-Penicillium italicum-Eurotiomycetes
OOF90907.1-602072-Aspergillus carbonarius I-Eurotiomycetes
XP 001271128.1-344612-Aspergillus clavatus NRRL-Eurotiomycetes
OJJ65769.1-767769-Aspergillus brasiliensis-Eurotiomycetes
CDM37172.1-1365484-Penicillium roqueforti FM-Eurotiomycetes
OOO03860.1-5062-Aspergillus oryzae-Eurotiomycetes
BAE66189.1-510516-Aspergillus oryzae RIB40-Eurotiomycetes
KOC17684.1-1392242-Aspergillus flavus AF70-Eurotiomycetes
BAE66187.1-510516-Aspergillus oryzae RIB40-Eurotiomycetes
XP 001827320.2-510516-Aspergillus oryzae RIB40-Eurotiomycetes
QUERY-g6970-746128-Aspergillus fumigatus F13619-Eurotiomycetes
XP 001263991.1-331117-Aspergillus fischeri NRRL-Eurotiomycetes
XP 001258122.1-331117-Aspergillus fischeri NRRL-Eurotiomycetes
XP 749859.1-330879-Aspergillus fumigatus Af2-Eurotiomycetes
XP 749857.1-330879-Aspergillus fumigatus Af2-Eurotiomycetes
KMK54477.1-1437362-Aspergillus fumigatus Z5-Eurotiomycetes

S4D

Tree scale: 0.1 ⊢—⊣

Taxonomy

- Leotiomycetes
- Sordariomycetes
- Eurotiomycetes
- Dothideomycetes
- other_Pezizomycotina
- other_Ascomycota
- other_Fungi
- other_Opisthokonta
- other_Eukaryota
- Bacteria
- Archaea
- Viruses

S4E

Tree scale: 0.1 ⊢──┤

Taxonomy

- ■ Leotiomycetes
- ■ Sordariomycetes
- ■ Eurotiomycetes
- ■ Dothideomycetes
- ■ other_Pezizomycotina
- ■ other_Ascomycota
- ■ other_Fungi
- ■ other_Opisthokonta
- ■ other_Eukaryota
- ■ Bacteria
- ■ Archaea
- ■ Viruses

S4F

Tree scale: 0.1

S4G

Tree scale: 0.1 ⊢———⊣

Taxonomy

- 🟥 Leotiomycetes
- 🟪 Sordariomycetes
- 🟧 Eurotiomycetes
- 🟧 Dothideomycetes
- 🟫 other_Pezizomycotina
- 🟪 other_Ascomycota
- 🟪 other_Fungi
- 🟦 other_Opisthokonta
- 🟦 other_Eukaryota
- 🟩 Bacteria
- 🟨 Archaea
- ⬜ Viruses

XP 018184752.1-1328760-Xylona heveae TC161-other Pezizomycotina
KFY21354.1-1420906-Pseudogymnoascus sp VKM-Leotiomycetes
XP 002841013.1-656061-Tuber melanosporum Mel28-other Pezizomycotina
QUERY-g6974-746128-Aspergillus fumigatus F13619-Eurotiomycetes
KJK74880.1-1291518-Metarhizium anisopliae BR-Sordariomycetes
GAT55036.1-658473-Mycena chlorophos-Basidiomycota
KTB38372.1-221103-Moniliophthora roreri-Basidiomycota
XP 007845553.1-1381753-Moniliophthora roreri MCA-Basidiomycota
AAF64435.2-40559-Botrytis cinerea-Leotiomycetes
ESZ95339.1-1432307-Sclerotinia borealis F 41-Leotiomycetes
XP 001586264.1-665079-Sclerotinia sclerotiorum-Leotiomycetes
KIN05050.1-913774-Oidiodendron maius Zn-Leotiomycetes
OGE55021.1-1835702-Penicillium arizonense-Eurotiomycetes
KFY94063.1-1420913-Pseudogymnoascus sp VKM-Leotiomycetes
EHL03343.1-1104152-Glarea lozoyensis 74030-Leotiomycetes
XP 008083356.1-1116229-Glarea lozoyensis ATCC 20-Leotiomycetes
CZR50636.1-576137-Phialocephala subalpina-Leotiomycetes
XP 018076654.1-149040-Phialocephala scopiformis-Leotiomycetes
XP 007292794.1-1072389-Marssonina brunnea f sp-Leotiomycetes
CZT47972.1-38038-Rhynchosporium secalis-Leotiomycetes
CZS94684.1-914237-Rhynchosporium commune-Leotiomycetes
XP 007915264.1-1286976-Phaeoacremonium minimum U-Sordariomycetes
OIW34274.1-1408157-Coniochaeta ligniaria NRR-Sordariomycetes
XP 009852902.1-510951-Neurospora tetrasperma FG-Sordariomycetes
XP 011395298.1-367110-Neurospora crassa OR74A-Sordariomycetes
OQE41467.1-36646-Penicillium coprophilum-Eurotiomycetes
OQD89879.1-416450-Penicillium antarcticum-Eurotiomycetes
OGE48087.1-1835702-Penicillium arizonense-Eurotiomycetes
GAM38946.1-1472165-Talaromyces cellulolyticu-Eurotiomycetes
XP 002479167.1-441959-Talaromyces stipitatus AT-Eurotiomycetes
XP 002144369.1-441960-Talaromyces marneffei ATC-Eurotiomycetes
KFX53693.1-1077442-Talaromyces marneffei PM1-Eurotiomycetes
ORY66156.1-1141098-Pseudomassariella vexata-Sordariomycetes
AEX31648.1-1491472-Trichoderma lixii-Sordariomycetes
XP 008031645.1-671987-Setosphaeria turcica Et28-Dothideomycetes
OTA93836.1-1001938-Hypoxylon sp CO27 5-Sordariomycetes
OTA69676.1-1001937-Hypoxylon sp EC38-Sordariomycetes
GAP91503.1-77044-Rosellinia necatrix-Sordariomycetes
ABR28367.1-167374-Xylaria sp BCC 1067-Sordariomycetes
GAW14624.1-1813822-fungal sp No 14919-other Fungi
XP 007780309.1-1168221-Coniosporium apollinis CB-Dothideomycetes
EWC44372.1-1043628-Drechslerella stenobrocha-other Pezizomycotina
XP 013344082.1-1043005-Aureobasidium subglaciale-Dothideomycetes
XP 013432457.1-1043004-Aureobasidium namibiae CB-Dothideomycetes
KEQ88423.1-1043002-Aureobasidium pullulans E-Dothideomycetes
XP 013325594.1-1408163-Rasamsonia emersonii CBS-Eurotiomycetes
XP 007581427.1-1287680-Neofusicoccum parvum UCRN-Dothideomycetes
EKG14558.1-1126212-Macrophomina phaseolina M-Dothideomycetes
OQD94582.1-60172-Penicillium solitum-Eurotiomycetes
OKP14090.1-1316194-Penicillium subrubescens-Eurotiomycetes
CEJ59761.1-104259-Penicillium brasilianum-Eurotiomycetes
OOQ82870.1-104259-Penicillium brasilianum-Eurotiomycetes
OCK75213.1-1314670-Lepidopterella palustris-Dothideomycetes
OCK98761.1-794803-Cenococcum geophilum 1 58-Dothideomycetes
OCL13154.1-574774-Glonium stellatum-Dothideomycetes
OAL53265.1-765867-Pyrenochaeta sp DS3sAY3a-Dothideomycetes
OAL06695.1-765868-Stagonospora sp SRC1lsM3-Dothideomycetes
XP 001794321.1-321614-Parastagonospora nodorum-Dothideomycetes
KNG49244.1-183478-Stemphylium lycopersici-Dothideomycetes
EME39528.1-675120-Dothistroma septosporum N-Dothideomycetes
XP 007928999.1-383855-Pseudocercospora fijiensi-Dothideomycetes
XP 016758133.1-692275-Sphaerulina musiva SO2202-Dothideomycetes
KJX98136.1-1047168-Zymoseptoria brevis-Dothideomycetes
XP 003850560.1-336722-Zymoseptoria tritici IPO3-Dothideomycetes
AJG42821.1-1047171-Zymoseptoria tritici-Dothideomycetes
OTA35326.1-1157616-Hortaea werneckii EXF 200-Dothideomycetes
KXL50548.1-245562-Acidomyces richmondensis-Dothideomycetes
XP 007681432.1-717646-Baudoinia panamericana UA-Dothideomycetes
ODQ69136.1-675824-Lipomyces starkeyi NRRL Y-Saccharomycotina
XP 002797035.1-502779-Paracoccidioides lutzii P-Eurotiomycetes
EGE77149.2-653446-Blastomyces dermatitidis-Eurotiomycetes
KLJ08977.1-1246674-Emmonsia parva UAMH 139-Eurotiomycetes
XP 001243393.1-246410-Coccidioides immitis RS-Eurotiomycetes
KMU88747.1-396776-Coccidioides immitis H538-Eurotiomycetes
KMM68322.1-454284-Coccidioides posadasii RM-Eurotiomycetes
XP 003066894.1-222929-Coccidioides posadasii C7-Eurotiomycetes
XP 002843435.1-554155-Arthroderma otae CBS 1134-Eurotiomycetes
XP 003172129.1-535722-Nannizzia gypsea CBS 1188-Eurotiomycetes
EGD96741.1-647933-Trichophyton tonsurans CB-Eurotiomycetes
XP 003012776.1-663331-Trichophyton benhamiae CB-Eurotiomycetes
XP 003236897.1-559305-Trichophyton rubrum CBS 1-Eurotiomycetes
XP 003023086.1-663202-Trichophyton verrucosum H-Eurotiomycetes
XP 007787617.1-1263415-Endocarpon pusillum Z0702-Eurotiomycetes
XP 013255212.1-1182545-Exophiala aquamarina CBS-Eurotiomycetes
XP 013267468.1-1442369-Rhinocladiella mackenziei-Eurotiomycetes
XP 007720660.1-1182541-Capronia coronata CBS 617-Eurotiomycetes
XP 007736804.1-1182542-Capronia epimyces CBS 606-Eurotiomycetes
XP 009152324.1-858893-Exophiala dermatitidis NI-Eurotiomycetes
XP 016237739.1-91928-Exophiala spinifera-Eurotiomycetes
XP 016262934.1-215243-Exophiala oligosperma-Eurotiomycetes
OCT46042.1-86049-Cladophialophora carrioni-Eurotiomycetes
XP 008724188.1-1279043-Cladophialophora carrioni-Eurotiomycetes
XP 018689397.1-1367422-Fonsecaea erecta-Eurotiomycetes
XP 016631957.1-1442371-Fonsecaea multimorphosa C-Eurotiomycetes
XP 016246567.1-569365-Cladophialophora immunda-Eurotiomycetes
OQV06046.1-569365-Cladophialophora immunda-Eurotiomycetes
XP 013286744.1-1442368-Fonsecaea pedrosoi CBS 27-Eurotiomycetes
OAL34481.1-856822-Fonsecaea nubica-Eurotiomycetes
XP 007751748.1-1182543-Cladophialophora psammoph-Eurotiomycetes
XP 016621099.1-1442370-Cladophialophora bantiana-Eurotiomycetes

S4H

XP 018138618.1-1380566-Pochonia chlamydosporia 1-Sordariomycetes
GAQ08559.1-293939-Aspergillus lentulus-Eurotiomycetes
XP 002482893.1-441959-Talaromyces stipitatus AT-Eurotiomycetes
XP 018138617.1-1380566-Pochonia chlamydosporia 1-Sordariomycetes
ORY69640.1-1141098-Pseudomassariella vexata-Sordariomycetes
XP 001217075.1-341663-Aspergillus terreus NIH26-Eurotiomycetes
XP 018067965.1-149040-Phialocephala scopiformis-Leotiomycetes
XP 018066031.1-149040-Phialocephala scopiformis-Leotiomycetes
CRG86679.1-28573-Talaromyces islandicus-Eurotiomycetes
OJJ40786.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
OJJ77434.1-767769-Aspergillus brasiliensis-Eurotiomycetes
KKK25665.1-308745-Aspergillus rambellii-Eurotiomycetes
XP 007731038.1-1182542-Capronia epimyces CBS 606-Eurotiomycetes
XP 008020155.1-671987-Setosphaeria turcica Et28-Dothideomycetes
CRG90999.1-28573-Talaromyces islandicus-Eurotiomycetes
XP 018188015.1-1328760-Xylona heveae TC161-other Pezizomycotina
XP 008026822.1-671987-Setosphaeria turcica Et28-Dothideomycetes
XP 002485357.1-441959-Talaromyces stipitatus AT-Eurotiomycetes
XP 001208364.1-341663-Aspergillus terreus NIH26-Eurotiomycetes
XP 020115508.1-1441469-Talaromyces atroroseus-Eurotiomycetes
ESZ98975.1-1432307-Sclerotinia borealis F 41-Leotiomycetes
OCL10940.1-574774-Glonium stellatum-Dothideomycetes
ORX95400.1-1231657-Clohesyomyces aquaticus-Dothideomycetes
KFZ05412.1-1420914-Pseudogymnoascus sp VKM-Leotiomycetes
KFY75262.1-1420912-Pseudogymnoascus sp VKM-Leotiomycetes
OCK83279.1-1314670-Lepidopterella palustris-Dothideomycetes
XP 007685917.1-930090-Bipolaris oryzae ATCC 445-Dothideomycetes
XP 006669619.1-983644-Cordyceps militaris CM01-Sordariomycetes
OAQ99551.1-1105325-Cordyceps confragosa-Sordariomycetes
XP 001258910.1-331117-Aspergillus fischeri NRRL-Eurotiomycetes
XP 018702657.1-1081104-Isaria fumosorosea ARSEF-Sordariomycetes
QUERY-g6975-746128-Aspergillus fumigatus F13619-Eurotiomycetes
KJK74888.1-1291518-Metarhizium anisopliae BR-Sordariomycetes
XP 002478259.1-441959-Talaromyces stipitatus AT-Eurotiomycetes
OJJ08318.1-1036611-Aspergillus versicolor CB-Eurotiomycetes
OJJ53427.1-1036612-Aspergillus sydowii CBS 5-Eurotiomycetes
KFA69539.1-1283841-Stachybotrys chlorohalona-Sordariomycetes
KFA74443.1-1283842-Stachybotrys chartarum IB-Sordariomycetes
KEY72230.1-1280523-Stachybotrys chartarum IB-Sordariomycetes
KFY13626.1-1420901-Pseudogymnoascus sp VKM-Leotiomycetes
KFY50143.1-1420909-Pseudogymnoascus sp VKM-Leotiomycetes
KFY81276.1-1420913-Pseudogymnoascus sp VKM-Leotiomycetes
KFZ03029.1-1420915-Pseudogymnoascus sp VKM-Leotiomycetes
XP 003836151.1-985895-Leptosphaeria maculans JN-Dothideomycetes
OTA33688.1-1157616-Hortaea werneckii EXF 200-Dothideomycetes
ELQ40620.1-1143189-Magnaporthe oryzae Y34-Sordariomycetes
XP 003709964.1-242507-Magnaporthe oryzae 70 15-Sordariomycetes
XP 020127315.1-236234-Diplodia corticola-Dothideomycetes
XP 007696884.1-665912-Bipolaris sorokiniana ND9-Dothideomycetes
XP 014081844.1-665024-Bipolaris maydis ATCC 483-Dothideomycetes
XP 018387357.1-5599-Alternaria alternata-Dothideomycetes
XP 014558804.1-930091-Bipolaris victoriae FI3-Dothideomycetes
XP 007717426.1-930089-Bipolaris zeicola 26 R 13-Dothideomycetes
KFX98711.1-1437433-Pseudogymnoascus sp VKM-Leotiomycetes
KFY00079.1-1391699-Pseudogymnoascus sp VKM-Leotiomycetes
GAO90615.1-91492-Aspergillus udagawae-Eurotiomycetes
XP 003173259.1-535722-Nannizzia gypsea CBS 1188-Eurotiomycetes
XP 002846809.1-554155-Arthroderma otae CBS 1134-Eurotiomycetes
OTB10323.1-1001832-Daldinia sp EC12-Sordariomycetes
OQE43316.1-36646-Penicillium coprophilum-Eurotiomycetes
OQD66534.1-60169-Penicillium polonicum-Eurotiomycetes
OQE01217.1-29845-Penicillium vulpinum-Eurotiomycetes
XP 016594713.1-27334-Penicillium expansum-Eurotiomycetes
CRG89870.1-28573-Talaromyces islandicus-Eurotiomycetes
XP 002144860.1-441960-Talaromyces marneffei ATC-Eurotiomycetes
KFX52360.1-1077442-Talaromyces marneffei PM1-Eurotiomycetes
OBT69657.1-1524831-Pseudogymnoascus sp 2334-Leotiomycetes
KFY41662.1-1420907-Pseudogymnoascus sp VKM-Leotiomycetes
KFY14206.1-1420902-Pseudogymnoascus sp VKM-Leotiomycetes
OBT82103.1-1622148-Pseudogymnoascus sp 03VT-Leotiomycetes
OBT74270.1-1622149-Pseudogymnoascus sp 05NY-Leotiomycetes
KFY39125.1-1420908-Pseudogymnoascus sp VKM-Leotiomycetes
KFY04764.1-1391699-Pseudogymnoascus sp VKM-Leotiomycetes
KFY01833.1-1437433-Pseudogymnoascus sp VKM-Leotiomycetes
OBT49351.1-1622147-Pseudogymnoascus sp WSF-Leotiomycetes
XP 012741383.1-658429-Pseudogymnoascus destruct-Leotiomycetes
OAF54784.1-655981-Pseudogymnoascus destruct-Leotiomycetes
XP 018130572.1-342668-Pseudogymnoascus verrucos-Leotiomycetes
KFZ09854.1-1420914-Pseudogymnoascus sp VKM-Leotiomycetes
OBT58818.1-1622150-Pseudogymnoascus sp 24MN-Leotiomycetes
KHJ33454.1-52586-Erysiphe necator-Leotiomycetes
EPQ66190.1-1268274-Blumeria graminis f sp-Leotiomycetes
XP 001225799.1-306901-Chaetomium globosum CBS 1-Sordariomycetes
CEJ60096.1-104259-Penicillium brasilianum-Eurotiomycetes
XP 002846710.1-554155-Arthroderma otae CBS 1134-Eurotiomycetes
XP 006969239.1-431241-Trichoderma reesei QM6a-Sordariomycetes
KDN62805.1-1173701-Colletotrichum sublineola-Sordariomycetes
KJK64082.1-1403190-Aspergillus parasiticus S-Eurotiomycetes
XP 002378740.1-332952-Aspergillus flavus NRRL33-Eurotiomycetes
KJJ30093.1-1392242-Aspergillus flavus AF70-Eurotiomycetes
XP 003190316.1-510516-Aspergillus oryzae RIB40-Eurotiomycetes
OQE91512.1-60175-Penicillium nalgiovense-Eurotiomycetes
XP 016595100.1-27334-Penicillium expansum-Eurotiomycetes
OQE39328.1-36646-Penicillium coprophilum-Eurotiomycetes
OQE11539.1-29845-Penicillium vulpinum-Eurotiomycetes
XP 002561285.1-500485-Penicillium rubens Wiscon-Eurotiomycetes
OQE14832.1-254877-Penicillium flavigenum-Eurotiomycetes
OQE02657.1-60172-Penicillium solitum-Eurotiomycetes
CRL27040.1-5075-Penicillium camemberti-Eurotiomycetes
OQD64769.1-60169-Penicillium polonicum-Eurotiomycetes

S4I

Chr 4 — MFS transporter — Gypsy transposon — CYP450 — PKS — Transporter — Protein kinase — O-methyltransferase — Aldehyde reductase — Amidohydrolase — CYP450 — Mariner transposon — Gypsy transposon — Chr 4

Afu4g01560   AFUB_102040

Tree scale: 0.1

Taxonomy

- Leotiomycetes
- Sordariomycetes
- Eurotiomycetes
- Dothideomycetes
- other_Pezizomycotina
- other_Ascomycota
- other_Fungi
- other_Opisthokonta
- other_Eukaryota
- Bacteria
- Archaea
- Viruses

XP 003067208.1-222929-Coccidioides posadasii C7-Eurotiomycetes
XP 001239644.2-246410-Coccidioides immitis RS-Eurotiomycetes
QUERY-g6976-746128-Aspergillus fumigatus F13619-Eurotiomycetes
CDM31026.1-1365484-Penicillium roqueforti FM-Eurotiomycetes
OQE07377.1-29845-Penicillium vulpinum-Eurotiomycetes
OGM50384.1-109264-Aspergillus bombycis-Eurotiomycetes
GAD99701.1-1356009-Byssochlamys spectabilis-Eurotiomycetes
CCX29578.1-1076935-Pyronema omphalodes CBS 1-other Pezizomycotina
XP 018190638.1-1328760-Xylona heveae TC161-other Pezizomycotina
XP 016265392.1-215243-Exophiala oligosperma-Eurotiomycetes
XP 007719706.1-1182541-Capronia coronata CBS 617-Eurotiomycetes
XP 002542375.1-336963-Uncinocarpus reesii 1704-Eurotiomycetes
XP 001245277.1-246410-Coccidioides immitis RS-Eurotiomycetes
XP 003071570.1-222929-Coccidioides posadasii C7-Eurotiomycetes
XP 020119833.1-1441469-Talaromyces atroroseus-Eurotiomycetes
XP 002484266.1-441959-Talaromyces stipitatus AT-Eurotiomycetes
CRG86649.1-28573-Talaromyces islandicus-Eurotiomycetes
GAM37920.1-1472165-Talaromyces cellulolyticu-Eurotiomycetes
XP 002150005.1-441960-Talaromyces marneffei ATC-Eurotiomycetes
OJD24174.1-1658174-Blastomyces percursus-Eurotiomycetes
XP 002627299.1-559298-Blastomyces gilchristii S-Eurotiomycetes
KLJ05464.1-1246674-Emmonsia parva UAMH 139-Eurotiomycetes
OAX83378.1-1658172-Emmonsia sp CAC 2015a-Eurotiomycetes
KKZ67067.1-1247875-Emmonsia crescens UAMH 30-Eurotiomycetes
EGE01764.1-559882-Trichophyton equinum CBS-Eurotiomycetes
EZF34609.1-1215336-Trichophyton interdigital-Eurotiomycetes
XP 003232978.1-559305-Trichophyton rubrum CBS 1-Eurotiomycetes
EZF25477.1-1215337-Trichophyton rubrum MR850-Eurotiomycetes
DAA74814.1-663331-Trichophyton benhamiae CB-Eurotiomycetes
EGD97296.1-647933-Trichophyton tonsurans CB-Eurotiomycetes
XP 003175358.1-535722-Nannizzia gypsea CBS 1188-Eurotiomycetes
KIM95126.1-913774-Oidiodendron maius Zn-Leotiomycetes
ESZ95100.1-1432307-Sclerotinia borealis F 41-Leotiomycetes
XP 001591780.1-665079-Sclerotinia sclerotiorum-Leotiomycetes
XP 001555725.1-332648-Botrytis cinerea B05 10-Leotiomycetes
CZT50538.1-38038-Rhynchosporium secalis-Leotiomycetes
XP 018075789.1-149040-Phialocephala scopiformis-Leotiomycetes
CZR67391.1-576137-Phialocephala subalpina-Leotiomycetes
OCK85450.1-1314670-Lepidopterella palustris-Dothideomycetes
XP 013431178.1-1043004-Aureobasidium namibiae CB-Dothideomycetes
XP 013341200.1-1043005-Aureobasidium subglaciale-Dothideomycetes
XP 018040703.1-1460663-Paraphaeosphaeria sporulo-Dothideomycetes
XP 003300409.1-861557-Pyrenophora teres f tere-Dothideomycetes
OAL06049.1-765868-Stagonospora sp SRC1lsM3-Dothideomycetes
OSS43606.1-105696-Epicoccum nigrum-Dothideomycetes
KKY19094.1-420778-Diplodia seriata-Dothideomycetes
XP 020133117.1-236234-Diplodia corticola-Dothideomycetes
EKG17053.1-1126212-Macrophomina phaseolina M-Dothideomycetes
XP 007580094.1-1287680-Neofusicoccum parvum UCRN-Dothideomycetes
KKZ63467.1-1247875-Emmonsia crescens UAMH 30-Eurotiomycetes
OAX77823.1-1658172-Emmonsia sp CAC 2015a-Eurotiomycetes
OJD11794.1-1447872-Emergomyces pasteuriana E-Eurotiomycetes
EGC46119.1-544711-Histoplasma capsulatum H8-Eurotiomycetes
XP 001540840.1-339724-Histoplasma capsulatum NA-Eurotiomycetes
EEH11142.1-447093-Histoplasma capsulatum G1-Eurotiomycetes
KLJ09996.1-1246674-Emmonsia parva UAMH 139-Eurotiomycetes
XP 002624675.1-559298-Blastomyces gilchristii S-Eurotiomycetes
ODH53684.1-121759-Paracoccidioides brasilie-Eurotiomycetes
ODH39903.1-121759-Paracoccidioides brasilie-Eurotiomycetes
EEH16171.1-482561-Paracoccidioides brasilie-Eurotiomycetes
XP 010763114.1-502780-Paracoccidioides brasilie-Eurotiomycetes
XP 015699878.1-502779-Paracoccidioides lutzii P-Eurotiomycetes
OOF94419.1-602072-Aspergillus carbonarius I-Eurotiomycetes
XP 001216622.1-341663-Aspergillus terreus NIH26-Eurotiomycetes
XP 020051878.1-690307-Aspergillus aculeatus ATC-Eurotiomycetes
CEN60170.1-454130-Aspergillus calidoustus-Eurotiomycetes
OJJ08067.1-1036611-Aspergillus versicolor CB-Eurotiomycetes
XP 747931.1-330879-Aspergillus fumigatus Af2-Eurotiomycetes
GAQ08997.1-293939-Aspergillus lentulus-Eurotiomycetes
XP 001266122.1-331117-Aspergillus fischeri NRRL-Eurotiomycetes
XP 001276120.1-344612-Aspergillus clavatus NRRL-Eurotiomycetes
KEY75694.1-1476192-Aspergillus fumigatus var-Eurotiomycetes
GAO84760.1-91492-Aspergillus udagawae-Eurotiomycetes
OQD72719.1-69771-Penicillium decumbens-Eurotiomycetes
OJJ33071.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
KKK12354.1-138278-Aspergillus ochraceoroseu-Eurotiomycetes
KKK15958.1-308745-Aspergillus rambellii-Eurotiomycetes
ODM19926.1-573508-Aspergillus cristatus-Eurotiomycetes
EYE95855.1-1388766-Aspergillus ruber CBS 135-Eurotiomycetes
OQD80537.1-416450-Penicillium antarcticum-Eurotiomycetes
OGE57334.1-1835702-Penicillium arizonense-Eurotiomycetes
OKP11883.1-1316194-Penicillium subrubescens-Eurotiomycetes
EPS34191.1-933388-Penicillium oxalicum 114-Eurotiomycetes
OOQ83889.1-104259-Penicillium brasilianum-Eurotiomycetes
CEJ62117.1-104259-Penicillium brasilianum-Eurotiomycetes
OJJ69307.1-767769-Aspergillus brasiliensis-Eurotiomycetes
XP 001393968.1-425011-Aspergillus niger CBS 513-Eurotiomycetes
EHA28595.1-380704-Aspergillus niger ATCC 10-Eurotiomycetes
OJI88301.1-767770-Aspergillus tubingensis C-Eurotiomycetes
OJZ82136.1-1137211-Aspergillus luchuensis CB-Eurotiomycetes
GAQ43896.1-5061-Aspergillus niger-Eurotiomycetes
GAA88176.1-1033177-Aspergillus kawachii IFO-Eurotiomycetes
OOO11703.1-5062-Aspergillus oryzae-Eurotiomycetes
EIT83343.1-1160506-Aspergillus oryzae 3 042-Eurotiomycetes
XP 001822762.2-510516-Aspergillus oryzae RIB40-Eurotiomycetes
XP 002382888.1-332952-Aspergillus flavus NRRL33-Eurotiomycetes
BAE61629.1-510516-Aspergillus oryzae RIB40-Eurotiomycetes
KOC15977.1-1392242-Aspergillus flavus AF70-Eurotiomycetes
OGM43570.1-109264-Aspergillus bombycis-Eurotiomycetes
XP 015401387.1-1509407-Aspergillus nomius NRRL 1-Eurotiomycetes

S4J

Tree scale: 0.1 ⊢——⊣

Taxonomy
- Leotiomycetes
- Sordariomycetes
- Eurotiomycetes
- Dothideomycetes
- other_Pezizomycotina
- other_Ascomycota
- other_Fungi
- other_Opisthokonta
- other_Eukaryota
- Bacteria
- Archaea
- Viruses

S4K

Tree scale: 0.1 ⊢─┤

Taxonomy

- 🟥 Leotiomycetes
- 🟪 Sordariomycetes
- 🟧 Eurotiomycetes
- 🟧 Dothideomycetes
- 🟫 other_Pezizomycotina
- 🟪 other_Ascomycota
- 🟪 other_Fungi
- 🟦 other_Opisthokonta
- 🟦 other_Eukaryota
- 🟩 Bacteria
- 🟨 Archaea
- ⬜ Viruses

S4L

Tree scale: 0.1 ⊢——⊣

Taxonomy

- Leotiomycetes
- Sordariomycetes
- Eurotiomycetes
- Dothideomycetes
- other_Pezizomycotina
- other_Ascomycota
- other_Fungi
- other_Opisthokonta
- other_Eukaryota
- Bacteria
- Archaea
- Viruses

OTB08227.1-1001833-Hypoxylon sp CI 4A-Sordariomycetes
KEY73627.1-1280523-Stachybotrys chartarum IB-Sordariomycetes
XP 018029975.1-1460663-Paraphaeosphaeria sporulo-Dothideomycetes
GAP88297.1-77044-Rosellinia necatrix-Sordariomycetes
OJJ67825.1-767769-Aspergillus brasiliensis-Eurotiomycetes
G3Y417.1-380704-Aspergillus niger ATCC 10-Eurotiomycetes
XP 001402406.2-425011-Aspergillus niger CBS 513-Eurotiomycetes
CAK49103.1-5061-Aspergillus niger-Eurotiomycetes
OQE12755.1-254877-Penicillium flavigenum-Eurotiomycetes
KXG51721.1-5078-Penicillium griseofulvum-Eurotiomycetes
OQD98988.1-29845-Penicillium vulpinum-Eurotiomycetes
CDM29146.1-1365484-Penicillium roqueforti FM-Eurotiomycetes
KGO67466.1-40296-Penicillium italicum-Eurotiomycetes
XP 002560462.1-500485-Penicillium rubens Wiscon-Eurotiomycetes
KZN83408.1-5076-Penicillium chrysogenum-Eurotiomycetes
ESZ91463.1-1432307-Sclerotinia borealis F 41-Leotiomycetes
QUERY-g6980-746128-Aspergillus fumigatus F13619-Eurotiomycetes
KJK74884.1-1291518-Metarhizium anisopliae BR-Sordariomycetes
KKP06688.1-5544-Trichoderma harzianum-Sordariomycetes
KJZ75228.1-1043627-Hirsutella minnesotensis-Sordariomycetes
OCK73150.1-1314670-Lepidopterella palustris-Dothideomycetes
CEN59761.1-454130-Aspergillus calidoustus-Eurotiomycetes
OJJ57536.1-1036612-Aspergillus sydowii CBS 5-Eurotiomycetes
OJJ05643.1-1036611-Aspergillus versicolor CB-Eurotiomycetes
XP 661205.1-227321-Aspergillus nidulans FGSC-Eurotiomycetes
CBF75802.1-227321-Aspergillus nidulans FGSC-Eurotiomycetes
KXX79207.1-100816-Madurella mycetomatis-Sordariomycetes
KFA69547.1-1283841-Stachybotrys chlorohalona-Sordariomycetes
KFA52212.1-1280524-Stachybotrys chartarum IB-Sordariomycetes
KEY72232.1-1280523-Stachybotrys chartarum IB-Sordariomycetes
KKY16576.1-420778-Diplodia seriata-Dothideomycetes
XP 020125930.1-236234-Diplodia corticola-Dothideomycetes
XP 003300441.1-861557-Pyrenophora teres f tere-Dothideomycetes
XP 001937756.1-426418-Pyrenophora tritici repen-Dothideomycetes
XP 008027821.1-671987-Setosphaeria turcica Et28-Dothideomycetes
XP 007689107.1-930090-Bipolaris oryzae ATCC 445-Dothideomycetes
XP 014073545.1-665024-Bipolaris maydis ATCC 483-Dothideomycetes
XP 011127785.1-756982-Arthrobotrys oligospora A-other Pezizomycotina
OJZ92826.1-1137211-Aspergillus luchuensis CB-Eurotiomycetes
XP 013282149.1-1442368-Fonsecaea pedrosoi CBS 27-Eurotiomycetes
KIJ36684.1-990650-Sphaerobolus stellatus SS-Basidiomycota
GAU94922.1-947166-Ramazzottius varieornatus-Metazoa
EKG18972.1-1126212-Macrophomina phaseolina M-Dothideomycetes
GAW15911.1-1813822-fungal sp No 14919-other Fungi
XP 007795392.1-1287681-Eutypa lata UCREL1-Sordariomycetes
XP 008712499.1-1220924-Cyphellophora europaea CB-Eurotiomycetes
OGM49889.1-109264-Aspergillus bombycis-Eurotiomycetes
OJJ66452.1-767769-Aspergillus brasiliensis-Eurotiomycetes
OGM45328.1-109264-Aspergillus bombycis-Eurotiomycetes
GAD97254.1-1356009-Byssochlamys spectabilis-Eurotiomycetes
OJJ37490.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
XP 018061083.1-149040-Phialocephala scopiformis-Leotiomycetes
XP 018131309.1-342668-Pseudogymnoascus verrucos-Leotiomycetes
OGM45263.1-109264-Aspergillus bombycis-Eurotiomycetes
XP 007790916.1-1287681-Eutypa lata UCREL1-Sordariomycetes
GAD91854.1-1356009-Byssochlamys spectabilis-Eurotiomycetes
XP 018700082.1-1081104-Isaria fumosorosea ARSEF-Sordariomycetes
XP 007914153.1-1286976-Phaeoacremonium minimum U-Sordariomycetes
XP 003043989.1-660122-Nectria haematococca mpVI-Sordariomycetes
XP 015406133.1-1509407-Aspergillus nomius NRRL 1-Eurotiomycetes
OGM49559.1-109264-Aspergillus bombycis-Eurotiomycetes
OPB36055.1-1491466-Trichoderma guizhouense-Sordariomycetes
XP 013938758.1-452589-Trichoderma atroviride IM-Sordariomycetes
XP 018664929.1-398673-Trichoderma gamsii-Sordariomycetes
KND94247.1-1163406-Tolypocladium ophioglosso-Sordariomycetes
KOM23882.1-268505-Ophiocordyceps unilateral-Sordariomycetes
CRG89981.1-28573-Talaromyces islandicus-Eurotiomycetes
KID82384.1-1276136-Metarhizium guizhouense A-Sordariomycetes
XP 014543987.1-1276141-Metarhizium brunneum ARSE-Sordariomycetes
XP 007826330.1-655844-Metarhizium robertsii ARS-Sordariomycetes
KJK73576.1-1291518-Metarhizium anisopliae BR-Sordariomycetes
KFG78219.1-5530-Metarhizium anisopliae-Sordariomycetes
KIL85477.1-40199-Fusarium avenaceum-Sordariomycetes
EMT72782.1-1229665-Fusarium oxysporum f sp-Sordariomycetes
EWZ45894.1-660027-Fusarium oxysporum Fo47-Sordariomycetes
EXK43331.1-1089452-Fusarium oxysporum f sp-Sordariomycetes
XP 018255266.1-426428-Fusarium oxysporum f sp-Sordariomycetes
CVK96608.1-192010-Fusarium mangiferae-Sordariomycetes
CZR45965.1-1227346-Fusarium proliferatum ET1-Sordariomycetes
CVL11161.1-948311-Fusarium proliferatum-Sordariomycetes
CCT71951.1-1279085-Fusarium fujikuroi IMI 58-Sordariomycetes
KLO92536.1-5127-Fusarium fujikuroi-Sordariomycetes
XP 007381777.1-741275-Punctularia strigosozonat-Basidiomycota
OAA80406.1-1081108-Cordyceps confragosa RCEF-Sordariomycetes
OJJ46451.1-1073090-Penicilliopsis zonata CBS-Eurotiomycetes
OQE00380.1-29845-Penicillium vulpinum-Eurotiomycetes
OJJ38752.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
XP 001273089.1-344612-Aspergillus clavatus NRRL-Eurotiomycetes
OQD99182.1-60172-Penicillium solitum-Eurotiomycetes
XP 014535842.1-1170230-Penicillium digitatum Pd1-Eurotiomycetes
OQE13089.1-254877-Penicillium flavigenum-Eurotiomycetes
OQD65548.1-60169-Penicillium polonicum-Eurotiomycetes
CRL29160.1-5075-Penicillium camemberti-Eurotiomycetes
KUM63385.1-48697-Penicillium freii-Eurotiomycetes
OQE09941.1-29845-Penicillium vulpinum-Eurotiomycetes
OQE43328.1-36646-Penicillium coprophilum-Eurotiomycetes
KXG50992.1-5078-Penicillium griseofulvum-Eurotiomycetes
OQD86089.1-416450-Penicillium antarcticum-Eurotiomycetes
AIG62133.1-27334-Penicillium expansum-Eurotiomycetes
XP 016595358.1-27334-Penicillium expansum-Eurotiomycetes

S4M

Taxonomy

- Leotiomycetes
- Sordariomycetes
- Eurotiomycetes
- Dothideomycetes
- other_Pezizomycotina
- other_Ascomycota
- other_Fungi
- other_Opisthokonta
- other_Eukaryota
- Bacteria
- Archaea
- Viruses

Figure S6. Gene phylogenies of the mobile SM gene cluster 1. These phylogenies are consistent with horizontal transfer between Eurotiomycete, Dothidiomycete, Leotiomycete, and Sordariomycete fungi.

S6B

Tree scale: 0.1 ⊢———⊣

Taxonomy

- ■ Leotiomycetes
- ■ Sordariomycetes
- ■ Eurotiomycetes
- ■ Dothideomycetes
- ■ other_Pezizomycotina
- ■ other_Ascomycota
- ■ other_Fungi
- ■ other_Opisthokonta
- ■ other_Eukaryota
- ■ Bacteria
- ■ Archaea
- ■ Viruses

XP 018190871.1-1328760-Xylona heveae TC161-other Pezizomycotina
KNG52497.1-183478-Stemphylium lycopersici-Dothideomycetes
ESZ91465.1-1432307-Sclerotinia borealis F 41-Leotiomycetes
KFA69534.1-1283841-Stachybotrys chlorohalona-Sordariomycetes
KFA74441.1-1283842-Stachybotrys chartarum IB-Sordariomycetes
XP 008027822.1-671987-Setosphaeria turcica Et28-Dothideomycetes
XP 014073570.1-665024-Bipolaris maydis ATCC 483-Dothideomycetes
XP 007689106.1-930090-Bipolaris oryzae ATCC 445-Dothideomycetes
XP 001937757.1-426418-Pyrenophora tritici repen-Dothideomycetes
XP 003300440.1-861557-Pyrenophora teres f tere-Dothideomycetes
KKY16577.1-420778-Diplodia seriata-Dothideomycetes
XP 020125929.1-236234-Diplodia corticola-Dothideomycetes
GAM36611.1-1472165-Talaromyces cellulolyticu-Eurotiomycetes
XP 003654926.1-578455-Thielavia terrestris NRRL-Sordariomycetes
GAM40107.1-1472165-Talaromyces cellulolyticu-Eurotiomycetes
XP 002479463.1-441959-Talaromyces stipitatus AT-Eurotiomycetes
EHA28243.1-380704-Aspergillus niger ATCC 10-Eurotiomycetes
XP 001393507.1-425011-Aspergillus niger CBS 513-Eurotiomycetes
XP 013942627.1-452589-Trichoderma atroviride IM-Sordariomycetes
XP 018661927.1-398673-Trichoderma gamsii-Sordariomycetes
OAF54392.1-655981-Pseudogymnoascus destruct-Leotiomycetes
KUI69403.1-105487-Valsa mali-Sordariomycetes
KUI53500.1-694573-Valsa mali var pyri-Sordariomycetes
OJJ06460.1-1036611-Aspergillus versicolor CB-Eurotiomycetes
SLM38091.1-136370-Umbilicaria pustulata-other Pezizomycotina
XP 007801361.1-1263415-Endocarpon pusillum Z0702-Eurotiomycetes
XP 008086617.1-1116229-Glarea lozoyensis ATCC 20-Leotiomycetes
XP 018142822.1-1380566-Pochonia chlamydosporia 1-Sordariomycetes
OCL02959.1-574774-Glonium stellatum-Dothideomycetes
OAK94599.1-765868-Stagonospora sp SRC1lsM3-Dothideomycetes
CRG83420.1-28573-Talaromyces islandicus-Eurotiomycetes
KUL90120.1-198730-Talaromyces verruculosus-Eurotiomycetes
OJJ67836.1-767769-Aspergillus brasiliensis-Eurotiomycetes
OJI89149.1-767770-Aspergillus tubingensis C-Eurotiomycetes
ODM21323.1-573508-Aspergillus cristatus-Eurotiomycetes
XP 018189398.1-1328760-Xylona heveae TC161-other Pezizomycotina
KFY42098.1-1420907-Pseudogymnoascus sp VKM-Leotiomycetes
KOS21702.1-150374-Escovopsis weberi-Sordariomycetes
XP 001803249.1-321614-Parastagonospora nodorum-Dothideomycetes
QUERY-Afu1g00980-330879-Aspergillus fumigatus Af2-Eurotiomycetes
KMK54467.1-1437362-Aspergillus fumigatus Z5-Eurotiomycetes
OTA99527.1-1001833-Hypoxylon sp CI 4A-Sordariomycetes
OTA97076.1-1001938-Hypoxylon sp CO27 5-Sordariomycetes
OTA68786.1-1001937-Hypoxylon sp EC38-Sordariomycetes
OQE10144.1-254877-Penicillium flavigenum-Eurotiomycetes
KXG46525.1-5078-Penicillium griseofulvum-Eurotiomycetes
XP 020058182.1-690307-Aspergillus aculeatus ATC-Eurotiomycetes
CDM37416.1-1365484-Penicillium roqueforti FM-Eurotiomycetes
OQD99008.1-29845-Penicillium vulpinum-Eurotiomycetes
KOS46297.1-229535-Penicillium nordicum-Eurotiomycetes
OQE96749.1-60175-Penicillium nalgiovense-Eurotiomycetes
XP 002564777.1-500485-Penicillium rubens Wiscon-Eurotiomycetes
OJZ86826.1-1137211-Aspergillus luchuensis CB-Eurotiomycetes
GAA84845.1-1033177-Aspergillus kawachii IFO-Eurotiomycetes
GAT24549.1-1069201-Aspergillus luchuensis-Eurotiomycetes
GAQ46578.1-5061-Aspergillus niger-Eurotiomycetes
OJI83692.1-767770-Aspergillus tubingensis C-Eurotiomycetes
CEN59740.1-454130-Aspergillus calidoustus-Eurotiomycetes
KIA75595.1-40382-Aspergillus ustus-Eurotiomycetes
KUL83490.1-198730-Talaromyces verruculosus-Eurotiomycetes
GAM39560.1-1472165-Talaromyces cellulolyticu-Eurotiomycetes
KFX41538.1-1077442-Talaromyces marneffei PM1-Eurotiomycetes
XP 002145721.1-441960-Talaromyces marneffei ATC-Eurotiomycetes
KKY17669.1-420778-Diplodia seriata-Dothideomycetes
OMP84237.1-420778-Diplodia seriata-Dothideomycetes
XP 007786093.1-1263415-Endocarpon pusillum Z0702-Eurotiomycetes
XP 008720114.1-1220924-Cyphellophora europaea CB-Eurotiomycetes
OJJ62960.1-1036612-Aspergillus sydowii CBS 5-Eurotiomycetes
OTA57984.1-1001937-Hypoxylon sp EC38-Sordariomycetes
OCW41151.1-158607-Diaporthe helianthi-Sordariomycetes
XP 007795874.1-1287681-Eutypa lata UCREL1-Sordariomycetes
OOQ89673.1-104259-Penicillium brasilianum-Eurotiomycetes
OQD81687.1-416450-Penicillium antarcticum-Eurotiomycetes
KFA79676.1-1283842-Stachybotrys chartarum IB-Sordariomycetes
KEY69109.1-1280523-Stachybotrys chartarum IB-Sordariomycetes
KFA45396.1-1280524-Stachybotrys chartarum IB-Sordariomycetes
OTB10571.1-1001832-Daldinia sp EC12-Sordariomycetes
KLO99253.1-5127-Fusarium fujikuroi-Sordariomycetes
XP 018754944.1-334819-Fusarium verticillioides-Sordariomycetes
ANH22767.1-696354-Hypocrella siamensis-Sordariomycetes
KIA75781.1-40382-Aspergillus ustus-Eurotiomycetes
KOM23411.1-268505-Ophiocordyceps unilateral-Sordariomycetes
KEY71722.1-1280523-Stachybotrys chartarum IB-Sordariomycetes
KFA79949.1-1283842-Stachybotrys chartarum IB-Sordariomycetes
KFA55988.1-1280524-Stachybotrys chartarum IB-Sordariomycetes
KFA68349.1-1283841-Stachybotrys chlorohalona-Sordariomycetes
KFA70203.1-1283841-Stachybotrys chlorohalona-Sordariomycetes
KFA71244.1-1283842-Stachybotrys chartarum IB-Sordariomycetes
KEY71773.1-1280523-Stachybotrys chartarum IB-Sordariomycetes
KFA54577.1-1280524-Stachybotrys chartarum IB-Sordariomycetes
ORY71423.1-1141098-Pseudomassariella vexata-Sordariomycetes
KLU89215.1-644358-Magnaporthiopsis poae ATC-Sordariomycetes
XP 003712304.1-242507-Magnaporthe oryzae 70 15-Sordariomycetes
ELQ69713.1-1143193-Magnaporthe oryzae P131-Sordariomycetes
XP 003351972.1-771870-Sordaria macrospora k hel-Sordariomycetes
CAQ58437.1-5147-Sordaria macrospora-Sordariomycetes
XP 965608.1-367110-Neurospora crassa OR74A-Sordariomycetes
EGZ76379.1-510952-Neurospora tetrasperma FG-Sordariomycetes
XP 009855463.1-510951-Neurospora tetrasperma FG-Sordariomycetes

S6C

Tree scale: 0.1

Taxonomy

- Leotiomycetes
- Sordariomycetes
- Eurotiomycetes
- Dothideomycetes
- other_Pezizomycotina
- other_Ascomycota
- other_Fungi
- other_Opisthokonta
- other_Eukaryota
- Bacteria
- Archaea
- Viruses

KZL76078.1-708197-Colletotrichum tofieldiae-Sordariomycetes
KZL85173.1-1573173-Colletotrichum incanum-Sordariomycetes
XP 008089668.1-645133-Colletotrichum graminicol-Sordariomycetes
CEJ62862.1-104259-Penicillium brasilianum-Eurotiomycetes
KUL90439.1-198730-Talaromyces verruculosus-Eurotiomycetes
GAA84860.1-1033177-Aspergillus kawachii IFO-Eurotiomycetes
GAQ46564.1-5061-Aspergillus niger-Eurotiomycetes
OJI83706.1-767770-Aspergillus tubingensis C-Eurotiomycetes
XP 007738022.1-1182542-Capronia epimyces CBS 606-Eurotiomycetes
KIV86245.1-1016849-Exophiala sideris-Eurotiomycetes
KIV78870.1-1016849-Exophiala sideris-Eurotiomycetes
XP 016635763.1-1442371-Fonsecaea multimorphosa C-Eurotiomycetes
XP 018689742.1-1367422-Fonsecaea erecta-Eurotiomycetes
KEQ57958.1-1043003-Aureobasidium melanogenum-Dothideomycetes
XP 007582626.1-1287680-Neofusicoccum parvum UCRN-Dothideomycetes
XP 007919710.1-383855-Pseudocercospora fijiensi-Dothideomycetes
KXT10441.1-113226-Pseudocercospora musae-Dothideomycetes
GAM33346.1-1472165-Talaromyces cellulolyticu-Eurotiomycetes
KUL83173.1-198730-Talaromyces verruculosus-Eurotiomycetes
OCL12203.1-574774-Glonium stellatum-Dothideomycetes
OCK78800.1-1314670-Lepidopterella palustris-Dothideomycetes
OQO09931.1-1507870-Rachicladosporium antarct-Dothideomycetes
OQO06724.1-1507870-Rachicladosporium antarct-Dothideomycetes
OQO25097.1-1974281-Rachicladosporium sp CCF-Dothideomycetes
KIN03586.1-913774-Oidiodendron maius Zn-Leotiomycetes
CZR67742.1-576137-Phialocephala subalpina-Leotiomycetes
XP 018070937.1-149040-Phialocephala scopiformis-Leotiomycetes
XP 013270009.1-1442369-Rhinocladiella mackenziei-Eurotiomycetes
KIV85980.1-1016849-Exophiala sideris-Eurotiomycetes
XP 013316661.1-348802-Exophiala xenobiotica-Eurotiomycetes
XP 016268262.1-215243-Exophiala oligosperma-Eurotiomycetes
XP 016240447.1-91928-Exophiala spinifera-Eurotiomycetes
OCL02960.1-574774-Glonium stellatum-Dothideomycetes
OCK79863.1-1314670-Lepidopterella palustris-Dothideomycetes
GAD94648.1-1356009-Byssochlamys spectabilis-Eurotiomycetes
XP 007361702.1-732165-Dichomitus squalens LYAD-Basidiomycota
XP 018068436.1-149040-Phialocephala scopiformis-Leotiomycetes
OJJ41604.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
XP 018067827.1-149040-Phialocephala scopiformis-Leotiomycetes
OCK97824.1-794803-Cenococcum geophilum 1 58-Dothideomycetes
OCL04641.1-574774-Glonium stellatum-Dothideomycetes
ODM21324.1-573508-Aspergillus cristatus-Eurotiomycetes
XP 018189399.1-1328760-Xylona heveae TC161-other Pezizomycotina
ODM17455.1-573508-Aspergillus cristatus-Eurotiomycetes
OJI89150.1-767770-Aspergillus tubingensis C-Eurotiomycetes
OJJ67835.1-767769-Aspergillus brasiliensis-Eurotiomycetes
GAQ11743.1-293939-Aspergillus lentulus-Eurotiomycetes
XP 008086616.1-1116229-Glarea lozoyensis ATCC 20-Leotiomycetes
XP 007798352.1-1287681-Eutypa lata UCREL1-Sordariomycetes
CRG83420.1-28573-Talaromyces islandicus-Eurotiomycetes
GAM35249.1-1472165-Talaromyces cellulolyticu-Eurotiomycetes
KUL90121.1-198730-Talaromyces verruculosus-Eurotiomycetes
KFY42097.1-1420907-Pseudogymnoascus sp VKM-Leotiomycetes
KOS21698.1-150374-Escovopsis weberi-Sordariomycetes
XP 003654012.1-578455-Thielavia terrestris NRRL-Sordariomycetes
OTA99528.1-1001833-Hypoxylon sp CI 4A-Sordariomycetes
OTA97075.1-1001938-Hypoxylon sp CO27 5-Sordariomycetes
OTA68787.1-1001937-Hypoxylon sp EC38-Sordariomycetes
XP 001803248.1-321614-Parastagonospora nodorum-Dothideomycetes
XP 014072629.1-665024-Bipolaris maydis ATCC 483-Dothideomycetes
QUERY-Afu1g00990-330879-Aspergillus fumigatus Af2-Eurotiomycetes
XP 018139917.1-1380566-Pochonia chlamydosporia 1-Sordariomycetes
XP 018175680.1-33203-Purpureocillium lilacinum-Sordariomycetes
GAW20535.1-1813822-fungal sp No 14919-other Fungi
XP 007286401.1-1213859-Colletotrichum gloeospori-Sordariomycetes
XP 007799272.1-1287681-Eutypa lata UCREL1-Sordariomycetes
OAA76696.1-1081108-Cordyceps confragosa RCEF-Sordariomycetes
XP 018703534.1-1081104-Isaria fumosorosea ARSEF-Sordariomycetes
XP 007599778.1-1445577-Colletotrichum fioriniae-Sordariomycetes
OHF00529.1-1209926-Colletotrichum orchidophi-Sordariomycetes
KXH66733.1-1209931-Colletotrichum salicis-Sordariomycetes
KXH41440.1-1460502-Colletotrichum nymphaeae-Sordariomycetes
KXH45168.1-703756-Colletotrichum simmondsii-Sordariomycetes
XP 013942446.1-452589-Trichoderma atroviride IM-Sordariomycetes
XP 013958002.1-413071-Trichoderma virens Gv29 8-Sordariomycetes
OPB36426.1-1491466-Trichoderma guizhouense-Sordariomycetes
KKP06960.1-5544-Trichoderma harzianum-Sordariomycetes
OTA07189.1-858221-Trichoderma parareesei-Sordariomycetes
ETR98295.1-1344414-Trichoderma reesei RUT C-Sordariomycetes
XP 006968870.1-431241-Trichoderma reesei QM6a-Sordariomycetes
KPM35926.1-78410-Neonectria ditissima-Sordariomycetes
XP 003041834.1-660122-Nectria haematococca mpVI-Sordariomycetes
KIL93926.1-40199-Fusarium avenaceum-Sordariomycetes
XP 018760089.1-334819-Fusarium verticillioides-Sordariomycetes
CVK86026.1-192010-Fusarium mangiferae-Sordariomycetes
KLP07008.1-5127-Fusarium fujikuroi-Sordariomycetes
KLO82373.1-5127-Fusarium fujikuroi-Sordariomycetes
CVK85442.1-948311-Fusarium proliferatum-Sordariomycetes
CZR37956.1-1227346-Fusarium proliferatum ET1-Sordariomycetes
EXM21264.1-1089449-Fusarium oxysporum f sp-Sordariomycetes
EXL74715.1-1089457-Fusarium oxysporum f sp-Sordariomycetes
EXL52372.1-1089448-Fusarium oxysporum f sp-Sordariomycetes
EWZ80636.1-1080343-Fusarium oxysporum f sp-Sordariomycetes
EWY82337.1-909455-Fusarium oxysporum FOSC 3-Sordariomycetes
EXA36705.1-1080344-Fusarium oxysporum f sp-Sordariomycetes
EWZ31047.1-660027-Fusarium oxysporum Fo47-Sordariomycetes
XP 018252497.1-426428-Fusarium oxysporum f sp-Sordariomycetes
EXK84015.1-1089458-Fusarium oxysporum f sp-Sordariomycetes
EGU83440.1-660025-Fusarium oxysporum Fo5176-Sordariomycetes

S6D

Tree scale: 0.1 ⊢——⊣

Taxonomy

- ▇ Leotiomycetes
- ▇ Sordariomycetes
- ▇ Eurotiomycetes
- ▇ Dothideomycetes
- ▇ other_Pezizomycotina
- ▇ other_Ascomycota
- ▇ other_Fungi
- ▇ other_Opisthokonta
- ▇ other_Eukaryota
- ▇ Bacteria
- ▇ Archaea
- ▇ Viruses

CRG90955.1-28573-Talaromyces islandicus-Eurotiomycetes
OCL11392.1-574774-Glonium stellatum-Dothideomycetes
OCL02963.1-574774-Glonium stellatum-Dothideomycetes
OQD95988.1-29845-Penicillium vulpinum-Eurotiomycetes
OAA51374.1-1081107-Cordyceps brongniartii RC-Sordariomycetes
XP 008086613.1-1116229-Glarea lozoyensis ATCC 20-Leotiomycetes
EHK97192.1-1104152-Glarea lozoyensis 74030-Leotiomycetes
OAK94601.1-765868-Stagonospora sp SRC1lsM3-Dothideomycetes
KUL90201.1-198730-Talaromyces verruculosus-Eurotiomycetes
CRG83421.1-28573-Talaromyces islandicus-Eurotiomycetes
ENH79514.1-1213857-Colletotrichum orbiculare-Sordariomycetes
EKG13588.1-1126212-Macrophomina phaseolina M-Dothideomycetes
OJJ67834.1-767769-Aspergillus brasiliensis-Eurotiomycetes
OJI89151.1-767770-Aspergillus tubingensis C-Eurotiomycetes
ODM21322.1-573508-Aspergillus cristatus-Eurotiomycetes
XP 018189397.1-1328760-Xylona heveae TC161-other Pezizomycotina
KOS21699.1-150374-Escovopsis weberi-Sordariomycetes
OTA99529.1-1001833-Hypoxylon sp CI 4A-Sordariomycetes
KFY42096.1-1420907-Pseudogymnoascus sp VKM-Leotiomycetes
XP 001803247.1-321614-Parastagonospora nodorum-Dothideomycetes
QUERY-Afu1g01000-330879-Aspergillus fumigatus Af2-Eurotiomycetes
KMK54470.1-1437362-Aspergillus fumigatus Z5-Eurotiomycetes
XP 013262530.1-1182545-Exophiala aquamarina CBS-Eurotiomycetes
XP 008714593.1-1220924-Cyphellophora europaea CB-Eurotiomycetes
XP 017995192.1-1664694-Phialophora attae-Eurotiomycetes
XP 016219388.1-253628-Verruconis gallopava-Dothideomycetes
XP 016626305.1-1442371-Fonsecaea multimorphosa C-Eurotiomycetes
XP 007755064.1-1182544-Cladophialophora yegresii-Eurotiomycetes
OCT50474.1-86049-Cladophialophora carrioni-Eurotiomycetes
XP 008728080.1-1279043-Cladophialophora carrioni-Eurotiomycetes
EQB55257.1-1237896-Colletotrichum gloeospori-Sordariomycetes
KKY39798.1-1214573-Diaporthe ampelina-Sordariomycetes
KUI61446.1-694573-Valsa mali var pyri-Sordariomycetes
KUI63333.1-105487-Valsa mali-Sordariomycetes
OTB10749.1-1001832-Daldinia sp EC12-Sordariomycetes
OQO03845.1-1507870-Rachicladosporium antarct-Dothideomycetes
OQO13276.1-1507870-Rachicladosporium antarct-Dothideomycetes
OQO10333.1-1974281-Rachicladosporium sp CCF-Dothideomycetes
OTA35077.1-1157616-Hortaea werneckii EXF 200-Dothideomycetes
XP 007673282.1-717646-Baudoinia panamericana UA-Dothideomycetes
EME49121.1-675120-Dothistroma septosporum N-Dothideomycetes
KJY00698.1-1047168-Zymoseptoria brevis-Dothideomycetes
XP 003853448.1-336722-Zymoseptoria tritici IPO3-Dothideomycetes
SMQ49856.1-1276538-Zymoseptoria tritici ST99-Dothideomycetes
XP 007922236.1-383855-Pseudocercospora fijiensi-Dothideomycetes
KXT13488.1-113226-Pseudocercospora musae-Dothideomycetes
KXS97951.1-321146-Mycosphaerella eumusae-Dothideomycetes
XP 009256835.1-1028729-Fusarium pseudograminearu-Sordariomycetes
XP 001216325.1-341663-Aspergillus terreus NIH26-Eurotiomycetes
OJJ37159.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
GAD94095.1-1356009-Byssochlamys spectabilis-Eurotiomycetes
XP 013324705.1-1408163-Rasamsonia emersonii CBS-Eurotiomycetes
KZT59844.1-1353952-Calocera cornea HHB12733-Basidiomycota
XP 007781013.1-1168221-Coniosporium apollinis CB-Dothideomycetes
GAA91749.1-1033177-Aspergillus kawachii IFO-Eurotiomycetes
GAQ33294.1-5061-Aspergillus niger-Eurotiomycetes
GAO82730.1-91492-Aspergillus udagawae-Eurotiomycetes
GAQ09606.1-293939-Aspergillus lentulus-Eurotiomycetes
OJI96179.1-1036611-Aspergillus versicolor CB-Eurotiomycetes
OJJ62072.1-1036612-Aspergillus sydowii CBS 5-Eurotiomycetes
XP 002145015.1-441960-Talaromyces marneffei ATC-Eurotiomycetes
XP 002340368.1-441959-Talaromyces stipitatus AT-Eurotiomycetes
XP 018063839.1-149040-Phialocephala scopiformis-Leotiomycetes
CZR63120.1-576137-Phialocephala subalpina-Leotiomycetes
OJZ86812.1-1137211-Aspergillus luchuensis CB-Eurotiomycetes
GAT24535.1-1069201-Aspergillus luchuensis-Eurotiomycetes
GAQ46565.1-5061-Aspergillus niger-Eurotiomycetes
OJI83705.1-767770-Aspergillus tubingensis C-Eurotiomycetes
OQE32221.1-303698-Penicillium steckii-Eurotiomycetes
OKP10497.1-1316194-Penicillium subrubescens-Eurotiomycetes
OOQ83193.1-104259-Penicillium brasilianum-Eurotiomycetes
CEJ58479.1-104259-Penicillium brasilianum-Eurotiomycetes
OQD90155.1-60172-Penicillium solitum-Eurotiomycetes
CRL25978.1-5075-Penicillium camemberti-Eurotiomycetes
OQE32653.1-254877-Penicillium flavigenum-Eurotiomycetes
OQE44909.1-36646-Penicillium coprophilum-Eurotiomycetes
CDM30815.1-1365484-Penicillium roqueforti FM-Eurotiomycetes
KUM63588.1-48697-Penicillium freii-Eurotiomycetes
OQD64120.1-60169-Penicillium polonicum-Eurotiomycetes
EXU97138.1-568076-Metarhizium robertsii-Sordariomycetes
KID85732.1-1276136-Metarhizium guizhouense A-Sordariomycetes
XP 018690608.1-1367422-Fonsecaea erecta-Eurotiomycetes
XP 016635219.1-1442371-Fonsecaea multimorphosa C-Eurotiomycetes
XP 016254311.1-569365-Cladophialophora immunda-Eurotiomycetes
XP 013267126.1-1442369-Rhinocladiella mackenziei-Eurotiomycetes
XP 007729240.1-1182542-Capronia epimyces CBS 606-Eurotiomycetes
XP 013256604.1-1182545-Exophiala aquamarina CBS-Eurotiomycetes
XP 016225318.1-212818-Exophiala mesophila-Eurotiomycetes
XP 007740292.1-1182543-Cladophialophora psammoph-Eurotiomycetes
XP 016614130.1-1442370-Cladophialophora bantiana-Eurotiomycetes
XP 008729795.1-1279043-Cladophialophora carrioni-Eurotiomycetes
XP 007760483.1-1182544-Cladophialophora yegresii-Eurotiomycetes
KIW63865.1-5601-Phialophora americana-Eurotiomycetes
XP 013285935.1-1442368-Fonsecaea pedrosoi CBS 27-Eurotiomycetes
OAG41568.1-254056-Fonsecaea monophora-Eurotiomycetes
OAL34722.1-856822-Fonsecaea nubica-Eurotiomycetes
XP 013321408.1-348802-Exophiala xenobiotica-Eurotiomycetes
XP 016258065.1-215243-Exophiala oligosperma-Eurotiomycetes
XP 016233892.1-91928-Exophiala spinifera-Eurotiomycetes

S6E

Tree scale: 0.1 ⊢─────┤

Taxonomy
- Leotiomycetes
- Sordariomycetes
- Eurotiomycetes
- Dothideomycetes
- other_Pezizomycotina
- other_Ascomycota
- other_Fungi
- other_Opisthokonta
- other_Eukaryota
- Bacteria
- Archaea
- Viruses

EGC42541.1-544711-Histoplasma capsulatum H8-Eurotiomycetes
EEH02664.1-447093-Histoplasma capsulatum G1-Eurotiomycetes
OOF98127.1-602072-Aspergillus carbonarius I-Eurotiomycetes
OCK93825.1-794803-Cenococcum geophilum 1 58-Dothideomycetes
CDM31411.1-1365484-Penicillium roqueforti FM-Eurotiomycetes
OQE22710.1-254877-Penicillium flavigenum-Eurotiomycetes
XP 008082817.1-1116229-Glarea lozoyensis ATCC 20-Leotiomycetes
XP 007581912.1-1287680-Neofusicoccum parvum UCRN-Dothideomycetes
XP 001942161.1-426418-Pyrenophora tritici repen-Dothideomycetes
XP 001904308.1-515849-Podospora anserina S mat-Sordariomycetes
XP 001216793.1-341663-Aspergillus terreus NIH26-Eurotiomycetes
OJJ30102.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
KIA75578.1-40382-Aspergillus ustus-Eurotiomycetes
XP 001799912.1-321614-Parastagonospora nodorum-Dothideomycetes
KDN65693.1-1173701-Colletotrichum sublineola-Sordariomycetes
KXH46496.1-1460502-Colletotrichum nymphaeae-Sordariomycetes
XP 007598704.1-1445577-Colletotrichum fioriniae-Sordariomycetes
OCK87084.1-794803-Cenococcum geophilum 1 58-Dothideomycetes
XP 014539096.1-1170230-Penicillium digitatum Pd1-Eurotiomycetes
XP 001258284.1-331117-Aspergillus fischeri NRRL-Eurotiomycetes
OKP11762.1-1316194-Penicillium subrubescens-Eurotiomycetes
SLM33859.1-136370-Umbilicaria pustulata-other Pezizomycotina
OCL02964.1-574774-Glonium stellatum-Dothideomycetes
XP 007919395.1-1286976-Phaeoacremonium minimum U-Sordariomycetes
OAL42975.1-765867-Pyrenochaeta sp DS3sAY3a-Dothideomycetes
CDP29883.1-515849-Podospora anserina S mat-Sordariomycetes
XP 001904579.1-515849-Podospora anserina S mat-Sordariomycetes
EHK97193.1-1104152-Glarea lozoyensis 74030-Leotiomycetes
XP 008086612.1-1116229-Glarea lozoyensis ATCC 20-Leotiomycetes
OTB10641.1-1001832-Daldinia sp EC12-Sordariomycetes
OAK94597.1-765868-Stagonospora sp SRC1lsM3-Dothideomycetes
CRG83422.1-28573-Talaromyces islandicus-Eurotiomycetes
KUL90105.1-198730-Talaromyces verruculosus-Eurotiomycetes
OTB20523.1-1001832-Daldinia sp EC12-Sordariomycetes
XP 018150745.1-759273-Colletotrichum higginsian-Sordariomycetes
XP 003041774.1-660122-Nectria haematococca mpVI-Sordariomycetes
ENH66034.1-1229664-Fusarium oxysporum f sp-Sordariomycetes
OJJ41603.1-1073089-Aspergillus wentii DTO 13-Eurotiomycetes
OQD95983.1-29845-Penicillium vulpinum-Eurotiomycetes
XP 016597047.1-27334-Penicillium expansum-Eurotiomycetes
KGO67343.1-27334-Penicillium expansum-Eurotiomycetes
KGO38831.1-27334-Penicillium expansum-Eurotiomycetes
KXX74576.1-100816-Madurella mycetomatis-Sordariomycetes
KZL82819.1-1573173-Colletotrichum incanum-Sordariomycetes
OHW95298.1-1573173-Colletotrichum incanum-Sordariomycetes
OLN85402.1-708187-Colletotrichum chlorophyt-Sordariomycetes
ENH79513.1-1213857-Colletotrichum orbiculare-Sordariomycetes
ODM21321.1-573508-Aspergillus cristatus-Eurotiomycetes
XP 018189396.1-1328760-Xylona heveae TC161-other Pezizomycotina
OJJ67833.1-767769-Aspergillus brasiliensis-Eurotiomycetes
OJI89152.1-767770-Aspergillus tubingensis C-Eurotiomycetes
XP 016640358.1-563466-Scedosporium apiospermum-Sordariomycetes
KJZ71022.1-1043627-Hirsutella minnesotensis-Sordariomycetes
XP 014540751.1-1276141-Metarhizium brunneum ARSE-Sordariomycetes
XP 013956168.1-413071-Trichoderma virens Gv29 8-Sordariomycetes
XP 008097951.1-645133-Colletotrichum graminicol-Sordariomycetes
XP 007278670.1-1213859-Colletotrichum gloeospori-Sordariomycetes
XP 007789027.1-1287681-Eutypa lata UCREL1-Sordariomycetes
XP 007828360.1-1229662-Pestalotiopsis fici W106-Sordariomycetes
OTB12112.1-1001832-Daldinia sp EC12-Sordariomycetes
OTB02208.1-1001833-Hypoxylon sp CI 4A-Sordariomycetes
OTA83804.1-1001938-Hypoxylon sp CO27 5-Sordariomycetes
OTA56259.1-1001937-Hypoxylon sp EC38-Sordariomycetes
XP 003665637.1-573729-Thermothelomyces thermoph-Sordariomycetes
OTA99530.1-1001833-Hypoxylon sp CI 4A-Sordariomycetes
KFY42107.1-1420907-Pseudogymnoascus sp VKM-Leotiomycetes
KOS21700.1-150374-Escovopsis weberi-Sordariomycetes
XP 001803246.1-321614-Parastagonospora nodorum-Dothideomycetes
QUERY-Afu1g01010-330879-Aspergillus fumigatus Af2-Eurotiomycetes
KMK54471.1-1437362-Aspergillus fumigatus Z5-Eurotiomycetes
KPM42830.1-78410-Neonectria ditissima-Sordariomycetes
KIM99782.1-913774-Oidiodendron maius Zn-Leotiomycetes
OQE27687.1-303698-Penicillium steckii-Eurotiomycetes
ALQ32841.1-1147111-Fusarium euwallaceae-Sordariomycetes
XP 003040326.1-660122-Nectria haematococca mpVI-Sordariomycetes
CEG03370.1-1318461-Fusarium acuminatum CS590-Sordariomycetes
ALQ32810.1-249398-Fusarium commune-Sordariomycetes
XP 018254836.1-426428-Fusarium oxysporum f sp-Sordariomycetes
EGU72765.1-660025-Fusarium oxysporum Fo5176-Sordariomycetes
XP 018256513.1-426428-Fusarium oxysporum f sp-Sordariomycetes
EXK26726.1-1089452-Fusarium oxysporum f sp-Sordariomycetes
KIL88133.1-40199-Fusarium avenaceum-Sordariomycetes
ALQ32988.1-1101274-Fusarium sp NRRL 52700-Sordariomycetes
ALQ32947.1-42676-Fusarium sacchari-Sordariomycetes
ALQ33027.1-42677-Fusarium subglutinans-Sordariomycetes
ALQ32867.1-282569-Fusarium gaditjirri-Sordariomycetes
ALQ32976.1-70751-Fusarium sp NRRL 25184-Sordariomycetes
ALQ32924.1-48865-Fusarium redolens-Sordariomycetes
XP 018761955.1-334819-Fusarium verticillioides-Sordariomycetes
AAR92219.1-117187-Fusarium verticillioides-Sordariomycetes
ALQ32798.1-79018-Fusarium bulbicola-Sordariomycetes
ALQ32745.1-48485-Fusarium anthophilum-Sordariomycetes
ALQ33019.1-42666-Fusarium succisae-Sordariomycetes
ALQ32886.1-75915-Fusarium miscanthi-Sordariomycetes
CVK94934.1-192010-Fusarium mangiferae-Sordariomycetes
KLO90557.1-5127-Fusarium fujikuroi-Sordariomycetes
KLP05512.1-5127-Fusarium fujikuroi-Sordariomycetes
KLP14354.1-5127-Fusarium fujikuroi-Sordariomycetes
CCT74300.1-1279085-Fusarium fujikuroi IMI 58-Sordariomycetes

Figure S5. Two alleles of the idiomorphic SM gene cluster 10 present in one strain each.
A) This allele contains an insertion of genes from Chromosome 6 immediately upstream of Allele C
(see main text Figure 4). None of these genes are likely SM gene cluster backbone genes. An additional
transposable element is found flanking this insertion. B) This allele contains an insertion of genes present in
the A1163 reference but not in the Af293 reference in the middle of Allele A (see main text Figure 4). None of these genes
are likely SM gene cluster backbone genes. One additional transposable element is contained in this insertion.

Figure S7. Genomic contexts of mobile SM gene cluster 5. (A) This gene cluster is located on Chromosome 5 in two strains, on Chromosome 4 in two strains, and on Chromosome 1 in one strain adjacent to a putative SM gene cluster. Multiple transposable elements flank the cluster in each strain. (B) Strains containing this cluster marked on SNP based phylogeny.

Figure S1. Marker gene phylogenies of all strains and *Aspergillus fischerianus*.
(A) Phylogeny of beta tubulin gene. (B) Phylogeny of calmodulin gene.