Distribution and inheritance of a gene cluster encoding a

sulfated tyrosine peptide in *Xanthomonas* spp.

Running title: *raxX-raxSTAB* gene cluster in *Xanthomonas* spp.

Megan McDonald,[*,1] **Benjamin Schwessinger,**[*,1,2]
Valley Stewart,[†,3] **and Pamela C. Ronald**[†,2,4]

* These authors contributed equally to this work

The authors declare no conflict of interest. Supported by grants from NIH and NSF.

[1] Research School of Biology, Australian National University, Canberra

[2] Department of Plant Pathology and the Genome Center, University of California, Davis

[3] Department of Microbiology & Molecular Genetics, University of California, Davis

[4] Joint Bioenergy Institute, Emeryville, California

[†] corresponding authors:

Valley Stewart	Pamela C. Ronald
Dpt. Microbiology & Molecular Genetics	Department of Plant Pathology
University of California, Davis	University of California, Davis
One Shields Avenue	One Shields Avenue
Davis CA 95616-8665 USA	Davis CA 95616-8665 USA
Email: vjstewart@ucdavis.edu	Email: pcronald@ucdavis.edu
Office: (530) 754-7994	Office: (530) 752-1654
Fax: (530) 754-8973	Fax: (530) 752-5674

ABSTRACT

Tyrosine sulfation is a post-translational modification that influences interaction

specificity between certain receptors and their protein ligands in diverse biological

processes. For example, rice XA21 receptor-mediated recognition of the sulfated

bacterial protein RaxX activates an immune response and triggers resistance to the

phytopathogen *Xanthomonas oryzae* pv. *oryzae* (*Xoo*). A five kb *raxX-raxSTAB* gene

cluster of *Xoo* encodes RaxX, the RaxST tyrosylprotein sulfotransferasea and the RaxA

and RaxB components of a predicted proteolytic maturation and ATP-dependent

peptide secretion complex. The complete *raxX-raxSTAB* gene cluster was found only in

Xanthomonas species, and its distribution is consistent with multiple gain and loss

events during *Xanthomonas* speciation. Homologs of the *raxST* gene are present in

genome sequences of diverse bacterial species. Together, these results establish a

foundation for investigating biological roles for tyrosine sulfation in bacteria.

ABSTRACT = 132 words (maximum 200)

INTRODUCTION

Host receptors activate innate immunity pathways upon pathogen recognition (Ronald

and Beutler 2010). The gene encoding the rice XA21 receptor kinase (Song *et al.,*

1995) confers broad spectrum resistance against the gamma-proteobacterium

Xanthomonas oryzae pv. *oryzae* (*Xoo*) (Wang *et al.,* 1996). This well-studied XA21-*Xoo*

model provides a basis from which to understand molecular and evolutionary

mechanisms of host-microbe interactions.

Several *Xoo rax* genes are <u>r</u>equired for <u>a</u>ctivation of <u>X</u>A21-mediated immunity (**Fig. 1a**).

The *raxX-raxSTAB* gene cluster encodes the 60-residue RaxX predicted precursor

protein that undergoes sulfation by the RaxST tyrosylprotein sulfotransferase at residue

Tyr-41 (Pruitt *et al.,* 2015). We hypothesize that the RaxABC proteolytic maturation and

ATP-dependent peptide secretion complex (da Silva *et al.,* 2004) further processes the

sulfated RaxX precursor by removing its double-glycine leader peptide prior to secretion

(Holland *et al.,* 2016). Located outside the *raxX-raxSTAB* gene cluster, the *raxC* gene,

an ortholog of the *tolC* gene, encodes the predicted outer membrane channel for this

complex (da Silva *et al.,* 2004). Finally, the *raxPQ* genes encode enzymes to assimilate

sulfate into 3'-phosphoadenosine 5'-phosphosulfate (Shen *et al.,* 2002), the sulfodonor

for the RaxST tyrosylprotein sulfotransferase (Han *et al.,* 2012).

Tyrosylprotein sulfotransferase is confined to the Golgi complex in both plants and

animals (Moore 2009). Thus, this post-translational modification is targeted to a subset

of cell surface and secreted proteins that influence a variety of eukaryotic physiological

processes (Matsubayashi 2014; Stone *et al.,* 2009). For example, tyrosine sulfation of

the chemokine receptors CCR5 and CXCR4 is necessary for high-affinity binding not

only to chemokines, but also to the HIV-1 surface glycoprotein (Farzan *et al.,* 1999;

Kleist *et al.,* 2016). In plants, sulfated tyrosine peptides influence xylem development,

root growth, and/or plant immune signaling (Matsubayashi 2014; Zhou *et al.,* 2017).

RaxST sulfation of RaxX residue Tyr-41 is the only example of tyrosine sulfation

reported in bacteria (da Silva *et al.,* 2004; Pruitt *et al.,* 2015). Strikingly, RaxX residues

40-52 share sequence similarity with mature active **p**lant peptide containing **s**ulfated

t**y**rosine (PSY) hormones (Amano *et al.,* 2007; Pruitt *et al.,* 2015; Pruitt *et al.,* 2017).

Indeed, RaxX, like PSY1, can enhance root growth in diverse plant species (Pruitt *et al.,*

2017). Moreover, RaxX also contributes to *Xoo* virulence in the absence of the XA21

immune receptor (Pruitt *et al.,* 2017). This apparent hormone mimickry by RaxX

therefore may serve broad functions in *Xoo* pathogenesis. To further elucidate the

biological role of bacterial tyrosine sulfation, we sought to identify the species

distribution and possible origin of genes in the *raxX-raxSTAB* gene cluster.

Here we show that the *raxX-raxSTAB* gene cluster is confined to a subset of

Xanthomonas species. In all cases examined, the *raxX-raxSTAB* gene cluster lies

between two core (housekeeping) genes, *gcvP* encoding a subunit of glycine

dehydrogenase, and a gene encoding a major facilitator subfamily transporter (*"mfsX"*).

Examination of nucleotide sequence conservation across the *raxX-raxSTAB* gene

cluster, and at its boundaries with the *gcvP* and *"mfsX"* genes, suggests that the *raxX-*

raxSTAB gene cluster was acquired through lateral transfer by *X. translucens*, a

pathogen of diverse cereal species (Langlois *et al.,* 2017), and separately by *X.*

maliensis, associated with but nonpathogenic for rice (Triplett *et al.,* 2015). Finally,

genes homologous to *raxST* are present in bacterial genomes from a wide range of

species, raising the possibility that RaxST-catalyzed tyrosine sulfation may occur in

other genomic and biological contexts in addition to RaxX.

RESULTS

The *raxX-raxSTAB* gene cluster is present in a subset of *Xanthomonas* spp.

We searched databases at the National Center for Biotechnology Information to identify

bacterial genomes with the *raxX-raxSTAB* gene cluster. We found the *raxX-raxSTAB*

gene cluster exclusively in *Xanthomonas* spp., and ultimately detected it in more than

200 unique genome sequences among 413 accessed through the RefSeq database

(O'Leary *et al.,* 2016).

Xanthomonas taxonomy has undergone substantial changes over the years (Vauterin *et*

al., 2000; Young 2008); see (Midha and Patil 2014) for a representative example). At

one point, many strains were denoted as pathovars of either *X. campestris* or *X.*

axonopodis, but today over 20 species are distinguished, many with multiple pathovars

(Rademaker *et al.,* 2005; Vauterin *et al.,* 1995). Because many of the genome

sequences we examined are from closely-related strains, in some cases associated

with different species designations, we constructed a phylogenetic tree in order to

organize these sequences by relatedness (**Fig. S1**).

The phylogenetic relationships among *Xanthomonas* spp. was assessed using the

entire genome assembly with Andi v0.10 (Haubold *et al.,* 2015; Klotzl and Haubold

2016). We compared our topology with several other *Xanthomonas* phylogenetic trees

published previously (Ferreira-Tonin *et al.,* 2012; Gardiner *et al.,* 2014; Hauben *et al.,*

1997; Midha and Patil 2014; Parkinson *et al.,* 2007; Parkinson *et al.,* 2009; Rademaker

et al., 2005; Triplett *et al.,* 2015; Young *et al.,* 2008). Most share broad similarity with

each other and with the whole-genome tree presented here in defining relationships

between well-sampled species. To facilitate discussion, we represent our phylogenetic

tree as a cladogram (**Fig. 2**)**.**

We detected the *raxX-raxSTAB* gene cluster in six lineages that consistently are

identified as being distinct from one another (Rademaker *et al.,* 2005; Vauterin *et al.,*

1995) (**Fig. 2**). One lineage includes the two *X. oryzae* pathovars, *oryzae* and *oryzicola*,

pathogenic on rice (Niño-Liu *et al.,* 2006). A second lineage includes *X. vasicola*,

strains of which are pathogenic on sugarcane, sorghum or maize, together with strains

denoted as *X. campestris* pv. *musacearum*, pathogenic on banana (Aritua *et al.,* 2008).

The third lineage includes *X. euvesicatoria* and *X. perforans*, pathogenic on pepper and

tomato (Potnis *et al.,* 2015), together with strains denoted as *X. alfalfa* subsp.

citrumelonis (pathogenic on citrus) and *X. dieffenbachiae* (anthuriums) (Rademaker

group 9.2; (Barak *et al.,* 2016; Rademaker *et al.,* 2005). The fourth lineage includes

strains denoted as *X. axonopodis* pathovars *manihotis* (pathogenic on cassava) and

phaseoli (bean) (Rademaker group 9.4; (Mhedbi-Hajri *et al.,* 2013; Rademaker *et al.,*

2005).

The fifth lineage includes *X. translucens*, different strains of which are pathogenic on

one or more cereal crops such as wheat and barley, and/or non-cereal forage and

turfgrass species (Langlois *et al.,* 2017). *X. translucens* is within the distinct cluster of

"early-branching" species whose divergence from the remainder apparently occurred

relatively early during *Xanthomonas* speciation (Parkinson *et al.,* 2007). The sixth

lineage comprises *X. maliensis*, associated with but nonpathogenic on rice (Triplett *et*

al., 2011); phylogenetic analyses place this species between the "early-branching"

species and the remainder (Triplett *et al.,* 2015).

Notably, the *raxX-raxSTAB* gene cluster is absent from the *X. citri* pathovar group,

pathogenic on a range of dicots including citrus. This group, which includes certain

strains denoted as *X. axonopodis* or *X. campestris* (Bansal *et al.,* 2017), clusters

phylogenetically among the *X. oryzae*, *X. euvesicatoria* and *X. axonopodis* pv.

manihotis groups (Midha and Patil 2014; Rademaker *et al.,* 2005; Vauterin *et al.,* 1995)

(**Fig. 2**).

Together, these observations suggest that the *raxX-raxSTAB* gene cluster experienced

multiple gains and/or losses during *Xanthomonas* speciation.

Sequence conservation of the *raxX-raxSTAB* gene cluster suggests lateral transfer between *Xanthomonas* spp.

From the initial analysis described above, we selected 15 species, representing the

phylogenetic range of *Xanthomonas*, for more detailed analyses of *rax* gene cluster

composition, organization, and inheritance (**Fig. 2; Table 1**). The corresponding

genome sequences are accompanied by published descriptions (**Table 1**). The close

relative *Stenotrophomonas maltophilia*, which does not contain the *raxX-raxSTAB* gene

cluster, serves as a reference (Moore *et al.,* 1997).

Both the organization and size of the *raxX-raxSTAB* gene cluster are conserved across

all six lineages in which it resides. To address hypotheses for patterns of *raxX-*

raxSTAB gene cluster inheritance, we compared individual phylogenetic trees for each

of the four *rax* genes to the overall *Xanthomona*s phylogenetic tree (**Fig. 3**) (Kuo and

Ochman 2009). For all four genes, sequences in *X. translucens*, in the early-branching

group, cluster separately from sequences in the other lineages. This finding is

congruent with the hypothesis, that *X. translucens* acquired the *raxX-raxSTAB* gene

cluster relatively early during *Xanthomonas* speciation. For *X. maliensis*, the *raxX-*

raxSTAB genes assort among those from *X. euvesicatoria* and the *X. axonopodis*

pathovars *manihotis* and *phaseoli* (**Fig. 3**)*,* even though the *X. maliensis* genome

sequence itself is more distantly related (**Fig. 2**). This finding suggests that *X. maliensis*

acquired the *raxX-raxSTAB* gene cluster relatively late during *Xanthomonas* speciation.

The *raxX-raxSTAB* gene cluster lies between two core (housekeeping) genes (**Fig. 1a**).

One, *gcvP*, encodes the pyridoxal-phosphate subunit of glycine dehydrogenase. An

approximately 170 nt riboswitch (*gcvR* in **Fig. 1a**) controls GcvP protein synthesis in

response to glycine (Mandal *et al.,* 2004). The other, "*mfsX*", encodes a **m**ajor

facilitator **s**ubfamily (MFS) transporter related to Bcr and CflA efflux proteins (da Silva *et*

al., 2004). Here, "*mfsX*" is only a provisional designation absent functional

characterization.

We further examined phylogenetic relationships by comparing nucleotide sequence

identity across the *gcvP, raxX, raxST, raxA, raxB* and *"mfsX"* coding regions, each from

the initiation through termination codon (**Table 2**). For comparison, values are

presented also for **g**enome-wide **a**verage **n**ucleotide **i**dentity (gANI) as well as the

alignment **f**raction (AF), which estimates the fraction of orthologous genes (Varghese *et*

al., 2015). For context, a widely-used criterion assigns 95% **a**verage **n**ucleotide **i**dentity

(ANI) as the cut-off point for species delineation (Goris *et al.,* 2007). Sequence from *X.*

euvesicatoria is the reference.

The *raxST*, *raxA* and *raxB* coding sequences from *X. axonopodis* pv. *manihotis* and *X.*

maliensis display the highest identity to those from *X. euvesicatoria*, at least 95% in

each case. Sequences from *Xoo* and *X. vasicola* (also known as *X. campestris* pv.

malvacearum) are about 90% identical, and those from *X. translucens* roughly 75%

identical (**Table 2**). The *raxX* coding sequences are more divergent, with identity to the

X. euvesicatoria sequence ranging from almost 90% for *X. maliensis* and *Xoo* to only

63% for *X. translucens* (**Table 2**).

Boundaries flanking the *raxX-raxSTAB* gene cluster and adjacent genes suggest lateral transfer through general recombination

Comparison of the *gcvP* - *[raxX-raxSTAB]* - "*mfsX*" region from all 16 reference species

reveals sharp boundaries flanking the position of the *raxX-raxSTAB* gene cluster. On

the left flank, substantial nucleotide identity spans the *gcvP* gene, the *gcvR* riboswitch,

and a presumptive promoter –10 element (Mitchell *et al.,* 2003) (**Fig. 1b**). On the right

flank, identity begins shortly after the "*mfsX*" initiation codon. Accordingly, upstream

signals for "*mfsX*" gene transcription (Mitchell *et al.,* 2003) and translation (Ma *et al.,*

2002) are conserved within, but not between, *raxX-raxSTAB*-positive and -negative

sequences (**Fig. 1b**).

Between these boundaries in *raxX-raxSTAB* gene cluster-negative species, the

compact (≤ 200 nt) *gcvP*-"*mfsX*" intergenic sequence is modestly conserved in most

genomes (about 60-80% overall identity) (**Fig. 1b**). Much of this identity comes from the

"*mfsX*" potential transcription and translation initiation sequences described above. The

overall intergenic sequence is less conserved in the early-branching species (*X.*

albilineans, *X. hyacinthi* and *X. sacchari*), displaying about 50-65% overall identity.

In *raxX-raxSTAB* gene cluster-positive genomes, sequence flanking these boundaries

appears unrelated to the *gcvP*-"*mfsX*" intergenic sequence from *raxX-raxSTAB* gene

cluster-negative genomes (**Fig. 1b**). Rather, it is well-conserved even in the early-

branching species, *X. translucens*. These results suggest that anomolous *raxX-*

raxSTAB gene cluster phylogenetic distribution results from lateral gene transfer.

Sequences of the adjacent *gcvP* gene display length polymorphisms (**Fig. 4**) that do not

align with overall *Xanthomonas* species relationships (**Fig. 2**). Polymorphisms of this

type are unusual, and indicate recombination (Nelson *et al.,* 1997). Their occurrence in

a gene adjacent to the *raxX-raxSTAB* gene cluster independently supports the model,

that this genomic region evolves through lateral gene transfer.

raxST homologs are present in genomes of diverse bacterial species

As we searched genome sequences available through GenBank for evidence of the

raxX-raxSTAB gene cluster outside of *Xanthomonas* spp., we identified sequences

encoding proteins with about 40% identity to, and approximately the same length as, the

Xoo RaxST protein. Sequence identity is high in residues that form the binding pocket

for the cofactor, 3'-phosphoadenosine 5'-phosphosulfate (da Silva *et al.,* 2004; Kakuta

et al., 1998), consistent with assignment of these encoded proteins as

sulfotransferases. It is not known if these genes encoding tyrosylprotein

sulfotransferases, as there are no defined sequence features that distinguish such

enzymes from other sulfotransferases that have non-protein substrates (Dong *et al.,*

2012; Teramoto *et al.,* 2013).

These *raxST* homologs are in a range of bacterial phyla including Proteobacteria and

Cyanobacteria (**Fig. 5**). Nevertheless, for most species represented by multiple

genome sequences, the *raxST* homolog was detected in a minority of individuals, so it

is not part of the core genome in these strains. Moreover, relationships between

species in a *raxST* gene phylogenetic tree bear no resemblance to those in the overall

tree of bacterial species. For example, in the *raxST* gene tree, sequences from

Cyanobacteria are flanked on both sides by sequences from Proteobacteria (**Fig. 5**).

Together, these findings provide evidence for lateral transfer of *raxST* homologs

transfer (Kuo and Ochman 2009).

DISCUSSION

We hypothesize that the *raxX-raxSTAB* gene cluster originated in an ancestor to the

lineage containing *X. oryzae*, *X. euvesicatoria*, and related species, with further gains or

loss through lateral transfer as described below (**Fig. 2**). Analysis suggests that

relatively few events were necessary to form the *raxX-raxSTAB* gene cluster. The

raxAB genes are homologous to those encoding proteolytic maturation and ATP-

dependent peptide secretion complexes (da Silva *et al.,* 2004; Lin *et al.,* 2015), related

to type 1 secretion systems but specialized for secreting small peptides such as

bacteriocins and peptide pheromones (Holland *et al.,* 2016). Frequently, the gene

encoding the secreted substrate is adjacent to genes encoding components of the

secretion complex (Dirix *et al.,* 2004). The *raxX* gene therefore might have evolved

from the gene for the seccreted peptide substrate of the RaxAB ancestor. Finally, as

we show here, homologs for the *raxST* gene are distributed broadly (**Fig. 5**).

The *raxX-raxSTAB* gene cluster does not exhibit features, such as a gene for a site-

specific recombinase, characteristic of self-mobile genomic islands (Hacker *et al.,*

1997). Moreover, variant-length alleles of the adjacent *gcvP* gene (**Fig. 4**) provide

evidence for general recombination in the vicinity (Nelson *et al.,* 1997). Thus, the

simplest model for *raxX-STAB* gene cluster lateral transfer is that it occurred through

general recombination between genes flanking each side of the *raxX-STAB* gene

cluster (**Fig. 1b**).

Three examples provide further evidence for lateral transfer. First, the *raxX-raxSTAB*

gene cluster from the early branching species *X. translucens* has essentially the same

size, composition and structure as the others. However, the *X. translucens raxX-*

raxSTAB sequences are more divergent (**Table 2**). This predicts that the *raxX-raxSTAB*

gene cluster has been part of the *X. translucens* genome sufficiently long for sequence

alterations to accumulate (Kuo and Ochman 2009).

In the second example of evidence for lateral transfer, the *X. maliensis raxSTAB*

sequences share strong similarity to those of *X. euvesicatoria* and the *X. axonopodis*

pathovars *manihotis* and *phaseoli*, whereas their genome sequences are more

divergent (**Table 2**). This suggests that *X. maliensis* acquired the *raxX-raxSTAB* gene

cluster relatively recently (Kuo and Ochman 2009).

The final example of evidence for lateral transfer considers apparent loss of the *raxX-*

raxSTAB gene cluster during differentiation of *X. citri* from the large group including

Xoo, X. euvesicatoria and related species (Midha and Patil 2014; Rademaker *et al.,*

2005; Vauterin *et al.,* 1995) (**Fig. 2**). Sequence in the *gcvP*-"*mfsX*" intergenic region is

conserved among *raxX-raxSTAB* gene cluster-negative species, including *X. citri* (**Fig.**

1b). This is consistent with loss from the *X. citri* lineage mediated by recombination,

with the *gcvP*-"*mfsX*" region from a *raxX-raxSTAB* gene cluster-negative species. The

result of this recombination would be replacement of the *raxX-raxSTAB* gene cluster

with a conserved *gcvP*-"*mfsX*" region.

In an alternative scenario, where the *X. citri raxX-raxSTAB* gene cluster was lost

through deletion, the remaining sequence in the intergenic region would more closely

resemble the *raxX-raxSTAB* gene cluster-positive boundary sequence. This conclusion

is not supported by the intergenic region found in *X. citri*. A second alternative scenario,

that the *raxX-raxSTAB* gene cluster formed after *X. citri* speciation, is not supported by

the analysis of *X. translucens* sequences described above.

Broad phylogenetic distribution of the *raxX-raxSTAB* gene cluster implies that its

associated phenotypes can contribute to host interactions with diverse *Xanthomonas*

spp. The *raxX-raxSTAB* gene cluster was identified in the context of the rice XA21-

mediated immune response (da Silva *et al.,* 2004; Pruitt *et al.,* 2015), but the sequence

and functional similarities between the bacterial RaxX and the plant PSY sulfopeptides

suggests that RaxX may mimic PSY phytohormone activies to facilitate infection (Pruitt

et al., 2017). Indeed, *Xoo* strains that cannot synthesize sulfated RaxX exhibit reduced

virulence (Pruitt *et al.,* 2017).

Evidence for lateral transfer to *X. translucens* and *X. maliensis* suggests that *raxX-*

raxSTAB gene cluster acquisition may contribute to emergence of new species or

pathovars. The potentially useful phenotype of PSY hormone mimickry conceivably

could introduce a particular strain to previously inaccessible hosts or niches. On the

other hand, loss of the *raxX-raxSTAB* gene cluster apparently occurred during formation

of the *X. citri* lineage (**Fig. 2**), perhaps indicating that the *raxX-raxSTAB* gene cluster did

not enhance fitness in this case.

Pathovar phenotypes that differentiate bacterium-plant interactions, characterized

extensively in members of the genus *Xanthomonas* (Jacques *et al.,* 2016), are not

predicted by the presence or absence of the *raxX-raxSTAB* gene cluster. Some

species that infect monocots exclusively contain the *raxX-raxSTAB* gene cluster (e.g.,

X. oryzae, X. translucens), whereas others do not (e.g., *X. arboricola, X. hyacinthi*).

Likewise, some species that infect dicots exclusively contain the *raxX-raxSTAB* gene

cluster (e.g., *X. euvesicatoria*), whereas others do not (e.g., *X. campestris* pv.

campestris; *X. citri*). Similarly, there is no association with tissue specificity; for

example, a single species contains both vascular (*X. oryzae* pv. *oryzae*) and

nonvascular (*X. oryzae* pv. *oryzicola*) pathogens.

MATERIALS AND METHODS

Survey of *raxX-STAB* gene clusters in *Xanthomonas* spp.

All available *Xanthomonas* genomes were downloaded from the NCBI ftp server on

January 29, 2016 (413 genome accessions). The genome fasta files were used to build

a local blast database using BLASTv2.27+ (Camacho *et al.,* 2009). For all genes in and

surrounding the *raxSTAB* operon blastn (evalue cutoff of 1e-3) was used to identify

homologs in the local blast database. Due to the small size of RaxX, tblastn was

required to identify homologs (evalue cutoff of 1e-3). Fasta files for each blast hit were

generated using a custom python script (available upon request). Alignments of all

genes were performed with Muscle v3.5 (Edgar 2004) implemented in the desktop tool

Geneious v9.1.8 (Kearse *et al.,* 2012). Alignment ends were trimmed so that each

sequence was equal in length and in the first coding frame. Maximum likelihood trees

were built with RaxML v8.2.4 (Stamatakis 2014) with the following settings: (-m

GTRGAMMA F -f a -x 3298589 -N 10000 -p 23). Trees shown in all figures are the

highest scoring ML tree and numbers shown on branches are the resampled bootstrap

values from 1000 replicates. Trees were drawn in FigTree v1.4.0

(http://tree.bio.ed.ac.uk/software/figtree/).

Whole genome phylogenies were generated using the entire genome assembly with the

program Andi v0.10 (Haubold *et al.,* 2015; Klotzl and Haubold 2016). These distance

matrices were plotted as neighbor-joining tree using Phylip v3.695 (Felsenstein 1981).

Numbers on the branches represent the proportion (0-100) that the branch appeared in

the "bootstrapped" distance matrices using Andi.

Sequence analyses

Nucleotide and deduced amino acid sequences were edited and analyzed with the

programs EditSeq^TM (version 14.1.0), MegAlign^TM (version 14.1.0) and SeqBuilder^TM

(version 14.1.0), DNASTAR, Madison, WI. The Integrated Microbial Genomes interface

(Chen *et al.,* 2017) was used to compare genome segments from different species, and

also to extract values for genome-wide Average Nucleotide Identity and genome

Alignment Fraction (Varghese *et al.,* 2015).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGMENTS

Research reported in this publication was supported by the National Institute of General Medical Sciences of the National Institutes of Health under award number R01GM59962, and by the Division of Integrative Organismal Systems of the National Science Foundation under award number IOS-1237975. B.S. is supported by a Discovery Early Career Research Award from the Australian Research Council under award number DE150101897.

REFERENCES

Amano Y, Tsubouchi H, Shinohara H, Ogawa M, Matsubayashi Y (2007). Tyrosine-sulfated glycopeptide involved in cellular proliferation and expansion in *Arabidopsis*. *Proc Natl Acad Sci U S A* **104:** 18333-18338.

Aritua V, Parkinson N, Thwaites R, Heeney JV, Jones DR, Tushemereirwe W *et al* (2008). Characterization of the *Xanthomonas* sp. causing wilt of enset and banana and its proposed reclassification as a strain of *X. vasicola*. *Plant Pathol* **57:** 170-177.

Bansal K, Midha S, Kumar S, Patil PB (2017). Ecological and evolutionary insights into pathovar diversity of Xanthomonas citri. *Appl Environ Microbiol* **83:** e02993-02916.

Barak JD, Vancheva T, Lefeuvre P, Jones JB, Timilsina S, Minsavage GV *et al* (2016). Whole-genome sequences of *Xanthomonas euvesicatoria* strains clarify taxonomy and reveal a stepwise erosion of Type 3 effectors. *Front Plant Sci* **7:** 1805.

Bart R, Cohn M, Kassen A, McCallum EJ, Shybut M, Petriello A *et al* (2012). High-throughput genomic sequencing of cassava bacterial blight strains identifies conserved effectors to target for durable resistance. *Proc Natl Acad Sci U S A* **109:** E1972-1979.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K *et al* (2009). BLAST+: architecture and applications. *BMC Bioinformatics* **10:** 421.

Chen IA, Markowitz VM, Chu K, Palaniappan K, Szeto E, Pillay M *et al* (2017). IMG/M: integrated genome and metagenome comparative data analysis system. *Nucleic Acids Res* **45:** D507-D516.

Crossman LC, Gould VC, Dow JM, Vernikos GS, Okazaki A, Sebaihia M *et al* (2008). The complete genome, comparative and functional analysis of *Stenotrophomonas maltophilia* reveals an organism heavily shielded by drug resistance determinants. *Genome Biol* **9:** R74.

da Silva AC, Ferro JA, Reinach FC, Farah CS, Furlan LR, Quaggio RB *et al* (2002). Comparison of the genomes of two *Xanthomonas* pathogens with differing host specificities. *Nature* **417:** 459-463.

da Silva FG, Shen Y, Dardick C, Burdman S, Yadav RC, de Leon AL *et al* (2004). Bacterial genes involved in type I secretion and sulfation are required to elicit the rice Xa21-mediated innate immune response. *Mol Plant Microbe Interact* **17:** 593-601.

Dirix G, Monsieurs P, Dombrecht B, Daniels R, Marchal K, Vanderleyden J *et al* (2004). Peptide signal molecules and bacteriocins in Gram-negative bacteria: a genome-wide in silico screening for peptides containing a double-glycine leader sequence and their cognate transporters. *Peptides* **25:** 1425-1440.

Dong D, Ako R, Wu B (2012). Crystal structures of human sulfotransferases: insights into the mechanisms of action and substrate selectivity. *Expert Opin Drug Metab Toxicol* **8:** 635-646.

Edgar RC (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32:** 1792-1797.

Farzan M, Mirzabekov T, Kolchinsky P, Wyatt R, Cayabyab M, Gerard NP *et al* (1999). Tyrosine sulfation of the amino terminus of CCR5 facilitates HIV-1 entry. *Cell* **96:** 667-676.

Felsenstein J (1981). Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* **17:** 368-376.

Ferreira-Tonin M, Rodrigues-Neto J, Harakava R, Destefano SA (2012). Phylogenetic analysis of *Xanthomonas* based on partial *rpoB* gene sequences and species differentiation by PCR-RFLP. *Int J Syst Evol Microbiol* **62:** 1419-1424.

Gardiner DM, Upadhyaya NM, Stiller J, Ellis JG, Dodds PN, Kazan K *et al* (2014). Genomic analysis of *Xanthomonas translucens* pathogenic on wheat and barley reveals cross-kingdom gene transfer events and diverse protein delivery systems. *PLoS One* **9:** e84995.

Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM (2007). DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* **57:** 81-91.

Hacker J, Blum-Oehler G, Mühldorfer I, Tschäpe H (1997). Pathogenicity islands of virulent bacteria: structure, function and impact on microbial evolution. *Mol Microbiol* **23:** 1089-1097.

Han SW, Lee SW, Bahar O, Schwessinger B, Robinson MR, Shaw JB *et al* (2012). Tyrosine sulfation in a Gram-negative bacterium. *Nat Commun* **3:** 1153.

Hauben L, Vauterin L, Swings J, Moore ER (1997). Comparison of 16S ribosomal DNA sequences of all *Xanthomonas* species. *Int J Syst Bacteriol* **47:** 328-335.

Haubold B, Klotzl F, Pfaffelhuber P (2015). andi: fast and accurate estimation of evolutionary distances between closely related genomes. *Bioinformatics* **31:** 1169-1175.

Holland IB, Peherstorfer S, Kanonenberg K, Lenders M, Reimann S, Schmitt L (2016). Type I protein secretion-deceptively simple yet with a wide range of mechanistic variability across the family. *EcoSal Plus* **7:** doi:10.1128/ecosalplus.ESP-0019-2015.

Jacobs JM, Pesce C, Lefeuvre P, Koebnik R (2015). Comparative genomics of a cannabis pathogen reveals insight into the evolution of pathogenicity in *Xanthomonas*. *Front Plant Sci* **6:** 431.

Jacques MA, Arlat M, Boulanger A, Boureau T, Carrere S, Cesbron S *et al* (2016). Using ecology, physiology, and genomics to understand host specificity in *Xanthomonas*. *Annu Rev Phytopathol* **54:** 163-187.

Kakuta Y, Pedersen LG, Pedersen LC, Negishi M (1998). Conserved structural motifs in the sulfotransferase family. *Trends Biochem Sci* **23:** 129-130.

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S *et al* (2012). Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28:** 1647-1649.

Kleist AB, Getschman AE, Ziarek JJ, Nevins AM, Gauthier PA, Chevigne A *et al* (2016). New paradigms in chemokine receptor signal transduction: Moving beyond the two-site model. *Biochem Pharmacol* **114:** 53-68.

Klotzl F, Haubold B (2016). Support values for genome phylogenies. *Life (Basel)* **6:** 11.

Kuo CH, Ochman H (2009). The fate of new bacterial genes. *FEMS Microbiol Rev* **33:** 38-43.

Langlois PA, Snelling J, Hamilton JP, Bragard C, Koebnik R, Verdier V *et al* (2017). Characterization of the *Xanthomonas translucen*s complex using draft genomes, comparative genomics, phylogenetic analysis, and diagnostic LAMP assays. *Phytopathology* **107:** 519-527.

Lin DY, Huang S, Chen J (2015). Crystal structures of a polypeptide processing and secretion transporter. *Nature* **523:** 425-430.

Ma J, Campbell A, Karlin S (2002). Correlations between Shine-Dalgarno sequences and gene features such as predicted expression levels and operon structures. *J Bacteriol* **184:** 5733-5745.

Mandal M, Lee M, Barrick JE, Weinberg Z, Emilsson GM, Ruzzo WL *et al* (2004). A glycine-dependent riboswitch that uses cooperative binding to control gene expression. *Science* **306:** 275-279.

Matsubayashi Y (2014). Posttranslationally modified small-peptide signals in plants. *Annu Rev Plant Biol* **65:** 385-413.

Mhedbi-Hajri N, Hajri A, Boureau T, Darrasse A, Durand K, Brin C *et al* (2013). Evolutionary history of the plant pathogenic bacterium *Xanthomonas axonopodis*. *PLoS One* **8:** e58474.

Midha S, Patil PB (2014). Genomic insights into the evolutionary origin of *Xanthomonas axonopodis* pv. *citri* and its ecological relatives. *Appl Environ Microbiol* **80:** 6266-6279.

Mitchell JE, Zheng D, Busby SJ, Minchin SD (2003). Identification and analysis of 'extended -10' promoters in *Escherichia coli*. *Nucleic Acids Res* **31:** 4689-4695.

Moore ER, Kruger AS, Hauben L, Seal SE, Daniels MJ, De Baere R *et al* (1997). 16S rRNA gene sequence analyses and inter- and intrageneric relationships of *Xanthomonas* species and *Stenotrophomonas maltophilia*. *FEMS Microbiol Lett* **151:** 145-153.

Moore KL (2009). Protein tyrosine sulfation: a critical posttranslation modification in plants and animals. *Proc Natl Acad Sci U S A* **106:** 14741-14742.

Naushad S, Adeolu M, Wong S, Sohail M, Schellhorn HE, Gupta RS (2015). A phylogenomic and molecular marker based taxonomic framework for the order *Xanthomonadales*: proposal to transfer the families *Algiphilacea*e and *Solimonadaceae* to the order *Nevskiales* ord. nov. and to create a new family within the order *Xanthomonadales*, the family *Rhodanobacteraceae* fam. nov., containing the genus *Rhodanobacter* and its closest relatives. *Antonie Van Leeuwenhoek* **107:** 467-485.

Nelson K, Wang FS, Boyd EF, Selander RK (1997). Size and sequence polymorphism in the isocitrate dehydrogenase kinase/phosphatase gene (*aceK*) and flanking regions in *Salmonella enterica* and *Escherichia coli*. *Genetics* **147:** 1509-1520.

Niño-Liu DO, Ronald PC, Bogdanove AJ (2006). *Xanthomonas oryzae* pathovars: model pathogens of a model crop. *Mol Plant Pathol* **7:** 303-324.

O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R *et al* (2016). Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res* **44:** D733-745.

Parkinson N, Aritua V, Heeney J, Cowie C, Bew J, Stead D (2007). Phylogenetic analysis of *Xanthomonas* species by comparison of partial gyrase B gene sequences. *Int J Syst Evol Microbiol* **57:** 2881-2887.

Parkinson N, Cowie C, Heeney J, Stead D (2009). Phylogenetic structure of *Xanthomonas* determined by comparison of *gyrB* sequences. *Int J Syst Evol Microbiol* **59:** 264-274.

Pereira UP, Gouran H, Nascimento R, Adaskaveg JE, Goulart LR, Dandekar AM (2015). Complete genome sequence of *Xanthomonas arboricola* pv. *juglandis* 417, a copper-resistant strain isolated from *Juglans regia* L. *Genome Announc* **3:** e01126-01115.

Pieretti I, Cociancich S, Bolot S, Carrere S, Morisset A, Rott P *et al* (2015). Full genome sequence analysis of two isolates reveals a novel Xanthomonas species close to the sugarcane pathogen *Xanthomonas albilineans*. *Genes (Basel)* **6:** 714-733.

Potnis N, Timilsina S, Strayer A, Shantharaj D, Barak JD, Paret ML *et al* (2015). Bacterial spot of tomato and pepper: diverse *Xanthomonas* species with a wide variety of virulence factors posing a worldwide challenge. *Mol Plant Pathol* **16:** 907-920.

Pruitt RN, Schwessinger B, Joe A, Thomas N, Liu F, Albert M *et al* (2015). The rice immune receptor XA21 recognizes a tyrosine-sulfated protein from a Gram-negative bacterium. *Sci Adv* **1:** e1500245.

Pruitt RN, Joe A, Zhang W, Feng W, Stewart V, Schwessinger B *et al* (2017). A microbially derived tyrosine-sulfated peptide mimics a plant peptide hormone. *New Phytol* **215:** in press.

Rademaker JLW, Louws FJ, Schultz MH, Rossbach U, Vauterin L, Swings J *et al* (2005). A comprehensive species to strain taxonomic framework for Xanthomonas. *Phytopathology* **95:** 1098-1111.

Ronald PC, Beutler B (2010). Plant and animal sensors of conserved microbial signatures. *Science* **330:** 1061-1064.

Salzberg SL, Sommer DD, Schatz MC, Phillippy AM, Rabinowicz PD, Tsuge S *et al* (2008). Genome sequence and rapid evolution of the rice pathogen Xanthomonas oryzae pv. oryzae PXO99A. *BMC Genomics* **9:** 204.

Shen Y, Sharma P, da Silva FG, Ronald P (2002). The *Xanthomonas oryzae* pv. *oryzae raxP* and *raxQ* genes encode an ATP sulphurylase and adenosine-5'-phosphosulphate kinase that are required for AvrXa21 avirulence activity. *Mol Microbiol* **44:** 37-48.

Song WY, Wang GL, Chen LL, Kim HS, Pi LY, Holsten T *et al* (1995). A receptor kinase-like protein encoded by the rice disease resistance gene, *Xa21*. *Science* **270:** 1804-1806.

Stamatakis A (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30:** 1312-1313.

Stone MJ, Chuang S, Hou X, Shoham M, Zhu JZ (2009). Tyrosine sulfation: an increasingly recognised post-translational modification of secreted proteins. *N Biotechnol* **25:** 299-317.

Studholme DJ, Wasukira A, Paszkiewicz K, Aritua V, Thwaites R, Smith J *et al* (2011). Draft genome sequences of Xanthomonas sacchari and two banana-associated Xanthomonads reveal insights into the *Xanthomonas* group 1 clade. *Genes (Basel)* **2:** 1050-1065.

Teramoto T, Fujikawa Y, Kawaguchi Y, Kurogi K, Soejima M, Adachi R *et al* (2013). Crystal structure of human tyrosylprotein sulfotransferase-2 reveals the mechanism of protein tyrosine sulfation reaction. *Nat Commun* **4:** 1572.

Thieme F, Koebnik R, Bekel T, Berger C, Boch J, Buttner D *et al* (2005). Insights into genome plasticity and pathogenicity of the plant pathogenic bacterium *Xanthomonas campestris* pv. *vesicatoria* revealed by the complete genome sequence. *J Bacteriol* **187:** 7254-7266.

Triplett LR, Hamilton JP, Buell CR, Tisserat NA, Verdier V, Zink F *et al* (2011). Genomic analysis of *Xanthomonas oryzae* isolates from rice grown in the United States reveals substantial divergence from known *X. oryzae* pathovars. *Appl Environ Microbiol* **77:** 3930-3937.

Triplett LR, Verdier V, Campillo T, Van Malderghem C, Cleenwerck I, Maes M *et al* (2015). Characterization of a novel clade of *Xanthomonas* isolated from rice leaves in Mali and proposal of *Xanthomonas maliensis* sp. nov. *Antonie Van Leeuwenhoek* **107:** 869-881.

Vancheva T, Bogatzevska N, Moncheva P, Lefeuvre P, Koebnik R (2015). Draft genome sequences of two *Xanthomonas vesicatoria* strains from the balkan peninsula. *Genome Announc* **3:** e01558-01514.

Vandroemme J, Cottyn B, Baeyen S, De Vos P, Maes M (2013). Draft genome sequence of *Xanthomonas fragariae* reveals reductive evolution and distinct virulence-related gene content. *BMC Genomics* **14:** 829.

Varghese NJ, Mukherjee S, Ivanova N, Konstantinidis KT, Mavrommatis K, Kyrpides NC *et al* (2015). Microbial species delineation using whole genome sequences. *Nucleic Acids Res* **43:** 6761-6771.

Vauterin L, Hoste B, Kersters K, Swings J (1995). Reclassification of *Xanthomonas*. *Int J Syst Bacteriol* **45**.

Vauterin L, Rademaker J, Swings J (2000). Synopsis on the taxonomy of the genus *Xanthomonas*. *Phytopathology* **90:** 677-682.

Wang GL, Song WY, Ruan DL, Sideris S, Ronald PC (1996). The cloned gene, Xa21, confers resistance to multiple *Xanthomonas oryzae* pv. *oryzae* isolates in transgenic plants. *Mol Plant Microbe Interact* **9:** 850-855.

Wasukira A, Tayebwa J, Thwaites R, Paszkiewicz K, Aritua V, Kubiriba J *et al* (2012). Genome-wide sequencing reveals two major sub-lineages in the genetically monomorphic pathogen *Xanthomonas campestris* pathovar *musacearum*. *Genes (Basel)* **3:** 361-377.

Young JM (2008). An overview of bacterial nomenclature with special reference to plant pathogens. *Syst Appl Microbiol* **31:** 405-424.

Young JM, Park DC, Shearman HM, Fargier E (2008). A multilocus sequence analysis of the genus *Xanthomonas*. *Syst Appl Microbiol* **31:** 366-377.

Zhou JM, Tang D, Wang G (2017). Receptor kinases in plant pathogen interactions: more than pattern recognition. *Plant Cell* **29:** 618-637.

Figure 1. The *raxX-raxSTAB* gene cluster. A. The *rax* genetic region, drawn to scale. B. Boundary sequences. Sequences conserved within a group but different from other groups are colored green, brown, or yellow. Black sequence is conserved in all lineages, and blue sequence represents matches to consensi for transcription and translation initiation sequences. An "*mfsX*" +1 frameshift in *Xoo* sequences is indicated by the vertical red line. Abbreviations: *S. maltophilia, Sm; X. albilineans, Xa; X. arboricola pv. juglandis, Xaj; X. axonopodis pv. manihotis, Xam; X. campestris pv. campestris, Xcc; X. campestris pv. musacearum, Xcm; X. cannabis, Xc; X. citri subsp. citri, Xac; X. euvesicatoria, Xe; X. fragariae, Xf; X. hyacinthi, Xh; X. maliensis, Xm; X. oryzae pv. oryzae,Xoo; X. sacchari,; Xs X. translucens, Xt; X. vesicatoria, Xv.*

Figure 2. Model for r*axX-raxSTAB* inheritance during *Xanthomonas* speciation. The *Xanthomonas* spp. cladogram is based on published phylogenetic trees; see text for references. Gray lines depict lineages for strains that lack the *raxX-raxSTAB* gene cluster, whereas black lines depict those that carry the cluster. Numbers indicate *gcvP* length polymorphism in each species (see **Fig. 4**). Hypothetical events are: A, formation of the *raxX-raxSTAB* gene cluster; B, lateral transfer to *X. translucens*; C, lateral transfer to *X. maliensis*; D, loss from *X. citri*.

Figure 3. Phylogenetic trees for *rax* gene nucleotide sequences. The best scoring maximum likelihood trees for (A) *raxA*, (B) *raxB*, (C) *raxX* and (D) *raxST* in *Xanthomona*s spp. Numbers shown on branches represent the proportion of branches supported by 10,000 bootstrap replicates (0-100). Bootstraps are not shown for branches with less than 50% support, nor for branches too short to easily distinguish.

Figure 4. GcvP length polymorphisms. The relevant portion of the GcvP amino acid sequence is shown for each of the reference strains. Species in red lack the *raxX-raxSTAB* gene cluster, whereas those in blue lines carry the cluster. Numbers denote different allelic types. The positions of residues Gly-733 and Val-738 (numbering for allelic type 1) are indicated. Abbreviations are as in Fig. 1b.

Figure 5. Phylogenetic tree for *raxST* homologs in diverse bacterial genera. Distribution of *raxST* homolgs across bacterial genera. The tree shown was constructed by neighbor-joining with 1000 bootstrap replicates; branches with < 50% bootstrap support are not drawn. The *raxST* sequence from *Xoo* strain PXO99[A] was used as query for tBLASTn.

Figure S1. Whole genome-based *Xanthomonas* phylogenetic tree. Phylogenetic tree constructed from comparison of whole genome sequences; see text for details. See **Fig. 2** for the corresponding cladogram. Red lines depict lineages for strains that lack the *raxX-raxSTAB* gene cluster, whereas blue lines depict those that carry the cluster.

Figure S1.

Table 1. Reference strains for sequence comparisons.

Species	Strain	*raxX-raxSTAB*	Accession	Reference
S. maltophilia	K279a	–	NC_010943.1	(Crossman *et al.,* 2008)
X. albilineans	GPE PC73	–	NC_013722.1	(Pieretti *et al.,* 2015)
X. arboricola pv. *juglandis*	Xaj 417	–	NZ_CP012251.1	(Pereira *et al.,* 2015)
X. axonopodis pv. *manihotis*	UA536	+	NZ_AKEQ00000000	(Bart *et al.,* 2012)
X. campestris pv. *campestris*	ATCC 33913	–	NC_003902.1	(da Silva *et al.,* 2002)
X. campestris pv. *musacearum*	NCPPB 4392	+	NZ_AKBI00000000.1	(Wasukira *et al.,* 2012)
X. cannabis	NCPPB 2877	–	NZ_JSZE00000000.1	(Jacobs *et al.,* 2015)
X. citri subsp. *citri*	306	–	NC_003919.1	(da Silva *et al.,* 2002)
X. euvesicatoria	85-10	+	NZ_CP017190.1	(Thieme *et al.,* 2005)
X. fragariae	LMG 25863	–	NZ_AJRZ00000000.1	(Vandroemme *et al.,* 2013)
X. hyacinthi	DSM 19077	–	JPLD00000000.1	(Naushad *et al.,* 2015)
X. maliensis	M97	+	NZ_AQPR00000000.1	(Triplett *et al.,* 2015)
X. oryzae pv. *oryzae*	PXO99[A]	+	NC_010717.2	(Salzberg *et al.,* 2008)
X. sacchari	R1	–	NZ_CP010409.1	(Studholme *et al.,* 2011)
X. translucens	DAR61454	+	GCA_000334075.1	(Gardiner *et al.,* 2014)
X. vesicatoria	15b	–	NZ_JSXZ00000000.1	(Vancheva *et al.,* 2015)

Table 2. Nucleotide sequence identity between *rax* genes.

Species	gcvP	raxX	raxST	raxA	raxB	"mfsX"	gANI [a]	AF [b]
			Comparison to *X. euvesicatoria* sequence					
			% Nucleotide identity				Genome	
X. citri subsp. *citri*	94.8	— [c]	—	—	—	97.3	94.9	0.80
X. axonopodis pv. *manihotis*	94.1	99.0	96.1	97.6	95.8	97.3	95.0	0.79
X. maliensis	93.5	89.5	96.4	95.0	96.4	**85.0**	**83.5**	**0.66**
X. campestris pv. *malvacearum*	91.3	82.2	91.4	90.9	93.2	93.6	91.2	0.72
X. oryzae pv. *oryzae*	92.8	88.5	91.4	87.6	89.5	91.6	91.2	0.61
X. translucens	88.7	63.1	80.7	71.0	77.3	80.4	80.4	0.54

[a] genome-wide Average Nucleotide Identity (Varghese *et al.,* 2015).

[b] genome Alignment Fraction, from *X. euvesicatoria* to subject species (Varghese *et al.,* 2015).

[c] —, gene not present

A

B

Left Boundary:

Right Boundary:

0.2