Ecophysiology of freshwater Verrucomicrobia inferred from

metagenome-assembled genomes

Shaomei He[1,2], Sarah LR Stevens[1], Leong-Keat Chan[4], Stefan Bertilsson[3], Tijana Glavina del

Rio[4], Susannah G Tringe[4], Rex R Malmstrom[4], and Katherine D McMahon[1,5,*]

[1]Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA

[2]Department of Geoscience, University of Wisconsin-Madison, Madison, WI, USA

[3]Department of Ecology and Genetics, Limnology and Science for Life Laboratory, Uppsala

University, Uppsala, Sweden

[4]DOE Joint Genome Institute, Walnut Creek, CA, USA

[5]Department of Civil and Environmental Engineering, University of Wisconsin-Madison,

Madison, WI, USA

*Corresponding author

Katherine D McMahon (trina.mcmahon@wisc.edu)

Running title: Freshwater Verrucomicrobia genomes

Keywords: Metagenome-assembled genome (MAG), Verrucomicrobia, Freshwater,

Glycoside hydrolase, Nutrient limitation, Cytochrome *c*

Word count for the Abstract: 226

Word count for the text (excluding references and figure legends): 5850

ABSTRACT

Microbes are critical in carbon and nutrient cycling in freshwater ecosystems. Members of

the Verrucomicrobia are ubiquitous in such systems, yet their roles and ecophysiology are

not well understood. In this study, we recovered 19 Verrucomicrobia draft genomes by

sequencing 184 time-series metagenomes from a eutrophic lake and a humic bog that differ

in carbon source and nutrient availabilities. These genomes span four of the seven

previously defined Verrucomicrobia subdivisions, and greatly expand the known genomic

diversity of freshwater Verrucomicrobia. Genome analysis revealed their potential role as

(poly)saccharide-degraders in freshwater, uncovered interesting genomic features for this

life style, and suggested their adaptation to nutrient availabilities in their environments.

Between the two lakes, Verrucomicrobia populations differ significantly in glycoside

hydrolase gene abundance and functional profiles, reflecting the autochthonous and

terrestrially-derived allochthonous carbon sources of the two ecosystems respectively.

Interestingly, a number of genomes recovered from the bog contained gene clusters that

potentially encode a novel porin-multiheme cytochrome c complex and might be involved

in extracellular electron transfer in the anoxic humic-rich environment. Notably, most

epilimnion genomes have large numbers of so-called "Planctomycete-specific" cytochrome

c-containing genes, which exhibited nearly opposite distribution patterns with glycoside

hydrolase genes, probably associated with the different environmental oxygen availability

and carbohydrate complexity between lakes/layers. Overall, the recovered genomes are a

major step towards understanding the role, ecophysiology and distribution of

Verrucomicrobia in freshwater.

IMPORTANCE

Freshwater Verrucomicrobia are cosmopolitan in lakes and rivers, yet their roles and

ecophysiology are not well understood, as cultured freshwater Verrucomicrobia are

restricted to one subdivision of this phylum. Here, we greatly expand the known genomic

diversity of this freshwater lineage by recovering 19 Verrucomicrobia draft genomes from

184 metagenomes collected from a eutrophic lake and a humic bog across multiple years.

Most of these genomes represent first freshwater representatives of several

Verrucomicrobia subdivisions. Genomic analysis revealed Verrucomicrobia as potential

(poly)saccharide-degraders, and suggested their adaptation to carbon source of different

origins in the two contrasting ecosystems. We identified putative extracellular electron

transfer genes and so-called "Planctomycete-specific" cytochrome c-containing genes, and

found their distinct distribution patterns between the lakes/layers. Overall, our analysis

greatly advances the understanding of the function, ecophysiology and distribution of

freshwater Verrucomicrobia, while highlighting their potential role in freshwater carbon

cycling.

INTRODUCTION

Verrucomicrobia are ubiquitous in freshwater and exhibit a cosmopolitan distribution in

lakes and rivers. They are present in up to 90% of lakes (1), with abundances typically

between <1% and 6% of total microbial community (2-4), but as high as 19% in a humic

lake (5). Yet, in comparison to other freshwater bacterial groups, such as members of the

Actinobacteria, Cyanobacteria and Proteobacteria phyla, Verrucomicrobia have received

relatively less attention, and their functions and ecophysiology in freshwater are not well

understood.

As a phylum, Verrucomicrobia (V) was first proposed relatively recently, in 1997

(6). Together with Planctomycetes (P), Chlamydiae (C), and sister phyla such as

Lentisphaerae, they comprise the PVC superphylum. In addition to being cosmopolitan in

freshwater, Verrucomicrobia have been found in oceans (7, 8), soil (9, 10), wetlands (11),

rhizosphere (12), and animal guts (13, 14), as free-living organisms or symbionts of

eukaryotes. Verrucomicrobia isolates are metabolically diverse, including aerobes,

facultative anaerobes, and obligate anaerobes, and they are mostly heterotrophs, using

various mono-, oligo-, and poly-saccharides for growth (6, 7, 11, 14-20). Not long ago an

autotrophic verrucomicrobial methanotroph (*Methylacidiphilum fumariolicum* SolV) was

discovered in acidic thermophilic environments (21).

In marine environments, Verrucomicrobia are also ubiquitous (22) and suggested to

have a key role as polysaccharide degraders (23, 24). Genomic insights gained through

sequencing single cells (24) or extracting Verrucomicrobia bins from metagenomes (25)

have revealed high abundances of glycoside hydrolase genes, providing more evidence for

their critical roles in C cycling in marine environments.

In freshwater, Verrucomicrobia have been suggested to degrade glycolate (26) and

polysaccharides (24). The abundance of some phylum members was favored by high

nutrient availabilities (27, 28), cyanobacterial blooms (29), low pH, high temperature, high

hydraulic retention time (30), and more labile DOC (5). To date, there are very few

freshwater Verrucomicrobia isolates, including *Verrucomicrobium spinosum* (31) and

several *Prosthecobacter* spp. (6). Physiological studies showed that they are aerobes,

primarily using carbohydrates, but not amino acids, alcohols, or rarely organic acids for

growth. However, these few cultured isolates only represent a single clade within

subdivision 1. By contrast, 16S rRNA gene based studies discovered a much wider

phylogenic range of freshwater Verrucomicrobia, including subdivisions 1, 2, 3, 4, 5, and 6

(3-5, 24, 32). Due to the very few cultured representatives and few available genomes from

this freshwater lineage, the ecological functions of the vast uncultured freshwater

Verrucomicrobia are largely unknown.

In this study, we sequenced a total of 184 metagenomes in a time-series study of

two lakes with contrasting characteristics, particularly differing in C source, nutrient

availabilities, and pH. We recovered a total of 19 Verrucomicrobia draft genomes spanning

subdivision 1, 2, 3, and 4 of the seven previously defined Verrucomicrobia subdivisions. We

inferred their metabolisms, revealed their adaptation to C and nutrient conditions, and

uncovered some interesting and novel features, including a novel putative porin-

multiheme cytochrome c system that may be involved in extracellular electron transfer.

The gained insights advanced our understanding of the ecophysiology, and suggested

potential roles in C cycling and ecological niches of this ubiquitous freshwater bacterial

group.

RESULTS AND DISCUSSION

Comparison of the two lakes

The two studied lakes exhibited contrasting characteristics (**Table 1**). The most notable

difference is the primary C source and nutrient availabilities. Mendota is an urban

eutrophic lake with most of its C being autochthonous (in-lake produced through

photosynthesis). By contrast, Trout Bog is a nutrient-poor dystrophic lake, surrounded by

temperate forests and sphagnum mats, thus receiving large amounts of terrestrially-

derived allochthonous C that is rich in humic and fulvic acids. Compared to Mendota, Trout

Bog features higher DOC levels, but is more limited in nutrient availability, with much

higher DOC:TN and DOC:TP ratios (**Table 1**). Nutrient limitation in Trout Bog is even more

extreme than revealed by these ratios because much of the N and P is tied up in complex

dissolved organic matter. In addition, Trout Bog has lower oxygenic photosynthesis due to

decreased photosynthetically active radiation (PAR) as a result of absorption by DOC (33).

Together with the consumption of dissolved oxygen by heterotrophic respiration, oxygen

levels decrease quickly with depth in the water column in Trout Bog. Dissolved oxygen

levels are below detection in the hypolimnion nearly year-round (34). Due to these

contrasts, we expected to observe differences in bacterial C and nutrient use, as well as

differences reflecting the electron acceptor conditions between these two lakes. Hence, the

retrieval of numerous Verrucomicrobia draft genomes in the two lakes not only allows the

prediction of their general functions in freshwater, but also provides an opportunity to

study their ecophysiological adaptation to the local environmental differences.

Verrucomicrobia draft genome retrieval and their distribution patterns

A total of 184 metagenomes were generated from samples collected across multiple years,

including 94 from the top 12 m of Mendota (mostly consisting of the epilimnion layer,

therefor referred to as "ME"), 45 from Trout Bog epilimnion ("TE"), and 45 from Trout Bog

hypolimnion ("TH"). Three combined assemblies were generated by co-assembling reads

from all metagenomes within the ME, TE, and TH groups, respectively. Using the binning

facilitated by tetranucleotide frequency and relative abundance patterns over time, a total

of 19 Verrucomicrobia metagenome-assembled genomes (MAGs) were obtained, including

eight from the combined assembly of ME, three from the combined assembly of TE, and

eight from the combined assembly of TH (**Table 2**). The 19 MAGs exhibited a clustering of

their tetranucleotide frequency largely based on the two lakes (**Fig. S1**), suggesting distinct

overall genomic signatures associated with each system.

Genome completeness of the 19 MAGs ranged from 51% to 95%, as determined by

checkM (35). Phylogenetic analysis of these MAGs using a concatenated alignment of their

conserved genes indicates that they span a wide phylogenetic spectrum and distribute in

subdivisions 1, 2, 3, and 4 of the seven previously defined Verrucomicrobia subdivisions (5,

21, 36) (**Fig. 1**), as well as three unclassified Verrucomicrobia MAGs.

Presently available freshwater Verrucomicrobia isolates are restricted to

subdivision 1. The recovered MAGs allow the inference of metabolisms and ecology of a

considerable diversity within uncultured freshwater Verrucomicrobia. Notably, all MAGs

from subdivision 3 were recovered from TH, and all MAGs from subdivision 1, except

TH2746, were from the epilimnion (either ME or TE), indicating differences in phylogenetic

distribution between lakes and between layers within a lake.

We used normalized coverage depth of MAGs within individual metagenomes

collected at different sampling time points and different lakes/layers to comparatively infer

relative population abundance across time and space (see detailed coverage depth

estimation in **Supplementary Text**). Briefly, we mapped reads from each metagenome to

MAGs with a minimum identity of 95%, and used the number of mapped reads to calculate

the relative abundance for each MAG based on coverage depth per contig and several

normalization steps. Thus, we assume that each MAG represents a distinct population

within the lake-layer from which it was recovered (37, 38). This estimate does not directly

indicate the actual relative abundance of these populations within the total community per

se; rather it allows us to compare population abundance levels from different lakes and

sampling occasions within the set of 19 MAGs. This analysis indicates that Verrucomicrobia

populations in Trout Bog were proportionally more abundant and persistent over time

compared to those in Mendota in general (**Table 2**). Verrucomicrobia populations in

Mendota boosted their abundances once to a few times during the sampling season and

diminished to extremely low levels for the remainder of the sampling season (generally

May to November), as reflected by the low median coverage depth of Mendota MAGs and

their large coefficient of variation (**Table 2**)

Saccharolytic life style and adaptation to different C sources

Verrucomicrobia isolates from different environments are known to grow on various

mono-, oligo-, and poly-saccharides, but are unable to grow on amino acids, alcohols, or

most organic acids (6, 7, 11, 14-20, 39). Culture-independent research suggests marine

Verrucomicrobia as candidate polysaccharide degraders with large number of genes

involved in polysaccharide utilization (23-25).

In the 19 Verrucomicrobia MAGs, we observed rich arrays of glycoside hydrolase

(GH) genes, representing a total of 78 different GH families acting on diverse

polysaccharides (**Fig. S2**). Although these genomes have different degrees of completeness,

genome completeness was not correlated with the number of GH genes recovered

(correlation coefficient = 0.312, p-value = 0.194), or the number of GH families represented

in each MAG (i.e. GH diversity, correlation coefficient = 0.278, p-value = 0.250). To compare

GH abundance among MAGs, we normalized GH occurrence frequencies by the total

number of genes in each MAG to estimate the percentage of genes annotated as GHs (i.e. GH

coding density) to account for the different genome size and completeness. This

normalization assumes GH genes are randomly distributed between the recovered and the

missing parts of the genome, and it allows us to make some general comparison among

these MAGs. GH coding density ranged from 0.4% to 4.9% for these MAGs (**Fig. 2a**), and in

general, was higher in Trout Bog MAGs than in Mendota MAGs. Notably, six TH MAGs had

extremely high (~4%) GH coding densities (**Fig. 2a**), with each MAG harboring 119-239 GH

genes, representing 36-59 different GH families (**Fig. 3 and S2**). Although GH coding

density in most ME genomes in subdivisions 1 and 2 was relatively low (0.4-1.6%), it was

still higher than in many other bacterial groups (24).

The GH abundance and diversity within a genome may determine the width of the

substrate spectrum and/or the complexity of carbohydrates used by that organism. For

example, there are 20 GH genes in the *Rubritalea marina* genome, and this marine

verrucomicrobial aerobe only uses a limited spectrum of carbohydrate monomers and

dimers, but not the majority of (poly)saccharides tested (15). By contrast, 164 GH genes

are present in the *Opitutus terrae* genome, and this soil verrucomicrobial anaerobe can thus

grow on a wider range of mono-, di- and poly-saccharides (16). Therefore, it is plausible

that the GH-rich Trout Bog Verrucomicrobia populations may be able to use a wider range

of more complex polysaccharides than the Mendota populations.

The 10 most abundant GH families in these Verrucomicrobia MAGs include GH2, 29,

78, 95, and 106 (**Fig. 3**). These specific GHs were absent or at very low abundances in

marine Verrucomicrobia genomes (24, 25), suggesting a general difference in carbohydrate

substrate use between freshwater- and marine Verrucomicrobia. Hierarchical clustering of

MAGs based on overall GH abundance profiles indicated a grouping pattern largely separated by lake (**Fig. S3**). Prominently over-represented GHs in most Trout Bog MAGs include GH2, GH29, 78, 95, and 106. By contrast, over-represented GHs in the Mendota MAGs are GH13, 20, 33, 57, and 77, which have different substrate spectra from GHs over-represented in the Trout Bog MAGs. Therefore, the patterns in GH functional profiles may suggest varied carbohydrate substrate preferences and ecological niches occupied by Verrucomicrobia, probably reflecting the different carbohydrate composition derived from different sources between Mendota and Trout Bog.

Overall, GH diversity and abundance profile may reflect the DOC availability, chemical variety and complexity, and suggest microbial adaptation to different C sources in the two ecosystems. We speculate that the rich arrays of GH genes, and presumably broader substrate spectra of Trout Bog populations, partly contribute to their higher abundance and persistence over the sampling season (**Table 2**), as they are less likely impacted by fluctuations of individual carbohydrates. By contrast, Mendota populations with fewer GHs and presumably more specific substrate spectra are relying on autochthonous C and therefore exhibit a "bloom-and-bust" abundance pattern (**Table 2**) that might be associated with cyanobacterial blooms as previous suggested (29). On the other hand, bogs experience seasonal phytoplankton blooms (40, 41) that introduce brief pulses of autochthonous C to these otherwise allochthonous-driven systems. Clearly, much remains to be learned about the routes through which C is metabolized by bacteria in such lakes, and comparative genomics is a novel way to use the organisms to tell us about C flow through the ecosystem.

Other genome features of the saccharide-degrading life style

Seven Verrucomicrobia MAGs spanning subdivisions 1, 2, 3, and 4 possess genes needed to

construct bacterial microcompartments (BMCs), which are quite rare among studied

bacterial lineages. Such BMC genes in Planctomycetes are involved in the degradation of

plant and algal cell wall sugars, and are required for growth on L-fucose, L-rhamnose and

fucoidans (42). Genes involved in L-fucose and L-rhamnose degradation cluster with BMC

shell protein-coding genes in the seven Verrucomicrobia MAGs (**Fig. 4**). This is consistent

with the high abundance of α-L-fucosidase or α-L-rhamnosidase GH genes (represented

by GH29, 78, 95, 106) in most of these MAGs (**Fig. 3**), suggesting the importance of fucose-

and rhamnose-containing polysaccharides for these Verrucomicrobia populations.

TonB-dependent receptor (TBDR) genes were found in Verrucomicrobia MAGs, and

are present at over 20 copies in TE1800 and TH2519. TBDRs are located on the outer

cellular membrane of Gram-negative bacteria, usually mediating the transport of iron

siderophore complex and vitamin B_{12} across the outer membrane through an active

process. More recently, TBDRs were suggested to be involved in carbohydrate transport

across the outer membrane by some bacteria that consume complex carbohydrates, and in

their carbohydrate utilization (CUT) loci, TBDR genes usually cluster with genes encoding

inner membrane transporters, GHs and regulators for efficient carbohydrate

transportation and utilization (43). Such novel CUT loci are present in TE1800 and

TH2519, with TBDR genes clustering with genes encoding inner membrane sugar

transporters, monosaccharide utilization enzymes, and GHs involved in the degradation of

pectin, xylan, and fucose-containing polymers (**Fig. 5**). Notably, most GHs in the CUT loci

are predicted to be extracellular or outer membrane proteins (**Fig. 5**), catalyzing

extracellular hydrolysis reactions to release mono- and oligo-saccharides, which are

transported across the outer membrane by TBDR proteins. Therefore, such CUT loci may

allow these verrucomicrobial populations to coordinately and effectively scavenge the

hydrolysis products before they diffuse away.

Genes encoding for inner membrane carbohydrate transporters are abundant in

Verrucomicrobia MAGs (**Fig. S4**). The Embden-Meyerhof pathway for glucose degradation,

as well as pathways for degrading a variety of other sugar monomers, including galactose,

rhamnose, fucose, xylose, and mannose, were recovered (complete or partly-complete) in

most MAGs (**Fig. 6**). As these sugars are abundant carbohydrate monomers in plankton and

plant cell walls, the presence of these pathways together with GH genes suggest that these

Verrucomicrobia populations may use plankton- and plant-derived saccharides. Machinery

for pyruvate degradation to acetyl-CoA and the TCA cycle are also present in most MAGs.

These results are largely consistent with their hypothesized role in carbohydrate

degradation and previous studies on Verrucomicrobia isolates.

Notably, a large number of genes encoding proteins belonging to a sulfatase family

(pfam00884) are present in the majority of MAGs (**Fig. 2b**), similar to the high

representation of these genes in marine Verrucomicrobia genomes (24, 25). Sulfatases

hydrolyze sulfate esters, which are rich in sulfated polysaccharides. In general, sulfated

polysaccharides are abundant in marine algae and plants (mainly in seaweeds) (44), but

have also been found in some freshwater cyanobacteria (45) and plant species (46).

Sulfatase genes in our Verrucomicrobia MAGs were often located in the same neighborhood

as genes encoding for extracellular proteins with a putative pectin lyase activity, proteins

with a carbohydrate-binding module (pfam13385), GHs, and proteins with PSCyt domains

(**Fig. 2c** and discussed later). Their genome context lends support for the participation of

these genes in C and sulfur cycling by degrading sulfated polysaccharides, which can serve

as an abundant source of sulfur for cell biosynthesis as well as C for energy and growth.

Previously, freshwater Verrucomicrobia were suggested to use the algal exudate

glycolate in humic lakes, based on the retrieval of genes encoding subunit D (*glcD*) of

glycolate oxidase, which converts glycolate to glyoxylate (26). However, these recovered

genes might not be bona fide *glcD* due to the lack of other essential subunits as revealed in

our study (see **Supplementary Text**). Among the MAGs, only TE4605 possesses all three

essential subunits of glycolate oxidase (*glcDEF)* (**Fig. S5**). However, genetic context analysis

suggests that TE4605 likely uses glycolate for amino acid assimilation, instead of energy

generation (**Fig. S5** and **Supplementary Text**). These results are consistent with the

absence of the glyoxylate shunt and especially the malate synthase, which converts

glyoxylate to malate to be used through the TCA cycle for energy generation in the 19

Verrucomicrobia MAGs (**Fig. S6**). Therefore, Verrucomicrobia populations represented by

the 19 MAGs are not likely key players in glycolate degradation, but more likely important

(poly)saccharide-degraders in freshwater, as suggested by the high abundance of GH,

sulfatase, and carbohydrate transporter genes, metabolic pathways for degrading diverse

carbohydrate monomers, and other genome features adapted to the saccharolytic life style.

Nitrogen (N) metabolism and adaptation to different N availabilities

Most Verrucomicrobia MAGs in our study do not appear to reduce nitrate or other

nitrogenous compounds, and they seem to uptake and use ammonia (**Fig. 6**), and

occasionally amino acids (**Fig. S4**), as an N-source. Further, some Trout Bog populations

may have additional avenues to generate ammonia, including genetic machineries for

assimilatory nitrate reduction in TH2746, nitrogenase genes for nitrogen fixation and

urease genes in some of the Trout Bog MAGs (**Fig. 6**), probably as adaptions to N-limited

conditions in Trout Bog.

Although Mendota is a eutrophic lake, N can become temporarily limiting during the

high-biomass period when N is consumed by large amounts of phytoplankton and

bacterioplankton (47). For some bacteria, when N is temporarily limited while C is in

excess, cells convert and store the extra C as biopolymers. For example, the

verrucomicrobial methanotroph *M. fumariolicum* SolV accumulated a large amount of

glycogen (up to 36% of the total dry weight of cells) when the culture was N-limited (48).

Similar to this verrucomicrobial methanotroph, genes in glycogen biosynthesis are present

in most MAGs from Mendota and Trout Bog (**Fig. 6**). Indeed, a glycogen synthesis pathway

is also present in most genomes of cultivated Verrucomicrobia in the public database (data

not shown), suggesting that glycogen accumulation might be a common feature for this

phylum to cope with the changing pools of C and N in the environment and facilitate their

survival when either is temporally limited.

Phosphorus (P) metabolism and other metabolic features

Verrucomicrobia populations represented by these MAGs may be able to survive under low

P conditions, as suggested by the presence of genes responding to P limitation, such as the

two-component regulator (*phoRB*), alkaline phosphatase (*phoA*), phosphonoacetate

hydrolase (*phnA*), and high-affinity phosphate-specific transporter system (*pstABC*) (**Fig.**

6). Detailed discussion in P acquisition and metabolism and other metabolic aspects, such

as acetate metabolism, sulfur metabolism, oxygen tolerance, and the presence of the

alternative complex III and cytochrome *c* oxidase genes in the oxidative phosphorylation

pathway, are discussed in the Supplementary Text (**Fig. S6**).

Anaerobic respiration and a putative porin-multiheme cytochrome *c* system

Respiration using alternative electron acceptors is important for overall lake metabolism in

the DOC-rich humic Trout Bog, as the oxygen levels decrease quickly with depth in the

water column. We therefore searched for genes involved in anaerobic respiration, and

found that genes in the dissimilatory reduction of nitrate, nitrite, sulfate, sulfite, DMSO, and

TMAO are largely absent in all MAGs (**Supplementary Text, Fig. S6)**. Compared to those

anaerobic processes, genes for dissimilatory metal reduction are less well understood. In

more extensively studied cultured iron [Fe(III)] reducers, outer surface *c*-type cytochromes

(cyt*c*), such as OmcE and OmcS in *Geobacter sulfurreducens* are involved in Fe(III)

reduction at the cell outer surface (49). Further, a periplasmic multiheme cytochrome *c*

(MHC, e.g. MtrA in *Shewanella oneidensis* and OmaB/OmaC in *G. sulfurreducens*) can be

embedded into a porin (e.g. MtrB in *S. oneidensis* and OmbB/OmbC in *G. sulfurreducens*),

forming a porin-MHC complex as an extracellular electron transfer (EET) conduit to reduce

extracellular Fe(III) (50, 51). Such outer surface cyt*c* and porin-MHC systems involved in

Fe(III) reduction were also suggested to be important in reducing the quinone groups in

humic substances (HS) at the cell surface (52-54). The reduced HS can be re-oxidized by

Fe(III) or oxygen, thus HS can serve as electron shuttles to facilitate Fe(III) reduction (55,

56) or as regenerable electron acceptors at the anoxic-oxic interface or over redox cycles

(57).

Outer surface cyt*c* or porin-MHC systems homologous to the ones in *G.*

sulfurreducens and *S. oneidensis* are not present in Verrucomicrobia MAGs. Instead, we

identified a novel porin-coding gene clustering with MHC genes in six MAGs (**Fig. 7**). These

porins were predicted to have at least 20 transmembrane motifs, and their adjacent cyt*c*

were predicted to be periplasmic proteins with eight conserved heme-binding sites. In

several cases, a gene encoding an extracellular MHC is also located in the same gene cluster.

As their gene organization is analogous to the porin-MHC gene clusters in *G. sulfurreducens*

and *S. oneidensis*, we hypothesize that these genes in Verrucomicrobia may encode a novel

porin-MHC complex involved in EET.

As these porin-MHC gene clusters are novel, we further confirmed that they are

indeed from Verrucomicrobia. Their containing contigs were indeed classified to

Verrucomicrobia based on the consensus of the best BLASTP hits for genes on these

contigs. Notably, the porin-MHC gene cluster was only observed in MAGs recovered from

the HS-rich Trout Bog, especially from the anoxic hypolimnion environment. Searching the

NCBI and IMG databases for the porin-MHC gene clusters homologous to those in Trout

Bog, we identified homologs in genomes within the Verrucomicrobia phylum, including

Opitutus terrae PB90-1 isolated from rice paddy soil, *Opitutus* sp. GAS368 isolated from

forest soil, "*Candidatus* Udaeobacter copiosus" recovered from prairie soil, Opititae-40 and

Opititae-129 recovered from freshwater sediment, and Verrucomicrobia bacterium

IMCC26134 recovered from freshwater; some of their residing environments are also rich

in HS. Therefore, based on the occurrence pattern of porin-MHC among Verrucomicrobia

genomes, we hypothesize that such porin-MHCs might participate in EET to HS in anoxic

HS-rich environments, and HS may further shuttle electrons to poorly soluble metal oxides

or be regenerated at the anoxic-oxic interface, thereby diverting more C flux to respiration

instead of fermentation and methanogenesis, which could impact the overall energy

metabolism and green-house gas emission in the bog environment.

Occurrence of Planctomycete-specific cytochrome *c* and domains

One of the interesting features of Verrucomicrobia and its sister phyla in the PVC

superphylum is the presence of a number of novel protein domains in some of their

member genomes (58, 59). These domains were initially identified in marine

planctomycete *Rhodopirellula baltica* (58) and therefore, were referred to as

"Planctomycete-specific", although some of them were later identified in other PVC

members (59). In our Verrucomicrobia MAGs, most genes containing Planctomycete-

specific cytochrome *c* domains (PSCyt1 to PSCyt3) also contain other Planctomycete-

specific domains (PSD1 through PSD5) with various combinations and arrangements (**Fig.**

8 and **S7a**). Further, PSCyt2-containing and PSCyt3-containing genes are usually next to

two different families of unknown genes, respectively (**Fig. S7b**). Such conserved domain

architectures and gene organizations, as well as their high occurrence frequencies in some

of the Verrucomicrobia MAGs are intriguing, yet nothing is known about their functions.

However, some of the PSCyt-containing genes also contain protein domains identifiable as

carbohydrate-binding modules (CBMs), suggesting a role in carbohydrate metabolism (see

detailed discussion in **Supplementary Text**).

The coding density of PSCyt-containing genes indicates that they tend to be more

abundant in the epilimnion (either ME or TE) genomes (**Fig. 2c**) and exhibit an inverse

correlation with the GH coding density (r = -0.62). Interestingly, sulfatase-coding genes are

often in the neighborhood of PSCyt-containing genes in ME and TE genomes, whereas

sulfatase-coding genes often neighbor with GH genes in TH genomes. The genomic context

suggests PSCyt-containing gene functions somewhat mirror those of GHs (although their

reaction mechanisms likely differ fundamentally). However, these PSCyt-containing genes

were predicted to be periplasmic or cytoplasmic proteins rather than extracellular or outer

membrane proteins. Hence, if they are indeed involved in carbohydrate degradation, they

likely act on mono- or oligomers that can be transported into the cell. Further, the

distribution patterns of GH versus PSCyt-containing genes between the epilimnion and

hypolimnion may reflect the difference in oxygen availability and their carbohydrate

substrate complexity between the two layers, suggesting some niche differentiation within

Verrucomicrobia in freshwater systems. Therefore, we suggest that a combination of

carbohydrate composition, electron acceptor availability and C accessibility drive gene

distributions in these populations.

Summary

Verrucomicrobia MAGs recovered from the two contrasting lakes greatly expanded the

known genomic diversity of freshwater Verrucomicrobia, revealed the ecophysiology and

some interesting adaptive features of this ubiquitous yet less understood freshwater

lineage. The overrepresentation of GH, sulfatase, and carbohydrate transporter genes, the

genetic potential to use various sugars, and the microcompartments for fucose and

rhamnose degradation suggest that they are potentially (poly)saccharide degraders in

freshwater. Most of the MAGs encode machineries to cope with the changing availability of

N and P and can survive nutrient limitation. Despite these generalities, these

Verrucomicrobia differ significantly between lakes in the abundance and functional profiles

of their GH genes, which may reflect different C sources of the two lakes. Interestingly, a

number of MAGs in Trout Bog possess gene clusters potentially encoding a novel porin-

multiheme cytochrome c complex, and might be involved in extracellular electron transfer

in the anoxic humic-rich environment. Intriguingly, large numbers of Planctomycete-

specific cytochrome c-containing genes are present in MAGs from the epilimnion,

exhibiting nearly opposite distribution patterns with GH genes. Future studies are needed

to elucidate the functions of these novel and fascinating genomic features.

In this study, we focused on using genome information to infer ecophysiology of

Verrucomicrobia. The rich time-series metagenome dataset and the many diverse

microbial genomes recovered in these two lakes also provide an opportunity for the future

study of Verrucomicrobia population dynamics in the context of the total community and

their interactions with environmental variables and other microbial groups.

As some of the MAGs analyzed here represent first genome representatives of

several Verrucomicrobia subdivisions from freshwater, an interesting question is whether

populations represented by the MAGs are native aquatic residents and active in aquatic

environment, or merely present after having been washed into the lake from surrounding

soil. Previous studies on freshwater Verrucomicrobia were largely based on 16S rRNA

genes, yet 16S rRNA genes were not recovered in most MAGs, making it difficult to directly

link our MAGs to previously identified freshwater Verrucomicrobia. Notably, our MAGs

were only distantly related to the ubiquitous and abundant soil Verrucomicrobia,

"*Candidatus* Udaeobacter copiosus" (10) (**Fig. 1**). In addition, Verrucomicrobia were

abundant in Trout Bog and other bogs from a five-year bog lake bacterial community

composition and dynamics study (60), with average relative abundance of 7.1% and 8.6%,

and maximal relative abundance of 25.4% and 39.5% in Trout Bog epilimnion and

hypolimnion respectively. Since the MAGs were presumably from the most abundant

Verrucomicrobia populations, they were not likely soil immigrants due to their high

abundance in the aquatic environment. To confirm their aquatic origin, future experiments

should be designed to test their activities and physiology in the aquatic environment based

on the genomic insights gained in this study.

MATERIALS AND METHODS

Study sites. Samples for metagenome sequencing were collected from two temperate lakes

in Wisconsin, USA, Lake Mendota and Trout Bog Lake, during ice-off periods of each year

(May to November). Mendota is an urban eutrophic lake with most of its C being

autochthonous (in-lake produced), whereas Trout Bog is a small, acidic and nutrient-poor

dystrophic lake with mostly terrestrially-derived (allochthonous) C. General lake

characteristics are summarized in **Table 1**.

Sampling. For Mendota, we collected depth-integrated water samples from the surface 12

m (mostly consisting of the epilimnion layer) at 94 time points from 2008 to 2012, and

samples were referred to as "ME" (38). For Trout Bog, we collected the integrated

hypolimnion layer at 45 time points from 2007 to 2009 and the integrated epilimnion layer

at 45 time points from 2007 to 2009, and samples were referred to as "TH" and "TE",

respectively (37). All samples were filtered through 0.22 μ m polyethersulfone filters and

stored at -80°C until extraction. DNA was extracted from the filters using the FastDNA kit

(MP Biomedicals) according to manufacturer's instruction with some minor modifications

as described previously (34).

Metagenome sequencing, assembly, and draft genome recovery. Details of

metagenome sequencing, assembly, and binning were described in Bendall *et al.* (37) and

Hamilton et al (61). Briefly, shotgun Illumina HiSeq 2500 metagenome libraries were

constructed for each of the DNA samples. Three combined assemblies were generated by

co-assembling reads from all metagenomes within the ME, TE, and TH groups, respectively.

Binning was conducted on the three combined assemblies to recover "metagenome-

assembled genomes" (MAGs) based on the combination of contig tetranucleotide frequency

and differential coverage patterns across time points using MetaBAT (62). Subsequent

manual curation of MAGs was conducted to remove contigs that did not correlate well with

the median temporal abundance pattern of all contigs within a MAG, as described in

Bendall *et al.* (37).

Genome annotation and completeness estimation. MAGs were submitted to the DOE

Joint Genome Institute's Integrated Microbial Genome (IMG) database for gene prediction

and function annotation (63). The IMG Taxon Object IDs for Verrucomicrobia MAGs are

listed in **Table 2**. The completeness and contamination of each MAG was estimated using

checkM with both the lineage-specific and Verrucomicrobia-specific workflows (35). The

Verrucomicrobia-specific workflow provided more accurate estimates (i.e. higher genome

completeness and lower contamination) than the lineage-specific workflow when tested on

11 complete genomes of Verrucomicrobia isolates available at IMG during our method

validation. We therefore only reported the estimates from Verrucomicrobia-specific

workflow (**Table 2**). MAGs with an estimated completeness lower than 50% were not

included in this study.

Taxonomic and phylogenetic analysis. A total of 19 MAGs were classified to the

Verrucomicrobia phylum based on taxonomic assignment by PhyloSift using 37 conserved

phylogenetic marker genes (64), as described in Bendall *et al.* (37). A phylogenetic tree was

reconstructed from the 19 Verrucomicrobia MAGs and 24 reference genomes using an

alignment concatenated from individual protein alignments of five conserved essential

single-copy genes (represented by TIGR01391, TIGR01011, TIGR00663, TIGR00460, and

TIGR00362) that were recovered in all Verrucomicrobia MAGs. Individual alignments were

first generated with MUSCLE (65), concatenated, and trimmed to exclude columns that

contain gaps for more than 30% of all sequences. A maximum likelihood phylogenetic tree

was constructed using PhyML 3.0 (66), with the LG substitution model and the gamma

distribution parameter estimated by PhyML. Bootstrap values were calculated based on

100 replicates. *Kiritimatiella glycovorans* L21-Fru-AB was used as an outgroup in the

phylogenetic tree. This bacterium was initially designated as the first (and so far the only)

cultured representative of Verrucomicrobia subdivision 5. However, this subdivision was

later proposed as a novel sister phylum associated with Verrucomicrobia (67), making it an

ideal outgroup for this analysis.

Estimate of metabolic potential. IMG provides functional annotation based on KO (KEGG orthology) term, COG (cluster of orthologous group), pfam, and TIGRfam. To estimate metabolic potential, we primarily used KO terms due to their direct link to KEGG pathways. COG, pfam, and TIGRfam were also used when KO terms were not available for a function. Pathways are primarily reconstructed according to KEGG modules, and MetaCyc pathway is used if a KEGG module is not available for a pathway. As these MAGs are incomplete genomes, a fraction of genes in a pathway may be missing due to genome incompleteness. Therefore, we estimated the completeness of a pathway as the fraction of recovered enzymes in that pathway (e.g. a pathway is 100% complete if all enzymes in that pathway are encoded by genes recovered in a MAG). As some genes are shared by multiple pathways, signature genes specific for a pathway were used to indicate the presence of a pathway. If signature genes for a pathway were missing in all MAGs, that pathway was likely absent in all genomes. Based on this, we established criteria for estimating pathway completeness in each MAG. If a signature gene in a pathway was present, we report the percentage of genes in the pathway that we found. If a signature gene was absent in a MAG, but present in at least one third of all MAGs (i.e. >=7), we still report the pathway completeness for that MAG in order to account for genome incompleteness. Otherwise, we considered the pathway to be absent (i.e. completeness is 0%).

Glycoside hydrolase identification. Glycoside hydrolase (GH) genes were identified using the dbCAN annotation tool (http://csbl.bmb.uga.edu/dbCAN/annotate.php) (68) using HMMER search against hidden Markov models (HMMs) built for all GHs, with an E-value cutoff of 1e-7, except GH109, for which we found that the HMM used by dbCAN is

pfam01408, which is a small domain at the N-terminus of GH109 proteins, but is not

specific for GH109. Therefore, to identify verrucomicrobial GH109, BLASTP was performed

using the two GH109 sequences (GenBank accession ACD03864 and ACD04752) from

verrucomicrobial *Akkermansia muciniphila* ATCC BAA-835 listed in the CAZy database

(http://www.cazy.org), with E-value cutoff of 1e-6 and query sequence coverage cutoff of

50%.

Other bioinformatic analyses. Protein cellular location was predicted using CELLO v.2.5

(http://cello.life.nctu.edu.tw) (69) and PSORTb v.3.0 (http://www.psort.org/psortb) (70).

The beta-barrel structure of outer membrane proteins was predicted by PRED-TMBB

(http://bioinformatics.biol.uoa.gr//PRED-TMBB) (71).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank the North Temperate Lakes Microbial Observatory 2007-2012 field crews, UW-

Trout Lake Station, the UW Center for Limnology, and the Global Lakes Ecological

Observatory Network for field and logistical support. We give special thanks to past

McMahon lab graduate students Ashley Shade, Ryan Newton, Emily Read, and Lucas

Beversdorf. We acknowledge efforts by many McMahon Lab undergrads and technicians

related to sample collection and DNA extraction, particularly Georgia Wolfe. We personally

thank the individual program directors and leadership at the National Science Foundation

for their commitment to continued support of long term ecological research.

The project was supported by funding from the United States National Science Foundation

Microbial Observatories program (MCB-0702395), the Long Term Ecological Research

program (NTL-LTER DEB-1440297) and an INSPIRE award (DEB-1344254). This material

is also based upon work that supported by the National Institute of Food and Agriculture,

U.S. Department of Agriculture (Hatch Project 1002996). The work conducted by the U.S.

Department of Energy Joint Genome Institute, a DOE Office of Science User Facility, is

supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-

AC02-05CH11231.

565
566
567   **FUNDING INFORMATION**
568   U.S. NSF Microbial Observatories program (MCB-0702395)
569   Katherine D McMahon
570
571   U.S. NSF Long Term Ecological Research program (NTL-LTER DEB-1440297)
572   Katherine D McMahon
573
574   U.S. NSF INSPIRE award (DEB-1344254)
575   Katherine D McMahon
576
577   U.S. National Institute of Food and Agriculture, U.S. Department of Agriculture (Hatch
578   Project 1002996)
579   Katherine D McMahon
580

REFERENCE

1. **Zwart G, van Hannen EJ, Kamst-van Agterveld MP, Van der Gucht K, Lindström ES, Van Wichelen J, Lauridsen T, Crump BC, Han S-K, Declerck S**. 2003. Rapid Screening for Freshwater Bacterial Groups by Using Reverse Line Blot Hybridization. Applied and Environmental Microbiology **69**:5875-5883.

2. **Newton RJ, Jones SE, Eiler A, McMahon KD, Bertilsson S**. 2011. A Guide to the Natural History of Freshwater Lake Bacteria. Microbiology and Molecular Biology Reviews : MMBR **75**:14-49.

3. **Eiler A, Bertilsson S**. 2004. Composition of freshwater bacterial communities associated with cyanobacterial blooms in four Swedish lakes. Environ Microbiol **6**:1228-43.

4. **Parveen B, Mary I, Vellet A, Ravet V, Debroas D**. 2013. Temporal dynamics and phylogenetic diversity of free-living and particle-associated Verrucomicrobia communities in relation to environmental variables in a mesotrophic lake. FEMS Microbiol Ecol **83**:189-201.

5. **Arnds J, Knittel K, Buck U, Winkel M, Amann R**. 2010. Development of a 16S rRNA-targeted probe set for Verrucomicrobia and its application for fluorescence in situ hybridization in a humic lake. Syst Appl Microbiol **33**:139-48.

6. **Hedlund BP, Gosink JJ, Staley JT**. 1997. Verrucomicrobia div. nov., a new division of the bacteria containing three new species of Prosthecobacter. Antonie van Leeuwenhoek **72**:29-38.

7. **Yoon J, Yasumoto-Hirose M, Katsuta A, Sekiguchi H, Matsuda S, Kasai H, Yokota A**. 2007. Coraliomargarita akajimensis gen. nov., sp. nov., a novel member of the phylum 'Verrucomicrobia' isolated from seawater in Japan. Int J Syst Evol Microbiol **57**:959-63.

8. **Yoon J, Matsuo Y, Katsuta A, Jang JH, Matsuda S, Adachi K, Kasai H, Yokota A**. 2008. Haloferula rosea gen. nov., sp. nov., Haloferula harenae sp. nov., Haloferula phyci sp. nov., Haloferula helveola sp. nov. and Haloferula sargassicola sp. nov., five marine representatives of the family Verrucomicrobiaceae within the phylum 'Verrucomicrobia'. Int J Syst Evol Microbiol **58**:2491-500.

9. **Sangwan P, Kovac S, Davis KE, Sait M, Janssen PH**. 2005. Detection and cultivation of soil verrucomicrobia. Appl Environ Microbiol **71**:8402-10.

10. **Brewer TE, Handley KM, Carini P, Gilbert JA, Fierer N**. 2016. Genome reduction in an abundant and ubiquitous soil bacterium 'Candidatus Udaeobacter copiosus'. **2**:16198.

11. **Qiu YL, Kuang XZ, Shi XS, Yuan XZ, Guo RB**. 2014. Terrimicrobium sacchariphilum gen. nov., sp. nov., an anaerobic bacterium of the class 'Spartobacteria' in the phylum Verrucomicrobia, isolated from a rice paddy field. Int J Syst Evol Microbiol **64**:1718-23.

12. **da Rocha UN, van Elsas JD, van Overbeek LS**. 2010. Real-time PCR detection of Holophagae (Acidobacteria) and Verrucomicrobia subdivision 1 groups in bulk and leek (Allium porrum) rhizosphere soils. Journal of Microbiological Methods **83**:141-148.

13. **Wertz JT, Kim E, Breznak JA, Schmidt TM, Rodrigues JLM**. 2012. Genomic and
Physiological Characterization of the Verrucomicrobia Isolate Diplosphaera
colitermitum gen. nov., sp. nov., Reveals Microaerophily and Nitrogen Fixation
Genes. Applied and Environmental Microbiology **78**:1544-1555.
14. **Derrien M, Vaughan EE, Plugge CM, de Vos WM**. 2004. Akkermansia muciniphila
gen. nov., sp. nov., a human intestinal mucin-degrading bacterium. Int J Syst Evol
Microbiol **54**:1469-76.
15. **Scheuermayer M, Gulder TA, Bringmann G, Hentschel U**. 2006. Rubritalea
marina gen. nov., sp. nov., a marine representative of the phylum 'Verrucomicrobia',
isolated from a sponge (Porifera). Int J Syst Evol Microbiol **56**:2119-24.
16. **Chin KJ, Liesack W, Janssen PH**. 2001. Opitutus terrae gen. nov., sp. nov., to
accommodate novel strains of the division 'Verrucomicrobia' isolated from rice
paddy soil. Int J Syst Evol Microbiol **51**:1965-8.
17. **Otsuka S, Suenaga T, Vu HT, Ueda H, Yokota A, Senoo K**. 2013. Brevifollis
gellanilyticus gen. nov., sp. nov., a gellan-gum-degrading bacterium of the phylum
Verrucomicrobia. Int J Syst Evol Microbiol **63**:3075-8.
18. **Sangwan P, Chen X, Hugenholtz P, Janssen PH**. 2004. Chthoniobacter flavus gen.
nov., sp. nov., the first pure-culture representative of subdivision two,
Spartobacteria classis nov., of the phylum Verrucomicrobia. Appl Environ Microbiol
70:5875-81.
19. **Hedlund BP, Gosink JJ, Staley JT**. 1996. Phylogeny of Prosthecobacter, the fusiform
caulobacters: members of a recently discovered division of the bacteria. Int J Syst
Bacteriol **46**:960-6.
20. **Otsuka S, Ueda H, Suenaga T, Uchino Y, Hamada M, Yokota A, Senoo K**. 2013.
Roseimicrobium gellanilyticum gen. nov., sp. nov., a new member of the class
Verrucomicrobiae. Int J Syst Evol Microbiol **63**:1982-6.
21. **Pol A, Heijmans K, Harhangi HR, Tedesco D, Jetten MS, Op den Camp HJ**. 2007.
Methanotrophy below pH 1 by a new Verrucomicrobia species. Nature **450**:874-8.
22. **Freitas S, Hatosy S, Fuhrman JA, Huse SM, Welch DB, Sogin ML, Martiny AC**.
2012. Global distribution and diversity of marine Verrucomicrobia. ISME J **6**:1499-
505.
23. **Cardman Z, Arnosti C, Durbin A, Ziervogel K, Cox C, Steen AD, Teske A**. 2014.
Verrucomicrobia are candidates for polysaccharide-degrading bacterioplankton in
an arctic fjord of Svalbard. Appl Environ Microbiol **80**:3749-56.
24. **Martinez-Garcia M, Brazel DM, Swan BK, Arnosti C, Chain PS, Reitenga KG, Xie
G, Poulton NJ, Lluesma Gomez M, Masland DE, Thompson B, Bellows WK,
Ziervogel K, Lo CC, Ahmed S, Gleasner CD, Detter CJ, Stepanauskas R**. 2012.
Capturing single cell genomes of active polysaccharide degraders: an unexpected
contribution of Verrucomicrobia. PLoS One **7**:e35314.
25. **Herlemann DP, Lundin D, Labrenz M, Jurgens K, Zheng Z, Aspeborg H,
Andersson AF**. 2013. Metagenomic de novo assembly of an aquatic representative
of the verrucomicrobial class Spartobacteria. MBio **4**:e00569-12.
26. **Paver SF, Kent AD**. 2010. Temporal patterns in glycolate-utilizing bacterial
community composition correlate with phytoplankton population dynamics in
humic lakes. Microb Ecol **60**:406-18.

27. **Haukka K, Kolmonen E, Hyder R, Hietala J, Vakkilainen K, Kairesalo T, Haario**
H, Sivonen K. 2006. Effect of nutrient loading on bacterioplankton community
composition in lake mesocosms. Microb Ecol **51**:137-46.

28. **Lindström ES, Vrede K, Leskinen E**. 2004. Response of a member of the
Verrucomicrobia, among the dominating bacteria in a hypolimnion, to increased
phosphorus availability. Journal of Plankton Research **26**:241-246.

29. **Kolmonen E, Sivonen K, Rapala J, Haukka K**. 2004. Diversity of cyanobacteria and
heterotrophic bacteria in cyanobacterial blooms in Lake Joutikas, Finland. Aquatic
Microbial Ecology **36**:201-211.

30. **Lindström ES, Kamst-Van Agterveld MP, Zwart G**. 2005. Distribution of Typical
Freshwater Bacterial Groups Is Associated with pH, Temperature, and Lake Water
Retention Time. Applied and Environmental Microbiology **71**:8201-8206.

31. **Schlesner H**. 1987. Verrucomicrobium spinosum gen. nov., sp. nov.: a fimbriated
prosthecate bacterium. Systematic and applied microbiology **10**:54-56.

32. **Zwart G, Huismans R, van Agterveld MP, Van de Peer Y, De Rijk P, Eenhoorn H,**
Muyzer G, van Hannen EJ, Gons HJ, Laanbroek HJ. 1998. Divergent members of
the bacterial division Verrucomicrobiales in a temperate freshwater lake. FEMS
Microbiology Ecology **25**:159-169.

33. **Read JS, Rose KC**. 2013. Physical responses of small temperate lakes to variation in
dissolved organic carbon concentrations. Limnology and Oceanography **58**:921-931.

34. **Shade A, Jones SE, McMahon KD**. 2008. The influence of habitat heterogeneity on
freshwater bacterial community composition and dynamics. Environmental
Microbiology **10**:1057-1067.

35. **Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW**. 2015. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. Genome Res **25**:1043-55.

36. **Schlesner H, Jenkins C, Staley J**. 2006. The Phylum Verrucomicrobia: A
Phylogenetically Heterogeneous Bacterial Group, p 881-896. *In* Dworkin M, Falkow
S, Rosenberg E, Schleifer K-H, Stackebrandt E (ed), The Prokaryotes doi:10.1007/0-
387-30747-8_37. Springer New York.

37. **Bendall ML, Stevens SLR, Chan L-K, Malfatti S, Schwientek P, Tremblay J,**
Schackwitz W, Martin J, Pati A, Bushnell B, Froula J, Kang D, Tringe SG,
Bertilsson S, Moran MA, Shade A, Newton RJ, McMahon KD, Malmstrom RR.
2016. Genome-wide selective sweeps and gene-specific sweeps in natural bacterial
populations. ISME J doi:10.1038/ismej.2015.241.

38. **Garcia SL, Stevens SLR, Crary B, Martinez-Garcia M, Stepanauskas R, Woyke T,**
Tringe SG, Andersson S, Bertilsson S, Malmstrom RR, McMahon KD. 2016.
Contrasting patterns of genome-level diversity across distinct co-occurring bacterial
populations. bioRxiv doi:10.1101/080168.

39. **Shieh WY, Jean WD**. 1998. Alterococcus agarolyticus, gen.nov., sp.nov., a halophilic
thermophilic bacterium capable of agar degradation. Can J Microbiol **44**:637-45.

40. **Kent AD, Jones SE, Yannarell AC, Graham JM, Lauster GH, Kratz TK, Triplett EW**.
2004. Annual Patterns in Bacterioplankton Community Variability in a Humic Lake.
Microbial Ecology **48**:550-560.

41. **Kent AD, Yannarell AC, Rusak JA, Triplett EW, McMahon KD**. 2007. Synchrony in
aquatic microbial community dynamics. ISME J **1**:38-47.

42. **Erbilgin O, McDonald KL, Kerfeld CA**. 2014. Characterization of a planctomycetal
organelle: a novel bacterial microcompartment for the aerobic degradation of plant
saccharides. Appl Environ Microbiol **80**:2193-205.
43. **Blanvillain S, Meyer D, Boulanger A, Lautier M, Guynet C, Denance N, Vasse J,**
Lauber E, Arlat M. 2007. Plant carbohydrate scavenging through tonB-dependent
receptors: a feature shared by phytopathogenic and aquatic bacteria. PLoS One
2:e224.
44. **Jiao G, Yu G, Zhang J, Ewart HS**. 2011. Chemical structures and bioactivities of
sulfated polysaccharides from marine algae. Mar Drugs **9**:196-223.
45. **Filali Mouhim R, Cornet J-F, Fontane T, Fournet B, Dubertret G**. 1993.
Production, isolation and preliminary characterization of the exopolysaccharide of
the cyanobacterium Spirulina platensis. Biotechnology Letters **15**:567-572.
46. **Dantas-Santos N, Gomes DL, Costa LS, Cordeiro SL, Costa MS, Trindade ES,**
Franco CR, Scortecci KC, Leite EL, Rocha HA. 2012. Freshwater plants synthesize
sulfated polysaccharides: heterogalactans from Water Hyacinth (Eicchornia
crassipes). Int J Mol Sci **13**:961-76.
47. **Beversdorf LJ, Miller TR, McMahon KD**. 2013. The role of nitrogen fixation in
cyanobacterial bloom toxicity in a temperate, eutrophic lake. PLoS One **8**:e56103.
48. **Khadem AF, van Teeseling MC, van Niftrik L, Jetten MS, Op den Camp HJ, Pol A**.
2012. Genomic and Physiological Analysis of Carbon Storage in the
Verrucomicrobial Methanotroph "Ca. Methylacidiphilum Fumariolicum" SolV. Front
Microbiol **3**:345.
49. **Mehta T, Coppi MV, Childers SE, Lovley DR**. 2005. Outer membrane c-type
cytochromes required for Fe(III) and Mn(IV) oxide reduction in Geobacter
sulfurreducens. Appl Environ Microbiol **71**:8634-41.
50. **Liu Y, Wang Z, Liu J, Levar C, Edwards MJ, Babauta JT, Kennedy DW, Shi Z,**
Beyenal H, Bond DR, Clarke TA, Butt JN, Richardson DJ, Rosso KM, Zachara JM,
Fredrickson JK, Shi L. 2014. A trans-outer membrane porin-cytochrome protein
complex for extracellular electron transfer by Geobacter sulfurreducens PCA.
Environmental Microbiology Reports **6**:776-785.
51. **Shi L, Fredrickson JK, Zachara JM**. 2014. Genomic analyses of bacterial porin-
cytochrome gene clusters. Frontiers in Microbiology **5**:657.
52. **Bucking C, Piepenbrock A, Kappler A, Gescher J**. 2012. Outer-membrane
cytochrome-independent reduction of extracellular electron acceptors in
Shewanella oneidensis. Microbiology **158**:2144-57.
53. **Shyu JBH, Lies DP, Newman DK**. 2002. Protective Role of tolC in Efflux of the
Electron Shuttle Anthraquinone-2,6-Disulfonate. Journal of Bacteriology **184**:1806-
1810.
54. **Voordeckers JW, Kim BC, Izallalen M, Lovley DR**. 2010. Role of Geobacter
sulfurreducens outer surface c-type cytochromes in reduction of soil humic acid and
anthraquinone-2,6-disulfonate. Appl Environ Microbiol **76**:2371-5.
55. **Lovley DR, Blunt-Harris EL**. 1999. Role of humic-bound iron as an electron
transfer agent in dissimilatory Fe(III) reduction. Appl Environ Microbiol **65**:4252-4.
56. **Lovley DR, Coates JD, Blunt-Harris EL, Phillips EJP, Woodward JC**. 1996. Humic
substances as electron acceptors for microbial respiration. Nature **382**:445-448.

57.   **Klupfel L, Piepenbrock A, Kappler A, Sander M**. 2014. Humic substances as fully
762   regenerable electron acceptors in recurrently anoxic environments. Nature Geosci
763   **7**:195-200.
764   58.   **Studholme DJ, Fuerst JA, Bateman A**. 2004. Novel protein domains and motifs in
765   the marine planctomycete Rhodopirellula baltica. FEMS Microbiol Lett **236**:333-40.
766   59.   **Kamneva OK, Knight SJ, Liberles DA, Ward NL**. 2012. Analysis of genome content
767   evolution in pvc bacterial super-phylum: assessment of candidate genes associated
768   with cellular organization and lifestyle. Genome Biol Evol **4**:1375-90.
769   60.   **Linz AM, Crary BC, Shade A, Owens S, Gilbert JA, Knight R, McMahon KD**. 2017.
770   Bacterial Community Composition and Dynamics Spanning Five Years in Freshwater
771   Bog Lakes. mSphere **2**.
772   61.   **Hamilton JJ, Garcia SL, Brown BS, Oyserman BO, Moya-Flores F, Bertilsson S,
773   Malmstrom RR, Forest KT, McMahon KD**. 2007. High-Throughput Metabolic
774   Network Analysis and Metatranscriptomics of a Cosmopolitan and Streamlined
775   Freshwater Lineage. bioRxiv doi:10.1101/106856.
776   62.   **Kang DD, Froula J, Egan R, Wang Z**. 2015. MetaBAT, an efficient tool for accurately
777   reconstructing single genomes from complex microbial communities. PeerJ **3**:e1165.
778   63.   **Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Pillay M, Ratner A,
779   Huang J, Woyke T, Huntemann M, Anderson I, Billis K, Varghese N, Mavromatis
780   K, Pati A, Ivanova NN, Kyrpides NC**. 2013. IMG 4 version of the integrated
781   microbial genomes comparative analysis system. Nucleic Acids Research
782   doi:10.1093/nar/gkt963.
783   64.   **Darling AE, Jospin G, Lowe E, Matsen IV FA, Bik HM, Eisen JA**. 2014. PhyloSift:
784   phylogenetic analysis of genomes and metagenomes. PeerJ **2**:e243.
785   65.   **Edgar RC**. 2004. MUSCLE: a multiple sequence alignment method with reduced time
786   and space complexity. BMC Bioinformatics **5**:113-113.
787   66.   **Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O**. 2010.
788   New algorithms and methods to estimate maximum-likelihood phylogenies:
789   assessing the performance of PhyML 3.0. Syst Biol **59**:307-21.
790   67.   **Spring S, Bunk B, Sproer C, Schumann P, Rohde M, Tindall BJ, Klenk HP**. 2016.
791   Characterization of the first cultured representative of Verrucomicrobia subdivision
792   5 indicates the proposal of a novel phylum. ISME J **10**:2801-2816.
793   68.   **Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y**. 2012. dbCAN: a web resource for
794   automated carbohydrate-active enzyme annotation. Nucleic Acids Res **40**:W445-51.
795   69.   **Yu CS, Chen YC, Lu CH, Hwang JK**. 2006. Prediction of protein subcellular
796   localization. Proteins **64**:643-51.
797   70.   **Yu NY, Wagner JR, Laird MR, Melli G, Rey S, Lo R, Dao P, Sahinalp SC, Ester M,
798   Foster LJ, Brinkman FSL**. 2010. PSORTb 3.0: improved protein subcellular
799   localization prediction with refined localization subcategories and predictive
800   capabilities for all prokaryotes. Bioinformatics **26**:1608-1615.
801   71.   **Bagos PG, Liakopoulos TD, Spyropoulos IC, Hamodrakas SJ**. 2004. PRED-TMBB:
802   a web server for predicting the topology of beta-barrel outer membrane proteins.
803   Nucleic Acids Res **32**:W400-4.
804
805

FIGURE LEGENDS

Fig. 1. Phylogenetic tree constructed with a concatenated alignment of protein sequences from five conserved essential single-copy genes (represented by TIGR01391, TIGR01011, TIGR00663, TIGR00460, and TIGR00362) that were recovered in all Verrucomicrobia MAGs. ME, TE and TH MAGs are labeled with red, green and blue, respectively. Genome ID in IMG or NCBI is indicated in the bracket. The outgroup is *Kiritimatiella glycovorans* L21-Fru-AB, which was initially assigned to subdivision 5, but this subdivision was recently proposed as a novel sister phylum to Verrucomicrobia (67).

Fig. 2. A. Coding densities of glycoside hydrolase genes, **B.** sulfatase genes, and **C.** Planctomycete-specific cytochrome *c* (PSCyt)-containing genes. Data from ME, TE and TH MAGs are labeled with red, green and blue, respectively. The three plots share the same *x*-axis label as indicated by the genome clustering on the top, which is based on a subtree extracted from the phylogenetic tree in Fig. 1 to indicate the phylogenetic relatedness of the 19 MAGs. The vertical dashed lines divide these MAGs to different subdivisions.

Fig. 3. Gene counts for the top 10 most abundant GH families, total gene counts for all GH families, and the number of GH families represented by these genes. MAGs are ordered as in the clustering in Fig. 2.

Fig. 4. Gene clusters encoding bacterial microcompartments (BMCs) involved in L-fucose

and L-rhamnose degradation. The vertical line indicates the end of a contig, and IMG gene

locus tag for the first gene in each presented gene cluster is indicated in the parentheses.

The BMC is schematically represented by a hexagon with the two building blocks labeled in

red and green, respectively. The two building blocks and reactions inside the BMC are

colored according to their encoding genes' color labels on the left side.

Fig. 5. Gene clusters encoding putative tonB-dependent carbohydrate utilization (CUT) loci.

IMG gene locus tag for the first gene in each presented gene cluster is indicated in the

parentheses. The horizontal solid lines below genes indicate predicted extracellular or

outer membrane proteins.

Fig. 6. Completeness estimates of key metabolic pathways. MAGs are ordered as in the

clustering in Fig. 2. Completeness value of "1" indicates a pathway is complete; "0" indicates

no genes were found in that pathway; and "(0)" indicates that although some genes in a

pathway are present, the pathway is likely absent because signature genes for that pathway

were not found in that draft genome AND signature genes are missing in more than two

thirds of all draft genomes.

Fig. 7. Gene clusters encoding putative porin-multiheme cytochrome *c* complex (PCC). IMG

gene locus tag for the first gene in each presented gene cluster is indicated in the

parentheses. The vertical line indicates the end of a contig, and horizontal lines below

genes indicate predicted cellular locations of their encoded proteins. These putative PCC

genes are in 18.1, 9.0, 6.1, 18.4, 70.0, 10.6 and 10.8 kbp long contigs, respectively. A

hypothesized model of extracellular electron transfer is shown on the right with yellow

arrows indicating electron flows. "IM" and "OM" refer to inner and outer membranes,

respectively, "ET in IM" refers to electron transfer in the inner membrane, and "$EA_{(ox)}$" and

"$EA_{(red)}$" refer to oxidized and reduced forms of the electron acceptor, respectively.

Fig. 8. Domain architecture and occurrence of PSCyt-containing genes. Based on the

combination of specific PSCyt and PSD domains, these domain structures can be classified

into three groups (I, II, and III). "CBM" refers to carbohydrate-binding modules, which

include pfam13385 (Laminin_G_3), pfam08531 (Bac_rhamnosid_N), pfam08305 (NPCBM),

pfam03422 (CBM_6), and pfam07691 (PA14). "PPI" refers to protein-protein interaction

domains, which include pfam02368 (Big_2), pfam00400 (WD40), and pfam00754

(F5_F8_type_C).

LIST OF SUPPLEMENTARY MATERIAL

Supplementary Text

Supplementary Figures S1 through S8

Supplementary Figure Legends

Fig. S1. A tiled display of an emergent self-organizing map (ESOM) based on the

tetranucleotide frequency (TNF) of the 19 Verrucomicrobia MAGs. TNF was calculated with

a window size of 5 kbp, with each dot on the ESOM representing a 5-kbp fragment (or a

contig if its length is shorter than 5 kbp). Dots (i.e. fragments) are colored according to

MAGs. A numeric ID is assigned to each MAG, and IDs from Mendota are labeled in black

and IDs from Trout Bog labeled in white. A red outline was drawn to indicate the clustering

of MAGs from Mendota on the ESOM.

Fig. S2. Counts of GH genes among the 78 different GH families present in MAGs**.**

Fig. S3. Heat map based on GH abundance profile patterns showing the clustering of MAGs

by different lakes.

Fig. S4. Counts of carbohydrate and amino acid transporter genes.

Fig. S5. Comparison of glycolate oxidase gene operons in *E. coli*, *C. flavus* and TE4605.

Fig. S6. Summary of important metabolic genes and pathways.

Fig. S7. Occurrence and gene organization of Planctomycetes-specific domains, DUF1501,

and DUF1552. (**a**) Counts of PSCyt, PSD, DUF1501, and DUF1552 domains in the MAGs. (**b**)

Clustering of PUF1501- and PSCyt2-containing genes, and clustering of PUF1552- and

PSCyt3-containing genes in the genome.

TABLE 1. Lakes included in this study[a]

Lake	Mendota	Trout Bog
GPS location	43.100°N, 89.405°W	46.041°N, 89.686°W
Lake type	Drainage lake	Seepage lake
Surface area (ha)	3938	1.1
Mean depth (m)	12.8	5.6
Max depth (m)	25.3	7.9
pH	8.3	5.2
Primary carbon source	Phytoplankton	Terrestrial subsidies
DOC (mg/L)	5.0	20.0
Total N (mg/L)	1.5	1.3
Total P (µg/L)	131	71
DOC/N	3.3	15.6
DOC/P	38.0	281.9
Trophic state	Eutrophic	Dystrophic

896 [a]Data from NTL-LTER https://lter.limnology.wisc.edu, averaged from the study years. DOC = Dissolved
897 organic carbon. N = Nitrogen. P = Phosphorus.

TABLE 2. Summary of Verrucomicrobia MAGs[a]

Genome	IMG Taxon OID	Subdivision	Recovered MAG Size (Mbp)[b]	Genome Completeness Estimate (%)[c]	Genome Contamination Estimate (%)[c]	GC Content (%)	Coding Base (%)	Gene Count	Normalized Coverage Depth[d]		
									Median	Mean	Coefficient of Variation (%)
ME3880	2582580573	1	1.6	70	2	58	90.9	1585	0.2	2.9	217
TH2746	2582580664	1	6.5	81	3	62	86.7	5430	3.3	4.7	82
ME12612	2582580523	1	2.2	79	3	59	89.0	2335	0.0	0.9	261
ME12173	2582580521	1	2.1	63	3	52	91.3	2070	0.0	0.8	583
TE4605	2582580638	1	4.7	91	0	59	91.1	4380	1.0	4.8	198
ME6381	2582580593	1	2.4	62	0	57	92.5	2221	0.0	0.4	285
ME8366	2582580607	2	3.6	87	5	63	87.4	3450	0.0	1.2	326
TH2747	2582580665	3	5.2	93	8	58	89.6	4846	1.8	2.8	99
TH3004	2582580668	3	4.5	93	6	57	91.4	3798	1.9	5.8	139
TH0989	2556921153	3	7.2	91	8	62	90.3	5583	6.1	6.3	61
TH2519	2593339181	4	1.8	69	2	42	94.3	1654	6.2	6.7	60
TE1800	2593339189	4	2.2	84	2	42	94.3	1998	10.8	11.3	77
TH4590	2582580688	4	3.3	87	1	65	90.7	3132	2.4	3.6	99
ME2014	2582580546	4	1.9	77	5	66	93.7	1700	1.0	3.3	174
ME12657	2582580524	4	1.9	81	7	68	94.0	1838	0.0	0.8	344
TE1301	2582580616	4	2.0	95	0	54	94.7	1943	4.1	14.2	187
TH4093	2582580682	Unclassified	4.7	77	6	48	86.5	3982	4.3	3.9	61
ME30509	2582580559	Unclassified	1.2	51	2	63	92.3	1160	0.0	0.5	479
TH4820	2582580691	Unclassified	3.0	56	3	63	86.7	2794	1.0	2.1	110

[a]MAGs from Lake Mendota are shaded.
[b]Recovered MAG size is the sum of the length of all contigs within a MAG.
[c]Genome completeness and contamination was estimated with checkM using Verrucomicrobia-specific marker gene sets.

[d]Normalized coverage depths of MAGs were calculated from the 94, 45, or 45 individual ME, TE, or TH metagenomes respectively, and were used to comparatively infer relative population abundance at the different sampling points. In addition to the median and mean coverage depths, the coefficient of variation is also shown to indicate variation among the sampling points.

Fig. 1. Phylogenetic tree constructed with a concatenated alignment of protein sequences from five conserved essential single-copy genes (represented by TIGR01391, TIGR01011, TIGR00663, TIGR00460, and TIGR00362) that were recovered in all Verrucomicrobia MAGs. ME, TE and TH MAGs are labeled with red, green and blue, respectively. Genome ID in IMG or NCBI is indicated in the bracket. The outgroup is *Kiritimatiella glycovorans* L21-Fru-AB, which was initially assigned to subdivision 5, but this subdivision was recently proposed as a novel sister phylum to Verrucomicrobia (67).

Fig. 2. Coding densities of glycoside hydrolase genes (a), sulfatase genes (b) and Planctomycete-specific cytochrome c (PSCyt)-containing genes (c). Data from ME, TE and TH MAGs are labeled with red, green and blue, respectively. The three plots share the same x-axis label indicated by the genome clustering on the top, which is is based on a subtree extracted from the phylogenetic tree in Fig. 1 to indicate the phylogenetic relatedness of the 19 MAGs. The vertical dashed lines divide these MAGs to different subdivisions.

GH	Main Activities	ME3880	TH2746	ME12612	ME12173	TE4605	ME6381	ME8366	TH2747	TH3004	TH0989	TH2519	TE1800	TH4590	ME2014	ME12657	TE1301	TH4093	ME30509	TH4820
GH29	α-fucosidases	2	22	0	1	0	0	0	17	19	24	4	5	6	0	0	0	14	3	13
GH2	β-galactosidases and other β-linked dimers	1	24	0	1	2	0	0	18	14	20	5	5	5	0	0	0	6	2	7
GH78	α-L-rhamnosidases	1	31	0	2	0	0	0	11	11	13	3	5	9	0	0	0	5	2	7
GH95	1,2-α-L-fucosidase	0	29	0	1	0	0	0	7	9	17	1	2	3	0	0	0	4	0	7
GH106	α-L-rhamnosidase	0	21	1	0	0	0	0	4	9	12	1	2	1	0	0	0	3	1	10
GH13	α-amylase	3	3	2	2	3	2	1	7	6	7	2	3	2	2	2	2	3	0	3
GH20	β-hexosaminidase	0	14	0	4	0	0	0	3	5	2	2	2	3	1	2	4	10	1	3
GH5	endoglucanase, endomannanase, β-glucosidase, β-mannosidase	0	4	0	1	4	1	3	13	4	10	0	0	0	0	0	1	6	1	6
GH28	polygalacturonases, related to pectin degradation	0	1	0	0	0	0	0	7	11	2	8	8	3	1	0	1	0	1	3
GH43	α-L-arabinofuranosidases, endo-α-L-arabinanases, β-D-xylosidases	1	4	0	2	2	0	0	11	15	9	1	1	2	0	0	0	0	0	2
Counts of other GH genes		9	86	7	18	29	10	10	95	83	119	12	15	32	19	10	13	68	11	77
Total counts of all GH genes		17	239	10	32	40	13	14	193	186	235	39	48	66	23	14	21	119	22	138
Total number of GH families represented		12	48	7	23	23	7	10	53	49	58	19	21	26	10	10	13	35	15	45

Fig. 4. Gene counts for the top 10 most abundant GH families, total gene counts for all GH families, and the number of GH families represented by these genes. MAGs are ordered as in the clustering in Fig. 2.

Fig. 4. Gene clusters encoding bacterial microcompartments (BMCs) involved in L-fucose and L-rhamnose degradation. The vertical line indicates the end of a contig, and IMG gene locus tag for the first gene in each presented gene cluster is indicated in the parenthesis. The BMC is schematically represented by a hexagon with the two building blocks labeled in red and green, respectively. The two building blocks and reactions inside the BMC are colored according to their encoding genes' color labels on the left side.

Fig. 5. Gene clusters encoding putative tonB-dependent carbohydrate utilization (CUT) loci. IMG gene locus tag for the first gene in each presented gene cluster is indicated in the parenthesis. The horizontal solid lines below genes indicate predicted extracellular or outer membrane proteins.

Genes and Pathways	ME3880	TH2746	ME12612	ME12173	TE4605	ME6381	ME8366	TH2747	TH3004	TH0989	TH2519	TE1800	TH4590	ME2014	ME12657	TE1301	TH4093	ME30509	TH4820
Central carbon metabolism																			
Glycolysis (Embden-Meyerhof pathway), glucose => pyruvate	0.78	0.89	0.89	0.67	1	0.89	0.78	1	0.89	1	0.67	0.89	1	0.89	0.56	1	0.89	0.56	0.33
Glycolysis (Entner-Doudoroff pathway)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)
Pentose phosphate pathway (Pentose phosphate cycle)	0.86	0.71	0.71	0.71	0.86	0.57	0.71	1	0.86	1	0.71	0.86	1	0.86	0.57	1	0.71	0.29	0.43
Pyruvate oxidation, pyruvate => acetyl-CoA	1	1	1	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Citrate cycle (TCA cycle, Krebs cycle)	0.63	0.88	1	0.38	0.75	0.63	0.88	0.88	0.88	0.75	1	1	0.88	0.63	0.75	1	0.5	0.25	0.25
Other carbohydrate metabolism																			
Galactose degradation to glycerate-3P	0.5	0.75	0.75	0.25	0.75	0.5	0.5	1	0.75	1	0.5	0.5	0.75	0.25	0.25	0.5	0.75	0.25	0.5
Rhamnose degradation	1	1	0.75	0.5	0.5	0.5	0.5	1	1	1	0.75	1	0.75	0.5	0.5	0.5	1	0.5	0.75
Fucose degradation	1	0.75	0.75	0.25	0.75	0.5	0.75	1	0.75	0.75	0.5	0.75	0.75	0.75	0.5	0.75	1	0.75	0.75
L-Arabinose degradation to xylulose-P for pentose pathway	1	1	0	0	0.67	0.33	0.33	1	1	1	0	0	0.33	0	0	0	0.67	0.33	0.33
Xylose degradation	1	1	1	1	1	0.5	1	1	1	1	0.5	1	1	1	1	1	0.5	1	1
D-Galacturonate degradation to pyruvate & D-glyceraldehyde 3P	(0)	(0)	(0)	(0)	0.8	0.2	(0)	0.6	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)
D-Glucuronate degradation to pyruvate and D-glyceraldehyde 3P	0	0.6	0.4	0.4	0.6	0	0.2	0.6	0.8	0.6	0.6	0.8	0.6	0.2	0.4	0.4	0.6	0.4	0.8
Mannose degradation to glucose-P	0.5	1	1	1	1	1	1	1	1	1	1	1	1	0.5	0.5	1	1	0.5	1
Lactaldehyde degradation to pyruvate (Aerobic)	1	1	0.33	1	0.33	0.33	1	1	1	0.67	0.33	0.33	1	0.33	0.33	0.33	0.67	0.67	0.67
Glycogen biosynthesis from alpha-D-glucose-6P via ADP-D glucose	0.75	0.25	0.75	0.75	0.75	0.5	1	1	0.75	1	0.5	0.5	1	1	1	1	0.5	0	0.75
Fermentation																			
Pyruvate to acetate via acetyl-coA	1	1	0.67	0.67	0.33	0.33	1	1	1	0.67	0.33	0.33	1	0.67	0.67	0.67	1	1	1
Pyruvate to propanoate	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)
Pyruvate to succinate	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)
Pyruvate to butanoate	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)
Pyruvate to butanol	0.71	0.71	0.71	0.29	1	0.71	0.57	0.43	0.43	0.43	0.57	0.57	1	0.57	0.57	0.57	0.43	0.43	0.43
Pyruvate to ethanol	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)
Pyruvate to lactate	0	0	0	0	0	0	0	0	0	0	1	1	1	0	0	0	0	0	0
Pyruvate to acetone	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)
Nitrogen related																			
Dissimilatory nitrate reduction, nitrate => ammonia	0	0	0	0	0	0	0	0	0	0	0	0	(0)	0	0	0	0	0	(0)
Denitrification, nitrate => nitrogen gas	0	0	0	0	0	0	0	0.25	0.25	0	0	0	0	0	0	0	0	0	0
TMAO (trimethylamine-N-oxide) reductioin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nitrification, ammonia => nitrite	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nitrogen fixation, nitrogen => ammonia	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	0	0
Assimilatory nitrate reduction, nitrate => ammonia	0	1	0	0	(0)	0	0	0	0	0	0	0	0	0	0	0	(0)	0	(0)
ABC-type urea transporter	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0
Urease (Urea ==> CO2 + NH3)	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0
Ammonia permease	1	1	1	1	1	1	1	1	1	1	0	0	1	0	1	1	1	0	1
Phosphorus related																			
Alkaline phosphatase (PhoA)	1	1	1	1	0	1	1	0	0	0	0	0	0	1	1	1	1	0	0
ABC-type phosphate-specific transport (Pst) system, high-affinity	0	1	1	1	1	1	1	1	1	1	1	1	1	0	1	1	1	1	0
PiT inorganic phosphate transporter (PitA), permease, low-affinity	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	0	0	1
Polyphosphate storage and utilization (PPK)	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	0
ABC-type phosphonate transport system	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Phosphonoacetate degradation	0	1	0	1	1	0	1	1	1	0	0	0	1	0	0	0	0	0	0

Fig. 6. Completeness estimates of key metabolic pathways. MAGs are ordered as in the clustering in Fig. 2. Completeness value of "1" indicates a pathway is complete; "0" indicates no genes were found in that pathway; and "(0)" indicates that although some genes in a pathway are present, the pathway is likely absent because signature genes for that pathway were not found in that draft genome AND signature genes are missing in more than two thirds of all draft genomes.

Fig. 7. Gene clusters encoding putative porin-multiheme cytochrome c complex (PCC). IMG gene locus tag for the first gene in each presented gene cluster is indicated in the parenthesis. The vertical line indicates the end of a contig, and horizontal lines below genes indicate predicted cellular locations of their encoded proteins. These putative PCC genes are in 18.1, 9.0, 6.1, 18.4, 70.0, 10.6 and 10.8 kbp long contigs, respectively. A hypothesized model of extracellular electron transfer is shown on the right with yellow arrows indicating electron flows. "IM" and "OM" refer to inner and outer membranes, respectively, "ET in IM" refers to electron transfer in the inner membrane, and "EA$_{(ox)}$" and "EA$_{(red)}$" refer to oxidized and reduced forms of the electron acceptor, respectively.

Fig. 8. Domain architecture and occurrence of PSCyt-containing genes. Based on the combination of specific PSCyt and PSD domains, these domain structures can be classified into three groups (I, II, and III). "CBM" refers to carbohydrate-binding modules, which include pfam13385 (Laminin_G_3), pfam08531 (Bac_rhamnosid_N), pfam08305 (NPCBM), pfam03422 (CBM_6), and pfam07691 (PA14). "PPI" refers to protein-protein interaction domains, which include pfam02368 (Big_2), pfam00400 (WD40), and pfam00754 (F5_F8_type_C).