Addressing the looming identity crisis in single cell RNA-seq

Megan Crow, Anirban Paul, Sara Ballouz, Z. Josh Huang, Jesse Gillis[*]

Cold Spring Harbor Laboratory, One Bungtown Road, Cold Spring Harbor, NY 11724, USA

mcrow@cshl.edu, paula@cshl.edu, sballouz@cshl.edu, huangj@cshl.edu, jgillis@cshl.edu

*corresponding author

Abstract

Single cell RNA-sequencing technology (scRNA-seq) provides a new avenue to discover and characterize cell types, but the experiment-specific technical biases and analytic variability inherent to current pipelines may undermine the replicability of these studies. Meta-analysis of rapidly accumulating data is further hampered by the use of *ad hoc* naming conventions. Here we demonstrate our replication framework, MetaNeighbor, that allows researchers to quantify the degree to which cell types replicate across datasets, and to rapidly identify clusters with high similarity for further testing. We first measure the replicability of neuronal identity by comparing more than 13 thousand individual scRNA-seq transcriptomes, then assess cross-dataset evidence for novel pyramidal neuron and cortical interneuron subtypes identified by scRNA-seq. We find that 24/45 cortical interneuron subtypes and 10/48 pyramidal neuron subtypes have evidence of replication in at least one other study. Identifying these putative replicates allows us to re-analyze the data for differential expression and provide lists of robust candidate marker genes. Across tasks we find that large sets of variably expressed genes can identify replicable cell types and subtypes with high accuracy, indicating many of the transcriptional changes characterizing cell identity are pervasive and easily detected.

Introduction

Single cell RNA-sequencing (scRNA-seq) has emerged as an important new technology enabling the dissection of heterogeneous biological systems into ever more refined cellular components. One popular application of the technology has been to try to define novel cell subtypes within a given tissue or within an already refined cell class, as in the lung (Treutlein et al., 2014), pancreas (Baron et al., 2016; Muraro et al., 2016; Segerstolpe et al., 2016; Wang et al., 2016), retina (Macosko et al., 2015; Shekhar et al., 2016), or others (Grun et al., 2015; Klein et al., 2015; Min et al., 2015). Because they aim to discover completely new cell subtypes, the majority of this work relies on unsupervised clustering, with most studies using customized pipelines with many unconstrained parameters, particularly in their inclusion criteria and statistical models (Grun et al., 2015; Habib et al., 2016; Macosko et al., 2015; Zeisel et al., 2015). While there has been steady refinement of these techniques as the field has come to appreciate the biases inherent to current scRNA-seq methods, including prominent batch effects (Hicks et al., 2015), expression drop-outs (Lun et al., 2016; Pierson and Yau, 2015), and the complexities of normalization given differences in cell size or cell state (Buettner et al., 2015; Vallejos et al., 2015), the question remains: how well do novel transcriptomic cell subtypes replicate across studies?

In order to answer this, we turned to the issue of cell diversity in the brain, a prime target of scRNA-seq as neuron diversity is critical for construction of the intricate, exquisite circuits underlying brain function. The heterogeneity of brain tissue makes it particularly important that results be assessed for replicability, while its popularity as a target of study makes this goal particularly feasible. Because a primary aim of neuroscience has been to derive a taxonomy of cell types (Ascoli et al., 2008), already more than twenty single cell RNA-seq experiments have been performed using mouse nervous tissue (Poulin et al., 2016). Remarkable strides have been made to address fundamental questions about the diversity of cells in the nervous system,

including efforts to describe the cellular composition of the cortex and hippocampus (Tasic et

al., 2016; Zeisel et al., 2015), to exhaustively discover the subtypes of bipolar neurons in the

retina (Shekhar et al., 2016), and to characterize similarities between human and mouse

midbrain development (La Manno et al., 2016). In spite of this wealth of data, there have been

few attempts to compare, validate and substantiate cell type transcriptional profiles across

scRNA-seq datasets, and no systematic or formal method has been developed for

accomplishing this task.

To address this gap in the field, we propose a simple, supervised framework, MetaNeighbor

(**meta**-analysis via **neighbor** voting), to assess how well cell type-specific transcriptional profiles

replicate across datasets. Our basic rationale is that if a cell type has a biological identity rooted

in the transcriptome then knowing its expression features in one dataset will allow us to find

cells of the same type in another dataset. We make use of the cell type labels supplied by data

providers, and assess the correspondence of cell types across datasets by taking the following

approach (see schematic, Figure 1):

1) First we construct a kernel: we calculate correlations between all pairs of cells that we

aim to compare across datasets based on the expression pattern of a set of genes. This

generates a network where each cell is a node and the edges are the strength of the

correlations between them.

2) Next, we do cross-dataset validation: we hide all cell type labels ('identity') for one

dataset at a time. This dataset will be used as our test set. Cells from all other datasets

remain labeled, and are used as the training set.

3) Finally, we predict the cell type labels of the test set: we use a neighbor voting algorithm

to predict the identity of the held-out cells based on their similarity to the training data.

Conceptually, this resembles approaches for the validation of sample clustering (Dudoit et al.,

2002; Kapp and Tibshirani, 2007) but it has been adapted to operate from within a supervised

learning framework. This permits both systematic scoring and carefully defined control

experiments to investigate the data features that drive high performance. Our implementation is

extremely fast and robust to technical differences between experiments; because prediction is

performed only within an individual dataset at a time, we are able to keep many aspects of

technical variation constant. This essentially controls for any dataset specific effects that would

otherwise swamp the subtler cell identity signal. The method provides a score that indicates the

degree to which a cell type replicates for each gene set that is tested. This means that

MetaNeighbor doubles as a low-tech 'feature selection tool' that we can use to identify the

transcriptional features that are most discriminative between cell types. By comparing the

scores returned from using Gene Ontology (GO) functions ("functional gene sets") or sets of

randomly chosen genes ("random gene sets"), we can determine whether co-expression of

specific gene sets is characteristic of particular cell types, and thus important for cell function or

identity.

We evaluate cell identity by taking sequential steps according to the basic taxonomy of brain

cells: first classifying neurons vs. non-neuronal cells across eight single cell RNA-seq studies,

then classifying cortical inhibitory neurons vs. excitatory neurons, and for our final step, we align

interneuron and pyramidal cell subtypes across three studies. Critically, we discover that that

almost any sufficiently large and highly variable set of genes can be used to distinguish between

cell types, suggesting that cell identity is widely represented within the transcriptome.

Furthermore, we find that cross-dataset analysis of pyramidal neurons results in broad definition

of cortical vs. hippocampal types, and find evidence for the replication of five layer-restricted

subtypes. In contrast, we find that cortical interneuron subtypes show clear lineage-specific

structure, and we readily identify 11 subtypes that replicate across datasets, including

Chandelier cells and five novel subtypes defined by transcriptional clustering in previous work.

Meta-analysis of differential expression across these highly replicable cortical interneuron

subtypes revealed evidence for canonical marker genes such as parvalbumin and somatostatin,

as well as new candidates which may be used for improved molecular genetic targeting, and to

understand the diverse phenotypes and functions of these cells.

Assessing neuronal identity with MetaNeighbor

We aimed to measure the replicability of cell identity across tasks of varying specificity.

Broadly, these are divided into tasks where we are recapitulating known cell identities, and ones

where are measuring the replicability of novel cell identities discovered in recent research. The

former class of task is the focus of this subsection: first, by assessing how well we could

distinguish neurons from non-neuronal cells ("task one"), and next assessing the discriminability

of excitatory and inhibitory neurons ("task two"). As detailed in the methods, MetaNeighbor

outputs a performance score for each gene set and task. This score is the mean area under the

receiver operator characteristic curve (AUROC) across all folds of cross-dataset validation, and

it can be interpreted as the probability that we will rank a positive higher than a negative (e.g.

neuron vs. non-neuronal cell, when using neurons as the positive label set) based on the

expression of a set of genes. This varies between 0 and 1, with 1 being perfect classification,

0.5 meaning that we have performed as well as if we had randomly guessed the cell's identity,

and 0.9 or above being extremely high. Comparison of scores across gene sets allows us to

discover their relative importance for defining cell identity.

As described above, in task one we assessed how well we could identify neurons and non-

neuronal cells across eight datasets with a total of 13928 cells (Table S1). Although this was

designed to be fairly simple, we were surprised to find that AUROC scores were significantly

higher than chance for all gene sets tested, including all randomly chosen sets ($AUROC_{all}$

$_{sets}$=0.78 ± 0.1, Figure 2A). Reassuringly, a bootstrapped sampling of the datasets showed a

trend toward increased performance with the inclusion of additional training data, indicating that

we are recognizing an aggregate signal across datasets (Figure S1). However, the significant

improvement of random sets over the null means that prior knowledge about gene function is

not required to differentiate between these cell classes. Randomly chosen sets of genes have

decidedly non-random expression patterns that enable discrimination between cell types.

Task two aimed to assess how well we could discriminate between cortical excitatory and

inhibitory neurons across four studies with a total of 2809 excitatory and 1162 inhibitory neurons

(Dueck et al., 2015; Habib et al., 2016; Tasic et al., 2016; Zeisel et al., 2015). Similar to our

previous results, we saw that AUROC scores were significantly higher than chance

(AUROC=0.69 ± 0.1, Figure 2B), suggesting that transcriptional differences are likely to be

encoded in a large number of genes.

Consistent with the view that a large fraction of transcripts are useful for determining cell

identity, we found a positive dependency of AUROC scores on gene set size, regardless of

whether genes within the sets were randomly selected or shared some biological function

(Figure 2B). This was further supported by a comparison of scores for task one using 100 sets

of either 100 or 800 randomly chosen genes. AUROC score distributions and means were

significantly different, with sets of 100 genes having lower scores but higher variability in

performance, whereas sets of 800 genes were more restricted in variance and gave higher

performance on average (Figure 2C, $AUROC_{100}$=0.80 ± 0.05, $AUROC_{800}$=0.90 ± 0.03, p<2.2E-

16, Wilcoxon rank sum test). The variability in performance observed while keeping set size

constant suggests that even in random sets, there are transcriptional features that contribute to

cell identity. We delved into this further by comparing AUROC scores across gene sets chosen

based on their mean expression as we have previously shown that this is a critical factor to

control for in evaluating single cell gene co-expression (Crow et al., 2016). We performed task

one again using expression-level based gene sets and found a strong positive relationship

between expression level and our ability to classify cells (Figure 2D, r_s=0.9).

These results provide evidence that MetaNeighbor can readily identify cells of the same type

across datasets, without relying on specific knowledge of marker genes. In these two examples,

all cells could be classified as one of two types, making this a binary classification task. We find

that a gene set's size and mean expression level are the key features that allow for cell type

discrimination in this setting.

Investigating cortical interneuron subtypes using MetaNeighbor

Cortical inhibitory interneurons have diverse characteristics based on their morphology,

connectivity, electrophysiology and developmental origins, and it has been an ongoing goal to

define cell subtypes based on these properties (Ascoli et al., 2008). In a related paper (Paul et

al., submitted), we describe the transcriptional profiles of GABAergic interneuron types which

were targeted using a combinatorial strategy including intersectional marker gene expression,

cell lineage, laminar distribution and birth timing, and have been extensively phenotyped both

electrophysiologically and morphologically (He et al., 2016). Previously, two studies were

published in which new interneuron subtypes were defined based on scRNA-seq transcriptional

profiles (Tasic et al., 2016; Zeisel et al., 2015). These found different numbers of subtypes (16

in one and 23 in the other), and the authors of the later paper compared their outcomes by

looking at the expression of a handful of marker genes, which yielded mixed results: a small

number of cell types seemed to have a direct match but for others the results were more

conflicting, with multiple types matching to one another, and others having no match at all. Here

we aimed to more quantitatively assess the similarity of their results, and compare them with our

own data which derives from phenotypically characterized sub-populations; i.e., not from

unsupervised expression clustering (see Table S2 for sample information).

MetaNeighbor relies on coordinated variation in expression level to detect cell identity, which

means that genes with high variability are particularly useful. Our preceding binary

classifications showed that genes with high mean expression were more likely to have variation

that allowed MetaNeighbor to learn cell identities. In the following analyses, we are examining

both rare and common cell types across datasets. In this case, the mean expression level of

marker genes should be a proxy for cell incidence: we can expect that the marker expression for

a more abundant type would have a higher mean expression. Since variance scales with

expression, the most highly variable genes in the dataset would likely only be discriminative for

the abundant type. Because we would like to be able to identify both abundant and rare cell

types, we select the genes with the highest variance at each mean expression level.

We identified 638 genes with high variability given their expression levels (detailed in Methods)

and these were used as a 'high variability gene set' to measure AUROC scores between each

pair of cells across datasets. When AUROCs were measured using all genes, we saw that

clustering was subject to strong lab-specific effects (Figure S2). In contrast, the use of variable

genes reproduced the known subtype structure, with major branches for the three main

subtypes, Pv, Sst and Htr3a.

To examine how the previously identified interneuron subtypes are represented across the three

studies, we tested the similarity of each pair of subtypes both within and across datasets using

the high variability gene set. For each genetically-targeted interneuron type profiled by Paul et

al., we found at least one corresponding subtype from the other two studies, which were defined

by having a mean AUROC score across training/testing folds >0.95 (Figure 3). This includes

Chandelier cells, a subtype that could not be definitively identified by either Tasic or Zeisel.

Using our reciprocal testing and training protocol we find that the Tasic_Pvalb Cpne5 subtype

are likely to be Chandelier cells (AUROC=0.99). In addition, expanding our criteria to include all

reciprocal best matches in addition to those with ID scores >0.95, we found correspondence

among five subtypes that were assessed only in the Tasic and Zeisel data,

Tasic_Smad3/Zeisel_Int14 (AUROC=0.97), Tasic_Sncg/Zeisel_Int6 (AUROC=0.95),

Tasic_Ndnf-Car4/Zeisel_Int15 (AUROC=0.95), Tasic_Igtp/Zeisel_Int13 (AUROC=0.94) and

Tasic_Ndnf-Cxcl14/Zeisel_Int12 (AUROC=0.91). Overall, based on this high-variance gene set,

we could identify 11 subtypes representing 24/45 (53%) types (Figure 3A), with total n for each

subtype ranging from 25-189 out of 1583 interneurons across all datasets (1.5-11%). These

results were robust to differences in data processing. Tasic *et al.* provided data as both RPKM

and TPM values, and while thousands of genes had extremely divergent expression between

the two, including some key markers like Vip, reciprocal average AUROCs among

corresponding subtypes were nearly identical (Figure S3). Our corresponding subtypes also

confirm the marker gene analysis performed by Tasic *et al.* (Table S3), without requiring manual

gene curation. Because we quantify the similarity among types we can prioritize matches, and

use these as input to MetaNeighbor for further evaluation.

In the above, we identified overlaps using a single gene set. To assess cell identification more

broadly, we ran MetaNeighbor with these new across-dataset subtype labels, measuring

predictive validity across all gene sets in GO (Figure 3A, far right). The distribution of AUROC

scores varied across subtypes but we found that the score from the high variability gene set was

representative of overall trends, with high performing groups showing higher mean AUROC

scores over many gene sets. As detailed in the previous section, we note that AUROC scores

are sensitive both to the number of training samples (n) and to underlying data features (e.g.,

transcriptome complexity), which complicates direct comparison of ID score distributions. Both

the high mean AUROCs across all putative replicate subtypes (>0.6), and the similarity of

maximum performance suggest that distinctive gene co-expression can be observed in each

subtype (max AUROC=0.92 ± 0.04). As with previous tasks, we found little difference in average

AUROCs using functional gene sets compared to random sets (mean AUROC$_{Random}$=0.67 ±

0.06, mean AUROC$_{GO}$=0.68 ± 0.1).

These results indicate that highly variable gene sets can be used alongside pairwise testing and

training as a heuristic to identify replicable subtypes.

Investigating pyramidal neuron subtypes using MetaNeighbor

The heterogeneity of pyramidal neurons is undisputed, but the organizing principles are still

debated, with some suggesting that identity is discrete and modular (Habib et al., 2016; Zeisel

et al., 2015) and others purporting that identities are more likely to be described by expression

gradients or spectra (Cembrowski et al., 2016). With MetaNeighbor we are able to quantitatively

assess the degree to which pyramidal subtypes defined by scRNA-seq replicate across diverse

datasets. If cell types are discrete and modular, we would expect to see sharp differences, with

some types showing very strong similarity to one another, and strong dissimilarities to other

types.

To compare pyramidal neuron scRNA-seq datasets we permuted through all combinations of

subtypes as testing and training data based on a set of 743 genes with high variability given

their expression level (subtypes listed in Table S2). This was the same procedure that was used

for cortical interneurons and while there were similar numbers of subtypes in total, a smaller

fraction corresponded across datasets (10/48, ~21%) yielding five putative subtypes (Figure

3B). The AUROC score heatmap was generally less modular than the heatmap of interneuron

scores. The most prominent feature was that types from the hippocampus and cortex tended to

cluster separately from one another. Within each region-specific cluster some layer- or area-

specific clustering was observed but it was not completely consistent. Particular discrepancy

was observed between the cortical layer 5 subtypes which showed more similar AUROC score

profiles to the hippocampal subtypes than to other deep layer types (Tasic L5b_Cdh13,

L5_Chrna6, L5b_Tph). Note that these were also the same subtypes that Tasic *et al.* found no

match for in their marker gene analysis. We suggest that the inclusion of additional datasets

may help to resolve this inconsistency.

We assessed the five putative subtypes using MetaNeighbor. All subtypes were significantly

discernable compared to the null (Figure 3B) and as with the interneuron subtypes, AUROC

scores from the high variability gene set were well correlated with mean performance across all

of GO (3888 gene sets). In line with previous tasks, we found that functional gene sets

performed equally to random gene sets (mean $AUROC_{Random}=0.71 \pm 0.08$, $AUROC_{GO}=0.70 \pm$

0.09).

Comparing gene set performance across tasks

Finally, we compared gene set results from the 11 replicate interneuron subtypes and the 5

pyramidal neuron subtypes. In agreement with our previous results, we found that the top

groups were all related to neuronal function, which is unsurprising given the large size of these

gene sets and their likelihood of expression and variation in these cells (Figure 3C). AUROCs

were highly correlated across tasks (r~0.76), with slightly higher performance for identifying

interneuron types compared to pyramidal types (Figure 3D). The linearity of the trend across all

scores suggests that fundamental data features, like mean expression level and set size,

underlie the differential discriminative value of gene sets. The high performance across many

sets (mean AUROC ~0.7) also supports the notion that cell identity is encoded promiscuously

across the transcriptome, and is not restricted to a small set of functionally important genes.

Identifying subtype specific genes

ScRNA-seq experiments often seek to define marker genes for novel subtypes. Though ideally

marker genes are perfectly discriminative with respect to all cells, in practice marker genes are

often contextual and defined relative to a particular out-group. Here we aimed to identify

possible marker genes that would allow discrimination among interneuron subtypes or

pyramidal neuron subtypes. For each of our identified replicate subtypes we generated a ranked

list of possible marker genes by performing one-tailed, non-parametric differential expression

analysis within each study for all subtypes (e.g., Int1 vs. all other interneurons in the Zeisel

study, Int2 vs. all interneurons, etc.) and combining p-values for replicated types using Fisher's

method (Table S4). Figure 4A shows the FDR adjusted p-values for the top candidates based

on fold change for the ten replicated interneuron subtypes with overlapping differential

expression patterns. Figure 4B shows the same for the two pyramidal neuron subtypes with

overlapping differential expression patterns. The majority of these genes have previously been

characterized as having some degree of subtype- or layer-specific expression, for example we

readily identify genes that were used for the Cre-driver lines in the Tasic and Paul studies (*Sst,*

Pvalb, Vip, Cck, Htr3a, Ctgf). Even though we filtered for genes with high fold changes, we see

that many genes are differentially expressed in more than one subtype. Notably, considerable

overlap can be observed among the *Htr3a*-expressing types. For example, the Vip Sncg

subtype (Tasic_Vip Sncg/Paul_Vip Cck) is only subtly different from the Sncg subtype

(Tasic_Sncg/Zeisel_Int6) across this subset of genes, with the Sncg cells lacking differential

expression of *Cxcl14* and *Nr2f2*.

We also identify some novel candidates, including *Ptn,* or pleiotrophin, which is significantly

more expressed in the three Nos1-expressing subtypes than in the others (Figure 4B). It is thus

expected to be discriminative of Nos1-positive neurons compared to other interneuron types.

We validated *Ptn* expression with *in situ* hybridization and we show clear expression in neurons

that are positive for both *Sst* and *Nos1* (Figure 4C). *Ptn* is a growth factor, and we suggest that

its expression may be required for maintaining the long-range axonal connections that

characterize these cells. These cells are well described by current markers, however this

approach is likely to be of particular value for novel subtypes that lack markers, allowing

researchers to prioritize genes for follow-up by assessing robustness across multiple data

sources.

Discussion

Single-cell transcriptomics promises to have a revolutionary impact by enabling comprehensive

sampling of cellular heterogeneity; nowhere is this variability more profound than within the

brain, making it a particular focus of both single-cell transcriptomics and our own analysis into

its replicability. The substantial history of transcriptomic analysis and meta-analysis gives us

guidance about bottlenecks that will be critical to consider in order to characterize cellular

heterogeneity. The most prominent of these is laboratory-specific bias, likely deriving from the

adherence to a strict set of internal standards, which may filter for some classes of biological

signal (e.g., poly-A selection) or induce purely technical grouping (e.g., by sequencing depth).

Because of this, it is imperative to be able to align data across studies and determine what is

replicable. In this work, we have provided a formal means of determining replicable cell identity

by treating it as a quantitative prediction task. The essential premise of our method is that if a

cell type has a distinct transcriptional profile within a dataset, then an algorithm trained from that

data set will correctly identify the same type within an independent data set.

The currently available data allowed us to draw a number of conclusions. We validated the

discrete identity of eleven interneuron subtypes, and described replicate transcriptional profiles

to prioritize possible marker genes, including *Ptn*, a growth factor that is preferentially expressed

in Sst Chodl cells. We performed a similar assessment for pyramidal neurons but found less

correspondence among datasets, suggesting that additional data will be required to determine

whether there is evidence for discrete pyramidal neuron types. One major surprise of our

analysis is the degree of replicability in the current data. Our AUROC scores are exceptionally

high, particularly when considered in the context of the well-described technical confounds of

single-cell data. We suspect this reflects the fundamental nature of the biological problem we

are facing: discrete cell types can be identified by their transcriptional profiles, and the biological

clarity of the problem overcomes technical variation.

This is further suggested by our result that cell identity has promiscuous effects within

transcriptional data. While in-depth investigation of the most salient gene functions is required to

characterize cell types, to simply identify cell types is relatively straightforward. This is

necessarily a major factor in the apparent successes of unsupervised methods in determining

novel cell types and suggests that cell type identity is clearly defined by transcriptional profiles,

regardless of cell selection protocols, library preparation techniques or fine-tuning of clustering

algorithms. To us this result recalls the startling finding by Venet *et al.* that "Most random gene

expression signatures are related to breast cancer outcome" (Venet et al., 2011). Where, until

that point, research had often focused on demonstrating that highly specific genes or gene

clusters could predict breast cancer outcome, Venet et al. clearly demonstrated that this was a

more straightforward task than targeted analyses would reveal, and was due to the strength of

the underlying biological signal: more aggressive cancers divide more, and so anything

correlated with fast cycling times will be associated with poor clinical outcomes. Comparison of

transcriptional signatures between different cell types provides an equally clear lens. Many gene

sets show more correlated expression within than across types, and variation across types is

likely to be accounted for by simple important factors, like cell size. This is not to say that more

detailed characterization of cell types is not necessary: understanding the differences between

cells and how they work will require focused investigation into the precise molecular players that

are differentially utilized. However, we hope that this helps to demonstrate that the variations on

dimension reduction and clustering methods in single cell RNA-seq are 'working', inevitably by

taking advantage of this very clear signal.

In this work we opted to use the subtype or cluster labels provided by the original authors, in

essence to characterize both the underlying data as well as current analytic practices. However,

this has limitations where studies cluster to different levels of specificity. For example, the Tasic

paper defines multiple Parvalbumin subtypes but the Zeisel and Paul work do not. Our method

makes it extremely easy to identify highly overlapping types at the levels defined by each

author, facilitating downstream work to validate the sub-clusters through meta-analysis and at

the bench. Given the known noisiness of single-cell expression and the complex and

idiosyncratic character of approaches taken to assessing it, the degree of replicability that we

see is much higher than could have been expected were there not simple explanations for the

derived clusters from individual laboratories. Our work shows that with additional data,

comprehensive evaluation and replication is likely to be quantitatively straightforward, making it

possible to have high confidence in derived cell sub-types quite rapidly. As this additional data is

generated, our approach can provide consistent updates of the field-wide consensus.

The simplicity of our method makes it unlikely to be biased toward the exact cell identity tasks

assessed here. For example, because of the method's reliance on relative ranks, it is almost

entirely immune to normalization as a potential confound. On the one hand, this limits our

sensitivity to detect real signals of some type, but this cost is more than offset by the robustness

of signals identified. Its simplicity also means that it is scalable, and readily admits to the

incorporation of data from individual labs in their ongoing work. Ultimately we hope that by

defining what is replicable clearly, MetaNeighbor will allow future studies involving cell-cell

comparisons to build on a strong foundation toward a comprehensive delineation of cell types.

Experimental Procedures

Animals, manual cell sorting and scRNA-seq

Mice were bred and cared for in accordance with animal husbandry protocols at Cold Spring

Harbor Laboratory, with access to food and water ad libitum and a 12 hour light-dark cycle.

Transgenic animals bred to target the six phenotypically characterized subpopulations were

generated using the following breeding strategies (detailed in He et al): Nkx2.1-CreER, Pv-ires-

Cre animals were bred separately to Ai14 reporter to label ChC and Pv cells in the cortex. ChCs

were enriched in frontal cortex with tamoxifen induction at embryonic day 17.5. Intersectional

labeling was achieved by breeding (a) Sst-Flp, Nos1-CreER, (b) Sst-Flp, CR-Cre, (c) Vip-Flp,

CR-Cre and (d) Vip-Flp, Cck-Cre separately to the Ai65 intersectional reporter that labels cells

with tdTomato only when both the lox-STOP-lox and frt-STOP-frt cassettes are excised. Adult

animals (P28-35) were sacrificed by cervical dislocation to harvest brains for single cell sorting.

Cell sorting and scRNA-seq were performed as described previously (Crow et al., 2016). Single

cells were collected by manual sorting then placed into single tubes with 1μl total volume of

RNAseOUT (Invitrogen), 1:400K diluted ERCC spike-in RNA, and sample-specific RT primers.

Cells were flash frozen in liquid nitrogen then stored at -80ºC until processed. RNA was linearly

amplified using the MessageAmp-II kit (Life Technologies) according to the manufacturer's

recommended protocol. Reverse transcription of amplified aRNA was done with SuperScript-III

(Invitrogen) and cDNA libraries were prepared with the Illumina TruSeq small RNA library

preparation kit (7-11 cycles of PCR). Libraries were size-selected with SPRISelect magnetic

beads (Agencourt) and sequenced with paired-end 101bp reads using an Illumina HiSeq.

PolyA-primed reads were mapped to the mouse reference genome (mm9) with Bowtie (v

0.12.7), while paired sequences were used for varietal tag counting. A custom python script was

used map and tally sequences with unique tags for each mRNA in each cell (Crow et al., 2016).

All data is available to download from GEO (accession GSE92522).

Public expression data

Data analysis was performed in R using custom scripts (github.com/maggiecrow/MetaNeighbor,

2016). Processed expression data tables were downloaded from GEO directly, then subset to

genes appearing on both Affymetrix GeneChip Mouse Gene 2.0 ST array (902119) and the

UCSC known gene list to generate a merged matrix containing all samples from each

experiment. The mean value was taken for all genes with more than one expression value

assigned. Where no gene name match could be found, a value of 0 was input. We considered

only samples that were explicitly labeled as single cells, and removed cells that expressed fewer

than 1000 genes with expression >0. Cell type labels were manually curated using sample

labels and metadata from GEO (see Tables S1 and S2). Merged data and metadata are linked

through our Github page.

Gene sets

Gene annotations were obtained from the GO Consortium 'goslim_generic' (August 2015).

These were filtered for terms appearing in the GO Consortium mouse annotations

'gene_association.mgi.gz' (December 2014) and for gene sets with between 20-1000 genes,

leaving 106 GO groups with 9221 associated genes. Random gene sets were generated by

randomly choosing genes with the same set size distribution as GO slim. Sets of high variance

genes were generated by binning data from each dataset into deciles based on expression

level, then making lists of the top 25% of the most variable genes for each decile, excluding the

most highly expressed bin. The high variance set was then defined as the intersect of the high

variance gene lists across the relevant datasets.

MetaNeighbor

All scripts, sample data and detailed directions to run MetaNeighbor in R can be found on our

Github page (github.com/maggiecrow/MetaNeighbor, 2016).

The input to MetaNeighbor is a set of genes, a data matrix and two sets of labels: one set for

labeling each experiment, and one set for labeling the cell types of interest. For each gene set,

the method generates a cell-cell similarity network by measuring the Spearman correlation

between all cells across the genes within the set, then ranking and standardizing the network so

that all values lie between 0 and 1. The use of rank correlations means that the method is

robust to any rank-preserving normalization (i.e., log2, TPM, RPKM). Ranking and standardizing

the networks ensures that distributions remain uniform across gene sets, and diminishes the

role outlier similarities can play since values are constrained.

The node degree of each cell is defined as the sum of the weights of all edges connected to it

(i.e., the sum of the standardized correlation coefficients between each cell and all others), and

this is used as the null predictor in the neighbor voting algorithm to standardize for a cell's 'hub-

ness': cells that are generically linked to many cells are preferentially down-weighted, whereas

those with fewer connections are less penalized. For each cell type assessment, the neighbor

voting predictor produces a weighted matrix of predicted labels by performing matrix

multiplication between the network and the binary vector (0,1) indicating cell type membership,

then dividing each element by the null predictor (i.e., node degree). In other words, each cell is

given a score equal to the fraction of its neighbors, including itself, which are part of a given cell

type (Ballouz et al., 2016). For cross-validation, we permute through all possible combinations

of leave-one-dataset-out cross-validation, sequentially hiding each experiment's cell labels in

turn, and then reporting how well we can recover cells of the same type as the mean area under

the receiver operator characteristic curve (AUROC) across all folds. A key difference from

conventional cross-validation is that there is no labeled data within the dataset for which

predictions are being made. Labeled data comes only from external datasets, ensuring

predictions are driven by signals that are replicable across data sources. To improve speed,

AUROCs are calculated analytically, where the AUROC for each cell type *j*, is calculated based

on the sum of the ranks of the scores for each cell *i*, belonging to that cell type. This can be

expressed as follows:

$$AUROC_j = \sum_{i}^{N} \frac{Ranks_i}{N * N_{Neg}} - \frac{N + 1}{2 * N_{Neg}}$$

where N is the number of true positives, and N_{Neg} is the number of true negatives. Note that for

experiments with only one cell type this cannot be computed as there are no true negatives.

AUROCs are reported as averages across all folds of cross-validation for each gene set

(excluding NAs from experiments with no negatives), and the distribution across gene sets is

plotted.

To test the dependency of results on the amount of training and testing data we repeated the

neuron vs. non-neuronal cell discrimination task after randomly selecting between two and

seven datasets ten times each. This was done for 21 representative gene sets. Means for each

gene set and each number of included datasets were plotted.

Identifying putative replicates

In cases where cell identity was undefined across datasets (i.e., cortical interneuron and

pyramidal subtypes) we treated each subtype label as a positive for each other subtype, and

assessed similarity over the high variance gene set described above. For example, Int1 from the

Zeisel dataset was used as the positive (training) set, and all other subtypes were considered

the test set in turn. Mean AUROCs from both testing and training folds are plotted in the

heatmap in Figure 3. A stringent cut-off of mean AUROC >0.95 and/or mutual best matches

across datasets identified putative replicated types for further assessment with our supervised

framework (detailed above). While lowering this threshold could increase the number of

subtypes with some match, we found that reciprocal top hits alone provided an upper bound on

the number of replicated types (i.e., lowering the thresholds did not allow for a higher number of

subtypes). New cell type labels encompassing these replicate types (e.g. a combined Sst-Chodl

label containing Int1 (Zeisel), Sst Chodl (Tasic) and Sst Nos1 (Paul)) were generated for

MetaNeighbor across random and GO sets, and for meta-analysis of differential expression.

While only reciprocal top-hits across laboratories were used to define novel cell-types,

conventional cross-validation within laboratories was performed to fill in AUROC scores across

labels contained within each lab.

Differential expression

For each cell type within a dataset (defined by the authors' original labeling), differential gene

expression was calculated using a one-sided Wilcoxon rank-sum test, comparing gene

expression within a given cell type to all other cells within the dataset (e.g., Zeisel_Int1 vs all

other Zeisel interneurons). Meta-analytic p-values were calculated for each putative replicated

type using Fisher's method (Fisher, 1925) then a multiple hypothesis test correction was

performed with the Benjamini-Hochberg method (Benjamini and Hochberg, 1995). Top

differentially expressed genes were those with an adjusted meta-analytic p-value <0.001 and

with log2 fold change >2 in each dataset. All differential expression data for putative replicated

subtypes can be found in Table S4.

Supplementary Material

473   Supplementary tables and figures may be accessed at the following link: http://bit.ly/2s58zPd

Author Contributions

475   JG conceived the study. JG, MC, and JH designed experiments. MC and JG wrote the

476   manuscript. MC and SB performed computational experiments. AP performed cell sorting, and

477   generated and parsed the raw sequencing data. JH supervised wet-lab data collection. All

478   authors read and approved the final manuscript.

Acknowledgments

MC, SB and JG were supported by a gift from T. and V. Stanley. Z.J.H. was supported by NIH

5R01MH094705-04, R01MH109665-01 and the CSHL Robertson Neuroscience Fund. A.P. was

supported by a NARSAD Postdoctoral Fellowship. The authors would like to thank Paul

Pavlidis, Bo Li and Jessica Tollkuhn for their thoughtful feedback on earlier drafts of this

manuscript. We would also like to thank the dedicated researchers who have made their data

publicly available. Our work would not be possible without their valuable contributions.

References

Ascoli, G.A., Alonso-Nanclares, L., Anderson, S.A., Barrionuevo, G., Benavides-Piccione, R.,
Burkhalter, A., Buzsaki, G., Cauli, B., Defelipe, J., Fairen, A*., et al.* (2008). Petilla terminology:
nomenclature of features of GABAergic interneurons of the cerebral cortex. Nat Rev Neurosci *9*,
557-568.

Ballouz, S., Weber, M., Pavlidis, P., and Gillis, J. (2016). EGAD: ultra-fast functional analysis of
gene networks. Bioinformatics (Oxford, England).

Baron, M., Veres, A., Wolock, Samuel L., Faust, Aubrey L., Gaujoux, R., Vetere, A., Ryu,
Jennifer H., Wagner, Bridget K., Shen-Orr, Shai S., Klein, Allon M*., et al.* (2016). A Single-Cell
Transcriptomic Map of the Human and Mouse Pancreas Reveals Inter- and Intra-cell Population
Structure. Cell Systems *3*, 346-360.e344.

Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series B
(Methodological) *57*, 289-300.

Buettner, F., Natarajan, K.N., Casale, F.P., Proserpio, V., Scialdone, A., Theis, F.J., Teichmann,
S.A., Marioni, J.C., and Stegle, O. (2015). Computational analysis of cell-to-cell heterogeneity in
single-cell RNA-sequencing data reveals hidden subpopulations of cells. Nature biotechnology
33, 155-160.

Cembrowski, M.S., Bachman, J.L., Wang, L., Sugino, K., Shields, B.C., and Spruston, N.
(2016). Spatial Gene-Expression Gradients Underlie Prominent Heterogeneity of CA1 Pyramidal
Neurons. Neuron *89*, 351-368.

Crow, M., Paul, A., Ballouz, S., Huang, Z.J., and Gillis, J. (2016). Exploiting single-cell
expression to characterize co-expression replicability. Genome biology *17*, 101.

Dudoit, S., Fridlyand, J., and Speed, T.P. (2002). Comparison of Discrimination Methods for the
Classification of Tumors Using Gene Expression Data. Journal of the American Statistical
Association *97*, 77-87.

Dueck, H., Khaladkar, M., Kim, T.K., Spaethling, J.M., Francis, C., Suresh, S., Fisher, S.A.,
Seale, P., Beck, S.G., Bartfai, T*., et al.* (2015). Deep sequencing reveals cell-type-specific
patterns of single-cell transcriptome variation. Genome biology *16*, 122.

Fisher, R.A. (1925). Statistical methods for research workers (Edinburgh, London,: Oliver and
Boyd).

github.com/maggiecrow/MetaNeighbor (2016). MetaNeighbor: a method to rapidly assess cell
type identity using both functional and random gene sets.

Grun, D., Lyubimova, A., Kester, L., Wiebrands, K., Basak, O., Sasaki, N., Clevers, H., and van
Oudenaarden, A. (2015). Single-cell messenger RNA sequencing reveals rare intestinal cell
types. Nature *525*, 251-255.

Habib, N., Li, Y., Heidenreich, M., Swiech, L., Avraham-Davidi, I., Trombetta, J.J., Hession, C.,
Zhang, F., and Regev, A. (2016). Div-Seq: Single-nucleus RNA-Seq reveals dynamics of rare
adult newborn neurons. Science (New York, NY) *353*, 925-928.

He, M., Tucciarone, J., Lee, S., Nigro, M.J., Kim, Y., Levine, J.M., Kelly, S.M., Krugikov, I., Wu,
P., Chen, Y.*, et al.* (2016). Strategies and Tools for Combinatorial Targeting of GABAergic
Neurons in Mouse Cerebral Cortex. Neuron *91*, 1228-1243.

Hicks, S.C., Teng, M., and Irizarry, R.A. (2015). On the widespread and critical impact of
systematic bias and batch effects in single-cell RNA-Seq data.

Kapp, A.V., and Tibshirani, R. (2007). Are clusters found in one dataset present in another
dataset? Biostatistics (Oxford, England) *8*, 9-31.

Klein, A.M., Mazutis, L., Akartuna, I., Tallapragada, N., Veres, A., Li, V., Peshkin, L., Weitz,
D.A., and Kirschner, M.W. (2015). Droplet barcoding for single-cell transcriptomics applied to
embryonic stem cells. Cell *161*, 1187-1201.

La Manno, G., Gyllborg, D., Codeluppi, S., Nishimura, K., Salto, C., Zeisel, A., Borm, L.E., Stott,
S.R., Toledo, E.M., Villaescusa, J.C.*, et al.* (2016). Molecular Diversity of Midbrain Development
in Mouse, Human, and Stem Cells. Cell *167*, 566-580.e519.

Lun, A.T., Bach, K., and Marioni, J.C. (2016). Pooling across cells to normalize single-cell RNA
sequencing data with many zero counts. Genome biology *17*, 75.

Macosko, E.Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M., Tirosh, I., Bialas,
A.R., Kamitaki, N., Martersteck, E.M.*, et al.* (2015). Highly Parallel Genome-wide Expression
Profiling of Individual Cells Using Nanoliter Droplets. Cell *161*, 1202-1214.

Min, J.W., Kim, W.J., Han, J.A., Jung, Y.J., Kim, K.T., Park, W.Y., Lee, H.O., and Choi, S.S.
(2015). Identification of Distinct Tumor Subpopulations in Lung Adenocarcinoma via Single-Cell
RNA-seq. PLoS One *10*, e0135817.

Muraro, Mauro J., Dharmadhikari, G., Grün, D., Groen, N., Dielen, T., Jansen, E., van Gurp, L.,
Engelse, Marten A., Carlotti, F., de Koning, Eelco J.P.*, et al.* (2016). A Single-Cell
Transcriptome Atlas of the Human Pancreas. Cell Systems *3*, 385-394.e383.

Peng, R. (2016). A Simple Explanation for the Replication Crisis in Science. In Simply Statistics.

Pierson, E., and Yau, C. (2015). ZIFA: Dimensionality reduction for zero-inflated single-cell gene
expression analysis. Genome biology *16*, 241.

Poulin, J.-F., Tasic, B., Hjerling-Leffler, J., Trimarchi, J.M., and Awatramani, R. (2016).
Disentangling neural cell diversity using single-cell transcriptomics. Nature neuroscience *19*,
1131-1141.

Segerstolpe, A., Palasantza, A., Eliasson, P., Andersson, E.M., Andreasson, A.C., Sun, X.,
Picelli, S., Sabirsh, A., Clausen, M., Bjursell, M.K.*, et al.* (2016). Single-Cell Transcriptome
Profiling of Human Pancreatic Islets in Health and Type 2 Diabetes. Cell metabolism *24*, 593-
607.

Shekhar, K., Lapan, S.W., Whitney, I.E., Tran, N.M., Macosko, E.Z., Kowalczyk, M., Adiconis,
X., Levin, J.Z., Nemesh, J., Goldman, M.*, et al.* (2016). Comprehensive Classification of Retinal
Bipolar Neurons by Single-Cell Transcriptomics. Cell *166*, 1308-1323.e1330.

Tasic, B., Menon, V., Nguyen, T.N., Kim, T.K., Jarsky, T., Yao, Z., Levi, B., Gray, L.T.,
Sorensen, S.A., Dolbeare, T.*, et al.* (2016). Adult mouse cortical cell taxonomy revealed by
single cell transcriptomics. Nature neuroscience *19*, 335-346.

Treutlein, B., Brownfield, D.G., Wu, A.R., Neff, N.F., Mantalas, G.L., Espinoza, F.H., Desai, T.J.,
Krasnow, M.A., and Quake, S.R. (2014). Reconstructing lineage hierarchies of the distal lung
epithelium using single-cell RNA-seq. Nature *509*, 371-375.

Vallejos, C.A., Marioni, J.C., and Richardson, S. (2015). BASiCS: Bayesian Analysis of Single-
Cell Sequencing Data. PLoS Comput Biol *11*, e1004333.

Venet, D., Dumont, J.E., and Detours, V. (2011). Most Random Gene Expression Signatures
Are Significantly Associated with Breast Cancer Outcome. PLoS Comput Biol *7*, e1002240.

Wang, Y.J., Schug, J., Won, K.-J., Liu, C., Naji, A., Avrahami, D., Golson, M.L., and Kaestner,
K.H. (2016). Single cell transcriptomics of the human endocrine pancreas. Diabetes.

Zeisel, A., Munoz-Manchado, A.B., Codeluppi, S., Lonnerberg, P., La Manno, G., Jureus, A.,
Marques, S., Munguba, H., He, L., Betsholtz, C.*, et al.* (2015). Brain structure. Cell types in the
mouse cortex and hippocampus revealed by single-cell RNA-seq. Science (New York, NY) *347*,
1138-1142.

Figures

Figure 1 – MetaNeighbor quantifies cell type identity across experiments
A – Schematic representation of gene set co-expression across individual cells. Cell types are indicated by their color. **B –** Similarity between cells is measured by taking the correlation of gene set expression between individual cells. On the top left of the panel, gene set expression between two cells, A and B, is plotted. There is a weak correlation between these cells. On the bottom left of the panel we see the correlation between cells A and C, which are strongly correlated. By taking the correlations between all pairs of cells we can build a cell network (right), where every node is a cell and the edges represent how similar each cell is to each other cell. **C -** The cell network that was generated in B can be extended to include data from multiple experiments (multiple datasets). The generation of this multi-dataset network is the first step of MetaNeighbor. **D –** The cross-validation and scoring scheme of MetaNeighbor is demonstrated in this panel. To assess cell type identity across experiments we use neighbor voting in cross-validation, systematically hiding the labels from one dataset at a time. Cells within the hidden dataset are predicted as similar to the cell types from other datasets, using a neighbor voting formalism. Whether these scores prioritize cells as the correct type within the dataset determines the performance, expressed as the AUROC. In other words, comparative assessment of cells occurs only within a dataset, but this is based only on training information from outside that dataset. This is then repeated for all gene sets of interest.

Figure 2 – Cell type identity is widely represented in the transcriptome
A & B – Distribution of AUROC scores from MetaNeighbor for discriminating neurons from non-neuronal cells
("task one", A) and for distinguishing excitatory vs. inhibitory neurons ("task two", B). GO scores are in black
and random gene set scores are plotted in gray. Dashed grey lines indicate the null expectation for correctly
guessing cell identity (AUROC=0.5). For both tasks, almost any gene set can be used to improve performance
above the null, suggesting widespread encoding of cell identity across the transcriptome. **C** – Task one
AUROC scores for each gene set are plotted with respect to the number of genes. A strong, positive
relationship is observed between gene set size and AUROC score, regardless of whether genes were chosen
randomly or based on shared functions. **D** – Distribution of AUROC scores for task one using 100 sets of 100
randomly chosen genes, or 800 randomly chosen genes. The mean AUROC score is significantly improved
with the use of larger gene sets (mean 100 = 0.80 +/- 0.05, mean 800 = 0.90 +/- 0.03). **E** – Relationship
between AUROC score and expression level. Task one was re-run using sets of genes chosen based on mean
expression. A strong positive relationship was observed between expression level and performance (r_s ~0.9).

Figure 3 – Cross-dataset analysis of interneuron and pyramidal neuron diversity

A – (Left) Heatmap of AUROC scores between interneuron subtypes based on the highly variable gene kernel. Dendrograms were generated by hierarchical clustering of Euclidean distances using average linkage. Row colors indicate data origin and column colors show marker expression. Clustering of AUROC score profiles recapitulates known cell type structure, with major branches representing the Pv, Sst and Htr3a lineages. (Middle) Table of reciprocal best matches and subtype pairs with scores >0.95. (Right) Boxplots of GO performance (3888 sets) for each replicated subtype, ordered by their AUROC score from the highly variable gene set. Subtypes are labeled with the names from Tasic *et al*. A positive relationship is observed between AUROC scores from the highly variable set and the average AUROC score for each subtype. Mean AUROCs are all greater than chance (0.5) suggesting robust cross-dataset replication across gene sets. **B** – (Left) Heatmap of AUROC scores between pyramidal subtypes based on the highly variable gene kernel, clustered as in A. Row colors indicate datasets and column colors show brain region, cortical layer or hippocampal area. Clustering of AUROC score profiles shows a separation of cortical and hippocampal subtypes. (Middle) Table of reciprocal best matches. (Right) Boxplots of GO performance (3888 sets) for each replicated subtype, ordered by their AUROC score from the highly variable gene set. Subtypes are labeled by layer. A positive relationship is observed between ID scores from the highly variable set and the average AUROC for each subtype. **C** – The table shows the top GO terms that allow for cross-dataset subtype discrimination, listed by their mean AUROC across tasks. For both tasks, high scores are obtained for terms related to neuronal function. **D** – AUROC scores for each GO function are plotted, with pyramidal scores on the y-axis and interneuron scores on the x-axis. AUROCs are highly correlated across tasks (r_s~0.76), suggesting limited functional specificity.

Figure 4 – Replicated subtypes show consistent differential expression
A – (Top) Heatmap of FDR adjusted p-values of top differentially expressed genes among replicated interneuron subtypes (NB only ten subtypes are shown as no differentially expressed genes were found for the Ndnf Car4 subtype). Subtype names are listed at the top of the columns and are labeled as in Tasic *et al*. Many genes are commonly differentially expressed among multiple subtypes, but combinatorial patterns distinguish them. (Right) Heatmap of FDR adjusted p-values of top differentially expressed genes among replicated pyramidal neuron subtypes. (NB only the two with overlapping differential expression are shown). Subtypes are labeled by layer. **B** – Standardized Ptn expression is plotted across the three experiments, where each box represents an interneuron subtype. High, but variable expression is observed across the three Sst Chodl types. **C** – Fluorescent double in-situ of Ai14/tdTomato driven by Sst-Flp and Nos-Cre expression (green) and Ptn (red). Dotted box indicates the area shown in higher magnification on the right, arrowheads point to cells that express both transcripts.