1 **TET-mediated epimutagenesis of the *Arabidopsis thaliana* methylome**
2

3 Lexiang Ji[a,*], William T. Jordan[b,*], Xiuling Shi[b], Lulu Hu[c], Chuan He[c], Robert J. Schmitz[b]

4
5 [a]Institute of Bioinformatics, University of Georgia, Athens, GA, 30602, USA.
6
7 [b]Department of Genetics, University of Georgia, Athens, GA, 30602, USA.
8
9 [c]Department of Chemistry, Department of Biochemistry and Molecular Biology, and
10 Institute for Biophysical Dynamics, Howard Hughes Medical Institute, University of
11 Chicago, Chicago, IL 60637, USA
12
13

*Indicates these authors contributed equally to this work

Corresponding author:

Robert J. Schmitz

University of Georgia

120 East Green Street, Athens, GA 30602

Telephone: (706) 542-1887

Fax: (706) 542-3910

E-mail: schmitz@uga.edu

Keywords: DNA methylation, TET, epigenetics, epigenomics, epimutagenesis

25
26

DNA methylation in the promoters of plant genes sometimes leads to transcriptional repression, and the wholesale removal of DNA methylation as seen in methyltransferase mutants results in drastic changes in gene expression and severe developmental defects. However, many cases of naturally-occurring DNA methylation variations have been reported, whereby the altered expression of differentially methylated genes is responsible for agronomically important traits. The ability to manipulate plant methylomes to generate populations of epigenetically distinct individuals could provide invaluable resources for breeding and research purposes. Here we describe "epimutagenesis", a novel method to rapidly generate variation of DNA methylation through random demethylation of the *Arabidopsis thaliana* genome. This method involves the expression of a human Ten-eleven translocation (TET) enzyme, and results in widespread hypomethylation that can be inherited to subsequent generations, mimicking mutants in the maintenance DNA methyltransferase *met1*. Application of TET-mediated epimutagenesis to agriculturally significant plants may result in differential expression of alleles normally silenced by DNA methylation, uncovering previously hidden phenotypic variations.

Our ability to develop novel beneficial crop traits has significantly improved over the last 100 years, although the ability to maintain this trajectory is limited by allelic diversity. While genetic variation has been heavily exploited for crop improvement, utility of epigenetic variation has yet to be efficiently implemented. Epigenetic variation arises not from a change in the DNA sequence, but by changes in modifications to DNA such as cytosine methylation. This variation can result in emergence of novel and stably inherited phenotypes, as well as unique patterns of gene expression.

In plant genomes, cytosine methylation occurs at three major sequence contexts:

CG, CHG and CHH (where H = A, C or T)[1]. Methylation at these different contexts is

coordinated by distinct maintenance mechanisms during DNA replication. The

methylation of DNA in all three contexts is essential for transcriptional silencing of

transposons, repeat sequences and certain genes. Genes regulated by this mechanism

are stably repressed throughout the soma and represent an untapped source of hidden

genetic variation if transcriptionally re-activated, as revealed from pioneering studies in

the model plant *A. thaliana*[2, 3, 4]. However, the impact of this variation is not observed in

wild-type plants, as genes silenced by DNA methylation are not expressed. This novel

source of genetic variation was uncovered by creating epigenetic recombinant inbred

lines (epiRILs) from crosses between a wild-type individual and a mutant defective in

maintenance of DNA methylation[2, 3, 4]. EpiRILs, while genetically wild type, contain

mosaic DNA methylomes dependent on chromosomal inheritance patterns, as DNA

methylation is meiotically inherited in *A. thaliana*[2, 5, 6, 7, 8]. Phenotypic characterization of

epiRILs has revealed extensive morphological variation with respect to traits such as

flowering time, root length and resistance to bacterial infection[2, 3, 4]. The morphological

variation generated by the creation of epiRILs has revealed extensive hidden genetic

variation in plant genomes that can be observed due to expression of newly

unmethylated regions. However, the creation of epiRILs requires one founding parent to

be a null mutant in the maintenance DNA methylation pathway. Unfortunately, unlike in

A. thaliana, the loss of DNA methylation maintenance activity often results in lethality in

crops[9, 10]. Therefore, novel methodologies are required to realize the potential of these

hidden epialleles in crop genomes.

Epimutagenesis is an alternative method to generate epiRILs. Instead of relying

on the genome-wide demethylation of one of the two founding parents, epimutagenesis

introduces random methylation variation via the introduction of a transgene. Here, we

describe a novel epimutagenesis approach in *A. thaliana* using a human Ten-eleven

translocation (TET1) methylcytosine dioxygenase[11, 12, 13, 14], which catalyzes the

conversion of 5-methylcytosine (5mC) to 5-hydroxymethylcytosine (5hmC). Although

TET enzymes and their primary product 5hmC are not found in plant genomes[15], ectopic

expression of a human TET enzyme resulted in widespread DNA demethylation and

induced phenotypic variation in *A. thaliana*.

RESULTS

Overexpressing hTET1cd in Arabidopsis hypomethylates the genome

Transgenic *A. thaliana* plants were generated expressing the catalytic domain (residues

1418-2136) of the human TET1 protein (hTET1cd) under the control of the CaMV35S

promoter. To assess the impact of hTET1cd expression on the *A. thaliana* methylome,

whole genome bisulfite sequencing (WGBS) was performed on two independently

derived transgenic plants (35S:TET1-1 and 35S:TET1-2; **Supplementary Table 1**).

WGBS revealed a global reduction of CG methylation from 18.2% in two wild-type

individuals to 8.9% in 35S:TET1-1 and 6.9% in 35S:TET1-2 (compared to 0.5% in *met1-*

3). The effects of hTET1cd expression on CHG and CHH methylation were not as

severe compared to CG methylation (**Fig. 1a**). Importantly, different degrees of CG

hypomethylation were observed in different independent transgenic plants. This result

has important implications for epimutagenesis in economically and agriculturally

significant plant species, as it appears feasible to control the degree of DNA

hypomethylation by screening for plants with desired levels of demethylation. Taken

together, these results show that the expression of hTET1cd results in intermediate CG

methylation levels when compared to wild-type and *met1* individuals.

The primary product of TET1 oxidation is 5hmC, which is indistinguishable from

5mC by WGBS. We therefore performed Tet-assisted bisulfite sequencing (TAB-seq) to

profile 5hmC levels in 35S:TET1 plants[16]. No detectable levels of 5hmC were found in

the transgenic lines assayed (**Supplementary Fig. 1a-b)**. Thus, the widespread loss of

CG DNA methylation observed may result from a failure to maintain methylation at CG

sites that possess 5hmC, or through active removal of 5hmC or further oxidized products

via the base excision repair pathway.

To better understand the effects of hTET1cd expression, we determined changes

in the *A. thaliana* methylome at the chromosomal and local levels. Plotting methylation

levels across all five chromosomes revealed a strong depletion of CG methylation at the

pericentromeric region (**Fig. 1b**). CG hypomethylation occurred at both gene body

methylated (gbM) and select RNA-directed DNA methylated (RdDM) loci. (**Fig. 1c and**

d). To further quantify the observed hypomethylation, metaplots were created for genes

and transposons, respectively (**Fig. 1e and f, Supplementary Fig. 1c-f**). Strong

reduction of mCG and a mild reduction of mCHG/mCHH were observed at both genes

and transposons. On average, 97.9% of gbM genes and 56.7% of methylated

transposons (where these regions have at least 50% mCG in wild type) lost at least half

of their CG methylation in epimutagenized lines (**Supplementary Table 5 and 6**).

Collectively, these results indicate hypomethylation was more severe in genes than

transposons, possibly the result of *de novo* methylation by the RdDM pathway which is

primarily active at transposons.

Tet1-mediated DNA demethylation mimics *met1* mutants

An analysis of Differentially Methylated Regions (DMRs) was then carried out to assess

the genome-wide impact of hTET1cd expression. 56,283 CG DMRs ranging in size from

6 - 20,286 base pairs (bp) were identified (**Fig. 1g, Supplementary Table 7**). Of these,

38.7% were located in intergenic sequences, 53.7% overlapped with genes and 7.6%

were located in promoter regions (≤1kb upstream of a gene). As also seen in *met1*

mutants, the predominant effect of hTET1cd expression is CG hypomethylation (12,641

and 20,601 DMRs lost more than 50% mCG in 35S:TET1-1 and 35S:TET1-2,

respectively; no region gained more than 50% mCG). However, the extent of CG

methylation loss caused by hTETcd expression is lower than in *met1*: 31.8 Mb of the

genome significantly lost CG methylation in *met1*, whereas 9.9 Mb and 18.0 Mb were

lost in 35S:TET1-1 and 35S:TET1-2, respectively.

Previous studies of the *met1* methylome have revealed a loss of mCHG/mCHH

methylation in a subset of CG-hypomethylated regions [17]. At these loci, DNA methylation

is stably lost, in contrast to regions where DNA methylation is re-established by *de novo*

methylation pathways. These loci are ideal targets of epimutagenesis, as the co-

existence of all three types of methylation is more frequently correlated with

transcriptional repression of genes than CG methylation alone. This, coupled with the

long-term stability of hypomethylation, may facilitate inherited transcriptional changes.

An analysis of the interdependence of the loss of CG methylation on non-CG

methylation levels revealed that 39.7 Kb and 931.5 Kb of CHG methylated sequences

lost significant amounts of methylation in two independent epimutagenized lines,

compared to 4.0 Mb of sequence in *met1* mutants. A similar analysis for the loss of CHH

methylation revealed losses of 23.3 Kb and 492.5 Kb in epimutagenized individuals,

compared to 1.1 Mb lost in *met1* mutants. Of the 56,283 identified CG DMRs, 10,491

overlapped regions that contained at least 20% CHG methylation and 7,214 overlapped

regions that contained at least 10% CHH methylation in wild-type individuals

(Supplementary Table 7). To determine how many of these regions are susceptible to

losing non-CG methylation if CG methylation is first depleted, we created a frequency

distribution of mCHG and mCHH levels in wild-type and epimutagenized individuals (**Fig.**

1h and i). 2,341 and 3,447 regions lost more than 10% CHG methylation in 35S:TET1-1

and 35S:TET1-2, respectively, whereas 2,475 and 3,379 regions lost more than 5%

CHH methylation in 35S:TET1-1 and 35S:TET1-2, respectively. Regions that are

susceptible to losses of CG and non-CG methylation in lines expressing hTET1cd share

a substantial overlap with regions that lose non-CG methylation in *met1*

(**Supplementary Fig. 1g and h)**. 1,708 (73.0%) and 2,386 (69.2%) regions that have

lost more than 10% mCHG in 35S:TET1-1 and 35S:TET1-2 have reduced levels in

met1, whereas 2,013 (81.6%) and 2,563 (75.9%) regions that have lost more than 5%

mCHH in 35S:TET1-1 and 35S:TET1-2 have reduced levels in *met1*. As crop genomes

have a greater number of loci targeted for silencing by CG, CHG and CHH methylation

when compared to *A. thaliana*, ectopic expression of hTET1cd is likely a viable approach

for the creation epiRILs[18].

Tet1-mediated variation of CHG methylation

Mutations in *met1* also result in hypermethylation of CHG sites in gene bodies due to the

loss of methylation in the 7th intron of the histone 3 lysine 9 (H3K9) demethylase,

INCREASE IN BONSAI METHYLATION 1 (*IBM1*)[19, 20, 21, 22]. This results in alternative

splicing of *IBM1,* ultimately producing a non-functional gene product (IBM1-S), which

results in ectopic accumulation of di-methylation of H3K9 (H3K9me2) throughout the

genome[23]. As in *met1*, the 7th intron of *IBM1* was hypomethylated in 35S:TET1-1,

35S:TET1-2, and an additional two lines, 35S:TET1-2^{T5} and 35S:TET1-2^{T6}, which were

propagated for an additional two and three generations, respectively (**Fig. 2a**). The

increased abundance of *IBM1-S* and the reduction of functional *IBM1* (*IBM1-L)* transcript

was confirmed by RT-qPCR in line 35S:TET1-2^{T6} (**Supplementary Fig. 2a**) leading to

CHG hypermethylation at gbM loci (**Fig. 2b**). Further quantitative analysis revealed

extensive variation in genome-wide gains and losses of CHG methylation in these two

lines, approximately 1.8 Mb and 2.3 Mb of additional CHG methylation, respectively (**Fig.**

2c, Supplementary Fig. 2b and c). To test the impact of a reduction in functional *IBM1*

on H3K9me2, we performed chromatin immunoprecipitation (ChIP) against H3K9me2 in

35S:TET1-2[T6], which revealed a subtle increase in H3K9me2 in gbM loci that possessed

CHG hypermethylation (**Fig. 3d**).

To further characterize regions of differential CHG methylation, identified CHG

DMRs in line 35S:TET1-2[T5] were categorized into discrete groups based on their DNA

methylation status in wild-type individuals. Of the 9,917 CHG DMRs identified, 1,460

were in loci that are defined as gbM in wild-type individuals, 584 were in unmethylated

regions, and 6,940 of them were in RdDM-like regions (**Fig. 2e-g**). Interestingly, in line

35S:TET1-2[T5], 1,408 (96.4%) of the CHG DMRs in gbM-like loci gained CHG

hypermethylation, whereas 2,156 (31.1%) of the CHG DMRs in RdDM-like regions lost

CHG, in contrast to 595 (8.6%) RdDM-like regions that gained CHG methylation. Lastly,

there were 502 (86.0%) loci that are unmethylated in wild-type individuals that gain CHG

methylation as well as CG and CHH methylation in the epimutagenized lines (**Fig. 2e-g**).

These results reveal that methods for epimutagenesis can result in both losses and

gains in DNA methylation genome-wide.

To characterize the effect of hTET1cd-induced methylome changes on gene

expression, we performed RNA-sequencing (RNA-seq) on leaf tissue of wild-type,

35S:TET1-1 and 35S:TET1-2. Compared to wild-type plants, 629 and 736 up-regulated

genes were identified in 35S:TET1-1 and 35S:TET1-2, respectively, with 176 and 260

genes overlapping with identified CG DMRs. 1,277 and 1,428 down-regulated genes

were identified and 268 and 324 of them overlapped with CG DMRs. There was a high

level of overlap in transcriptome changes seen in 35S:TET1-1 and 35S:TET1-2

compared to *met1* and *ibm1* (**Fig. 2h and i**). Of the genes up-regulated in *met1*, 36.8%

and 38.7% overlapped with up-regulated genes in 35S:TET1-1 and 35S:TET1-2,

respectively (**Supplementary Fig. 2d**). An even greater overlap was observed with

down-regulated genes in *met1*, as 60.1% and 65.2% overlapped with down-regulated

genes in 35S:TET1-1 and 35S:TET1-2, respectively (**Supplementary Fig. 2e**). These

results reveal that hTET1cd expression in *A. thaliana* is a viable approach for accessing

hidden sources of allelic variation by inducing expression variation.

Tet1 expression leads to a delay in the floral transition

In the transgenic plants that were used for WGBS, we observed a delay in the

developmental transition from vegetative to reproductive growth (**Fig. 3a and b**). We

hypothesized that the observed late-flowering phenotype was associated with the

demethylation of the *FWA* (*FLOWERING WAGENINGEN*) locus, as is observed in *met1*

mutants[24, 25]. A closer inspection of the DNA methylation status of this locus revealed

that DNA methylation was completely abolished, as was methylation at adjacent CHG

and CHH sites (**Fig. 3c**). As in *met1*, the loss of methylation at the *FWA* locus was

associated with an increase in *FWA* expression (**Fig. 3d**), which is known to cause a

delay in flowering by restricting the movement of the florigen signal, *FT*, to the shoot

apex[26]. These results demonstrate that expression of hTET1cd leads to phenotypic

variation by abolishing methylation at some regions in all sequence contexts (CG, CHG

and CHH sites).

Tet1-mediated demethylation can be stably inherited over generations

To assess the stability and inheritance of Tet1-mediated demethylation, T1 individuals

expressing 35S:TET1 were self-fertilized, allowing for the loss of the hTET1cd transgene

due to allelic segregation. Unexpectedly, transgene-free 35S:TET1-1 T2 individuals

(35S:TET1-1.3[-TET1], 35S:TET1-1.4[-TET1] and 35S:TET1-1.5[-TET1]) exhibited a reversion to a

normal flowering phenotype, and genome-wide methylation levels closely resembled that

of wild-type individuals (**Fig. 4a**). WGBS on T2 individuals retaining the transgene

revealed similar levels of CG methylation as the T1 parent (35S:TET1-1.1[+TET1] and

35S:TET1-1.2[+TET1]). The methylation level of CGs at genes and transposons revealed

that demethylation of gbM loci was partially inherited in transgene-free T2 individuals,

whereas active remethylation was found at transposons in the same individuals (**Fig. 4b-**

f). These results indicate that an active process likely in the meristem and/or germline is

counteracting the activity of the hTET1cd transgene[27]. To quantify how many regions

were susceptible to the loss of non-CG methylation, a DMR analysis was conducted for

each 35S:TET1-1 T2 individual. 655 and 659 CHG hypomethylated regions were

identified in T2 lines retaining the transgene. In contrast, 155, 199 and 201

hypomethylated regions were identified in three transgene-free T2 individuals,

respectively (**Fig. 4g**).

244   To determine if increased expression of hTET1cd would increase the likelihood

of germline transmittance of demethylation to transgene-free progeny, transgenic *A.*

thaliana plants were generated expressing a previously described superfolder GFP

(sfGFP) hTET1cd fusion, under control of the *A. thaliana* ACTIN 2 (ACT2) promoter

(ACT2:sfGFP-hTET1cd) which is known to have activity in all tissues of juvenile plants,

including meristematic tissue[28]. Translation and nuclear localization of the sfGFP-

TET1cd fusion protein was confirmed in young cotyledons using confocal microscopy

(**Supplementary Fig 3a**). T1 populations transformed with ACT2:sfGFP-hTET1cd

exhibited a 27-fold increase in later flowering compared to 35S:hTET1cd, indicating high

activity of the sfGFP-TET1cd fusion protein in *A. thaliana* (**Fig. 5a**). To assess the

variation between lines, we performed WGBS on four independent ACT2:TET1 T1 lines

(**Supplementary Fig. 4a-c**). Differential levels of global CG demethylation were

observed in these four lines, further confirming that plants subjected to epimutagenesis

can possess different degrees of demethylation. Subsequent DMR analysis revealed

68,260 CG DMRs, 9,235 CHG DMRs and 2,793 CHH DMRs between these lines, a

drastic increase in DMRs compared to those within siblings (**Fig. 4g**).

To further assess the inheritance of the demethylation pattern as a result of

epimutagenesis, we selected T2 progeny of a late flowering ACT2:TET1-1 T1 individual

containing and lacking the transgene. Individuals retaining and lacking the transgene

due to allelic segregation both exhibited a late-flowering phenotype, ectopic expression

and loss of DNA methylation of *FWA* (**Supplementary Fig 3b and c**). WGBS on these

two individuals revealed a reduction in CG methylation that was maintained and stably

inherited irrespective of transgene presence, confirming the high activity of

demethylation in ACT2:TET1 lines compared to 35S:TET1 lines (**Fig. 5b-f**). It is unclear

exactly why the late-flowering phenotype was stably inherited in the T2 individuals

without the transgene in the ACT2 driven lines versus the 35S driven lines, although it is

likely a combination of promoter strength and cell type specificity. For stable inheritance

of demethylation and the late-flowering phenotype TET1 activity would be required in the

meristematic and/or germline cells. Collectively, these results demonstrate the stable

inheritance of TET1-mediated demethylation and a delayed floral transition in the

absence of the transgene.

DISCUSSION

The discovery that expression of the catalytic domain of the human TET1 protein in *A.*

thaliana leads to widespread loss of CG methylation and enables the creation

epimutants without the need for methyltransferase mutants, which often causes lethality

in crops. In addition to epimutagenesis, hTET1cd could be used in combination with

sequence-specific DNA binding proteins such as dCas9 to direct DNA demethylation in

plant genomes, as has been demonstrated in mammalian systems[29, 30, 31, 32, 33]. The

stable meiotic inheritance of DNA methylation states in flowering plant genomes

provides a stark contrast to the inheritance of DNA methylation in mammalian genomes,

where genome-wide erasure of DNA methylation and reprogramming occurs each

generation[34]. This property of flowering plant genomes makes them ideal targets of

induced-epialleles, as once a new methylation state occurs it is often inherited in

subsequent generations. Application of epimutagenesis and the use of TET-mediated

engineering of DNA methylation states in economically and agriculturally significant plant

species will be an interesting area of future investigation.

ACKNOWLEDGEMENTS

We thank Nathan Springer for comments and discussions on this study as well as the

Georgia Genomics Facility and the Georgia Advanced Computing Resource Center for

technical support. This work was supported by the National Science Foundation (MCB-

1650331), by The Pew Charitable Trusts and by the Office of the Vice President of

Research at UGA to R.J.S. C.H was supported by the National Institutes of Health (NIH

HG006827) and is an investigator of the Howard Hughes Medical Institute. W.T.J. was

supported by a Scholars of Excellence Fellowship from the University of Georgia and

National Institute of General Medical Sciences of the National Institutes of Health award

number T32GM007103. The content is solely the responsibility of the authors and does

not necessarily represent the official view of the National Institutes of Health.

AUTHOR CONTRIBUTIONS

R.J.S. conceived the project. L.J., W.T.J. and R.J.S. designed experiments. L.J., W.T.J.,

X.S., L.H., and C.H. performed research. L.J., W.T.J. and R.J.S. analyzed the data.

R.J.S. wrote the paper. All authors discussed the results and commented on the

manuscript.

COMPETING FINANCIAL INTERESTS

A provisional patent is pending for the development of this technology as it applies to

epigenome engineering of plants.

Figure 1. Overexpression of hTET1cd induced global CG demethylation in *A.*

***thaliana*.** (a) Bar graph of global methylation levels in two Col-0 WT replicates, two

35S:TET1 transgenic individuals and *met1*. (b) Metaplot of CG methylation levels (100

kb windows) across five *A. thaliana* chromosomes. Methylation level differences were

defined relative to Col-0 WT-1, and Col-0 WT-2 was used to assess background

interference. Genome browser view of methylome profile of two regions (c and d) of the

A. thaliana genome (purple vertical lines = CG methylation, blue vertical lines = CHG

methylation and gold vertical lines = CHH methylation). Metagene plots of CG

methylation level across (e) gene bodies and (f) transposable elements. (g) Heat map of

CG methylation level of CG DMRs. Bar plots of CHG (h) and CHH (i) methylation levels

of CG DMRs that possess non-CG methylation in wild-type individuals.

Figure 2. Global fluctuation of CHG methylation in 35S:TET1 plants. (a) Genome

browser view of *IBM1* (AT3G07610) in Col-0 WT, three 35S:TET1 transgenic plants and

met1. A decrease in CG methylation from coding regions was accompanied by an

increase in non-CG methylation. Both CG and non-CG methylation was lost from the

large intron (purple vertical lines = CG methylation, blue vertical lines = CHG methylation

and gold vertical lines = CHH methylation). (b) Genome browser view of a representative

CHG hypermethylated region. (c) The amount of the genome affected by differential

CHG methylation. These DMRs were defined relative to Col-0 WT-1, as Col-0 WT-2

DMRs were used to assess background interference. (d) Boxplot of H3K9me2

distribution in gbM loci (**, t-test, p-value < 0.01). (e) Heat map of CHG methylation

displaying CHG DMRs. Corresponding CG and CHH methylation levels are shown in (f)

and (g). Heat maps showing log$_2$ transformed FPKM profiles of up-regulated genes (h)

and down-regulated genes (i) in two 35S:TET1 transgenic individuals, *met1* and *ibm1*

mutants relative to WT.

Figure 3. 35S:TET1 plants have a delayed flowering phenotype. (a) Photographs of

one 35S:TET1-1 transgenic plant and Col-0 WT plant and (b) corresponding number of

rosette leaves (**, t-test, p-value < 0.01). (c) Genome browser view of *FWA*

(AT4G25530). Both CG and non-CG DNA methylation are depleted from the 5'UTR in

35S:TET1-2 plants (purple vertical lines = CG methylation, blue vertical lines = CHG

methylation and gold vertical lines = CHH methylation). (d) Expression level (FPKM) of

FWA.

349
350
Figure 4. Transgenerational demethylation profile of 35S:TET1 individuals. (a) Bar

plots of global methylation levels in two Col-0 WT replicates and 35S:TET1-1 plants.

Metagene plots of mCG level across (b) gene bodies and (c) transposable elements.

Heat map of mCG level of (d) all gbM genes and (e) transposable elements with >20%

mCG in wild type. (f) Genome browser view of a methylome profile of a representative

region of the *A. thaliana* genome in 35S:TET1-1 individuals. (g) Number of identified

CHG DMRs in ACT2:TET1 T2 plants. These DMRs were defined relative to Col-0 WT-1,

as Col-0 WT-2 DMRs were used to assess background.

360
Figure 5. Demethylation profile of ACT2:TET1 individuals. (a) Bar plot of number of

rosette leaves in 35S:TET1 and ACT2:TET1 T1 individuals upon flowering. (b) Bar plot

of global methylation levels in two Col-0 WT replicates and two ACT2:TET1-1 T2 plants.

Metagene plots of mCG level across (c) gene bodies and (d) transposable elements.

Heat map of mCG level of (e) all gbM genes and (f) transposable elements with >20%

mCG in wild type.

367
METHODS

Synthesis and cloning of the human TET1 catalytic domain. A human TET1 catalytic

domain (hTET1cd) sequence (residues 1418-2136) was synthesized by GenScript, and

moved to a plant transformation compatible vector (pMDC32) using LR clonase from Life

Technologies per the manufacturer's instructions (Catalog #11791100). ACT2:sfGFP-

hTET1cd was subcloned by Genscript in the pMDC32 vector background using the

sfGFP-TET1cd fragment from Addgene plasmid #82561. The ACT2 promoter sequence

was kindly provided by Dr. Richard Meagher.

Plant transformation and screening. The hTET1cd sequence in the pMDC32 vector

was transformed into *Agrobacterium tumefaciens* strain C58C1 and plated on LB-agar

supplemented with kanamycin (50 µg/mL), gentamicin (25 µg/mL), and rifampicin (50

µg/mL). A single kanamycin resistant colony was selected and used to start a 250-mL

culture in LB Broth Miller liquid media supplemented with gentamicin (25 µg/mL),

kanamycin (50 µg/mL), and rifampicin (50 µg/mL), which was incubated for two days at

30°C. Bacterial cells were pelleted by centrifugation at 4,000 RPM for 30 minutes and

the supernatant decanted. The remaining bacterial pellet was re-suspended in 200 mL of

5% sucrose with 0.05% Silwet L77. Plant transformation was performed using the floral

dip method described by Clough and Bent[35]. Seeds were harvested upon senescence

and transgenic plants were identified via selection on ½ LS plates supplemented with

Hygromycin B (25 µg/mL). 35S:TET1-1 is a T1 individual, 35S:TET1-2 is a T3 plant,

35S:TET1-3 is a T4 plant. All transgenic individuals chosen for analysis contain

independent insertions of hTET1cd and are not the result of single-seed decent unless

otherwise noted.

DNA and RNA isolation. *A. thaliana* leaf tissue was flash-frozen and finely ground to a

powder using a mortar and pestle. DNA extraction was carried out on all samples using

the DNeasy Plant Mini Kit (QIAGEN), and the DNA was sheered to approximately 200

bp by sonication. RNA was isolated from finely ground flash-frozen leaf tissue using

Trizol (Thermo Scientific). For RT-qPCR, RNA was further treated with TURBO™

DNase (Thermo Scientific) according to manufacturers instructions. One µg of RNA was

subsequently reverse transcribed with M-MuLV Reverse Transcriptase according to

manufacturers instructions (NEB). RT-qPCR was used to analyzed cDNA populations

using PP2AA-3 (AT1G13320) as an endogenous control, and was performed on a

Roche LIghtCycler 480 instrument using SYBR Green detection chemistry. The genes

assayed by this method were *IBM1-S*, *IBM1-L,* and *FWA*. Primers used for RT-qPCR

were designed using PrimerQuest from Integrated DNA Technologies

(www.idtdna.com/PrimerQuest/). Primers sequences used for RT-qPCR: PP2A-F: 5'

AATGAGGCAGAAGTTCGGATAG 3', PP2A-R: 5'CAGGGAAGAATGTGCTGGATAG 3',

ibm1s-F: 5' TCTTTCTTCTAAGTCTGTCCATTCT 3', ibm1s-R: 5'

GTGACCGATTAGGAAATGGTATCT 3', ibm1L-F: 5' CCGAAGCCAAAGTGGAGATA 3',

ibm1L-R: 5' CTTCCTCTTCCGTAGACTTCTTT 3', FWA-F: 5'

CAAGATGGTGGAAGGATGAGAA 3', FWA-R: 5' CTCTGTTCTTCAGTGGGATGAG 3'.

Library construction. Genomic DNA libraries were prepared following the MethylC-seq

protocol without use of the bisulfite conversion step. MethylC-seq libraries were

prepared as previously described in[36]. RNA-seq libraries were constructed using Illumina

TruSeq Stranded RNA LT Kit (Illumina, San Diego, CA) following the manufacturer's

instructions with limited modifications. The starting quantity of total RNA was adjusted to

1.3 µg, and all volumes were reduced to a third of the described quantity. TAB-seq

libraries were prepared as previously described in[16].

ChIP library preparation. Leaves were treated with formaldehyde to covalently link

protein to DNA, washed several times with distilled water, patted dry and ground into fine

powder in liquid nitrogen. Chromatin was extracted with a series of extraction buffers

and sonicated. The final chromatin solution was incubated overnight with anti-H3K9me2

antibody (Cell Signaling Technology, 9753S) coated Dynabeads protein A (Life

Technologies, 10002D) to precipitate the immune complex. After a few washes, the

immune complex was eluted and incubated at 65 degrees in the presence of a high

concentration of NaCl in a water-bath overnight. After degrading the proteins with

proteinase K, DNA was recovered by phenol/chloroform/isoamyl alcohol extraction

followed by ethanol precipitation. The DNA pellet was then dissolved in 30µl of nuclease-

free water.

Sequencing. Illumina sequencing was performed at the University of Georgia Genomics

Facility using an Illumina NextSeq 500 instrument. For MethylC-seq and TAB-seq, raw

reads were trimmed for adapters and preprocessed to remove low quality reads using

cutadapt 1.9.dev1[37]. For RNA-seq and ChIP-seq, these processes were carried out by

Trimmomatic v0.32[38]. The mutant allele of *met1* is *met1-3* and methylome data was

downloaded under accession GSE39901. The mutant allele of *ibm1* is *ibm1-6*.

MethylC-seq data processing. Qualified reads were aligned to the *A. thaliana* TAIR10

reference genome as described in[39]. Chloroplast DNA (which is fully unmethylated) was

used as a control to calculate the sodium bisulfite reaction non-conversion rate of

unmodified cytosines. All conversion rates were >99% (**Supplementary Table 1**). The

list of gbM genes used in this study was previously curated[18]. Heat maps were clustered

by Complete Linkage method conducted by R (https://www.r-project.org). All methylation

levels reported in all analyses are presented as differences in absolute values, including

defining DMRs and calculating hyper/hypomethylated regions. The only exception is in

the comparison of mCG loss between gbM, where we used a percentage difference.

RNA-seq data processing. Qualified reads were aligned to the *A. thaliana* TAIR10

reference genome using TopHat v2.0.13[40] (**Supplementary Table 2**). Gene expression

values were computed using Cufflinks v2.2.1[41]. Genes determined to have at least 2-fold

$\log_2$ expression changes by Cufflinks and passed tests were identified as differentially

expressed genes.

TAB-seq data processing. Qualified reads were aligned to the *A. thaliana* TAIR10

reference genome using Methylpy as described in[39]. A modified lambda DNA sequence

was used to assess the quality of prepared libraries. In the added lambda sequence, all

non-CG cytosines are unmethylated and CG cytosines are methylated to 5mC. The

'non-conversion' rate is used to measure the rate of non-CG cytosines failing to be

converted to thymines after bisulfite treatment. The '5mC non-conversion' is used to

estimate the 5mCG cytosines failing to be converted to thymines after TET treatment.

(**Supplementary Table 3**).

ChIP-Seq Data Analysis. Qualified reads were aligned to the *A. thaliana* TAIR10

reference genome using Bowtie 1.1.1 with following parameters: bowtie -m 1 -v 2 --best -

-strata --chunkmbs 1024 -S. Aligned reads were sorted using samtools v 1.2 and clonal

duplicates were removed using SAMtools version 0.1.19 (**Supplementary Table 4**).

Metaplot analysis. For metaplot analyses, twenty 50 bp bins were created for both

upstream and downstream regions of gene bodies/TEs. Gene bodies/TE regions were

evenly divided into 20 bins. Weighted methylation levels were computed for each bin as

described previously[42].

DMR analysis. Identification of DMRs was performed as described in[43] and adjusted p-

value (Benjamini-Hochberg correction) 0.05 was adopted as the cutoff. Only DMRs with

at least 5 DMSs (Differential Methylated Sites) and a 10% absolute methylation level

difference within each DMR were reported and used for subsequent analysis. For

coverage calculations, each sample was combined with two Col-0 WT replicates to

identify DMRs. Each sample was compared with both Col-0 WT replicates separately

and for a DMR to be identified it must have been identified in both comparisons.

Absolute methylation differences of +/- (50% for CG, 10% for CHG and CHH) were

defined as hyper/hypo methylation, respectively. DMRs overlapping regions with mCG

>= 5%, mCHG and mCHH >= 1% in both two Col-0 WT replicates were defined as

RdDM-like regions. DMRs overlapping regions with mCG >= 5%, mCHG and mCHH <

1% in both two Col-0 WT replicates were defined as gbM regions. DMRs overlapping

regions with all three contexts less methylated at less than 1% in both Col-0 WT

replicates were defined as unmethylated regions. Overlap comparisons were performed

using bedtools v2.26.0[44].

Data availability: The data generated from this study has been uploaded to Gene

Expression Omnibus (GEO) database and can be retrieved through accession number

GSE93024.

References

1. Law JA, Jacobsen SE. Establishing, maintaining and modifying DNA methylation patterns in plants and animals. *Nature reviews Genetics* **11**, 204-220 (2010).

2. Johannes F, *et al.* Assessing the impact of transgenerational epigenetic variation on complex traits. *PLoS Genet* **5**, e1000530 (2009).

3. Reinders J, *et al.* Compromised stability of DNA methylation and transposon immobilization in mosaic Arabidopsis epigenomes. *Genes Dev* **23**, 939-950 (2009).

4. Cortijo S, *et al.* Mapping the Epigenetic Basis of Complex Traits. *Science*, (2014).

5. Bewick AJ, *et al.* On the origin and evolutionary consequences of gene body DNA methylation. *Proc Natl Acad Sci U S A* **113**, 9111-9116 (2016).

6. Cortijo S, Wardenaar R, Colome-Tatche M, Johannes F, Colot V. Genome-Wide Analysis of DNA Methylation in Arabidopsis Using MeDIP-Chip. *Methods Mol Biol* **1112**, 125-149 (2014).

7. Reinders J, Delucinge Vivier C, Theiler G, Chollet D, Descombes P, Paszkowski J. Genome-wide, high-resolution DNA methylation profiling using bisulfite-mediated cytosine conversion. *Genome Res* **18**, 469-476 (2008).

8. Hofmeister BT, Lee K, Rohr NA, Hall DW, Schmitz RJ. Stable inheritance of DNA methylation allows creation of epigenotype maps and the study of epiallele inheritance patterns in the absence of genetic variation. *Genome biology* **18**, 155 (2017).

9. Li Q, *et al.* Genetic Perturbation of the Maize Methylome. *Plant Cell*, (2014).

10. Hu L, *et al.* Mutation of a major CG methylase in rice causes genome-wide hypomethylation, dysregulated genome expression, and seedling lethality. *Proc Natl Acad Sci U S A* **111**, 10642-10647 (2014).

11. Tahiliani M, *et al.* Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* **324**, 930-935 (2009).

12. Pastor WA, *et al.* Genome-wide mapping of 5-hydroxymethylcytosine in embryonic stem cells. *Nature* **473**, 394-397 (2011).

538   13.   Ito S, D'Alessio AC, Taranova OV, Hong K, Sowers LC, Zhang Y. Role of Tet
539   proteins in 5mC to 5hmC conversion, ES-cell self-renewal and inner cell mass
540   specification. *Nature* **466**, 1129-1133 (2010).
541

542   14.   Hollwey E, Watson M, Meyer P. Expression of the C-Terminal Domain of
543   Mammalian TET3 DNA Dioxygenase in Arabidopsis thaliana Induces
544   Heritable Methylation Changes at rDNA Loci. *Advances in Bioscience and*
545   *Biotechnology* **7**, 243-250 (2016).
546

547   15.   Erdmann RM, Souza AL, Clish CB, Gehring M. 5-hydroxymethylcytosine is not
548   present in appreciable quantities in Arabidopsis DNA. *G3 (Bethesda)* **5**, 1-8
549   (2014).
550

551   16.   Yu M, *et al.* Tet-assisted bisulfite sequencing of 5-hydroxymethylcytosine.
552   *Nature protocols* **7**, 2159-2170 (2012).
553

554   17.   Stroud H, Greenberg MV, Feng S, Bernatavichute YV, Jacobsen SE.
555   Comprehensive analysis of silencing mutants reveals complex regulation of
556   the Arabidopsis methylome. *Cell* **152**, 352-364 (2013).
557

558   18.   Niederhuth CE*, et al.* Widespread natural variation of DNA methylation
559   within angiosperms. *Genome biology* **17**, 194 (2016).
560

561   19.   Saze H, Shiraishi A, Miura A, Kakutani T. Control of genic DNA methylation by
562   a jmjC domain-containing protein in Arabidopsis thaliana. *Science* **319**, 462-
563   465 (2008).
564

565   20.   Miura A, Nakamura M, Inagaki S, Kobayashi A, Saze H, Kakutani T. An
566   Arabidopsis jmjC domain protein protects transcribed genes from DNA
567   methylation at CHG sites. *EMBO J* **28**, 1078-1086 (2009).
568

569   21.   Rigal M, Kevei Z, Pelissier T, Mathieu O. DNA methylation in an intron of the
570   IBM1 histone demethylase gene stabilizes chromatin modification patterns.
571   *EMBO J* **31**, 2981-2993 (2012).
572

573   22.   Lister R*, et al.* Highly integrated single-base resolution maps of the
574   epigenome in Arabidopsis. *Cell* **133**, 523-536 (2008).
575

576   23.   Rigal M, Kevei Z, Pélissier T, Mathieu O. DNA methylation in an intron of the
577   IBM1 histone demethylase gene stabilizes chromatin modification patterns.
578   *The EMBO journal* **31**, 2981-2993 (2012).
579

580   24.   Soppe WJ*, et al.* The late flowering phenotype of fwa mutants is caused by
581   gain-of-function epigenetic alleles of a homeodomain gene. *Mol Cell* **6**, 791-
582   802 (2000).
583

584   25.   Finnegan EJ, Peacock WJ, Dennis ES. Reduced DNA methylation in
585   Arabidopsis thaliana results in abnormal plant development. *Proceedings of*
586   *the National Academy of Sciences of the United States of America* **93**, 8449-
587   8454 (1996).
588
589   26.   Ikeda Y, Kobayashi Y, Yamaguchi A, Abe M, Araki T. Molecular basis of late-
590   flowering phenotype caused by dominant epi-alleles of the FWA locus in
591   Arabidopsis. *Plant & cell physiology* **48**, 205-220 (2007).
592
593   27.   Baubec T, Finke A, Scheid OM, Pecinka A. Meristem‐specific expression of
594   epigenetic regulators safeguards transposon silencing in Arabidopsis. *EMBO*
595   *reports* **15**, 446-452 (2014).
596
597   28.   An YQ, McDowell JM, Huang S, McKinney EC, Chambliss S, Meagher RB.
598   Strong, constitutive expression of the Arabidopsis ACT2/ACT8 actin subclass
599   in vegetative tissues. *The Plant Journal* **10**, 107-121 (1996).
600
601   29.   Choudhury SR, Cui Y, Lubecka K, Stefanska B, Irudayaraj J. CRISPR-dCas9
602   mediated TET1 targeting for selective DNA demethylation at BRCA1
603   promoter. *Oncotarget*, (2016).
604
605   30.   Maeder ML*, et al.* Targeted DNA demethylation and activation of endogenous
606   genes using programmable TALE-TET1 fusion proteins. *Nat Biotechnol* **31**,
607   1137-1142 (2013).
608
609   31.   Vojta A*, et al.* Repurposing the CRISPR-Cas9 system for targeted DNA
610   methylation. *Nucleic Acids Res*, (2016).
611
612   32.   Mendenhall EM*, et al.* Locus-specific editing of histone modifications at
613   endogenous enhancers. *Nat Biotechnol* **31**, 1133-1136 (2013).
614
615   33.   Liu XS*, et al.* Editing DNA Methylation in the Mammalian Genome. *Cell* **167**,
616   233-247 e217 (2016).
617
618   34.   Heard E, Martienssen RA. Transgenerational epigenetic inheritance: myths
619   and mechanisms. *Cell* **157**, 95-109 (2014).
620
621   35.   Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-
622   mediated transformation of Arabidopsis thaliana. *The Plant journal : for cell*
623   *and molecular biology* **16**, 735-743 (1998).
624
625   36.   Urich MA, Nery JR, Lister R, Schmitz RJ, Ecker JR. MethylC-seq library
626   preparation for base-resolution whole-genome bisulfite sequencing. *Nature*
627   *protocols* **10**, 475-483 (2015).
628

629　37.　Martin M. Cutadapt removes adapter sequences from high-throughput
630　　　　sequencing reads. *EMBnet journal* **17**, pp. 10-12 (2011).
631
632　38.　Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
633　　　　sequence data. *Bioinformatics* **30**, 2114-2120 (2014).
634
635　39.　Schmitz RJ*, et al.* Epigenome-wide inheritance of cytosine methylation
636　　　　variants in a recombinant inbred population. *Genome research* **23**, 1663-
637　　　　1674 (2013).
638
639　40.　Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2:
640　　　　accurate alignment of transcriptomes in the presence of insertions, deletions
641　　　　and gene fusions. *Genome biology* **14**, R36 (2013).
642
643　41.　Trapnell C*, et al.* Transcript assembly and quantification by RNA-Seq reveals
644　　　　unannotated transcripts and isoform switching during cell differentiation.
645　　　　*Nature biotechnology* **28**, 511-515 (2010).
646
647　42.　Schultz MD, Schmitz RJ, Ecker JR. 'Leveling' the playing field for analyses of
648　　　　single-base resolution DNA methylomes. *Trends in genetics : TIG* **28**, 583-585
649　　　　(2012).
650
651　43.　Schultz MD*, et al.* Human body epigenome maps reveal noncanonical DNA
652　　　　methylation variation. *Nature* **523**, 212-216 (2015).
653
654　44.　Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing
655　　　　genomic features. *Bioinformatics* **26**, 841-842 (2010).
656
657
658

a *IBM1* (AT3G07610)

b AT4G31570

Col-0 WT-1
Col-0 WT-2
35S:TET1-1
35S:TET1-2
35S:TET1-2^{T5}
35S:TET1-2^{T6}
met1

h Relative FPKM level
log2 (Sample/wild type)

−4 −2 0 2 4

Up-regulated

c

CHG DMR coverage (Mb)

- Hypermethylation
- Hypomethylation

Col-0 WT-2   1   2   2^{T5} 2^{T6} *met1*

35S:TET1

d

H3K9me2 RPKM

**

0   1   5   10   15   20   100

mCHG level (%)

i Down-regulated

35S:TET1-1   35S:TET1-2   *met1*   *ibm1*

e CHG   **f** CG   **g** CHH

gbM-like (n=1,460)

unmethylatd (n=584)

RdDM-like (n=6,940)

Col-0 WT-1
Col-0 WT-2   1   2   2^{T5} 2^{T6} *met1*

35S:TET1

methylation level (%)

0   20   40   60   80   100

a

35S:TET1-2   Col-0 WT

b

c

FWA (AT4G25530)

d

a

b

c **Gene**

d **TE**

e

f