Evolutionarily younger and faster diversifying plants are more threatened by extinction

Andrew J. Tanentzap*, Javier Igea, Matthew G. Johnston

Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, UK

CB2 3EA

*Correspondence: ajt65@cam.ac.uk

Abstract

Many species are threatened by extinction, yet few factors have been identified to explain this

risk across the plant Tree of Life (ToL). Lineages that are evolutionarily younger or

occurring within rapidly diversifying clades may have elevated extinction risk because they

occupy smaller geographic ranges and adaptive zones as compared with older lineages that

have tolerated longer periods of environmental change. Here we find that faster diversifying

plant genera had more species at risk of extinction. Evolutionary age had no effect in the 297

genera that we sampled, potentially because of sampling older genera, on average, from

across the ToL. Repeating our analyses in two well-sampled and large groups, we found that

extinction risk decreased with evolutionary age in conifer species but not palms. Small

sample sizes limited our power to detect effects of lineage diversification in these groups.

Our results suggest that contrasting modes of speciation may explain differing patterns of

extinction risk across the broader ToL and have consequences for biodiversity conservation.

Introduction

Much of the world's biodiversity is threatened to varying degrees by extinction [1]. In

addition to present-day environmental pressures, some species may intrinsically face a higher

risk of extinction because of at least two historical reasons. The first is that traits that were

once advantageous became less adaptive as environments diverged from previous selection

regimes [2], thereby making older species more likely to be threatened by extinction. A

classic example is the loss of large-bodied Eurasian mammals, such as woolly rhinoceros

(*Coelodonta antiquitatis*), at the end of the most recent glacial period. Woolly rhinoceros

were presumably maladapted for changes to their steppe habitat arising from warmer

temperatures [3]. However, older species that survived long-term environmental changes

may also be less at risk of extinction because they tolerate a greater range of conditions and

are ecologically more generalist [4,5]. A second historical explanation for variation in

extinction risk is that younger species, which often arise from rapidly diversifying lineages

[6], initially tend to have smaller geographic ranges. Small ranges are especially likely if

speciation started from small reproductively isolated populations. These small populations

can also occupy a narrower adaptive space. Together, small geographic and adaptive ranges

can increase extinction risk in both younger species and rapidly diversifying clades with

many young species [7-9].

Previous work in birds [4] and marsupials [10] found that older lineages were more

threatened by extinction, whilst the reverse was shown across non-lemur primates [8], but

there is no reason for these patterns to hold at a large taxonomic scale across other divisions

in the Tree of Life. The only study on plants, to our knowledge, found a higher extinction

risk in younger, rapidly diversifying clades of the South African Cape [7]. Broader

generalisations across plants have not been possible until now because of poor taxonomic

sampling coverage that prevents reliable divergence times from being estimated.

Here, we combined the largest time-calibrated phylogenetic tree presently estimated

for vascular plants with peer-reviewed assessments of conservation status from the

International Union for Conservation of Nature (IUCN) *Red List* [11] to test whether younger

and faster-evolving clades were associated with greater extinction risk across 297 genera.

We complemented our findings with analyses for two large, ancient, and widespread plant

clades (conifers and palms). These analyses allowed us to address concerns around

estimating divergence times and threat status from the larger but under-sampled phylogenetic

tree.

Methods

Data assembly

We first selected genera for which we could confidently estimate the time of divergence from

their sister clades (i.e. 'stem age') in the time-calibrated, species-level phylogenetic tree of

Qian & Jin [12]. The selected genera came from densely sampled clades (i.e. families) to

circumvent low sampling across the broader tree both at a species- and genus-level. For each

family, we calculated the proportion of genera that were sampled in the phylogeny from the

taxonomic database curated by the *taxonlookup* v1.1.1 *R* package and retained those with

$\geq$60% coverage. We also used stem ages because they only require one species to be

sampled within each genus and reflect the entire evolutionary history of clades unlike crown

ages that can have young age biases because they consider only extant species [13]. Taxa

outside of an established "core clade" for each genus, as determined using *MonoPhy* in R,

were removed prior to all calculations. After calculating ages from the large tree, we

intersected the selected genera with 20,473 IUCN assessments [11] and calculated the

proportion of species in each genus threatened with extinction. We further restricted our

analysis to genera with >1 species, of which ≥20% had sufficient data to be assessed for

extinction risk. Overall, n=297 genera had both reliable age and risk status data spanning

2,679 IUCN species-level assessments.

We also estimated net diversification rates for each of our selected genera. We used a

well-established method-of-moments estimator that assumed diversification rates were

constant over time within genera given a known stem age and species richness [14].

Following standard practice, we assumed three values of relative extinction ε of 0, 0.5 and 0.9

when estimating diversification [14]. Taxonomy was standardised to The Plant List

nomenclature using the *Taxonstand* R package prior to processing all datasets.

We also repeated our genus-level diversification analysis as above with two large

clades that were well sampled at a species-level in separate time-calibrated phylogenies.

These clades included 70% of all 651 accepted Pinales (extant conifers) [15] and all 2,539

Arecaceae (palms) [16]. Using these data, we also intersected risk statuses with species stem

ages, giving n=340 and 346, respectively. For the palms, we used the maximum clade

credibility tree that we computed from the posterior distribution of trees that was generated

using topological constraints based on Govaerts taxonomy recommended in ref. [16].

Statistical analyses

We separately tested whether genera with a greater proportion of threatened taxa had younger

ages and faster diversification rates using phylogenetic least squares regression (PGLS) in the

R *caper* package. Following standard practice, we fitted the PGLS with maximum-likelihood

transformations of branch lengths based on the strength of phylogenetic covariance estimated

by Pagel's λ [17]. Both ages and diversification rates were log-transformed. Models were

not fitted with both predictors simultaneously as they were highly correlated (Spearman's $r <$

-0.79). We repeated this analysis in conifers and palms, and again did not simultaneously fit

age and diversification rates given high correlations ($r < $ -0.71 when $\varepsilon = 0.0$ or 0.5).

For conifers and palms, we also tested whether threatened status was associated with

younger species ages using logistic regression. Models were fitted with penalised maximum-

likelihood that accounted for the phylogenetic non-independence of species using the *phylolm*

R package. Predictors were scaled to a mean of 0 and standard deviation of 1 to compare

effects.

Results

We found that relatively more species were threatened with extinction in faster diversifying

genera (for ε of 0.0, 0.5, 0.9: $t_{295} = 2.25, 2.32, 2.40$ and $p = 0.025, 0.021, 0.017$; respectively).

The mean proportion of a genus threatened with extinction doubled from 32% to 65%

between the slowest and fastest diversifying genera (figure 1). These results did not arise

simply because faster diversifying genera were younger, as genus age had no effect on risk

status ($t_{295} = $ -1.48, $p = 0.141$). However, we found an age bias in our dataset. Sampled

genera were older and more diverse on average than obtained by applying our sampling

criteria to the initial tree (i.e. before intersecting with threat status; table S1), which may

explain why we failed to detect an association between younger genera and extinction risk.

Diversification was also slower in our sample, but the sampling criteria did not lead to

differences in risk status (table S1). Repeating our analyses with only the genera from the

more complete species-level datasets was inconclusive, likely because of small sample sizes

($n \leq 36$; table S2).

Analyses with the more complete species-level datasets also revealed that younger

conifers but not palms were relatively more threatened by extinction ($z_{344} = $ -2.67, $p = 0.008$

and $z_{338} = -1.46$, $p = 0.145$, respectively; figure 2a). The mean effect ± SE was more than

twice as large in the conifers (-0.36 ± 0.14 vs -0.15 ± 0.10 on log-scale), leading to nearly a

50% absolute reduction in the probability of being threatened over the range of observed ages

(figure 2b).

Discussion

Our finding that faster diversifying lineages faced greater extinction risk, which the IUCN

associates with smaller range and/or population sizes [11], may arise because most plant

speciation is allopatric or peripatric [18,19]. Therefore, where these modes occur more often,

species will ultimately have smaller geographic ranges and greater extinction risk [7]. Age

effects on extinction risk should also prevail for allopatric speciation, which requires post-

speciation range expansions that necessarily take time to accumulate [20]. Under peripatry,

time-dependent range expansions are unnecessary for ancestral taxa to continue generating

many new species with small range sizes [20]. These dynamics can also help explain varying

age effects on extinction risk in other studies that span disparate taxonomic groups [4,7-10]

and between conifers and palms. Conifer diversification is presumed to have been more

dependent on range contraction and expansion [15,21] than in palms, where dispersal into

isolated populations, consistent with peripatry, might better explain speciation [22].

Palms and conifers have contrasting biogeographic histories that can explain our

differing results between the groups. Conifers mostly occupy temperate habitats as opposed

to the predominately tropical distribution of palms. The greater ecological stability of the

tropics may therefore result in little influence of evolutionary age on palm extinction [23].

For example, older palm species would have not been exposed to the repeated glacial cycles

that fragmented temperate conifer populations [5] and could have elevated extinction risk.

Another explanation is that palm species that were susceptible to environmental change have

already gone extinct or traits that make species more prone to extinction are not

taxonomically conserved [8]. Our results with conifers and palms were also not simply an

artefact of biased sampling as ages and rates did not markedly differ from observations across

entire clades, i.e. before filtering with IUCN data (table s3). Similar results to the larger

phylogeny also suggest that our conclusions were robust to biases in IUCN assessments [24].

Our results suggest that macro-evolutionary dynamics have some value for

biodiversity conservation. Specifically, we found that these dynamics provided an indicator

of contemporary extinction risk that might be easier to derive for large numbers of taxa than

detailed species-level assessments. Macro-evolutionary dynamics might also offer insight

into the vulnerability of species to future change, as the smaller population and range sizes

that make some species prone to extinction are likely to be carried into the future [25].

Although our results must be interpreted with caution, given several biases inherent to the

size of our datasets, they provide new evidence that lineages span a continuum from little

species turnover to producing fast diversifying and extinction-prone taxa [9]. The consistent

results we found at different taxonomic scales also suggests that similar patterns should be

uncovered in the future when the plant Tree of Life is more densely sampled.

Ethics

Does not apply.

Data accessibility

Data will be deposited in the Dryad repository upon acceptance.

Authors' contributions

AJT conceived the study. All authors designed the study. AJT carried out the data analysis

and drafted the manuscript with input from JI and MGJ. All authors gave final approval for

publication and agree to be held accountable for the work performed therein.

Competing interests

We have no competing interests.

Funding

None declared.

Acknowledgements

We thank Guy Tanentzapf for the question that inspired this study and Greg Jordan for

sharing the conifer phylogeny.

References

1. Barnosky AD, et al. 2011 Has the Earth's sixth mass extinction already arrived? *Nature*

471, 51-57.

2. Wilson EO. 1959 Adaptive shift and dispersal in a tropical ant fauna. *Evolution* **13**, 122-

144.

3. Lorenzen ED, et al. 2011 Species-specific responses of Late Quaternary megafauna to

climate and humans. *Nature* **479**, 359-364.

4. Gaston KJ, Blackburn TM. 1997 Evolutionary age and risk of extinction in the global

avifauna. *Evol Ecol* **11**, 557-565.

5. Liow LH. 2007 Lineages with long durations are old and morphologically average: an

analysis using multiple datasets. *Evolution* **61**, 885-901.

6.   Valente LM, Savolainen V, Vargas P. 2010 Unparalleled rates of species diversification

in Europe. *Proc Roy Soc B* **277**, 1489-96.

7.   Davies TJ, et al. 2011 Extinction risk and diversification are linked in a plant biodiversity

hotspot. *PLoS Biol* **9**, e1000620.

8.   Arregoitia LDV, Blomberg SP, Fisher DO. 2013 Phylogenetic correlates of extinction

risk in mammals: species in older lineages are not at greater risk. *Proc Roy Soc B* **280**,

20131092.

9.   Greenberg DA, Mooers AØ. 2017 Linking speciation to extinction: diversification raises

contemporary extinction risk in amphibians. *Evol Lett* **1**, 40-48.

10.   Johnson CN, Delean S, Balmford A. 2002 Phylogeny and the selectivity of extinction in

Australian marsupials. *Anim Conserv* **5**, 135-142.

11.   IUCN. 2016 *The IUCN Red List of Threatened Species.* (http://www.iucnredlist.org)

12.   Qian H, Jin Y. 2016 An updated megaphylogeny of plants, a tool for generating plant

phylogenies and an analysis of phylogenetic community structure. *J Plant Ecol* **9**, 233-

239.

13.   Scholl JP, Wiens JJ. 2016 Diversification rates and species richness across the Tree of

Life. *Proc. R. Soc. B* **283**, 20161334.

14.   Magallon S, Sanderson MJ. 2001 Absolute diversification rates in angiosperm clades.

Evolution **55**, 1762-1780.

15.   Leslie AB, Beaulieu JM, Rai HS, Crane PR, Donoghue MJ, Mathews S. 2012

Hemisphere-scale differences in conifer evolutionary dynamics. *PNAS* **109**, 16217-

16221.

16.   Faurby S, Eiserhardt WL, Baker WJ, Svenning J-C. 2016 An all-evidence species-level

supertree for the palms (Arecaceae). *Mol Phylogenet Evol* **100**, 57-69.

17. Orme D. 2013 *The caper Package: Comparative Analysis of Phylogenetics and*

Evolution in R. (https://cran.r-project.org/web/packages/caper/vignettes/caper.pdf)

18. Anacker BL, Strauss SY. 2014 The geography and ecology of plant speciation: range

overlap and niche divergence in sister species. *Proc. R. Soc. B* **281**, 20132980.

19. Igea J, Bogarín D, Papadopulos AST, Savolainen V. 2015 A comparative analysis of

island floras challenges taxonomy-based biogeographical models of speciation.

Evolution **69**, 482-491.

20. Pigot AL, Phillimore AB, Owens IPF, Orme CDL. 2010 The shape and temporal

dynamics of phylogenetic trees arising from geographic speciation. *Syst Biol* **59**, 660-

673.

21. Condamine FL, Leslie AB, Antonelli A. 2017 Ancient islands acted as refugia and

pumps for conifer diversity. *Cladistics* **33**, 69-92.

22. Roncal J, et al. 2015 Palm diversification in two geologically contrasting regions of

western Amazonia. *J. Biogeogr.* **42**, 1503-1513.

23. Couvreur TL, Forest F, Baker WJ. 2011 Origin and global diversification patterns of

tropical rain forests: inferences from a complete genus-level phylogeny of palms. *BMC*

Biology **9**, 44.

24. Webb TJ, Mindel BL. 2015 Global patterns of extinction risk in marine and non-marine

systems. *Curr Biol* **24**, 506-511.

25. Condamine FL, Rolland J, Morlon H. 2013 Macroevolutionary perspectives to

environmental change *Ecol Lett* **16**, 72-85.

Figure 1. More species are threatened with extinction in faster diversifying genera. Solid

line is mean association estimated by PGLS for $\varepsilon = 0.50$.

Figure 2. Younger conifers but not palms have a greater probability of being threatened by extinction. (**a**) Boxplot for stem ages of conifer (white, *n*=346) and palm (grey, *n*=340) species that were classified as either threatened or non-threatened. Solid line is median, box is inter-quartile range, whiskers extend 1.5-times the interquartile range, and points are outliers. (**b**) Change in probability of a conifer being classified as threatened with species age. Solid line is mean association estimated by phylogenetic logistic regression.