Range size dynamics can explain why evolutionarily age and diversification rate

correlate with contemporary extinction risk in plants

Andrew J. Tanentzap*[1], Javier Igea[1], Matthew G. Johnston[1], Matthew J. Larcombe[2]

[1]Ecosystems and Global Change Group, Department of Plant Sciences, University of

Cambridge, Downing Street, Cambridge, CB2 3EA, UK

[2]Department of Botany, University of Otago, Dunedin, 9054, New Zealand

*Correspondence: ajt65@cam.ac.uk

Abstract

Extinction threatens many species, yet few factors predict this risk across the plant Tree of

Life (ToL). Taxon age is one factor that may associate with extinction if occupancy of

geographic and adaptive zones varies with time, but evidence for such an association has

been equivocal. Age-dependent occupancy can also influence diversification rates and thus

extinction risk where new taxa have small range and population sizes. Here we analysed 509

well-sampled genera from across the plant ToL. We found that a greater proportion of

species were threatened by extinction in younger and faster-diversifying genera. Repeating

our analyses in two large, well-sampled groups, we found that extinction risk increased with

evolutionary age in conifer species but not palms. Potential range size decreased in older,

non-threatened conifers more strongly than in threatened taxa, suggesting that range size

dynamics may explain differing patterns of extinction risk across the ToL with consequences

for biodiversity conservation.

Keywords: conservation, environmental change, macroecology, macroevolution

phylogenetics   **Article type**: Note   **Number of words in the text**: 3215

Introduction

Much of the world's biodiversity is threatened by extinction because of small geographic ranges and/or population sizes (Pimm et al. 2014). In addition to having traits that promote small ranges and population sizes independent of phylogeny, such as those associated with life history and resource use, some species may be more threatened by extinction because of their evolutionary history (Bennett and Owens 1997; Purvis et al. 2000; Johnson et al. 2002; Arregoitia et al. 2013). Extinction is consequently non-randomly distributed across the Tree of Life (ToL), suggesting that chance events and human activities alone may not be fully responsible for explaining species losses (Bennett and Owens 1997; Purvis et al. 2000; Vamosi and Wilson 2008). Identifying macro-evolutionary predictors of extinction risk can therefore help to assess future conservation status where range and population data are lacking and identify reasons for its non-randomness across the ToL (Jetz and Freckleton 2015).

Taxon age is one measure of the amount of environmental and evolutionary change that species have experienced and may be associated with extinction risk for at least two reasons. The first relates to the idea that older taxa should be less at risk of extinction because they have had more time to disperse across a greater range (Paul et al. 2009; Ceolin and Giehl 2017), consistent with the age-and-area hypothesis (Willis 1926). While differences in the time for dispersal may weaken over long time scales (i.e. millions of years), younger taxa may also face less available space and resources as niches fill through time irrespective of dispersal ability (Tanentzap et al. 2015). As new taxa initially tend to have small range and population sizes, especially if speciation started from small reproductively isolated populations that occupy narrow adaptive spaces (Valente et al. 2010; Castiglione et al. 2017), younger species in rapidly diversifying clades should face a greater risk of extinction (Davies et al. 2011; Greenberg & Mooers 2017).

A second historical explanation for variation in extinction risk relates to differences in

niche breadth among species of different ages. Older species may have survived long-term

environmental changes because they are more generalist (Liow 2007). As broader niches are

positively associated with larger ranges (Slatyer et al. 2013), this explanation would result in

another positive age-and-area association. By contrast, there may be a negative correlation

between age and extinction risk if older species are more specialised and have smaller ranges.

We term this idea the evolutionary specialism hypothesis. Older species can appear more

specialised because traits that were once advantageous became less adaptive as environments

diverged from past selection regimes (Wilson 1959; Žliobaitė et al. 2017). More specialist

species with narrower niches and geographic ranges may only persist over long time periods

in refugia or by having large local population sizes (Williams et al. 2009).

The potential for species to expand their range and reduce extinction risk with time

may ultimately depend on their mode of speciation. Repeated range expansion and

contraction (i.e. "taxon cycles") that isolate peripheral populations consistent with centrifugal

or peripatric speciation can produce small ranges in descendent taxa (Gaston 1998).

Consequently, older species may have a lower extinction risk because they have had more

time to disperse and expand their range, and experience less niche pre-emption from earlier

evolving competitors (Tanentzap et al. 2015). Lineages with high diversification rates under

this mode of speciation can similarly face greater extinction by producing species that have

small ranges (Schwartz and Simberloff 2001). By contrast, any signature of time in

extinction risk distributions may be absent with vicariant speciation because asymmetry in

the ranges of ancestors and daughter species is consistently smaller and ancestral species

often disappear via cladogenesis (Gaston 1998).

Evidence that taxon age is associated with extinction varies among lineages, so testing

correlations in relation to range size can help explain this variation and make generalisations

across different divisions in the ToL. Previous work in birds (Gaston and Blackburn 1997)

and marsupials (Johnson et al. 2002) found that older lineages were more threatened by

extinction, whilst the reverse was shown across non-lemur primates (Arregoitia et al. 2013).

The only study on plants, to our knowledge, found a higher extinction risk in younger, rapidly

diversifying clades of the South African Cape (Davies et al. 2011). Broader generalisations

across plants have not been possible until now because of poor taxonomic sampling that

prevents reliable divergence times from being estimated.

Here, we tested whether younger and faster-evolving lineages were associated with

greater extinction risk across 509 genera representing 7,497 species. We did so by combining

the largest time-calibrated phylogenetic tree presently estimated for vascular plants with all

available peer-reviewed assessments of conservation status from the International Union for

Conservation of Nature (IUCN) *Red List* (2016). We complemented our findings with

analyses for two large, ancient, and widespread plant clades (conifers and palms). These

analyses allowed us to address concerns around estimating divergence times from the larger

but under-sampled phylogenetic tree and threat status from incompletely sampled genera. By

working at the species-level, we could also collate geographic distribution data to test the

age-and-area and specialism hypotheses, and how they might explain differences in age-

extinction correlations between taxonomic groups with contrasting histories. Positive

correlations between taxon age and range size would support the idea that older species have

had more time for dispersal (i.e. age-and-area hypothesis), whereas a negative correlation

would support the idea that older species are more specialist.

Methods

Data assembly

We first selected genera for which we could confidently estimate the time of divergence from

their sister genera, i.e. 'stem age'. Genera were selected from the time-calibrated, species-

level phylogenetic tree of Qian and Jin (2016), which was an updated version of Zanne et al.

(2014). The selected genera came from densely sampled clades (i.e. families) to circumvent

low sampling across the broader tree both at a species- and genus-level. For each family, we

calculated the proportion of genera that were sampled in the phylogeny from the taxonomic

database curated by *taxonlookup* v1.1.1 (Pennell et al. 2016) in R v3.2 and retained those

with ≥60% coverage. We also used stem ages because they only require one species to be

sampled within each genus and reflect the entire evolutionary history of clades unlike crown

ages that can have young age biases because they consider only extant species (Scholl and

Wiens 2016). Taxa outside of an established "core clade" for each genus, as determined

using *MonoPhy* in R (Schwery and O'Meara 2016), were removed prior to all calculations.

After calculating ages from the large tree, we intersected the selected genera with

25,452 IUCN assessments and calculated the proportion of species in each genus threatened

with extinction. Threat status is jointly determined from abundance, recent temporal change

in population size, and various measures of geographic distribution, such as occupancy and

fragmentation (IUCN 2016). Therefore, metrics of range size alone may not entirely predict

extinction risk despite the potential to use these terms interchangeably. We further restricted

our analysis to genera with >1 species, of which ≥20% had sufficient data to be assessed for

extinction risk. Overall, 509 genera had both reliable age and risk status data spanning 7,497

IUCN species-level assessments.

We also estimated net diversification rates for the 509 genera. We used a well-

established method-of-moments estimator that assumed diversification rates were constant

over time within genera given a known stem age and species richness (Magallon and

Sanderson 2001). Following standard practice, we assumed three values of relative

extinction ε of 0.0, 0.5 and 0.9 when estimating diversification (Magallon and Sanderson

2001). All taxonomy was standardised to The Plant List nomenclature using the *Taxonstand*

R package prior (Cayuela et al. 2012).

We also repeated our diversification analysis as above with two large clades that were

well sampled at a species-level in separate time-calibrated phylogenies. These clades

included 70% of all 651 accepted Pinales (extant conifers) (Leslie et al. 2012) and all 2,539

Arecaceae (palms) (Faurby et al. 2016). We intersected risk statuses of the two clades with

species stem ages, giving $n = 433$ and 547, respectively. For the palms, we used the

maximum clade credibility tree that we computed from the posterior distribution of trees that

was generated using topological constraints based on Govaerts taxonomy recommended in

Faurby et al. (2016).

Finally, we assembled range data for our two large clades. Georeferenced records

with no flagged issues were downloaded from the Global Biodiversity Information Facility

(www.gbif.org). Conifer data were supplemented by published records absent from GBIF

(table A1). All duplicate and spatially invalid records (e.g. non-numerical, exceeding global

extent, located in the ocean, urban areas, or country centroids) were removed with the R

package *sampbias*. Using the occurrences, we estimated potential range size with a

mechanistic species distribution model (SDM) that predicted the physiological tolerances of

species for growth from distribution data (Higgins et al. 2012). Absence points for the SDM

were generated using standard approaches (details given in Appendix A). We then summed

the total number of equal-area (Mollweide projected) 0.25 decimal degree grid cells occupied

by each species. We found no evidence that sampling varied systematically with species age

in a way that would bias our subsequent analyses (table B1).

 Statistical analyses

We separately tested whether genera with a greater proportion of threatened taxa were

correlated with younger ages and faster diversification rates using phylogenetic least squares

(PGLS) regression. Although the least squares model assumed normally distributed errors,

and the response variable was a proportion with binomial errors, PGLS is appropriate for

testing the null hypothesis of no statistically significant effect of an independent variable on a

non-Gaussian response (Ives 2015). We also fitted the PGLS regression using the *gls*

function in R because this approach, unlike other model fitting functions that incorporated

phylogenetic information (e.g. *phyloglm*), could account for different sample sizes across

genera by weighting observations with the inverse square-root of the number of IUCN

assessments that they received (Garamszegi and Møller 2010). Following standard practice,

the PGLS was fitted with maximum-likelihood transformations of branch lengths based on

the strength of phylogenetic covariance estimated by Pagel's λ (Orme 2013). Both ages and

diversification rates were log-transformed. Models were not fitted with both predictors

simultaneously as they were highly correlated (Spearman's $r < -0.79$). We repeated this

analysis in conifers and palms, and again did not simultaneously fit age and diversification

rates given high correlations ($r = -0.78$ to -0.91). Fit of the PGLS was summarised by the

correlation coefficient *r* between predicted and observed values.

For conifers and palms, we also tested whether extinction risk was associated with

younger species and how this was influenced by range dynamics. We first fitted logistic

regression models to threat status as a function of species age using penalised maximum-

likelihood and accounted for phylogenetic non-independence of species with the *phylolm* R

package (Ho et al. 2014). Predictors were scaled to a mean of 0 and standard deviation of 1

to compare effects.

170   We also tested how potential range size was associated with species age in both

171  conifers and palms. First, we used PGLS to test whether older ages correlated with larger

172  range sizes, which by definition reduce extinction risk (IUCN 2016), and allowed the effect

173  to vary with threat status (i.e. statistical interaction). We expected threatened species would,

174  by definition, always have relatively small ranges, producing an invariant or weak age-range

175  association. By contrast, non-threatened species should reach larger ranges with time if the

176  age-and-area hypotheses was supported, whereas the reverse could be expected under the

177  specialism hypothesis. One limitation with this analysis is that it does not compare

178  threatened and non-threatened species of the same age, and so can introduce biases if there

179  are systematic differences in the ages of these two groups.

180   To further analyse how potential range size was associated with species age, we

181  undertook a second comparison that focused on pairs of sister species with contrasting threat

182  status. For each pair, we calculated the difference in potential range size between the sisters,

183  so as to avoid pseudoreplication, and correlated this with their age. We compared this

184  association to when sisters had the same threat status to test the null hypothesis that being

185  threatened with extinction does not change age-range associations. Focusing on sister pairs

186  was desirable because it can minimize factors that confound age-range associations, such as

187  unobserved extinctions (Hodge and Bellwood 2015). Range differences can also shed light

188  on the underlying mode of speciation. For example, there may be greater disparity in the

189  ranges of young species pairs under peripatric as opposed to allopatric speciation (Gaston

190  1998; Hodge and Bellwood 2015), resulting in a negative correlation between age and range

191  asymmetry. We tested if this correlation was different from randomly sampling the same

192  number of sister pairs 1,000 times, but choosing those where both members of the pair had

193  the same threat status. We chose both species to be non-threatened for conifers and both to

be threatened for palms as most identical species pairs in the two clades fell into these two

categories (70/85 and 51/68, respectively). R code to perform our analyses is in Data S1.

Results

We found that relatively more species were threatened with extinction in faster diversifying

genera (for ε of 0.0, 0.5, 0.9: $t_{507} = 3.64, 3.73, 3.83$, respectively; $p < 0.001$ and $r = 0.15$ for

all). The mean proportion of a genus threatened with extinction more than doubled from 36%

to 84% between the slowest and fastest diversifying genera (fig. 1a). Although these results

could have arisen because faster diversifying genera were younger (fig. 1b), as genus age was

negatively associated with risk status ($t_{507} = -2.82, p = 0.005, r = 0.14$), diversification rate

was not a simple proxy for age as it had larger effect sizes. A caveat is that we did find some

bias in our dataset. Sampled genera were older and slower diversifying, on average, than

obtained by applying our sampling criteria to the initial tree (i.e. before intersecting with

threat status; table B2). Repeating our analyses with only the genera from the more complete

conifer and palm species-level datasets was also inconclusive (table B3), potentially because

of small sample sizes ($n < 70$; fig. B1). Many conifer genera were also highly threatened

despite being old and slowly diversifying (fig. B2).

In contrast to our finding across the plant ToL, analyses with the more complete

species-level datasets revealed that older conifers but not palms were relatively more

threatened by extinction ($z_{431} = 2.17, p = 0.030$ and $z_{545} = -1.70, p = 0.089$, respectively; fig.

2a). The absolute mean effect $\pm$ SE was nearly double in the conifers (0.27 ± 0.12 vs $-0.14 \pm$

0.08 on log-scale), leading to a 31% absolute increase in the probability of being threatened

over the range of observed ages (fig. 2b).

A smaller potential range size increased the extinction risk of older conifers,

supporting the specialism hypothesis. We specifically found that non-threatened conifers had

narrower ranges as their age increased relative to sister species that were threatened (fig. 3);

ranges in neither threat status independently changed with age (table B1). As the age of

conifers increased, this difference between sister-species pairs of contrasting threat status was

larger than expected if sisters had the same threat status ($r = -0.27$, $p = 0.025$; fig. 3a).

Contrasting threat status did not alter correlations between age and potential range size in

palms, consistent with the lack of an age-extinction association ($r = -0.14$, $p = 0.222$), and

there was no correlation between species age and absolute range size (table B1). Our results

with conifers and palms were also not simply an artefact of biased sampling as ages and rates

did not markedly differ from observations across entire clades, i.e. before filtering with IUCN

data (table B4).

Discussion

Our results implicated range size as a proximate explanation for why clade age and

diversification rate were associated with extinction risk in plants. Although our findings

across the wider plant ToL contrasted those in conifers, they were consistent with the age-

and-area hypothesis in at least two ways. First, young species tend to occupy narrower

geographic and adaptive spaces (Castiglione et al. 2017), particularly as most plant speciation

involves vicarance (Davies et al. 2011; Anacker and Strauss 2014; Igea et al. 2015). Time

may consequently be required for post-speciation range expansions despite much of the

available area remaining favourable for establishment (Pigot et al. 2010; Pigot and Tobias

2013; Anacker and Strauss 2014). Second, if species diversification is density-dependent,

such as because of limited resources, then younger lineages that occupy smaller ranges will

tend to leave more niche space available that can be filled by more young species (Rabosky

and Hurlbert 2015). The consequent increase in rates of species diversification will again

elevate extinction risk in younger lineages if reproductive isolation arises within small

geographic and adaptive spaces. Time-dependent range expansions may be unnecessary

under other modes of speciation, e.g. parapatry or sympatry (Pigot et al. 2010), and if range

expansion is not limited post-speciation (Schurr et al. 2007). These differences in modes of

speciation can also help explain the lack of consistent evidence for age-dependent extinction

across the large taxonomic scale in our study and across animals (Gaston and Blackburn

1997; Johnson et al. 2002; Davies et al. 2011; Arregoitia et al. 2013; Greenberg and Mooers

2017).

Older conifer species were more threatened with extinction at least partly because

they had smaller potential ranges, supporting the evolutionary specialism hypothesis. Many

old conifers evolved during warmer wetter climates, where they occupied larger ranges than

in the present day (Farjon 1996; Jordan et al. 2016). Cycadales, which are closely related to

conifers, have undergone similar range contractions because of global cooling, resulting in

high extinction risk (Yessoufou et al. 2017). By contrast, wide ranging conifers can appear

young if they have successively given rise to other small-ranged species at the periphery of

their range (Hodge and Bellwood 2015). This scenario of range contraction and budding in

conifers contrasts with palms (Leslie et al. 2012; Condamine et al. 2017), which may have

formed new species by dispersing into new habitats that were subject to divergent adaptation

rather than vicariance (Savolainen et al. 2006; Roncal et al. 2015). Consequently, palms may

lack age-range associations that influence extinction risk. We also cannot exclude the

possibility that palm species that were susceptible to environmental change have already gone

extinct or traits that make them more prone to extinction are not taxonomically conserved,

resulting in no signature of taxon age on extinction (Arregoitia et al. 2013).

Our findings suggest that macro-evolutionary dynamics have some value for

biodiversity conservation. Specifically, we found that these dynamics provided an indicator

of contemporary extinction risk that might be easier to derive for large numbers of taxa than

detailed species-level assessments. Macro-evolutionary dynamics might also offer insight

into the vulnerability of species to future change, as the smaller population and range sizes

that make some species prone to extinction are likely to be carried into the future (Condamine

et al. 2013). Although our results must be interpreted with caution, given biases inherent to

our datasets, they provide new evidence that lineages span a continuum from little species

turnover to producing fast diversifying and extinction-prone taxa (Greenberg and Mooers

2017). The effect of age that we found at different taxonomic scales also suggests similar

patterns should emerge when the plant Tree of Life becomes more densely sampled.

Data Accessibility

Data will be deposited in the Dryad repository upon acceptance.

Acknowledgements

We thank G. Tanentzapf for the question that inspired this study and G. Jordan for sharing the

conifer phylogeny. A Mooers, D. Greenberg, and one anonymous reviewer provided

comments that improved an earlier version of this manuscript.

Literature Cited

Anacker, B. L., and S. Y. Strauss. 2014. The geography and ecology of plant speciation:

range overlap and niche divergence in sister species. Proceedings of the Royal Society

B 281:20132980.

Arregoitia, L. D. V., S. P. Blomberg, and D. O. Fisher. 2013. Phylogenetic correlates of

extinction risk in mammals: species in older lineages are not at greater risk.

Proceedings of the Royal Society B 280:20131092.

Baldwin, B. G. 2014. Origins of plant diversity in the California Floristic Province. Annual

Review of Ecology, Evolution, and Systematics 45:347–369.

Bennett, P. M., and I. P. F. Owens. 1997. Variation in extinction risk among birds: chance or

evolutionary predisposition? Proceedings of the Royal Society B 264:401–408.

Castiglione, S., A. Mondanaro, M. Melchionna, C. Serio, M. Di Febbraro, F. Carotenuto, and

P. Raia. 2017. Diversification rates and the evolution of species range size frequency

distribution. Frontiers in Ecology and Evolution 5:147.

Cayuela, L., Í. G. la Cerda, F. S. Albuquerque, and D. J. Golicher. 2012. taxonstand: An r

package for species names standardisation in vegetation databases. Methods in

Ecology and Evolution 3:1078–1083.

Ceolin, G. B., and E. L. H. Giehl. 2017. A little bit everyday: range size determinants in

Arachis (Fabaceae), a dispersal-limited group. Journal of Biogeography 44:2798–

2807.

Condamine, F. L., A. B. Leslie, and A. Antonelli. 2017. Ancient islands acted as refugia and

pumps for conifer diversity. Cladistics 33:69–92.

Davies, T. J., G. F. Smith, D. U. Bellstedt, J. S. Boatwright, B. Bytebier, R. M. Cowling, F.

Forest, et al. 2011. Extinction risk and diversification are linked in a plant biodiversity

hotspot. PLOS Biology 9:e1000620.

Farjon, A. 1996. Biodiversity of *Pinus* (Pinaceae) in Mexico: speciation and palaeo-

endemism. Botanical Journal of the Linnean Society 121:365–384.

Faurby, S., W. L. Eiserhardt, W. J. Baker, and J.-C. Svenning. 2016. An all-evidence species-

level supertree for the palms (Arecaceae). Molecular Phylogenetics and Evolution

100:57–69.

Garamszegi, L. Z., and A. P. Møller. 2010. Effects of sample size and intraspecific variation

in phylogenetic comparative studies: a meta-analytic review. Biological Reviews

85:797–805.

Gaston, K. J. 1998. Species-range size distributions: products of speciation, extinction and

transformation. Philosophical Transactions of the Royal Society B 353:219–230.

Gaston, K. J., and T. M. Blackburn. 1997. Evolutionary age and risk of extinction in the

global avifauna. Evolutionary Ecology 11:557–565.

Greenberg, D. A., and A. Ø. Mooers. 2017. Linking speciation to extinction: Diversification

raises contemporary extinction risk in amphibians. Evolution Letters 1:40–48.

Higgins, S. I., R. B. O'Hara, O. Bykova, M. D. Cramer, I. Chuine, E.-M. Gerstner, T.

Hickler, et al. 2012. A physiological analogy of the niche for projecting the potential

distribution of plants. Journal of Biogeography 39:2132–2145.

Ho, T., L. Si, and C. Ané. 2014. A linear-time algorithm for Gaussian and non-Gaussian trait

evolution models. Systematic Biology 63:397–408.

Hodge, J., and D. R. Bellwood. 2015. On the relationship between species age and

geographical range in reef fishes: are widespread species older than they seem?

Global Ecology and Biogeography 24:495–505.

Igea, J., D. Bogarín, A. S. T. Papadopulos, and V. Savolainen. 2015. A comparative analysis

of island floras challenges taxonomy-based biogeographical models of speciation.

Evolution 69:482–491.

International Union for Conservation of Nature [IUCN]. 2016. The IUCN Red List of

Threatened Species. Available at: http://www.iucnredlist.org

Ives, A. R. 2015. For testing the significance of regression coefficients, go ahead and log-

transform count data. Methods in Ecology and Evolution 6:828–835.

Jetz, W., and R. P. Freckleton. 2015. Towards a general framework for predicting threat

status of data-deficient species from phylogenetic, spatial and environmental

information. Philosophical Transactions of the Royal Society B 370:20140016.

Johnson, C. N., S. Delean, and A. Balmford. 2002. Phylogeny and the selectivity of

extinction in Australian marsupials. Animal Conservation 5:135–142.

Jordan, G. J., P. A. Harrison, J. R. P. Worth, G. J. Williamson, and J. B. Kirkpatrick. 2016.

Palaeoendemic plants provide evidence for persistence of open, well-watered

vegetation since the Cretaceous. Global Ecology and Biogeography 25:127–140.

Leslie, A. B., J. M. Beaulieu, H. S. Rai, P. R. Crane, M. J. Donoghue, and S. Mathews. 2012.

Hemisphere-scale differences in conifer evolutionary dynamics. Proceedings of the

National Academy of Sciences 109:16217–16221.

Liow, L. H. 2007. Lineages with long durations are old and morphologically average: an

analysis using multiple datasets. Evolution 61:885–901.

Magallon, S., and M. J. Sanderson. 2001. Absolute diversification rates in angiosperm clades.

Evolution 55:1762–1780.

Orme, D. 2013. The caper Package: Comparative Analysis of Phylogenetics and Evolution in

R. Available at: https://cran.r-project.org/web/packages/caper/vignettes/caper.pdf

Paul, J. R., C. Morton, C. M. Taylor, and S. J. Tonsor. 2009. Evolutionary time for dispersal

limits the extent but not the occupancy of species' potential ranges in the tropical

plant genus *Psychotria* (Rubiaceae). American Naturalist 173:188–199.

Pennell, M. W., R. G. FitzJohn, and W. K. Cornwell. 2016. A simple approach for

maximizing the overlap of phylogenetic and comparative data. Methods in Ecology

and Evolution 7:751–758.

Pigot, A. L., A. B. Phillimore, I. P. F. Owens, and C. D. L. Orme. 2010. The shape and

temporal dynamics of phylogenetic trees arising from geographic speciation.

Systematic Biology 59:660–673.

Pigot, A. L., and J. A. Tobias. 2013. Species interactions constrain geographic range

expansion over evolutionary time. Ecology Letters 16:330–338.

Pimm, S. L., C. N. Jenkins, R. Abell, T. M. Brooks, J. L. Gittleman, L. N. Joppa, P. H.

Raven, et al. 2014. The biodiversity of species and their rates of extinction,

distribution, and protection. Science 344:1246752.

Purvis, A., J. L. Gittleman, G. Cowlishaw, and G. M. Mace. 2000. Predicting extinction risk

in declining species. Proceedings of the Royal Society B 267:1947–1952.

Qian, H., and Y. Jin. 2016. An updated megaphylogeny of plants, a tool for generating plant

phylogenies and an analysis of phylogenetic community structure. Journal of Plant

Ecology 9:233–239.

Rabosky, D. L., and A. H. Hurlbert. 2015. Species richness at continental scales is dominated

by ecological limits. American Naturalist 185:572–583.

Roncal, J., M. Couderc, P. Baby, F. Kahn, B. Millán, A. W. Meerow, and J.-C. Pintaud. 2015.

Palm diversification in two geologically contrasting regions of western Amazonia.

Journal of Biogeography 42:1503–1513.

Savolainen, V., M.-C. Anstett, C. Lexer, I. Hutton, J. J. Clarkson, M. V. Norup, M. P. Powell,

et al. 2006. Sympatric speciation in palms on an oceanic island. Nature 441:210–213.

Scholl, J. P., and J. J. Wiens. 2016. Diversification rates and species richness across the Tree

of Life. Proceedings of the Royal Society B 283:20161334.

Schwery, O., and B. C. O'Meara. 2016. MonoPhy: a simple R package to find and visualize

monophyly issues. PeerJ Computer Science 2:e56.

Schwartz, M. W., and D. Simberloff. Taxon size predicts rates of rarity in vascular plants.

Ecology Letters 4:464–469.

Slatyer, R. A., M. Hirst, and J. P. Sexton. 2013. Niche breadth predicts geographical range

size: a general ecological pattern. Ecology Letters 16:1104–1114.

Tanentzap, A. J., A. J. Brandt, R. D. Smissen, P. B. Heenan, T. Fukami, and W. G. Lee.

2015. When do plant radiations influence community assembly? The importance of

historical contingency in the race for niche space. New Phytologist 207:468–479.

Valente, L. M., V. Savolainen, and P. Vargas. 2010. Unparalleled rates of species

diversification in Europe. Proceedings of the Royal Society B 277:rspb20092163.

Vamosi, J. C., and J. R. U. Wilson. 2008. Nonrandom extinction leads to elevated loss of

angiosperm evolutionary history. Ecology Letters 11:1047–1053.

Williams, S. E., Williams, Y. M., VanDerWal, J., Isaac, J. L., Shoo, L. P., and C. N. Johnson.

2009. Ecological specialization and population size in a biodiversity hotspot: How

rare species avoid extinction. Proceedings of the National Academy of Sciences 106:

19737–19741.

Willis, J. C. 1926. Age and area. Quarterly Review of Biology 1:553–571.

Wilson, E. O. 1959. Adaptive shift and dispersal in a tropical ant fauna. Evolution 13:122–

144.

Yessoufou, K., B. H. Daru, R. Tafirei, H. O. Elansary, and I. Rampedi. 2017. Integrating

biogeography, threat and evolutionary data to explore extinction crisis in the

taxonomic group of cycads. Ecology and Evolution 7:2735–2746.

Zanne A. E., D. C. Tank, W. K. Cornwell, J. M. Eastman, S. A. Smith, R. G. FitzJohn, D J.

McGlinn, et al. 2014. Three keys to the radiation of angiosperms into freezing

environments. Nature 506:89–92.

Žliobaitė, I., M. Fortelius, and N. C. Stenseth. 2017. Reconciling taxon senescence with the

Red Queen's hypothesis. Nature 552:92–95.

Figure 1. More species are threatened with extinction in (**a**) faster diversifying and (**b**)

younger genera. Diversification was estimated for $\varepsilon = 0.50$. Solid lines are mean

associations estimated by PGLS.

Figure 2. Older conifers but not palms have a greater probability of being threatened by

extinction. (**a**) Boxplot for stem ages of conifer (white, *n* = 433) and palm (grey, *n* = 547)

species that were classified as either threatened or non-threatened. Solid line is median, box

is inter-quartile range, whiskers extend 1.5-times the interquartile range, and points are

outliers. (**b**) Change in probability of a conifer being classified as threatened with species

age. Solid line is mean association estimated by phylogenetic logistic regression.

Figure 3. Differences in range size between sister conifers of contrasting threat status decrease with their age. For each sister pair of non-threatened (NT) and threatened (T) taxa we calculated the difference in log-transformed potential range sizes. Solid line is the slope for the corresponding correlation coefficient *r*. Inset shows frequency distribution of *r* calculated for 1,000 random simulations of sister pairs of the same threat category, with vertical line denoting observed correlation for contrasting threat status, i.e. corresponding to plotted data points.