De novo draft assembly of the *Botrylloides leachii* genome provides further insight into tunicate evolution.

Simon Blanchoud[1#], Kim Rutherford[2], Lisa Zondag[1], Neil Gemmell[2] and Megan J Wilson[1*]

1 Developmental Biology and Genomics Laboratory

Department of Anatomy, School of Biomedical Sciences, University of Otago, P.O. Box 56,

Dunedin 9054, New Zealand

Current address: Department of Zoology, University of Fribourg, Switzerland

* Corresponding author:

Email: meganj.wilson@otago.ac.nz

Ph. +64 3 4704695

Fax: +64 479 7254

Keywords: chordate, regeneration, *Botrylloides leachii*, ascidian, tunicate, genome, evolution

Abstract (250 words)

Tunicates are marine invertebrates that compose the closest phylogenetic group to the vertebrates. This chordate subphylum contains a particularly diverse range of reproductive methods, regenerative abilities and life-history strategies. Consequently, tunicates provide an extraordinary perspective into the emergence and diversity of chordate traits. Currently published tunicate genomes include three Phlebobranchiae, one Thaliacean, one Larvacean and one Stolidobranchian. To gain further insights into the evolution of the tunicate phylum, we have sequenced the genome of the colonial Stolidobranchian *Botrylloides leachii*.

We have produced a high-quality (90% BUSCO genes) 159 Mb assembly, containing 82 % of the predicted 194 Mb. The *B. leachii* genome is much smaller than that of *Botryllus schlosseri (*725 Mb), but comparable to those of *C. intestinalis* and *M. oculata* (both 160 Mb). This difference is largely due to an increase in repetitive DNA content in *B. schlosseri*. By analyzing the structure and composition of the conserved homeobox gene clusters, we identified many examples of multiple cluster breaks and gene dispersion, suggesting that several lineage-specific genome rearrangements occurred during tunicate evolution.

In addition, we investigate molecular pathways commonly associated with regeneration and development. We found lineage-specific gene gain and loss within the Wnt, Notch and retinoic acid pathways. Such examples of genetic changes to key evolutionary conserved pathways may underlie some of the diverse regenerative abilities observed in the tunicate subphylum. These results, combined with the relatively recent separation from their last common ancestor (630 MYA), supports the widely held view that tunicate genomes are evolving particularly rapidly.

Introduction

Tunicates are a group of marine suspension-feeding hermaphrodites found worldwide in the inter- or sub-tidal region of the seas. This subphylum of invertebrates is phylogenetically located in the Chordata phylum, between the more basal Cephalochordata and the higher Vertebrata, of which they are considered the closest relatives (Fig. 1A; (Delsuc *et al.*, 2006)). These organisms include a wide range of reproductive methods, regenerative abilities, developmental strategies and life cycles (Lemaire *et al.*, 2008). Importantly, and despite a drastically different body plan during their adult life cycle, tunicates have organs as well as a tissue complexity incipient to that of vertebrates (Fig. 1A), including a heart, a notochord, an endostyle and a vascular system (Millar, 1971). In addition, this group of animals is undergoing faster genomic evolution compared to vertebrates through a higher nucleotide substitution rate in both their nuclear and mitochondrial genomes (Tsagkogeorga *et al.*, 2010, 2012; Rubinstein *et al.*, 2013; Berna and Alvarez-Valin, 2014). Therefore, this chordate subphylum provides an excellent opportunity to study the origin of vertebrates, the emergence of clade specific traits and the function of conserved molecular mechanisms. Biological features that can be investigated in tunicates include, among others, the evolution of colonialism, neoteny, sessilness, and budding. However, there are currently only seven Tunicata genomes publicly available, of which three have been well annotated. There is thus a paucity in the sampling of this very diverse subphylum.

Tunicates are separated into eight clades: Phlebobranchia (class Ascidiacea), Molgulida (class Ascidiacea), Copelata (class Appendicularia), Pyrosomida (class Thaliacea), Salpida (class Thaliacea), Doliolida (class Thaliacea), Aplousobranchia (class Ascidiacea), and Stolidobranchia (class Ascidiacea). Phlebobranchiae and Molgulida are clades of sessile solitary benthic ascidians. These animals reproduce sexually, releasing eggs through their atrial siphon for external fertilization, hence producing a motile larva. These larvae will

colonize novel environments, attach to a submersed substrate and undergo metamorphosis

into a sessile filter-feeding adult. These ascidians are capable of regenerating a limited set of

organs, including their oral siphon (Auger *et al.*, 2010) although regeneration capability

reduces as they age (Jeffery, 2015). These clades are currently the most sampled ones with

five published genomes (*Ciona intestinalis*, *Ciona savigny, Molgula oculata*, *Molgula occulta*,

Molgula occidentalis; (Dehal *et al.*, 2002a; Small *et al.*, 2007; Stolfi *et al.*, 2014), and two yet

unpublished species (*Phallusia mamilata*, *Phallusia fumigata*; (Brozovic *et al.*, 2016)). *C.*

intestinalis and *O. dioika* where the first Tunicate species to have been sequenced and are the

best characterized.

Copelata is a class of planktonic free-swimming organisms that possess chordate traits

common to all tunicate larvae including a notochord, neural tube and pharyngeal slits. These

social neotenic animals form dioecious communities where each individual lives inside a

special external mucous structure, termed house, which concentrates and funnels their food.

Oikopleura dioika is the sole example of the Copelatan to have its genome sequenced, showing

exceptional compaction (70 Mb; (Seo *et al.*, 2001)). Whether these animals can undergo

regeneration has not yet been assessed.

Pyrosomida, Salpida and Doliolida are clades of planktonic pelagic animals forming

cylindrical free-floating compound colonies (Piette and Lemaire, 2015). These organisms can

reproduce both sexually, through autogamy to initiate novel colonies, as well as asexually,

through stolonial budding to increase the size of the colony. Owing to their peculiar life cycle

and habitat, these tunicates have not been extensively studied, no genome has been

sequenced and whether they can undergo regeneration remains unknown.

Aplousobranchia and Stolidobranchia consist of both solitary and colonial sessile

benthic organisms. Colonial tunicates are capable of both sexual, through autogamy, and

asexual reproduction, through a wide range of budding types (palleal, vascular, stolonial,

pyloric and strobilation; (Brown and Swalla, 2012)). In addition, these compound organisms can undergo whole-body regeneration (WBR; reviewed in (Kürn *et al.*, 2011)). Colonial ascidians are emerging as unique and increasingly popular model organisms for a variety of studies including immunobiology, allorecognition, angiogenesis and WBR (Rinkevich *et al.*, 1995, 2013; Ballarin *et al.*, 2001; Manni *et al.*, 2007; Rinkevich, Douek, *et al.*, 2007; Gasparini *et al.*, 2008; Franchi *et al.*, 2011; Lauzon *et al.*, 2013). While the genome of two solitary Aplousobranchian are currently being assembled (*Halocynthia rorezi*, *Halocynthia aurantium*; (Brozovic *et al.*, 2016)), colonial tunicates were only exemplified by *Botryllus schlosseri*. Furthermore, this ascidian which a significant expantion of its genome size when compared to the other available Tunicate genomes (725 Mb). To further investigate this fascinating subphylum and assess whether genome expansion is a prerequisite for coloniality, we have assembled and analysed the genome sequence of *Botrylloides leachii* (class Ascidiacea, order Stolidobranchia; (Savigny, 1816)).

The viviparous colonial ascidian *B. leachii* can reproduce sexually through a tadpole stage that allows the settlement of a novel colony onto a new substrate (Fig. 1B). Each colony is composed of genetically identical adults (termed zooids) organized in ladder-like systems and embedded in gelatinous matrix (tunic). While each adult has its own heart, they all share a common vascular system embedded within the tunic. In the presence of sufficient food supply, the size of the colony doubles approximately every 20 days through synchronized asexual reproduction, known as palleal budding. During this process each adult produces two daughter zooids that ultimately replace the mother, which is then resorbed by the colony. In addition, upon loss of all zooids from the colony, *B. leachii* can undergo whole-body regeneration and restore a single fully-functional adult in as little as 10 days from a small piece of its vascular system (Rinkevich *et al.*, 1995). Furthermore, when facing unfavorable environmental conditions, these colonial tunicates can enter into hibernation, whereby all

zooids undergo regression and are resorbed by the remaining vascular system. When a favorable environment is restored, mature adults will be restored to reestablish the colony (Burighel *et al.*, 1976).

We have assembled and annotated the first *de novo* draft genome of the *B. leachii* by taking advantage of our recently published transcriptomes (Zondag *et al.*, 2016). Using this genome, we have then undertaken a large-scale comparison of the four best-annotated ascidian genomes *(B. schlosser*i, *C. intestinalis, M. oculat*a and *O. dioika*) to gain insights into some of the diverse biological abilities that have evolved within the Tunicata.

Results

Genome assembly and annotation

To minimize contamination from marine algae and bacteria typically present in the pharyngeal basket of feeding *B. leachii*, we isolated genomic DNA from embryos of a single wild *B. leachii* colony. Genomic DNA was used to produce two libraries: one short-range consisting of 19090212 fragments (300 bp) of which 100 bp were paired-end sequenced - important for obtaining high coverage - and a second long-range mate pair with 31780788 fragments (1.5-15 kb size range, median ~3 kb) of which 250 bp were paired-end sequenced – important for scaffolding the assembly. Following quality checks, low quality reads were removed and sequencing adaptors were trimmed, thus resulting in a high-quality dataset of 86644308 paired-end and 12112004 single-end sequences (100% with a mean Phred score >= 30, <1% with an adapter sequence, Fig. S1).

We then followed a reference-free genome characterization (Simpson, 2014); provided with statistics from the human, fish (*Maylandia zebra*; (Bradnam *et al.*, 2013)), bird (*Melopsittacus undulatus;* (Bradnam *et al.*, 2013)) and oyster (*Crassostrea gigas*, (Zhang *et al.*, 2012)) genomes for comparison; to estimate three properties of the *B. leachii* genome. First, k-mer count statistics were used to estimate the genome size to be 194.2 Mb (194153277 bp). This size is thus comparable to that of the solitary *C. intestinalis, C. savigny and M. oculata* (160 Mb, 190 Mb and 160 Mb, respectively; (Dehal *et al.*, 2002a; Small *et al.*, 2007; Stolfi *et al.*, 2014), larger than the compacted 70 Mb genome of *O. dioika* but appreciably smaller than the predicted 725 Mb genome of the related colonial ascidian *B. schlosseri*, of which 580 Mb have been sequenced (Voskoboynik *et al.*, 2013). Second, by quantifying the structure of the de Brujin graph obtained using the k-mer counts, the computational complexity of the assembly

was estimated (sequencing errors 1/213, allelic differences 1/233, genomic repeats 1/2439). With a cumulative occurrence of 1/106, the *B. leachii* genome is similar to that of bird, more variable than those of fish and human, but still quite less complex than the notably difficult oyster genome (Fig. S1). Third, sequence coverage was estimated using the distribution of 51-mers counts, showing a well-separated bimodal distribution with a true-genomic k-mers maximum at 31x coverage, similar to the human genome but higher than both the fish and the bird. Overall, these metrics suggest that *B. leachii* has a genome well suited for *de novo* assembly and that our sequencing could result in a high quality assembly .

De novo assembly using Metassembler (Wences and Schatz, 2015) produced a genome of 159132706 bp (estimated percentage of genome assembled is 82%), with an average sequencing coverage of 66x (after adaptor trimming). The assembly is composed of 1778 scaffolds, with a N50 scaffold length of 209776 and a L50 scaffold count of 223. The 7783 contigs, with a N50 length of 48085, and a L50 count of 781, represent a total size of 146061259 (92%, Table 1). To evaluate the completeness of our assembly, we used the Benchmarking Universal Single-Copy Orthologs (BUSCO; (Simão *et al.*, 2015)). This tool provides a quantitative measure of genome completeness by verifying the presence of a set of manually curated and highly conserved genes. Out of the 843 orthologues selected in metazoans, 760 (90%) were found in our assembly of the *B. leachii* genome (File S1), a relatively high score when compared to the BUSCO score of frequently used genome assemblies such as *Homo sapiens* (89%, GCA_000001405.15) and *Mus musculus* (78%, GCA_000001635.4). In addition, we took advantage of our previous assembly of the *B. leachii* transcriptome (Zondag *et al.*, 2016) to further assess the quality of our genome. Using BLAT (Kent, 2002), we were able to map 93 % of transcript sequences (48510/52004) onto our assembly. Finally, we further estimated the quality of our assembly by mapping the proteomes of the available tunicate genomes using tBLASTn (Camacho *et al.*, 2009): *C.*

intestinalis 71% (4233/14740), *C. savigny* 81% (3835/20155), *M. oculata* 77% (3828/16616) and *B. schlosseri* 71% (13575/46519). Overall, these results indicate that our *de novo* genome is largely complete and suitable for annotation.

Ab initio genome annotation was performed using MAKER2 (Holt and Yandell, 2011) and predicted 15839 coding genes, of which 13507 could be classified using InterProScan (Jones *et al.*, 2014). Comparing these predictions with our mapping of the transcriptome, we found out that 83% of our aligned cDNA (40188/48510) mapped to a predicted gene locus thus spanning 78 % of the annotated genes (12395/15839). In addition, a total of 4213 non-coding sequences were predicted using Infernal (Nawrocki and Eddy, 2013), Snoscan (Lowe, 1999) and tRNAscan-SE (Lowe, 1999). Finally, repetitive elements were annotated using RepeatMasker (Smit *et al.*, 2015) and a specie-specific library created using the RepeatModeler module (Smit and Hubley, 2015). 18% of the genome was identified as containing repetitive elements (Table 2), a majority (17%) of these being interspersed repeats. This proportion is similar to that found in other tunicates including *C. intestinalis* (17%), M. oculata (22 %) and *O. dioica* (15%), while being lower than that in *B. schlosseri* (60%) and *C. savigny* (33 %).

Ancient gene linkages are fragmented in tunicate genomes

Ancient gene linkages are spatially restricted and highly conserved sets of genes. These clusters arose in a common ancestor and were preserved because of a common regulatory mechanism such as cis-regulatory elements located within the cluster The homeobox-containing *Hox* gene family, typically composed of 13 members in vertebrates (Hoegg and Meyer, 2005), is among the best-studied examples of such ancient gene cluster and is critical for the correct embryonic development (Pearson *et al.*, 2005). In particular, the linear genomic arrangement within the *Hox* cluster reflects their spatial expression along the

anterior-posterior body axis (Pascual-Anaya *et al.*, 2013), which establishes regional identity across this axis. The basal cephalochordate *B. floridae* has all 13 *hox* genes located in a single stereotypical cluster, along with an additional 14th gene (Fig. 2B; (Takatori *et al.*, 2008)), suggesting that the chordate ancestor also had an intact cluster. However in tunicates, this clustering appears to be lost. In *C. intestinalis*, the nine identified *Hox* genes are distributed across five scaffolds, with linkages preserved only between *Hox2, Hox3* and *Hox4; Hox5* and *Hox6; Hox12* and *Hox13* (Fig. 2; (Spagnuolo *et al.*, 2003; Wada *et al.*, 2003)). In *O.dioica,* the total number of *Hox* genes is further reduced to eight, split between 6 scaffolds, including a duplication of *Hox9* (Fig. 2A; (Edvardsen *et al.*, 2005)). In *M. oculata* we could identify only six *Hox* genes, divided between 4 scaffolds, with clustering retained for the *Hox10*, *Hox11* and *Hox12* genes (Fig. 2). In Botryllidae genomes, the same seven *Hox* genes are conserved (Fig 2B), with a preserved linkage between *Hox10*, *Hox12* and *Hox13* in *B. leachii* and a three copies of *Hox5* present in *B. schlosseri*. Altogether these genomic distributions of the *Hox* cluster genes supports the hypothesis that reduction and separation of this ancient gene linkage occurred at the base of tunicate lineage (Edvardsen *et al.*, 2005). In addition, *Hox9* appears to be specifically retained in neotenic Tunicates while there is no pattern of consereved *Hox* cluster genes specific to colonial ascidians.

A second ancient homeobox-containing gene linkage is the *NK* cluster. This cluster, which was seemingly inherited from the last common ancestor of bilaterians (Luke *et al.*, 2003), consists of *Msx, Lbx, Tlx, NKx1, NKx3, NKx4* and *NKx5* (Fig. 3). In *B. floridae*, linkages between *Msx, NKx4* and *NKx3*; as well as between *Lbx* and *Tlx* provide evidence of retained ancestral clustering while *NKx5* was lost (Fig. 3; (Luke *et al.*, 2003)). However in vertebrates, NKx5 is still present while only the gene linkages between *Lbx* and *Tlx* as well as between *Nkx4* and *Nkx3* remain (Fig. 3; (Garcia-Fernàndez, 2005)). To further clarify the evolution of this ancestral cluster in tunicates, we determined the structure of the NK cluster within five

ascidian genomes. In all these species, *NKx1* is absent and no evidence of clustering could be found with all identified orthologues located on different scaffolds (Fig. 3). In *C. intestinalis*, *M. oculata* and *O. dioica* only five members of this cluster remain, with the loss of either *Lbx* or *Tlx* as well as of *NKx3* and the duplication of the orthologue of *NKx4* (Fig. 3). By contrast, in the colonial tunicates *B. leachii* and *B. schlosseri*, *Tbx*, *Lbx* and *NKx3* are all present. In *B. schlosseri*, *Msx1* is absent and *NKx4* duplicated. In the *B. leachii* genome, *NK1* is the only ancestral cluster member to be missing and *Nk5* has been duplicated (Fig. 3). Altogether, these results suggest that there has been a loss of *NKx5* in Cephalochordate, one of *NKx1* in Tunicate and that the conjunction of *NKx3*, *Lbx* and *Tbx* is specific to colonial ascidians.

A third ancient linkage that we investigated is the pharyngeal cluster, a gene group present in hemichordates, echinoderm and vertebrates genomes that is considered to be Deuterosome specific (Simakov *et al.*, 2015). The cluster groups *foxhead domain protein* (*FoxA*), *NKx2 (NKx2.2 and Nkx2.1)*, *Pax1/9*, *mitochondrial solute carrier family 25 member 21* (*slc25A21)*, *mirror-image polydactyly 1 protein* (*mipol1*), *egl nine homolog 3* (*egln3*) and *dehydrogenase/reductase member 7* (*dhrs7*). Among these, s*lc25a21, Pax1/9*, *mipol1* and *FoxA* pairs are also found in protostomes suggesting an even more ancient origin (Simakov *et al.*, 2015). The pharyngeal cluster is thought to have arisen due to the location of the regulatory elements of *Pax1/9* and *FoxA* within the introns of *slc25A21* and *mipol1* (Santagati *et al.*, 2003; Wang *et al.*, 2007), constraining the genes to remain in tight association with each other. In the *B. floridae* genome, the entire cluster is located on the same scaffold, with the exception of the *Nkx2.1* and *Nk2.2* gene pair located on separate scaffold. In *C. intestinalis*, only orthologs of *FoxA*, slc25a29, *Pax1* and *Pax9* could be identified. Nevertheless, all of them are located on the same chromosome (Fig. 4). In *O. dioica*, the cluster appears even further reduced as, while orthologues of *FoxA*, *Pax1/9* and *Nkx2.2* genes were found on different scaffolds, only one rather distant linkage (> 1 Mb) between *a Pax-like* gene and *slc25A21* is retained. Copies of

Pax1/9, slc25a29 and Nk2.2 genes were found in *B. schlosseri*, however all on different scaffolds. In the *B. leachii* genome, *mipol1* is the sole missing gene from this cluster. However, only the pairing of a *Pax-like* and *slc25A21* genes remains (Fig. 4). For both *B. schlosseri* and *M. oculata*, there was no evidence of clustering between genes (Fig. 4) Altogether, these results suggest that most of the Tunicates did not conserve the structure of this ancient linkage, but it is unknown what consequences this would have to their expression and function.

Lineage-specific changes to cell-signaling pathways in Botryllidae genomes.

To dissect more specifically the evolution of colonial ascidians, we examined the genomes of *B. leachii* and *B. schlosseri*, looking for key components of signaling pathways required for metazoan development and regeneration. Of particular interest, we focused on the Wingless-related integration site (Wnt), Notch and Retinoic acid (RA) signaling pathways.

Wnt pathway

Wnt ligands are secreted glycoproteins that have roles in axis patterning, morphogenesis and cell specification (Loh *et al.*, 2016). The ancestral gene cluster appears to originate very early on during muti-cellular evolution and to be composed of eleven members (Kusserow *et al.*, 2005; Guder *et al.*, 2006). The *Wnt* gene family expanded to 19 members in the human genome, while independent gene loss has reduced this family reduced to 7 genes in *Drosophila melanogaster* and *Caenorhabditis elegans* (Prud'homme *et al.*, 2002). Consequently, we set out to investigate whether the *Wnt* gene family has either expanded or reduced during Tunicata speciation.

Most strikingly, we found among tunicate genomes an increase in the number of *Wnt5a* genes. *C. intestinalis* has a total of 11 *Wnt* genes, including a single *Wnt5a* gene (Fig. 5, Table S2; (Hino *et al.*, 2003)). In the compact *O. dioica* genome, this number has reduced to 6 (Wnts

3, 4, 7, 11 and 16), none of which are *Wnt5a* orthologues (Table S2). *M. oculata* has only 7 Wnt ligand genes , including three *Wnt5a-like* genes (Fig. 5, Table S2). In *B. schlosseri*, we identified 15 *Wnt* members, including seven *Wnt5a-like* genes on multiple scaffolds (Fig. 5, Table S2). Finally, in the *B. leachii* genome, fourteen *Wnt* ligand genes were identified, including four *Wnt5a* genes located on the same scaffold near *Wnt4* (Fig. 5). Overall, this suggests that an expansion through gene duplication of the Wnt5 family occurred during tunicate evolution, but was lost in some lineages.

To assess the functionality of the Wnt pathway in Tunicates, we set out to assess whether its downstream effectors are themselves present in the available genomic data. The downstream pathways activated by Wnt ligands are divided into canonical, non-canonical calcium and non-canonical planar cell polarity. The Wnt5a ligand is associated with both of the non-canonical pathways through binding of membrane receptors that include *frizzled* (*Fzd4*), *receptor tyrosine kinase-like orphan receptor 1/2* (*Ror1/2*) and *atypical tyrosine kinase receptor* (*Ryk*). Further downstream, disheveled (dsh), β-catenin (Cnntb), Axin, low-density lipoprotein receptor-related protein 5/6 (LRP5/6) and nuclear factor of activated T-cells (NFAT) are proteins essential for triggering intracellular responses to Wnt signaling (MacDonald *et al.*, 2009). We identified orthologues for each of these signaling transduction molecules in all Tunicata genomes (Table S2), with no evidence of further gene duplication events. This supports the interpretation that signaling through the Wnt pathway is functional in tunicates.

Notch pathway

Notch receptors are transmembrane proteins that are involved in cell-cell signaling during development, morphogenesis and regeneration (Hamada *et al.*, 2015). Following activation through the binding of the delta or jagged/serrate ligands, the intracellular domain

of Notch is cleaved and induces the expression of downstream target genes including the *hes (hairy and enhancer of split)* gene family members (Guruharsha *et al.*, 2012). The presence of both Notch and the Delta/Serrate/lag-2 (DSL) proteins in most metazoan genomes suggests that their last common ancestor had a single copy of each gene (Gazave *et al.*, 2009). To establish how this pathway has evolved in tunicates, we screened these genomes for the Notch receptor using the conserved lin-Notch repeat (LNR) domain, and for genes encoding probable Notch ligands such as genes from the DSL family .

In all examined genomes, only a single *Notch* receptor gene was identified while the number of ligand genes varied (Table S3). *C. intestinalis* genome contains two *DSL* genes; *O. dioica M. oculata* and *B. schlosseri* a single *DSL* one. By contrast, we found three DSL genes in *B. leachii* (Table S3). To determine the relationships between these identified tunicate DSL-like genes, a phylogeny was constructed along with other chordate DSL proteins. All three *B. leachii* genes are Delta orthologues, two of them related to the *B. schlosseri* and *Cionidae.*copy; the third one closer to the *M. oculata* and *H. roretzi* variant. The mouse, human and zebrafish delta and delta-like (DLL) proteins form a discrete clade loosely related to the genes found in Cephalochordate and Tunicate (Fig. 6, shaded box). Jagged proteins form a separate and less conserved clade (Fig. 6). The tunicate DSL-like proteins present long phylogenetic branches, suggestive of greater diversity, which is also observed in the protein alignment (Fig. S3). This suggests that the tunicate DSL proteins are diverging rapidly from each other, indicative of lineage specific evolution of DSL-like genes.

Retinoic acid signaling

Retinoic acid (RA) is an extracellular metabolite that is essential for chordate embryonic development. RA is synthesized from retinol (vitamin A) by two successive oxidation steps. In the first step, retinol dehydrogenase (RDH) transforms retinol into retinal.

Then RA is produced by aldehyde dehydrogenase (ALDH), a superfamily of enzymes with essential roles in detoxification and metabolism (Jackson *et al.*, 2011). RA influences the expression of downstream target genes by binding to the RA receptors, RAR and RXR (Fig. 7A (Cunningham and Duester, 2015)). Finally, RA is metabolized by the enzymes cytochrome P450 family 26 (Cyp26), a process key to restricting RA-induced responses to specific tissues or cell types (Ross and Zolfaghari, 2011). Components of this pathway have been found in non-chordate animals, suggesting a more ancient origin (Canestro *et al.*, 2006). This pathway has previously been shown to be required for *B. leachii* WBR and *Ciona* development, yet several genes required for RA signaling appear to be missing in *O. dioica* (Martí-Solans *et al.*, 2016).

Rdh10 is the major dehydrogenase associated with the first steps of RA production, although the Rdh16 and RdhE2 enzymes can also substitute this function (Belyaeva *et al.*, 2009, 2015; Lee *et al.*, 2009). The *O. dioica* genome has no orthologues for either *Rdh10* or *Rdh16* but it does have four genes that encode for RdhE2 proteins (Martí-Solans *et al.*, 2016). *O. dioica* also lacks both an *Aldh1*-type gene as well as a *Cyp26* gene but has a single RXR-orthologue (Table S4, (Martí-Solans *et al.*, 2016)). In contrast, the *C. intestinalis* genome, contains single copies of *Rdh10*, *Rdh16* and *RdhE2* genes and a total of four *Aldh1* genes, located on two chromosomes (Canestro et al., 2006). Consistent with *C. intestinalis*; *M. oculata*, *B. leachii* and *B. schlosseri* genomes all have single copies of *Rdh10, Rdh16* and *RdhE2* genes, as well as three *Aldh1a/b* genes on separate scaffolds (Table S4).

Three retinoic acid receptor genes were identified within the *B. leachii* genome, one of which had been cloned previously (*g03013,* (Rinkevich, Paz, *et al.*, 2007).   All three were also found in *C. intestinalis*, *M. oculata* and *B. schlosseri* genomes(Table S4). While there is only one potential *Cyp26* gene in *M. oculata*, four paralogues were identified in *B. leachii* and *B. schlosseri*. A phylogenetic analysis showed that these 4 genes group with CYP26 proteins (Fig.

347   7B, Table S4). Altogether, these results show a loss of key RA-pathway genes in *O. dioica*

348   (*Rdh10, Rdh16, Cyp26* and *Aldh1a*) , while increased copy numbers in other tunicate genomes.

Discussion

Genomic diversity within the Solidobranchia

The *B. leachii* genome, along with previous genomic analyses of other ascidian species, support the widely held view that ascidian genomes are diverse and rapidly evolving, which is particularly evident in the Solidobranchia group (Seo *et al.*, 2001; Dehal *et al.*, 2002b; Tsagkogeorga *et al.*, 2010, 2012; Bock *et al.*, 2012; Rubinstein *et al.*, 2013; Voskoboynik *et al.*, 2013; Griggio *et al.*, 2014; Stolfi *et al.*, 2014) . Nevertheless, botryllids are sufficiently similar in external appearance and morphology for early researchers to have suggested that *Botrylloides* could be a subgenus of *Botryllus* (Saito *et al.*, 2001; Nydam *et al.*, 2017) . Strikingly however, the *B. schlosseri* genome differs from that of *B. leachii*, as well as from other sequenced tunicate genomes (Table 2). In particular, the comparison between the *B. leachii* and *B. schlosseri* genome sizes (194 Mb vs 725 Mb), their fraction of repetitive sequences (18 % vs 60 %; 65 % in (Voskoboynik *et al.*, 2013)) and their predicted gene number (15'839 vs 27'000; (Voskoboynik *et al.*, 2013)) suggest a different genomic architecture. Altogether, these comparisons indicate that the *B. schlosseri* genome has undergone a significant increase in its genomic content, including retrotransposons (Table S1). In particular, there are at least two additional families in the *B. schlosseri* hAT transposon superfamily and counts of common hAT element, such as hAT-Charlie, differ dramatically (for hAT-Charlie: 366 in *B. leachii* vs 46,661 in *B. schlosseri*). DNA methylation is a key suppressor of transposon activity, changes to the methylation of transposable elements is a known driver of increased tranposition (O'Neill *et al.*, 1998; Simmen *et al.*, 1999; Suzuki *et al.*, 2007; Maumus and Quesneville, 2014). However, DNA methylation in tunicate species has been only studied in *C. intestinalis,* and is described as mosaic, gene body methylation, whereas non-coding regions, including transposons, remain unmethylated (Suzuki et al. 2007). However,

DNA methylation has not been studied in other tunicates, and it It is unknown how

retrotransposons are suppressed in tunicate genomes. Nevertheless, the observed increase in

transpositions could be a consequence of low non-coding DNA methylation, which may

contribute to the rapid genome evolution observed in tunicate species, even between closely

related species such as *B. schlosseri* and *B. leachii*.

Rapid genome evolution, and active transposable elements in particular, are proposed

to aid adaptation to new environments for invasive species (Stapley *et al.*, 2015). Differences

in the colonization ability of tunicates has been noted, not only between related species such

as *B. leachii* and *B. schlosseri* (Brunetti, 1974, 1976; Brunetti *et al.*, 1980), but even at the

molecular level within *B. schlosseri* populations (Bock *et al.*, 2012; Nydam *et al.*, 2017). It is

thus possible that the observed success in tunicate invasion (Zhan *et al.*, 2015) is supported

by their plasticity in genome characteristics like transposon diversity and gene number.

Ancient homeobox gene clusters whose structure has been retained over millions of

years of evolution in many organisms are fragmented in tunicate genomes. Because, the

expression of each *Hox* genes across the anterior-posterior axis relates to their location within

the *Hox* gene cluster (Pascual-Anaya *et al.*, 2013), cluster breaks are predicted to have

consequences for patterning processes. However, an adult body plan with correct spatial

orientation of its body axes during tissue development in ascidians also needs to be

established during sexual, asexual and WBR. Early patterning events in tunicate species have

only been characterized during sexual reproduction in *Ciona*. Early stages of development

(prior to gastrulation) follow a mosaic pattern of developmental axis formation, where

inheritance of maternally provided factors establishes the body axes (Nishida, 2005). *Hox*

gene knockdown experiments in *C. intestinalis* revealed that they have very limited roles, with

defects only observed in larval neuronal and tail development upon loss of *Ci-Hox11* and *Ci-*

Hox12 function (Ikuta *et al.*, 2010). It thus appears that patterning events in *C. intestinalis* are

less dependent upon anterior-posterior spatial expression of *Hox* genes to establish regional

identity. Previously, in *B. schlosseri*, the entry point of the connective test vessel into the

developing bud determines the posterior end of the new zooid (Sabbadin *et al.*, 1975).

Therefore it is possible that ascidians incorporate environmental and physical cues to

compensate for the lost gene cluster during polarity establishment. A wider analysis

comprising multiple tunicate species will be necessary to investigate the exact consequences

of homeobox cluster dispersion and whether the compensatory mechanism observed in *C.*

intestinalis is the norm or an exception.

Lineage-specific changes to evolutionarily conserved cell communication pathways

Cell signaling pathways are critical for morphogenesis, development and adult

physiology. In particular, we have focused our analysis on three highly conserved pathways:

Wnt, Notch and Retinoic Acid signaling. Representatives of all twelve *Wnt* genes subfamilies

are found in metazoans, suggesting that they evolved before evolution of the bilaterians

(Janssen *et al.*, 2010). We identified members of each Wnt subfamily in tunicate genomes,

along with numerous examples of lineage-specific gene loss and/or duplication. The most

striking of these events was an increase in *Wnt5a* gene copy number in *B. leachii*, *B. schlosseri*

and *M. oculata* genomes. Indeed, most invertebrates genomes, including the basal chordate *B.*

floridae, contain a single *Wnt5* gene while most vertebrate genomes have two *Wnt5a*

paralogues, believed to be a result of whole genome duplication (Martin *et al.*, 2012).

However, in the analyzed tunicate genomes, up to 15 copies of this gene were identified,

potentially these additional genes may have been co-opted into novel roles and were retained

during tunicate evolution. Wnt5a ligands have numerous biological roles, including a

suppressive one during zebrafish regeneration (Stoick-Cooper *et al.*, 2007) and a promotive

one during amphioxus regeneration (Somorjai *et al.*, 2012). Furthermore, components of both

Wnt signaling pathways are differentially expressed during WBR (Zondag *et al.*, 2016). Altogether, it is thus conceivable that *Wnt5a* gene number has expanded in colonial tunicates to sustain WBR. A functional characterization of the role of these numerous copies of Wnt5a would thus be highly interesting and potentially give evolutionary insights into chordate regeneration.

All components of the Notch pathway are present in the genomes we investigated. Of particular interest, the DSL Notch ligandappears to be rapidly evolving in the tunicates. This indicates that tunicate DSL proteins are under less pressure, than vertebrate orthologous proteins, to conserve their protein sequence. Given that the concentration and activity of the DSL ligands are typically the rate limiting step in this key signaling pathway, it will be interesting to assess whether the functional properties of tunicate proteins have adapted accordingly.

Components of the RA signaling pathway have also been identified in all the tunicate genomes. However, *Oikopleura* has seemingly lost a functional RA synthesis pathway, while still forming a functional body plan. This suggests that either uniquely RA is not involved in critical developmental events in this species, that the RA signaling function has been replaced or that *O. dioica* utilizes an alternative synthesis approach. Conversely, lineage specific increases in RA pathway gene numbers have been observed in *C. intestinalis* (Aldh1, (Sobreira *et al.*, 2011)) and botryllids (*CYP26* genes).

RA, Notch and Wnt pathways play roles in regeneration and development in many species, including Solidobranchian tunicates (Rinkevich, Paz, *et al.*, 2007; Rinkevich *et al.*, 2008; Zondag *et al.*, 2016) and *Cionidae (Hamada et al., 2015; Jeffery, 2015)*. The observed loss of RA signaling genes may result in a poorer regeneration ability for *O. dioica*, however it's regenerative has not been characterized. Given the unique chordate WBR potential developed by colonial tunicates, it is conceivable that there is selective pressure on their genomes to

retain these pathways. In particular given the likely role these pathways also play in colony

reactivation following hibernation, as well as in asexual reproduction.

In addition, among tunicates there are significant differences in both life cycle,

reproduction and regeneration ability, even between closely related species of the same

family, that likely reflect an underlying diversity in genomic content. For instance, differences

in both asexual and sexual reproduction have been observed between within the Botryllidae

family (Berrill, 1941, 1947, 1951; Oka and Watanabe, 1957; Brunetti, 1974, 1976).

Furthermore, *B. schlosseri* can only undergo WBR during a short time frame of their asexual

reproductive cycle when the adults are reabsorbed by the colony (Voskoboynik *et al.*, 2007;

Kürn *et al.*, 2011) while *B. leachii* can undergo WBR throughout their adult life (Rinkevich,

Paz, *et al.*, 2007). Overall, this indicates that despite a generally similar appearance, the rapid

evolution of the Tunicata subphylum has provided diversity and innovations within its

species. It will thus be particularly interesting to investigate in future studies how such

genomic plasticity balances between adaptation to new challenges and constraint, preserving

common morphological features.

In conclusion, our assembly of the *B. leachii* genome provides an essential resource for

the study of this colonial ascidian as well as a crucial point of comparison to gain further

insights into the remarkable genetic diversity among tunicate species. In addition, the

genome of *B. leachii* will be most useful for dissecting WBR in chordates, in particular by

comparing it with that of *B. schlosseri* to understand how the initiation of WBR can be blocked

during specific periods of their life cycle. Furthermore, given the key phylogenetic position of

Tunicates with respect to vertebrates, the analysis of their genomes will provide important

insights in the emergence of chordate traits and the origin of vertebrates.

Methods

Sampling, library preparation and sequencing

   B. leachii colonies were collected from Nelson harbour (latitude 41.26°S, longitude 173.28°E) in New Zealand. To reduce the likelihood of contamination, embryos were dissected out of a colony and pooled before carrying out DNA extraction using E.Z.N.A SP Plant DNA Mini Kit. A total of 2 μg each was sent to New Zealand Genomics Limited (NZGL) for library preparation and sequencing. A short read (TruSeq illumina HiSeq) generated 19090212 paired-end reads of 100 bp (average fragment size: 450bp, adaptor length: 120bp). A second library (Illumina Nextera MiSeq Mate Pair) not size-selected library (fragment size: 1.5-15 kb, median size: ~3 kb, adaptor length: 38bp).generated 31780788 paired-end sequences of 250 bp.

   PreQC report was generated using the String Graph Assembler software package (Simpson, 2014) and quality metrics before assembly with both FastQC (Andrews, 2010) as well as MultiQC (Ewels *et al.*, 2016) (Fig. S1). These analyses revealed that 91 % of sequences had a mean Phred quality score >= 30, 96 % of bases a mean Phred quality score >= 30, and 39 % of sequences an adapter sequence (either Illumina or Nextera). Adaptor trimming was performed with NxTrim (O'Connell *et al.*, 2015) for the mate pair library, followed by Trimmomatic (Bolger *et al.*, 2014) with the following options: MINLEN:40 ILLUMINACLIP:2:30:12:1:true LEADING:3 TRAILING:3 MAXINFO:40:0.4 MINLEN:40 for both libraries. After trimming, 86644308 paired-end (85 %) and 12112004 (12 %) single-end sequences remained (100% with a mean Phred quality score >= 30, <1% with an adapter sequence).

Genome assembly

De novo assembly was performed in three consecutive iterations following a Meta-assembly approach (Table S5). First, both libraries were assembled together in parallel, using a k-mer size of 63 following the results from KmerGenie (Chikhi and Medvedev, 2014) whenever available, by five assemblers: AbySS (Simpson *et al.*, 2009), Velvet (Zerbino and Birney, 2008), SOAPdenovo2 (Luo *et al.*, 2012), ALLPATHS-LG (Gnerre *et al.*, 2011), MaSuRCA (Zimin *et al.*, 2013). The MaSuRCA assembler was run twice, once running the adapter filtering function (here termed "MaSuRCA-filtered"), the other without (termed simply "MaSuRCA"). Their respective quality was then estimated using three different metrics: the N50 length, the BUSCO core-genes completion (Simão *et al.*, 2015) and the Glimmer number of predicted genes (Delcher *et al.*, 1999). Second, these drafts were combined by following each ranking using Metassembler (Wences and Schatz, 2015), hence producing three new assemblies (limiting the maximum insert size at 15 kb). Third, the *B. leachii* transcriptome (Zondag *et al.*, 2016) was aligned to each meta-assembly using STAR (Dobin *et al.*, 2013), which were then combined thrice more using Metassembler following their alignment percentage and limiting the maximum insert size at 3 kb, 8 kb and 15 kb. Finally, the quality of the meta-meta-assemblies was estimated using the BUSCO score and the best one (Table S5) selected as the reference *de novo* assembly.

Data access

B. leachiii: ANISEED http://www.aniseed.cnrs.fr/fgb2/gbrowse/boleac_v3/

C. intestinalis: Ciona_intestinalis.KH.cds.all.fa, Ciona_intestinalis.joinedscaffold.fa Ciona unmasked v2.0, https://www.aniseed.cnrs.fr/fgb2/gbrowse/ciona_intestinalis/

C. savigny: *C. savigni*_paired_scaffolds.fa,

https://www.aniseed.cnrs.fr/fgb2/gbrowse/cisavi_ens81/,

M. oculata: , https://www.aniseed.cnrs.fr/fgb2/gbrowse/moocul_elv12/,

B. schlosseri: botznik-transcripts.fa, botznik-chr.fa, botznik-ctg.fa,

https://www.aniseed.cnrs.fr/fgb2/gbrowse/boschl_botznik2013/,

O. dioica: Odioica_reference_v3.fa, Oikopleura_transcripts_reference_v1.0.fa,

http://www.genoscope.cns.fr/externe/GenomeBrowser/Oikopleura/.

Repeat region analysis

A *de novo* repeat library was build for each tunicate genome using RepeatModeler

(Smit and Hubley, 2015). This utilizes the RECON tandem repeats finder from the RepeatScout

packages to identify species-specific repeats in a genome assembly. RepeatMasker (Smit *et al.*,

2015) was then used to mask those repeats.

Gene annotation

Ab initio genome annotation was performed using MAKER2 (Holt and Yandell, 2011)

with Augustus (Stanke and Waack, 2003) and SNAP (Korf, 2004) for gene prediction. In

addition, we used our previously published transcriptome (Zondag *et al.*, 2016) and a

concatenation of UniProtKB (UniProt Consortium, 2015), *C. intestinalis* and *B. schlosseri*

proteins into a custom proteome as evidence of gene product. Using the predicted genes,

Augustus and SNAP were then trained to the specificity of *B. leachii* genome. A second round

of predictions was then performed, followed by a second round of training. The final

annotation of the genome was obtained after running a third round of predictions, and the

provided trained Augustus and SNAP configurations after a third round of training.

The following databases were used for homology identification: InterProScan (Jones *et al.*,

2014) tRNA (Lowe and Eddy, 1997) snoRNA (Lowe, 1999) Infernal (Nawrocki and Eddy,

2013) Rfam (Nawrocki *et al.*, 2015).

Analysis of specific gene families

Genes and transcripts for each examined genome were identified by a tBLASTn search with an e-value cut-off at 10^{-5} using the SequencerServer software (Priyam *et al.*, 2015). This was followed by a reciprocal BLAST using SMART blast (http://blast.ncbi.nlm.nih.gov/smartblast/smartBlast.cgi?CMD=Web) to confirm their identity.

Delta serrate ligand conserved protein domain (PF01414) was used to identify the corresponding proteins in tunicate genomes. To identify *Notch* receptor genes the conserved LNR (lin-notch repeat) domain (PF00066) was used. ALDH-like genes were identified by tBLASTn search (PF00171) and classified using SMART blast.

Phylogenetics

Sequences were aligned with ClustalX (Jeanmougin *et al.*, 1998) before using ProtTest 3 (Abascal *et al.*, 2005) to determine the best-fit model of evolution. The best-fit model for the DSL phylogeny was WAG+I+G and, for CYP26 proteins, was LG+I+G.

Bayesian inference (BI) phylogenies were constructed by MrBayes (Ronquist and Huelsenbeck, 2003) with a mixed model for 100,000 generations and summarized using a Sump burnin of 200. Maximum Likelihood (ML) phylogenies were generated by PhyML (Guindon *et al.*, 2010), using the appropriate model, estimating the amino acid frequences.

Accession numbers are provided in File S3 and sequence alignments are provided in Figure S3. Analyses carried out with BI and ML produced identical tree topologies. Trees were displayed using FigTree v1.4.2 [REF] (http://tree.bio.ed.ac.uk/software/figtree/).

Acknowledgements

Funding support was provided to M.J.W. by the Otago BMS Deans Bequest and Department of Anatomy. S.B. by the Swiss National Science Foundation (SNSF) [grant number P2ELP3_158873]. We would like to thank Peter Maxwell and the New Zealand eScience Infrastructure (NeSI); Christelle Dantec and ANISEED for help and advice during the annotation process, as well as for the accompanying *B. leachii* genome browser.

 Figure Legends

Figure 1 *B. leachii* **phylogenetic position and life cycle.**

A. Schematic showing phylogeny of tunicates with respect to the chordate clade. **B.** Life cycle of *B. leachii*. The colony expands and grows by asexual reproduction. During favorable conditions such as warmer water temperatures, members of the colonies start sexual reproduction. The embryos remain with the colony in brood pouches until release. Hatched larvae attach to nearby substrates and begin metamorphosis into a zooid.

Figure 2. *Hox* genes are dispersed and reduced in number within tunicate genomes.
Schematic depicting remaining *Hox* gene linkages in five tunicate genomes in comparison to
the ancestral *Hox* complex, which included thirteen genes. Orthologous genes are indicated by
common colours. Chromosome or scaffold number is shown, along with gene ID when
available for newly annotated genomes.

Figure 3. NK homeobox cluster genes are fragmented within tunicate genomes.

Schematic depicting the organization of the *NK homeobox* cluster genes among the studied chordate genomes. Double-parallel lines indicate > 1Mb distance between genes. Chromosome or scaffold number is shown, along with gene ID when available for newly annotated genomes. Orthologous genes are indicated by common colours.

Figure 4. Ancestral gene linkages remain between a few pharyngeal cluster genes in tunicate genomes. Gene order of the six pharyngeal cluster genes, *Nkx2.1, Nk2.2, Pax1/9* and *FoxA* in chordate genomes. Double-parallel lines indicate > 1Mb distance between genes. Chromosome or scaffold number is shown, along with gene ID when available for newly annotated genomes. Orthologous genes are indicated by common colours.

Figure 5. Duplication of *Wnt5a* genes in Tunicate genomes.

Schematic showing the genomic location of *B. leachii* (A), *B. schlosseri* (B), *M. oculata* (C) and *C. intestinalis* (D) *Wnt5*-like genes within each genome. Double-parallel lines indicate > 1Mb distance between genes.

Figure 6. *B. leachii* **Notch pathway**

Bayesian phylogenetic tree depicting the relationship between tunicate and vertebrate DSL

proteins, using *Drosophila* Delta to root the tree. *B. leachii* proteins are shown in red and

shaded areas correspond to clade groupings. Branch support values (probabilities) are

indicated.

Abbreviations: Bl, *Botrylliodes leachii***; BsS,** *Botryllus schlosseri***; Od,** *Oikopleura odioica***;**

Hs, *Homo sapiens***; Ci,** *Ciona intestinalis***, Dm,** *Drosophila melanogaster***; Dr,** *Danio rerio;*

Mo*, M. oculata;* **Hr** *Halocynthia roretzi***;** *Blan, Branchiostoma lanceolatum; Bf,*

Branchiostoma floridae.

Figure 7. **Evolution of the RA pathway in tunicates**

(A) Overview of the RA synthesis and degradation pathway. In bold are the major proteins that contribute to RA signaling during animal development. Indicated below these are changes to the number of copies present in examined genomes. **(B)** ML phylogenetic tree depicting the relationship between invertebrate and vertebrate CYP26 proteins using CYP4 and CYP51 proteins as an outgroup. *B. schlosseri* and *B. leachii* proteins are shown in red, *C. intestinalis* and *M. oculata* are indicated in blue. No *Cyp26* gene has been identifed in the *O. dioica* genome. Values for the approximate likelihood-ratio test (aLRT) are indicated.

Abbreviations: Bl, *Botrylliodes leachii*; **Bs**, *Botryllus schlosseri*; **Hs**, *Homo sapiens*; **Ci**, *Ciona intestinalis;* **Mo**, *M. oculata.*

References

Abascal, F., R. Zardoya and D. Posada. **2005**. ProtTest: selection of best-fit models of protein evolution. *Bioinformatics* **21**: 2104–2105.

Andrews, S. 2010. FastQC: a quality control tool for high throughput sequence data.

Auger, H., Y. Sasakura, J.-S. Joly and W. R. Jeffery. **2010**. Regeneration of oral siphon pigment organs in the ascidian Ciona intestinalis. *Dev. Biol.* **339**: 374–389.

Ballarin, L., A. Franchini, E. Ottaviani and A. Sabbadin. **2001**. Morula cells as the major immunomodulatory hemocytes in ascidians: Evidences from the colonial species *Botryllus schlosseri*. *Biol. Bull.* **201**: 59–64.

Belyaeva, O. V., C. Chang, M. C. Berlett and N. Y. Kedishvili. **2015**. Evolutionary origins of retinoid active short-chain dehydrogenases/reductases of SDR16C family. *Chem. Biol. Interact.* **234**: 135–143.

Belyaeva, O. V., S.-A. Lee, O. V. Kolupaev and N. Y. Kedishvili. **2009**. Identification and characterization of retinoid-active short-chain dehydrogenases/reductases in Drosophila melanogaster. *Biochim. Biophys. Acta - Gen. Subj.* **1790**: 1266–1273.

Berna, L. and F. Alvarez-Valin. **2014**. Evolutionary genomics of fast evolving tunicates. *Genome Biol. Evol.* **6**: 1724–1738.

Berrill, N. J. 1941. The development of the bud in *Botryllus*. *Biol. Bull.* **80**: 169.

Berrill, N. J. 1947. The developmental cycle of *Botrylloides*. *Q. J. Microsc. Sci.* **88**: 393–407.

Berrill, N. J. 1951. Regeneration and Budding in Tunicates. *Biol. Rev.* **26**: 456–475.

Bock, D. G., H. J. MacIsaac and M. E. Cristescu. **2012**. Multilocus genetic analyses differentiate between widespread and spatially restricted cryptic species in a model ascidian. *Proc. R. Soc. B Biol. Sci.* **279**: 2377–2385.

Bolger, A. M., M. Lohse and B. Usadel. **2014**. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.

Bradnam, K. R., J. N. Fass, A. Alexandrov, P. Baranay, M. Bechner, I. Birol, S. Boisvert, J. A. Chapman, G. Chapuis, R. Chikhi, *et al.* 2013. Assemblathon 2: evaluating de novo methods of genome assembly in three vertebrate species. *Gigascience* **2**: 10.

Brown, F. D. and B. J. Swalla. **2012**. Evolution and development of budding by stem cells: Ascidian coloniality as a case study. *Dev. Biol.* **369**: 151–162. Elsevier.

Brozovic, M., C. Martin, C. Dantec, D. Dauga, M. Mendez, P. Simion, M. Percher, B. Laporte, C. Scornavacca, A. Di Gregorio, *et al.* 2016. ANISEED 2015: A digital framework for the comparative developmental biology of ascidians. *Nucleic Acids Res.* **44**: D808–D818.

Brunetti, R. 1974. *Observations on the Life Cycle of Botryllus Schlosseri (Pallas) (Ascidiacea) in*
the Venetian Lagoon.

Brunetti, R. 1976. Biological cycle of *Botrylloides leachi* (Savigny) (Ascidiacea) in the Venetian
lagoon. *Vie Milieu* **XXVI**: 105–122.

Brunetti, R., L. Beghi, M. Bressan and M. Marin. **1980**. Combined effects of temperature and
salinity on colonies of *Botryllus schlosseri* and *Botrylloides leachi* (Ascidiacea) from the
Venetian Lagoon. *Mar. Ecol. Prog. Ser.* **2**: 303–314.

Burighel, P., R. Brunetti and G. Zaniolo. **1976**. Hibernation of the colonial ascidian *Botrylloides*
leachi (Savigny): histological observations. *Ital. J. Zool.* **43**: 293–301.

Camacho, C., G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer and T. L.
Madden. **2009**. BLAST+: architecture and applications. *BMC Bioinformatics* **10**: 421.

Canestro, C., J. H. Postlethwait, R. Gonzalez-Duarte and R. Albalat. **2006**. Is retinoic acid
genetic machinery a chordate innovation? *Evol. Dev.* **8**: 394–406.

Chikhi, R. and P. Medvedev. **2014**. Informed and automated k-mer size selection for genome
assembly. *Bioinformatics* **30**: 31–37.

Cunningham, T. J. and G. Duester. **2015**. Mechanisms of retinoic acid signalling and its roles in
organ and limb development. *Nat. Rev. Mol. Cell Biol.* **16**: 110–123.

Dehal, P., Y. Satou, R. K. Campbell, J. Chapman, B. Degnan, A. De Tomaso, B. Davidson, A.
Di Gregorio, M. Gelpke, D. M. Goodstein, *et al.* 2002a. The Draft Genome of Ciona
intestinalis: Insights into Chordate and Vertebrate Origins. *Science (80-.).* **298**: 2157–2167.

Dehal, P., Y. Satou, R. K. Campbell, J. Chapman, B. Degnan, A. W. De Tomaso, B. Davidson,
A. Di Gregorio, M. Gelpke, D. M. Goodstein, *et al.* 2002b. The draft genome of *Ciona*
intestinalis: insights into chordate and vertebrate origins. *Science* **298**: 2157–67.

Delcher, A. L., D. Harmon, S. Kasif, O. White and S. L. Salzberg. **1999**. Improved microbial
gene identification with GLIMMER. *Nucleic Acids Res.* **27**: 4636–41.

Delsuc, F., H. Brinkmann, D. Chourrout and H. Philippe. **2006**. Tunicates and not
cephalochordates are the closest living relatives of vertebrates. *Nature* **439**: 965–968.

Dobin, A., C. a Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson
and T. R. Gingeras. **2013**. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**:
15–21.

Edvardsen, R. B., H.-C. Seo, M. F. Jensen, A. Mialon, J. Mikhaleva, M. Bjordal, J. Cartry, R.
Reinhardt, J. Weissenbach, P. Wincker, *et al.* 2005. Remodelling of the homeobox gene
complement in the tunicate Oikopleura dioica. *Curr. Biol.* **15**: R12–R13.

Ewels, P., M. Magnusson, S. Lundin and M. Käller. **2016**. MultiQC: Summarize analysis results
for multiple tools and samples in a single report. *Bioinformatics* btw354.

Franchi, N., F. Schiavon, M. Carletto, F. Gasparini, G. Bertoloni, S. C. E. Tosatto and L. Ballarin. **2011**. Immune roles of a rhamnose-binding lectin in the colonial ascidian Botryllus schlosseri. *Immunobiology* **216**: 725–736.

Garcia-Fernàndez, J. 2005. The genesis and evolution of homeobox gene clusters. *Nat. Rev. Genet.* **6**: 881–92.

Gasparini, F., P. Burighel, L. Manni and G. Zaniolo. **2008**. Vascular regeneration and angiogenic-like sprouting mechanism in a compound ascidian is similar to vertebrates. *Evol. Dev.* **10**: 591–605.

Gazave, E., P. Lapébie, G. S. Richards, F. Brunet, A. V Ereskovsky, B. M. Degnan, C. Borchiellini, M. Vervoort and E. Renard. **2009**. Origin and evolution of the Notch signalling pathway: an overview from eukaryotic genomes. *BMC Evol. Biol.* **9**: 249.

Gnerre, S., I. MacCallum, D. Przybylski, F. J. Ribeiro, J. N. Burton, B. J. Walker, T. Sharpe, G. Hall, T. P. Shea, S. Sykes, *et al.* 2011. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc. Natl. Acad. Sci.* **108**: 1513–1518.

Griggio, F., A. Voskoboynik, F. Iannelli, F. Justy, M.-K. M.-K. Tilak, T. Xavier, G. Pesole, E. J. P. Douzery, F. Mastrototaro and C. Gissi. **2014**. Ascidian Mitogenomics: Comparison of Evolutionary Rates in Closely Related Taxa Provides Evidence of Ongoing Speciation Events. *Genome Biol. Evol.* **6**: 591–605.

Guder, C., I. Philipp, T. Lengfeld, H. Watanabe, B. Hobmayer and T. W. Holstein. **2006**. The Wnt code: cnidarians signal the way. *Oncogene* **25**: 7450–7460.

Guindon, S., J. F. Dufayard, V. Lefort, M. Anisimova, W. Hordijk and O. Gascuel. **2010**. New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. *Syst. Biol.* **59**: 307–321.

Guruharsha, K. G., M. W. Kankel and S. Artavanis-Tsakonas. **2012**. The Notch signalling system: recent insights into the complexity of a conserved pathway. *Nat. Rev. Genet.* **13**: 654–666.

Hamada, M., S. Goricki, M. S. Byerly, N. Satoh and W. R. Jeffery. **2015**. Evolution of the chordate regeneration blastema: Differential gene expression and conserved role of notch signaling during siphon regeneration in the ascidian Ciona. *Dev. Biol.* **405**: 304–315.

Hino, K., Y. Satou, K. Yagi and N. Satoh. **2003**. A genomewide survey of developmentally relevant genes in Ciona intestinalis. *Dev. Genes Evol.* **213**: 264–272.

Hoegg, S. and A. Meyer. **2005**. Hox clusters as models for vertebrate genome evolution. *Trends Genet.* **21**: 421–424.

Holt, C. and M. Yandell. **2011**. MAKER2: an annotation pipeline and genome-database management tool for second-generation genome projects. *BMC Bioinformatics* **12**: 491.

Ikuta, T., N. Satoh and H. Saiga. **2010**. Limited functions of Hox genes in the larval development of the ascidian Ciona intestinalis. *Development* **137**: 1505–1513.

Jackson, B., C. Brocker, D. C. Thompson, W. Black, K. Vasiliou, D. W. Nebert and V. Vasiliou. **2011**. Update on the aldehyde dehydrogenase gene (ALDH) superfamily. *Hum. Genomics* **5**: 283–303.

Janssen, R., M. Le Gouar, M. Pechmann, F. Poulin, R. Bolognesi, E. E. Schwager, C. Hopfen, J. K. Colbourne, G. E. Budd, S. J. Brown, *et al.* **2010**. Conservation, loss, and redeployment of Wnt ligands in protostomes: implications for understanding the evolution of segment formation. *BMC Evol. Biol.* **10**: 374.

Jeanmougin, F., J. D. Thompson, M. Gouy, D. G. Higgins and T. J. Gibson. **1998**. Multiple sequence alignment with Clustal X. *Trends Biochem. Sci.* **23**: 403–5.

Jeffery, W. R. **2015**. Regeneration, Stem Cells, and Aging in the Tunicate Ciona: Insights from the Oral Siphon. *Int. Rev. Cell Mol. Biol.* **319**: 255–82.

Jones, P., D. Binns, H. Y. Chang, M. Fraser, W. Li, C. McAnulla, H. McWilliam, J. Maslen, A. Mitchell, G. Nuka, *et al.* **2014**. InterProScan 5: Genome-scale protein function classification. *Bioinformatics* **30**: 1236–1240.

Kent, W. J. **2002**. BLAT--the BLAST-like alignment tool. *Genome Res.* **12**: 656–64.

Korf, I. **2004**. Gene finding in novel genomes. *BMC Bioinformatics* **5**: 59.

Kürn, U., S. Rendulic, S. Tiozzo and R. J. Lauzon. **2011**. Asexual propagation and regeneration in colonial ascidians. *Biol. Bull.* **221**: 43–61.

Kusserow, A., K. Pang, C. Sturm, M. Hrouda, J. Lentfer, H. A. Schmidt, U. Technau, A. von Haeseler, B. Hobmayer, M. Q. Martindale, *et al.* **2005**. Unexpected complexity of the Wnt gene family in a sea anemone. *Nature* **433**: 156–160.

Lauzon, R. J., C. Brown, L. Kerr and S. Tiozzo. **2013**. Phagocyte dynamics in a highly regenerative urochordate: insights into development and host defense. *Dev. Biol.* **374**: 357–73. Elsevier.

Lee, S.-A., O. V. Belyaeva and N. Y. Kedishvili. **2009**. Biochemical characterization of human epidermal retinol dehydrogenase 2. *Chem. Biol. Interact.* **178**: 182–7.

Lemaire, P., W. C. Smith and H. Nishida. **2008**. Ascidians and the plasticity of the chordate developmental program. *Curr. Biol.* **18**: R620-31.

Loh, K. M., R. van Amerongen and R. Nusse. **2016**. Generating Cellular Diversity and Spatial Form: Wnt Signaling and the Evolution of Multicellular Animals. *Dev. Cell* **38**: 643–655.

Lowe, T. M. **1999**. A Computational Screen for Methylation Guide snoRNAs in Yeast. *Science (80-.).* **283**: 1168–1171.

Lowe, T. M. and S. R. Eddy. **1997**. tRNAscan-SE: A program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* **25**: 955–964.

Luke, G. N., L. F. C. Castro, K. McLay, C. Bird, A. Coulson and P. W. H. Holland. **2003**.
Dispersal of NK homeobox gene clusters in amphioxus and humans. *Proc. Natl. Acad. Sci.*
100: 5292–5295.

Luo, R., B. Liu, Y. Xie, Z. Li, W. Huang, J. Yuan, G. He, Y. Chen, Q. Pan, Y. Liu, *et al.* 2012.
SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler.
Gigascience **1**: 18.

MacDonald, B. T., K. Tamai and X. He. **2009**. Wnt/β-Catenin Signaling: Components,
Mechanisms, and Diseases. *Dev. Cell* **17**: 9–26.

Manni, L., G. Zaniolo, F. Cima, P. Burighel and L. Ballarin. **2007**. *Botryllus schlosseri*: a model
ascidian for the study of asexual reproduction. *Dev. Dyn.* **236**: 335–52.

Martí-Solans, J., O. V. Belyaeva, N. P. Torres-Aguila, N. Y. Kedishvili, R. Albalat and C.
Cañestro. **2016**. Coelimination and Survival in Gene Network Evolution: Dismantling the
RA-Signaling in a Chordate. *Mol. Biol. Evol.* **33**: 2401–2416.

Martin, A., S. Maher, K. Summerhurst, D. Davidson and P. Murphy. **2012**. Differential
deployment of paralogous Wnt genes in the mouse and chick embryo during development.
Evol. Dev. **14**: 178–195.

Maumus, F. and H. Quesneville. **2014**. Ancestral repeats have shaped epigenome and genome
composition for millions of years in Arabidopsis thaliana. *Nat. Commun.* **5**.

Millar, R. H. 1971. The biology of ascidians. *Adv. Mar. Biol.* **9**: 1–100.

Nawrocki, E. P., S. W. Burge, A. Bateman, J. Daub, R. Y. Eberhardt, S. R. Eddy, E. W.
Floden, P. P. Gardner, T. A. Jones, J. Tate, *et al.* 2015. Rfam 12.0: Updates to the RNA
families database. *Nucleic Acids Res.* **43**: D130–D137.

Nawrocki, E. P. and S. R. Eddy. **2013**. Infernal 1.1: 100-fold faster RNA homology searches.
Bioinformatics **29**: 2933–2935.

Nishida, H. 2005. Specification of embryonic axis and mosaic development in ascidians. *Dev. Dyn.*
233: 1177–1193.

Nydam, M. L., K. B. Giesbrecht and E. E. Stephenson. **2017**. Origin and Dispersal History of
Two Colonial Ascidian Clades in the Botryllus schlosseri Species Complex. *PLoS One* **12**:
e0169944.

O'Connell, J., O. Schulz-Trieglaff, E. Carlson, M. M. Hims, N. a. Gormley and a. J. Cox.
2015. NxTrim: optimized trimming of Illumina mate pair reads. *Bioinformatics* **31**: btv057.

O'Neill, R. J., M. J. O'Neill and J. A. Graves. **1998**. Undermethylation associated with
retroelement activation and chromosome remodelling in an interspecific mammalian hybrid.
Nature **393**: 68–72.

Oka, H. and H. Watanabe. **1957**. Vascular budding, a new type of budding in *Botryllus*. *Biol.*
Bull. **112**: 225.

Pascual-Anaya, J., S. D'Aniello, S. Kuratani and J. Garcia-Fernàndez. **2013**. Evolution of Hox
gene clusters in deuterostomes. *BMC Dev. Biol.* **13**: 26.

Pearson, J. C., D. Lemons and W. McGinnis. **2005**. Modulating Hox gene functions during
animal body patterning. *Nat. Rev. Genet.* **6**: 893–904.

Piette, J. and P. Lemaire. **2015**. Thaliaceans, The Neglected Pelagic Relatives of Ascidians: A
Developmental and Evolutionary Enigma. *Q. Rev. Biol.* **90**: 117–145.

Priyam, A., B. J. Woodcroft, V. Rai, A. Munagala, I. Moghul, F. Ter, M. A. Gibbins, H.
Moon, G. Leonard, W. Rumpf, *et al.* **2015**. Sequenceserver: a modern graphical user
interface for custom BLAST databases. *bioRxiv*.

Prud'homme, B., N. Lartillot, G. Balavoine, A. Adoutte and M. Vervoort. **2002**. Phylogenetic
analysis of the Wnt gene family. Insights from lophotrochozoan members. *Curr. Biol.* **12**:
1395.

Rinkevich, B., Z. Shlemberg and L. Fishelson. **1995**. Whole-body protochordate regeneration
from totipotent blood cells. *Proc. Natl. Acad. Sci. U. S. A.* **92**: 7695–9.

Rinkevich, Y., J. Douek, O. Haber, B. Rinkevich and R. Reshef. **2007**. Urochordate whole body
regeneration inaugurates a diverse innate immune signaling profile. *Dev. Biol.* **312**: 131–46.

Rinkevich, Y., G. Paz, B. Rinkevich and R. Reshef. **2007**. Systemic bud induction and retinoic
acid signaling underlie whole body regeneration in the urochordate *Botrylloides leachi. PLoS*
Biol. **5**: e71.

Rinkevich, Y., B. Rinkevich and R. Reshef. **2008**. Cell signaling and transcription factor genes
expressed during whole body regeneration in a colonial chordate. *BMC Dev. Biol.* **8**: 100.

Rinkevich, Y., A. Voskoboynik, A. Rosner, C. Rabinowitz, G. Paz, M. Oren, J. Douek, G.
Alfassi, E. Moiseeva, K. J. Ishizuka, *et al.* **2013**. Repeated, long-term cycling of putative
stem cells between niches in a basal chordate. *Dev. Cell* **24**: 76–88. Elsevier Inc.

Ronquist, F. and J. P. Huelsenbeck. **2003**. MrBayes 3: Bayesian phylogenetic inference under
mixed models. *Bioinformatics* **19**: 1572–4.

Ross, A. C. and R. Zolfaghari. **2011**. Cytochrome P450s in the regulation of cellular retinoic acid
metabolism. *Annu. Rev. Nutr.* **31**: 65–87.

Rubinstein, N. D., T. Feldstein, N. Shenkar, F. Botero-Castro, F. Griggio, F. Mastrototaro, F.
Delsuc, E. J. P. Douzery, C. Gissi and D. Huchon. **2013**. Deep sequencing of mixed total
DNA without barcodes allows efficient assembly of highly plastic Ascidian mitochondrial
genomes. *Genome Biol. Evol.* **5**: 1185–1199.

Sabbadin, A., G. Zaniolo and F. Majone. **1975**. Determination of polarity and bilateral
asymmetry in palleal and vascular buds of the ascidian Botryllus schlosseri. *Dev. Biol.* **46**: 79–
87.

Saito, Y., M. Shirae, M. Okuyama and S. Cohen. **2001**. Phylogeny of Botryllid Ascidians. In:
The Biology of Ascidians, pp. 315–320. Springer Japan, Tokyo.

Santagati, F., K. Abe, V. Schmidt, T. Schmitt-John, M. Suzuki, K.-I. Yamamura and K. Imai.
2003. Identification of Cis-regulatory elements in the mouse Pax9/Nkx2-9 genomic region:
implication for evolutionary conserved synteny. *Genetics* **165**: 235–42.

Savigny, J.-C. 1816. *Mémoires sur les animaux sans vertèbres*. Dufour, G., Paris.

Seo, H.-C. C., M. Kube, R. B. Edvardsen, M. F. Jensen, A. Beck, E. Spriet, G. Gorsky, E. M.
Thompson, H. Lehrach, R. Reinhardt, *et al.* 2001. Miniature genome in the marine chordate
Oikopleura dioica. *Science* **294**: 2506.

Simakov, O., T. Kawashima, F. Marlétaz, J. Jenkins, R. Koyanagi, T. Mitros, K. Hisata, J.
Bredeson, E. Shoguchi, F. Gyoja, *et al.* 2015. Hemichordate genomes and deuterostome
origins. *Nature* **527**: 459–465.

Simão, F. A., R. M. Waterhouse, P. Ioannidis and E. V Kriventseva. **2015**. BUSCO : assessing
genome assembly and annotation completeness with single-copy orthologs. *Genome Anal.* 9–
10.

Simmen, M. W., S. Leitgeb, J. Charlton, S. J. Jones, B. R. Harris, V. H. Clark and A. Bird.
1999. Nonmethylated transposable elements and methylated genes in a chordate genome.
Science **283**: 1164–7.

Simpson, J. T. 2014. Exploring genome characteristics and sequence quality without a reference.
Bioinformatics **30**: 1228–1235.

Simpson, J. T., K. Wong, S. D. Jackman, J. E. Schein, S. J. M. Jones and I. Birol. **2009**.
ABySS: A parallel assembler for short read sequence data. *Genome Res.* **19**: 1117–1123.

Small, K. S., M. Brudno, M. M. Hill and A. Sidow. **2007**. A haplome alignment and reference
sequence of the highly polymorphic Ciona savignyi genome. *Genome Biol.* **8**: R41.

Smit, A. and R. Hubley. **2015**. RepeatModeler Open-1.0.

Smit, A., R. Hubley and P. Green. **2015**. RepeatMasker Open-4.0.

Sobreira, T. J. P., F. Marletaz, M. Simoes-Costa, D. Schechtman, A. C. Pereira, F. Brunet, S.
Sweeney, A. Pani, J. Aronowicz, C. J. Lowe, *et al.* 2011. Structural shifts of aldehyde
dehydrogenase enzymes were instrumental for the early evolution of retinoid-dependent axial
patterning in metazoans. *Proc. Natl. Acad. Sci.* **108**: 226–231.

Somorjai, I. M. L., H. Escrivà and J. Garcia-Fernàndez. **2012**. Amphioxus makes the cut—
Again. *Commun. Integr. Biol.* **5**: 499–502.

Spagnuolo, A., F. Ristoratore, A. Di Gregorio, F. Aniello, M. Branno and R. Di Lauro. **2003**.
Unusual number and genomic organization of Hox genes in the tunicate Ciona intestinalis.
Gene **309**: 71–9.

Stanke, M. and S. Waack. **2003**. Gene prediction with a hidden Markov model and a new intron
submodel. *Bioinformatics* **19**: 215–225.

Stapley, J., A. W. Santure and S. R. Dennis. **2015**. Transposable elements as agents of rapid
adaptation may explain the genetic paradox of invasive species. *Mol. Ecol.* **24**: 2241–2252.

Stoick-Cooper, C. L., G. Weidinger, K. J. Riehle, C. Hubbert, M. B. Major, N. Fausto and R.
T. Moon. **2007**. Distinct Wnt signaling pathways have opposing roles in appendage
regeneration. *Development* **134**: 479–89.

Stolfi, A., E. K. Lowe, C. Racioppi, F. Ristoratore, C. T. Brown, B. J. Swalla and L.
Christiaen. **2014**. Divergent mechanisms regulate conserved cardiopharyngeal development
and gene expression in distantly related ascidians. *Elife* **3**: e03728.

Suzuki, M. M., A. R. W. Kerr, D. De Sousa and A. Bird. **2007**. CpG methylation is targeted to
transcription units in an invertebrate genome. *Genome Res.* **17**: 625–631.

Takatori, N., T. Butts, S. Candiani, M. Pestarino, D. E. K. Ferrier, H. Saiga and P. W. H.
Holland. **2008**. Comprehensive survey and classification of homeobox genes in the genome of
amphioxus, Branchiostoma floridae. *Dev. Genes Evol.* **218**: 579–590.

Tsagkogeorga, G., V. Cahais and N. Galtier. **2012**. The Population Genomics of a Fast Evolver:
High Levels of Diversity, Functional Constraint, and Molecular Adaptation in the Tunicate
Ciona intestinalis. *Genome Biol. Evol.* **4**: 852–861.

Tsagkogeorga, G., X. Turon, N. Galtier, E. J. P. Douzery and F. Delsuc. **2010**. Accelerated
Evolutionary Rate of Housekeeping Genes in Tunicates. *J. Mol. Evol.* **71**: 153–167.

UniProt Consortium. **2015**. UniProt: a hub for protein information. *Nucleic Acids Res.* **43**: D204–
D212.

Voskoboynik, A., N. F. Neff, D. Sahoo, A. M. Newman, D. Pushkarev, W. Koh, B. Passarelli,
H. C. Fan, G. L. Mantalas, K. J. Palmeri, *et al.* **2013**. The genome sequence of the colonial
chordate, *Botryllus schlosseri*. *Elife* **2**: 1–24.

Voskoboynik, A., N. Simon-Blecher, Y. Soen, B. Rinkevich, A. W. De Tomaso, K. J. Ishizuka
and I. L. Weissman. **2007**. Striving for normality: whole body regeneration through a series
of abnormal generations. *FASEB J.* **21**: 1335–44.

Wada, S., M. Tokuoka, E. Shoguchi, K. Kobayashi, A. Di Gregorio, A. Spagnuolo, M. Branno,
Y. Kohara, D. Rokhsar, M. Levine, *et al.* **2003**. A genomewide survey of developmentally
relevant genes in Ciona intestinalis. *Dev. Genes Evol.* **213**: 222–234.

Wang, X.-P., M. Suomalainen, S. Felszeghy, L. C. Zelarayan, M. T. Alonso, M. V Plikus, R. L.
Maas, C.-M. Chuong, T. Schimmang and I. Thesleff. **2007**. An integrated gene regulatory
network controls stem cell proliferation in teeth. *PLoS Biol.* **5**: e159.

Wences, A. H. and M. C. Schatz. **2015**. Metassembler: merging and optimizing de novo genome
assemblies. *Genome Biol.* **16**: 207. Genome Biology.

Zerbino, D. R. and E. Birney. **2008**. Velvet: algorithms for de novo short read assembly using de
Bruijn graphs. *Genome Res.* **18**: 821–9.

Zhan, A., E. Briski, D. G. Bock, S. Ghabooli and H. J. MacIsaac. **2015**. Ascidians as models for
studying invasion success. *Mar. Biol.* **162**.

Zhang, G., X. Fang, X. Guo, L. Li, R. Luo, F. Xu, P. Yang, L. Zhang, X. Wang, H. Qi, *et al.*
2012. The oyster genome reveals stress adaptation and complexity of shell formation. *Nature*
490: 49–54.

Zimin, A. V., G. Marcais, D. Puiu, M. Roberts, S. L. Salzberg and J. A. Yorke. **2013**. The
MaSuRCA genome assembler. *Bioinformatics* **29**: 2669–2677.

Zondag, L. E., K. Rutherford, N. J. Gemmell and M. J. Wilson. **2016**. Uncovering the pathways
underlying whole body regeneration in a chordate model, *Botrylloides leachi* using de novo
transcriptome analysis. *BMC Genomics* **17**: 114. BMC Genomics.