Lack of dynamic balance between vesicle transport and microtubule growth in neurite outgrowth
enables formation of dystrophic bulbs

Arjun Singh Yadaw, Mustafa M. Siddiq, Vera Rabinovich, Rosa Tolentino, Ravi Iyengar* and
Jens Hansen*

Department of Pharmacological Sciences and Systems Biology Center New York, Icahn School
of Medicine at Mount Sinai, New York, NY

*- joint senior authors
Jens.Hansen@mssm.edu or Ravi.Iyengar@mssm.edu

Address editorial correspondence to
Ravi Iyengar
Dept. of Pharmacological Sciences, Box 1215
1425 Madison Rm 12-70
New York NY 10029
Ravi.iyengar@mssm.edu

Abstract

Whole cell responses involve multiple subcellular processes (SCPs). To understand how balance
between SCPs controls the dynamics of whole cell responses we studied neurite outgrowth in rat
primary cortical neurons in culture. We used a combination of dynamical models and experiments
to understand the conditions that permitted growth at a specified velocity and when aberrant
growth could lead to the formation of dystrophic bulbs. We hypothesized that dystrophic bulb
formation is due to quantitative imbalances between SCPs. Simulations predict redundancies
between lower level sibling SCPs within each type of high level SCP. In contrast, higher level
SCPs, such as vesicle transport and exocytosis or microtubule growth characteristic of each type
need to be strictly coordinated with each other and imbalances result in stalling of neurite
outgrowth. From these simulations, we predicted the effect of changing the activities of SCPs
involved in vesicle exocytosis or microtubule growth could lead to formation of dystrophic bulbs.
siRNA ablation experiments verified these predictions. We conclude that whole cell dynamics
requires balance between the higher-level SCPs involved and imbalances can terminate whole cell
responses such as neurite outgrowth.

Keywords: Dystrophic bulbs, neurite outgrowth, Vesicles transport, microtubule dynamics,
neurite outgrowth, multi compartment model, ordinary differential equations

Introduction

Neurite outgrowth is an early event that changes the state of neurons (Arimura and Kaibuchi, 2007) and allows neurons to develop axons and dendritic trees that connect to other neurons and become electrically active. When there is nerve injury and the axons are severed, the process of regeneration which is similar to neurite outgrowth often fails, resulting in the formation of bulbs at the ends of regenerating axons. These bulbs are called dystrophic bulbs (Hill, 2017). Although some of the cellular pathways involved in axonal regeneration, failure and formation of the dystrophic bulbs are known (Blanquie and Bradke, 2018; Filious and Schwab, 2018) the mechanisms by which these lead to failure are not well understood. In this study we have used an integrated computational and experimental approach to understand the origins of dystrophic bulbs and more generally the conditions under which whole cell responses such as neurite outgrowth can be maintained. The central question we asked in this study is whether quantitative imbalances between subcellular processes can lead to the formation of dystrophic bulbs.

Whole cell responses that involve, both morphological and physiological changes, are complex because they can engage many subcellular processes (SCPs) that are together responsible for whole cell functions. SCPs include biochemical pathways and cell biological processes involved the functions of different organelles. Coordinated changes in the activities of SCPs often lead to a change in cell state, such as moving to a more differentiated phenotype. Detailed understanding of how different SCPs collaborate to produce an integrated whole cell response requires experiments, along with mathematical models that capture the dynamics of individual SCPs as well as the interrelationship between the dynamics of multiple SCPs. Based on the biological detail that is represented by the SCPs they can be classified as Level-1, Level-2 and Level-3 SCPs. Higher level SCPs (i.e. SCPs with smaller numbers) represent a cell biological function that combines the functions of their lower level SCP children (Hansen et al., 2017). To understand whole cell responses, it is necessary to delineate the quantitative coordination between the dynamics of the SCPs. In the case of neurite outgrowth, three Level-1 SCPs, *Production of Membrane Components, Vesicle Transport & Exocytosis* and *Microtubule Growth* need to be coordinated. Membrane components such as membrane lipids and proteins are synthesized in the cell body and transported to the growth cone via vesicle transport. Vesicles that fuse with the growth cone plasma membrane deliver their membrane lipids to the neurite tip, causing the neurite shaft to grow. Microtubules form the basis of the neurite scaffold that needs to extent in coordination with the neurite shaft. Tsaneva-Atanasova et al (Tsaneva-Atanasova et al., 2009) developed a model of the interaction between microtubules and membrane vesicles in regulating neurite outgrowth. The framework of this model serves as a useful starting point for more extensive models where we can consider how a group of distinct but related SCPs such as *Vesicle Budding, Vesicle Transport* and *Vesicle Fusion* along with a second group of SCPs that involves *Microtubule Nucleation, Microtubule Growth* and *Conversion of Dynamic to Stable Microtubules* interact to enable the growth of neurites. Such detailed models could serve as basis for understanding how a single network can regulate the relationship between kinetic parameters and changes in the levels or activities of molecular

components within the different SCPs. To develop a detailed model of neurite outgrowth based on interactions between SCPs of different levels, we constructed a multi compartment ordinary differential equation model to extend and integrate the microtubule growth model developed by Margolin et al. (Margolin et al., 2012) with the vesicle transport model by Heinrich and Rapoport (Heinrich and Rapoport, 2005). We have also incorporated more recent experimental data to make the model contemporary.

Many mathematical models of neurite outgrowth have been developed (Kiddie et al., 2005; Wissner-Gross et al., 2011) and these models have provided valuable insights into how complex biological processes can be modeled at various levels of description. Often these models abstract the details of the underlying mechanisms and consequently it is difficult to decipher how the balance between the various SCPs is achieved to generate whole cell responses. Some studies have focused on specific facets of the neurite outgrowth process such the role of signaling and regulatory motifs in determining how particular neurites are selected to become axons (Fivaz et al., 2008). Such models are useful in understanding the regulation of complex cellular responses. In developing our dynamical model of neurite outgrowth, we have made a critical change from our previous approaches to dynamical modeling in developing this model. We had typically used a bottom-up approach where individual reactions are assembled into modules that have emergent behaviors (Azeloglu et al., 2014; Bhalla and Iyengar, 1999). In contrast, the approach we use here can be called top-down as we made a starting postulate that the three Level-1 SCPs involved in neurite outgrowth need to be appropriately coordinated to produce a steady velocity of neurite outgrowth. We start with experimental observations of neurite outgrowth velocities and use numerical simulations to determine how their Level-2 children and Level-3 grandchildren SCPs need to be coordinated to allow neurite outgrowth with the observed velocities and under kinetic parameter constellations that match additional experimental data. Our simulations show redundancy between Level-3 sibling SCP activities. Changes in function of one Level-3 sibling SCP can be compensated by alteration of function in another. However, such generalized reciprocal relationships do not exist between Level-2 SCPs. Change in activity in one Level-2 SCP cannot be simply compensated by activity adaptation of one of its sibling Level-2 SCPs but requires adaptations of multiple sibling SCPs. Consequently, loss of coordinated whole cell function is more likely. We used the findings from analyses of the computational models to develop predictions regarding effects of modulating higher SCPs on whole cell function. We tested the predictions experimentally and found them to be valid.

Results

Model Description: Integration of subcellular processes involved in vesicle production, transport and exocytosis and microtubule dynamics to model neurite outgrowth

Neurite outgrowth and axonal regeneration share common SCPs such as vesicle production &
transport and microtubule   growth in an orderly fashion leading to a growing shaft and a growth
cone at the tip as shown schematically in Figure 1A left panel. Failure of growth and the subsequent
formation of dystrophic bulb (Figure 1A right panel) can be hypothesized as failure of SCPs to
function "properly" with respect to each other. To understand how this lack of coordination
between SCPs occurs it is necessary to have a detailed description of the SCPs and their
relationships as part of setting up the computational model we provide a detailed description of
SCPs and their relationships. Among the major SCPs that are involved in neurite outgrowth (NOG)
are the Level-1 SCPs *Production of Membrane Components*, *Vesicle Transport and Exocytosis*
and *Microtubule Growth* (Figure 1B). Each of these SCPs is composed of multiple Level-2
children SCPs whose activities need to be coordinated to ensure NOG without complication. Since
our major focus was on vesicle transport and microtubule growth, we further categorized Level-2
SCP function by introducing their Level-3 children SCP. SCPs with the same parent SCPs are
defined to be siblings. This limited scope enabled us to develop detailed, yet manageable
computational models. It is important to note that at this stage our model does not consider local
actin dynamics at the growth cone, since we particularly focus on SCPs that are involved in
extension of the neurite shaft; i.e. membrane lipid and protein synthesis and delivery and
microtubule scaffold growth.

The level-1 SCP *Production of Membrane Components* has two Level-2 children SCPs, *Membrane*
Lipid Production and *Membrane Protein Production*, that are incorporated into our model via
constant synthesis rates of the individual components at the Trans-Golgi Network (TGN) (Figure
1C; Reaction 1 in Table S1) (Gracias et al., 2014; Wang et al., 2011).

The Level-2 children SCPs of the Level-1 SCP *Vesicle Transport & Exocytosis* are *Vesicle*
Budding at Trans-Golgi Network and *Vesicle Endocytosis at Growth Cone Plasma Membrane*
(GC-PM), *Anterograde* and *Retrograde Microtubule-based Vesicle Transport, Vesicle Fusion with*
Trans-Golgi Network and *Vesicle Exocytosis at Growth Cone Plasma Membrane*. Each of these
six Level-2 SCPs has two Level-3 children SCPs. To simulate the dynamics of the vesicle transport
SCPs we extend a dynamical model of bidirectional membrane lipid and protein transport between
the endoplasmic reticulum and the cis-Golgi (Heinrich and Rapoport, 2005). In our model the two
organelles are the TGN and the GC-PM (Figure 1C). The level-2 SCP *Vesicle Budding at the TGN*
consists of the Level-2 SCPs *Coat Protein Recruitment and Formation at the TGN* and *Vesicle*
Invagination and Scission at TGN. The former is mediated by Recruitment Factor 1 that resides at
the TGN, the latter is mediated by a cell body specific budding rate (Reaction 2 in Table S1).
Membrane proteins at the TGN are recruited into the budding vesicle via association with the coat
proteins (Table S2). The Level-2 SCP *Anterograde Microtubule-Based Vesicle Transport &*
Exocytosis that moves vesicles from the cell body cytoplasm into the neurite shaft cytoplasm and
from there into the growth cone cytoplasm involves the Level-3 SCPs *Kinesin Recruitment to*
Vesicle and *Kinesin Mediated Vesicle Transport along Microtubule*. Kinesin motor proteins are
recruited to the vesicle via kinesin receptors that are part of the vesicle membrane. Here, we assume

that kinesin is available under saturating conditions, so that the kinesin concentration per vesicle
is determined by the number of kinesin receptors. Based on the number of kinesin bound kinesin
receptors we calculate the likelihood that a vesicle is bound to the microtubule (Table S1,
Reactions 6/7). This likelihood also depends on the affinity of kinesin to the microtubule,
incorporated into our model as the fraction of Microtubule (MT)-bound kinesin. The fraction of
MT-bound kinesin decreases from the cell body cytoplasm to the neurite shaft cytoplasm and
growth cone cytoplasm (Table S3). A low fraction of MT-bound kinesin in the neurite shaft
cytoplasm ensures that only a few vesicles (~9%) are moving forward, in agreement with
experimental data (Ahmed and Saif, 2014). The large pool of stationary vesicles in the neurite
shaft cytoplasm could serve as a membrane reservoir that allows quick recruitment of additional
membrane to the GC-PM on demand. In the growth cone cytoplasm almost all vesicles are
unbound, allowing their fusion with the GC-PM.

The Level-2 SCP *Vesicle exocytosis at GC-PM* is mediated by the interplay of the Level-3 SCPs
Vesicle Tethering and *Vesicle Fusion at GC-PM*. The former is realized in our model via a growth
cone specific tethering rate, the latter via complex formation between vesicle(v)-SNARE V and
target(t)-SNARE Y that resides at the GC-PM (Table S1, Reaction 8). Retrograde vesicle transport
from the GC-PM to the TGN is incorporated by a similar set of Level-2 and Level-3 SCPs that
consist of a symmetric set of membrane proteins and rate constants. We assume a high fraction of
dynein is attached to the MT in the neurite shaft cytoplasm, so that most of the retrograde vesicles
are actively transported towards the TGN, without formation of a vesicle reservoir.

Vesicles that fuse with the GC-PM, increase its surface area. In our model, we set the GC-PM
surface area to 50 μm^2 (Kunda et al., 2001; Ren and Suter, 2016). Any membrane surface area that
is added to the GC-PM and would increase its size beyond 50 μm^2 is added to the neurite shaft
(Table S1, Reaction 9). Membrane proteins are not added to the neurite shaft, since our model is
based on the assumption that the diffusion of the membrane proteins into the neurite shaft is
prevented by intra-membranous diffusion barriers or cortical cytoskeleton proteins (Ye and Zhang,
2013) that ensure a highly specialized growth cone plasma membrane. Neurite shaft length arises
from its membrane surface area (Table S1, Reaction 10). For simplicity, we do not consider
increase of the neurite diameter at this stage. The neurite shaft cytoplasm lies within the neurite
shaft and grows with the lengthening neurite shaft. Most of the anterograde vesicles in the neurite
shaft cytoplasm are not actively transported along the microtubule but are stationary components
of the membrane reservoir. Consequently, the growing neurite shaft cytoplasm acts as a sink for
vesicles and vesicle membrane proteins.

Microtubule (MT) growth is mediated by the interplay of the Level-2 SCPs *Dynamic Microtubule*
Growth and *Stable Microtubule Growth*. (Brady et al., 1984; Kollins et al., 2009; Song et al., 2013;
Witte et al., 2008). Dynamic MTs are characterized by periods of MT growth and catastrophic
breakdown, the latter can either lead to MT disappearance or rescue, followed by a new growth
period. The time frame of these periods is in seconds or minutes. Stable MTs do not show such

periodic growth behavior and we assume that there is no degradation of stable MTs. Dynamic MTs
are continuously generated at a specified nucleation rate (Level-3 SCP *Dynamic MT Nucleation*)
(Reaction 11 in Table S4, Figure 1D). The length of dynamic MTs (Reaction 12, Table S4) as well
as the rate of MT catastrophic breakdown depends on the free tubulin concentration that is
regulated by the SCP *Tubulin Sequestration* (Reaction 13 Table S4) (Curmi et al., 1997; Jourdain
et al., 1997; Manna et al., 2009) and on the hydrolysis rate of tubulin-bound GTP (SCP *GTP*
Hydrolysis). The latter SCP determines the size of the GTP cap at the tip of the dynamic MT that
protects it from catastrophic breakdown (Seetapun et al., 2012).

Conceptual frame work for modeling whole cell dynamics arising from interactions between
SCPs

Our model uses whole cell responses in a top-down based manner that started with the literature
curation of SCPs involved and their assignment to different levels of biological detail (Figure 2,
Step 1a), as described above. Like the organization of our Molecular Biology of the Cell Ontology
(Hansen et al., 2017), Level-1 and Level-2 SCPs describe more general, Level-3 SCPs more
specific cell biological or biochemical functions. SCPs of the different levels are in parent-child
relationships. SCPs are represented as coupled biochemical reactions that are transformed into
coupled ordinary differential equations (ODEs) (Figure 2, Steps 1b/c). We use experimental data
to define model constraints that the simulations must match (Figure 2, Step 1d). For systematic
analyses we develop analytical solutions for the prediction of kinetic parameters of SCPs at steady
state (Figure 2, Step 2a). Here, we use the term steady-state to describe a continuous whole cell
response without variation of amplitude. We use the analytical solutions to predict steady-state
kinetic parameters and values that match the predefined model constraints (Figure 1, Step 2b).
Through systematic parameter variation, we generate multiple steady-state solutions that allow for
the classification of dependencies between SCPs at different levels (Figure 2, Step 2c). To
investigate the effect of SCP dysfunction on whole-cell responses, we generate numerical
simulations based on steady-state parameter sets after arbitrary modification of kinetic parameters
of a single SCP followed by experimental verification (Figure 2, Step 2d).

Simulations, Analytical Analyses and Predictions

To characterize the coordinated activities of these Level-3 SCPs, we ran an initial set of simulations
of the growth of a single dynamic MT using a stochastic growth model (Margolin et al., 2012)
(Figure S1). Since we simulate NOG over hours and days, while dynamic MT growth behavior
changes in seconds and minutes, we do not simulate individual dynamic MTs, but the whole
population of dynamic MTs. For this population simulation, we developed two formulas based on
the results of the model by Margolin et al. They describe the dependence of the average growth
length of dynamic MTs and of the catastrophic breakdown rate that leads to complete

disappearance of the dynamic MT on the effective tubulin concentration and the GTP hydrolysis
rate (Reactions 12/13 in Table S4; Figure S1). The count of dynamic MTs depends on the
nucleation rate, the degradation rate due to catastrophic breakdown and the conversion rate into
stable MTs (Reaction 14 in Table S4; Figure S1). The activity of the Level-2 SCP *Stable*
Microtubule Growth is regulated by the Level-3 SCP *MT Stabilization* (Table S4, Reaction 15).
To calculate the length of the microtubule scaffold within the neurite, we sum the length of all
dynamic and stable MTs and divide it by the number of MTs per neurite cross-section (Reaction
16/17 in Table S4).

Experimental determination of varying neurite growth rate velocities

In primary cultures, neurons put out neurites at varying rates. To quantify the variability and
determine the velocities with which neurites grow, we used a high content imaging system (IN
Cell Analyzer) to track neurite outgrowth in primary rat cortical neurons. The experimentally
observed velocities of neurite outgrowth were used as a top-down constraint for our model. We
incubated rat cortical neurons for 16h and then documented neurite outgrowth every 6h for an
additional 48h. For each neuron, we quantified the length of its longest neurite (the one most likely
to become an axon). Representative images are shown in Figure 3A and 3B, and a complete set of
images is shown in Supplementary Figure 2. At each time point the neurites significantly differed
in their lengths (Figure 3B). We identified the longest neurite for each neuron in a field
(Supplementary Figure 2) and calculated their length distribution at every time point (Figure 3C),
identified the length of selected quantiles and calculated the growth velocity for these quantiles
(Figure 3D). The neurite growth velocities ranged from 0 to ~20 µm/h. We used these velocities
as a top-down whole- cell response constraint in modeling the activities of the Level-1, -2 and -3
SCPs. Increase in the whole-cell response, i.e. neurite outgrowth velocity, depends on the
coordinated increase of membrane production and delivery of membrane vesicles to the growth
cone as well as microtubule growth. Both types of SCPs need to facilitate growth with the same
velocity, since they are interdependent. Vesicle transport is microtubule dependent and
microtubule growth is limited by the length of the neurite shaft. Similarly, the increase in
membrane production needs to be coordinated with increased vesicle transport. More vesicles
would need to bud from the TGN, to be actively transported through the neurite shaft cytoplasm
and to fuse with the GC-PM. These interdependencies form the basis for our top-down constraint.

Postulates governing SCP dynamics

In addition to the experimentally determined NOG velocities, we used experimental data from the
literature for additional constraints at the level of SCPs. These constraints arise from a set of
postulates of permissible SCP functions that are described in (Table S5). We postulate that about
~10% of the anterograde vesicles in the neurite shaft cytoplasm are bound to the microtubule and
actively moving along the microtubule (Ahmed and Saif, 2014). The other 90% constitute a
membrane pool that can be recruited on demand to adapt to short-term increases in outgrowth
velocity. Similarly, we assume that the vesicles that reside in the growth cone (Erturk et al., 2007)

are the first membrane reservoir for the quick recruitment of additional membrane and also define
- in the same way - that 10% of the anterograde vesicles in the growth cone cytoplasm move, i.e.
fuse with the growth cone membrane. This assumption is based on the experimental observation
that GC membrane precursor vesicles reside in the GC cytoplasm before fusion with the membrane
with a half-life of ≥14min (Pfenninger et al., 2003), which would suggest that around 5% of the
vesicles fuse with the membrane. Similarly, we assume that 90% of the retrograde vesicles in the
neurite shaft cytoplasm and the cell body cytoplasm are moving. The total growth cone membrane
is completely internalized within 30 - 60 min (Diefenbach et al., 1999) which suggest an
endocytosis rate between 0.833 μm^2/min - 1.66 μm^2/min for a growth cone with a surface area of
50 μm^2 (Kunda et al., 2001; Ren and Suter, 2016). Based on these observations and the assumption
that some of the endocytosed vesicles might be part of transcytotic re-arrangements during growth
cone steering (Tojima and Kamiguchi, 2015) or are transported to other growth cones and not to
the cell body (Denburg et al., 2005), we set the rate of membrane back transport from the growth
cone to the TGN to 0.5 μm^2/min. The backward transported membrane can also be called cycling
membrane, since it continuously cycles between the TGN and the GC-PM without being added to
the neurite shaft. For the microtubules we postulate that dynamic MTs should fill up 20 μm of the
growing neurite. This is the length of the minor processes (Yu and Baas, 1994) that are
characterized by continuous retraction and extension periods (Arimura and Kaibuchi, 2007). We
assume that such periods are committed by continuous MT growth and retraction, so that the MT
scaffold in these processes should mainly consist of dynamic MTs.

Using these constraints, we ran simulations to identify an initial solution at one intermediate rate
of neurite outgrowth. From these initial simulations, we developed an overall analytical approach
that allows us to predict the relationships that produce neurite outgrowth at a specified velocity.

We identified a set of parameters that generated neurite outgrowth with a velocity of 10 μm/h
(Figure 4 and S3). Under steady-state conditions neurite, surface area continuously increased at a
rate of 0.5236 μm^2/min in agreement with the amount of membrane that is needed to increase the
length of the growing neurite (Figure 4A). The length of the microtubule bundles also increase
proportionally to maintain the shaft growth rate (Figure 4B). For this we selected an effective
tubulin concentration of 9 μM to achieve a length increase of the microtubule bundles with a
velocity of 10 μm/h. The vesicles in the neurite shaft cytoplasm increased in parallel with a rate of
~0.57 vesicles/min assuming a membrane area of 0.05 μm^2 per vesicle (Figure 4C). In considering
the values of the ordinate for the different SCPs (Figure 4) it became clear that there are multiple
combinations of kinetic parameters and amounts of components that allow neurite outgrowth at
the same velocity without violating the constraints we imposed on dynamics of the SCPs.

To enable a systematic analysis of the redundancies within and between SCPs, we generated an
analytical solution (Figure 5, and S4) for the prediction of kinetic parameters at steady state that
enable neurite outgrowth at a given velocity without violating constraints on SCP dynamics (Table
S5). In our solution, we consider that an increase in neurite outgrowth depends on varying sets of
protein amounts and kinetic parameters that allow increased net anterograde transport of

membrane vesicles and continuous synthesis of new proteins and membrane to compensate for the
accumulation within the growing neurite shaft cytoplasm. In addition to the postulated constraints
on SCP dynamics, we pre-define independent protein amounts and kinetic parameters (Figure S4,
right side) to calculate the dependent protein amounts and kinetic parameters. To achieve a given
NOG velocity a certain amount of lipid membrane needs to be synthesized and transported from
the TGN to the GC. In addition to this membrane that will be added to the neurite shaft three other
membrane types can be classified based on their transport destination, allowing us to distinguish
four membrane types (Figure 5A). These membrane types should not be confused with the four-
different vesicle sets that transport the membrane between the TGN and GC-PM. Type I membrane
buds from the TGN and is added to the vesicle reservoir in the growing neurite shaft cytoplasm.
Type II membrane is transported from the TGN to the GC-PM and added to the neurite shaft as
anterograde vesicles. Type III membrane is transported anterogradely from the TGN to the GC-
PM, endocytosed at GC-PM and added to the NSC as retrograde vesicles. Type IV membrane is
transported from TGN to GC-PM and from there back to TGN, and therefore is labeled cycling
membrane. Initial and final anterograde and retrograde rates that describe budding and fusion
processes at the TGN and GC-PM can be calculated by summing up the involved membrane types.
The fluxes of the individual membrane types can be calculated based on the anticipated NOG
velocity and model constraints (Figure 5B, (1)). To allow the fusion of sufficient anterograde
vesicles with the GC-PM, a certain number of SNARE complexes need to be formed between v-
SNAREs V and t-SNAREs Y (Figure 5B, (2)). Consideration of the tethering rate allows the
calculation of the needed amount and concentration of v-SNARE Vs. The concentration allows
the determination of the initial and final forward fluxes of SNARE V ((Figure 5B, (3)). Similarly,
concentrations and fluxes can be calculated for the v-SNARE U that mediates vesicle fusion at the
TGN ((Figure 5B, (4)/ (5)). During vesicle budding at the TGN and GC-PM, SNAREs compete
with each other for the vesicle binding spots. Knowledge of the fluxes of both v-SNAREs allows
the calculation of their concentrations at the TGN and GC-PM (Figure 5, (6)/ (7)). The production
rates of v-SNARES (and any other membrane vesicle membrane proteins) can be calculated based
on their vesicle concentrations and the number of vesicles that accumulate in the growing neurite
shaft cytoplasm (Figure 5B, (8)).

MT growth at a given velocity (Figure 5C) depends on the conversion rate of dynamic into stable
MTs (Figure 5D, (9)). By pre-simulation results using the model of Margolin et al, we obtained
the dependence of the average length of dynamic MTs on the free (i.e. not sequestered) tubulin
concentration and GTP hydrolysis rate. The average length allows the calculation of the number
of dynamic MTs (Figure 5D, (10)), followed by calculation of the stabilization rate (Figure 5D,
(11)) and subsequently the nucleation rate (Figure 5D, (12)).

The analytical solution was validated by comparing the analytically predicted outcome with the
results of the numerical simulation for example sets of parameter and protein amounts (Figure 5E
and S5). We used the predicted parameter and protein amount sets as starting points in our model
and ran the numerical simulation for 5000 minutes (~3.5 days). Every 500 min we compared the

anticipated model constraints with the numerically obtained values to estimate the accuracy of our
analytical prediction. The analytical solution matched the anticipated model constraints with high
accuracy, except the fusion rate for retrograde vesicles with the TGN that for was too high for the
velocities 0 and 2.5 μm/h.

Complementary Compensatory Relationships between Level-3 sibling SCPs can contribute
to robustness of response

We used the analytical solution to systematically analyze the relationship between different SCPs
(Figure 1B). For our studies we selected growth velocities that covered the whole range of
experimentally observed velocities (Fig 6). We predicted that Level-3 sibling SCPs would have
complementary activities. An increase in the activity of one Level-3 SCP can be compensated by
a decrease in the activity of at least one of its sibling Level-3 SCPs. To acquire a specified NOG
velocity without violation of the model constraints there are multiple combinations of the activities
of the two Level-3 SCPs *Coat Recruitment and Formation* and *Vesicle Invagination and Scission*
(Figure 6A), the two children SCPs of the Level-2 SCP *Vesicle budding at the TGN*. Such SCP
redundancies are also observed between the Level-3 SCPs *Kinesin Recruitment to the Vesicle* and
Kinesin Mediated Vesicle Transport along the MT (Figure 6B) that define the activity of the Level-
2 SCP *Anterograde Microtubule-Based Vesicle Transport*. Similarly, the SCPs *Vesicle Tethering*
and *Vesicle Fusion* (Figure 6C), sub processes of the SCP *Vesicle Exocytosis* are characterized by
an interaction that is based on an inverse relationship between their activities.

The balance of the Level-3 SCP activities to fulfill the Level-2 SCP function is not isolated at
Level -3 but influences the activity of other SCPs as well. The more the activity of a Level-2 SCP
arises from a Level-3 SCP that involves vesicle proteins, the higher the need for continuous
membrane protein supply to compensate for the consumption of membrane proteins in the neurite
shaft cytoplasm sink. This accounts for the kinesin receptor (Figure 6E) as well as for the v-
SNARE V (Figure 6F). Here, one might consider that the complementary SCP *Kinesin-Mediated*
Vesicle Transport along the MT is determined by components along the whole microtubule while
the complementary SCP *Vesicle Tethering* at GC-PM is determined by components of the tethering
machinery that are locally restricted to the growth cone. The maintenance of the activity of the
former SCP might therefore depend on a continuous synthesis of cytoplasmic protein that is not
considered in our model.
392
Similar compensatory relationships exist between Level-3 sibling SCPs involved in Microtubule
Growth. The combined length of dynamic MTs that are available for conversion into stable MTs
is regulated by an inverse relation between the activities of the Level-3 SCPs *Dynamic MT*
Nucleation, *GTP Hydrolysis* and *Tubulin Sequestration*. The average dynamic MT length (Figure
7B) as well as the degradation rate of dynamic MTs (Figure 7C) depends upon the Level-3 SCPs
Tubulin Sequestration (which regulates effective tubulin concentration) and *GTP Hydrolysis*. The
balanced activities of these two Level-3 SCPs and the Level-3 SCP *Dynamic Microtubule*
Nucleation (Figure 7D/E) determines the activity of the Level-2 SCP *Dynamic Microtubule*

Growth. Similarly, to the vesicle transport Level-3 SCPs, any loss of function in one of these MT
Level-3 SCPs can be compensated by a gain of function in one of the other.

Sibling Level 2 SCPS do not readily complement each other leading to dystrophic bulbs

The relationships between Level-2 SCP activities are more complicated. A decrease in the activity
of one Level-2 SCP cannot simply be compensated by increase in the activity of one of its Level-
2 sibling SCPs. To simulate mismatching activity of a Level-2 SCP on NOG, we first predicted a
set of kinetic parameters that allow growth under physiological conditions as defined by our model
constraints. Then we selected one of its Level-3 SCP children, followed by the perturbation of one
kinetic parameter that belonged to this Level-3 SCP, so that this parameter no longer matched
anymore to the predicted parameter set. The resulting unbalanced activities between the redundant
Level-3 SCP siblings propagated to impaired Level-2 SCP activity that did not match with the
activities of the other Level-2 SCPs. We then searched for adaptations in other parameters that
allow restoration of neurite outgrowth under physiological conditions as defined by our model
constraints. As shown for the example Level-2 SCP *Vesicle Budding at the Trans-Golgi Network*,
multiple Level-2 SCP activities needed to be altered in a coordinated manner (Figure 8), e.g.
Membrane Lipid and *Protein Production* (simulated via reduced production rates), *Anterograde*
Vesicle Transport along the Microtubule (simulated via reduced amount of Kinesin receptor) and
Vesicle Exocytosis at the GC-PM (simulated via reduced amount of v-SNARE V). These activity
changes were associated with a reduction of NOG velocity (from 10 μm/h to 5.5 μm/h), since the
cycling rate was kept constant at 0.5 $μm^2$/min.

To simulate the effect of mismatching Level-2 SCP activities on NOG, we reduced the activity of
one Level-2 SCP (as described above, but without adaptation of any other Level-2 SCP activities).
We selected one *Vesicle Transport & Exocytosis* SCP and one *Microtubule Growth* SCP for further
investigation. To simulate the effect of the mismatching activity of the Level-2 SCP *Vesicle*
Exocytosis at GC-PM, we selected the v-SNARE V that is involved in the activity of its Level-2
child SCP *Vesicle Fusion at the Growth Cone*. We predicted that a reduction in its activity does
not influence NOG velocity, but overloads the growth cone with anterograde vesicles, a feature
associated with dystrophic bulbs (Bradke et al., 2012) (Figure 9A/B). To test our predictions, we
selected the v-SNARE VAMP7 that is involved in vesicle fusion at the GC-PM (Burgo et al.,
2012). In agreement with our predictions, VAMP7 knock down (KD) did not significantly decrease
NOG velocity, as shown by the total length of neurites 48h after axotonomy (Figure 9C/D). Its
main effect was the generation of dystrophic bulbs (Figure 9C/E).

To simulate impaired activity of the Level-2 SCP *Stable Microtubule Growth* we selected its
Level-3 SCP child *Microtubule Stabilization* and perturbed the kinetic parameter stabilization rate.
Our simulations predicted a significant decrease in the total MTB length that affects NOG (Figure
9F). We validated our predictions by knocking down the microtubule crosslinking protein MTCL1
that mediates MT stabilization (Satake et al., 2017). MTCL1 KO significantly reduced NOG
velocity and increased the number of dystrophic bulbs (Figure 9G/H/I).

Discussion

Whole cell responses depend on subcellular activities at different levels

Neurite outgrowth depends on the coordinated activities of various sub-cellular processes (SCPs). Of the different SCPs involved in our model, the types of Level-1 SCPs are critical. Three major sub-cellular functions that are covered by these SCPs are *Production of Membrane Components, Vesicle Transport & Exocytosis* and *Microtubule Growth.* We curated SCPs that are involved in these Level-1 sub-cellular functions and classified them as Level-2 and Level-3 SCPs that are in parent-child relationships. Based on the hierarchical relationships between these SCPs, we developed a dynamical model that incorporates the three types of Level-1 SCPs and documented that the activity of the SCPs involved need to be highly coordinated to allow neurite outgrowth at a certain velocity. To make the model more manageable, our simulations were done under several assumptions: (1) the size of the TGN & GC-PM (50 µm^2) is independent of the outgrowth velocity; (2) the rate of back transported membrane from the growth cone to the TGN is 0.5 µm^2/min; (3) 90 % of the anterograde vesicles in the neurite shaft cytoplasm and the growth cone cytoplasm serve as membrane reservoir (i.e. these vesicles are not actively transported along the MT or fusing with the GC-PM) and (4) 20 µm of the neurite microtubule scaffold consists of dynamic MTs and the rest of stable MTs. All these assumptions are based on previous experimental data and allowed us to identify the key features of the balance between SCP types required for a dynamic whole cell response. We developed an analytical prediction for steady state dynamics at the specified NOG velocities that allowed us to investigate SCP relationships in a systematic manner.

Complementary dependence in lower level SCPs enables robustness of response, while imbalances in higher level SCPs are not readily compensated

We could show that the activities of Level-3 SCP siblings needed to generate the function of their Level-2 SCP parent inversely depends on each other. This applied to both Level-3 SCPs involved in *Vesicular Transport & Exocytosis* as well as Level-3 SCPs involved in *Microtubule Growth*. This complementary compensatory capability will confer the cell to ability to mount a response in a robust manner

 In contrast, the activities of Level-2 SCPs must be highly coordinated to maintain the Level-1 SCP function. The impaired activity of one Level-2 SCP cannot be simply compensated by an increased activity of another Level-2 SCP. Dependencies between Level-2 SCPs are more complex. In case of *Vesicle Transport & Exocytosis*, loss of function of one Level-2 SCP might only be compensated, if multiple other Level-2 SCPs also reduce their function. Impaired Level-2 SCP activities were simulated by reduction of the activity of one of its Level-3 children SCPs without compensatory activity modification of the other.

Our predictions proposed that reduced exocytosis does not influence NOG velocity. We predicted that this results in an excess of anterograde vesicle in the GC that can be one of the reasons for the

formation of dystrophic bulbs. In agreement with our prediction, knockdown of v-SNARE
VAMP7 did not change NOG velocity but rather, it creates a pathological phenotype with multiple
dystrophic bulbs instead of healthy growth cones.

A reduction in the growth of stable MTs was predicted to cause a reduction in NOG length, and it
was experimentally confirmed by knock down of MTCL1. As the activities of the SCPs related to
Vesicle Transport & Exocytosis are unchanged, in this case the reduction in neurite outgrowth will
be accompanied by the appearance of dystrophic bulbs. This prediction was also experimentally
confirmed.

Different subcellular processes can give rise to similar whole cell phenotypes

The observations in this study both computational and experimental demonstrate the critical need
for balance and coordination between different types of subcellular processes required to mount a
whole cell response. It is noteworthy that robustness of whole cell responses arises from the
relative redundancy of lower level subcellular processes within each type. However at least for
neurite outgrowth, such redundancies do not occur at higher level SCPs where, when imbalance
occurs the whole cell function shuts down even though each high level SCP might be operational
for a certain time period. It is the uncoordinated functioning of the higher level SCPs that result in
the morphological abnormalities like the dystrophic bulbs. Since lack of coordination can arise to
due to dysfunction of different higher level SCPs we get similar dystrophic bulbs when we partially
ablate a microtubule stabilizing protein or a vesicle fusion protein. This provides a striking
example of phenotypic convergence in spite of molecular divergence.

Overall the major conclusions of this study are that 1) elements of both robustness and fragility
coexist within subcellular processes needed to produce whole cell responses; 2) since multiple
types of subcellular processes are involved in mounting a whole cell responses changes in in
different molecular entities or subcellular processes can result in the same aberrant phenotype; 3)
quantitative imbalances between different types of subcellular processes are sufficient for
alteration in cellular dynamics causing abnormal phenotype.

Acknowledgements: This research was supported by NIH grant GM54508 and the Systems Biology Center grant GM 071558. We thank Alan Stern and Marc Birtwistle with help in image acquisition using the InCell analyzer and Frank Bradke, Tatiana Stepanova and Ali Erturk for providing experimental data for constraining our model. We thank Nikolaos Tzavaras for helping us in image quantification in MetaMorph. We thank Drs Joseph Goldfarb and Robert D Blitzer for critical reading of the manuscript.

Author contributions: R.I. and J.H. conceived the project. A.Y. and J.H. developed and implemented the model. J.H. developed the analytical prediction for steady state dynamics. M.S, V.R. and R.T. prepared rat cortical neurons. A.Y. and M.S. analyzed the images. A.Y., R.I. and J.H. wrote the manuscript.

Declaration of Interests: The authors declare no competing interests.

References

Ahmed, W.W., and Saif, T.A. (2014). Active transport of vesicles in neurons is modulated by mechanical tension. Scientific reports *4*, 4481.

Arimura, N., and Kaibuchi, K. (2007). Neuronal polarity: from extracellular signals to intracellular mechanisms. Nature reviews Neuroscience *8*, 194-205.

Azeloglu, E.U., Hardy, S.V., Eungdamrong, N.J., Chen, Y., Jayaraman, G., Chuang, P.Y., Fang, W., Xiong, H., Neves, S.R., Jain, M.R.*, et al.* (2014). Interconnected network motifs control podocyte morphology and kidney function. Science signaling *7*, ra12.

Bhalla, U.S., and Iyengar, R. (1999). Emergent properties of networks of biological signaling pathways. Science *283*, 381-387.

Blanquie, O. and Bradke, F. (2018), Cytoskeleton dynamics in axon regeneration, current opinion in neurobiology, 51, 60-69.

Bradke, F., Fawcett, J.W. and Spira, M.E. (2012). Assembly of a new growth cone after axotomy: the precursor to axon regeneration. Nature reviews neuroscience 13, 183-193.

Brady, S.T., Tytell, M., and Lasek, R.J. (1984). Axonal tubulin and axonal microtubules: biochemical evidence for cold stability. J Cell Biol *99*, 1716-1724.

Burgo, A., Proux-Gillardeaux, V., Sotirakis, E., Bun, P., Casano, A., Verraes, A., Liem, R.K., Formstecher, E., Coppey-Moisan, M., and Galli, T. (2012). A molecular network for the transport of the TI-VAMP/VAMP7 vesicles from cell center to periphery. Dev Cell *23*, 166-180.

Curmi, P.A., Andersen, S.S., Lachkar, S., Gavet, O., Karsenti, E., Knossow, M., and Sobel, A. (1997). The stathmin/tubulin interaction in vitro. The Journal of biological chemistry *272*, 25029-25036.

Denburg, J.L., Hughen, R.W., Tucker, D., and Kater, S.B. (2005). Fate of constitutive endocytic vesicles formed in the growth cone: transport of vesicles from one growth cone to another in the same neuron. Journal of neurobiology *62*, 262-277.

Diefenbach, T.J., Guthrie, P.B., Stier, H., Billups, B., and Kater, S.B. (1999). Membrane recycling in the
neuronal growth cone revealed by FM1-43 labeling. The Journal of neuroscience : the official journal of
the Society for Neuroscience *19*, 9436-9444.
Erturk, A., Hellal, F., Enes, J., and Bradke, F. (2007). Disorganized microtubules underlie the formation of
retraction bulbs and the failure of axonal regeneration. The Journal of neuroscience : the official journal
of the Society for Neuroscience *27*, 9169-9180.
Filious, A. R., and Schwab, J.M. (2018), Determinants of Axon Growth, Plasticity, and Regeneration in the
Context of Spinal Cord Injury, The American Journal of Pathology, 188, 53-62.
Fivaz, M., Bandara, S., Inoue, T., and Meyer, T. (2008). Robust neuronal symmetry breaking by Ras-
triggered local positive feedback. Current biology : CB *18*, 44-50.
Gracias, N.G., Shirkey-Son, N.J., and Hengst, U. (2014). Local translation of TC10 is required for membrane
expansion during axon outgrowth. Nature communications *5*, 3506.
Hansen, J., Meretzky, D., Woldesenbet, S., Stolovitzky, G., and Iyengar, R. (2017). A flexible ontology for
inference of emergent whole cell function from relationships between subcellular processes. Sci Rep *7*,
17689.
Heinrich, R., and Rapoport, T.A. (2005). Generation of nonidentical compartments in vesicular transport
systems. The Journal of cell biology *168*, 271-280.
Hill, C. E. (2017), A view from the ending: Axonal dieback and regeneration following SCI, Neuroscience
Letters, 652, 11-24.
Jourdain, L., Curmi, P., Sobel, A., Pantaloni, D., and Carlier, M.F. (1997). Stathmin: a tubulin-sequestering
protein which forms a ternary T2S complex with two tubulin molecules. Biochemistry *36*, 10817-10821.
Kiddie, G., McLean, D., Van Ooyen, A., and Graham, B. (2005). Biologically plausible models of neurite
outgrowth. Progress in brain research *147*, 67-80.
Kollins, K.M., Bell, R.L., Butts, M., and Withers, G.S. (2009). Dendrites differ from axons in patterns of
microtubule stability and polymerization during development. Neural Dev *4*, 26.
Kunda, P., Paglini, G., Quiroga, S., Kosik, K., and Caceres, A. (2001). Evidence for the involvement of Tiam1
in axon formation. The Journal of neuroscience : the official journal of the Society for Neuroscience *21*,
2361-2372.
Manna, T., Thrower, D.A., Honnappa, S., Steinmetz, M.O., and Wilson, L. (2009). Regulation of microtubule
dynamic instability in vitro by differentially phosphorylated stathmin. The Journal of biological chemistry
284, 15640-15649.
Margolin, G., Gregoretti, I.V., Cickovski, T.M., Li, C., Shi, W., Alber, M.S., and Goodson, H.V. (2012). The
mechanisms of microtubule catastrophe and rescue: implications from analysis of a dimer-scale
computational model. Molecular biology of the cell *23*, 642-656.
Pfenninger, K.H., Laurino, L., Peretti, D., Wang, X., Rosso, S., Morfini, G., Caceres, A., and Quiroga, S.
(2003). Regulation of membrane expansion at the nerve growth cone. Journal of cell science *116*, 1209-
1217.
Ren, Y., and Suter, D.M. (2016). Increase in Growth Cone Size Correlates with Decrease in Neurite Growth
Rate. Neural plasticity *2016*, 3497901.
Satake, T., Yamashita, K., Hayashi, K., Miyatake, S., Tamura-Nakano, M., Doi, H., Furuta, Y., Shioi, G., Miura,
E., Takeo, Y.H*., et al.* (2017). MTCL1 plays an essential role in maintaining Purkinje neuron axon initial
segment. EMBO J *36*, 1227-1242.
Seetapun, D., Castle, B.T., McIntyre, A.J., Tran, P.T., and Odde, D.J. (2012). Estimating the microtubule GTP
cap size in vivo. Curr Biol *22*, 1681-1687.
Song, Y., Kirkpatrick, L.L., Schilling, A.B., Helseth, D.L., Chabot, N., Keillor, J.W., Johnson, G.V., and Brady,
S.T. (2013). Transglutaminase and polyamination of tubulin: posttranslational modification for stabilizing
axonal microtubules. Neuron *78*, 109-123.

Tojima, T., and Kamiguchi, H. (2015). Exocytic and endocytic membrane trafficking in axon development.
Development, growth & differentiation *57*, 291-304.
Tsaneva-Atanasova, K., Burgo, A., Galli, T., and Holcman, D. (2009). Quantifying neurite growth mediated
by interactions among secretory vesicles, microtubules, and actin networks. Biophysical journal *96*, 840-
857.
Wang, T., Liu, Y., Xu, X.H., Deng, C.Y., Wu, K.Y., Zhu, J., Fu, X.Q., He, M., and Luo, Z.G. (2011). Lgl1 activation
of rab10 promotes axonal membrane trafficking underlying neuronal polarization. Developmental cell *21*,
431-444.
Wissner-Gross, Z.D., Scott, M.A., Ku, D., Ramaswamy, P., and Fatih Yanik, M. (2011). Large-scale analysis
of neurite growth dynamics on micropatterned substrates. Integrative biology : quantitative biosciences
from nano to macro *3*, 65-74.
Witte, H., Neukirchen, D., and Bradke, F. (2008). Microtubule stabilization specifies initial neuronal
polarization. J Cell Biol *180*, 619-632.
Ye, F., and Zhang, M. (2013). Structures and target recognition modes of PDZ domains: recurring themes
and emerging pictures. The Biochemical journal *455*, 1-14.
Yu, W., and Baas, P.W. (1994). Changes in microtubule number and length during axon differentiation.
The Journal of neuroscience : the official journal of the Society for Neuroscience *14*, 2818-2829.

Figure 1A

Growth Cone

Dystrophic Bulb

Figure 1B

| Whole cell function | Level-1 SCPs | Level-2 SCPs | Level-3 SCPs |

The figure shows a hierarchical network diagram connecting whole cell function to SCPs at different levels.

Whole cell function: Neurite Outgrowth

Level-1 SCPs: Production Membrane Components; Vesicle Transport & Exocytosis; Microtubule Growth

Level-2 SCPs: Membrane Lipid Production; Membrane Protein Production; Vesicle Budding at Trans-Golgi Network; Anterograde Microtubule Based Vesicle Transport; Vesicle Exocytosis at Growth Cone PM; Vesicle Endocytosis at Growth Cone PM; Retrograde Microtubule Based Vesicle Transport; Vesicle Fusion with Trans-Golgi Network; Dynamic Microtubule Growth; Stable Microtubule Growth

Level-3 SCPs: Coat Protein Recruitment and Formation at TGN; Vesicle Invagination and Scission at TGN; Kinesin Recruitment to Vesicle; Kinesin Mediated Vesicle Transport along Microtubule; Vesicle Tethering at Growth Cone PM; Vesicle Fusion with Growth Cone PM; Coat Protein Recruitment and Formation at GC-PM; Vesicle Invagination and Scission at GC-PM; Dynein Recruitment to Vesicle; Dynein Mediated Vesicle Transport along Microtubule; Vesicle Tethering at Trans-Golgi Network; Vesicle Fusion with Trans-Golgi Network; Dynamic Microtubule Nucleation; Tubulin Sequestration; GTP Hydrolysis; MT Stabilization

678 **Figure 1C**

Figure 1D

Stable Microtubule pool

Dynamic Microtubule pool

$[f_{12}]$ **Degradation of dynamic MTs**

$[f_{10}]$ **Nucleation of dynamic MTs**

$[f_{14}]$ **Conversion of dynamic into stable MTs**

Figure 2

Top down based SCP modelling of whole cell dynamics

Build ODE model of SCP dynamics

Generate analytical representation for systematic analysis of SCP relationships

Step 1a: List different classes of SCPs and their relationships at various levels required for whole cell responses

Step 2a: Develop analytical representations for SCP relationships

Step 1b: Represent SCPs as coupled biochemical reactions

Step 2b: Use analytical prediction to identify parameter sets that allow simulations in agreement with experimental data

Step 1c: Develop coupled ODE models for different SCPs

Step 2c: Classify SCP sets at different levels as redundant or non-redundant for whole cell function

Step 1d: Generate and identify experimental data that ODE model-based simulations have to match

Step 2d: Predict effect of SCP imbalances for experimental testing

Figure 3

Figure 4

Figure 5

A

B

---

1) Calculate Type I – IV membrane transport rates and initial and final transport rates

    1.1) $\boldsymbol{Neurite\ shaft\ growth} = \boldsymbol{Type\ II\ membrane\ transport\ rate} = \boldsymbol{NOG\ velocity} \times 2\boldsymbol{\pi} \times \boldsymbol{r_{neurite}}$

    Calculate type I, III and IV transport rates based on NOG velocity, model constraints and Type II membrane transport rate

    Calculate initial and final forward and backward transport rates as indicated in (A)

2) Calculate v – SNARE V protein amount and concentration in GCC that allows final forward transport rate

    2.1) $YV\ SNARE\ complex\ in\ GC = \dfrac{Final\ forward\ transport\ rate\ \times SNARE\ complex\ per\ vesicle\ fusion}{Anterograde\ vesicle\ surface\ area}$

    2.2) $VV_{GCC}^{B_G} = \dfrac{YV\ SNARE\ complexes\ GC}{tethering\ rate\ at\ GC\ \times\ YY_{PM}}$,   2.3) $V_{GCC}^{B_G} = \dfrac{VV_{GCC}^{B_G}}{n_{GCC}^{B_G}}$

3) Calculate v – SNARE V forward fluxes

    3.1) $VV\ initial\ forward = V_{GCC}^{B_G} \times initial\ forward\ transport\ rate$

    3.2) $VV\ final\ forward = V_{GCC}^{B_G} \times final\ forward\ transport\ rate$

4) Calculate v – SNARE U protein amount and backward fluxes similarly

5) Calculate v – SNARE U concentration at GC that allows v – SNARE U backward fluxes

    5.1) $VV\ final\ forward = V_{GCC}^{B_G} \times final\ forward\ transport\ rate$

6) Calculate v – SNARE U concentration at GC that allows v – SNARE U backward fluxes

    6.1) $const_{GC} = 1 + \dfrac{Y_{GC}}{kya} + \dfrac{X_{GC}}{kxa}$ 6.2) $b_s = initial\ backward\ transport\ rate\ \times SNARE\ binding\ spots\ per\ vesicle\ area$

    6.3) $U_{GC} = \dfrac{UU\ initial\ backward\ \times const_{GC} \times kva}{(b_s - VV\ initial\ backward\ \times UU\ initial\ backward)}$

7) Calculate v – SNARE V concentration at GC

    7.1) $V_{GC} = \dfrac{VV\ initial\ backward \times (\frac{U_{GC}}{kua} + const_{GC}) \times kva}{(b_s - VV\ initial\ backward\ )}$

8) Calculate v – SNARE V production rate at TGN to compensate for v – SNARE V accumulation in MTC

    8.1) $VV\ production\ rate = membrane\ type\ I \times V\ forward\ rate + membrane\ type\ III \times V\ backward$

C

793 **D**

---

(9) Calculate MTB growth velocity

$$NOG\ velocity = \ MTB\ growth\ velocity = \frac{Length\ increase\ of\ stable \times MTs}{MTs\ per\ crossection}$$

(10) Calculate number of dynamic MTs

$$Number\ dynamic\ MTs = \frac{Total\ dyn\ MTs\ length}{(Average\ dynamic\ MT\ length)}$$

(11) Calculate MT stabilization rate

$$stablization\ rate = \frac{Length\ increase\ of\ stable\ MTs}{(Average\ dynamic\ MT\ length\ \times Number\ dynamic\ MTs)}$$

(12) Calculate MT nucleation rate

$$Nucleation\ rate\ = (Stabilization\ rate + Degradation\ rate) \times Total\ number\ dynamic\ MTs$$

E

804 **Figure 6**

 Figure 7

851 **Figure 8**

863 **Figure 9**

865 **A**   **B**

C

D

E

F

G

H

I

Figure Legends

Figure 1: Identification of different level SCPs involved in neurite outgrowth and their
incorporation into multicompartmental ODE model. (A) Based on our observations dystrophic
bulbs (right panel) revealed by tubulin staining are 8-10 times larger than physiological growth
cones (left panel). Characteristics of dystrophic bulbs are an accumulation of anterograde vesicles
(Bradke et al., 2012) as well as disorganized microtubules (Blanquie and Bradke, 2018)**. (B)** The
three major SCP types, *Production of Membrane Components, Vesicle Transport & Exocytosis*
and Microtubule Growth, describe essential sub-cellular functions involved in neurite outgrowth.
We labeled them as Level-1 SCPs and identified their Level-2 children and Level-3 grandchildren
SCPs. **(C)** To model membrane production at the Trans-Golgi network (TGN) and delivery to the
growth cone plasma membrane (GC-PM) via vesicle transport, we extended a dynamical model
that simulates vesicle transport between the endoplasmic reticulum and the Golgi (Heinrich and
Rapoport 2005). In our version, new synthesized membrane is added to the Trans-Golgi network
from where it is transported to the growth cone plasma membrane by vesicular transport. Vesicles
bud from the TGN, move along the microtubules via active kinesin mediated transport and fuse
with the growth cone plasma membrane, leading to neurite shaft growth. The vesicles pass through
three intermediate compartments: the cell body cytoplasm (CBC), the neurite shaft cytoplasm
(NSC) and the growth cone cytoplasm (GCC). In each compartment kinesin has a different affinity
for microtubules, leading to varying fractions of vesicles that are actively transported via kinesin
along the MT. Retrograde vesicles are generated via endocytosis at the growth cone plasma
membrane and move along the microtubule towards TGN through dynein mediated active
transport. Vesicle budding at the TGN or GC-PM is mediated by the interaction of recruitment
factors and coat proteins, and vesicle fusion with the TGN or the growth cone membrane by the
formation of SNARE complexes between vesicle(v)-SNAREs and target(t)-SNAREs, catalyzed
by local tethering machineries. Motor proteins are bound to vesicles through motor protein
receptors. A_G/B_G label vesicles that bud from the TGN with coat protein A/B, A_{PM}/B_{PM} vesicles
that bud from the GC-PM with coat protein A/B. Since at steady state almost all anterograde
vesicles bud with coat protein B from the TGN, and almost all retrograde vesicles bud with coat
protein A from the GC-PM, we highlighted their transport routes in brighter colors. **(D)** To
simulate MT growth, we consider two different MT pools, stable and dynamicMTs. After
nucleation new MTs are added to the pool of dynamic MTs that is characterized by alternating
phases of growth and catastrophic breakdown. The frequency and duration of these phases depend
on the tubulin concentration (regulated by the SCP *Tubulin Sequestration*) and GTP hydrolysis
rate (SCP *GTP Hydrolysis*). Consequently, the length distribution of the dynamic MT pool and the
degradation rate of dynamic MTs depends on the activity of both SCPs (supplementary figure 2).
Dynamic MTs are either degraded or converted into stable MTs that form the growing neurite
scaffold.

Figure 2: Top-down based SCP modelling of whole cell dynamics. Our pipeline for the generation of dynamical models that allow the analysis of relationships between sub-cellular processes (SCPs) during whole cell responses consists of two major steps. First, we build multicompartmental ODE models for the simulation of SCP activities that are based on literature curated data (Step 1a-d). In the second step we develop analytical predictions for kinetic parameters at steady state that allow a systematic analysis of SCP interactions (Step 2a-d).

Figure 3: Neurites grow with different outgrowth velocities. Neurons were dissected from rat cortical brain, incubated for 16h to allow initial growth, and followed by image acquisition every 6h up to 70h after plating. **(A-B)** The growth dynamics of two example neurons, one with a low (A) and one with a high (B) outgrowth velocity, are shown over given time-period. **(C)** At each timepoint we quantified the length of the longest neurite of each neuron. Histograms show the length distribution of all longest neurites at the indicated timepoints. **(D)** At each timepoint we identified the top and bottom 10% and 25% quantile length as well as the median length. Linear interpolation of the lengths for each selected quantile documented outgrowth velocities ranging from 0 to ~20 µm/h.

Figure 4: Simulation of vesicle transport and microtubule growth during neurite outgrowth. **(A-B)** We identified a set of kinetic parameters (Supplementary table S7) that allows neurite outgrowth via coordinated neurite shaft (A) and microtubule bundle growth (B) with a velocity of 10 µm/h. **(C)** The growing neurite shaft cytoplasm (NSC) within the growing neurite acts as a sink for anterograde and retrograde vesicles, causing an accumulation of vesicles within this compartment. **(D)** Our identified set of kinetic parameters allows NOG without violation of the model constraints that we selected to define physiological outgrowth conditions.

Figure 5: Generation of an analytical solution for the prediction of kinetic parameters at steady state. **(A)** The development of the analytical solution depends on the categorization of membrane fluxes based on their destination compartment. Such classification distinguishes four different membrane types (I-IV) (that should not be confused with the four different vesicle types we show in figure 2B). Initial and final anterograde and retrograde transport rates (which refer to budding and fusion rates at the TGN and GC) are the sum of the transport rates of the encircled membranes types. **(B)** Calculation of kinetic parameter for v-SNARE V. VV/YY/UU refer to total count of SNAREs, V/Y/U describe concentration of SNAREs (per membrane area). See supplementary methods for complete derivation of analytical solution. **(C)** Microtubule bundle (MTB) growth is mediated by the conversion of dynamic MTs into stable MTs. Dynamic MTs nucleate de-novo, followed by their degradation or stabilization. **(D)** Nucleation, stabilization and degradation rates for dynamic MTs are calculated as shown. **(E)** To verify our analytical solution we analyzed, whether our predicted parameter sets allow neurite outgrowth with the selected velocity and without violation of model constraints. For each velocity, we selected 6 different parameter sets that differ in the number of v-SNARE V and the tethering rate at the growth cone

plasma membrane (supplementary figure 5A). Every 500 min over a simulation period of 5,000 min (~3.5days), we documented the indicated outputs. Averages and standard deviations were calculated for each velocity. As shown for a velocity of 15 μm/h, our analytical solution predicts steady state dynamics that match the anticipated model constraints (horizontal lines) with high accuracy.

Figure 6: Redundancies among vesicle transport level-3 sibling SCPs. To analyze the relationships between Level-3 SCPs, we used our analytical solution to predict steady state dynamics that enable neurite outgrowth for specified velocities. **(A), (B) & (C)** Level-3 SCPs show complementary relationships between their activities to generate higher level SCP function without violation of the model constraints. **(A)** Different activities of the SCPs *Coat Recruitment and Formation at TGN* and *Vesicle Invagination and Scission at TGN* were modeled by changing the amount of Recruitment Factor 1 and the budding rate at the TGN. SCP activities were inversely related and increased with increasing velocity. Line colors indicate selected velocities as shown in (G). **(B)** A similar SCP redundancy was observed for the SCPs *Kinesin Recruitment to Vesicle* (simulated by # number of kinesin receptors) and *Kinesin Mediated Vesicle Transport along the MT* (simulated by fraction of bound kinesin). **(C)** Inverse relationship between SCP activities was also observed for the SCPs *Vesicle Tethering* (simulated via the tethering rate at the GC) and *Vesicle Fusion* (simulated by amount # of v-SNARE V). **(D)** Velocities were color coded in each figure as indicated. **(E) & (F)** The more a Level-2 SCP activity is generated by a Level-3 SCP that contains a vesicle membrane protein, the higher the SCP activity of membrane protein production to compensate for the loss of membrane protein in the growing NSC reservoir. **(E)** In dependence of the selected fraction of MT-bound kinesin, vesicles need a different amount of kinesin receptors to ensure NOG without violation of the model constraints. A higher kinesin receptor concentration per vesicle directly translates into the need for a higher kinesin production rate. Lines refer to pre-defined fractions of MT-bound kinesin: 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0 (from top to bottom). **(F)** Different tethering rates demand different v-SNARE V concentrations per vesicle and consequently different V production rates to ensure NOG without violation of the model constraints (t-SNARE Y is kept constant). Lines refer to different tethering rates: 2.5×10^{-6}, 3×10^{-6}, 3.5×10^{-6}, 4×10^{-6}, 4.5×10^{-6}, 5×10^{-6} (from top to bottom).

Figure 7: Redundancies among microtubule growth Level-3 sibling SCPs. (A) Summary of the basic reaction for the simulation of MT growth. See Figure 2C for details. **(B), (C), (D) & (E)** Level-3 SCPs that define sub-functions of the Level-2 SCP *Dynamic MT growth* show complementary activities to allow NOG without violation of the model constraints. Multiple combinations of the SCP activities *Tubulin Sequestration*, *GTP Hydrolysis* and *Nucleation of Dynamic MTs* enable NOG with a given velocity. **(B-C)** The influence of the SCPs *Tubulin Sequestration* (which determines the effective tubulin concentration) and *GTP Hydrolysis* on the length distribution (B) and degradation rate (C) of the dynamic MTs were determined in a pre-simulation using a stochastic model for the simulation of single dynamic MT growth (Margolin et al). Colored lines refer to the selected GTP hydrolysis rates shown in (B) and (C) (pink: 0.72/sec,

d) and (yellow: 0.675/sec, e). **(D-E)** Multiple combinations of dynamic MT nucleation rates and
effective tubulin concentrations allow NOG under a given velocity for each GTP hydrolysis rate,
as shown for the hydrolysis rate of 0.72/sec (d) and 0.675/sec (e). **(F)** Velocities in (D) and (E)
were color coded as indicated.

Figure 8: Loss of function of Level-2 SCPs requires multiple adaptations to allow NOG under
physiological conditions. We predicted a parameter set that allows steady neurite outgrowth with
a velocity of 10 μm/h. To investigate Level-2 SCP dependencies we reduced the activity of the
SCP Vesicle budding at the TGN by reducing the amount of Recruitment Factor 1 without
compensatory increase of the budding rate. In consequence, the activities of multiple Level-2 SCPs
needed to be coordinatively adapted to secure neurite outgrowth under physiological conditions
(as defined by our model constraints). One possible adaptation involves the reduction in the
activity of the SCPs *Membrane Lipid Production*, of *Anterograde Microtubule-Based Vesicle*
Transport (via reduced kinesin receptors), of *Vesicle Exocytosis* (via reduced v-SNARE V) and of
Membrane Protein Production of the involved transmembrane molecules. Since this option
includes a constant cycling rate (i.e. rate of back transported membrane), it causes a reduction in
neurite outgrowth velocity.

Figure 9: Imbalanced SCP dynamics disturb neurite outgrowth. (A-B) We predicted four sets
of kinetic parameters that allow coordinated NOG at the velocities 5, 10, 15 and 20 μm/h. To
simulate the effect of uncoordinated vesicle SCP dynamics we decreased the count and production
rate of v-SNARE V to 25%, 50% and 75% of the original prediction, without modifying any other
parameters. Numerical simulations predict that reduced exocytosis causes an increase in the count
of vesicles in the growth cone cytoplasm (A) without reducing NOG velocity (B). **(C)** We validated
our prediction by analyzing neurite outgrowth after knockdown of the v-SNARE VAMP7. P1 rat
cortical neurons transfected with siRNA against VAMP7 or scrambled siRNAs were stained for
β-III tubulin to document neurite growth across the microgroove of the chamber. In agreement
with our prediction, VAMP7 knock down did not decrease total outgrowth length. It caused a
significant number of physiological growth cones to turn into dystrophic bulbs. **(D)** Analysis of
quantified neurite lengths confirmed that there was no significant difference in outgrowth length.
Shown are average values and standard deviations (three independent experiments). **(E)** Neurites
growing from neurons after VAMP7 knock down showed a significant increase in the number of
dystrophic bulbs, indicative for impaired NOG. **(F)** To simulate the effect of reduced MT
stabilization, we reduced the stabilization rate that turns dynamic into stable MTs to 25%, 50%
and 75% of the original prediction, without changing the predicted values for all other kinetic
parameters. Numerical simulations propose that decreasing stabilization leads to a significant
decrease in NOG outgrowth over the investigated time. **(G)** We validated our predictions by knock
down of the microtubule crosslinking protein MTCL1. MTCL1 knock down caused reduced NOG
outgrowth and dystrophic bulb formation. **(H)** Outgrowth quantification confirms significant
decrease in NOG (3 independent experiments). **(I)** Mtcl1 also significantly increased the number
of dystrophic bulbs. *$p<0.05$, **$p<0.01$, ***$p<0.001$ based on unpaired t-test for all panels.