Combining next-generation sequencing and mtDNA data to uncover cryptic lineages of

Mexican highland frogs

Eugenia Zarza[1], Elizabeth M. Connors[1], James M. Maley[1], Whitney L.E. Tsai[1], Peter Heimes[2],

Moises Kaplan[3] & John E. McCormack[1,*]

[1] Moore Laboratory of Zoology, Occidental College, Los Angeles, CA 90041

[2] Playa Miramar 441, Colonia Marte 08830, Mexico City, Mexico

[3] 1508 Brooklyn Ave, Ann Arbor, Michigan 48104

EZ email: eugenia.zarza@gmail.com

EMC email: beth.connors.94@gmail.com

JMM email: jmaley@oxy.edu

WLET email: wtsai@oxy.edu

PH email: heimesp@hotmail.com

MK email: moises.kaplan@gmail.com

*Corresponding author: mccormack@oxy.edu

Abstract

Recently, molecular studies have uncovered significant cryptic diversity in the Mexican Highlands, leading to the description of many new endemic species. DNA approaches to this kind of species discovery have included both mitochondrial DNA (mtDNA) sequencing and multilocus genomic methods. While these marker types have often been pitted against one another, there are benefits to deploying them together, as linked mtDNA data can provide the bridge between uncovering lineages through rigorous multilocus genomic analysis and identifying lineages through comparison to existing mtDNA databases. Here, we apply one class of multilocus genomic marker, ultraconserved elements (UCEs), and linked mtDNA data to a species complex of frogs (*Sarcohyla bistincta*) found in the Mexican Highlands. We generated data from 1,891 UCEs, which contained 1,742 informative SNPs for *S. bistincta* and closely related species and captured mitochondrial genomes for most samples. Genetic analyses based on both whole loci and SNPs agree there are numerous distinct and divergent lineages within *S. bistincta*. The SNP-based species tree provides the most conservative estimate of 8 well-supported lineages in three major clades. Having linked mtDNA data allowed us to tap into the large number of mtDNA sequences available on GenBank and identify one of these lineages as an already-described species, *S. pentheter*. One identified clade (containing 2 of the 8 lineages) was 10% divergent in mtDNA and paraphyletic with other *S. bistincta*, making this clade a clear candidate for species status. Phylogenies from UCEs and mtDNA mostly agreed in their topologies, but differed in that mtDNA suggested a more complex evolutionary history perhaps influenced by gene flow between some neighboring lineages. Our study demonstrates that the Mexican Highlands still hold substantial undescribed diversity. Combining multilocus genomic data with linked mtDNA data is a useful approach for identifying potential new species and associating them with already described taxa, which is especially important in groups with undescribed subadult phenotypes, where geographic ranges are unclear, or where phenotypes are conserved.

47   Keywords: ultraconserved elements, genomics, population genetics, phylogeography,

48   phylogenetics, systematics, species limits, Hylinae

Introduction

The Mexican Highlands are a global biodiversity hotspot (Myers et al. 2000). Recent molecular studies have uncovered significant cryptic diversity in the Mexican Highlands, leading to the description of new endemic species or the elevation of former subspecies to species rank (Bryson et al. 2010; Bryson et al. 2012; Bryson et al. 2011; McCormack et al. 2008; Rovito et al. 2015). At the same time, habitat loss threatens many of these species before they have even been described (Ponce-Reyes et al. 2012). Amphibians are particularly sensitive to habitat alterations, and many are threatened by habitat loss and invasive diseases (Stuart et al. 2008). Because most amphibians have reduced dispersal, they also show patterns of microendemism (Parra-Olea et al. 2014), meaning that important pockets of diversity and new species are still being uncovered in the Mexican Highlands (Campbell et al. 2009; Meik et al. 2006).

We investigate potential cryptic diversity within the *Sarcohyla bistincta* species complex in the Mexican Highlands. The genus *Sarcohyla*, which is considered distinct from *Plectrohyla* by some authors to reflect those species west of the Isthmus of Tehuantepec (Duellman et al. 2016), contains 24 species of stream-dwelling frogs, many of them critically endangered and many that have never been seen after their original discovery (references compiled in Stuart et al. 2008). Some species are thought to be in serious decline or extinct (Lips et al. 2004). The actual number of species, their relationships, and geographic ranges are not well known (Duellman 2001; Duellman et al. 2016; Faivovich et al. 2009). *S. bistincta* is the most broadly distributed member of the genus. It occurs in several mountain ranges separated by lowland barriers and might therefore be comprised of multiple species.

We sought to assess potential cryptic diversity in *S. bistincta* using multilocus genomic markers collected via next-generation sequencing (NGS) as well as mitochondrial DNA (mtDNA) data. Often, mtDNA and multilocus nuclear markers have been pitted against one another in biodiversity discovery (Edwards & Bensch 2009; Moritz & Cicero 2004; Zink & Barrowclough 2008). Mitochondrial DNA offers many benefits, including economy, efficiency, and wide comparative potential across species and studies. Consequently, there is now a vast

trove of mtDNA sequences on GenBank. A drawback of using only mtDNA is that a single

marker will often fail to accurately depict the speciation process (Edwards & Bensch 2009). In

response, multilocus methods have multiplied (Edwards 2009; Fujita et al. 2012). While multiple

loci help model a more realistic speciation process, multilocus studies suffer from the lack of a

standardized marker set, which limits the ability to link uncovered lineages with species

already identified and described in prior studies and in public databases. This is especially

important in groups with multiple subadult phenotypes (e.g., insects and frogs) and where

adult phenotypes are conserved across species. While many studies have used both types of

markers, few have explicitly explored the benefits of linked mtDNA and NGS data at the level

of the individual for lineage discovery and identification.

For our multilocus genomic markers, we use ultraconserved elements (UCEs). UCEs are

an appealing multilocus marker set because the same loci are found across major branches of

the tree of life, where they act as anchors for variable DNA in flanking regions (Faircloth et al.

2012). For instance, it is possible to capture the same set of 1000 or more UCEs across all

mammals (McCormack et al. 2012), all reptiles (Crawford et al. 2012), or hundreds of UCEs

across arachnid lineages separated by hundreds of millions of years (Starrett et al. 2016). While

the power of UCEs for deep-level systematics is clear, their utility at shallower scales around the

species level is still coming into focus (McCormack et al. 2016; Smith et al. 2014; Zarza et al.

2016). An added benefit of the UCE enrichment process (and all so-called "sequence capture"

methods) is that whole mtDNA genomes are often captured as off-target "bycatch" (do Amaral

et al. 2015), meaning each individual has associated nuclear and mtDNA data (e.g., Zarza et al.

2016). Our specific goals were to determine if UCEs are useful for addressing potential cryptic

diversity within a species complex of frogs. Then, we assess whether linked mtDNA data, by

providing a bridge to Genbank data, allows for more refined conclusions about the

identification of any discovered lineages.

Methods

Sampling and Ingroup Determination

MK and PH collected tadpoles from January to June 2004 across most of the range where *Sarcohyla bistincta* are known to exist (Duellman 2001) in the Transvolcanic Belt of Michoacán, Morelos, and the state of México, the Sierra Madre del Sur of Guerrero, and the highlands of Oaxaca stretching into Puebla and Veracruz (Fig. 1; Table 1). Unsampled parts of the *S. bistincta* range include the far west Transvolcanic Belt in Michoacán and Jalisco, the far northwest in the Sierra Madre Occidental (Nayarit, Durango, and Sinaloa), and the far northeast in Hidalgo (see Fig. S1 for sampled and unsampled locations and known ranges of all *Sarcohyla* species).

Tadpoles were targeted to improve sampling efficiency, which allowed for a larger sampling range and sample density. After collection from a sampling location with a dip net, tadpoles were, to the extent possible, separated by species based on morphology and reared to subadults in the laboratory prior to vouchering. Species identification was based on the most recent diagnosis of *S. bistincta* and other closely related species (Duellman 2001). One tadpole was chosen for the tissue voucher, while the other individuals became physical vouchers with museum catalog numbers. Thus, we provide both field numbers and catalog numbers in Table 1 to provide a link to both the exact genetic material (field number) and the associated phenotype voucher representing that genotype (museum catalog number). Before limiting our taxonomic sampling to 38 individuals in a clade thought to represent *S. bistincta* (as we discuss later, some turned out to be already-described species nested within *S. bistincta*), we ran preliminary phylogenetic analyses including broader sampling of 45 *Sarcohyla* individuals and an outgroup genus *Exerodonta* to ensure we had correctly identified the ingroup (Table S1; Fig. S2).

Figure 1. (A) Map of central Mexico showing sampling sites for *S. bistincta* and close outgroups, with numbers matching localities listed in Table 1 and colors matching Structure results below. Unsampled parts of the distribution of *S. bistincta* are shown in Fig. S1; (B) *S. bistincta* individual from near site 1; (C) Composite results of repeated Structure runs at $K = 2$ showing the finest detectable structure in the genetic data. Each vertical line represents an individual labeled with its UMMZ catalog number above and, in descending order below, the site number and the lineage number based on the SNAPP analysis. One tissue voucher MK 730 (2) could not be linked to a specimen voucher and thus its geographic locality is unknown.

Sequence capture and next-generation sequencing

We extracted genomic DNA from tissue using a Qiagen (Valencia, CA) DNAeasy Blood and Tissue extraction kit. We visualized extractions on an agarose gel to ensure fragments were larger than 200 base pairs (bp) and quantified the resulting double-stranded DNA using a Qubit 2.0 Fluorometer (Carlsbad, CA). For each sample, we sheared 100 μl of 20 ng/μl concentration DNA to a size distribution with its peak between 400 and 600 bp using a Bioruptor ultrasonicator (Diagenode). We prepared libraries for each sheared sample with a KAPA (Boston, MA) LTP library preparation kit for the Illumina platform, attaching custom indexing tags (Faircloth & Glenn 2012) to each sample to allow sample pooling.

We enriched pools of eight samples using a set of synthetic RNA probes that target 5,060 tetrapod UCEs (MYbaits_Tetrapods-UCE-5K kit, Mycroarray) following the standard UCE enrichment protocol (Faircloth et al. 2012) with one modification. Amphibians have large and variable genome sizes with a high percentage of repetitive DNA (Olmo 1991). While we do not have information about the genome size and composition of *Sarcohyla* specifically, we wanted to decrease the potential risk of the probes hybridizing to repetitive elements (McCartney-Melstad et al. 2016). We thus increased by 6X the amount of the Cot-1 blocker, a synthetic DNA derived from chicken that binds to repetitive regions. After enrichment and recovery PCR, we verified the library size range with an Agilent 2100 Bioanalyzer (Palo Alto, CA). We quantified the enriched pools using qPCR and combined them in equimolar ratios before sequencing on an Illumina HiSeq 2000 lane (100-bp paired-end cycle) at the University of California Santa Cruz Genome Technology Center.

Bioinformatics of next-generation sequencing data

We demultiplexed the Illumina raw reads and converted them to FASTQ format with the program bcl2fastq v.1.8.4 (Illumina, Inc.). To eliminate adapter contamination and low quality bases, we trimmed the FASTQC output with illumiprocessor (Faircloth 2013). We trimmed and

assembled these reads into contigs with Trinity (Haas et al. 2013) and ABySS (Simpson et al.

2009), both of which are built into the PHYLUCE pipeline (Faircloth 2015). PHYLUCE uses

LASTZ (Harris 2007) to align all assembled contigs to UCE probes in order to isolate only UCE

contigs and to identify and eliminate paralogs.

Phylogenetic trees from concatenated UCE data

We extracted UCE contigs into a single FASTA file and aligned the output for each locus using

MAFFT (Katoh et al. 2005). We required that 75% of the taxa needed to have data for a given

locus to be included in the final concatenated matrix, which led to dropping loci that did not

meet this threshold. We then constructed a maximum-likelihood (ML) tree in RAxML v8.0.19

(Stamatakis 2014) under the GTRGAMMA model of evolution with 100 bootstrap searches,

followed by a search for the tree with the highest likelihood.

Mitochondrial DNA assembly and analysis

We identified and assembled mtDNA genomes from off-target, trimmed Illumina reads using

the reference genome of a closely related species, *Hyla annectans* (Genebank accession number

KM271781; Ye et al. 2016). We used MITObim 1.7 (Hahn et al. 2013), a Perl wrapper for MIRA

4.0.2 (Chevreux et al. 1999), that takes an iterative mapping approach for assembly. We

conducted *de novo* annotation of the assembled mtDNA regions with the MITOchondrial

genome annotation Server, MITOS (Bernt et al. 2013). We selected for analysis only those

individual genomes with MIRA quality score greater than 30. We aligned each protein-coding

region separately in Geneious vR8 (Kearse et al. 2012) using MUSCLE (Edgar 2004) and

corrected the alignments manually when necessary and constructed a concatenated mtDNA

matrix, which we also ran in RAxML v8.0.19.

We melded this mtDNA data with existing *Sarcohyla* and *Plectrohyla* mtDNA data on

Genbank to determine whether any of the lineages we uncovered in *S. bistincta* relate to already-

described species. We determined that *cytochrome b* is the best-represented mtDNA gene on

Genbank for this group. We downloaded all existing *cytochrome b* sequences from *Sarcohyla* and

Plectrohyla taxa. We combined these sequences with those from a subset of our *S. bistincta*

samples, choosing the individual with the most raw reads from each major genetic lineage in

the UCE tree. We aligned the trimmed, filtered reads for each individual to a *Sarcohyla*

cytochrome b reference sequence and formed a consensus sequence for each individual from the

mapped reads. We then created an alignment and generated a phylogeny using BEAST v2.4.2

(Bouckaert et al. 2014). We repeated this process with 16S data because we suspected *S. pentheter*

was closely related to our ingroup, but *S. pentheter* had no *cyt b* sequence on Genbank.

Calling SNPs from UCE loci

We called SNPs from UCE loci so that we could run genetic clustering tests and infer a species

tree using SNAPP (Bryant et al. 2012), which uses SNPs as input data. Calling SNPs requires a

reference sequence, and we chose the sample with the most UCE contigs recovered within the

ingroup (UMMZ 239727). We then used BWA (Li & Durbin 2009) to map the reads of each

sample to this reference. We used SAMtools (Li et al. 2009) to sort the reads, and Picard

(http://broadinstitute.github.io/picard) to identify and remove PCR duplicates. We realigned

the mapped reads to minimize mismatched bases due to indels, and we removed indels using

the Genome Analysis Toolkit 3.2 (GATK; McKenna et al. 2010) and a custom script

(indelrealigner.sh), as suggested by the Best Practices workflow (DePristo et al. 2011; van der

Auwera et al. 2013).

There is no SNP database available for treefrogs, so we followed best practices for base

recalibration for non-model organisms suggested by GATK (McKenna et al. 2010). This consists

of (1) doing an initial round of calling SNPs on the original, uncalibrated data, (2) selecting the

SNPs with the highest confidence (a minimum emission and call quality of 40 or more), and (3)

using these SNPs as the reference database of known SNPs. We executed four rounds of base

recalibration on the original data to filter out systematic error using a custom script. We called

genotypes on the last recalibrated BAM file. We used vcf-tools (Danecek et al. 2011) to select one

SNP per UCE and produce two data sets, one allowing 25% missing data for STRUCTURE v

2.3.4 (Pritchard et al. 2000), and one with no missing data – a requirement for SNAPP as

implemented in BEAST v2.2.1 (Bouckaert et al. 2014).

STRUCTURE analyses

We used STRUCTURE v2.3.4 (Pritchard et al. 2000) as an unbiased way to assess the limits of

fine-scale genetic structure in our data. Our intent with using STRUCTURE was not to

determine the single most likely number of genetic clusters; thus, we did not use a method for

identifying the "true K" (Evanno et al. 2005), which can underestimate fine-scale population

structure (Janes et al. 2017). Rather, our goal was to determine the maximum possible number of

genetic clusters in our data (e.g., Brown et al. 2007; Gowen et al. 2014). We began by analyzing

all individuals of *S. bistincta* plus two outgroup species *S. chryses* and *S. hazelae* under $K = 4$,

reasoning that this would likely split out the two outgroups as well as reveal at most one

division within *S. bistincta*. After this, each identified genetic cluster within *S. bistincta* was

further analyzed at $K = 2$ until no coherent geographically-based structure was evident in the

plots. All runs were completed twice and each used an admixture model and 10M generations

with 1M generations as burn-in, which led to convergence for all analyses.

SNAPP tree and species delimitation

We generated a species tree from the SNP matrix using SNAPP 1.1.10 (Bryant et al. 2012). This

analysis included putative *S. bistincta* samples (again, later shown to include other species

nested within) and one outgroup, *S. chryses.* For this run, we made no *a priori* assumptions

about how individuals grouped into species and allowed each individual to be considered its

own "species" (i.e., terminal tip). We ran two instances of SNAPP for seven million generations

using default priors. We combined tree and parameter files from both runs with LogCombiner

2.1.3 and displayed the full set of likely species trees with Densitree v2.2.1 (Bouckaert et al.

2014).

Results

NGS summary statistics

Detailed summary statistics for each of the 38 ingroup samples and two outgroups are

described in Table 1. ABySS produced longer contigs than Trinity, and a higher number of UCE

loci, so we used ABYSS contigs in all downstream analyses. Reads per sample ranged from

17,052 to 3,423,330 with an average of 1,185,165 reads. The number of identified UCEs ranged

from 381 to 2,444 with an average of 1,976 UCEs. The mean length of individual UCE loci per

individual ranged from 222 to 717 bp with an average of 522 bp. On average, 18% of the

assembled contigs corresponded to unique UCE loci.

For SNP calling, across 40 samples of *S. bistincta* and outgroups, 9% of the trimmed

reads mapped to our designated reference individual. SNP read depth ranged from 2.4 to 35.0

with an average depth of 21.2. The recalibration and quality control steps resulted in an initial

matrix of 16,578 SNPs. After removing non-biallelic loci, selecting one SNP for every UCE, and

allowing 25% missing data, there were 1,742 SNPs left in the STRUCTURE data set, while the

100% complete data matrix for SNAPP contained 399 SNPs.

UCE phylogeny from concatenated data

Our more taxonomically inclusive data set with all available *Sarcohyla* and outgroup *Exerodonta*

xera (Table S1) contained 1,866 UCE loci and 1,030,450 bp for a concatenated analysis. The

resulting ML tree (Fig. S2) showed strong support for monophyly of *Sarcohyla*, and identified *S.*

arborescandens and *S. cyclada* as sister species that together form a clade sister to the rest of the

Sarcohyla included in the study. We limited further analyses to a smaller data set of 40 samples

with *S. hazelae* as the outgroup (Table 1). This focal data set contained 1,891 UCE loci and

1,038,600 bp.

The ML tree of these 40 samples found strong support for many clades within the

species currently described as *S. bistincta*, conforming to distinct geographic areas (Fig. 2a).

Figure 2. (A) UCE tree; (B) mtDNA tree. Colors match Structure groups identified in Fig. 1.

Tips are labeled with their UMMZ catalog number.

In brief, there were three clades on relatively long branches: (1) a clade distributed across the

Transvolcanic Belt (shaded blue in figures); (2) a clade inhabiting two disjunct areas along the

coastal slopes of the Sierra Madre del Sur in Guerrero and Oaxaca (shaded gray); and (3) a clade

broadly distributed in the Sierra Madre del Sur (shaded red and pink), the Oaxaca Highlands

(shaded yellow and orange), and one individual in the southern portion of the Transvolcanic

Belt (shaded purple). One individual that nested within *S. bistincta* was labeled as a different

species, *S. mykter*, from Guerrero. We suspect that this sample was mislabeled and is actually a

duplicate of an *S. bistincta* sample already included in the study because their field numbers are

similar (last two digits transposed) and the two samples grouped together in all analyses. We

have left this sample in all analyses, but have labeled it as a duplicate of *S. bistincta* UMMZ

239749. Another tissue voucher, MK 730 (2), could not be linked definitively to a physical

voucher, and thus its geographic location is unknown. Its tip label has been left as the field

number. Connecting UCE with mtDNA data, discussed later, revealed that one specimen in the

gray clade (UMMZ 239772) is an already-described species, *S. pentheter*.

mtDNA tree

Our final concatenated mtDNA matrix (individual *cyt b* and 16S tree are discussed later) was

11,269 base pairs including gaps, as coverage of the mtDNA genome varied from sample to

sample in accordance with the non-targeted nature of the DNA collection (Table 1).

Relationships in the ML tree (Fig. 2b) among the 29 individuals with high quality scores were

similar to the concatenated UCE tree with two key differences, both within the broadly

distributed clade in Guerrero and Oaxaca: (1) in the mtDNA tree, individuals from eastern and

western Guerrero (shaded pink and red in figures) formed a clade, whereas they were more

distantly related in the UCE tree; (2) in the mtDNA tree, individual UMMZ 239731 (shaded

purple) was nested within the Guerrero clade instead of being sister to a much more inclusive

clade, as in the UCE tree.

Structure analysis

As expected, the first run of STRUCTURE at K = 4 split the two outgroup species into distinct

clusters and split the remaining individuals into two clusters. Further analysis of each cluster at

K = 2 revealed ten genetic clusters in all (Fig. 1c), which are largely concordant with clades

observed in the UCE and mtDNA phylogeny. Most individuals did not share assignment

among clusters, although a few individuals showed mixed assignment among clusters.

SNAPP species tree

The SNAPP tree and its cloudogram of posterior species trees (Fig. 3) revealed eight well-

supported lineages consistent with many of the genetic clusters in the Structure analysis and

with relationships in the UCE and mtDNA whole-genome phylogenies. With respect to the

discrepancies between UCE and mtDNA trees detailed above, the SNAPP tree agreed with the

mtDNA tree that eastern and western Guerrero individuals form a clade, but agreed with the

UCE tree in the placement of individual UMMZ 239731.

 Figure 3. Cloudogram of the posterior distribution of SNAPP trees from 399 high-quality

SNPs mined from UCE loci. Tip labels are UMMZ catalog numbers. Lineages and clades are

discussed in text. Colors match genetic clusters from Figure 1.

mtDNA phylogeny combining new data with Genbank sequences

Using 16S sequences, we determined that one lineage from Fig. 3 matched an *S. pentheter*

sequence on Genbank. This individual (UMMZ 239772) had one of the lowest read counts of

any samples and very few mtDNA reads. However, five Illumina reads mapped to 16S covering

421 bp of the 681 bp reference sequence (Genbank *S. pentheter* accession number DQ055825).

Over this stretch, UMMZ 239772 was identical to the *S. pentheter* reference. As a point of

comparison, UMMZ 239679 (a member of the blue *S. bistincta* Lineage 1 in the Transvolcanic

Belt) had 70 differences across the 681 bp (10.3% divergence). This DNA identification of

UMMZ 239772 as *S. pentheter* was later confirmed by re-examining the subadult specimen.

After confirming UMMZ 239679 as *S. pentheter*, we generated a Bayesian tree of

cytochrome b combining the samples from this study with Genbank sequences (Fig. 4). This tree

revealed not only that *S. pentheter* was nested within the current *S. bistincta*, but so was another

species not included in our sampling, *S. calthula*. The tree also helped clarify relationships

outside of *S. bistincta* by supporting *S. chryses + S. mykter* to be sister to the *S. bistincta + S.*

pentheter + S. calthula clade.

Figure 4. Phylogeny of mtDNA for a subset of the individuals from this study (labeled with red UMMZ catalog numbers) combined with existing sequences from Genbank (labeled with accession numbers). Colored boxes relate to genetic lineages in prior figures. The tree was rooted with *Exerodonta xera*. Photo credit: Peter Heimes.

Discussion

Bridging multilocus and mtDNA data for cryptic lineage discovery

Multilocus NGS data identified numerous divergent lineages within the *Sarcohyla bistincta* complex, supporting similar patterns observed in a recent study of this complex based on mtDNA and a few nuclear genes (Caviedes-Solis & Nieto-Montes 2017). We discuss these uncovered lineages in detail below, but note here that linking NGS with mtDNA data allowed us to query our lineages – first uncovered with multilocus genomic data – against GenBank to see if any of them corresponded to already-described species. Doing so revealed that *S. bistincta* is paraphyletic, with two already-described species nested within its current taxonomic limits (Fig. 4). The approach of discovering lineages with multilocus data and identifying them with the help of linked mtDNA data should be especially useful in understudied groups where basic natural history information is lacking.

UCEs as a universal genomic marker set for species discovery?

Although it requires further study, UCEs could answer the call for an multilocus DNA marker set that satisfies criteria for both ease-of-use, universality, and genomic coverage (Coissac et al. 2016). UCE probe sets are now available for many taxonomic groups (Faircloth et al. 2015; Faircloth et al. 2013; Starrett et al. 2016). They capture a discrete and replicable portion of the genome, in this case a set of around 2,000 loci in frogs (from a larger set of ~5,000 vertebrate loci) that query approximately 1,000,000 base pairs, or 0.02% of the frog genome. The replicable nature of UCEs sets them apart from other types of genomic markers, like RAD loci, which can vary from experiment to experiment (DaCosta & Sorenson 2014) and find fewer orthologs with increasing phylogenetic distance (Cruaud et al. 2014). Other alternatives to UCEs exist, like exons, but in mammals exons had fewer loci conserved over broad taxonomic scales than UCEs, making them less able to be universally applied (McCormack et al. 2012). A recent study successfully applied UCEs in conjunction with species delimitation methods to two frog genera, *Melanophryniscus* and *Brachycephalus* (Pie et al. 2017). As more studies are conducted, one future research avenue would be determine how much locus-sharing occurs among studies, and whether objective benchmarks for species-level divergence can be identified.

Implications for species limits within Sarcohyla bistincta

The monophyletic group containing *Sarcohyla bistincta* is comprised of many distinct lineages as well as two already-described species (*S. pentheter* and *S. calthula*). These lineages are grouped into three fairly divergent clades, which agree with clades uncovered in a recent study based on fewer markers (Caviedes-Solis & Nieto-Montes 2017). In discussing these clades and lineages below, we use the most conservative estimate of 8 lineages, supported by the species tree, as a framework. Within that framework, we discuss the potential for further genetic structuring, as suggested by genetic clustering results and individual UCE and mtDNA phylogenies. We do not attempt to estimate the "true" number of species through species delimitation or other

methods, because we feel species delimitation is best carried out through integrative taxonomy

(Will et al. 2005), including at minimum analysis of the phenotype.

Clade 1 – Transvolcanic Belt of central Mexico. This clade is sister to the rest of *S. bistincta* plus two other species, *S. pentheter* and *S. calthula*, and is 10% divergent in mtDNA from other *S. bistincta*, making it an obvious candidate for species status. This clade might itself contain multiple species in the form of the lineages below. Unsampled populations of *S. bistincta* in the Sierra Madre Occidental (Fig. S1) are very likely related to Lineage 1 of this clade, but should be included in future studies, as they might be distinctive.

Lineage 1 (light blue in Fig. 1) – Michoacán to western Mexico state. The Structure results show fine-scale genetic structure across this range, and the presence of a geographic and genetic intermediate hints at continuity of gene flow along the distribution from sites 1 to 4 in Figure 1. In addition to unsampled populations mentioned above, some populations in far western Michoacán (Fig. S1) are also unsampled and could reveal further genetic structure.

Lineage 2 (dark blue in Fig. 1) – Morelos. Denser sampling between sites 4 and 5 could help determine whether the genetic distinctness of this individual in the mtDNA and UCE trees is a true discontinuity or the result of a sampling gap.

Clade 2 – Guerrero to Puebla and Veracruz and south through Oaxaca. This clade was also found to be distinct in a recent study (Caviedes-Solis & Nieto-Montes 2017) and forms the core *S. bistincta* (*sensu stricto*).

Lineage 3 (red in Fig. 1) – Central and eastern Guerrero. Members of this lineage are distinct from Lineage 4 and are monophyletic in the UCE-based trees (though not in the mtDNA tree). Further sampling in between site 6 and site 7 would clarify whether the genetic discontinuity between Lineages 3 and 4 results from a sampling gap.

<u>Lineage 4</u> (pink in Fig. 1) – Western Guerrero. This lineage is monophyletic in all trees,
although only a few individuals were sampled from a single locality.

<u>Lineage 5</u> (orange and yellow in Fig. 1) – Puebla, Veracruz, and Oaxaca. This lineage
contains the type locality for *S. bistincta*. Central and southern Oaxaca individuals
(orange) are genetically distinct from individuals to the north (yellow). One genetic
intermediate in central Oaxaca suggests genetic continuity across this range. An
unsampled northern population in Hidalgo is most likely related to this lineage,
and should be included in future studies.

<u>Lineage 6</u> (purple in Fig. 1) – far northern Guerrero. This lineage is distinct but
represented by only a single individual. However, in the mtDNA tree, this
individual is nested within Lineages 4 and 5 above. Sampling more individuals is
needed to determine how distinct this lineage might be.

Clade 3 – Pacific slope of Guerrero and Oaxaca. This clade contains two species: one already
described (*S. pentheter*) and one currently being described (*sp. nov.*; Kaplan et al. in
prep). The relationship of *S. calthula* to this clade is unclear in our results, as
bootstrap support was low in the combined mtDNA tree (Fig. 4). Caviedes-Solis et
al. (2017) placed *S. calthula* sister to *S. pentheter* based on five genes. Thus *S. calthula*
is likely also a member of Clade 3.

<u>Lineage 7</u> (dark gray in Fig. 1) – Pacific slope of Guerrero. This lineage was being
described as a new species (we call it *sp. nov.*) on the basis of phenotypic differences
before this genetic study was begun (Kaplan et al. in prep). Thus, our study lends
support to species status for this lineage.

<u>Lineage 8</u> (light gray in Fig. 1) – *S. pentheter*. Pacific slope of Oaxaca.

The three clades and nearly all of the lineages were distinct in the mtDNA tree as well as the
UCE tree (Fig. 2). The mtDNA tree, however, supports a more complicated history for Lineages
3, 4, and 6 in Guerrero. It is unclear why UCE and mtDNA results differed in this regard, but

some reticulate processes might have influenced the mtDNA genomes of these lineages, perhaps, given their close geographic proximity, mtDNA capture of one lineage by another through gene flow (e.g., Bryson Jr et al. 2010).

Additional insights into broader *Sarcohyla* relationships offered by the mtDNA tree include support for a previously hypothesized close relationship between *S. hazelae* and *S. thorectes* (Caviedes-Solis & Nieto-Montes 2017; Faivovich et al. 2009), and a sister relationship between *S. mykter* and *S. chryses* (also found by Caviedes-Solis et al. (2017)). As *Sarcohyla* is very poorly represented by voucher specimens and DNA sequences (Fig. S1), a complete understanding of the history of this genus must await more complete taxonomic and genomic sampling. Unfortunately, there appear to be some microendemic *Sarcohyla* that might have already gone extinct (Lips et al. 2004), especially in the Oaxacan highlands, although recent resurveys provide hope for rediscoveries (Delia et al. 2013).

This and other recent studies show there is still substantial diversity remaining to be described in the Mexican Highlands. Genetic studies to uncover this diversity might use different approaches and marker types, but these efforts need not be in opposition. As our study shows, NGS and mtDNA data work well together, when lineages uncovered via multilocus methods (but with linked mtDNA data) can be checked against and combined with existing mtDNA databases like Genbank. Identification of vouchered material might not be a problem with some well-studied groups, like birds, but it is invaluable for groups with undescribed larval stages, unclear range boundaries, and difficult or highly conserved phenotypes. As the destruction of native habitats continues apace, it is important that we identify distinctive lineages and geographic centers of diversity before they are lost.

List of abbreviations

UCEs: ultraconserved elements   **SNPs:** single nucleotide polymorphisms   **mtDNA:** mitochondrial DNA **bp:** base pairs **ML:** maximum-likelihood **UMMZ:** University of Michigan Museum of Zoology **RAD loci:** restriction digest-associated loci

Declarations

Ethics approval and consent to participate

This study complied with standard ethical guidelines for the rearing and collection of tadpoles.

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated and analyzed in the current study will be made available on Genbank (BioProject ID PRJNA393258) and Dryad upon publication.

Competing Interests

The authors declare that they have no competing interests.

Funding

The project was partially funded by National Science Foundation grant DEB-1258205 to JM.

Authors' contributions

MK and PH collected samples in the field; MK and JEM designed the study; EZ and WLET produced the genomic data; EZ, EMC, and JMM analyzed the data; JEM and EZ wrote the manuscript; JEM created the figures; All authors read and approved the final manuscript.

Acknowledgments

We thank Greg Schneider and Ron Nussbaum at UMMZ for providing access to tissue samples and vouchers, and Oscar Flores for help with permits.

Literature Cited

Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsch G, Pütz J, Middendorf M, and Stadler PF. 2013. MITOS: Improved de novo metazoan mitochondrial genome annotation. *Molecular Phylogenetics and Evolution* 69:313-319.

Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, Suchard MA, Rambaut A, and Drummond AJ. 2014. BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS Computational Biology* 10:e1003537.

Brown DM, Brenneman RA, Koepfli K-P, Pollinger JP, Milá B, Georgiadis NJ, Louis EE, Grether GF, Jacobs DK, and Wayne RK. 2007. Extensive population genetic structure in the giraffe. *BMC biology* 5:57.

Bryant D, Bouckaert R, Felsenstein J, Rosenberg NA, and RoyChoudhury A. 2012. Inferring species trees directly from biallelic genetic markers: bypassing gene trees in a full coalescent analysis. *Molecular Biology and Evolution* 29:1917-1932.

Bryson Jr RW, De Oca ANM, Jaeger JR, and Riddle BR. 2010. Elucidation of cryptic diversity in a widespread Nearctic treefrog reveals episodes of mitochondrial gene capture as frogs diversified across a dynamic landscape. *Evolution* 64:2315-2330.

Bryson RW, De Oca ANM, Jaeger JR, and Riddle BR. 2010. Elucidation of cryptic diversity in a widespread Nearctic treefrog reveals episodes of mitochondrial gene capture as frogs diversified across a dynamic landscape. *Evolution* 64:2315-2330.

Bryson RW, García-Vázquez UO, and Riddle BR. 2012. Diversification in the Mexican horned lizard *Phrynosoma orbiculare* across a dynamic landscape. *Molecular Phylogenetics and Evolution* 62:87-96.

Bryson RW, Murphy RW, Lathrop A, and Lazcano-Villareal D. 2011. Evolutionary drivers of phylogeographical diversity in the highlands of Mexico: a case study of the *Crotalus triseriatus* species group of montane rattlesnakes. *Journal of Biogeography* 38:697-710.

Campbell JA, Blancas-Hernández JC, and Smith EN. 2009. A new species of stream-breeding treefrog of the genus Charadrahyla (Hylidae) from the Sierra Madre del Sur of Guerrero, Mexico. *Copeia* 2009:287-295.

Caviedes-Solis I, and Nieto-Montes dOA. 2017. A multilocus phylogeny of the genus *Sarcohyla* (Anura: Hylidae), and an investigation of species boundaries using statistical species delimitation. *Molecular Phylogenetics and Evolution* 118:184-193.

Chevreux B, Wetter T, and Suhai S. 1999. Genome sequence assembly using trace signals and
additional sequence information. *Computer Science and Biology: Proceedings of the German*
Conference on Bioinformatics 99:45-56.
Coissac E, Hollingsworth PM, Lavergne S, and Taberlet P. 2016. From barcodes to genomes:
extending the concept of DNA barcoding. *Molecular ecology* 25:1423-1428.
Crawford NG, Faircloth BC, McCormack JE, Brumfield RT, Winker K, and Glenn TC. 2012.
More than 1000 ultraconserved elements provide evidence that turtles are the sister
group of archosaurs. *Biology Letters* 8:783-786. 10.1098/rsbl.2012.0331
Cruaud A, Gautier M, Galan M, Foucaud J, Sauné L, Genson G, Dubois E, Nidelet S, Deuve T,
and Rasplus J-Y. 2014. Empirical assessment of RAD sequencing for interspecific
phylogeny. *Molecular Biology and Evolution* 31:1272-1274.
DaCosta JM, and Sorenson MD. 2014. Amplification biases and consistent recovery of loci in a
double-digest RAD-seq protocol. *PloS one* 9:e106713.
Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,
Marth GT, and Sherry ST. 2011. The variant call format and VCFtools. *Bioinformatics*
27:2156-2158.
Delia JR, Whitney JL, and Burkhardt T. 2013. Rediscovery of 'lost' treefrogs from the Oaxacan
highlands of Mexico. *Biodiversity and conservation* 22:1405-1414.
DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, Del Angel
G, Rivas MA, and Hanna M. 2011. A framework for variation discovery and genotyping
using next-generation DNA sequencing data. *Nature genetics* 43:491-498.
do Amaral FR, Neves LG, Resende Jr MF, Mobili F, Miyaki CY, Pellegrino KC, and Biondo C.
2015. Ultraconserved elements sequencing as a low-cost source of complete
mitochondrial genomes and microsatellite markers in non-model amniotes. *PloS one*
10:e0138446.
Duellman WE. 2001. *Hylid frogs of Middle America*. Ithaca, New York: Society for the Study of
Amphibians and Reptiles.
Duellman WE, Marion AB, and Hedges SB. 2016. Phylogenetics, classification, and
biogeography of the treefrogs (Amphibia: Anura: Arboranae). *Zootaxa* 4104:1-109.
Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic acids research* 32:1792-1797.
Edwards S, and Bensch S. 2009. Looking forwards or looking backwards in avian
phylogeography? A comment on. *Molecular ecology* 18:2930-2933.
Edwards SV. 2009. Is a new and general theory of molecular systematics emerging? *Evolution*
63:1-19.

Evanno G, Regnaut S, and Goudet J. 2005. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular ecology* 14:2611-2620.

Faircloth BC. 2013. illumiprocessor: a trimmomatic wrapper for parallel adapter and quality trimming.https://github.com/faircloth-lab/illumiprocessor/.

Faircloth BC. 2015. PHYLUCE is a software package for the analysis of conserved genomic loci. *Bioinformatics* 32:786-788.

Faircloth BC, Branstetter MG, White ND, and Brady SG. 2015. Target enrichment of ultraconserved elements from arthropods provides a genomic perspective on relationships among Hymenoptera. *Molecular Ecology Resources* 15:489-501.

Faircloth BC, and Glenn TC. 2012. Not all sequence tags are created equal: designing and validating sequence identification tags robust to indels. *PloS one* 7:e42543.

Faircloth BC, McCormack JE, Crawford NG, Harvey MG, Brumfield RT, and Glenn TC. 2012. Ultraconserved elements anchor thousands of genetic markers spanning multiple evolutionary timescales. *Systematic Biology* 61:717-726. 10.1093/sysbio/sys004

Faircloth BC, Sorenson L, Santini F, and Alfaro ME. 2013. A phylogenomic perspective on the radiation of ray-finned fishes based upon targeted sequencing of ultraconserved elements (UCEs). *PloS one* 8:e65923.

Faivovich J, Haddad CF, Garcia PC, Frost DR, Campbell JA, and Wheeler WC. 2009. Systematic review of the frog family Hylidae, with special reference to Hylinae: phylogenetic analysis and taxonomic revision. *Bulletin of the American Museum of Natural History* 294:1-240.

Fujita MK, Leaché AD, Burbrink FT, McGuire JA, and Moritz C. 2012. Coalescent-based species delimitation in an integrative taxonomy. *Trends in Ecology & Evolution* 27:480-488.

Gowen FC, Maley JM, Cicero C, Peterson AT, Faircloth BC, Warr TC, and McCormack JE. 2014. Speciation in Western Scrub-Jays, Haldane's rule, and genetic clines in secondary contact. *BMC evolutionary biology* 14:135.

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B, and Lieber M. 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature protocols* 8:1494-1512.

Hahn C, Bachmann L, and Chevreux B. 2013. Reconstructing mitochondrial genomes directly from genomic next-generation sequencing reads—a baiting and iterative mapping approach. *Nucleic acids research* 41:e129.

Harris RS. 2007. *Improved pairwise alignment of genomic DNA*. PhD Thesis: Pennsylvania State University.

Janes JK, Miller JM, Dupuis JR, Malenfant RM, Gorrell JC, Cullingham CI, and Andrew RL.
2017. The K= 2 conundrum. *Molecular ecology* 14:3594–3602.

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A,
Markowitz S, and Duran C. 2012. Geneious Basic: an integrated and extendable desktop
software platform for the organization and analysis of sequence data. *Bioinformatics*
28:1647-1649.

Li H, and Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler
transform. *Bioinformatics* 25:1754-1760.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, and Durbin
R. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics* 25:2078-2079.

Lips KR, Mendelson Iii JR, Munoz-Alonso A, Canseco-Márquez L, and Mulcahy DG. 2004.
Amphibian population declines in montane southern Mexico: resurveys of historical
localities. *Biological Conservation* 119:555-564.

McCartney-Melstad E, Mount GG, and Shaffer HB. 2016. Exon capture optimization in
amphibians with large genomes. *Molecular Ecology Resources* 16:1084-1094.

McCormack J, Tsai WLE, and Faircloth BC. 2016. Sequence capture of ultraconserved elements
from bird museum specimens. *Molecular Ecology Resources* 16:1189-1203.

McCormack JE, Faircloth BC, Crawford NG, Gowaty PA, Brumfield RT, and Glenn TC. 2012.
Ultraconserved elements are novel phylogenomic markers that resolve placental
mammal phylogeny when combined with species-tree analysis. *Genome Research* 22:746-
754. 10.1101/gr.125864.111

McCormack JE, Peterson A, Bonaccorso E, and Smith TB. 2008. Speciation in the highlands of
Mexico: genetic and phenotypic divergence in the Mexican jay (*Aphelocoma ultramarina*).
Molecular ecology 17:2505-2521.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K,
Altshuler D, Gabriel S, and Daly M. 2010. The Genome Analysis Toolkit: a MapReduce
framework for analyzing next-generation DNA sequencing data. *Genome Research*
20:1297-1303.

Meik JM, Smith EN, Canseco-Márquez L, and Campbell JA. 2006. New species of the
Plectrohyla bistincta group (Hylidae: Hylinae: Hylini) from Oaxaca, Mexico. *Journal of*
Herpetology 40:304-309.

Moritz C, and Cicero C. 2004. DNA barcoding: promise and pitfalls. *PLoS Biology* 2:e354.

Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GA, and Kent J. 2000. Biodiversity
hotspots for conservation priorities. *Nature* 403:853-858.

Olmo E. 1991. Genome variations in the transition from amphibians to reptiles. *J Mol Evol* 33:68-
75.

Parra-Olea G, Flores-Villela O, and Mendoza-Almeralla C. 2014. Biodiversidad de anfibios en
México. *Revista mexicana de biodiversidad* 85:460-466.

Pie MR, Bornschein MR, Ribiero LF, Faircloth BC, and McCormack JE. 2017. Phylogenomic
species delimitation in microendemic frogs Of the Brazilian Atlantic forest. *bioRxiv*:doi:
https://doi.org/10.1101/143735.

Ponce-Reyes R, Reynoso-Rosales V-H, Watson JE, VanDerWal J, Fuller RA, Pressey RL, and
Possingham HP. 2012. Vulnerability of cloud forest reserves in Mexico to climate
change. *Nature Climate Change* 2:448.

Pritchard JK, Stephens M, and Donnelly P. 2000. Inference of population structure using
multilocus genotype data. *Genetics* 155:945-959.

Rovito SM, Parra-Olea G, Recuero E, and Wake DB. 2015. Diversification and biogeographical
history of Neotropical plethodontid salamanders. *Zoological Journal of the Linnean Society*
175:167-188.

Simpson JT, Wong K, Jackman SD, Schein JE, Jones SJ, and Birol I. 2009. ABySS: a parallel
assembler for short read sequence data. *Genome Research* 19:1117-1123.

Smith BT, Harvey MG, Faircloth BC, Glenn TC, and Brumfield RT. 2014. Target capture and
massively parallel sequencing of ultraconserved elements for comparative studies at
shallow evolutionary time scales. *Systematic Biology* 63:83-95.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* 30:1312-1313.

Starrett J, Derkarabetian S, Hedin M, Bryson RW, McCormack JE, and Faircloth BC. 2016. High
phylogenetic utility of an ultraconserved element probe set designed for Arachnida.
Molecular Ecology Resources 17:812-823.

Stuart SN, Hoffmann M, Chanson J, Cox N, Berridge R, Ramani P, and Young B. 2008.
Threatened amphibians of the world. Barcelona, Spain: Lynx Edicions.

van der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, Jordan
T, Shakir K, Roazen D, and Thibault J. 2013. From FastQ data to high-confidence variant
calls: the genome analysis toolkit best practices pipeline. *Current protocols in*
bioinformatics 11: 11.10:11.10. 11-11.10. 33.

Will KW, Mishler BD, and Wheeler QD. 2005. The perils of DNA barcoding and the need for
integrative taxonomy. *Systematic Biology* 54:844-851.

Ye L, Zhu C, Yu D, Zhang Y, and Zhang J. 2016. The complete mitochondrial genome of *Hyla*
annectans (Anura: Hylidae). *Mitochondrial DNA Part A* 27:1593-1594.

Zarza E, Faircloth BC, Tsai WL, Bryson RW, Klicka J, and McCormack JE. 2016. Hidden histories
of gene flow in highland birds revealed with genomic markers. *Molecular ecology*
25:5144-5157.
Zink RM, and Barrowclough GF. 2008. Mitochondrial DNA under siege in avian
phylogeography. *Molecular ecology* 17:2107-2121.

Tables

Table 1. Specimen information and summary statistics for *Sarcohyla bistincta* and closely related

species.

Map Number[1]	Field Number[2]	Catalog Number[3]	Current Taxonomy	State	Latitude	Longitude	Trimmed Paired Reads	UCEs	Average UCE Length	SNP Read Depth	mtDNA Reads	mtDNA Average Coverage	mtDNA Average Quality
1	MK 618	UMMZ 239796	*S. bistincta*	MICHOACAN	19.7911	-100.6605	1,889,670	2,323	529.7	31.8	3,381	20.8	86
2	MK 627-31	UMMZ 239683	*S. bistincta*	MICHOACAN	19.4266	-102.0736	1,933,019	2,254	370.9	21.6	3,415	20.8	85
3	MK 666	UMMZ 239745	*S. bistincta*	MICHOACAN	19.3452	-100.3128	2,186,871	2,350	621.3	33.9	3,900	23.4	83
4	MK 600	UMMZ 239679	*S. bistincta*	MEXICO	19.1501	-100.1469	932,448	2,174	500.8	24.4	2,377	15.7	83
4	MK 600 (1)	UMMZ 239678	*S. bistincta*	MEXICO	19.1501	-100.1469	1,217,987	2,267	532.8	21.9	2,570	16.4	84
5	MK 645	UMMZ 239749	*S. bistincta*	MORELOS	18.9224	-99.2442	2,217,054	2,406	532.5	34.2	3,494	21.8	86
5	MK 645 dup	UMMZ 239749	*S. bistincta*	MORELOS	18.9224	-99.2442	548,545	1,964	513.0	17.7	841	7.4	60
6	MK 759	UMMZ 239701	*S. bistincta*	GUERRERO	18.0013	-101.1716	1,224,835	2,199	557.6	25.0	2,522	16.2	84
6	MK 760	UMMZ 239705	*S. bistincta*	GUERRERO	18.0013	-101.1716	1,074,315	2,203	559.4	24.3	743	6.9	58
6	MK 760 (2)	UMMZ 239704	*S. bistincta*	GUERRERO	18.0013	-101.1716	927,774	2,246	579.2	26.3	906	9.4	6
7	MK 691 (5)	UMMZ 239744	*S. bistincta*	GUERRERO	17.5324	-99.9994	2,084,203	2,353	672.5	27.2	1,678	11.9	80
8	MK 650 (1)	UMMZ 239725	*S. bistincta*	GUERRERO	17.6843	-99.8034	941,101	2,124	556.6	22.6	1,278	9.7	74
8	MK 650 (2)	UMMZ 239726	*S. bistincta*	GUERRERO	17.6843	-99.8034	2,224,898	2,394	506.4	28.4	1,737	12.2	79
8	MK 652	UMMZ 239727	*S. bistincta*	GUERRERO	17.6843	-99.8034	3,423,330	2,444	526.0	29.8	1,396	10.4	74
9	MK 671 (4)	UMMZ 239733	*S. bistincta*	GUERRERO	17.6407	-99.6797	1,012,300	2,107	557.0	22.5	842	7.4	62
9	MK 672	UMMZ 239738	*S. bistincta*	GUERRERO	17.6407	-99.6797	297,782	1,667	439.8	11.9	400	5.1	41
10	MK 656 (1)	UMMZ 239729	*S. bistincta*	GUERRERO	17.5526	-99.6626	580,194	1,950	497.6	18.4	547	5.9	42
11	MK 674 (1)	UMMZ 239690	*S. bistincta*	GUERRERO	17.5087	-99.1258	659,418	1,941	565.4	18.8	1,086	8.7	68
11	MK 675 (2)	UMMZ 239879	*S. bistincta*	GUERRERO	17.5087	-99.1258	336,414	1,565	519.1	10.4	227	4.2	24
12	MK 662	UMMZ 239731	*S. bistincta*	GUERRERO	18.6359	-99.6491	1,137,742	2,130	548.7	23.5	796	7.2	56
13	MK 697 (3)	UMMZ 239789	*S. bistincta*	VERACRUZ	18.6585	-97.1574	1,524,182	2,214	625.6	24.3	2,479	16.0	82
13	MK 699 (1)	UMMZ 239791	*S. bistincta*	VERACRUZ	18.6477	-97.1574	1,342,010	2,206	610.9	24.4	1,769	12.3	79
14	MK 700 (2)	UMMZ 239750	*S. bistincta*	PUEBLA	18.3220	-97.0285	2,203,360	2,298	697.8	27.6	4,155	24.8	86
15	MK 705 (1)	UMMZ 239862	*S. bistincta*	OAXACA	18.1576	-96.8684	2,529,703	2,417	644.6	30.3	3,164	19.6	85
16	MK 715	UMMZ 239755	*S. bistincta*	OAXACA	17.2390	-97.0032	347,152	1,712	437.7	12.1	254	4.3	31
17	MK 716 (1)	UMMZ 239758	*S. bistincta*	OAXACA	17.3036	-96.7930	52,461	594	267.4	2.7	8	3.0	1
18	MK 718 (2)	UMMZ 239765	*S. bistincta*	OAXACA	17.4211	-96.6876	606,978	2,014	568.6	19.7	366	4.9	37
19	MK 755 (1)	UMMZ 239786	*S. bistincta*	OAXACA	17.4153	-96.5671	871,034	2,062	563.2	22.0	1,022	8.4	69
20	MK 751	UMMZ 239785	*S. bistincta*	OAXACA	17.3160	-96.4435	1,577,190	2,342	621.5	28.5	2,376	15.4	84
22	MK 748 (2)	UMMZ 239780	*S. bistincta*	OAXACA	16.9791	-96.1364	91,912	1,009	302.5	4.2	23	3.1	3
22	MK 748 (4)	UMMZ 239782	*S. bistincta*	OAXACA	16.9791	-96.1364	1,058,426	2,170	553.8	23.7	499	5.7	42
22	MK 767	UMMZ 239788	*S. bistincta*	OAXACA	16.9859	-96.1358	1,083,058	2,206	524.7	24.3	617	6.3	52
23	MK 721	UMMZ 239767	*S. bistincta*	OAXACA	16.7377	-97.0384	146,471	1,215	376.0	6.1	62	3.3	10
24	MK 766	UMMZ 239794	*S. bistincta*	OAXACA	16.2522	-97.1536	994,521	2,166	577.4	23.3	467	5.5	43
	MK 730 (2)	?	*P. bistincta*	?	?	?	2,967,630	2,404	716.9	31.1	3,581	21.7	84
25	MK 685 (2)	UMMZ 239739	*S. sp. nov.*	GUERRERO	17.3812	-100.2009	269,729	1,645	450.6	10.8	184	4.0	20
26	MK 689 (2)	UMMZ 239740	*S. sp. nov.*	GUERRERO	17.3000	-100.2792	67,313	502	221.6	2.6	32	3.1	6
27	MK 727 (2)	UMMZ 239772	*S. pentheter*	OAXACA	16.1916	-97.0958	17,052	381	247.1	2.4	10	3.0	2
28	MK 691 (3)	UMMZ 239651	*S. chryses*	GUERRERO	17.5324	-99.8994	2,267,946	2,347	663.6	35.0	4,919	33.4	83
29	MK 770	UMMZ 239802	*S. hazelae*	OAXACA	17.2216	-96.5839	538,500	2,066	526.2	19.8	557	5.9	44

[1] Map number in Figure 1
[2] The first three-digit number corresponds to a sampling location. If there is a second number in parentheses, this corresponds to different aquaria where tadpoles were sorted by species before
[3] All specimens are from the University of Michigan Museum of Zoology. One specimen could not be linked to a catalog number.

Table S1. Information and summary statistics on all 45 samples used to determine the ingroup

for this study.

Field Number	UMMZ Number	Current Taxonomy	State	Locality	Latitude	Longitude	Trimmed Paired Reads	UCEs	Average UCE length	mtDNA reads	mtDNA average coverage	mtDNA average quality	Fig. 1 Map Number	Accession number
MK 618	UMMZ 239796	*Sarcohyla bistincta*	MICHOACAN	LOS AZUFRES / SAN PEDRO ROAD	19.791051	-100.660542	1,889,670	2323	529.7	3381	20.8	86	1	
MK 627-31	UMMZ 239683	*Sarcohyla bistincta*	MICHOACAN	PARQUE BARRANCA DEL CUPATITZIO URUAPAN	19.426646	-102.073574	1,933,019	2254	370.9	3415	20.8	85	2	
MK 666	UMMZ 239745	*Sarcohyla bistincta*	MICHOACAN	13 KMS ROAD ZITACUARO MACHERO (LITTLE TOWN WERE THERE IS A MONARCH BUTTERFLY SANCTUARY)	19.3452	-100.3128	2,186,871	2350	621.3	3900	23.4	83	3	
MK 600	UMMZ 239679	*Sarcohyla bistincta*	MEXICO	CARRETERA VALLE DE BRAVO-SAN PEDRO TENEYAC ARROYO 1 (6860 FEET) (POR ALBARRADA) BOSQUE BIEN CONSERVADO Y HUMEDO	19.150083	-100.1469	932,448	2174	500.8	2377	15.7	83	4	
MK 600 (1)	UMMZ 239678	*Sarcohyla bistincta*	MEXICO	CARRETERA VALLE DE BRAVO-SAN PEDRO TENEYAC ARROYO 1 (6860 FEET) (POR ALBARRADA) BOSQUE BIEN CONSERVADO Y HUMEDO	19.150083	-100.1469	1,217,987	2267	532.8	2570	16.4	84	4	
MK 645	UMMZ 239749	*Sarcohyla bistincta*	MORELOS	2 KMS OUT OF CUERNAVACA ON THE CUERNAVACA CHALMA RD. (PASSING CONJUNTO CERRADO) OF LA BARRANCA WHERE THE SALTO SAN ANTONIO IS	18.922402	-99.244151	2,217,054	2406	532.5	3494	21.8	86	5	
MK 645 dupe	UMMZ 239749	*Sarcohyla bistincta*	MORELOS	2 KMS OUT OF CUERNAVACA ON THE CUERNAVACA CHALMA RD. (PASSING CONJUNTO CERRADO) OF LA BARRANCA WHERE THE SALTO SAN ANTONIO IS	18.922402	-99.244151	548,585	1964	513.0	841	7.4	60	5	
MK 759	UMMZ 239701	*Sarcohyla bistincta*	GUERRERO	CA 100 DEG KMS FROM CIUDAD ALTAMIRANO VIA IXTAPA ZIHUATANEJO	18.0013	-101.1716	1,224,835	2199	557.6	2522	16.2	84	6	
MK 760	UMMZ 239705	*Sarcohyla bistincta*	GUERRERO	CA 100 DEG KMS FROM CIUDAD ALTAMIRANO VIA IXTAPA ZIHUATANEJO	18.0013	-101.1716	1,074,315	2203	559.4	743	6.9	58	6	
MK 760 (2)	UMMZ 239704	*Sarcohyla bistincta*	GUERRERO	CA 100 DEG KMS FROM CIUDAD ALTAMIRANO VIA IXTAPA ZIHUATANEJO	18.0013	-101.1716	927,774	2246	579.2	906	9.4	6	6	
MK 691 (5)	UMMZ 239744	*Sarcohyla bistincta*	GUERRERO	2-3 KMS IN THE ROAD TO JALEACA FROM POINT WHERE ROAD TRIFURCATE TO PTO. DEL GALLO / YEXTLA / ANDJALEACA.COMMING FROM CARRIZAL DE BRAVO / IN RIVER UNDER THE BRIDGE	17.5324	-99.8994	2,084,203	2353	672.5	1678	11.9	80	7	
MK 650 (1)	UMMZ 239725	*Sarcohyla bistincta*	GUERRERO	MOUNTAINS W. OF CHILPANCINGO / TOWN LOS MORROS	17.6843	-99.80339	941,101	2124	556.6	1278	9.7	74	8	
MK 650 (2)	UMMZ 239726	*Sarcohyla bistincta*	GUERRERO	MOUNTAINS W. OF CHILPANCINGO / TOWN LOS MORROS	17.6843	-99.80339	2,224,898	2394	506.4	1737	12.2	79	8	
MK 652	UMMZ 239727	*Sarcohyla bistincta*	GUERRERO	MOUNTAINS W. OF CHILPANCINGO / TOWN LOS MORROS	17.6843	-99.80339	3,423,330	2444	526.0	1396	10.4	74	8	
MK 671 (4)	UMMZ 239733	*Sarcohyla bistincta*	GUERRERO	BEHIND CHICHIHUALCO / ON ROAD TO CARRIZAL DE BRAVO / 2 KMS FROM ENTRONQUE	17.6407	-99.6797	1,012,300	2107	557.0	842	7.4	62	9	
MK 672	UMMZ 239738	*Sarcohyla bistincta*	GUERRERO	BEHIND CHICHIHUALCO / ON ROAD TO CARRIZAL DE BRAVO / 2 KMS FROM ENTRONQUE	17.6407	-99.6797	297,782	1667	439.8	400	5.1	41	9	
MK 656 (1)	UMMZ 239729	*Sarcohyla bistincta*	GUERRERO	3 KMS FROM THE TOWN OF OMILTEMI IN THE ROAD OMILTEMI CHILPANCINGO	17.552603	-99.662569	580,194	1950	497.6	547	5.9	42	10	
MK 674 (1)	UMMZ 239690	*Sarcohyla bistincta*	GUERRERO	ON ATZACUALOYA HUEYCATENANGO RD.	17.5087	-99.1258	1,086	1941	565.4	1086	8.7	68	11	
MK 675 (2)	UMMZ 239879	*Sarcohyla bistincta*	GUERRERO	ON ATZACUALOYA HUEYCATENANGO RD.	17.5087	-99.1258	336,474	1565	519.1	227	4.2	24	11	
MK 662	UMMZ 239731	*Sarcohyla bistincta*	GUERRERO	1.6 KMS FROM THE TOWN OF TETIPAC ON THE TETIPAC TAXCO ROAD (MAYBE ARROYO LAS DAMAS)	18.635895	-99.6491	1,137,742	2130	548.7	796	7.2	56	12	
MK 697 (3)	UMMZ 239789	*Sarcohyla bistincta*	VERACRUZ	ON ATZOMPA XOXOCOTLA RD. 1.5 KMS FROM XOXOCOTLA	18.6585	-97.1574	1,524,182	2214	625.6	2479	16.0	82	13	
MK 699 (1)	UMMZ 239791	*Sarcohyla bistincta*	VERACRUZ	ON STREAM CROSSING THE TOWN OF XOXOCOTLA	18.6477	-97.1574	1,342,010	2206	610.9	1769	12.3	79	13	
MK 700 (2)	UMMZ 239750	*Sarcohyla bistincta*	PUEBLA	IN THE STREAM LOCATED AFTER THE TOWN OF ZOQUITLAN TURNING DOWN AT THE CENTRO DE SALUD	18.322	-97.0285	2,203,360	2298	697.8	4155	24.8	86	14	
MK 705 (1)	UMMZ 239862	*Sarcohyla bistincta*	OAXACA	56 KMS FROM TEOTITLAN VIA HUAUTLA	18.1576	-96.8684	2,529,703	2417	644.6	3164	19.6	85	15	
MK 715	UMMZ 239755	*Sarcohyla bistincta*	OAXACA	NEAR "EL TEJOCOTE" (FRENTE AL KINDER)	17.239	-97.0032	347,152	1712	437.7	254	4.3	31	16	
MK 716 (1)	UMMZ 239758	*Sarcohyla bistincta*	OAXACA	ON ROAD BETWEEN THE TOWNS OF SAN JUAN DEL ESTADO AND SAN MIGUEL ALOAPAN	17.3036	-96.793	52,461	594	267.4	8	3.0	1	17	
MK 718 (2)	UMMZ 239765	*Sarcohyla bistincta*	OAXACA	PASSING SAN MIGUEL ALOAPAN	17.4211	-96.6876	606,978	2014	568.6	366	4.9	37	18	
MK 755 (1)	UMMZ 239786	*Sarcohyla bistincta*	OAXACA	ON ROAD SAN JUAN ATEPEC SAN MIGUEL ABEJONES	17.4153	-96.5671	871,034	2062	563.2	1022	8.4	69	19	
MK 751	UMMZ 239785	*Sarcohyla bistincta*	OAXACA	3.8 KMS PASSING "RANCHO TEXAS" ON THE ROAD FROM THE TOWN OF IXTLAN DE JUAREZ	17.316	-96.4435	1,577,190	2342	621.5	2376	15.4	84	20	
MK 748 (2)	UMMZ 239780	*Sarcohyla bistincta*	OAXACA	CA 37 KMS FROM MITLA ON THE ROAD MITLA AYUTLA	16.9791	-96.1364	91,912	1009	302.5	23	3.1	3	22	
MK 748 (4)	UMMZ 239782	*Sarcohyla bistincta*	OAXACA	CA 37 KMS FROM MITLA ON THE ROAD MITLA AYUTLA	16.9791	-96.1364	1,058,426	2170	553.8	499	5.7	42	22	
MK 767	UMMZ 239788	*Sarcohyla bistincta*	OAXACA	9.2 KMS E STA MARIA ALBARRADAS / SIERRA MIXE	16.985888	-96.135816	1,083,058	2206	524.7	617	6.3	52	22	
MK 721	UMMZ 239767	*Sarcohyla bistincta*	OAXACA	2 KMS N. THE TOWN OF STA. MARIA LAXICHIO VIA THE TOWN OF SAN SEBASTIAN RIO DULCE	16.7377	-97.0384	146,471	1215	376.0	62	3.3	10	23	
MK 766	UMMZ 239794	*Sarcohyla bistincta*	OAXACA	CERRO DE VIDRIO VIA A PUERTO ESCONDIDO	16.25216	-97.15359	994,521	2166	577.4	467	5.5	43	24	
MK 730 (2)	?	*Sarcohyla bistincta*	?	?			2,967,630	2404	716.9	3581	21.7	84		
MK 685 (2)	UMMZ 239739	*Sarcohyla sp. nov.*	GUERRERO	ON ATOYAC PTO DE GALLO RD / BETWEEN 10 - 20 KMS NORTH OF THE TOWN "EL PARAISO"	17.3812	-100.2009	269,729	1645	450.6	184	4.0	20	25	
MK 689 (2)	UMMZ 239740	*Sarcohyla sp. nov.*	GUERRERO	ON ATOYAC PTO DEL GALLO RD. / 500 M. NORTH OF THE TOWN OF SAN VICENTE	17.3	-100.2792	67,313	502	221.6	32	3.1	6	26	
MK 727 (2)	UMMZ 239772	*Sarcohyla pentheter*	OAXACA	RIO "EL SALADO" / 8 KMS N SAN JUAN LACHAO ON HWY 135	16.1916	-97.0958	17,052	381	247.1	10	3.0	2	27	
MK 691 (3)	UMMZ 239651	*Plectrohyla chryses*	GUERRERO	2-3 KMS IN THE ROAD TO JALEACA FROM POINT WHERE ROAD TRIFURCATE TO PTO. DEL GALLO / YEXTLA / ANDJALEACA.COMMING FROM CARRIZAL DE BRAVO / IN RIVER UNDER THE BRIDGE	17.5324	-99.8994	2,267,946	2347	663.6	4919	33.4	83	28	
MK 770	UMMZ 239802	*Plectrohyla hazelae*	OAXACA	EL PUNTO SIERRA JUAREZ	17.22156	-96.58386	538,500	2066	526.2	557	5.9	44	29	
MK 667	UMMZ 239952	*Sarcohyla arborescandens*	VERACRUZ	PUERTO DEL AIRE (ARRIBA DE ALCUTZINGO)	18.6787	-97.3485	923,330	2116	511.8	1154	9.0	69		
MK 700 (1)	UMMZ 239813	*Sarcohyla cyclada*	PUEBLA	IN THE STREAM LOCATED AFTER THE TOWN OF ZOQUITLAN TURNING DOWN AT THE CENTRO DE SALUD	18.322	-97.0285	2,490,616	2385	661.4	2961	18.4	82		
MK 701	UMMZ 239814	*Sarcohyla cyclada*	OAXACA	24 KMS FROM THE TOWN OF TEOTITLAN DE FLORES MAGON VIA HUAUTLA DE JIMENEZ	18.1781	-97.0054	1,166,574	2129	565.5	1061	8.7	69		
MK 742 (1)	UMMZ 239954	*Sarcohyla arborescandens*	VERACRUZ	LEFT ROAD BIFURCATING FROM ROAD TO THE TOWN OF "LAS MINAS"	19.6758	-97.1751	1,074,225	2240	553.8	1750	12.0	79		
MK 768	UMMZ 239833	*Exerodonta xera*	PUEBLA	5 KM SW ZAPOTITLAN DE SALINAS	18.311958	-97.51266	1,257,938	2288	581.5	3506	21.4	84		

Additional Files

Fig. S1. Sampled and unsampled parts of *S. bistincta* range in relation to known distributions (or localities, where distributional information is lacking) of other *Sarcohyla* species.

698   Fig. S2. UCE tree of 45 samples of *Sarcohyla* and outgroup *Exerodonta xera* used to determine the

699   ingroup.