Whole genome amplification and sequencing of low cell numbers directly from a

bacteria spiked blood model

Catherine Anscombe [*1,2], Raju.V Misra[1] and Saheer Gharbia[1]

[1] Genomics Research Unit, Public Health England, London, United Kingdom

[2] Oxford University Clinical Research Unit, Ho Chi Minh City, Vietnam

* Corresponding Author: Catherine Anscombe, the Hospital for Tropical Diseases, 764 Vo Van Kiet, Dist.
5, Ho Chi Minh City, Vietnam. canscombe@oucru.org

Abstract

Whilst next generation sequencing is frequently used to whole genome sequence bacteria from
cultures, it's rarely applied directly to clinical samples. Therefore, this study addresses the issue of
applying NGS microbial diagnostics directly to blood samples. To demonstrate the potential of direct
from blood sequencing a bacteria spiked blood model was developed. Horse blood was spiked with
clinical samples of *E. coli* and *S. aureus*, and a process developed to isolate bacterial cells whilst
removing the majority of host DNA. One sample of each isolate was then amplified using ϕ29
multiple displacement amplification (MDA) and sequenced. The total processing time, from sample to
amplified DNA ready for sequencing was 3.5 hours, significantly faster than the 18-hour overnight
culture step which is typically required. Both bacteria showed 100% survival through the processing.
The direct from sample sequencing resulted in greater than 92% genome coverage of the pathogens
whilst limiting the sequencing of host genome (less than 7% of all reads). Analysis of *de novo*
assembled reads allowed accurate genotypic antibiotic resistance prediction. The sample processing is
easily applicable to multiple sequencing platforms. Overall this model demonstrates potential to
rapidly generate whole genome bacterial data directly from blood.

Introduction

Bacterial sequencing of clinical isolates has been used in many settings, including for virulence determinants[1–4], population structure[5][6] and outbreak investigations [7–10]. Almost exclusively current applications of sequencing rely either upon culture or targeted amplification imparting a diagnostic bias towards known pathogens and bacteria with known and relatively simple growth requirements. Bacterial whole genome sequencing is most often performed at reference or research centres to study population structures and evolution, with results rarely available in a clinically actionable time. One exception to this is the application of next generation sequencing to the diagnosis of tuberculosis[11,12], where WGS was shown, in some cases, to be quicker than current routine methods for predicting resistance; however this process still relies on culturing of the bacteria prior to sequencing.

Direct from sample sequencing of pathogens using an untargeted (metagenomic) approach has the potential to lower turnaround times and lower diagnostic bias. Metagenomic sequencing techniques have already been applied to ecological studies [13–16], gut microbiomes [17–19] and investigation and identification of viruses [20–23].

Bacteraemia is a major global cause of morbidity and mortality, with a large range of aetiologies and is a particular problem in healthcare settings[24,25]. The current microbiological diagnostic process is culture based often involving specialised equipment, with time to positivity varying due to aetiology and pathogen load. Although recent advances in MALDI-TOF processing has allowed direct identification of bacteria from positive blood cultures [26,27,28], antibiotic sensitivities take a further 18 hours from pure bacterial cultures. The direct application of whole genome sequencing to blood samples would allow rapid pathogen diagnosis, along with simultaneous pathogen typing and genotypic resistance prediction. Furthermore, by applying an unbiased pathogen detection method the diagnostic bias would be lowered.

A two-stage host cell removal was performed, firstly red blood cells were removed using HetaSep®, previously used to isolate nucleated cells in the blood, particularly granulocytes. Secondly a selective white blood cell lysis was undertaken using saponin[29] in order to release intracellular pathogens and aid in host nucleic acid removal.

To directly sequence pathogens, present at low levels (as few as 1 bacterial cell per ml blood) such as those found in sterile site infections an unbiased and high-fidelity amplification enzyme is needed. Multiple displacement amplification (MDA) using ϕ 29 is an alternative method to PCR for the production of DNA in high enough amounts for sequencing. ϕ 29 MDA has been applied to samples with very low starting DNA amounts from single cells (both prokaryotic and eukaryotic) and provided DNA in levels high enough to perform sequencing [30,31]. It has been demonstrated that this method is

60　suitable for use with single bacterial cells[30,31] and more recently malaria parasites directly from blood

61　samples [32]. The aim of this study was to devise a process to allow same day untargeted bacterial

62　sequencing to be performed on DNA directly isolated from a bacteria spiked blood model.

Methods

Isolates

65　Clinical isolates of *E. coli* and *S. aureus* were collected from the Royal Free Hospital Hampstead,

66　where they had been stored at -80°C after isolation from septic patients. *E. coli* and *S. aureus* are

67　common aetiologies of blood stream infections, representing different cell wall types and cell

68　morphologies. Phenotypic data was produced using the BD phoenix and was retrieved from final

69　hospital reports. Bacteria were cultured on Horse blood agar at 37°C overnight.

Spiking and isolation process

Spiked samples were set up by adding an estimated ten bacterial cells of the clinical isolate of *S.*

aureus or *E. coli* calculated using serial dilutions to 1ml horse blood. The workflow depicted

in**Error! Reference source not found.** Figure 1 was then applied, 200 μl HetaSep® was added and

the sample vortexed and incubated at 37°C for 10 minutes. 550 μl supernatant was removed and 200

μl 5% saponin was added to a final 2% solution, and incubated at room temperature for 5 minutes.

700μl sterile water was added for a water shock and incubated at room temperature for 30 seconds

before salt restoration with the addition of 21μl 5M NaCl. The sample was centrifuged at 4000xg for

5 minutes and the supernatant discarded before addition of 2 μl turbo DNase1 and 5 μl of 10x buffer

(Ambion). The sample was vortexed and incubated at 37°C for 15 minutes EDTA was added to a final

concentration of 15nM. The sample was then centrifuged for 5 minutes at 4000xg and the supernatant

removed and discarded. The pellet was washed in decreasing volumes of PBS, initially 200 μl then

100 μl followed by 20 μl with each stage being centrifuged at 6000xg for three minutes. The final

pellet was suspended in 4 μl sterile PBS.

Bacterial survival

85　To check for bacterial survival at each stage of processing 13 spiked samples were set up and

86　processed. At the end of each processing stage (Table 1) one sample was cultured and colony counts

87　performed. This was repeated three times for the *E. coli* and *S. aureus* isolate.

DNA extraction and Amplification

Extraction was performed using alkaline method; briefly cell suspensions were added to 200 mM

potassium hydroxide (Qiagen) and 50mM dithiothreitol (Qiagen) and incubated at 65°C for 10

minutes. The reaction was then neutralised using neutralisation buffer (Qiagen), briefly vortexed and

placed on ice.

Amplification was performed using φ29 MDA (Repli-g Single Cell Kit Qiagen). A master mix was

prepared on ice in a total volume of 40 µl, with 29 µl reaction buffer, containing endonuclease

resistant hexamer primers and 2 µl (40 U) of φ29 polymerase (Qiagen, REPLI-g Single Cell Kit). The

extracted DNA was then added to the master mix and the sample was then incubated at 30^0C for 2

hours, and the reaction stopped by heating to 65^0C for 3 minutes.

Sequencing

DNA was quantified using Qubit BR kit and 3 µg of φ29 MDA DNA was de-branched using S1

nuclease in a 90 µl reaction as follows, 3 µl 10x buffer, 3 µl 0.5M NaCl, 10 µl S1 nuclease (1U/ µl)

with water to make the volume to 90µl. The digestion reaction was left at room temperature for 30

minutes and the enzyme deactivated by incubating at 70ºC with 6 µl 0.5M EDTA. The DNA was then

fragmented by nebulisation at 30psi for 180 seconds. The DNA was sequenced using the 454 Junior

using the manufacturer's recommended methods.

Data analysis

Data analysis was performed in three stages, first host and contaminating reads were removed before

abundance trimming, secondly reads were classified using lowest common ancestor (LCA) analysis

and finally classified reads were assembled and analysed.

Initially reads were mapped using Newbler (Roche diagnostics) standard parameters against host and

a local contamination library consisting of sequenced negative controls and reads identified as

contamination in previous runs. Unmapped reads were written into a new fastq file using a custom

python script and taken forward for further analysis. The remaining reads were abundance trimmed

using Khmer[33] using two passes to a maximum depth of 50, to remove over represented reads

produce by the φ29 MDA method. Following this the reads were error trimmed using Prinseq[34], with

ends being trimmed to a Q20 cut off.

Blastn was then used to assign the reads against the NCBI nt database. Once completed lowest

common ancestor (LCA) analysis was performed using MEGAN[35], and reads associated with the

bacterial species of interest were extracted to a new fastq file. These reads were then *de novo*

assembled using standard parameters in SPAdes[36] assembly outputs were assessed using QUAST

(Quality Assessment Tool for Genome Assemblies)[34]. Reference assemblies were performed using the

closest reference sequence identified using the LCA analysis. Antibiotic prediction was performed

using Mykrobe[11] for *S. aureus* and ResFinder[37] for *E. Coli* using the *de novo* assembled reads.

Results

Sample processing

When the full blood processing method was applied to *E. coli* and *S. aureus*, good survival rates were

found for both bacteria through-out all stages. Details of survival at each stage of processing can be

found in Table 1

Sequencing results S. aureus

After processing and DNA amplification the concentration of DNA was 139ng / µl , after sequencing

the number of reads passing basic filter was 128,500. Once known contaminants were removed and

error and abundance trimming complete 124,145 reads remained. After reads were assigned using

Blastn and LCA analysis performed using MEGAN, 92.3% of reads were classified. 72.22% of reads

were classified as Staphylococcus species and 62.1% reads were identified as *S. aureus*. 6.76% of

the reads were identified as the genus Equus, and 1.72% were identified as *Parascaris equorum.*

Remaining reads were not classified above the Class level in LCA analysis. When examining the *S.*

aureus reads closer 4254 reads were identified to the subspecies level, (*Staphylococcus aureus subsp.*

aureus HO 5096 0412), this subspecies has a complete genome available (GenBank

GCA_000284535.1) and was used as a reference for reference mapping. Reference mapping covered

92% of the reference genome. When the reads identified as *Staphylococcus* using the LCA analysis

were extracted and the reads *de novo* assembled 1212 contigs were produced. When this was

compared to the same reference as the reference assembly 83% of the genome was covered with 10

misassemblies and a N50 of 3882. Mykrobe analysis was performed using the *de novo* assembly gave

genotypic result for 12 antibiotics. When comparing the genotypic and phenotypic (BD Phoenix™)

results matched in 11 of the 12 antibiotics (Table 2). The results for ciprofloxacin were inconclusive

in genotypic tests, but resistant by phenotypic methods.

Sequencing results E. coli

After processing and amplification, the DNA concentration of the sample was 143 ng/ µl. Post

sequencing 173,597 reads passed the initial filter, once contaminants were removed and the reads

error and abundance trimmed 170,243 reads remained. Overall 73% of all reads remaining after the

analysis pipeline were identified as *E. coli.* 31959 (18%) reads had no identity. Unlike *S. aureus* it

was not possible to type the *E. coli*, as 150 reads were assigned to O7:K1 and 211 reads were assigned

to JJ1886. Reads which were identified as *Enterobacteriaceae, Escherichia* and *E. coli* by BLAST

and LCA analysis were extracted from the fastq produced after pipeline completion. The genome of

JJ1886 was available from the Integrated Microbial Genomes database (ID 2558309052), and the

chromosomal sequence was used as the reference against which the extracted reads were assembled.

After reference mapping assembly 93.5% of the genome was covered. When the reads were *de novo*

assembled, 89% of the same reference was covered in 1334 contigs. The *de novo* assembly was used

to identify several resistance markers using ResFinder. The resistance markers included *dfrA*,

conferring trimethoprim resistance, *gyrA* conferring resistance to fluoroquinolones. *mdtK* is an efflux

pump conferring resistance to norfloxacin and *blaCMY-2* which is an AmpC, conferring resistance to

beta-lactams including cephalosporins. Additionally, eight drug efflux systems were identified. These

results were concordant to the phenotypic data (Table 2).

Discussion

Direct diagnostics using WGS from clinical samples would, in many ways, provide the ideal

diagnostic method. By providing all the information from whole genome data with the speed of direct

sample testing. There are increasing references to using metagenomic approaches to detect pathogens

directly from clinical samples. A recent study, using the Oxford Nanopore Technologies MinION to

sequence directly from TB smear positive respiratory samples, demonstrated it was possible to get

phenotypically concordant results from sequencing in 12.5 hours[38] . Direct sequencing from

orthopaedic device infections showed that a metagenomic sequencing approach gave species level

specificity and sensitivity of 88% compared to traditional culture techniques[39]. In a study of ocular

eye fluid samples, it was possible to match 9/12 bacteria using metagenomic sequencing compared to

traditional culture techniques, with the additional advantage of being able to obtain sequence types

and antibiograms for some of the identified pathogens[40]. Whilst these studies provide good evidence

of the application of metagenomic sequencing, applying these methods directly to blood still has

various are various obstacles, including low pathogen numbers, high host background and difficulty

in interpreting genotypic data.

Here, a model was prepared to demonstrate the potential for direct from sample sequencing from

whole blood. This method allows DNA to be ready within 3.5 hours of sample receipt, including

bacterial isolation and DNA amplification. Once the DNA is prepared it could be processed on any

sequencing platform. Final time to results would depend on the platform used, and the scale of

batching required to make the method cost effective. A recent study looked at real time analysis of

sequencing produced on the Oxford Nanopore MinION platform, showed it was possible to identify

the bacterial species in a sample within 30 minutes of starting the run. Initial resistance predictions

were ready within two hours, and all analysis was completed within 10 hours[41]. Combining good

sample preparation and real-time analysis could allow results to be obtained in a clinically actionable

time frame. In addition to the wet lab model, an analysis pipeline was developed to allow accurate

interpretation of the results. This included removal of contamination, normalisation of reads and LCA

analysis. Reference mapping and *de novo* assembly programmes were selected to give the best results

for the data produced, in terms of amplification using ϕ29 MDA and sequencing on the 454 Junior.

Fresh whole horse blood was used to model bacteraemia, as it was readily available. The process was

developed to remove RBCs early in the process, as they represent the largest proportion of the

cellular makeup of blood (up to 96%[42]), debulking the sample, and preventing release of oxidative

agents. Selective lysis on WBCs allowed the release of any intracellular pathogens and exposed host

nucleic acid to nucleases. *S. aureus* and *E. coli* were chosen due to different cell wall types and cell

morphology; both demonstrated good survival during the developed sample processing.

The use of φ29 MDA was selected due to its ability to quickly and accurately amplify DNA

without the need for specific primers, allowing its application to a wide range of pathogens. φ29

MDA has been associated with amplification bias, especially when starting from low input samples,

previously this was thought to be random, but has now been shown to be systemic[43]. Lower reaction

volumes have been shown to lower the effect of the bias, however significant scaling down from the

current 5 μl starting material would be technically difficult. Use of digital droplet MDA has been

shown to give a more uniform coverage across the genome and lower none specific amplifcation[44],

however this requires specialised equipment. For this study amplification bias was compensated for

bioinformatically, using a process of digital normalisation[33], which removed over represented reads,

and added an additional error removal stage by removing very low level k-mers. This aided in

assembly of reads, as most assemblers assume even coverage across the genome, it also allowed the

data size to be reduced without loss of information. Despite the bias in the amplification, good

genome coverage and characterisation of the bacteria were achieved (89% and 93% gnome coverage

using *de novo* methods), with missing sections often being repeat elements within the genome.

The majority of sequencing reads from spiked horse blood were associated with the spiked bacteria

(62.1% S. *aureus* and 73% *E. coli*), considering the horse genome is over 500 times larger than the *E.*

coli genome this shows that the vast majority of the host material was removed. Overall there was

good concordance of phenotypic and genotypic results showing the potential for rapid genotypic

prediction of antibiotic resistance from 10 bacterial cells in 1 ml of host blood. In the *S.aureus*

sequencing the closest sub-species identified was *Staphylococcus aureus subsp. aureus* HO 5096

0412, this most likely represents the closest sequence in the reference database, rather than the actual

type of this organism. Antibiotic resistance predictions were concordant with the phenotypic results

except for ciprofloxacin. The inconclusive ciprofloxacin results demonstrate the need for improved

understanding of the mechanisms of resistance. Ciprofloxacin resistance is harder to predict as it isn't

associated with a single gene acquisition. Three mechanisms of fluoroquinolone resistance have been

proposed in *S. aureus*, Topoisomerase IV gene mutations, DNA gyrase gene mutations and an active

efflux pump (NorA)[45]. The complexity of predicting ciprofloxacin resistance suggests that the

database may be lacking in its ability to predict ciprofloxacin resistance, and so this is the most likely

cause of the inconclusive result for ciprofloxacin resistance. The developers of Mykrobe, a commonly

used antibiotic resistance predictor (Bradley et al[11]) found a false negativity rate of 4.6% for

ciprofloxacin resistance.

In addition to identifying the isolates resistance to beta-lactams the database was able to identify the

blaZ gene and *MecA* gene. Genotypic testing will never entirely replace phenotypic susceptibility

testing, due to its inability to identify novel resistance determinants and the comprehensive nature of

phenotypic testing. However, in the case of invasive sepsis, the gain in speed provided by not having

to culture the organism to determine susceptibility could be life-saving.

Multiple antibiotic resistance factors were identified in the *E. coli*, which gave good concordance with

the phenotypic data. Using the genotypic data, it was possible to rapidly identify the beta-lactamase

present as *BlaCMY-2*. The rapid identification of the specific resistance genes in bacteria could help

identify outbreaks by providing more information than a simple antibiogram. Additionally, it could

help monitor resistance genes, to identify genes that are increasing in incidence. Large amounts of

horizontal genome transfer amongst Gram negative bacteria has the potential to cause outbreaks of

resistant bacteria through genes or plasmids[46], which would be more complex to track. Rapid

identification of genes causing the resistance in isolates could help inform epidemiological and

outbreak studies which could involve several species of bacteria. A study of NGS sequencing data

from bacteraemia isolates of *E. coli* have shown resistance prediction specificity of 97%[2], if this was

coupled with direct from sample sequencing genotypic prediction could inform treatment more

rapidly than phenotypic test.

This study demonstrates that *de novo* assembly methods can be used to identify bacterial species and

accurately predict antibiotic resistance. Lower coverage of the genomes after reference assembly

(92% and 93%), could reflect that references used were close but not exact matches to the clinical

isolates in the study, or the inability of short read assembly methods to assemble repeated elements

which are longer than the read length. The inability to subtype the *E. coli* strain most likely reflects

the large variety in *E. coli* genomes with the core genome representing only 13% of the pangenome[47].

An earlier study by the authors, using *E. coli* K12 MG1655 gave genome coverage with 100%

sequence identity with the E. coli K12 MG1655 reference (data not shown). Clinical isolates rather

than fully characterised laboratory strains were selected to test the applicability of the analysis to

unknown samples. The ability to identify bacteria in a sample will be very dependent on the quality of

the database used, and results will be biased towards common pathogens, but this method should be

applicable to most bacteria found in clinical samples.

Limitations of this model include differences between horse and human blood, and difference in

sample between spiked bacteria and bacteria causing a true infection. Additionally, factors in the

blood such as immune and inflammatory reactions could alter the efficiency of the preparation

method. As the bacteria was only quantified using dilution methods it is not possible to know the

exact number of bacteria spiked into the sample. Whilst three replicates of each bacteria were checked

for survival through the model, it was only possible to sequence one representative of each of the

bacteria. However, this model shows proof of principle that direct unbiased sequencing directly from

blood is possible and could be used to diagnose and inform treatment of bacterial bloodstream

infections.

Acknowledgements

This project was funded by Roche Diagnostics as a scientific studentship award and completed at

Public Health England Centre for Infection.

Declarations of interest.

None

References

1.   Gordon, N. C. *et al.* Prediction of staphylococcus aureus antimicrobial resistance by whole-
275   genome sequencing. *J. Clin. Microbiol.* **52,** 1182–1191 (2014).

276   2.   Stoesser, N. *et al.* Predicting antimicrobial susceptibilities for Escherichia coli and Klebsiella
277   pneumoniae isolates using whole genomic sequence data. *J. Antimicrob. Chemother.* **68,**
278   2234–44 (2013).

279   3.   Leopold, S. R., Goering, R. V., Witten, A., Harmsen, D. & Mellmann, A. Bacterial whole-
280   genome sequencing revisited: Portable, scalable, and standardized analysis for typing and
281   detection of virulence and antibiotic resistance genes. *J. Clin. Microbiol.* **52,** 2365–2370
282   (2014).

283   4.   Ashton, P. M. *et al.* Insight into Shiga toxin genes encoded by Escherichia coli O157 from
284   whole genome sequencing. *PeerJ* **3,** e739 (2015).

285   5.   Kennedy, A. D. *et al.* Epidemic community-associated methicillin-resistant Staphylococcus
286   aureus: recent clonal expansion and diversification. *Proc. Natl. Acad. Sci.* **105,** 1327–1332
287   (2008).

288   6.   Harris, S. R. *et al.* Evolution of MRSA During Hospital Transmission and Intercontinental
289   Spread. *Science* **327,** 469–474 (2010).

290   7.   Eyre, D. W. *et al.* A pilot study of rapid benchtop sequencing of Staphylococcus aureus and
291   Clostridium difficile for outbreak detection and surveillance. *BMJ Open* **2,** e001124–e001124
292   (2012).

293   8.   Reuter, S. *et al.* A pilot study of rapid whole-genome sequencing for the investigation of a
294   Legionella outbreak. *BMJ Open* **3,** e002175 (2013).

295   9.   Snitkin, E. S. *et al.* Tracking a Hospital Outbreak of Carbapenem-Resistant Klebsiella
296   pneumoniae with Whole-Genome Sequencing. *Sci. Transl. Med.* **4,** 148ra116-148ra116
297   (2012).

298   10.   Köser, C. U. *et al.* Rapid whole-genome sequencing for investigation of a neonatal MRSA
299   outbreak. *N.Engl.J.Med.* **366,** 2267–2275 (2012).

300   11.   Bradley, P. *et al.* Rapid antibiotic resistance predictions from genome sequence data for S.
301   aureus and M. tuberculosis. *bioRxiv* **6,** 18564 (2015).

12. Pankhurst, L. J. *et al.* Rapid, comprehensive, and affordable mycobacterial diagnosis with
whole-genome sequencing: A prospective study. *Lancet Respir. Med.* **4,** 49–58 (2016).

13. Mason, O. U. *et al.* Metagenomics reveals sediment microbial community response to
Deepwater Horizon oil spill. *ISME J.* **8,** 1464–75 (2014).

14. Vieira-silva, S. & Rocha, E. P. C. The systemic imprint of growth and its uses in ecological
(meta)genomics. *PLoS Genet.* **6,** (2010).

15. Sharon, I. *et al.* Comparative metagenomics of microbial traits within oceanic viral
communities. *ISME J.* **5,** 1178–1190 (2011).

16. Gilbert, J. a. & Dupont, C. L. Microbial Metagenomics: Beyond the Genome. *Ann. Rev. Mar.*
Sci. **3,** 347–371 (2011).

17. Erickson, A. R. *et al.* Integrated Metagenomics/Metaproteomics Reveals Human Host-
Microbiota Signatures of Crohn's Disease. *PLoS One* **7,** (2012).

18. Lamendella, R., Domingo, J. W. S., Ghosh, S., Martinson, J. & Oerther, D. B. Comparative
fecal metagenomics unveils unique functional capacity of the swine gut. *BMC Microbiol.* **11,**
103 (2011).

19. Walker, A. W., Duncan, S. H., Louis, P. & Flint, H. J. Phylogeny, culturing, and
metagenomics of the human gut microbiota. *Trends Microbiol.* **22,** 267–274 (2014).

20. Kristensen, D. M., Mushegian, A. R., Dolja, V. V. & Koonin, E. V. New dimensions of the
virus world discovered through metagenomics. *Trends Microbiol.* **18,** 11–19 (2010).

21. Mokili, J. L., Rohwer, F. & Dutilh, B. E. Metagenomics and future perspectives in virus
discovery. *Curr. Opin. Virol.* **2,** 63–77 (2012).

22. Do, L. A. H. *et al.* Respiratory Syncytial Virus and Other Viral Infections among Children
under Two Years Old in Southern Vietnam 2009-2010: Clinical Characteristics and Disease
Severity. *PLoS One* **11,** e0160606 (2016).

23. Dacheux, L. *et al.* A preliminary study of viral metagenomics of french bat species in contact
with humans: Identification of new mammalian viruses. *PLoS One* **9,** (2014).

24. Pien, B. C. *et al.* The Clinical and Prognostic Importance of Positive Blood Cultures in Adults.
Am. J. Med. **123,** 819–828 (2010).

25.   Prasad, N., Murdoch, D. R., Reyburn, H. & Crump, J. A. Etiology of Severe Febrile Illness in
331   Low- and Middle-Income Countries : A Systematic Review. *PLoS One* 1–25 (2015).
332   doi:10.1371/journal.pone.0127962

333   26.   Schubert, S. *et al.* Novel, improved sample preparation for rapid, direct identification from
334   positive blood cultures using matrix-assisted laser desorption/ionization time-of-flight
335   (MALDI-TOF) mass spectrometry. *J. Mol. Diagn.* **13,** 701–6 (2011).

336   27.   Prod'Hom, G., Bizzini, A., Durussel, C., Bille, J. & Greub, G. Matrix-assisted laser desorption
337   ionization-time of flight mass spectrometry for direct bacterial identification From positive
338   blood culture pellets. *J. Clin. Microbiol.* **48,** 1481–1483 (2010).

339   28.   Stevenson, L. G., Drake, S. K. & Murray, P. R. Rapid identification of bacteria in positive
340   blood culture broths by matrix-assisted laser desorption ionization-time of flight mass
341   spectrometry. *J. Clin. Microbiol.* **48,** 444–447 (2010).

342   29.   Zelenin, S. *et al.* Microfluidic-based isolation of bacteria from whole blood for sepsis
343   diagnostics. *Biotechnol. Lett.* **37,** 825–830 (2015).

344   30.   Lasken, R. S. Single-cell genomic sequencing using Multiple Displacement Amplification.
345   *Curr. Opin. Microbiol.* **10,** 510–516 (2007).

346   31.   Raghunathan, A. *et al.* Genomic DNA Amplification from a Single Bacterium Genomic DNA
347   Amplification from a Single Bacterium. *Appl. Environ. Microbiol.* **71,** 3342–3347 (2005).

348   32.   Cowell, A. N. *et al.* Selective Whole-Genome Amplification Is a Robust Method That Enables
349   Scalable Whole-Genome Sequencing of Plasmodium vivax from Unprocessed Clinical
350   Samples. *MBio* **8,** e02257-16 (2017).

351   33.   Brown, C. T., Howe, A., Zhang, Q., Pyrkosz, A. B. & Brom, T. H. A Reference-Free
352   Algorithm for Computational Normalization of Shotgun Sequencing Data. *arXiv* **1203.4802,**
353   1–18 (2012).

354   34.   Schmieder, R. & Edwards, R. Quality control and preprocessing of metagenomic datasets.
355   *Bioinformatics* **27,** 863–864 (2011).

356   35.   Huson, D., Auch, A., Qi, J. & Schuster, S. MEGAN analysis of metagenome data. *Gennome
357   Res.* **17,** 377–386 (2007).

358   36.   Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications to
359   Single-Cell Sequencing. *J. Comput. Biol.* **19,** 455–477 (2012).

37.   Zankari, E. *et al.* Identification of acquired antimicrobial resistance genes. *J. Antimicrob.*
Chemother. **67,** 2640–2644 (2012).

38.   Votintseva, A. A. *et al.* Same-day diagnostic and surveillance data for tuberculosis via whole-
genome sequencing of direct respiratory samples. *J. Clin. Microbiol.* **55,** 1285–1298 (2017).

39.   Street, T. L. *et al.* Molecular diagnosis of orthopaedic device infection direct from sonication
fluid by metagenomic sequencing. *J. Clin. Microbiol.* **55,** JCM.00462-17 (2017).

40.   Kirstahler, P. *et al.* Genomics-Based Identification of Microorganisms in Human Ocular Body
Fluid. *bioRxiv* 1–14 (2017). doi:10.1101/176529

41.   Cao, M. D. *et al.* Streaming algorithms for identification of pathogens and antibiotic resistance
potential from real-time MinION™ sequencing. *Gigascience* **5,** 1–16 (2016).

42.   Alberts B, Johnson A, L. J. *Molecular Biology of the Cell. 4th edition.* (2002).

43.   Woyke, T. *et al.* One bacterial cell, one complete genome. *PLoS One* **5,** (2010).

44.   Rhee, M., Light, Y. K., Meagher, R. J. & Singh, A. K. Digital Droplet Multiple Displacement
Amplification (ddMDA) for Whole Genome Sequencing of Limited DNA Samples. *PLoS One*
11, e0153699 (2016).

45.   Tanaka, M., Wang, T., Onodera, Y., Uchida, Y. & Sato, K. Mechanism of quinolone resistance
in Staphylococcus aureus. *J. Infect. Chemother.* **6,** 131–139 (2000).

46.   Tofteland, S., Naseer, U., Lislevand, J. H., Sundsfjord, A. & Samuelsen, ??rjan. A Long-Term
Low-Frequency Hospital Outbreak of KPC-Producing Klebsiella pneumoniae Involving
Intergenus Plasmid Diffusion and a Persisting Environmental Reservoir. *PLoS One* **8,** 1–8
(2013).

47.   Vieira, G. *et al.* Core and panmetabolism in Escherichia coli. *J. Bacteriol.* **193,** 1461–1472
(2011).

***Tables and Figures*- Whole genome amplification and sequencing of low cell numbers directly from a bacteria spiked blood model**

Figure 1 Illustration of the process for bacterial isolation from whole blood using HetaSep and selective lysis with Saponin. Numbers refer to sampling points where bacterial recovery was investigated. Total processing time for bacterial isolation is one hour.

Stage		*S. aureus*	*E. coli*
1	Spiked cells	10	11
2	RBC separation-bottom	0	0
3	RBC separation-top	10	9
4	Addition of Saponin	12	9
5	Water shock	12	12
6	Salt restoration	10	10
7	Cell pelleting-top	0	0
	Cell pelleting-bottom	9	9
8	DNAse treatment	11	12
9	EDTA and PBS	10	12
10	Wash 1-top	0	0
	Wash 1-bottom	9	11
11	Wash 2-top	0	0
	Wash 2–bottom	10	10
12	Wash 3-top	0	0
	Wash 3–bottom	8	9
13	Final pellet	11	9

Table 1 the average number of CFU isolated at each processing stage for bacterial isolation from whole blood using HetaSep and selective lysis with Saponin after workflow improvements. Numbers refer to sampling points where bacterial recovery was investigated as illustrated in **Figure 1**

	Phenotypic-(MIC)	Genotypic	Comment
	E. coli		
Ampicillin	R (>8)	R	*blaCMY-2*
Ceftriaxone	R (>4)	R	*blaCMY-2*
Ciprofloxacin	R (>1)	R	*gyrA*
Colistin	S (<=0.5)	S	
Gentamicin	S (<=0.25)	S	
Levofloxacin	R (>2)	R	*gyrA*
Meropenem	S (<=0.25)	S	
Nalidixic acid	(>16)	R	*ermB*
Nitrofurantoin	S (<=16)	S	
Trimethoprim	R (>4)	R	*dfrA*
	S. aureus		
Ciprofloxacin	R (>2)	Inconclusive	
Clindamycin	S (<=0.25)	S	
Erythromycin	S (<=0.25)	S	
Fusidic acid	S (<=1)	S	
Gentamicin	S (<1)	S	
Methicillin	R (oxacillin 2)	R	*MecA*
Mupirocin	S (<=1)	S	
Penicillin G	R (> 0.25)	R	*blaZ*
Rifampicin	S (<=0.25)	S	
Tetracycline	S (<=0.5)	S	
Trimethoprim	S (<=1)	S	
Vancomycin	S (1)	S	

Table 2 phenotypic and genotypic result for antibiograms of *E. coli* (ResFinder) and *S. aureus* (Mykrobe)