Melanism patches up the defective cuticular morphological traits through

promoting the up-regulation of cuticular protein-coding genes in *Bombyx mori*

Liang Qiao[*1], Ri-xin Wang[§], You-jin Hao[*], Hai Hu[§], Gao Xiong[§], Song-zhen He[§],

Jiang-bo Song[§], Kun-peng Lu[§], James Mallet[†], Ya-qun Xin[‡], Bin Chen[*], Fang-yin

Dai[§1]

* Chongqing Key Laboratory of Vector Insects, Institute of Entomology and

Molecular Biology, College of Life Sciences, Chongqing Normal University,

Chongqing, 401331, China

[§] State Key Laboratory of Silkworm Genome Biology; Key Laboratory of Sericultural

Biology and Genetic Breeding, Ministry of Agriculture; College of Biotechnology,

Southwest University, Chongqing 400716, China.

† Department of Organismic and Evolutionary Biology, Harvard University,

Cambridge, MA 02138, USA

‡Department of Evolutionary Ecology, LMU Munich Großhaderner Str. 2, 82152,

Germany

Running title: Role of melanin on cuticle defects

Keywords: Melanic coloring; Cuticle features; Cuticular protein-encoding genes; Induction; *Bombyx mori*

[1] **Corresponding Authors:**

Liang Qiao, Chongqing Key Laboratory of Vector Insects, Institute of Entomology and Molecular Biology, College of Life Sciences, Chongqing Normal University, No. 37 Daxuecheng Road, Chongqing, 401331, China; E-mail: qiaoliangswu@163.com; Tel: 86-23-65910315

Fang-yin Dai, State Key Laboratory of Silkworm Genome Biology, Key Laboratory for Sericulture Functional Genomics and Biotechnology of Agricultural Ministry, Southwest University, No. 2 Tiansheng Road, Chongqing, +86-23-68400715, China; E-mail: fydai@swu.edu.cn; Tel: 86-23-68250346

Abstract

Melanin and cuticular proteins are important cuticle components in insect. The cuticle defects caused by the loss function of cuticular protein-encoding genes could hinder melanin deposition. However, the effects of melanin variation on cuticular protein-encoding genes and the corresponding morphological traits associated with these genes remain largely unknown. Using *Bombyx mori* as a model, we showed that the level of melanism during larval cuticle pigmentation correlated positively with the expression of cuticular protein-encoding genes. This correlation stemmed from the simultaneous induction of these genes by the melanin precursors. More importantly, the effect of the melanism background on the cuticle induced the up-regulation of other functionally redundant cuticular protein-encoding genes to rescue the morphological and adaptive defects caused by the dysfunction of some mutated cuticular proteins, and the restorative ability increased with increasing melanism levels, which gives a novel evidence that melanism enhances insect adaptability. These findings deepen our understanding of the interactions among cuticle components, as well as their importance in the stabilization of the normal morphology and function of the cuticle.

Introduction

The prerequisite for the benefits of melanism to insect is not only the integrity of the melanin biosynthesis and regulatory pathway (Wilson *et al.* 2001; Liu *et al.* 2015; Mallet and Hoekstra 2016), but also the normal presence of the platform it relied on

(Wittkopp *et al.* 2003; Wittkopp and Beldade 2009; Andersen 2010; Moussian 2010;

Van Belleghem *et al.* 2017). For the insects, the most important fundamental platform

for the color pattern drawing is the cuticle (Hopkins and Kramer 1992; Andersen 2010;

Moussian 2010). During shaping the cuticle, the maintenance and stability of the

cuticle features depends on normal functional cuticular proteins and the interactions

with other components (Hopkins and Kramer 1992; Guan *et al.* 2006; Suderman *et al.*

2006; Andersen 2010; Moussian 2010; Chaudhari *et al.* 2011; Noh *et al.* 2016). Due

to the crucial roles of cuticular proteins on cuticle development, when their coding

genes are loss of function, the abnormal or defective cuticle will likely affect the

deposition and attachment of melanin, which is not conducive to the performance of

color pattern (Kanekatsu *et al.* 1988; Oota and Kanekatsu 1993; Arakane *et al.* 2012;

Jasrapuria *et al.* 2012; Wang 2013; Noh *et al.* 2015). Yet little is known about the

corresponding response mode of cuticular protein-encoding genes via the alteration of

the melanin biosynthesis or regulatory pathway.

Recently, several high throughput expression surveys showed that the abundance of

cuticular protein-encoding genes in different colored integuments varied in some

insect species, especially with evidently up-regulated in the melanic regions

(Futahashi *et al.* 2012; He *et al.* 2016; Wu *et al.* 2016; Tajiri 2017), and some of those

shared very similar expression patterns and functions (Nakato *et al.* 1994; Nakato *et*

al. 1997; Shofuda *et al.* 1999; Okamoto *et al.* 2008; Liang *et al.* 2010; Tang *et al.*

2010; Qiao *et al.* 2014).These studies implied that there are probably some

relationships between the promotion of melanism and the expression of cuticular

protein-encoding genes. Prior to this study, further exploring and understanding of the

potential relationships were unclear. Additionally, when melanism-promoting

instructions and defective cuticle proteins occur simultaneously, what are the effects

of their relationships on the presence of the corresponding morphological traits？

In the Lepidoptera model, *Bombyx mori*, an intriguing phenomenon has been

reported that a larval melanic mutant, *Striped* (p^S, 2-0.0) is able to rehabilitate the

malformed body shape, as well as the adaptability defects of the *stony* mutant (*st*,

8-0.0) (Xiang 1995; Banno *et al.* 2005). A recent study clarified that a transcription

factor, *Apontic-like*, which boosts the expression of melanin synthesis genes in

epidermal cells, is responsible for the p^S mutant (Yoda *et al.* 2014). Besides this, there

are also multiple alleles with different melanism degrees at the *p* locus, including p^B,

p^M, *etc* (Xiang 1995; Banno *et al.* 2005; Yoda *et al.* 2014). And the *stony* mutant (*st*,

8-0.0) which precisely caused by the deletion of a RR1-type larval cuticular

protein-encoding gene, *BmLcp17* (or *BmorCPR2*) showed hard and tight touch feeling,

uncoordinated ratios of the length of the internodes and the intersegment folds (I/IF),

bulges at intersegment folds, and severely defective locomotion and behavioral

activities in the larval stage (Qiao *et al.* 2014). In addition, the similarity of the gene

expression patterns and functional characteristics among some members of the

RR1-type larval cuticular protein-encoding gene family, such as *BmLcp18*, *BmLcp22*,

BmLcp30 also suggest that they may play very similar roles as *BmLcp17* in shaping

the larvae cuticle (Nakato *et al.* 1994; Nakato *et al.* 1997; Shofuda *et al.* 1999;

Okamoto *et al.* 2008; Liang *et al.* 2010; Tang *et al.* 2010; Qiao *et al.* 2014). These

dispersed findings are linked through the epistasis of p^S to *stony*, then provide the

breakthrough and the exceptional genetic resources for exploring the interactions

between melanin and cuticular protein-encoding genes.

Here we illustrated that the transcripts levels of four cuticular protein-encoding

genes were positively correlated with the melanism degree of larval cuticle, which

were due to the simultaneous induction these genes by the intracellular melanin

precursors. Moreover, by importing melanism-promoting instruction into the *stony*

mutant, the cuticle deficiency rescued through functionally redundant compensation

by some other up-regulated cuticular protein-encoding genes, which a new evidence

that melanism as a beneficial trait. These findings deepen our understanding of the

interactions among genetic factors which shape morphological features in

lepidopteran, and emphasize the ecological and evolutionary significance of these

mutual interactions.

Materials and Methods

Silkworm strains

Wild-type strains Dazao ($+^p$) and melanic mutant strains p^M, p^S and p^B (Xiang 1995;

Banno *et al.* 2005; Yoda *et al.* 2014) were analyzed in this study. The darkness of

pigment was measured as mean OD value using Image J (https://imagej.nih.gov/ij/).

In terms of melanism degree, the body color of an individual that is homozygous at

one of the aforementioned melanic loci is darker than that of a heterozygous

individual (Xiang 1995; Banno *et al.* 2005). The albinism mutant *albino* (*al*) (Banno

et al. 2005; Fujii *et al.* 2013), non-diapause wild-type strain N4 (used for melanin

inhibition treatment) and *BmLcp17* deletion strain Dazao-*stony* (near isogeneic line of

Dazao, which have been backcrossed with Dazao over 26 generations) were supplied

by the Silkworm Gene Bank in Southwest University. The *al* mutant was fed with

artificial die at 28℃, and the others were fed fresh mulberry leaves under a

12 h/12 h light/dark photoperiod at 24 °C.

Chemicals and cell lines

140 L-dopa (D9628), dopamine (H8502), tetrahydrofolic acid (BH_4) (T3125) and

141 2,4-Diamino-6-hydroxypyrimidine (DAHP) (D19206) were purchased from

142 Sigma-Aldrich (St. Louis, MO, USA). *BmNs* cell line was kept in our laboratory.

Mating combinations and progeny phenotypes identification

The p^S and p^M strains were crossed with the Dazao-*stony* strain to generate the F_1

generation, respectively. The F_2 generation were produced by an F_1 self-cross, and

individuals of day 5 of the fifth-instar were collected to further use. The p^B strain was

crossed with the Dazao-*stony* strain to generate F_1 progeny, which mated with the

Dazao-*stony* strain to generate the BC_1 generation, and fed them until at day 5 of the

fifth-instar. Firstly, individuals of the F_2 or BC_1 generations were separated by their

cuticle pigmentation. Subsequently, their phenotypes were further classified by

morphological characteristics, touch feeling, and the ratios between the number of

internodes and intersegmental folds in the second, third, and fourth abdominal

segments based on a earlier method (Qiao *et al.* 2014).

Genotyping

155 Because the p^S, p^M and p^B mutations are multiple alleles at the p locus, they should

156 be located in proximity each other on the chromosome 2 (Xiang 1995; Banno *et al.*

157 2005; Yoda *et al.* 2014). Based on the reported sequence of the gene corresponding to

158 the p^S allele, a polymerase chain reaction (PCR)-based molecular marker within of the

159 genomic region of the *Apontic-like* was designed. PCR screening were performed in

160 the p^X (X=M, S and B) and Dazao-*stony* to obtain the polymorphism molecular marker

161 for p locus. Similarly, molecular marker was also designed within genomic region of

162 the *BmLcp17* for polymorphism screening of the *stony* locus. The primers used in this

163 study are listed in Table S1.

164 *Association analysis of gene expression, phenotype and genotype*

165 Day 5 fifth-instar larvae of the Dazao or Dazao-*stony* strains were selected for

166 cuticle dissection. The cuticles of the semi-lunar marking region and the non-melanic

167 portion between the paired semi-lunar marking were finely dissected. Then, gene

168 expression levels of *BmLcp17*, *BmLcp18*, *BmLcp22* and *BmLcp30* in these regions

169 were compared. Gene expression patterns in the dorsal epidermis of abdominal

170 segments (from semi-lunar marking to star marking) from day 5 of fifth instar larvae

171 (Dazao, p^S/+, p^M/+, p^B/+). Similarly, day 1 of second instar larvae of the *al* and Dazao

172 strains were also investigated. In addition, the same regions of dorsal epidermis were

173 collected from F_2 generation individuals with the p^S/p^S, st/st, and p^S/+, st/st genotypes,

174 as well as the p^M/p^M, st/st and p^M/+, st/st genotypes for gene expression analyses. For

175 all genotyped individuals, the ratios of the length of the intersegment and the

176 intersegment fold were also analysed using image J.

Melanin precursors-promoting and inhibition treatments

The preparation and concentration of L-dopa and dopamine solutions were slightly

modified according to the description by Futahashi (Futahashi and Fujiwara 2005),

and filtered with a 0.22 μm filter prior to use. Cells were washed three times with

Grace medium without melanin precursors to remove metabolites and other

contaminants on the cell surface. Then, 0.8mL medium containing L-dopa or

dopamine was added separately into each well, and the medium without melanin

precursors was used as the control. The plate was sealed with tape, wrapped with foil

and incubated at 28℃ for 24 h for a gene expression analysis. In BH_4 feeding assays,

the 30mM working solution was prepared by dissolving tetrahydrofolic acid into

ddH_2O, and then smeared on the artificial diet to feed the *al* mutant, and ddH_2O was

used as the control. Phenotype were observed and recorded from the second instar and

the expression of cuticular protein-encoding genes was also analysed. For the

melanism-inhibition experiment, the wild-type strain N4 was selected. Newly-hatched

larvae were divided into treatment and control groups. Individuals in the treatment

group fed with DAHP dissolved in 0.1M NaOH (15g/L), and individuals in the

control group fed 0.1M with NaOH. Phenotypes observation and gene expressions

detection were performed on day 1 of the second-instar larvae.

Quantitative RT–PCR

Total RNA extraction, reverse transcription and qRT-PCR conducted as described

previously (Qiao *et al.* 2014). Three biological replicates were prepared for each

condition, and *BmRPL3* was used as the internal control. Primers are listed in Table

S1.

Results

Entirely distinct expression patterns of cuticular protein-encoding genes and the

cuticle appearance between melanic and non-melanic regions

The expression patterns of *BmLcp17*, *BmLcp18*, *BmLcp22* and *BmLcp30* in melanic

or non-melanic epidermal regions are shown in Figure 1. These results, together with

earlier studies (Futahashi *et al.* 2012; He *et al.* 2016; Wu *et al.* 2016), revealed that the

gene expressions were significantly higher in melanic parts of the cuticle than in

non-melanic parts. Moreover, micro protrusions were more intensive in the melanic

regions than in the non-melanic regions, and accompanied by a higher chitin content

(another important cuticle component (Hopkins and Kramer 1992; Moussian *et al.*

2006; Andersen 2010; Moussian 2010; Chaudhari *et al.* 2011; Qiao *et al.* 2014), and

the reduction chitin content was reported to impede cuticle melanism (Moussian *et al.*

2005)) (Figure S1A). These results showed that, regardless of the different genetic

basis of the melanic mutants or the melanic markings in the non-melanic strains,

excessive accumulation of melanin in the cuticle (accompany by the changes in the

surface structure and the chitin content of the melanic cuticle) was closely related to

the high expression levels of the cuticular protein-encoding genes.

Expression level of larval cuticular protein-encoding genes positively correlated

with the degree of cuticle melanism

To obtain further insights into the relationships between the accumulation of

melanin and the expression of cuticular protein-encoding genes, we investigated gene

expression patterns using four different mutations at the p locus (Dazao ($+^p$), p^M, p^S

and p^B), which showed gradually increasing melanin accumulation (Figure 2A). Our

results revealed that the gene expression levels were gradually and significantly

up-regulated with the increase in the degree of melanism of the cuticle (Figure 2B).

They further showed that the expression levels correlated positively with the degree of

melanism. Thus, the quantities of cuticular proteins with similar or redundant

functions could be increased greatly by the increasing the degree of melanism.

Typical stony phenotyped individuals masked after the introduction of melanic loci

into the BmLcp17 defection strain

We assessed the effects of modulating the melanic background on the phenotypic

defects caused by the deletion of *BmLcp17*. After mating the p^B and *stony* parental,

the percentage of BC_1 individuals with melanism and the normal body shape in the

backcrossed population of the $p^B \times stony$ cross was 52 % (290/553; and theoretically, it

was 25 %), yet individuals with the melanism cuticle and *stony*-type body shape were

not found (theoretically should be almost equivalent to the number of individuals with

melanism cuticle and normal body shape) (Figure 3). In the cross of $p^S \times stony$, ~10.8%

(36/331) of F_2 progeny had an lighter melanism body color and smaller body size

(Figure 3). These individuals exhibited a little hard and tight touched body, but the

body was much softer than that of the *stony* mutant. Their intersegment folds bulged

slightly, and the length were significantly shorter than that of the internodes;

accordingly, their phenotypes slightly resembled the morphological features of the

stony mutant (Figure 3 and 4A). Even so, We did not find individuals with the typical

stony-type body shape and also defective adaptability under the melanism background

(theoretical ratio is 3/16). (Figure 3). Similarly, in the $p^M \times stony$ corss, only

approximately ~11.7% (51/437) of the individuals of the F_2 population were very

lightly melanic, but they exhibited obviously unusual morphological features (Figure

3, Figure 4A). When compared with the ~10.8% F_2 individuals (aforementioned) from

the $p^S \times stony$ cross, the touch feelings of individuals from the $p^M \times stony$ cross were

tighter and firmer, and the intersegment folds bulged more obviously and had a higher

proportion among the segments, meaning that their body features were closer to the

phenotype of the *stony* mutant (Figure 3 and 4A) (Qiao *et al.* 2014). Nevertheless,

melanic individuals showing the typical *stony*-type body features and defective

adaptability still did not appear (theoretical ratio is 3/16) in the progeny from the $p^M \times$

stony cross. Therefore, induction of the melanic mutation into individuals with a

defective cuticular protein-encoding gene could mask the adverse phenotypes, and the

masking effect was more remarkable with the increasing degree of melanism.

Other larval cuticular protein-encoding genes up-regulated evidently in the melanic

and non-stony phenotypic, but with mutated BmLcp17 genotypic offspring

Using the molecular markers (closely linked to the *p* and *st* loci, respectively), we

further genotyped the progenies with melanic color pattern and non-*stony* (including

ambiguous *stony*-like) body shape. The results revealed that approximately 49% of

the individuals showing a melanic color and normal body shape in the BC_1 population

from the $p^B \times stony$ cross was the $p^B/+^{pB}$, *st/st* genotype (Figure 4A). The ratios of the

length of the internodes and the intersegment folds (I/IF) were ~4, which is very

similar to that in $p^B/+^{pB}$, $+^{st}/st$ individuals, and also no significant differences as that in

the wild-type individuals (Figure 4B) (Qiao *et al.* 2014). In the F$_2$ generation from the

p^S×*stony* cross, approximately 9.3% of the individuals with the p^S/p^S, st/st genotype

and very few individuals (~1%) with the genotype $p^S/+^{pS}$, st/st exhibited a melanic

color and the normal body shape (Figure 4A). For p^S/p^S, st/st genotyped individuals,

the I/IF value was approximately 3.3 (Figure 4B). Although the I/IF value was lower

than that in $p^S/\_$, $+^{st}/\_$ individuals (approximately 4, Figure 4B), it was much higher

than that in the *stony* mutant (approximately 1.6 (Qiao *et al.* 2014)). Despite the

slightly smaller body size of the p^S/p^S, st/st individuals, their body shape was

essentially normal (Figure 4A). The genotypes of those lightly melanic individuals,

whose body shape was slightly like that of the *stony* phenotype (mentioned in Result

3), were all the $p^S/+^{pS}$, st/st, and the I/IF value of these individuals was approximately

2.7 (Figures 4A and 4B). In progeny of the p^M×*stony* cross, ~10.5% of the individuals

with the p^M/p^M, st/st genotype and ~0.7 % of the individuals with the $p^M/+^{pM}$, st/st

genotype showed a melanic color and subtle *stony* features (just very slight bulges)

(Figure 4A). The body size of the p^M/p^M, st/st individuals were smaller than those of

the $p^M/\_$, $+^{st}/\_$ individuals. They exhibited a certain sense of hardness, and the I/IF

ratio was approximately 2.8, which is in good agreement with their phenotypes

(Figure 4B). Additionally, individuals showing very slight melanism and

morphological features that were more similar to that of the *stony* mutant (mentioned

in Result 3) were all the $p^M/+^{pM}$, st/st genotype, and their I/IF values were

approximately 1.8, which is closer to that of the *stony* mutant (Figures 4A and 4B). In

addition, the expression of cuticular protein-encoding genes in p^S/p^S, *st/st* individuals

were significantly higher than that in $p^S/+^{pS}$, *st/st* individuals (Figure. 4C); a similar

result was also obtained from the p^M/p^M, *st/st* and $p^M/+^{pM}$, *st/st* individuals (Figure 4C).

Taken together, these results revealed that more cuticular proteins with similar

functions were accumulated in the cuticle of melanic homologous individuals at the *p*

locus. Based on the comprehensive and association analysis, we infer that melanic

background effectively drove the expressions of cuticular protein-encoding genes

with similar expression patterns and redundant functions, which compensated for the

morphological and adaptability defects caused by the dysfunctional *BmLcp17* gene;

and the law of compensatory abilities was $p^B/+^{pB}$, *st/st* > p^S/p^S, *st/st* > p^M/p^M, *st/st* ξ

$p^S/+^{pS}$, *st/st* > $p^M/+^{pM}$, *st/st*, which corresponds well with the gradual weakening of the

degree of melanism (Figures 4 and Figure S3).

Content variations of melanin precursors affect the transcript amount of the

cuticular protein-encoding genes

Due to the crucial material basis for the cuticle melanism (no matter what kind of

genetic basis caused by) is the extensive accumulation of melanin precursors in the

epidermal cells; thus, the essence that melanism tend to increase the expression of

some cuticular protein-encoding genes should be driven by the relations between the

accumulation of melanin precursors and the transcripts amount of the cuticular

protein-encoding genes. The basal expressions of four cuticular protein-encoding

genes were detected in *BmNs* cells (Figure S5), indicating that there are regulatory

pathways controlling the expression of cuticular protein-coding genes in this cell line;

therefore, this cell line can be used to examine the effect of melanin precursors on

gene expressions. After incubating *BmNs* cells with melanin precursors, the

expression of cuticular protein-encoding genes was significantly higher in cells

treated either by L-dopa or dopamine, compare with that in the control group (Figure

5 top (left)). In addition, when treated with BH_4, the second-instar *al* mutant (which is

characterized by albinism and a porous cuticle due to a mutation of sepiapterin

reductase, which leads to the insufficient synthesis of the co-factor BH_4 during the

synthesis of melanin precursors (Banno *et al.* 2005; Fujii *et al.* 2013) was rescued to a

melanic body color (Lab unpublished contribution) (Fujii *et al.* 2013), and the gene

expressions were obviously higher than that in the control group (Figure 5 top (right)).

These results suggest that the expression of cuticular protein-encoding genes can be

induced by increasing amounts of melanin precursors. Furthermore, adding DAHP (an

inhibitor of guanylate cyclase hydrolase (GTPCHI), which is an important

rate-limiting enzyme in the synthesis of BH_4 (Hamadate *et al.* 2008), inhibited the

synthesis of BH_4 in wild-type second-instar larvae by blocking the synthesis of

melanin precursors in epidermal cells, which caused individuals to lose their original

melanic body color (Lab unpublished contribution). The gene expressions were also

significantly reduced when compared with that in the control group (Figure 5 bottom).

Thus, the content and variation of the intracellular melanin precursors are important

factors regulating the expression of cuticular protein-encoding genes. We concluded

that the inducing effect of the melanin precursors on the expression of cuticular

protein-encoding genes is the basis for melanism promoting genes' transcription.

Discussion

Melanin is deposited in the cuticle, in which many of cuticular proteins also occur;

thus, overlapping distribution of cuticular proteins make it possible for them to

interact with melanin, thereby contributing to the phenotype of the cuticle (Hopkins

and Kramer 1992; Suderman *et al.* 2006; Andersen 2010; Moussian 2010; Hu *et al.*

2013). We clearly observed that extensive star-like protrusions were present on the

cuticle when a large amount of melanin accumulated (Figure S1). Similar correlations

between cuticle structure and body color have been reported multiple times (Futahashi

et al. 2012; Noh *et al.* 2016; Tan *et al.* 2016), suggest that there should be some

interactions between cuticular proteins and melanin. Although the exact interaction

pattern between these two cuticular components is unknown, microscopic observation

clearly shows that the deposition of an excessive amount of melanin affected the

cuticle characteristics.

The expression profile of cuticular protein-encoding genes in melanic silkworm

mutants and the black markings of *Papilio* larvae supported the view that

overexpressed cuticular proteins probably participate in the transport or maintenance

of the corresponding pigments in a specific painted cuticle (Figures 1, 2 and Figure S1)

(Futahashi *et al.* 2012; He *et al.* 2016; Wu *et al.* 2016). Yet over-expression of

cuticular protein-encoding genes *in vivo* cannot trigger (or induce) the formation of

the melanic cuticle (Tajiri *et al.* 2017). Therefore, we reasoned that once the

instructions for melanin accumulation are included in the developmental program

(melanism being the original factor), other cuticle features and structures should adapt

to the level of melanin accumulation, regardless of genetic background. The

up-regulation of cuticular protein-encoding genes should be a necessary adaptation for

the maintenance and stability of the structural characteristics and physical properties

of melanic cuticles. The interactions of melanin and cuticular proteins at the

ultrastructural level deserves special attention in follow-up studies.

Despite the elaborate regulatory mechanism by which melanin precursors affect the

expression of cuticular protein-encoding genes is unclear, this study really revealed

the existence of this regulatory phenomenon (Figure 5). Cuticle formation is regulated

stringently in temporal and spatial patterns, the accumulation and oxidization of

melanin precursors, and the interactions (such as cross-linking) among other

components should occur after the formation of the initial formation of the cuticle

(Moussian 2010; Sobala and Adler 2016; Tajiri 2017). Therefore, we propose that the

cuticular proteins induced by melanin precursors are used to set up a platform for

further accumulation and oxidization of melanin precursors. When melanin-associated

protein-encoding genes have similar expression patterns and functions (Nakato *et al.*

1994; Nakato *et al.* 1997; Shofuda *et al.* 1999; Okamoto *et al.* 2008; Liang *et al.* 2010;

Tang *et al.* 2010; Qiao *et al.* 2014), the production of a large amount of functionally

similar cuticular proteins would be driven by the melanic background to guarantee

construction and stability of the melanic cuticle. During this process, melanism

unlocks the complementary features of melanin-associated cuticular proteins (such as

BmLcp18, *BmLcp22*, *BmLcp30*), rescued cuticular malformation caused by the loss of

function of some cuticular proteins (such as defected *BmLcp17* in *stony* mutant).

Because it appears that a special cuticle-forming pattern is regulated by melanin

accumulation, melanism may enhance insect adaptability to avoid the impairment of

survival caused by the mutation and/or functional loss of some cuticular proteins, and

these results also add new evidence to explain how melanism can be a beneficial trait

(Wilson *et al.* 2001; True 2003; Wittkopp *et al.* 2003; Wittkopp and Beldade 2009;

Liu *et al.* 2015; Mallet and Hoekstra 2016).

As far as we known, as structural proteins, there is no evidence to suggest that the

four larval cuticular protein-encoding genes and their orthologous can enter the

nucleus and regulate gene expression by acting as transcription factors. In addition,

our findings and some other studies showed that changes in the expression of one

cuticular protein-encoding genes of R&R family did not affect the expression of other

members (Figure S6, S7) (Arakane *et al.* 2012; Noh *et al.* 2015). Moreover, several

studies revealed that organisms optimize resources use at gene or protein expression

level (Liebermeister *et al.* 2004; Dekel and Alon 2005); thus, the expression of

melanin associated cuticle protein-encoding genes might be appropriately and

simultaneously coordinated with the sufficient accumulation of intracellular melanin

precursors, as a relatively direct, economical, efficient and convenient strategy.

Furthermore, DAHP inhibits GTPCHI activity, but it does not directly impair the

extracellular accumulation of melanin and protein–protein interactions in the cuticle.

Finally, there is some evidence that melanin precursors regulate gene expressions via

the receptors (Konradi *et al.* 1996; Berke *et al.* 1998; Westin *et al.* 2001). In summary,

we hypothesized that the up-regulation of the four *BmLcp* genes were induced

simultaneously by excessive amounts of melanin precursors, but should not be the

interaction among these four genes on regulation level. To test our hypothesis, further

analyses will be performed to determine the expression of the remaining *BmLcp* genes

in some *BmLcps* mutant (such as defective *BmLcp17*) cell line by increasing or

decreasing the content of melanin precursors.

The markings in the *stony* mutant were lighter than those in the wild-type strain,

and accompanied by the down-regulation of melanin synthase genes (Figure S4). If

the dysfunction of some cuticular proteins cannot be effectively supplemented,

abnormalities of the cuticle structure and interactions of various cuticular components

may occur, which probably result in barriers for melanin deposition and metabolism.

Perhaps this is the reason why the $p^S/+^{pS}$, *st/st* (or $p^M/+^{pM}$, *st/st*) genotypes are lighter

than the $p^S/+^{pS}$, $+^{st}/\_$ (or $p^M/+^{pM}$, $+^{st}/\_$) genotypes, besides to their different body shape.

Because the homozygous of p^B mutation is lethal (Xiang 1995; Banno *et al.* 2005), we

were unable to obtain F_2 progeny with the p^B/p^B, *st/st* genotype. However, it is worth

noting that the body shape and the melanism degree of $p^B/+^{pB}$, $+^{st}/st$ and $p^B/+^{pB}$, *st/st*

genotype individuals were almost same (Figure 4A). Therefore, during cuticle

formation, if the signal promoting melanism is sufficiently strong, plenty of melanin

precursors should be generated. The contents of functionally redundant proteins

induced by melanin precursors are enough to fill the gap generated by the dysfunction

of some cuticular proteins in theoretically, which will be able to guarantees the normal

accumulation of melanin. Therefore, we believe that there should be a threshold of the

melanism-promoting effect. When the accumulation of melanin precursors spans this

threshold, the requirements for cuticular proteins with similar function are not

weakened even when one of them loses the function, which ensuring the formation of

a normal cuticle structure the later pigmentation.

We proposed a possible regulatory model as follows: when large amounts of

melanin precursors induced by endogenous- and/or exogenous melanism-promoting

factors, the melanin pathway may directly or indirectly induce the up-regulation of

some cuticular protein-encoding genes to guarantee the formation of normal structural

features of the melanic cuticle. During this process, if some cuticular proteins lose

their functions, other functionally redundant cuticular proteins induced by melanin

precursors can compensate for the functional defects. The compensatory intensity is

increased with increasing melanin accumulation. When melanin accumulation spans a

certain threshold, this compensation can totally mask the defective phenotype caused

by the malfunctioning genes, which adds to growing evidence that melanism may

have pleiotropic effects that enhance fitness over and above the effect of melanin

accumulation itself (Figure 6). Due to the coexistence of excess melanin and cuticular

proteins is common in other insects, and homologues the four *BmLcp* gene are widely

distributed in the Lepidoptera (Table S2), we presume that the above reciprocal action

and its corresponding biological significance are conserved in other Lepidoptera

insects to that in *Bombyx mori*.

In summary, we used crucial components of the cuticle to elucidate the mutual

effects among some key genetic factors, and the physiological significance of these

mutual effects during the development of the morphological features. Our findings not

only contribute a realistic basis to in-depth study of the interaction patterns of melanin

and cuticular proteins, but they will also inspire relevant studies of other Lepidopteran

insects or other insect species.

Literature Cited

Andersen, S. O., 2010 Insect cuticular sclerotization: a review. Insect Biochem Mol
Biol 40**:** 166-178.

Arakane, Y., J. Lomakin, S. H. Gehrke, Y. Hiromasa, J. M. Tomich *et al.*, 2012
Formation of rigid, non-flight forewings (elytra) of a beetle requires two major
cuticular proteins. PLoS Genet 8**:** e1002682.

Banno, Y., H. Fujii, Y. Kawaguchi, K. Yamamoto, K. Nishikawa *et al.*, 2005 *A guide*
to the silkworm mutants: 2005 gene name and gene symbol. Kyusyu
University, Fukuoka, Japan.

Berke, J. D., R. F. Paletzki, G. J. Aronson, S. E. Hyman and C. R. Gerfen, 1998 A
complex program of striatal gene expression induced by dopaminergic
stimulation. J Neurosci 18**:** 5301-5310.

Chaudhari, S. S., Y. Arakane, C. A. Specht, B. Moussian, D. L. Boyle *et al.*, 2011
Knickkopf protein protects and organizes chitin in the newly synthesized
insect exoskeleton. P Natl Acad Sci USA 108**:** 17028-17033.

Dekel, E., and U. Alon, 2005 Optimality and evolutionary tuning of the expression
level of a protein. Nature 436**:** 588-592.

Fujii, T., H. Abe, M. Kawamoto, S. Katsuma, Y. Banno *et al.*, 2013 Albino (al) is a
tetrahydrobiopterin (BH$_4$)-deficient mutant of the silkworm *Bombyx mori*.
Insect Biochem Mol Biol 43**:** 594-600.

Futahashi, R., and H. Fujiwara, 2005 Melanin-synthesis enzymes coregulate
stage-specific larval cuticular markings in the swallowtail butterfly, *Papilio*
xuthus. Dev Genes Evol 215**:** 519-529.

Futahashi, R., H. Shirataki, T. Narita, K. Mita and H. Fujiwara, 2012 Comprehensive
microarray-based analysis for stage-specific larval camouflage
pattern-associated genes in the swallowtail butterfly, *Papilio xuthus*. BMC
Biol 10**:** 46.

Guan, X., B. W. Middlebrooks, S. Alexander and S. A. Wasserman, 2006 Mutation of
TweedleD, a member of an unconventional cuticle protein family, alters body

shape in *Drosophila*. P Natl Acad Sci USA 103**:** 16794-16799.

Hamadate, N., K. Noguchi, M. Sakanashi, T. Matsuzaki, J. Nakasone *et al.*, 2008
Effect of decreased levels of intrinsic tetrahydrobiopterin on endothelial
function in anesthetized rats. J Pharmacol Sci 107**:** 49-56.

He, S. Z., X. L. Tong, K. P. Lu, Y. R. Lu, J. W. Luo *et al.*, 2016 Comparative analysis
of transcriptomes among *Bombyx mori* strains and sexes reveals the genes
regulating melanic morph and the related phenotypes. Plos One 11.

Hopkins, T. L., and K. J. Kramer, 1992 Insect cuticle sclerotization. Annu Rev
Entomol 37**:** 273-302.

Hu, Y. G., Y. H. Shen, Z. Zhang and G. Q. Shi, 2013 Melanin and urate act to prevent
ultraviolet in the integument of the silkworm, *Bombyx mori*. Arch Insect
Biochem 83**:** 41-55.

Jasrapuria, S., C. A. Specht, K. J. Kramer, R. W. Beeman and S. Muthukrishnan, 2012
Gene families of cuticular proteins analogous to peritrophins (CPAPs) in
Tribolium castaneum have diverse functions. Plos One 7.

Kanekatsu, R., Y. Banno, E. Nagashima and T. Miyashita, 1988 Genetical studies on a
new spontaneous mutant "*Bamboo*" (*Bo*) of the silkworm. The Journal of
Sericultural Science of Japan 57**:** 151-156.

Konradi, C., J. C. Leveque and S. E. Hyman, 1996 Amphetamine and
dopamine-induced immediate early gene expression in striatal neurons
depends on postsynaptic NMDA receptors and calcium. J Neurosci 16**:**
4231-4239.

Liang, J., L. Zhang, Z. Xiang and N. He, 2010 Expression profile of cuticular genes of
silkworm, *Bombyx mori*. BMC Genomics 11**:** 173.

Liebermeister, W., E. Klipp, S. Schuster and R. Heinrich, 2004 A theory of optimal
differential gene expression. Biosystems 76**:** 261-278.

Liu, S. S., M. Wang and X. C. Li, 2015 Pupal melanization is associated with higher
fitness in *Spodoptera exigua*. Sci Rep-Uk 5.

Mallet, J., and H. E. Hoekstra, 2016 Ecological genetics: a key gene for mimicry and
melanism. Curr Biol 26**:** R802-R804.

Min, C., J. B. Song, L. I. Zhi-Quan, D. M. Tang, X. L. Tong *et al.*, 2016 Progress and
perspective of silkworm as a model of human diseases for drug screening.
Acta Pharmaceutica Sinica.

Moussian, B., 2010 Recent advances in understanding mechanisms of insect cuticle
differentiation. Insect Biochem Mol Biol 40**:** 363-375.

Moussian, B., H. Schwarz, S. Bartoszewski and C. Nusslein-Volhard, 2005
Involvement of chitin in exoskeleton morphogenesis in *Drosophila*
melanogaster. J Morphol 264**:** 117-130.

Moussian, B., E. Tang, A. Tonning, S. Helms, H. Schwarz *et al.*, 2006 *Drosophila*
Knickkopf and Retroactive are needed for epithelial tube growth and cuticle
differentiation through their specific requirement for chitin filament
organization. Development 133**:** 163-171.

Nakato, H., K. Shofuda, S. Izumi and S. Tomino, 1994 Structure and developmental
expression of a larval cuticle protein gene of the silkworm, *Bombyx mori*.

521   Biochim Biophys Acta 1218**:** 64-74.

522   Nakato, H., M. Takekoshi, T. Togawa, S. Izumi and S. Tomino, 1997 Purification and
523   cDNA cloning of evolutionarily conserved larval cuticle proteins of the
524   silkworm, *Bombyx mori*. Insect Biochem Mol Biol 27**:** 701-709.

525   Noh, M. Y., S. Muthukrishnan, K. J. Kramer and Y. Arakane, 2015 *Tribolium*
526   *castaneum* RR-1 cuticular protein TcCPR4 is required for formation of pore
527   canals in rigid cuticle. Plos Genetics 11.

528   Noh, M. Y., S. Muthukrishnan, K. J. Kramer and Y. Arakane, 2016 Cuticle formation
529   and pigmentation in beetles. Curr Opin Insect Sci 17**:** 1-9.

530   Okamoto, S., R. Futahashi, T. Kojima, K. Mita and H. Fujiwara, 2008 Catalogue of
531   epidermal genes: genes expressed in the epidermis during larval molt of the
532   silkworm *Bombyx mori*. BMC Genomics 9**:** 396.

533   Oota, K., and R. Kanekatsu, 1993 Morphological studies on the *Bamboo* (*Bo*) mutant
534   of the silkworm, *Bombyx mori*. The Journal of Sericultural Science of Japan
535   62**:** 448-454.

536   Qiao, L., G. Xiong, R. X. Wang, S. Z. He, J. Chen *et al.*, 2014 Mutation of a cuticular
537   protein, BmorCPR2, alters larval body shape and adaptability in silkworm,
538   *Bombyx mori*. Genetics 196**:** 1103-1115.

539   Shofuda, K., T. Togawa, H. Nakato, S. Tomino and S. Izumi, 1999 Molecular cloning
540   and characterization of a cDNA encoding a larval cuticle protein of *Bombyx*
541   *mori*. Comp Biochem Physiol B Biochem Mol Biol 122**:** 105-109.

542   Sobala, L. F., and P. N. Adler, 2016 The gene expression program for the formation of
543   wing cuticle in *Drosophila*. Plos Genetics 12.

544   Suderman, R. J., N. T. Dittmer, M. R. Kanost and K. J. Kramer, 2006 Model reactions
545   for insect cuticle sclerotization: cross-linking of recombinant cuticular proteins
546   upon their laccase-catalyzed oxidative conjugation with catechols. Insect
547   Biochem Molec 36**:** 610-611.

548   Tajiri, R., 2017 Cuticle itself as a central and dynamic player in shaping cuticle. Curr
549   Opin Insect Sci 19**:** 30-35.

550   Tajiri, R., N. Ogawa, H. Fujiwara and T. Kojima, 2017 Mechanical control of whole
551   body shape by a single cuticular protein obstructor-E in *Drosophila*
552   *melanogaster*. Plos Genetics 13.

553   Tan, D., X. L. Tong, H. Hu, S. Y. Wu, C. L. Li *et al.*, 2016 Morphological
554   characterization and molecular mapping of an irradiation-induced Speckled
555   mutant in the silkworm, *Bombyx mori*. Insect Mol Biol 25**:** 93-104.

556   Tang, L., J. Liang, Z. Zhan, Z. Xiang and N. He, 2010 Identification of the
557   chitin-binding proteins from the larval proteins of silkworm, *Bombyx mori*.
558   Insect Biochem Mol Biol 40**:** 228-234.

559   True, J. R., 2003 Insect melanism: the molecules matter. Trends Ecol Evol 18**:**
560   640-647.

561   Van Belleghem, S. M., P. Rastas, A. Papanicolaou, S. H. Martin, C. F. Arias *et al.*,
562   2017 Complex modular architecture around a simple toolkit of wing pattern
563   genes. Nat Ecol Evol 1.

564   Wang, L.-Y., 2013 Molecular genetic studies on the oily mutants in the silkworm,

Bombyx mori, pp. 43 in *Department of Agricultural and Environmental Biology, Graduate School of Agricultural and Life Sciences*. University of Tokyo.

Westin, J. E., M. Andersson, M. Lundblad and M. A. Cenci, 2001 Persistent changes in striatal gene expression induced by long-term L-DOPA treatment in a rat model of Parkinson's disease. Eur J Neurosci 14**:** 1171-1176.

Wilson, K., S. C. Cotter, A. F. Reeson and J. K. Pell, 2001 Melanism and disease resistance in insects. Ecol Lett 4**:** 637-649.

Wittkopp, P. J., and P. Beldade, 2009 Development and evolution of insect pigmentation: genetic mechanisms and the potential consequences of pleiotropy. Semin Cell Dev Biol 20**:** 65-71.

Wittkopp, P. J., S. B. Carroll and A. Kopp, 2003 Evolution in black and white: genetic control of pigment patterns in *Drosophila*. Trends Genet 19**:** 495-504.

Wu, S. Y., X. L. Tong, C. X. Peng, G. Xiong, K. P. Lu *et al.*, 2016 Comparative analysis of the integument transcriptomes of the black dilute mutant and the wild-type silkworm *Bombyx mori*. Sci Rep-Uk 6.

Xiang, Z., 1995 *Genetics and breeding of the silkworm*. Chinese Agriculture Press, Beijing, PR China.

Yoda, S., J. Yamaguchi, K. Mita, K. Yamamoto, Y. Banno *et al.*, 2014 The transcription factor Apontic-like controls diverse colouration pattern in caterpillars. Nat Commun 5**:** 4936.

Acknowledgments

We thank Dr. Tianzhu Xiong, Dr. Nathan Edelman, Dr. Jorge Eduardo Amaya Romero and Ms. Yan Yao for the valuable advices. This work was supported by the Hi-Tech Research and Development 863 Program of China (Grant No. 2013AA102507), the National Natural Science Foundation of China (No. 31302038; 31372379), the Natural Science Foundation Project of ChongQing (CSTC) (No. cstc2013jcyjA80022) and Par-Eu Scholars Program (20136666).

Figure legends

Figure 1. Expression of four larval cuticular protein-encoding genes in melanic and non-melanic integuments. A and B represent the gene expressions between the

semi-lunar marking (black box) and the non-melanic region (between the semi-lunar

marking, red box) in Dazao or Dazao-*stony*, respectively. Scale bar: 2 mm. C.

Comparative analysis of gene expressions in the dorsal side of abdominal segments

(from the third to the fourth segment, red box) between the p^S and Dazao strains.

Scale bar: 1 cm. D. Comparison of gene expressions between the second-instar *al*

mutant and the Dazao strain (melanic). The red hashtag symbol indicates the Fig. 1 D

we are showing is cited from the previous study of our lab group (Min *et al.* 2016)

with modification. Scale bar: 2 mm. *t*-test, n=3; * $p < 0.05$; ** $p < 0.01$; *** $p <$

0.001.

Figure 2. Expression of cuticular protein-encoding genes in integuments showing

different degree of melanism. A and B display comparisons of degree of melanism

and cuticular protein-encoding gene expression levels among four strains with mutant

alleles at the *p* locus. Scale bar: 1 cm. Ratios represent the ratios of gene expression

levels between two strains. Symbols (−, +, ++, and +++) represent the increment of

the degree of melanism. Star represents the melanin-associated cuticular

protein-encoding genes. *t*-test, n=3; * $p < 0.05$; ** $p < 0.01$.

Figure 3. Segregation patterns of the phenotypes of progenies from different crosses

of melanic mutant strains and the *stony* strain. In the segregated progenies, the first

item in the list (B,N), (B,st), (S,N), (S,st), (M,N), (M,st), (N,N) and (N,st) represents

p^B-, p^S- p^M- and Normal color patterns, respectively. The second item indicates body

shape features marked with non-*stony* type (N) and the *stony* type (st). It is

noteworthy that in (S, st-am+) and (M, st-am+), the second item represents the

ambiguous *stony* body shape. The size of "+" symbol represents the corresponding

degree of the ambiguous *stony* body shape. Superscript "T"s represent theoretical

values. Superscript "A"s represent actual values. Backslashes indicates that a value

was not obtained. Chi-squared test, $* p < 0.05; ** p < 0.01$.

Figure 4. Association analysis of the genotypes, phenotypes and gene expression

levels in segregated progenies from different crosses. A. Correlation analysis between

the genotypes and phenotypes in self-crossed or backcrossed progenies. Scale bar: 1

cm. White and red stars represent polymorphic bands at the $+^p$ and p^X loci ($X = B$, S or

M), respectively. Red and white hash-tag represent polymorphic bands at the *st* and $+^{st}$

locus, respectively. Solid and dotted red arrows indicate the relative degree of bulging

(solid > dotted), respectively. Slashes show the genotypes within one phenotypic

category. The thickness of the slash represents the proportion of the corresponding

genotypes. Dotted backslashes indicate that the number of individuals with the

corresponding genotype is quite low. B. Ratios of the length of internodes and

intersegmental folds in the second, third, and fourth abdominal segments of

individuals with different genotypes (showing melanic body color) in the self-crossed

or backcrossed progenies. n≥3, *t*-test, $** p < 0.01$. C. Gene expression analysis of

cuticular protein-encoding genes in homogeneous and heterogeneous individuals at

the *p* locus from self-crossed progenies of $p^S \times stony$, and $p^M \times stony$ under the

condition that the cuticle was melanic and the genotype was homozygous recessive at

st locus. n = 3, *t* – test, * $p < 0.05$; ** $p \leq 0.01$.

Figure 5. Effect of melanin precursors (top left: in cells) and BH_4 (top right: in *vivo*)

treatments on the expression of cuticular protein-enconding genes (*t*-test; n = 3, * $p <$

0.05; ** $p < 0.01$), and variations of gene expression levels in larval cuticle treated

with the inhibitor DAHP (bottom). n = 3, *t*-test, ** $p < 0.01$.

Figure 6. Schematic overview of the effect of melanin precursors on the expressions

of cuticular protein-encoding genes. Black solid circles represent the melanin.

Triangles represent the BmLcps with similar expression patterns and functions. Solid

and hollow triangles represent the normal and defective functions, respectively. Red

rhombi represent other components (such as chitin) in the cuticle. Solid and dotted

double-headed arrows indicate probable interactions and loss of interactions due to

the functional deficiency of some *BmLcps* (such as dysfunctional *BmLcp17*),

respectively. Purple arrows show the direction in which the melanin precursors or

cuticular proteins flow from the haemolymph to the epidermal cells as well as from

the epidermal cells to the cuticle. Red arrows indicate the increased contents of

melanin precursors or the up-regulation of cuticular protein-encoding genes. Purple

polyline arrows indicate melanism-promoting factors produced by other genetic

instructions. Double dovetail arrows indicate the effect of melanin precursors on

cuticular protein-encoding genes. The question mark indicates that the details of

induction (directly or indirectly) are unclear. Here the cuticle especially means the

exo-cuticles in which the melanin and cuticular proteins deposited in (Hopkins and

Kramer 1992; Suderman *et al.* 2006; Andersen 2010; Moussian 2010).