Title: A rate threshold mechanism regulates MAPK stress signaling and

survival

Authors: Amanda N. Johnson[1], Guoliang Li[1], Hossein Jashnsaz[1], Alexander Thiemicke[1],

Benjamin K. Kesler[1], Dustin C. Rogers[1], and Gregor Neuert[1-3*]

Affiliations:

[1] Department of Molecular Physiology and Biophysics, School of Medicine, Vanderbilt

University, Nashville, TN, 37232, USA

[2] Department of Pharmacology, School of Medicine, Vanderbilt University, Nashville, TN, 37232,

USA

[3] Department of Biomedical Engineering, School of Engineering, Vanderbilt University,

Nashville, TN, 37232, USA

*Correspondence to: gregor.neuert@vanderbilt.edu

<u>Abstract</u>

Cells are exposed to changes in extracellular stimulus concentration that vary as a

function of rate. However, the effect of stimulation rate on cell behavior and signaling remains

poorly understood. Here, we examined how varying the rate of stress application alters budding

yeast cell viability and mitogen-activated protein kinase (MAPK) signaling at the single-cell

level. We show that cell survival and signaling depend on a rate threshold that operates in

conjunction with a concentration threshold to determine the timing of MAPK signaling during

rate-varying stimulus treatments. We also discovered that the stimulation rate threshold is

sensitive to changes in the expression levels of the Ptp2 phosphatase, but not of another

phosphatase that similarly regulates osmostress signaling during switch-like treatments. Our

results demonstrate that stimulation rate is a regulated determinant of signaling output and

provide a paradigm to guide the dissection of major stimulation rate-dependent mechanisms in

other systems.

Main Text

All cells employ signal transduction pathways to respond to physiologically relevant

changes in extracellular stressors, nutrient levels, hormones, morphogens, and other stimuli that

vary as functions of both concentration and rate in healthy and diseased states[1–7]. Switch-like

"instantaneous" changes in the concentrations of stimuli in the extracellular environment have

been widely used to show that the strength of signaling and overall cellular response are

dependent on the stimulus concentration, which in many cases needs to exceed a certain

threshold[8,9]. Although studies have shown that the rate of stimulation can also influence

signaling output in a variety of pathways[10–17], it is not clear how cells integrate information

conveyed by changes in both the stimulation rate and concentration in determining whether or

not to commence signaling.

Recent investigations have demonstrated that stimulation rate can be a determining factor

in signal transduction. In contrast to switch-like perturbations, which trigger a broad set of

stress response pathways, slow stimulation rates activate a specific response to the stress applied

in *Bacillus subtilis* cells[10]. Meanwhile, shallow morphogen gradient stimulation fails to activate

developmental pathways in mouse myoblast cells in culture, even when concentrations sufficient

for activation during pulsed treatment are delivered[12]. These observations raise the possibility

that stimulation profiles must exceed a set minimum rate or rate threshold to achieve signaling

activation. Although such rate thresholds would help cells decide if and how to respond to

dynamic changes in stimulus concentration, the possibility of signaling regulation by a rate

threshold has never been directly investigated in any system. Further, no study has

experimentally examined how stimulation rate requirements impact cell phenotype or how cells

molecularly regulate the stimulation rate required for signaling activation. As such, the

biological significance of any existing rate threshold regulation of signaling remains unknown.

The budding yeast *Saccharomyces cerevisiae* high osmolarity glycerol (HOG) pathway

provides an ideal model system for addressing these issues (Fig. 1a). The evolutionarily

conserved mitogen-activated protein kinase (MAPK) Hog1 serves as the central signaling

mediator of this pathway[18–20]. It is well established that instantaneous increases in osmotic stress

concentration induce Hog1 phosphorylation, activation, and translocation to the nucleus[19,21–29].

Activated Hog1 governs the majority of the cellular osmoadaptation response that enables cells

to survive[21,30,31]. Multiple apparently redundant MAPK phosphatases dephosphorylate and

inactivate Hog1, which, along with the termination of upstream signaling after adaptation, results

in its return to the cytosol (Fig. 1a)[21,23,24,32–38]. Because of this behavior, time-lapse analysis of

Hog1 nuclear enrichment in single cells has proven an excellent way to monitor signaling

responses to dynamic stimulation patterns in real-time[25–29,39,40]. Further, such assays have been

readily combined with traditional growth and molecular genetic approaches to link observed

signaling responses with cell behavior and signaling pathway architecture[26–28].

Here, we use systematically-designed osmotic stress treatments imposed at varying rates

of increase to show that a rate threshold condition regulates yeast high-stress survival and Hog1

MAPK signaling. We demonstrate that only stimulus profiles that satisfy both this rate threshold

condition and a concentration threshold condition result in robust signaling. We go on to show

that the protein tyrosine phosphatase Ptp2, but not the related Ptp3 phosphatase, serves as a

major rate threshold regulator. By expressing *PTP2* under the control of a series of different

enhancer-promoter DNA constructs, we demonstrate that changes in the level of Ptp2 expression

can alter the stimulation rate required for signaling induction. These findings establish rate

thresholds as a critical and regulated component of signaling biology akin to concentration

thresholds.

Results

Stimulus treatment rate affects yeast high osmotic stress survival and Hog1 nuclear

translocation pattern. We hypothesized that a rate threshold governs the budding yeast HOG

pathway. The existence of such a threshold would place a minimum stimulation rate on

osmoadaptation. Shallow rates of stimulation that fall below this minimum would not result in a

robust osmoadaptive response, even if the same stimulus concentration triggered a response

when applied at a faster rate.

To test this hypothesis, we first took advantage of the fact that pre-exposure to a mild

pulsed stress treatment increases the stress tolerance of yeast through induction of the

osmoadaptive response[41–43]. This increased tolerance enables cells to survive a second,

otherwise lethal, high-stress pulse treatment[41,42]. This preconditioning protection paradigm

enables us to examine the rate-dependence of mild stress responses independent of whether the

stresses themselves differentially impact cell fitness.

Cells were initially exposed to preconditioning osmostresses (400 mM NaCl final

concentration, equivalent integrated osmolarity) delivered at different rates. The impact of these

preconditioning treatments on high-stress susceptibility was determined by comparing the

viability of cells after a second transient high-stress treatment and no stress treatment (2 M and 0

M NaCl) (Fig. 1b). No substantial difference in growth was observed among the preconditioned

cultures immediately following the initial treatments (Supplemental Fig. 1). However, the rate of

preconditioning stress delivery profoundly influenced cell survival after the second high-stress;

cells exposed to a slow rate treatment [4 mM NaCl/min (grey)] were significantly more sensitive

to a high second stress than cells exposed to faster rate treatments [16 mM NaCl/min (cyan) or

pulse (blue)] (Fig. 1c).

99   To investigate the mechanism(s) underlying the priming of high second stress survival in

response to variations in preconditioning stimulation rate, we tracked the nuclear enrichment of

Hog1-yellow fluorescent protein (YFP), a proxy for canonical salt-stress signaling activation,

during rate-varying stress treatments via single-cell time-lapse microscopy. Because previously

published studies show that Hog1 nuclear translocation follows changing external stress patterns

in real-time[28,29,39], we began probing the relationship between stimulation rate and translocation

using a stress treatment applied as a quadratic function (linearly increasing rate over time) (Fig.

1d, e). Compared to pulse treatment, quadratic treatment resulted in an unprecedented delayed

and diminished Hog1 nuclear accumulation response (Fig. 1f). Stimulation rate thus strongly

influences signaling as well as high-stress survival phenotype, consistent with expectations based

on rate threshold regulation of the HOG pathway.

A rate threshold regulates Hog1 nuclear enrichment. The Hog1 quadratic stimulation

response pattern could be related to either known stimulus concentration threshold

requirements[23] or stimulation rate, since both are changing over time (Fig. 2a, b). To investigate

these relationships, Hog1 nuclear localization during the quadratic stress gradient was reanalyzed

separately as a function of either the treatment concentration or rate. The concentration and rate

corresponding with nuclear enrichment were extrapolated using two intersecting lines fitting

Hog1 nuclear localization baseline and accumulation increase for each independent experiment.

The thresholds for Hog1 translocation were 117.5 ± 34.5 mM NaCl and 12.1 ± 1.6 mM

NaCl/min, respectively (Fig. 2a, b). Surprisingly, the concentration associated with nuclear

accumulation was much higher than the 50-70 mM salt expected based on published switch-like

treatment responses[23,26], a result that raises the possibility that the rate of stress change drives

Hog1 nuclear enrichment pattern during quadratic stimulation.

123   To systematically test if ~12 mM NaCl/min represents an unreported rate threshold

requirement for signaling, we exposed cells to linear ramp treatments delivered at rates above, at,

and below 12 mM NaCl/min (Fig. 2c, d). Only ramps with rates greater than 12 mM NaCl/min

resulted in detectable, albeit slightly delayed, Hog1 translocation (Fig. 2e). This result directly

demonstrates that stimulus profiles must exceed a rate threshold to induce Hog1 nuclear

enrichment.

Hog1 translocation requires both the rate and concentration thresholds. If stimulation rate

solely determines signaling response, all ramps with gradients steeper than the rate threshold

should bring about Hog1 translocation at the onset of treatment. Instead, we observed a delay in

translocation in the 16 mM NaCl/min and 20 mM NaCl/min ramp treatments (Fig. 2e).

Previously published data also shows a lag in nuclear translocation during ramp treatment[27],

although the corresponding report does not discuss this observation. Reanalyzing translocation

during our ramp treatments that resulted in Hog1 nuclear localization as a function of

concentration revealed that the timing of translocation in both activating ramps was associated

with 57 ± 14.5 mM NaCl (Fig. 2f). This value agrees with the anticipated concentration

threshold of 50-70 mM salt.

These results suggest that Hog1 accumulates in the nucleus only when stress stimulations

exceed both the rate and concentration thresholds (Fig. 2g). Yeast cell stress signaling responses

thus appear to follow AND logic where the output (signaling) occurs only if both input

conditions (the rate and concentration thresholds) are met for a given stimulation profile. In this

model, neither steps that occur at an infinite rate but fall below the concentration threshold nor

slow increases in NaCl that exceed the concentration threshold effectively trigger signaling. On

the other hand, for those stimulations where there is a difference in the time each threshold

condition is satisfied, Hog1 translocation corresponds with whichever condition was met last.

The activating ramps used in these studies exceed the rate threshold before the concentration

threshold such that the timing of Hog1 translocation corresponds with the concentration

threshold (Fig. 2c-g). In our quadratic stimulation, the order in which these conditions are met is

flipped and nuclear accumulation corresponds with the rate threshold instead (Fig. 3a-c, i). The

AND logic model thus explains how quadratic stimulations can result in translocation at

comparable stimulation rates but different stimulus concentrations (Fig. 3d-i), and how fast rate

treatments (e.g. pulses and very fast ramps) which promptly satisfy both threshold conditions

induce rapid activation (Fig. 4a-g).

Identification of a rate threshold regulator. The existence of a rate threshold raises the

possibility of a rate threshold regulator. The ideal regulator would prevent Hog1 from

responding when stimulation patterns fail to reach the rate threshold without impinging on the

concentration threshold. To identify such a regulator, we first considered key modulators of

Hog1 function (Fig. 5a). We specifically focused on the protein tyrosine phosphatases Ptp2 and

Ptp3, which act as partially redundant Hog1 off-switches during instantaneous changes in stress

concentration[21,32,34], because these factors counteract phosphorylation of the Hog1 tyrosine

residue (Tyr-176) required for salt-induced Hog1 activity[44].

We hypothesized that the deletion of Ptp2 and/or Ptp3 would remove a barrier to stimuli-

induced Hog1 activation in stress treatments that exceed the concentration threshold but fall

below the rate threshold. To test this hypothesis, we examined Hog1 nuclear accumulation

patterns in wild-type (*WT*), *ptp2Δ,* and *ptp3Δ* yeast strains during our quadratic stimulation, in

which reduction or removal of the rate threshold would enable translocation earlier in the

treatment time course according to the AND logic model (Fig. 5b-d). While the pattern of Hog1

nuclear accumulation in *ptp3Δ* cells appeared similar to *WT* during quadratic stimulation,

accumulation in *ptp2Δ* cells occurred much earlier (Fig. 5e).

Reanalyzing accumulation as a function of rate revealed that accumulation in *ptp2Δ*

corresponds with a substantially reduced rate (Fig. 6a) compared to *WT*. Further, reanalyzing

accumulation as a function of concentration revealed that the timing of nuclear enrichment in

ptp2Δ corresponds with our measured concentration threshold (Fig. 6b). Thus, in *ptp2Δ* cells,

signaling apparently depends only on the concentration threshold condition (Fig. 6c), an effect

invisible to pulse treatments in which little difference in translocation pattern was observed

among *WT*, *ptp2Δ*, and *ptp3Δ* strains (Fig. 6d-e).

Tuning the rate threshold. If Ptp2 indeed serves as a major rate threshold regulator, its over-

expression should increase the threshold rate required for Hog1 nuclear enrichment. To test this

hypothesis, we constitutively expressed *PTP2* in the *ptp2Δ* strain under the control of the

upstream regulatory DNA sequences for other genes [*ADH1* and *TEF1*], creating strains with

highly increased Ptp2 expression levels (Fig. 7a), and studied Hog1 translocation during

quadratic stimulations. A quadratic treatment delivering rates up to 16 mM/min failed to induce

activation in the over-expression strains (Fig. 7b-d), while showing activation correspondence

with the expected stimulation rates in the empty vector and *PTP2* expression strains (Fig. 7e).

However, a quadratic treatment that included rates up to 30 mM/min (Fig. 8a, b) induced Hog1

translocation in all strains (Fig. 8c). Translocation in this second quadratic treatment followed

the pattern expected based on the relative timing of the concentration and rate thresholds in the

empty vector and native *PTP2* expression strains (Fig. 8d, Supplementary Fig.2), whereas

translocation in the over-expression strains corresponded with higher stimulation rates (Fig. 8d).

Ptp2 expression levels thus fine-tune the rate required to activate signaling.

In summary, the experiments we have presented directly describe regulation of the HOG

pathway by a rate threshold whose set point can be modulated by Ptp2 expression levels. Our

rate threshold discoveries represent a paradigm shift in the logic of signaling regulation.

Discussion

Because stimulation patterns vary in terms of both stimulus concentration and rate of

addition in vivo[1-7], it is important to thoroughly understand how the rate of stimulation affects

signal transduction and resulting cell behavior. Here, we have investigated how stimulation rate

impacts the budding yeast Hog1 model MAPK pathway. We find that shallow rates of

stimulation that fall below a threshold under-prepare cells for future high-stress survival,

corresponding with a failure to induce signaling. We also find that this rate threshold

collaborates with the concentration threshold in that both conditions must be met for signaling to

occur.

These findings demonstrate that the timing of osmostress signaling activation depends as

much upon stimulation rate as it does upon stimulus concentration change. Thus, stimulation

profile matters at least as much as stimulus identity in determining whether, when, and to what

extent signaling and associated cellular activity occur. Using our innovative application of rate-

varying stimulus treatments, we find that signal transduction can correspond with either the rate

or the concentration threshold depending on the type of stimulus profile applied (Fig. 9).

Meanwhile, signaling pathway mutations, such as the *PTP2* deletion and expression level

variants, may eliminate or alter the rate threshold requirement.

The signaling logic paradigm that controls the yeast osmostress pathway also appears to

govern other pathways given the failure of shallow stimulus gradients to trigger *B. subtilus*

general stress and mouse myoblast morphogen signaling even when concentrations sufficient for

activation during pulsed treatment[10,12]. We therefore speculate that rate thresholds are a general

operating principle of signaling systems, particularly those where known physiological variation

in stimulation rate exists[1–7]. A general rate threshold logic mechanism carries with it

implications for: (1) how cells make signaling decisions, (2) experimental design, and (3)

apparent redundancy in signaling networks.

Rate threshold logic mechanisms likely set important limits on pathway sensitivity to

extracellular stimulation. In the case of the HOG pathway studied here, these limits likely serve

the same purpose as the sensitivity limits set by the concentration threshold: conserve cellular

resources specifically for significant osmostresses. Like pulsed concentration changes below the

concentration threshold, slow increases in osmostress may not immediately threaten cell fitness.

Consistent with this idea, we do not observe any substantial growth difference between yeast

treated with single ramp stimulations above and below the rate threshold (see Supplementary

Fig. 1). Rate threshold regulation of osmostress signaling thus appears to ensure an

advantageous level of cell sensitivity to stimulation rate. In other pathways, rate threshold

mechanisms would provide a helpful way to differentiate signal from noise. The likely rate

threshold in the *B. subtilis* general stress pathway[10] probably also helps conserve resources by

activating only when the type of stress the cells experience is unclear. Meanwhile, having cells

effectively ignore very shallow morphogen gradients, as appears to be the case in the mouse

myoblast morphogen signaling pathway, likely helps direct cells into separate fates during

development[11,12].

Because both rate and concentration thresholds set the limits of cellular sensitivity to

extracellular stimuli, measuring these conditions in other signaling pathways serves as an

essential goal in signaling biology. Having these measurements in hand would enable better

understanding and prediction of how cells will respond to a given dynamic stimulation profile.

Further, defining the limits of cellular sensitivity would enable the detection of changes in these

limits caused by signaling pathway mutations such as those common in cancer and

developmental disorders[45]. Identifying these aberrant sensitivities would improve our

understanding of disease etiology and aid in the design of better treatments. Importantly, the

signaling logic paradigm uncovered by our study provides the necessary guide for designing

dynamic stimulation profiles that rapidly detect and measure threshold conditions. These profiles

are easy to implement and can be readily applied to make threshold condition measurements in

any signaling pathway and/or cell type[4,14].

The existence of any detected threshold condition begs the question of what regulates the

condition. We found that in the yeast HOG pathway, the protein tyrosine phosphatase Ptp2

serves as one major regulator of our newly discovered rate threshold condition. Excitingly, this

function provides Ptp2, which is otherwise partially redundant with Ptp3[21,32,34], with an

apparently unique role. Additionally, it expands the list of possible roles for phosphatases,

which remain much less understood than kinases, in spite of the fact that both types of factors

contribute to balancing cellular phosphorylation levels[46]. While phosphatases have generally

been thought of as pathway on/off switches, our findings and those of another pioneering study

performed using bacterial two-component systems[47] show that these important signaling factors

can regulate cellular sensitivity to extracellular stimulation. Further investigations using

dynamic stimulation profiles are needed to determine whether phosphatases in other signaling

systems regulate cellular sensitivity to stimulus addition. Such studies may also enable the

identification of unique functions for specific phosphatases as well as other factors that appear

redundant or partially redundant during switch-like changes in stimulus concentration, thereby

facilitating the disentanglement of complex signaling networks.

In conclusion, we have demonstrated a rate threshold mechanism in the HOG pathway

along with a highly generalizable strategy for investigating threshold conditions and the

signaling pathway components that regulate them. We strongly anticipate that the application of

this strategy to other systems will lead to improved disease diagnosis and treatment while

transforming our understanding of the logic underlying the complex signaling networks

prevalent in higher eukaryotes.

Main References

1.   Polonsky, K. S., Given, B. D. & Van Cauter, E. Twenty-four-hour profiles and pulsatile

patterns of insulin secretion in normal and obese subjects. *J. Clin. Invest.* **81**, 442–448

(1988).

2.   Hoffmann, E. K., Lambert, I. H. & Pedersen, S. F. Physiology of Cell Volume Regulation

in Vertebrates. *Physiol. Rev.* **89**, 193–277 (2009).

3.   Chovatiya, R. & Medzhitov, R. Stress, Inflammation, and Defense of Homeostasis. *Mol.*

Cell **54**, 281–288 (2014).

4.   Shimizu, T. S., Tu, Y. & Berg, H. C. A modular gradient-sensing network for chemotaxis

in Escherichia coli revealed by responses to time-varying stimuli. *Mol. Syst. Biol.* **6**, 382

(2010).

5.   Wang, C. J., Bergmann, A., Lin, B., Kim, K. & Levchenko, A. Diverse Sensitivity

Thresholds in Dynamic Signaling Responses by Social Amoebae. *Sci. Signal.* **5**, 213

(2012).

6.   Brabant, G., Prank, K. & Schofl, C. Pulsatile patterns in hormone secretion. *Trends*

Endocrinol. Metab. **3**, 183–190 (1992).

7.   Sagner, A. & Briscoe, J. Morphogen interpretation: concentration, time, competence, and

signaling dynamics. *Wiley Interdiscip. Rev. Dev. Biol.* **6**, e271 (2017).

doi:10.1002/wdev.271

8.   Forsten, K. E. & Lauffenburger, D. A. Autocrine ligand binding to cell receptors.

Mathematical analysis of competition by solution 'decoys'. *Biophys. J.* **61**, 518–529

(1992).

9.   Ang, J., Harris, E., Hussey, B. J., Kil, R. & McMillen, D. R. Tuning response curves for

296   synthetic biology. *ACS Synth. Biol.* **2**, 547–567 (2013).

297   10.   Young, J. W., Locke, J. C. W. & Elowitz, M. B. Rate of environmental change determines

298   stress response specificity. *Proc. Natl. Acad. Sci.* **110**, 4140–4145 (2013).

299   11.   Heemskerk, I. *et al.* Rapid changes in morphogen concentration control self-organized

300   patterning in human embryonic stem cells. *Elife* **8**, e40526 (2019).

301   12.   Sorre, B., Warmflash, A., Brivanlou, A. H. H. & Siggia, E. E. D. Encoding of temporal

302   signals by the TGF-β Pathway and implications for embryonic patterning. *Dev. Cell* **30**,

303   334–342 (2014).

304   13.   Kubota, H. *et al.* Temporal Coding of Insulin Action through Multiplexing of the AKT

305   Pathway. *Mol. Cell* **46**, 820–832 (2012).

306   14.   Thiemicke, A., Jashnsaz, H., Li, G. & Neuert, G. Generating kinetic environments to

307   study dynamic cellular processes in single cells. *Sci. Rep.* **9**, 10129 (2019).

308   15.   Fujita, K. a *et al.* Decoupling of Receptor and Downstream Signals in the Akt Pathway by

309   Its Low-Pass Filter Characteristics. *Sci. Signal.* **3**, ra56 (2010).

310   16.   Sasagawa, S., Ozaki, Y., Fujita, K. & Kuroda, S. Prediction and validation of the distinct

311   dynamics of transient and sustained ERK activation. *Nat. Cell Biol.* **7**, 365–373 (2005).

312   17.   Goulev, Y. *et al.* Nonlinear feedback drives homeostatic plasticity in H_2O_2 stress

313   response. *Elife* **6**, e2397 (2017).

314   18.   Brewster, J. L. & Gustin, M. C. Hog1: 20 years of discovery and impact. *Sci. Signal.* **7**,

315   re7 (2014).

19. Brewster, J., de Valoir, T., Dwyer, N., Winter, E. & Gustin, M. An osmosensing signal

transduction pathway in yeast. *Science* **259**, 1760–1763 (1993).

20. Westfall, P. J. When the Stress of Your Environment Makes You Go HOG Wild. *Science*

306, 1511–1512 (2004).

21. Saito, H. & Posas, F. Response to Hyperosmotic Stress. *Genetics* **192**, 289–318 (2012).

22. Ferrigno, P., Posas, F., Koepp, D., Saito, H. & Silver, P. A. Regulated nucleo/cytoplasmic

exchange of HOG1 MAPK requires the importin beta homologs NMD5 and XPO1.

EMBO J. **17**, 5606–14 (1998).

23. English, J. G. *et al.* MAPK feedback encodes a switch and timer for tunable stress

adaptation in yeast. *Sci. Signal.* **8**, ra5 (2015).

24. Reiser, V., Ruis, H. & Ammerer, G. Kinase activity-dependent nuclear export opposes

stress-induced nuclear accumulation and retention of Hog1 mitogen-activated protein

kinase in the budding yeast Saccharomyces cerevisiae. *Mol. Biol. Cell* **10**, 1147–1161

(1999).

25. Muzzey, D., Gómez-Uribe, C. A., Mettetal, J. T. & van Oudenaarden, A. A Systems-

Level Analysis of Perfect Adaptation in Yeast Osmoregulation. *Cell* **138**, 160–171 (2009).

26. Macia, J. *et al.* Dynamic Signaling in the Hog1 MAPK Pathway Relies on High Basal

Signal Transduction. *Sci. Signal.* **2**, ra13 (2009).

27. Granados, A. A. *et al.* Distributing tasks via multiple input pathways increase cellular

survival in stress. *Elife* **6**, e21415 (2017).

28. Mitchell, A., Wei, P. & Lim, W. A. Oscillatory stress stimulation uncovers an Achilles

heel of the yeast MAPK signaling network. *Science* **350**, 1379–1383 (2015).

29.   Hersen, P., McClean, M. N., Mahadevan, L. & Ramanathan, S. Signal processing by the

HOG MAP kinase pathway. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 7165–70 (2008).

30.   Westfall, P. J., Patterson, J. C., Chen, R. E. & Thorner, J. Stress resistance and signal

fidelity independent of nuclear MAPK function. *Proc. Natl. Acad. Sci. U. S. A.* **105**,

12212–12217 (2008).

31.   Westfall, P. J. & Thorner, J. Analysis of Mitogen-Activated Protein Kinase Signaling

Specificity in Response to Hyperosmotic Stress: Use of an Analog-Sensitive HOG1

Allele. *Eukaryot. Cell* **5**, 1215–1228 (2006).

32.   Wurgler-Murphy, S. M., Maeda, T., Witten, E. A. & Saito, H. Regulation of the

Saccharomyces cerevisiae HOG1 mitogen-activated protein kinase by the PTP2 and PTP3

protein tyrosine phosphatases. *Mol. Cell. Biol.* **17**, 1289–97 (1997).

33.   Warmka, J., Hanneman, J., Lee, J., Amin, D. & Ota, I. Ptc1, a Type 2C Ser/Thr

Phosphatase, Inactivates the HOG Pathway by Dephosphorylating the Mitogen-Activated

Protein Kinase Hog1. *Mol. Cell. Biol.* **21**, 51–60 (2001).

34.   Mattison, C. P. & Ota, I. M. Two protein tyrosine phosphatases, Ptp2 and Ptp3, modulate

the subcellular localization of the Hog1 MAP kinase in yeast. *Genes Dev.* **14**, 1229–1235

(2000).

35.   Maeda, T., Wurgler-Murphy, S. M. & Saito, H. A two-component system that regulates an

osmosensing MAP kinase cascade in yeast. *Nature* **369**, 242–245 (1994).

36.   Young, C., Mapes, J., Hanneman, J., Al-Zarban, S. & Ota, I. Role of Ptc2 type 2C Ser/Thr

phosphatase in yeast high-osmolarity glycerol pathway inactivation. *Eukaryot. Cell* **1**,

1032–40 (2002).

37. Jacoby, T. *et al.* Two protein-tyrosine phosphatases inactivate the osmotic stress response

pathway in yeast by targeting the mitogen-activated protein kinase, Hog1. *J. Biol. Chem.*

272, 17749–17755 (1997).

38. Martín, H., Flández, M., Nombela, C. & Molina, M. Protein phosphatases in MAPK

signalling: we keep learning from yeast. *Mol. Microbiol.* **58**, 6–16 (2005).

39. Mettetal, J. T. *et al.* The Frequency Dependence of Osmo-Adaptation in Saccharomyces

cerevisiae. *Science* **319**, 482–484 (2008).

40. Neuert, G. *et al.* Systematic Identification of Signal-Activated Stochastic Gene

Regulation. *Science* **339**, 584–587 (2013).

41. Lewis, J. G., Learmonth, R. P. & Watson, K. Induction of heat, freezing and salt tolerance

by heat and salt shock in Saccharomyces cerevisiae. *Microbiology* **141**, 687–694 (1995).

42. Berry, D. B. & Gasch, A. P. Stress-activated Genomic Expression Changes Serve a

Preparative Role for Impending Stress in Yeast. *Mol. Biol. Cell* **19**, 4580–4587 (2008).

43. Berry, D. B. *et al.* Multiple Means to the Same End: The Genetic Basis of Acquired Stress

Resistance in Yeast. *PLoS Genet.* **7**, e1002353 (2011).

44. Bell, M. & Engelberg, D. Phosphorylation of Tyr-176 of the Yeast MAPK Hog1/p38 Is

Not Vital for Hog1 Biological Activity. *J. Biol. Chem.* **278**, 14603–14606 (2003).

45. Lim, W., Mayer, B. & Pawson, T. *Cell signaling : principles and mechanisms*. (Garland

Science, 2015).

46. Fahs, S., Lujan, P. & Köhn, M. Approaches to Study Phosphatases. *ACS Chem. Biol.* **11**,

2944–2961 (2016).

47. Landry, B. P., Palanki, R., Dyulgyarov, N., Hartsough, L. A. & Tabor, J. J. Phosphatase

activity tunes two-component system sensor detection threshold. *Nat. Commun.* **9**, 1433

(2018).

48. Brachmann, C. B. *et al.* Designer deletion strains derived from Saccharomyces cerevisiae

S288C: A useful set of strains and plasmids for PCR-mediated gene disruption and other

applications *Yeast* **14**, 115–132 (1998).

49. Winzeler, E. A. *et al.* Functional characterization of the S. cerevisiae genome by gene

deletion and parallel analysis. *Science* **285**, 901–906 (1999).

50. Gietz, R. D. & Woods, R. A. Transformation of yeast by lithium acetate/single-stranded

carrier DNA/polyethylene glycol method. *Methods Enzymol.* **350**, 87–96 (2002).

51. Kesler, B., Li, G., Thiemicke, A., Venkat, R. & Neuert, G. Automated cell boundary and

3D nuclear segmentation of cells in suspension. *Sci. Rep.* **9**, 10237 (2019).

52. Munsky, B., Li, G., Fox, Z. R., Shepherd, D. P. & Neuert, G. Distribution shapes govern

the discovery of predictive models for gene regulation. *Proc. Natl. Acad. Sci.* **115**, 7533–

7538 (2018).

53. Kushnirov, V. V. Rapid and reliable protein extraction from yeast. *Yeast* **16**, 857–860

(2000).

Figures

Fig. 1. Hog1 signaling and cell survival are sensitive to the rate of preconditioning osmotic stress application. (**a**) Schematic of the budding yeast high osmolarity glycerol (HOG) response. (**b**) Preconditioning protection assay workflow indicating first stress treatments (left plot), high-stress exposure (middle panel), and colony formation readout (right panel). (**c**) High-stress survival as a function of each first treatment relative to the untreated first stress condition. Bars and errors are means and standard deviation from three biological replicates. * = statistically significant by Kolmogorov-Smirnov (KS) test ($p < 0.05$). NS = not significant. (**d**)

408 Treatment concentration over time. (**e**) Treatment rate over time. (**f**) Hog1 nuclear localization

409 during the treatments depicted in "d" and "e". Insert highlights localization pattern in the

410 quadratic treated sample. Lines represent means and shaded error represents the standard

411 deviation from 3-4 biological replicates.

Fig. 2. A rate threshold condition regulates signaling nuclear localization. (**a**) Quadratic

signaling response from Fig. 1f plotted as a function of treatment concentration. The intersection

points between the dashed lines representing Hog1 nuclear localization baseline (black) and

accumulation pattern either expected for step treatments (blue) or measured in quadratic

treatment (red) indicate concentration thresholds (CTh). The apparent quadratic CTh is listed in

bold red font. (**b**) Quadratic treatment response from Fig. 1F plotted as a function of treatment

rate. A rate threshold (RTh) (bold red font) was identified via the approach used for CTh

determination. (**c**) NaCl concentration change during ramp treatments. CTh is marked. (**d**) NaCl

rate during ramp treatments. RTh is indicated. (**e**) Signaling response to ramp treatments plotted

as a function of time. (**f**) Signaling response to ramp treatments plotted as a function of treatment

concentration. The concentrations corresponding with Hog1 nuclear localization in the ramps

where significant activation was detected are indicated in bold font in the top left corner of the

plot color-coded to match the corresponding treatment. The concentration corresponding with

Hog1 nuclear localization calculated by combining the data from all the biological replicates in

both activating ramps was 57 +/- 14.5 mM. (**g**) AND logic model indicating the order of

condition completion for the ramps exceeding RTh.

Fig. 3. Hog1 nuclear localization timing depends on the rate threshold in quadratic

treatments. (**a**) Quadratic NaCl concentration change indicating the time that the concentration

threshold (t_{CTh}) is satisfied. (**b**) NaCl rate change indicating the time the treatment reaches the

rate threshold (t_{RTh}). (**c**) Signaling response to the quadratic treatment depicted in "a" and "b"

indicating t_{CTh} and t_{RTh}. (**d**) Two additional quadratic treatments used to study how treatment rate

affects Hog1 translocation. A 25 min quadratic to 0.4 M NaCl is shown in blue and a 40 min

quadratic to 0.4 M is shown in red. The concentration threshold (C_{Th}) is indicated by the black

dotted line. (**e**) Treatment rate during the quadratic treatments depicted in "d". The rate threshold

(R_{Th}) is indicated in the black dotted line. (**f**) Hog1 nuclear localization during the quadratic

treatments depicted in "d" and "e" plotted as a function of time. The time that each treatment

satisfies the C_{Th} and R_{Th} is indicated by the blue dotted lines (for the 25 min quadratic to 0.4 M

NaCl) and the red dotted lines (for the 40 min quadratic to 0.4 M NaCl). (**g**) Hog1 activation

during the two additional quadratic treatments plotted as a function of concentration. The

concentration thresholds correlating with Hog1 nuclear localization during these treatments are

indicated in font color coded to match the corresponding treatment. (**h**) Hog1 translocation

during the additional quadratic treatments plotted as a function of rate. The rate thresholds

correlating with Hog1 activation during these treatments are indicated as in **d**. (**i**) AND logic

model for quadratic stimulation.

Fig. 4. Step and fast ramp stimulations do not enable the detection of threshold conditions.

(a) Step treatment concentration over time indicating C_{Th}. (b) Treatment rate during step

treatment indicating R_{Th}. (c) Hog1 nuclear localization during step stimulation indicating the

time step treatment satisfied the C_{Th} and R_{Th}. (d) Fast ramp treatment concentration over time

483   indicating C_{Th}. **(e)** Treatment rate during fast ramp treatment indicating R_{Th}. **(f)** Hog1

484   translocation during fast ramp stimulation indicating the time step treatment satisfied the C_{Th} and

485   R_{Th}. **(g)** AND logic model indicating the order in which the step and fast ramp treatments satisfy

486   the C_{Th} and R_{Th} conditions.

 Fig. 5. Hog1 localizes to the nucleus at an earlier time point during quadratic treatment in

488 **the *ptp2Δ* strain.** (**a**) Simplified model of the HOG pathway. Hypothesized regulators of the rate

threshold are indicated in red and cyan. (**b**) Quadratic NaCl concentration treatment indicating C_{Th} and $t_{C_{Th}}$. (**c**) NaCl rate during quadratic treatment with R_{Th} and $t_{R_{Th}}$ marked. The treatment rate at the time C_{Th} is satisfied (R at C_{Th}) is also indicated. (**d**) Signaling response to quadratic stimulation in WT (measured, blue) and rate threshold knock out (R_{Th} KO) mutant (predicted, red). $t_{C_{Th}}$ and $t_{R_{Th}}$ are marked. (**e**) Signaling response to quadratic treatment in *WT*, *ptp2Δ*, and *ptp3Δ* cells.

509
510 **Fig. 6. The rate threshold for Hog1 nuclear localization is specifically dependent on the**

511 **Ptp2 phosphatase.** (**a**) Signaling response in the *WT*, *ptp2Δ*, and *ptp3Δ* strains plotted as a

512 function of NaCl treatment rate. The fitted RTh in each strain is listed in bold type. (**b**) Response

513 in each strain plotted as a function of NaCl concentration. The apparent C_{Th} in each strain is

514 listed. **(c)** Step stimulation indicating treatment concentration over time. **(d)** Hog1 translocation

515 in response to the stimulation in "c" in each strain. **(e)** Model of Hog1 activation in the *ptp2Δ*

516 strain indicating the apparent loss of regulation via the rate threshold.

Fig. 7. 25 minute quadratic treatment to 0.2M NaCl fails to induce Hog1 nuclear localization in the *PTP2* over-expressions strains. (a) HA$_3$-tagged Ptp2 expression levels in the *ptp2Δ* strain carrying either an empty vector (empty, red) or one of three plasmids named based on the gene upstream regulatory DNA driving *PTP2* expression [*PTP2* (blue), *ADH1* (cyan), and *TEF1* (black)] were scored via anti-HA immunoblotting relative to loading control (Actin). *Top:* 5 min imaging exposure, *bottom:* 15 min imaging exposure. **(b)** Ptp2 expression levels and predicted rate thresholds in the *ptp2Δ* strain carrying either an empty vector ("empty") or expressing *PTP2* from the listed expression plasmid. **(c)** Quadratic treatment concentration

over time. **(d)** Rate over time for the treatment profile depicted in "b". **(e)** Hog1 nuclear

localization in the *ptp2Δ* strain containing an empty vector (red) or expressing *PTP2* from the

upstream regulatory DNA sequences of one of three genes [*PTP2* (blue), *ADH1* (cyan), or *TEF1*

(black)]. **(f)** The rate threshold corresponding with Hog1 nuclear localization in each *PTP2*

expression variant strain determined using the 25 min quadratic treatment to 0.2M NaCl. The

rates corresponding with Hog1 nuclear localization are indicated in bold font in the top left

corner of the plot color-coded to the match the corresponding *PTP2* expression variant strain.

Fig. 8. Ptp2 expression levels fine-tune the stimulation rate required to activate Hog1

signaling. (**a**) NaCl concentration during a 0.4M NaCl quadratic stimulation indicating the C_{Th}

and t_{CTh}. (**b**) NaCl rate during these stimulations indicating R_{Th} and t_{RTh}. (**c**) Hog1 nuclear

localization response to the quadratic stimulation in "a" and "b" depicted as a function of time in

each *PTP2* expression strain. (**d**) Signaling response plotted as a function of NaCl treatment rate.

The R_{Th} in each strain are indicated in bold.

Fig. 9. Model of signaling regulation by threshold conditions. The timing of signaling

activation (bottom) can correspond with both of the threshold conditions or either the C_{Th} or R_{Th}

alone depending on stimulation type (top).

Methods

Yeast constructs and strains

All plasmids used in this study (Supplementary Table 1) were constructed via standard restriction enzyme cloning. Appropriate restriction sites were either included in custom synthesized oligonucleotides (IDT) or introduced by PCR with Q5 High-Fidelity Polymerase (NEB) according to the manufacturer's instructions. All constructs were sequence verified.

The yeast strains used in this study (Supplementary Table 2) were derived from the haploid *Saccharomyces cerevisiae* BY4741 strain[48]. All yeast genomic deletions and integrations were confirmed via PCR. The *PTP2* and *PTP3* genes have been replaced with the kanamycin resistance cassette (*KANMX4*) in the *ptp2Δ* and *ptp3Δ* strains[49]. Strains used to track Hog1-YFP nuclear localization were obtained from a previous study[39] or generated by C-terminally tagging the endogenous *HOG1* with sequences encoding yellow fluorescent protein (YFP) [33] through homologous DNA recombination. Ptp2 expression and empty vector control plasmids were introduced into yeast via high efficiency yeast transformation[50].

Yeast growth

Three days before an experiment, yeast cells from a glycerol stock stored at −80 °C were streaked out on Complete Synthetic Media (CSM) (Formedium) or CSM lacking Uracil (CSM-Ura) (Formedium) for selection as needed. The day before the experiment, a single colony from the plate was inoculated in liquid media (pre-culture). After 2 - 6 hr, the optical density at 600 nm (OD_{600}) of the pre-culture was measured and diluted into fresh liquid culture to reach an OD_{600} of 0.3-0.5 (depending on the experimental application) the next morning. These OD_{600} values represent 1.5-2.5 x 10^6 cells/mL.

Pump profile generation

622   Pump profiles designed to deliver varying rate salt treatments were generated using an in-

623   house developed MATLAB script that accounts for stock NaCl concentration, pump withdraw

624   rate, volume in the container in which the profile is generated, and culture volume change[14].

625   Profiles were loaded onto programmable syringe pumps (New Era Pump Systems) prior to

626   treatment.

High-stress preconditioning assay

High-stress preconditioning assays were designed based on previous work[42,43]. Initial

stress treatments began when cells in culture grown overnight reached an OD_{600} of 0.3. The

untreated and 1 hr 0.4 M NaCl pulse 1^{st} stress treatments were chosen as negative and positive

controls. Ramp treatments were delivered directly to cells in culture via tubing connecting

culture flasks to stock CSM NaCl solution placed in programmable syringe pumps (New Era

Pump Systems). These treatments were designed in order to deliver salt at a rate of either 4 mM

NaCl/min or 16 mM NaCl/min to a final concentration of 0.4 M NaCl. The time each ramp

treatment was held at 0.4 M NaCl was determined such that all first stress treatments that

included NaCl received the same cumulative salt exposure (see calculations below). Because

treatment duration varied, first stress treatment start times were adjusted such that all treatments

ended at the same time. Upon completion of the first treatment, culture OD_{600} was measured and

14mL of each culture was transferred to 15 mL conical tubes, and cells were harvested by

centrifugation for 5 min at 3,000 x g. The supernatant was removed, and cells were resuspended

in 1 mL sterile H_2O, transferred to 1.5 mL microcentrifuge tubes (Eppendorf) and pelleted by

centrifugation for 5 min at 21,300 x g. Supernatant was removed and pellets were resuspended

in 100 µL sterile H₂O. 5 µL of each resuspended culture was transferred to 145 µL of either

CSM (0 M NaCl second stress) or 2 M NaCl CSM (high second stress) in the first well of each

row of a 96-well plate. Once all cultures were transferred, each was diluted 1:2 across the wells

of each plate column containing 100 µL of each second stress treatment. 96-well plates were

incubated at 30 °C with shaking at 250 rpm for 2.5 hrs. After this time, 150 µL CSM was added

to each well. Culture in the wells was spotted using a replica plater (Sigma R2508) onto a 15 cm

2% agar CSM plate to enable the determination of relative high-stress survival. To quantitate

differences in high second stress survival, culture from the first well of each row (containing the

highest concentration of cells from each sample) was diluted 1:2,000, and 600 µL of each

treatment was spread onto a 15 cm 2% agar CSM plate.

Plates were incubated for 2-3 days at 30 °C and imaged using a ChemiDoc MP (BioRad).

Individual colonies from the plates generated to enable high second stress survival quantitation

were counted using in-house developed MATLAB scripts. In brief, after the user is prompted to

define the plate diameter by clicking, the scripts process the plate images to generate a uniform

background. The image is converted to binary with the background and colonies converted into

opposite values and the number of centroids within a user-defined size range in the binary colony

value are recorded.

For each first stress treatment, survival was calculated manually by dividing the number of

colonies formed from culture exposed to the 2 M NaCl second stress by the number formed from

exposure to 0 M NaCl second stress. Relative survival mean and standard deviation were

calculated by normalizing the survival resulting from each first stress to mean survival in the

untreated first stress culture for three independent biological replicates. Statistical significance

was calculated using the Kolmogorov-Smirnov (K-S) test.Example Cumulative Salt Exposure

Calculation

-1 hr 0.4 M NaCl step = 60 min X 400 mM NaCl = 24,000 mM NaClmin cumulative salt exposure

-4 mM NaCl/min ramp to 400 mM = (400 mM X 100 min)/2 = 20,0000 mM NaCl min cumulative

salt exposure

24,0000 mM NaCl min – 20,000 mM NaClmin = 4,000 mM NaCl min more salt exposure

needed

4,000 mMNaCl min/400 mM = 10 min hold after ramp

Time-lapse microscopy

Treatments

3 ml of yeast cells expressing *HOG1-YFP* from the chromosome grown to an OD_{600} of

0.45-0.55 (2.25-2.75 x 10^6 cells/mL) were pelleted by centrifugation at 500 x g for 6 min. The

HOG1-YFP strain obtained from a previous study[39] was used for the experiments presented in

Figure 2E, Figure 2F, Figure 3F-H, and Figure 4. The *HOG1-YFP* strains generated for this

study were used to collect the data in all other figures. Cell pellets were resuspended in 50 µl

CSM and loaded into a flow chamber as previously described[25].

Three types of perturbations were used in time-lapse microscopy treatments: pulse, linear

ramp and quadratic gradients. To deliver the pulse treatments, flow chambers containing cells

were connected via tubing to a conical tube containing the CSM NaCl treatment to be delivered

on one end and a programmable syringe pump (New Era Pump Systems) set to withdraw media

on the other. Ramp and quadratic perturbations were set up as described previously[14]. In brief,

these perturbations were generated in a beaker containing CSM subjected to constant mixing on

a stir plate by delivering CSM NaCl stock solution in pre-programmed steps via syringe pump

(New Era Pump Systems). Media was drawn from this beaker to the flow chamber over the time

course of treatment profile generation using a syringe pump programmed to withdraw media at a

rate of 0.1 mL/min. At the completion of all NaCl treatments, cells were returned to CSM by

switching the beaker or conical containing CSM NaCl to a conical containing CSM only.

Image acquisition

The Micro-Manager program (https://micro-manager.org/) version 1.4 was used to

control a Nikon T*i* Eclipse epi-fluorescent microscope equipped with perfect focus (Nikon), a

100x VC DIC lens (Nikon), a fluorescent filter for YFP (Semrock), an X-cite fluorescent light

source (Excelitas) and an Orca Flash 4v2 CMOS camera (Hamamatsu). YFP images were

collected every 1 min with an exposure of 20 ms. Bright-field (TRANS) images with an

exposure of 10 ms were collected every 10 sec during each time-lapse experiment. 3-4 YFP

images of cells loaded on the flow chamber were taken prior to the start of each treatment to

enable Hog1 nuclear localization baseline level determination.

Image analysis

Image segmentation was performed in two-steps. First, YFP images (detecting Hog1-

YFP) were smoothed and background corrected. A threshold was set to identify the brightest

pixels for each cell. The region containing the 100 brightest pixels in each cell was then used as

an intracellular marker. Second, TRANS images were smoothed, background corrected and

overlain with the corresponding YFP image (or the YFP image from the previous time point in

the case were no YFP image was taken). A watershed algorithm was applied to these overlain

images to segment and label the objects (cells). After segmentation, objects that deviate

significantly in size from the average cell or on the border of the image are removed, resulting in

an image with segmented cells. This process was repeated for each image. After segmentation,

the centroid of each cell was computed and stored. The distance between the centroids of each

cell in consecutive images were compared. The centroids that show the smallest distance

difference between two consecutive images are considered the same cell at two different time

points. This whole procedure is repeated for each image resulting in single cell trajectories.

Each single cell trace was visually inspected, and cells exhibiting large fluctuations in YFP or

TRANS signal over the time course were removed. In a typical experiment, 20-30% of a starting

population of 200-450 single cells remained after this filtering step for further processing.

To determine Hog1 nuclear localization in each cell, the average per pixel fluorescent

intensity of the whole cell (I_w) and of the top 100 brightest fluorescent pixels (I_t) was calculated.

In addition, fluorescent signal per pixel of the camera background (I_b) was recorded. Hog1

nuclear localization was calculated as $Hog1(t) = [(I_t(t) - I_b) / (I_w(t) - I_b)]$. The value for Hog1(t =

0) defines Hog1 nuclear localization baseline level. Single cell traces were smoothed and

nuclear localization over the time course was determined by subtracting Hog1(t=0) from

Hog1(t). For cell volume measurements, volume change relative to the volume at the beginning

of the experiment was calculated. For both the single cell volume and Hog1(t) fluorescent

traces, the median and the average median distance (the equivalent of the standard deviation if

the median is used instead of the mean) were computed to put less weight on sporadic outlier

cells due to the image segmentation process.

The in-house developed codes used for image segmentation[51] and time-lapse image

analysis[14,40,52] have been previously described.

Determination of concentration and rate thresholds

The rate and concentration thresholds were determined by plotting the median Hog1

nuclear localization in each biological replicate as a function of treatment concentration or rate,

respectively. Two points were manually selected from each of these plots demarcating the

beginning and end point of robust Hog1 nuclear enrichment. The data between these points was

automatically fit to the linear function ($y = mx + b$). The "x" value (concentration or rate) at the

intersection point of a horizontal line at Hog1 = 0 (baseline) and the line fitting Hog1 nuclear

enrichment pattern (activation) was determined. The average and standard deviation of the "x"

values from the replicates for each treatment profile represent the concentration and rate

thresholds corresponding with Hog1 nuclear localization in those treatment profiles. The average

slope ("m") and y-intercept ("b") from the linear function fit to the Hog1 activation data for each

biological replicate obtained from a particular treatment profile were used to generate the dotted

lines overlaying Hog1 nuclear enrichment in the threshold determination figures.

Immunoblotting

Mild alkali treatment and heating in standard electrophoresis buffer[53] were used to extract

total protein from yeast cell pellets containing 1 OD_{600} unit (5×10^6 cells). Proteins were

fractionated on a NuPAGE 4-12% Bis-Tris denaturing polyacrylamide gel (Thermo Fisher

Scientific). Gels were equilibrated in transfer buffer (30 mM Bicine, 25 mM Bis-Tris, 1 mM

EDTA, 60 µM chlorobutanol) and electrotransferred to PVDF membranes pre-equilibrated in

transfer buffer (GE, Immobilon-P. 0.45μM). 5% w/v non-fat milk in 1 X Tris-buffered saline

(TBS) (100mM Tris-Cl (pH 7.5), 150 mM NaCl) was used for blocking. HAx3-Ptp2 was

detected using an anti-HA HRP conjugate 3F10 (Roche) at a 1:5,000 dilution. Endogenous actin

(used as a loading control) was detected using a 1:5,000 dilution of anti-ß-actin antibody (Abcam

#ab8224) followed by incubation with a 1:2,500 dilution of HRP-conjugated horse anti-mouse

IgG antibody (Cell Signaling #7076S). All antibody incubations were performed in 1 % w/v non-

fat milk blocking in 1X TBS. Antibody signals were visualized using. Enhanced

chemiluminescence (ECL) reagent (GE) was used to detect antibody signals. Signals were

visualized using a ChemiDoc MP Imaging System (BioRad).

Data availability

The datasets generated during and/or analysed during the current study are available from

the corresponding author on reasonable request.

Code availability

Cell segmentation codes have been previously described[51] and are available in the public

dropbox repository:

https://www.dropbox.com/sh/egb27tsgk6fpixf/AADaJ8DSjab_c0gU7N7ZF0Zba?dl=0.

Image processing codes have also been previously described[14,40,52]. All codes used for

image processing and analysis in this study will be made available upon reasonable request.

Acknowledgements

We would like to thank Mr. Cleland, Mr. Hughes, Mr. Venkat and Drs. Weil, Tansey,

Wadzinski, Carrasco, and Colbran for feedback on the study and manuscript. We would also like

to thank the Vanderbilt Molecular Biology core facility for providing access to and technical

assistance with instrumentation necessary to complete this work. This work was supported by

the NIH Director's New Innovator Award (1DP2GM114849), the NIGMS (5R01GM115892),

the Advanced Computing Center for Research and Education (ACCRE) (1S10OD023680-01),

and Vanderbilt University Startup Funds. AJ was supported by the Ion Channels and

Transporters Training Grant (T32NS007491), BK was supported by the Molecular Biophysics

Training Grant (5T32GM008320) and AT was supported by an AHA predoctoral fellowship

(18PRE34050016). The content of this publication is solely the responsibility of the authors and

does not necessarily represent the official views of the National Institutes of Health.

Author contributions

Conceptualization: A.N.J. and G.N. Data Curation: A.N.J., H.J., and G.N., Formal

Analysis: A.N.J., H.J., and G.N., Funding Acquisition: G.N. Investigation: A.N.J., G.L., H.J.,

A.T., Methodology: A.N.J., G.L., H.J., and A.T., Project administration: A.N.J. and G.N.,

Resources: A.N.J., D.R., and G.N., Software: B.K., H.J., and G.N., Supervision: G.N.,

Validation: A.N.J. and H.J., Visualization: A.N.J., Writing-original draft: A.N.J., Writing-

reviewing and editing: all authors.

<u>Competing interests</u>

802   The authors declare no competing financial interest.

803   **<u>Additional Information</u>**

804   **Supplementary Information** is available for this paper

805   **Correspondence and requests for materials** should be addressed to G.N.

808