Binding Thermodynamics and Kinetics Calculations Using Chemical Host and Guest: A Comprehensive Picture of Molecular Recognition

Zhiye Tang and Chia-en A. Chang*

Department of Chemistry, University of California, Riverside, CA92521

Telephone: (951) 827-7263

Email: chiaenc@ucr.edu

Understanding the fine balance between changes of entropy and enthalpy and the competition between a guest and water molecules in molecular binding is crucial in fundamental studies and practical applications. Experiments provide measurements. However, illustrating the binding/unbinding processes gives a complete picture of molecular recognition not directly available from experiments, and computational methods bridge the gaps. Here, we investigated guest association/dissociation with β-cyclodextrin (β-CD) by using microsecond-timescale molecular dynamics (MD) simulations, post-analysis and numerical calculations. We computed association and dissociation rate constants, enthalpy, and solvent and solute entropy of binding. All the computed values of k_{on}, k_{off}, ΔH, ΔS, and ΔG using GAFF-CD and q4MD-CD force fields for β-CD could be compared with experimental data directly and agreed reasonably with experiment findings. Both force fields resulted in similar computed ΔG from independently computed kinetics rates, $\Delta G = -RT\ln(k_{on} \cdot C° / k_{off})$, and thermodynamics properties, $\Delta G = \Delta H - T\Delta S$. The water entropy calculations show that entropy gain of desolvating water molecules are a major driving force, and both force fields have the same strength of non-polar attractions between solutes and β-CD as well. Water molecules play a crucial role in guest binding to β-CD. However, collective water/β-CD motions could contribute to different computed k_{on} and ΔH values by different force fields, mainly because the parameters of β-CD provide different motions of β-CD, hydrogen-bond networks of water molecules in the cavity of free β-CD and the strength of desolvation penalty. As a result, q4MD-CD suggests that guest binding is mostly driven by enthalpy, while GAFF-CD shows that gaining entropy is the major driven force of binding. The study further interprets experiments, deepens our understanding of ligand binding, and suggests strategies for force field parameterization.

Keywords: protein-ligand binding affinity; cell method; drug design; H-bond, hydration shell dynamics; host-guest

1. Introduction

Molecular recognition determines the binding of two molecules — a common phenomenon in chemical and biological processes. Thus, understanding molecular recognition is of interest in both fundamental studies and has practical applications in chemical industries and drug discovery. Molecular binding can be characterized by thermodynamic and kinetic properties, which are governed by factors such as polar and non-polar interactions between solutes, solvent effects, molecular diffusion and conformational fluctuations. Molecular binding affinity is a straightforward characterizer of recognition and can be obtained experimentally by enthalpimetry such as calorimetry [1-2]. However, it is difficult to measure the binding entropy directly from experiments, and it is impossible to measure separate entropy contributions from the solvent and solute. Recent studies have revealed the importance of kinetic properties [3-5] and showed that drug efficacy is sometimes correlated with the kinetic properties better than binding affinity [6-7]. Investigating residence time [8-10] was proposed in the last decade and has become a useful criterion in predicting drug efficacy. Efforts have been devoted to studies of kinetic–structure relations [11-14]. Computational methods, which have the advantage of atomistic resolution, are useful tools to investigate the fundamentals of thermodynamics and kinetics. With the recent breakthrough in computational power, molecular dynamics (MD) is able to simulate up to microsecond or millisecond timescale molecular motions [15-16]. Therefore, computational methods can be used to sample a larger time scale of dynamics of host–guest systems and take advantage of numerical methods to directly extract the enthalpic and entropic profiles of both the solvent and solute, as well as the kinetics from unbiased long-time-scale MD. New findings observed from chemical host–guest systems with theoretically sound computational methods has advanced our knowledge of molecular recognition and brought new insights into ligand–protein systems.

Cyclodextrins (CDs) are a class of cyclic oligosaccharide compounds that can be obtained by degradation of starch by α-1,4-glucan-glycosyltransferases. Depending on the number of glucopyranose units (D-glucose), these compounds can be classified into 6 units (α-CD), 7 units (β-CD), 8 units (γ-CD) and CDs of more glucopyranose units. In β-CD (Figure 1), the 7 glucopyranose units (D-glucose) enclose a cavity with a diameter of about 6.5 Å. The glucopyranose units bring the center of the cavity a hydrophobic surface of the carbon chains in the glucopyranose, whereas the rims of β-CD consist of hydrophilic hydroxyl groups. Notably, the wide and narrow rims of β-CD are asymmetrical. Because of the structure and size of the cavity, β-CD can host a wide variety of guest molecules via nonpolar or polar attractions. With these properties, β-CD and its derivatives have many applications in many fields, such

as the cosmetic industry, pharmaceuticals, catalysis, and the food and agricultural industries [17-23]. Thus, the experimental thermodynamics and kinetics data are available for a variety of β-CD complexes [24-30].

β-CD consists of 147 atoms, so we can directly investigate the fundamentals of binding thermodynamics and kinetics by MD simulations. Short MD simulations have been used to study ligand binding affinity, dynamics, H-bonds and hydration properties of β-CD complexes [31-38]. Replica Exchange Molecular Dynamics (REMD) and the Mining Minima 2 method were used to thoroughly sample the bound state of various β-CD complexes and accurately evaluate their binding affinities by using an implicit solvent model [39-40]. QM/MM methods were also used to investigate the binding pose, binding enthalpy, and properties of the β-CD cavity [41-44]. In most of these studies, the conformations of β-CD resemble the crystal structures via shorter than 20 ns MD simulations, and no further dynamic information from experiments or long-time-scale MD of β-CD complexes. In addition, although the binding affinity decomposition into enthalpy and entropy has been achieved for some ligands, the use of an implicit solvent model and previous computation techniques cannot obtain both solvent and solute entropy contributions that allow direct comparison with experimental measured ΔH and ΔS. It is still difficult to sample multiple association/dissociation events for kinetic rate calculations [45].

The present study applied microsecond-timescale MD simulations with GPU acceleration to compute binding kinetics and thermodynamics of 7 guests with β-CD to reveal the binding mechanisms. Using two force fields for β-CD, GAFF-CD and q4MD-CD, we performed the simulations that sampled multiple association/dissociation events of the complexes to compute k_{on} and k_{off}, together with MD of free β-CD and guests and a water box to compute entropy and enthalpy from both the contribution of solvent and solutes. The non-polar interactions between solutes and the water entropy gains on binding were the major driving forces of association, regardless of the use of force fields. However, GAFF-CD provided slightly more flexible β-CD, which produced more complicated H-bond networks between the free β-CD and water molecules in the first hydration shell, thereby resulting in slower k_{on} and larger desolvation penalty. The study also gives details regarding multiple guest association and dissociation pathways and explains how each decomposed energetic and entropic term contributes to binding kinetics/thermodynamics and binding mechanisms.

2. Method

Molecular Systems and Force Fields. We selected 4 weak binders (1-butanol, t-butanol, 1-propanol, methyl butyrate) and 3 strong binders (aspirin, 1-naphthyl ethanol, 2-naphthyl ethanol) as the guest

molecules (ligands) for the β-CD host–guest complex. The set of data have similar experimental binding affinity (ΔG) by both kinetics and thermodynamic measurement, and the available rate constants are accessible by MD for sampling multiple association and dissociation events. Structures of guests were manually created by using Vega ZZ [46] and the structure of β-CD was obtained from the Cambridge Crystal Data Center (PDB ID: WEWTOJ) (Figure 1). We used two different force fields for β-CD: Amber general force field (GAFF),[47] termed GAFF-CD, and q4MD-CD force field [48]. For simulations with GAFF-CD, we took the partial charge of β-CD from a previous work [40]. For simulations using q4MD-CD, the partial charge of β-CD was provided in the force field. The guests were modeled by only GAFF and the partial charges were computed by using optimization and then ChelpG atomic charge calculation with the Gaussian package [49] at 6-31+g(d,p)/B3LYP level. Initial conformations of the complexes were obtained by manually locating the ligand in the center of the cavity of β-CD. All MD simulations and calculations were repeated in the same setting for the free and bound β-CD with GAFF-CD and q4MD-CD.

Molecular Dynamics Simulations. We performed microsecond-timescale MD runs for a total of 1 to 11 μs simulations (Table SI 1) for each β-CD-ligand system, that is, on the complex, the free β-CD, the free ligand and an empty water box [50]. For each run, the system was first solvated with exactly 1737 TIP3P water molecules (about 12Å away from the solute) and then optimized to eliminate possible clashes. The water molecules were then equilibrated at 298K for 1 ns, followed by an equilibration of the entire system from 200K to 298K for 150 ps. By using Amber 14 with GPU acceleration [47, 51], long production runs were performed at 298K maintained by a Langevin thermostat [52-53] in an NPT (isothermal-isobaric) ensemble. A frame was saved every 1 ps for each run. For post-analysis and numerical calculations, we resaved a frame every 10 ps, and the trajectories were visualized by VMD [54]. Table SI 1 summarizes the total lengths of all MD simulations and numbers of frames used in thermodynamics calculations. We also performed 50-ns MD runs for free β-CD in a vacuum by using GAFF-CD and q4MD-CD to examine the force field parameters, as detailed in SI Section 2.

Binding Enthalpy Calculation. The binding enthalpy (ΔH) was calculated by $\Delta H = <E>_{Complex} + <E>_{Water} - <E>_{Host} - <E>_{Guest}$ [50] with the periodicity of the water box, where $<E>_{Complex}$, $<E>_{Water}$, $<E>_{Host}$ and $<E>_{Guest}$ are the averaged potential energy of all molecules from the trajectories of the complex, water, host and guest, respectively. Note that only the bound state conformations from the trajectories of the complexes were used to compute $<E>_{Complex}$. The kinetic energy will be cancelled, so it is not evaluated

here. The pressure-volume term is negligible in a solvated system. The computed decomposed enthalpy terms are reported in SI Section 3. We also computed the surface area, number of H-bonds, and number of solvation water (detailed in SI section 4) to explain our decomposed enthalpy terms.

Solute Entropy Calculation. Solute entropy change (ΔS_{Solute}) and water entropy change (ΔS_{Water}) are computed separately for binding entropy, $\Delta S = \Delta S_{Solute} + \Delta S_{Water}$, where ΔS_{Solute} was decomposed into host internal (conformational/vibrational) entropy change (ΔS_{Host}), guest internal entropy change ($\Delta S_{Guest\ Int}$) and guest external (translational/rotational) entropy change ($\Delta S_{Guest\ Ext}$), illustrated as $\Delta S_{Solute} = \Delta S_{Host} + \Delta S_{Guest\ Int} + \Delta S_{Guest\ Ext}$.

Notably, the host retains its external degrees of freedom, so there is no change of external entropy for the host. The internal entropy changes of the host ($\Delta S_{Host} = S_{Host\ Complex} - S_{Host\ Free}$) and guest ($\Delta S_{Guest\ Int} = S_{Guest\ Int\ Complex} - S_{Guest\ Int\ Free}$) were calculated by taking the difference in internal entropies of their free state and bound state by using the Gibbs entropy formula (eq. 1).

$$S_{Int} = -R\int P(\mathbf{\Omega_{Solute}})\ln(P(\mathbf{\Omega_{Solute}}))d\mathbf{\Omega_{Solute}} \qquad \text{eq. 1}$$

where R is the gas constant and $P(\mathbf{\Omega_{Solute}})$ is the probability distribution of conformations ($\mathbf{\Omega_{Solute}}$) defined by key dihedrals in the species, as detailed in SI Section 5.

In the free states, both β-CD and ligands can diffuse freely in solution, but ligand diffusion may be restricted after binding into the cavity of β-CD. The external entropy loss of the ligand was calculated as $\Delta S_{Guest\ Ext} = S_{Guest\ Ext\ Complex} - S_{Guest\ Ext\ Free}$, where $S_{Guest\ Ext\ Free} = R\ln(8\pi^2/C^0)$ is approximately 6.98 kcal/mol at 298 K [55-56]. R is the gas constant and C^0 is the standard concentration (1M). $S_{Guest\ Ext\ Complex}$ is the external entropy of the ligands evaluated by numerical integration after aligning the MD trajectories of the complexes to the crystal structure of β-CD,

$$S_{Guest\ Ext\ Complex} = R\ln(\int \sin(\theta)dxdydzd\phi d\varphi d\theta) \qquad \text{eq. 2}$$

where x, y, z are the Cartesian coordinates of the ligand translation and ϕ, φ, θ are the Euler angles of the ligand rotation. The bin size of x, y, z was 2Å and the bin size of ϕ, φ, θ was 30°. The ϕ and φ were integrated from -π to π and θ was integrated from -π/2 to π/2.

Water Entropy Calculation. We evaluated the water entropy change on ligand binding (ΔS_{Water}) from solvation entropy ($\Delta S_{Solv.}$) of the free β-CD ($\Delta S_{Solv.\ Host}$), free guests ($\Delta S_{Solv.\ Guest}$) and the complexes ($\Delta S_{Solv.\ Complex}$) by $\Delta S_{Solv.} = \Delta S_{Solv.\ Trans} + \Delta S_{Solv.\ Rot} + \Delta S_{Solv.\ Conf}$. The $\Delta S_{Solv.}$ was calculated by the grid cell method [57-59]. In the grid cell method, the water molecules in the water box are treated as if they are vibrating in a local cell and confined by the forces from surrounding water molecules and solutes. In the cell method, ΔS_{Solv} decomposes into vibrational ($\Delta S_{Solv.\ Vib}$) and conformational ($\Delta S_{Solv.\ Conf}$) entropies, and $\Delta S_{Solv.\ Vib}$ further decomposes into translational ($\Delta S_{Solv.\ Trans}$) and rotational ($\Delta S_{Solv.\ Rot}$) entropies. The entropy terms were calculated by using eqs. 6.8 to 6.11 in SI, and an in-house script was developed to perform the calculations. The changes in water entropy on ligand binding were evaluated by $\Delta S_{Water} = \Delta S_{Solv.\ Complex} - \Delta S_{Solv.\ Host} - \Delta S_{Solv.\ Guest}$. The translational ($\Delta S_{Water\ Trans}$), rotational ($\Delta S_{Water\ Rot}$) and conformational ($\Delta S_{Water\ Conf}$) terms of changes of water entropy on ligand binding were evaluated by the same equation. The setting of solvation water entropy calculation is provided in SI Section 6. We counted the number of solvation water molecules in the free guest, free host, and complexes to explain the water entropy calculations, as detailed in SI section 7.

Calculations of the Association and Dissociation Rate Constants. As mentioned previously, our MD simulations can sample multiple association and dissociation events in one long run for each system. We define the bound state for a complex when the center of mass of the ligand is within 7.5 Å from the center of mass of β-CD. Any bound/unbound period that lasted longer than 1.0 ns was counted as one bound/unbound event. Because molecules fluctuate during ligand binding, if a ligand stayed or left β-CD for less than 1.0 ns, we did not consider the motion as a bound/unbound event. The average bound/unbound period lengths were calculated and plugged into eqs. 3 and 4 to compute k_{on} and k_{off}, where *[solute]* is the concentration of solute estimated by the averaged size of the water box, as detailed in SI Section 8. After obtaining k_{on} and k_{off}, we calculated the equilibrium constant $K_{eq} = k_{on} / k_{off}$. We also estimated the diffusion-controlled association rate constants by using diffusion coefficients and radii of host and guest molecules for comparison, as detailed in SI section 9.

$$k_{on} = \frac{1}{\text{average unbound time} \times [\text{solute}]} \qquad\qquad \text{eq. 3}$$

$$k_{off} = \frac{1}{\text{average bound time}} \qquad\qquad \text{eq. 4}$$

Binding Free Energy Calculation. The binding free energies can be computed by using the classical thermodynamics and statistical mechanics equations. We evaluated binding free energies (ΔG_{Comp1}) from computed binding enthalpy and binding entropy, as in eq. 5. We also evaluated binding free energies (ΔG_{Comp2}) from computed association and dissociation rate constants (k_{on} and k_{off}), as in eq. 6.

$$\Delta G_{Comp1} = \Delta H - T\Delta S \qquad\qquad \text{eq. 5}$$

$$\Delta G_{Comp2} = -RT\ln(k_{on} \cdot C^{o} / k_{off}) \qquad\qquad \text{eq. 6}$$

Uncertainty Evaluation of Computed Properties. We evaluated the uncertainties of binding enthalpy (ΔH), the entropy change of water (ΔS_{Water}), internal entropy change of β-cyclodextrin (ΔS_{Host}), internal ($\Delta S_{Guest\ Int}$) and external ($\Delta S_{Guest\ Ext}$) entropy change of guests, total binding entropy (ΔS), binding free energy from eq. 5 (ΔG_{Comp1}), association rate constant (k_{on}), dissociation rate constant (k_{off}), and equilibrium constant (K_{eq}). The details are provided in SI section 10.

3. Results

We used microsecond-timescale MD simulations with an explicit solvent model to compute the binding enthalpies, entropies, and association/dissociation rate constants for β-CD with 7 guest molecules (Figure 1). Binding free energy ΔG was computed by using eq. 5 (ΔG_{Comp1}), from the thermodynamic properties and eq. 6 (ΔG_{Comp2}), from the kinetics binding rate constants. All computed results yielded reasonable agreement with experimental data (Tables 1 and 2, and Figure 2). Two force fields, GAFF-CD and q4MD-CD, were used to assign parameters for β-CD, and all ligands used the GAFF force field. In general, ΔG computed with GAFF-CD is slightly more positive than experimental data, and binding is driving by entropy. In contrast, q4MD-CD yields slightly more negative computed ΔG than experimental data, and binding is driving by enthalpy. Here we were able to separate enthalpy and entropy calculations, and we therefore use directly computed ΔH and -TΔS to investigate their contributions of a guest and β-CD binding.

Binding Enthalpy and Entropy Calculations

The calculated ΔH, -TΔS and ΔG_{Comp1} by using eq. 5 with the GAFF-CD and q4MD-CD force fields are compared with the experimental data in Table 1. The computed ΔG is mostly within 2 kcal/mol of experiments, and they provide a correct trend. Although results from the two force fields reasonably agree with experiment results, GAFF-CD generally underestimated and q4MD-CD overestimated the binding

free energy. The intermolecular van der Waals (vdW) attraction between the β-CD and guest is one major driving force for the complex formation (Tables 3 and 4). The water entropy gain on binding is also a major driving force for the complex formation (Table 5). Interestingly, compared to q4MD-CD, the less negative binding affinities using GAFF-CD are primarily from larger desolvation penalty, resulting in the less negative binding enthalpy. The host became more flexible and gained configuration entropy on binding, which is another favorable factor in guest binding (Table 5). In contrast, as compared with GAFF-CD, q4MD-CD modeled a significantly smaller desolvation penalty and more negative ΔH, which become the main driving force for binding. However, the systems need to pay a higher cost in entropy (–TΔS) because the host became more rigid in the bound state.

Unlike for experiments and most calculations that measured ΔG and ΔH, we independently computed ΔH and –TΔS. Nevertheless, we still observed entropy-enthalpy compensation in our systems with different force fields, so the compensation in these systems indeed has a physical implication and is not the artifact from mathematics of eq. 5. We compared the calculated ΔH and -TΔS for 1-propanol and 1-butanol for which experimental data are available. Both experiments and calculations with GAFF-CD showed a small positive ΔH, which slightly opposed binding, but the systems gained more entropy to compensate losing enthalpy on binding. Although only 2 guests studied here have experimental ΔH and -TΔS, the binding data for other alcohols of β-CD with calorimetry (ITC) and UV experiments commonly show gaining entropy, and enthalpy was not the predominant determinant for binding in all cases (Figure SI 7). The following subsections present results of binding enthalpy of solute-solute and solute-solvent and binding entropy of solutes and solvent.

***Changes of Enthalpy from Different Components*.** We evaluated the absolute values of binding enthalpy by using potential energies <E> from microsecond-timescale MD trajectories of the four species, and the convergences of the enthalpy calculations were first examined (Figures SI 8 to 10). For all simulations, the potential energy reaches a stable value within a fluctuation of 0.4 kcal/mol, which is also within reported experimental uncertainty [24, 28-29, 60-61]. The fluctuations of potential energies of β-CD-1–propanol, β-CD-1–butanol and β-CD-t–butanol complexes are larger with GAFF-CD than the other systems. The fractions of the bound states of these systems are lower than with the other systems; thus, the smaller numbers of sampling result in larger fluctuations.

To understand binding enthalpies, we decomposed the calculated values into various contributions (Table 3). We also provide the decompositions into vdW and Coulombic interactions for ΔH$_{Solute\ Inter}$,

$\Delta H_{Host\ Conf}$, $\Delta H_{Host\text{-}Water}$ and $\Delta H_{Guest\text{-}Water}$ (Tables 4, 6, 7). Regardless of type of force field used, we immediately noticed that $\Delta H_{Host\text{-}Guest}$ data are all favorable (negative), and stronger binders such as naphthyl ethanol and aspirin have large negative $\Delta H_{Host\text{-}Guest}$ value, ~ -30 kcal/mol (Table 3). However, $\Delta H_{Desolvation}$ largely compensates the solute-solute attraction, for significantly smaller ΔH, ranging from 3.0 to -1.4 kcal/mol with GAFF-CD and -0.9 to -5.7 kcal/mol with q4MD-CD. The computed ΔH with GAFF-CD yielded positive values for weak binders (ΔG > -2.0 kcal/mol), which has been seen in experiments. In contrast, ΔH values are all negative with q4MD-CD. On binding, both β-CD and the guest desolvate water molecules and lose interaction with water molecules; thus, $\Delta H_{Host\text{-}Water}$ and $\Delta H_{Guest\text{-}Water}$ values are all positive. The values are larger for stronger binders; presumably, guests such as naphthyl ethanols have larger size as compared with weaker/small binders such as propanol. The water molecules released after binding regain interactions with other water molecules, thereby resulting in all negative $\Delta H_{Water\text{-}Water}$ values. However, this term is not negative enough to counterbalance the loss from water-solute interactions. As a result, desolvation enthalpy is inevitably all positive and becomes the major force that opposes binding. Of note, the desolvation penalty from the polar contribution is the major force that opposes binding with GAFF-CD, which results from losing more intermolecular H-bonds between water molecules and β-CD (Table 7). The partial charges and the non-polar attractions between guests and β-CD modeled with both force fields are highly similar. However, different parameters for the bonded terms result in different modeled β-CD conformations, which significantly changes the H-bond networks between water molecules and β-CD. As illustrated in Figure 3 and Table SI 2, the conformation results in breaking more intermolecular H-bonding and larger desolvation penalty in GAFF-CD on guest binding.

Strong enough intermolecular attraction $\Delta H_{Solute\ Inter}$ is always required for molecular recognition, and vdW attraction was the major driving force for all the guests (Table 4). The loss of intramolecular H-bonds of β-CD on guest binding is compensated by the intermolecular Columbic attraction in the bound state ($\Delta H_{Host\ Conf\ (Coul)}$ in Table 6). Although the values of $\Delta H_{Host\text{-}Guest}$ with both force fields are similar, the decomposition shows significantly larger numbers of each contribution to $\Delta H_{Host\ Conf}$, such as $\Delta H_{Host\ Conf\ (vdW)}$ and $\Delta H_{Host\ Conf\ (Coul)}$ of β-CD, with GAFF-CD (Table 6) because of larger conformational changes after ligand binding. The free β-CD prefers flipping 2 glucopyranose units instead of holding an open cavity, as shown in the crystal structure (Figure 4). The glucose rings flipped outward during ligand binding, which lost more water molecules to allow the binding site of β-CD to be accessible for guests to bind. In contrast, q4MD-CD appeared to have crystal-like host structures in both the free and bound states. Note that in vacuum, both GAFF-CD and q4MD-CD sampled predominantly crystal-like host structures

(Figure SI 1), which indicates that the glucose ring flipping is largely induced by the hydration shell. GAFF-CD not only changed host conformations upon ligand binding but also makes the host more flexible.

Changes of Solute Entropy on Ligand Binding. Solute entropy, also termed configuration entropy, reflects the flexibility of a molecular system. Here we used numerical integration to compute each entropy term, by eq. 1 for internal (conformational/vibrational) and eq. 2 for external (translational/rotational) solute entropy. We used the well-defined dihedral distribution analyzed from our MD trajectories to compute internal solute entropy, as detailed in SI Section 5. For the small and weak binding guests studied here, the internal solute entropy of a guest is nearly identical when the guest is in the free solution or bound to β-CD, suggesting a weak correlation between internal degrees of freedom and external translational and rotational degrees of freedom (Table 5 and SI Movie). The external and internal entropy terms therefore can be computed separately in our systems. The calculated entropy values are shown in Table 5. A system is well known to lose configuration entropy because the intermolecular attractions inevitably rigidify the 2 molecules on binding [39-40, 62-63]. For example, a drug binding to HIV protease can result in >10 kcal/mol entropic penalty from rigidifying conformational and vibrational degrees of freedom [64]. Additional loss of the configuration entropy also comes from translational and rotational entropy of a ligand, > 7 kcal/mol, by confining itself in a snug binding site [55-56]. Post-analyzing our MD trajectories showed that all guests and β-CD were not markedly rigidified in the bound state. The guests lost ~ 1.5–2 kcal/mol external entropy, and β-CD was slightly more flexible, gaining 0.5–1.8 kcal/mol internal entropy with GAFF-CD or being unchanged with q4MD-CD on guest binding (Table 5). Even though different force fields for the host yield considerable differences in β-CD fluctuations, the variation in internal entropy of guests (-T$\Delta S_{Guest\ Int}$) is negligible. Also, the interactions between β-CD and guests are not large enough to strongly confine a guest to a handful of well-defined bound guest conformations, thereby allowing a guest to freely tumble and diffuse in the cavity of β-CD. Figure SI 11 illustrates that ligands can pose a wide collection of locations and orientations covering the center and top regions of the binding site, thereby resulting in a small reduction of -T$\Delta S_{Guest\ Ext}$.

Changes of Water Entropy on Ligand Binding. Water entropy is one major driving force in ligand binding in these systems, contributing to -2 to -4 kcal/mol to the free energy of binding (Tables 1 and 5). Gaining water entropy dominates in the binding of all guests to β-CD with GAFF-CD and the first 3 weaker binders to β-CD with q4MD-CD. The change in water entropy (-TΔS_{water}) is a combined effect

from rearranging water molecules, which affects their vibrational and conformational entropy, and from releasing the water molecules residing in the cavity of β-CD or interacting with the guest after the complex is formed (Tables 8 and SI 3 to 6). We used a grid cell theory approach to calculate the water entropy over the space near the solute, as compared with the entropy estimated for the bulk solvent. We also used an alternative method that computes the molar entropy of water when different solutes are present to approximate -TΔS$_{water}$ (detailed in SI Section 6). Both methods agree with each other. The theory and implementation were first validated by comparing our computed molar entropy of bulk water at 298K, 5.26 kcal/mol (Tables SI 3 to 5), with standard molar entropy of water at 298K, 4.98kcal/mol. The rigid water TIP3P model performed very well in reproducing standard molar entropy, as reported here and previously, with a two-phase thermodynamic method [65-66].

Table 5 lists the changes in water entropy on each guest binding to β-CD, and the contribution from each decomposed entropy term is in Tables SI 3 to 5. Using different force fields to model β-CD did not change the computed -TΔS$_{Water}$, so solute flexibility does not play an important role in water entropy calculations. Aspirin and methyl butyrate have more water entropy gains on binding than do other guests. The 2 guests have more polar functional groups to capture nearby water molecules in their free state, and after forming the complex with β-CD, these water molecules are released, which results in gaining more entropy on binding.

Figure 5 reveals the fluctuation of water molecules presented in the grid space in a pure water box and when β-CD, 2-naphthyl ethanol, or the complex is present using GAFF-CD. The behavior of water molar entropy in q4MD-CD is similar to GAFF-CD, so we only one force field is presented in this figure. The averaged jiggling of water molecules is quantified by computed translational, rotational and conformational entropy, as shown in columns 2-4 of Figure 5, and the darker grid indicates that water molecules in this cell are less mobile. We used an in-house multi-layer visualization program to display the decomposed entropy terms for each grid, and the layer through the center of the solute was selected in the figure. In general, the translational entropy decreases at the surface of the solute because the existence of the solute hinders free diffusion of water molecules. In free β-CD and 2-naphthyl ethanol, the translational entropy of water decreases ~ 0.2–0.4 kcal/mol around the solutes, as shown by darker cells in the vicinity of each solute in Figure 5B, 5C and 5D.

The rotational entropy increases on the hydrophobic surface and decreases near hydrophilic regions. For example, with free 2-naphthyl ethanol, the rotational entropy increases ~ 0.2–0.3 kcal/mol near the carbon chains or benzene ring and in the hydrophobic cavity of free and bound β-CD (see lighter

cells in Figures 5B and 5C). The non-polar surface may promote water tumbling, but forming H-bonds with a polar group decreases rotational entropy by ~ 0.2–0.5 kcal/mol near the hydroxyl group of 2-naphthyl ethanol. The conformational entropy is correlated with the density of water molecules (eq. 6.6 in SI). The cavity of β-CD is large enough to hold ~ 5–6 water molecules, on average, and the waters slightly decrease their conformational entropy by 0.2-0.3 kcal/mol because the cavity restrains the rearrangement of water molecules. However, the water is not frozen in the cavity and it keeps the fluid-like property, for no significant decrease in translational water entropy. Because the density of water fluctuates during the MD simulations, except for water molecules in the vicinity of solutes, conformational entropy of waters in the bulk solvent or surrounding a solute does not noticeably differ.

Binding Kinetics: Calculations of Association and Dissociation Rate Constants

The fast kinetics of guest binding to β-CD allowed for directly assessing the association (k_{on}) and dissociation (k_{off}) rate constants from the bound and unbound lengths during microsecond-long unguided MD simulations. Table 2 summarizes the calculated k_{on} and k_{off} for systems modeled by GAFF-CD and q4MD-CD. The diffusion-controlled limit for the molecular systems is approximated by $k_{on\_diffuse} = 4\pi DR$, where D is the diffusion coefficient obtained by our MD simulations (4×10^{-9} m^2/s for a guest such as aspirin and 7×10^{-10} m^2/s for β-CD, regardless of force field used) and R is the sum of the radius of β-CD and a guest (~ 10Å). The estimated diffusion-controlled association rate constants for all systems are $k_{on\_diffuse}$ ~ 3–4$\times10^{10}$ 1/Ms by using the approximated size of each molecule and the diffusion coefficient obtained by MD (Table SI 8). The modeled k_{on} by using GAFF-CD agrees very well with experimental data, and all guests showed 2 orders of magnitude slower k_{on} than $k_{on\_diffuse}$. Using q4MD-CD slightly overestimated k_{on} for all guest binding, and the value is one order of magnitude slower than $k_{on\_diffuse}$ because of the spatial factor. Because β-CD does not require considerable conformational changes or slow transition to acquire all the guests, experiments revealed no differences in k_{on} for different guests. However, k_{on} modeled with GAFF-CD shows that larger guests such as aspirin and naphthyl ethanols associate marginally faster to β-CD, with k_{on} values close to 10^9 1/Ms, as compared with smaller guests such as butanols, with k_{on} ~ 2–3$\times10^8$ 1/Ms. The faster association from modeling may result from stronger intermolecular attraction that can more easily retain the guest once 2 molecules collide. In contrast, because k_{on} modeled with q4MD-CD is already fast and close to $k_{on\_diffuse}$, a small difference is observed, with k_{on} ~ 1–4$\times10^9$ 1/Ms.

The difference in computed k_{on} with the two force fields and from experimental k_{on} result from the intermolecular attractions and the desolvation process. Guest diffusing on the cyclodextrin surface is unlikely due to weak intermolecular interactions, and the β-CD surface features a restricted target area. Assuming no intermolecular attraction, successful binding occurs only with the first collision occurring in the window areas of the cavity (~ 2.5% of the entire surface), which are the areas covering the top and bottom entrance of the cavity. With q4MD-CD, modeled k_{on} is 20 times slower than $k_{on\_diffuse}$, and ~ 5% of molecular encounters result in successful binding. Therefore, the restricted target area is the main contribution to a slower k_{on}. Using the same concept, less than 1% of the initial association results in a stable complex modeled with GAFF-CD, except for aspirin and 2-naphthyl ethanol. The chief reason that causes further slowdown in k_{on} with GAFF-CD is the desolvation process during association. Although the tilt glucose rings in the free β-CD may partly occlude the cavity, rotating the 2 dihedrals in C-O-C for different glucose ring tilting is nearly barrier-less. Replacing water molecules that formed the H-bond network with the free β-CD creates an energy barrier and results in unsuccessful binding, even for a guest already diffused to the cavity. Note, successful binding was considered only when a complex larger than 1 ns was formed in MD simulations.

In contrast to k_{on}, k_{off} modeled by q4MD-CD agrees very well with experiments; however, with GAFF-CD, all guests left β-CD approximately one order faster than the measured k_{off} values. Because of the faster k_{off}, the equilibrium constants (K_{eq}) are systematically smaller than the experimental values, except for 1-naphthyl ethanol. The dissociation rate constants are directly proportional to how long a guest can stay in the pocket of β-CD, also termed residence time in the drug discovery community. Different force field parameters can largely affect k_{off}. The longer average bound time indicates more negative ΔH (Table 3), in which ΔH computed with q4MD-CD is stronger than that with GAFF-CD. However, longer bound time does not always require stronger intermolecular attractions, and Table 3 shows that the water effects can be the major differentiating factors. Of note, we sampled hundreds of bound/free states during long simulations (Tables SI 10 and 11) but these are still less than the real experiments.

For the weak binders, we observed one direct association/dissociation pathway in which a guest diffused into the window of the cavity and then stayed with β-CD. The association perturbed the conformations of β-CD to get rid of hydrated waters and flip glucopyranose. We term this pathway the direct binding pathway (SI Movies 1,2,7,8). For the strong binders such as aspirin, for which k_{on} is 3- to 10-fold faster than weak binders modeled by both force fields, we observed one more association/dissociation pathway, termed the sticky binding pathway (SI Movies 3-6, 9-12). The stronger

intermolecular attractions allow the guest to stay on the surface of β-CD for surface diffusion to reach the cavity. This situation largely increases the possibility of binding events because the guests can overcome the limitation of a restricted target area of the surface. Note that unlike some ligand–protein binding in which the large biomolecular system needs longer than a microsecond timescale for both molecules to arrange to form a complex, binding processes of guest–β-CD are very fast, in the sub-nanosecond range, without large binding energy barriers. Nevertheless, the intermolecular attractions, possible surface diffusion and desolvation process still play a key role in controlling binding kinetics.

4. Discussion

This study demonstrates that unbiased MD simulations can be used to compute CD–guest binding kinetics and thermodynamics. The kinetic properties and binding enthalpies can be extracted directly from the simulations, but post-processing of the MD trajectories by using numerical integration and the grid cell method is needed to compute solute and solvent entropy of binding. Computed values with both force fields, GAFF-CD and q4MD-CD, agree well with experimental measures; however, values strikingly differ depending on whether the binding is driving by enthalpy or entropy.

Binding free energy computed from the kinetic data (ΔG_{Comp2}) correlate better with experiment (Figure 2), compared with ΔG_{Comp1} computed from ΔH-$T\Delta S$. The calculation only needs one long MD run that includes both solutes in one water box. Eq. 6 may be rewritten as ΔG_{Comp2} = -RTln($\frac{1}{\text{average unbound time} \times [\text{solute}]} \cdot C° / \frac{1}{\text{average bound time}}$). The numbers of bound and unbound events for computing the average time are nearly identical from the same one MD run; thus, another equation to describe binding free energy may be ΔG_{Comp3} = -RTln($\frac{\text{total bound time}}{\text{total unbound time} \times [\text{solute}]} \cdot C°$). From classical statistical mechanics, the probability of observing a bound complex during this long MD run is $P_{bound} = \frac{\int_{bound} e^{-E_{bound}(r)/RT} dr}{\int_{all} e^{-E(r)/RT} dr}$. As a result, $\Delta G = G_{bound} - G_{unbound}$ = -RTln($\frac{P_{bound}}{P_{unbound}}$) which is the same as equation ΔG_{Comp3}. The enthalpy change is also available from the average energy of the bound and unbound states in the trajectories of the complexes, $\Delta H_{Bound-unbound} = <H_{bound}> - <H_{unbound}>$. The computed enthalpy changes are highly similar to the values computed using four separated MD runs (Figure SI 12 and Table 1). This study aims to investigate the kinetic properties in addition to binding affinities; thus, ΔG_{Comp3} was not used to obtain the results. However, for other weak binding systems that k_{on} and k_{off} measurements are not available, and for the research that focuses on binding free energies, we suggest that one can run a

long MD simulation to obtain ΔG and ΔH. It is worth mentioning that although directly computing entropy changes is not needed if ΔG and ΔH are available, the calculation allows researchers to investigate the water effects in a great detail. The calculations also bring further insights into contributions regarding water entropy and solute configuration entropy to binding. However, because configuration entropy calculations inevitably need multi-dimension integral, a larger error than ΔG computed from k_{on} and k_{off} is anticipated.

Intuitively, a more negative binding enthalpy ΔH modeled by q4MD-CD is likely a result of the Lennard-Jones parameters, which could result in more optimized $\Delta H_{Host\text{-}Guest}$ [67]. Nevertheless, in the decomposed term $\Delta H_{Solute\ Inter}$ of $\Delta H_{Host\text{-}Guest}$, the computed $\Delta H_{Solute\ Inter\ (vdW)}$ and $\Delta H_{Solute\ Inter\ (Coul)}$ show that both force fields model highly similar inter-solute attractions, and the main determinant is from water. Both force field parameters allow the sugar ring to flip in the free states, and β-CD can easily adjust to an open cavity conformation when forming a complex with a guest. Therefore, unlike existing study showing that substituents attached to decorated β-CDs block a guest from binding [68], the ring flipping itself in our study did not hinder guest binding. However, more flipped sugar rings modeled by GAFF-CD allow the formation of more H-bonds between waters and β-CD and a more structured H-bond network as compared with conformations modeled by q4MD-CD. Therefore, the cavity more energetically accommodates stable water molecules, which results in large enthalpy penalty on desolvating those water molecules. The Coulombic term $\Delta H_{Host\text{-}Water\ (Coul)}$ becomes the essential energy term that opposes binding due to breakage of β-CD–water H-bonds, and the value is significantly larger with GAFF-CD. We suspected that the bonded parameters with GAFF-CD may highly prefer sugar ring flipping in the free states. However, the MD simulations in vacuum showed that both GAFF-CD and q4MD-CD highly prefer wide-open, crystal structure-like conformations in the free states (Figure SI 1). Because β-CD is reasonably flexible, in particular with GAFF-CD modeling, adding explicit water molecules easily induces the conformation changes. A "slaving" model in which water drives protein fluctuations was proposed [69]. Recently, a direct measurement of hydration water dynamics in protein systems illustrated that the surface hydration-shell fluctuation drives protein side-chain motions [70]. Here we showed that the water molecules are highly responsible for molecular recognition in both thermodynamics and kinetics.

Without considering other factors, the theoretical k_{on} may be estimated by multiplying the restricted target area by the diffusion-controlled limit $k_{on\_diffuse}$[71], 2.5% $\times 4 \times 10^{10}$ 1/Ms $= 10^9$ 1/Ms, which is close to the modeled k_{on} of most guests binding with q4MD-CD. To increase an association rate faster

than the theoretical value 10^9 1/Ms, when the initially collision did not bring a guest to the restricted target area, intermolecular attractions kept the guest in close proximity to β-CD until the guest reached the cavity. This situation can be observed during MD simulations, as illustrated in the sticky binding pathways with the tighter binder aspirin. Note no long-range electrostatic attractions for rate enhancement because of the neutral molecules. A guest always needs to compete with water molecules during binding. Therefore, k_{on} can be slower than the theoretical value 10^9 1/Ms if there exist barriers from removing stable water molecules in the hydration shell. The desolvation barrier increases when the free β-CD forms a larger number of H-bonds between water molecules modeled by GAFF-CD, which results in more unsuccessful guest binding and slower k_{on}. As a result, although a less wide-open cavity conformation in free β-CD may seem to directly lead to a slower k_{on} in GAFF-CD, the direct cause is from paying a higher cost to disturb its solvation shell. After forming a complex with a guest, β-CD gained a few kcal/mol, showing a more negative $\Delta H_{Host\ Conf}$ on binding. A similar finding from investigating a binding free energy barriers for a drug binding a protein showed that desolvation of the binding pocket contributes the most to the free energy cost, as opposed to reorganizing the protein binding pocket [72]. However, for molecular systems that encounter large-scale conformational changes and/or induce fit during ligand binding, rearranging conformations may still significantly affect the association rate constants [73], and the kinetic property can be highly system-dependent. With q4MD-CD, because water molecules form a less stable H-bond network, the role of desolvation in binding kinetics is not as important as with GAFF-CD.

Unlike desolvation effects, which are quite different from the 2 force fields, another dominant but similar driving force for binding from both force fields is the attractive component of the vdW energy, ranging from -6 to -23 kcal/mol. This driving force may be expected because of the nonpolar property of the β-CD cavity and the neutral guests. This term mainly accounts for dispersion forces between β-CD and guests in the force-field parameters and is similar for both force fields, with a trend that larger guests have more negative $\Delta H_{Solute\ Inter\ (vdW)}$. In experiments, measuring the separate contributions for binding from dispersive interactions and classical hydrophobic effects in aqueous environment is challenging, and the absolute values from the dispersive interactions are not available experimentally [74-75]. As compared with the vdW attraction, the Coulombic attraction between β-CD and guests is significantly weaker because the guests are not highly charged molecules and few intermolecular H-bonds are formed in the complex. The intermolecular attractions are balanced by the enthalpy penalty from disrupting attractions between waters and solutes and gaining water-water enthalpy for those water molecules replaced by a guest, which results in merely a few kcal/mol net enthalpy changes on binding. One may consider

hydrophobic effects as the major contributions to β-CD and guest recognition [76]. Of note, although the pocket of β-CD is non-polar, it is a very tiny cavity, and the rims of β-CD consist of several hydroxyl groups. On binding, ~ 20–25 water molecules were replaced by a larger guest, which agrees with experimental measurement [77] (Table SI 6). However, not all replaced water molecules are "unhappy". Therefore, although the replaced water molecules also regain water-water attractions in the bulk solvent, the costs to replace the stable water molecules result in the large desolvation penalty.

The systems did not encounter large solute entropy loss, which contrasts with several existing publications that suggested loss of configuration entropy when a drug binds its target protein [63-64, 78-80]. Unlike most drug-like compounds, which fit tightly to their target protein pocket, our guests only loosely fit in the cavity of β-CD. Therefore, the mobility of β-CD is not reduced considerably by a guest. With GAFF-CD, the hydrated water molecules in the cavity of free β-CD showed an ordered H-bond network and slaved the conformational motions of β-CD. On ligand binding, a bound guest did not form a stable H-bond network with β-CD; thus, β-CD showed a slightly increased flexibility. The guests were also able to form various contacts with β-CD. Similar to alternative contacts provided by the hydrophobic binding pocket of protein systems [81], we did not observe rigidity of β-CD with GAFF-CD.

The enthalpy and entropy balance may follow immediately from eq. 5, $\Delta G_{comp1}=\Delta H-T\Delta S$ [62, 82-83]. Therefore, it has been suggested that the entropy-enthalpy compensation is from a much smaller range of experimentally measured ΔG for a series of ligands than the range of ΔH. Different from most experimental techniques, we computed the entropy and enthalpy terms separately, and still observed the entropy-enthalpy compensation. Our guests were all weak binders and did not have a wide spectrum of ΔG. The computed range of ΔH is in a similar ballpark as ΔG, and the range of $-T\Delta S$ is relatively smaller than ΔG and ΔH. As a result, the enthalpy change mostly governs if a guest is a stronger or weaker binder. Our calculations reveal the physical basis of larger range of ΔH and more similar $-T\Delta S$. The enthalpy calculations are based on energy functions in the force fields, but the Gibb's entropy formula is based on the distribution of the microstates (eq. 1). Unlike protein systems with numerous rotatable bonds and a larger binding site to mostly enclose a ligand, a guest is not completely confined within the cavity of β-CD, and the host remains highly flexible. Interestingly, $-T\Delta S_{water}$ is similar in both force fields, and $-T\Delta S_{water}$ is not simply proportional to the size of a guest. Instead, $-T\Delta S_{water}$ relates more to the hydrophilicity of a guest such as 1-propanol, methyl butyrate and aspirin when forming a complex with β-CD. The free guests reduce more entropy of water in their solvation shell, and these solvation waters gain more entropy on guest binding. For q4MD-CD, because the free β-CD generally has a more open

cavity, more waters were released on binding (Table SI 6). However, the ring flipping conformation modelled by GAFF-CD produces a more structured H-bond network for the first hydration shell. As a result, although fewer water molecules were released on binding, those waters gained more entropy than those of β-CD with q4MD-CD, which resulted in a similar computed -TΔS$_{water}$ from both force fields. The results suggest that as in enthalpy, entropy calculations feature a fine balance.

Force-field parameters are critical for accurate modeling and successful prediction [84-87]. In this study, we used GAFF for β-CD (GAFF-CD) and for all guests, and q4md-CD, a specialized force field for CDs that combines Amber99SB and GLYCAM04 to match experimental geometries from crystal structures and NMR [48]. It is common practice to seek agreement between the calculated and experimental binding affinities/binding free energies for validating and improving the parameters of force fields or solvent models. Using computed thermodynamics and kinetics (eqs. 5 and 6), both force fields for β-CD showed reasonable agreement with experimental binding affinities, which validated the parameters used. Interestingly, different force fields concluded different driving forces, with GAFF-CD and q4MD-CD showing an entropy- and enthalpy-driven binding, respectively. In addition, q4MD-CD yielded better agreement between computed and experimental association rate constants. Using only binding free energy in the training set for parameterization was suggested to risk an incorrect entropy-enthalpy balance; therefore, binding enthalpy needs to be considered for optimizing parameters [67]. With continuing growth in computer power, for molecular systems with fast association/dissociation rate constants, we suggest considering computed binding kinetics for validating and optimizing force-field parameters as well. Our studies also showed the importance of and challenge in correctly modeling multiple conformations in which solvent effects may be remarkable and experimental structures are not available. Although our preliminary studies indicated that using TIP3P and TIP4P water models did not yield different sampled conformations during MD simulations, other molecular systems may be more sensitive to the solvent effects with different water models. In the future, we envision a more careful force-field optimization that considers binding free energy, enthalpy-entropy balance and kinetic properties. We also anticipate further investigation into the role of water in the binding kinetics of various guests to a pocket with different polar and/or nonpolar properties [88-89].

5. Conclusion

In this work, we performed microsecond-timescale MD simulations with GPU acceleration for 7 chosen β-CD complexes. The computed thermodynamic and kinetic properties agree with experimental values

reasonably well. The binding of β-CD complexes is mainly driven by the non-polar attraction between β-CD and the guest and the entropy gain of desolvated water molecules, regardless of the force field used. With both force fields, the ligands have only a small entropy penalty, and the entropy term also contributes to the binding process. GAFF-CD reproduced more favorable binding entropy and less favorable binding enthalpy due to stronger desolvation penalty than did q4MD-CD. As compared with GAFF-CD, q4MD-CD produced more rigid dynamics of free β-CD. However, the conformational rearrangement did not contribute to the differences in thermodynamics and kinetics modeled by the 2 force fields, and the real determinant was the different H-bond networks between the solvation water molecules and free β-CD. With GAFF-CD, free β-CD forms more H-bonds with solvation waters in a more distorted conformation; on ligand binding, the ligand needs to pay more enthalpic penalties to remove these stable H-bonded waters, which results in the stronger desolvation enthalpy. Removing the H-bonded waters also slows the association process. In the bound state, free β-CD forms fewer H-bonds with its environment and thus becomes more flexible and gains configuration entropy. Although we computed ΔH and $-T\Delta S$ separately, the compensation is observed. Our study also showed that different force-field parameters can yield the same computed ΔG but different entropy and entropy balance.

Acknowledgements
We thank support from the US National Institute of Health (GM-109045), US National Science Foundation (MCB-1350401), and NSF national super computer centers (TG-CHE130009). We also thank Dr. Michael Gilson and Dr. Niel Henriksen for insightful discussions on β-cyclodextrin systems and Dr. Ron Levy for sharing OPLS force field parameters for β-cyclodextrin for our tests.

Figure 1. The structure of β-cyclodextrin (β-CD) and the 7 guest molecules. In the structure of β-CD, hydrogen atoms are not shown. For the β-CD internal (vibrational/conformational) entropy calculation, the 14 dihedral angles used to define the conformations of β-CD are in blue.

Figure 2. The correlations between ΔG_{Comp1}, ΔG_{Comp2} and experimental values. The results from GAFF-CD are shown in blue triangles, and results from q4MD-CD are shown in orange rectangles. The correlations are label correspondingly.

GAFF-CD   q4MD-CD

Figure 3. The hydrogen bond (H-bond) patterns of representative free β-CD conformations with GAFF-CD and q4MD-CD. With GAFF-CD, there are 18 water molecules forming H-bonds with β-CD, 24 H-bonds with water (blue dotted lines), and 1 intramolecular H-bond (orange dotted lines). With q4MD-CD, there are 11 water molecules forming H-bonds with β-CD, 16 H-bonds with water, and 5 intramolecular H-bonds.

Figure 4. RMSD plots and representative conformations of β-CD for free β-CD, β-CD-2-naphthyl ethanol complex and β-CD-t-butanol complex with GAFF-CD and q4MD-CD. RMSD (Å) are computed against the crystal structure by using conformations chosen every 100 ps from all conformations of free β-CD and bound-state conformations of complexes. Representative conformations are shown near the labels and circles on the plots. In the representative conformations, ligands are in green.

Figure 5. The water entropy decompositions of pure water and in the vicinities of free β-CD, free 2-napthyl ethanol and β-CD-2-napthyl ethanol complex in GAFF-CD. From top to bottom are pure water (A), free β-CD (B), free 2-napthyl ethanol (C), and β-CD-2–naphthyl ethanol complex (D). The plane of the spatial grid in the figures is shown in green in the first column. From left to right are total entropy (1, S_{Water}), translational entropy (2, $S_{Water\ Trans}$), rotational entropy (3, $S_{Water\ Rot}$) and conformational entropy (4, $S_{Water\ Conf}$). Each column has a separate color bar (values in kcal/mol). The red areas are regions occupied by the solute molecules, and water entropy cannot be calculated.

Calculated (GAFF-CD)			
Guest	ΔG_{Comp1}	ΔH	$-T\Delta S$
1-Propanol	-1.54±0.93	0.54±0.74	-2.08±0.19
1-Butanol	1.01±0.59	2.99±0.52	-1.98±0.06
Methyl Butyrate	-2.40±0.52	1.88±0.47	-4.29±0.05
t-Butanol	0.38±0.46	2.86±0.40	-2.48±0.06
1-Naphthyl Ethanol	-3.01±0.31	-1.41±0.28	-1.60±0.03
Aspirin	-3.84±0.35	-0.01±0.31	-3.83±0.04
2-Naphthyl Ethanol	-2.93±0.30	-0.88±0.27	-2.05±0.03
Calculated (q4MD-CD)			
Guest	ΔG_{Comp1}	ΔH	$-T\Delta S$
1-Propanol	-2.21±0.47	-0.66±0.41	-1.55±0.06
1-Butanol	-1.33±0.34	-0.85±0.30	-0.48±0.04
Methyl Butyrate	-4.17±0.38	-2.00±0.33	-2.17±0.05
t-Butanol	-2.50±0.38	-2.87±0.33	0.37±0.05
1-Naphthyl Ethanol	-5.53±0.32	-5.44±0.29	-0.09±0.04
Aspirin	-6.33±0.37	-4.30±0.33	-2.03±0.05
2-Naphthyl Ethanol	-4.32±0.31	-4.03±0.28	-0.29±0.03
Experimental			
Guest	ΔG	ΔH	$-T\Delta S$
1-Propanol [24]	-0.88±0.09	1.43±0.48	-2.34±0.57
1-Butanol [24]	-1.67±0.19	0.69	-2.34
Methyl Butyrate [61]	-1.99±0.02	/	/
t-Butanol [90]	-2.22±0.01	/	/
1-Naphthyl Ethanol [29]	-4.08±0.01	/	/
Aspirin [28]	-3.74±0.00	/	/
2-Naphthyl Ethanol [29]	-3.97±0.07	/	/
			Units: kcal/mol

Table 1. The binding free energies (ΔG), enthalpies (ΔH) and entropies ($-T\Delta S$) of β-cyclodextrin (β-CD) with 7 guests from experiments and calculations by using GAFF-CD and q4MD-CD force fields. Unavailable experimental values are marked by /. All values are in kcal/mol.

Calculated (GAFF-CD)				
Guest	k_{on} (1/s·M)	k_{off} (1/s)	K_{eq} (1/M)	ΔG_{Comp2} (kcal/mol)
1-Propanol	$2.0\pm0.1\times10^8$	$3.9\pm0.3\times10^8$	0.5 ± 0.06	0.40 ± 0.08
1-Butanol	$2.2\pm0.1\times10^8$	$1.1\pm0.2\times10^8$	2.0 ± 0.3	-0.41 ± 0.10
Methyl Butyrate	$3.9\pm0.1\times10^8$	$8.0\pm0.6\times10^7$	4.9 ± 0.4	-0.94 ± 0.05
t-Butanol	$2.3\pm0.1\times10^8$	$8.5\pm1.2\times10^7$	3.2 ± 0.4	-0.69 ± 0.08
1-Naphthyl Ethanol	$8.6\pm1.1\times10^8$	$1.5\pm1.9\times10^6$	558.9 ± 100	-3.75 ± 0.12
Aspirin	$1.1\pm0.01\times10^9$	$2.4\pm0.3\times10^7$	46 ± 4.3	-2.27 ± 0.06
2-Naphthyl Ethanol	$9.5\pm0.4\times10^8$	$5.2\pm0.9\times10^6$	182 ± 23	-3.08 ± 0.08
Calculated (q4MD-CD)				
Guest	k_{on} (1/s·M)	k_{off} (1/s)	K_{eq} (1/M)	ΔG_{Comp2} (kcal/mol)
1-Propanol	$1.2\pm0.02\times10^9$	$1.2\pm0.02\times10^8$	10.1 ± 0.2	-1.37 ± 0.01
1-Butanol	$1.5\pm0.03\times10^9$	$3.3\pm0.07\times10^7$	45.9 ± 1.2	-2.27 ± 0.02
Methyl Butyrate	$1.6\pm0.1\times10^9$	$1.1\pm0.1\times10^7$	146.8 ± 15	-2.95 ± 0.06
t-Butanol	$1.1\pm0.07\times10^9$	$7.2\pm1.0\times10^6$	157.5 ± 18	-3.00 ± 0.07
1-Naphthyl Ethanol	$1.2\pm0.5\times10^9$	$1.4\pm0.5\times10^6$	876.0 ± 42	-4.01 ± 0.03
Aspirin	$3.2\pm0.3\times10^9$	$3.1\pm0.9\times10^6$	1035.3 ± 83	-4.11 ± 0.05
2-Naphthyl Ethanol	3.91×10^9	5.0×10^5	7788.3	-5.31
Experimental				
Guest	k_{on} (1/s·M)	k_{off} (1/s)	K_{eq} (1/M)	ΔG (kcal/mol)
1-Propanol [90]	$5.1\pm0.7\times10^8$	$1.2\pm0.07\times10^8$	4.2 ± 0.6	-0.88 ± 0.09
1-Butanol [90]	$2.8\pm0.8\times10^8$	$3.8\pm0.6\times10^7$	7.2 ± 2.0	-1.67 ± 0.19
Methyl Butyrate [61]	$3.7\pm0.3\times10^8$	$1.28\pm0.03\times10^7$	29 ± 1	-1.99 ± 0.02
t-Butanol [90]	$3.6\pm0.1\times10^8$	$0.85\pm0.01\times10^7$	42.6 ± 1.0	-2.22 ± 0.01
1-Naphthyl Ethanol [29]	$4.7\pm1.9\times10^8$	$4.8\pm1.8\times10^5$	979 ± 10	-4.08 ± 0.01
Aspirin [28]	$7.2\pm0.04\times10^8$	$1.3\pm0.03\times10^6$	549 ± 2	-3.74 ± 0.00
2-Naphthyl Ethanol [29]	$2.9\pm1.6\times10^8$	$1.8\pm0.7\times10^5$	820 ± 90	-3.97 ± 0.07

Table 2. The association and dissociation rate constants of the 7 β-CD complexes from experiments and calculations by using GAFF-CD and q4MD-CD force fields. The standard deviations of rate constants of β-CD-2-naphthyl ethanol with q4MD-CD are not available because of lack of adequate binding events.

Calculated (GAFF-CD)								
Guest	ΔH	$\Delta H_{\text{Host-Guest}}$	$\Delta H_{\text{Host Conf}}$	$\Delta H_{\text{Guest Conf}}$	$\Delta H_{\text{Solute Inter}}$	$\Delta H_{\text{Desolvation}}$	$\Delta H_{\text{Host-Water}}$	$\Delta H_{\text{Guest-Water}}$	$\Delta H_{\text{Water-Water}}$
1-Propanol	0.54	-10.64	-0.87	-0.01	-9.76	11.18	12.20	6.92	-7.93
1-Butanol	2.99	-15.28	-2.25	0.00	-13.03	18.27	20.55	8.34	-10.62
Methyl Butyrate	1.88	-19.31	-2.85	0.02	-16.48	21.19	25.58	10.28	-14.67
t-Butanol	2.86	-15.71	-2.55	-0.04	-13.13	18.57	20.71	8.39	-10.53
1-Naphthyl Ethanol	-1.41	-30.60	-4.45	0.03	-26.18	29.19	35.45	14.93	-21.19
Aspirin	-0.01	-31.53	-3.77	0.23	-27.99	31.52	34.60	19.15	-22.22
2-Naphthyl Ethanol	-0.88	-28.45	-3.97	0.13	-24.61	27.57	33.21	13.76	-19.40
Calculated (q4MD-CD)								
Guest	ΔH	$\Delta H_{\text{Host-Guest}}$	$\Delta H_{\text{Host Conf}}$	$\Delta H_{\text{Guest Conf}}$	$\Delta H_{\text{Solute Inter}}$	$\Delta H_{\text{Desolvation}}$	$\Delta H_{\text{Host-Water}}$	$\Delta H_{\text{Guest-Water}}$	$\Delta H_{\text{Water-Water}}$
1-Propanol	-0.66	-13.61	-2.05	0.04	-11.60	12.95	13.72	7.74	-8.50
1-Butanol	-0.85	-16.66	-2.66	0.09	-14.10	15.82	17.72	9.21	-11.10
Methyl Butyrate	-2.00	-20.61	-3.53	0.18	-17.26	18.61	22.05	11.10	-14.55
t-Butanol	-2.87	-17.38	-3.50	-0.06	-13.82	14.51	16.31	7.87	-9.67
1-Naphthyl Ethanol	-5.44	-28.56	-3.42	0.20	-25.34	23.12	29.65	14.51	-21.04
Aspirin	-4.30	-29.85	-3.39	0.57	-27.03	25.55	28.91	18.17	-21.53
2-Naphthyl Ethanol	-4.03	-27.41	-3.38	0.16	-24.20	23.38	28.82	13.95	-19.39
									Units: kcal/mol

Table 3. The binding enthalpy decompositions of β-CD complexes by using GAFF-CD and q4MD-CD force fields. The total binding enthalpy (ΔH) is decomposed into solute-solute interaction change ($\Delta H_{\text{Host-Guest}}$) and desolvation energy ($\Delta H_{\text{Desolvation}}$). $\Delta H_{\text{Host-Guest}}$ includes conformational enthalpy changes of host ($\Delta H_{\text{Host Conf}}$) and guest ($\Delta H_{\text{Guest Conf}}$), and host–guest interaction change ($\Delta H_{\text{Solute Inter}}$). $\Delta H_{\text{Desolvation}}$ decomposes into host–water ($\Delta H_{\text{Host-Water}}$), guest–water ($\Delta H_{\text{Guest-Water}}$) and water–water ($\Delta H_{\text{Water-Water}}$) interaction changes. All values are in kcal/mol.

Calculated (GAFF-CD)			
Guest	$\Delta H_{\text{Solute Inter}}$	$\Delta H_{\text{Solute Inter (vdW)}}$	$\Delta H_{\text{Solute Inter (Coul)}}$
1-Propanol	-9.76	-6.08	-3.68
1-Butanol	-13.03	-10.33	-2.7
Methyl Ethanol	-16.48	-14.33	-2.16
t-Butanol	-13.13	-10.8	-2.32
1-Naphthyl Ethanol	-26.18	-23.57	-2.61
Aspirin	-27.99	-22.76	-5.22
2-Naphthyl Ethanol	-24.61	-21.72	-2.89
Calculated (q4MD-CD)			
Guest	$\Delta H_{\text{Solute Inter}}$	$\Delta H_{\text{Solute Inter (vdW)}}$	$\Delta H_{\text{Solute Inter (Coul)}}$
1-Propanol	-11.6	-9.67	-1.93
1-Butanol	-14.1	-12.09	-2.01
Methyl Ethanol	-17.26	-15.12	-2.14
t-Butanol	-13.82	-12.39	-1.43
1-Naphthyl Ethanol	-25.34	-22.76	-2.57
Aspirin	-27.03	-22.15	-4.88
2-Naphthyl Ethanol	-24.2	-21.45	-2.75
			Units: kcal/mol

Table 4. Decompositions of host-guest interaction change ($\Delta H_{\text{Solute Inter}}$) into van der Waals and Coulombic energies by using GAFF-CD and q4MD-CD force fields. $\Delta H_{\text{Solute Inter}}$ is decomposed into van der Waals energy ($\Delta H_{\text{Solute Inter (vdW)}}$), and Coulombic energy ($\Delta H_{\text{Solute Inter (Coul)}}$). All values are in kcal/mol.

Calculated (GAFF-CD)					
Guest	$-T\Delta S_{Water}$	$-T\Delta S_{Host}$	$-T\Delta S_{Guest\ Int}$	$-T\Delta S_{Guest\ Ext}$	$-T\Delta S$
1-Propanol	-3.32±0.18	-0.52±0.01	0.00±0.00	1.75±0.00	-2.08±0.19
1-Butanol	-2.37±0.06	-1.20±0.00	-0.05±0.00	1.64±0.00	-1.98±0.06
Methyl Butyrate	-4.00±0.04	-1.58±0.00	-0.10±0.00	1.39±0.00	-4.29±0.05
t-Butanol	-2.22±0.05	-1.84±0.00	0.00±0.00	1.57±0.00	-2.48±0.06
1-Naphthyl Ethanol	-2.28±0.02	-1.24±0.00	0.00±0.00	1.92±0.00	-1.60±0.03
Aspirin	-4.08±0.03	-1.53±0.00	0.12±0.00	1.67±0.00	-3.83±0.04
2-Naphthyl Ethanol	-2.33±0.02	-1.33±0.00	0.00±0.00	1.61±0.01	-2.05±0.03
Calculated (q4MD-CD)					
Guest	$-T\Delta S_{Water}$	$-T\Delta S_{Host}$	$-T\Delta S_{Guest\ Int}$	$-T\Delta S_{Guest\ Ext}$	$-T\Delta S$
1-Propanol	-3.36±0.05	0.21±0.00	0.02±0.00	1.57±0.00	-1.55±0.06
1-Butanol	-2.31±0.03	0.25±0.00	-0.17±0.00	1.74±0.00	-0.48±0.04
Methyl Butyrate	-4.12±0.04	0.22±0.00	-0.11±0.00	1.84±0.00	-2.17±0.05
t-Butanol	-1.84±0.05	0.45±0.00	0.00±0.00	1.76±0.00	0.37±0.05
1-Naphthyl Ethanol	-2.17±0.03	0.12±0.00	0.00±0.00	1.95±0.01	-0.09±0.04
Aspirin	-3.81±0.04	0.11±0.00	-0.16±0.00	1.82±0.00	-2.03±0.05
2-Naphthyl Ethanol	-2.31±0.03	0.03±0.00	0.00±0.00	1.99±0.01	-0.29±0.03
					Units: kcal/mol

Table 5. The binding entropy decomposition of β-CD complexes by using GAFF-CD and q4MD-CD force fields. $-T\Delta S_{Water}$, $-T\Delta S_{Host}$, $-T\Delta S_{Guest\ Int}$, $-T\Delta S_{Guest\ Ext}$ and $-T\Delta S$ are the entropy change of water, internal entropy change of β-cyclodextrin, internal and external entropy change of guests, and the total binding entropy at 298K. All values are in kcal/mol.

Calculated (GAFF-CD)				
Guest	$\Delta H_{Host\ Conf}$	$\Delta H_{Host\ Conf\ (Bonded)}$	$\Delta H_{Host\ Conf\ (vdW)}$	$\Delta H_{Host\ Conf\ (Coul)}$
1-Propanol	-0.87	1.37	1.76	-4.00
1-Butanol	-2.25	3.97	5.22	-11.44
Methyl Ethanol	-2.85	4.65	6.01	-13.51
t-Butanol	-2.55	5.05	6.07	-13.67
1-Naphthyl Ethanol	-4.45	6.23	8.71	-19.39
Aspirin	-3.77	5.89	7.95	-17.62
2-Naphthyl Ethanol	-3.97	5.54	8.15	-17.66
Calculated (q4MD-CD)				
Guest	$\Delta H_{Host\ Conf}$	$\Delta H_{Host\ Conf\ (Bonded)}$	$\Delta H_{Host\ Conf\ (vdW)}$	$\Delta H_{Host\ Conf\ (Coul)}$
1-Propanol	-2.05	-0.38	1.79	-3.46
1-Butanol	-2.66	-0.38	1.95	-4.23
Methyl Ethanol	-3.53	-0.47	2.29	-5.34
t-Butanol	-3.50	-0.83	2.18	-4.86
1-Naphthyl Ethanol	-3.42	0.62	2.57	-6.62
Aspirin	-3.39	0.34	2.44	-6.16
2-Naphthyl Ethanol	-3.38	0.21	2.57	-6.16
				Units: kcal/mol

Table 6. Decompositions of β-CD conformational enthalpy change ($\Delta H_{Host\ Conf}$) into bonded, van der Waals and Coulombic energies by using GAFF-CD and q4MD-CD force fields. $\Delta H_{Host\ Conf}$ is decomposed into bonded energy ($\Delta H_{Host\ Conf\ (Bonded)}$), van der Waals energy ($\Delta H_{Host\ Conf\ (vdW)}$), and Coulombic energy ($\Delta H_{Host\ Conf\ (Coul)}$). All values are in kcal/mol.

Calculated (GAFF-CD)						
Guest	$\Delta H_{Host\text{-}Water}$	$\Delta H_{Host\text{-}Water}$ (vdW)	$\Delta H_{Host\text{-}Water}$ (Coul)	$\Delta H_{Guest\text{-}Water}$	$\Delta H_{Guest\text{-}Water}$ (vdW)	$\Delta H_{Guest\text{-}Water}$ (Coul)
1-Propanol	12.20	1.34	10.85	6.92	2.12	4.80
1-Butanol	20.55	0.12	20.44	8.34	4.15	4.19
Methyl Ethanol	25.58	0.64	24.94	10.28	5.79	4.49
t-Butanol	20.71	-1.38	22.09	8.39	4.16	4.23
1-Naphthyl Ethanol	35.45	0.91	34.54	14.93	9.62	5.32
Aspirin	34.60	0.40	34.20	19.15	9.13	10.02
2-Naphthyl Ethanol	33.21	1.54	31.66	13.76	9.29	4.47
Calculated (q4MD-CD)						
Guest	$\Delta H_{Host\text{-}Water}$	$\Delta H_{Host\text{-}Water}$ (vdW)	$\Delta H_{Host\text{-}Water}$ (Coul)	$\Delta H_{Guest\text{-}Water}$	$\Delta H_{Guest\text{-}Water}$ (vdW)	$\Delta H_{Guest\text{-}Water}$ (Coul)
1-Propanol	13.72	4.08	9.63	7.74	3.85	3.89
1-Butanol	17.72	5.45	12.27	9.21	4.99	4.22
Methyl Ethanol	22.05	6.51	15.54	11.10	6.34	4.77
t-Butanol	16.31	4.75	11.56	7.87	5.12	2.75
1-Naphthyl Ethanol	29.65	8.63	21.02	14.51	9.60	4.91
Aspirin	28.91	7.75	21.16	18.17	9.35	8.82
2-Naphthyl Ethanol	28.82	8.95	19.87	13.95	9.37	4.59

Units: kcal/mol

Table 7. Decompositions of host-water ($\Delta H_{Host\text{-}Water}$) and guest-water ($\Delta H_{Guest\text{-}Water}$) interaction enthalpy changes into van der Waals and Coulombic energies by using GAFF-CD and q4MD-CD force fields. $\Delta H_{Host\text{-}Water}$ and $\Delta H_{Guest\text{-}Water}$ decompose into van der Waals energy ($\Delta H_{Host\text{-}Water\,(vdW)}$ and $\Delta H_{Guest\text{-}Water\,(vdW)}$) and Coulombic energy ($\Delta H_{Host\text{-}Water\,(Coul)}$ and $\Delta H_{Guest\text{-}Water\,(Coul)}$) terms. All values are in kcal/mol.

Calculated (GAFF-CD)				
Guest	$-T\Delta S_{\text{Water Trans}}$	$-T\Delta S_{\text{Water Rot}}$	$-T\Delta S_{\text{Water Vib}}$	$-T\Delta S_{\text{Water Conf}}$	$-T\Delta S_{\text{Water}}$
1-Propanol	-2.02	-0.91	-2.92	-0.39	-3.32
1-Butanol	-1.18	-0.73	-1.90	-0.47	-2.37
Methyl Butyrate	-1.92	-1.36	-3.28	-0.72	-4.00
t-Butanol	-1.16	-0.46	-1.63	-0.59	-2.22
1-Naphthyl Ethanol	-1.12	-0.18	-1.30	-0.98	-2.28
Aspirin	-2.36	-0.91	-3.26	-0.82	-4.08
2-Naphthyl Ethanol	-1.15	-0.24	-1.38	-0.95	-2.33
Calculated (q4MD-CD)				
Guest	$-T\Delta S_{\text{Water Trans}}$	$-T\Delta S_{\text{Water Rot}}$	$-T\Delta S_{\text{Water Vib}}$	$-T\Delta S_{\text{Water Conf}}$	$-T\Delta S_{\text{Water}}$
1-Propanol	-1.93	-0.90	-2.83	-0.53	-3.36
1-Butanol	-1.13	-0.47	-1.60	-0.71	-2.31
Methyl Butyrate	-1.80	-1.46	-3.26	-0.86	-4.12
t-Butanol	-0.77	-0.33	-1.10	-0.74	-1.84
1-Naphthyl Ethanol	-0.70	-0.37	-1.07	-1.10	-2.17
Aspirin	-1.99	-0.86	-2.85	-0.96	-3.81
2-Naphthyl Ethanol	-0.80	-0.47	-1.26	-1.04	-2.31
					Units: kcal/mol

Table 8. The decompositions of water entropy change by using GAFF-CD and q4MD-CD force fields. The binding water entropy ($-T\Delta S_{\text{Water}}$) decomposes into translational ($-T\Delta S_{\text{Water Trans}}$), rotational ($-T\Delta S_{\text{Water Rot}}$), vibrational ($-T\Delta S_{\text{Water Vib}} = -T\Delta S_{\text{Water Trans}} + -T\Delta S_{\text{Water Rot}}$), and conformational ($-T\Delta S_{\text{Water Conf}}$) terms. All values are in kcal/mol.

References

(1)   Leavitt, S.; Freire, E., *Curr. Opin. Struct. Biol.* **2001,** *11* (5), 560-566.
(2)   Bouchemal, K.; Mazzaferro, S., *Drug Discov. Today* **2012,** *17* (11-12), 623-629.
(3)   Swinney, D. C., *Curr. Opin. Drug Discovery Dev.* **2009,** *12* (1), 31-39.
(4)   Schreiber, G.; Haran, G.; Zhou, H. X., *Chem. Rev.* **2009,** *109* (3), 839-860.
(5)   Zhang, R.; Monsma, F., *Curr. Opin. Drug Discovery Dev.* **2009,** *12* (4), 488-496.
(6)   Guo, D.; Mulder-Krieger, T.; Ijzerman, A. P.; Heitman, L. H., *Br. J. Pharmacol.* **2012,** *166* (6), 1846-1859.
(7)   Sykes, D. A.; Dowling, M. R.; Charlton, S. J., *Mol. Pharmacol.* **2009,** *76* (3), 543-551.
(8)   Tummino, P. J.; Copeland, R. A., *Biochemistry* **2008,** *47* (32), 8465-8465.
(9)   Copeland, R. A.; Pompliano, D. L.; Meek, T. D., *Nat. Rev. Drug Discov.* **2007,** *6* (3), 249-249.
(10)   Copeland, R. A., *Nat. Rev. Drug Discov.* **2016,** *15* (2), 87-95.
(11)   Schneider, E. V.; Boettcher, J.; Huber, R.; Maskos, K.; Neumann, L., *Proc. Natl. Acad. Sci. U.S.A.* **2013,** *110* (20), 8081-8086.
(12)   Huang, Y.-m. M.; Kang, M.; Chang, C.-e. A., *J. Mol. Recognit.* **2014,** *27* (9), 537-548.
(13)   Cusack, K. P.; Wang, Y.; Hoemann, M. Z.; Marjanovic, J.; Heym, R. G.; Vasudevan, A., *Bioorg. Med. Chem. Lett.* **2015,** *25* (10), 2019-2027.
(14)   Seow, V.; Lim, J.; Cotterell, A. J.; Yau, M. K.; Xu, W. J.; Lohman, R. J.; Kok, W. M.; Stoermer, M. J.; Sweet, M. J.; Reid, R. C.; Suen, J. Y.; Fairlie, D. P., *Sci. Rep.* **2016,** *6*.
(15)   Friedrichs, M. S.; Eastman, P.; Vaidyanathan, V.; Houston, M.; Legrand, S.; Beberg, A. L.; Ensign, D. L.; Bruns, C. M.; Pande, V. S., *J. Comput. Chem.* **2009,** *30* (6), 864-872.
(16)   Shaw, D. E.; Deneroff, M. M.; Dror, R. O.; Kuskin, J. S.; Larson, R. H.; Salmon, J. K.; Young, C.; Batson, B.; Bowers, K. J.; Chao, J. C.; Eastwood, M. P.; Gagliardo, J.; Grossman, J. P.; Ho, C. R.; Ierardi, D. J.; Kolossvary, I.; Klepeis, J. L.; Layman, T.; McLeavey, C.; Moraes, M. A.; Mueller, R.; Priest, E. C.; Shan, Y.; Spengler, J.; Theobald, M.; Towles, B.; Wang, S. C., *Commun. ACM* **2008,** *51* (7), 91-97.
(17)   Challa, R.; Ahuja, A.; Ali, J.; Khar, R. K., *AAPS PharmSciTech* **2005,** *6* (2).
(18)   Del Valle, E. M. M., *Process Biochem.* **2004,** *39* (9), 1033-1046.
(19)   Singh, M.; Sharma, R.; Banerjee, U. C., *Biotechnol. Adv.* **2002,** *20* (5-6), 341-359.
(20)   Davis, M. E.; Brewster, M. E., *Nat. Rev. Drug Discov.* **2004,** *3* (12), 1023-1035.
(21)   Marchetti, L.; Levine, M., *ACS Catal.* **2011,** *1* (9), 1090-1118.
(22)   Breslow, R.; Dong, S. D., *Chem. Rev.* **1998,** *98* (5), 1997-2011.
(23)   Lai, W. F., *Biomaterials* **2014,** *35* (1), 401-411.
(24)   Rekharsky, M. V.; Inoue, Y., *Chem. Rev.* **1998,** *98* (5), 1875-1917.
(25)   Izatt, R. M.; Pawlak, K.; Bradshaw, J. S.; Bruening, R. L., *Chem. Rev.* **1995,** *95* (7), 2529-2586.
(26)   Yim, C. T.; Zhu, X. X.; Brown, G. R., *J. Phys. Chem. B* **1999,** *103* (3), 597-602.
(27)   Nilsson, M.; Valente, A. J. M.; Olofsson, G.; Soderman, O.; Bonini, M., *J. Phys. Chem. B* **2008,** *112* (36), 11310-11316.
(28)   Fukahori, T.; Kondo, M.; Nishikawa, S., *J. Phys. Chem. B* **2006,** *110* (9), 4487-4491.
(29)   Barros, T. C.; Stefaniak, K.; Holzwarth, J. F.; Bohne, C., *J. Phys. Chem. A* **1998,** *102* (28), 5639-5651.
(30)   Schneider, H. J.; Hacket, F.; Rudiger, V.; Ikeda, H., *Chem. Rev.* **1998,** *98* (5), 1755-1785.
(31)   Jana, M.; Bandyopadhyay, S., *J. Phys. Chem. B* **2011,** *115* (19), 6347-6357.
(32)   Zhang, H.; Ge, C.; van der Spoel, D.; Feng, W.; Tan, T., *J. Phys. Chem. B* **2012,** *116* (12), 3880-3889.
(33)   Zoppi, A.; Quevedo, M. A.; Delrivo, A.; Longhi, M. R., *J. Pharm. Sci.* **2010,** *99* (7), 3166-3176.

(34)   Zhang, H.; Feng, W.; Li, C.; Tan, T., *J. Phys. Chem. B* **2010,** *114* (14), 4876-4883.

(35)   Yeguas, V.; Altarsha, M.; Monard, G.; Lopez, R.; Ruiz-Lopez, M. F., *J. Phys. Chem. A* **2011,** *115* (42), 11810-11817.

(36)   Henriksen, N. M.; Fenley, A. T.; Gilson, M. K., *J. Chem. Theory Comput.* **2015,** *11* (9), 4377-4394.

(37)   Zhang, H.; Yin, C.; Yan, H.; van der Spoel, D., *J. Chem. Inf. Model.* **2016,** (56), 2080−2092

(38)   Sellner, B.; Zifferer, G.; Kornherr, A.; Krois, D.; Brinker, U. H., *J. Phys. Chem. B* **2008,** *112* (3), 710-714.

(39)   Wickstrom, L.; He, P.; Gallicchio, E.; Levy, R. M., *J. Chem. Theory Comput.* **2013,** *9* (7), 3136-3150.

(40)   Chen, W.; Chang, C. E.; Gilson, M. K., *Biophys. J.* **2004,** *87* (5), 3035-3049.

(41)   Koehler, J. E. H.; Grczelschak-Mick, N., *Beilstein J. Org. Chem.* **2013,** *9*, 118-134.

(42)   Dinar, K.; Sahra, K.; Seridi, A.; Kadri, M., *Chem. Phys. Lett.* **2014,** *595*, 113-120.

(43)   Liu, P.; Zhang, D.; Zhan, J., *J. Phys. Chem. A* **2010,** *114* (50), 13122-13128.

(44)   Lambert, A.; Yeguas, V.; Monard, G.; Ruiz-Lopez, M. F., *Comp. Theor. Chem.* **2011,** *968* (1-3), 71-76.

(45)   Pan, A. C.; Xu, H.; Palpant, T.; Shaw, D. E., *J. Chem. Theory Comput.* **2017,** *13*, 3372-3377.

(46)   Pedretti, A.; Villa, L.; Vistoli, G., *J. Comput. Aided Mol. Des.* **2004,** *18* (3), 167-173.

(47)   Wang, J. M.; Wolf, R. M.; Caldwell, J. W.; Kollman, P. A.; Case, D. A., *J. Comput. Chem.* **2004,** *25* (9), 1157-1174.

(48)   Cezard, C.; Trivelli, X.; Aubry, F.; Djedaini-Pilard, F.; Dupradeau, F.-Y., *PCCP* **2011,** *13* (33), 15103-15121.

(49)   Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery Jr., J. A.; Peralta, J. E.; Ogliaro, F.; Bearpark, M. J.; Heyd, J.; Brothers, E. N.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A. P.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, N. J.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. *Gaussian 09*, Gaussian, Inc.: Wallingford, CT, USA, 2009.

(50)   Fenley, A. T.; Henriksen, N. M.; Muddana, H. S.; Gilson, M. K., *J. Chem. Theory Comput.* **2014,** *10* (9), 4069-4078.

(51)   Wang, J.; Wang, W.; Kollman, P. A.; Case, D. A., *J. Mol. Graph. Model.* **2006,** *25* (2), 247-260.

(52)   Doll, J. D.; Dion, D. R., *J. Chem. Phys.* **1976,** *65* (9), 3762-3766.

(53)   Adelman, S. A., *J. Chem. Phys.* **1979,** *71* (11), 4471-4486.

(54)   Humphrey, W.; Dalke, A.; Schulten, K., *J. Mol. Graph. Model.* **1996,** *14* (1), 33-38.

(55)   Chang, C. E.; Gilson, M. K., *J. Am. Chem. Soc.* **2004,** *126* (40), 13156-13164.

(56)   Chang, C. E.; Potter, M. J.; Gilson, M. K., *J. Phys. Chem. B* **2003,** *107* (4), 1048-1055.

(57)   Henchman, R. H., *J. Chem. Phys.* **2007,** *126* (6).

(58)   Irudayam, S. J.; Henchman, R. H., *J. Phys. Condens. Matter* **2010,** *22* (28).

(59)   Gerogiokas, G.; Calabro, G.; Henchman, R. H.; Southey, M. W. Y.; Law, R. J.; Michel, J., *J. Chem. Theory Comput.* **2014,** *10* (1), 35-48.

(60)   Fukahori, T.; Nishikawa, S.; Yamaguchi, K., *Bull. Chem. Soc. Jpn.* **2004,** *77* (12), 2193-2198.

(61)   Nishikawa, S.; Fukahori, T.; Ishikawa, K., *J. Phys. Chem. A* **2002,** *106* (12), 3029-3033.

(62)   Chodera, J. D.; Mobley, D. L., *Annu. Rev. Biophys.* **2013,** *42*, 121-142.

(63)   Huang, Y.-m. M.; Chen, W.; Potter, M. J.; Chang, C.-e. A., *Biophys. J.* **2012,** *103* (2), 342-351.

(64)   Chang, C. E. A.; Chen, W.; Gilson, M. K., *Proc. Natl. Acad. Sci. U.S.A.* **2007,** *104* (5), 1534-1539.

(65)   Lin, S.-T.; Maiti, P. K.; Goddard, W. A., III, *J. Phys. Chem. B* **2010,** *114* (24), 8191-8198.

(66)   Pascal, T. A.; Schaerf, D.; Jung, Y.; Kuehne, T. D., *J. Chem. Phys.* **2012,** *137* (24).

(67)   Yin, J.; Fenley, A. T.; Henriksen, N. M.; Gilson, M. K., *J. Phys. Chem. B* **2015,** *119* (32), 10145-10155.

(68)   Tidemand, K. D.; Schonbeck, C.; Holm, R.; Westh, P.; Peters, G. H., *J. Phys. Chem. B* **2014,** *118* (37), 10889-10897.

(69)   Bellissent-Funel, M. C.; Hassanali, A.; Havenith, M.; Henchman, R.; Pohl, P.; Sterpone, F.; van der Spoel, D.; Xu, Y.; Garcia, A. E., *Chem. Rev.* **2016,** *116* (13), 7673-7697.

(70)   Qin, Y. Z.; Wang, L. J.; Zhong, D. P., *Proc. Natl. Acad. Sci. U.S.A.* **2016,** *113* (30), 8424-8429.

(71)   Berg, O. G.; Vonhippel, P. H., *Annu. Rev. Biophys. Biophys. Chem.* **1985,** *14*, 131-160.

(72)   Mondal, J.; Friesner, R. A.; Berne, B. J., *J. Chem. Theory Comput.* **2014,** *10* (12), 5696-5705.

(73)   Wilson, C.; Agafonov, R. V.; Hoemberger, M.; Kutter, S.; Zorba, A.; Halpin, J.; Buosi, V.; Otten, R.; Waterman, D.; Theobald, D. L.; Kern, D., *Science* **2015,** *347* (6224), 882-886.

(74)   Persch, E.; Dumele, O.; Diederich, F., *Angew. Chem. Int. Ed.* **2015,** *54* (11), 3290-3327.

(75)   Hobza, P., *Acc. Chem. Res.* **2012,** *45* (4), 663-672.

(76)   Snyder, P. W.; Lockett, M. R.; Moustakas, D. T.; Whitesides, G. M., *European Physical Journal-Special Topics* **2014,** *223* (5), 853-891.

(77)   Taulier, N.; Chalikian, T. V., *J. Phys. Chem. B* **2006,** *110* (25), 12222-12224.

(78)   Mobley, D. L.; Dill, K. A., *Structure* **2009,** *17* (4), 489-498.

(79)   Silver, N. W.; King, B. M.; Nalam, M. N. L.; Cao, H.; Ali, A.; Reddy, G.; Rana, T. M.; Schiffer, C. A.; Tidor, B., *J. Chem. Theory Comput.* **2013,** *9* (11), 5098-5115.

(80)   Forti, F.; Cavasotto, C. N.; Orozco, M.; Barril, X.; Luque, F. J., *J. Chem. Theory Comput.* **2012,** *8* (5), 1808-1819.

(81)   Chang, C. E. A.; McLaughlin, W. A.; Baron, R.; Wang, W.; McCammon, J. A., *Proc. Natl. Acad. Sci. U.S.A.* **2008,** *105* (21), 7456-7461.

(82)   Sharp, K., *Protein Sci.* **2001,** *10* (3), 661-667.

(83)   Faver, J. C.; Yang, W.; Merz, K. M., *J. Chem. Theory Comput.* **2012,** *8* (10), 3769-3776.

(84)   Beauchamp, K. A.; Lin, Y. S.; Das, R.; Pande, V. S., *J. Chem. Theory Comput.* **2012,** *8* (4), 1409-1414.

(85)   Lindorff-Larsen, K.; Maragakis, P.; Piana, S.; Eastwood, M. P.; Dror, R. O.; Shaw, D. E., *PLoS One* **2012,** *7* (2).

(86)   Rauscher, S.; Gapsys, V.; Gajda, M. J.; Zweckstetter, M.; de Groot, B. L.; Grubmuller, H., *J. Chem. Theory Comput.* **2015,** *11* (11), 5513-5524.

(87)   Wickstrom, L.; Deng, N. J.; He, P.; Mentes, A.; Nguyen, C.; Gilson, M. K.; Kurtzman, T.; Gallicchio, E.; Levy, R. M., *J. Mol. Recognit.* **2016,** *29* (1), 10-21.

(88)   Setny, P.; Baron, R.; Kekenes-Huskey, P. M.; McCammon, J. A.; Dzubiella, J., *Proc. Natl. Acad. Sci. U.S.A.* **2013,** *110* (4), 1197-1202.

(89)   Weiss, R. G.; Setny, P.; Dzubiella, J., *J. Phys. Chem. B* **2016,** *120* (33), 8127-8136.

(90)   Nishikawa, S.; Ugawa, T.; Fukahori, T., *J. Phys. Chem. B* **2001,** *105* (31), 7594-7597.

TOC

Binding

Desolvation

Supporting Information

Binding Thermodynamics and Kinetics Calculations Using Chemical Host and Guest: A Comprehensive Picture of Molecular Recognition

Zhiye Tang and Chia-en A. Chang*

Department of Chemistry, University of California, Riverside, CA92521

Telephone: (951) 827-7263

Email: chiaenc@ucr.edu

1.   Lengths of MD Simulations

Force Field	System	Total (μs)	Length Used (μs)	# of Bound State Conformations
GAFF	1-Propanol	1.00	1.00	100000
	1-Butanol	2.56	2.56	256000
	Methyl Butyrate	1.00	1.00	100000
	t-Butanol	1.00	1.00	100000
	1-Naphthyl Ethanol	2.00	2.00	200000
	Aspirin	1.00	1.00	100000
	2-Naphthyl Ethanol	2.50	2.50	250000
TIP3P	Water	2.50	2.50	250000
GAFF-CD	β-CD-1-Propanol	6.50	6.50	23256
	β-CD-1-Butanol	6.50	6.50	48869
	β-CD-Methyl Butyrate	7.50	7.50	111496
	β-CD-t-Butanol	8.00	8.00	86200
	β-CD-1-Naphthyl Ethanol	11.00	4.00	369283
	β-CD-Aspirin	9.50	5.50	453449
	β-CD-2-Naphthyl Ethanol	11.00	4.50	319300
	β-CD	2.50	2.50	250000
q4MD-CD	β-CD-1-Propanol	4.50	2.50	60225
	β-CD-1-Butanol	5.00	2.50	152864
	β-CD-Methyl Butyrate	4.50	2.50	214756
	β-CD-t-Butanol	4.50	2.50	214310
	β-CD-1-Naphthyl Ethanol	6.00	2.50	236130
	β-CD-Aspirin	6.00	2.50	241138
	β-CD-2-Naphthyl Ethanol	6.00	4.50	442508
	β-CD	2.60	2.60	260000

Table SI 1. The lengths of MD simulations of β-cyclodextrin (β-CD) complexes and numbers of frames used in calculations of thermodynamic properties. We used GAFF-CD and q4MD-CD for free β-CD and the complexes. We used GAFF for the free ligands and TIP3P for water in all simulations. All trajectories are resaved every 10 ps for thermodynamics calculations. The lengths used in calculations of thermodynamic properties are shown in column 4. Column 5 shows the total numbers of frames in free β-CD, free ligands, and water trajectories, and the numbers of bound state conformations in complex trajectories corresponding to the length indicated in column 4. Only bound state complex conformations were used in thermodynamics calculations. All full lengths of trajectories were used in association and dissociation rate constant calculations.

2.   Conformations of β-CD in Vacuum MD Simulations in GAFF-CD and q4MD-CD

We performed MD simulations of free β-CD in vacuum for 50 ns using GAFF-CD and q4MD-CD respectively. Periodic condition was not considered. The trajectories were saved every 10 ps and totally 5000 frames were saved for the two MD runs. The RMSDs were computed against the crystal structures and plotted in Figure SI 1.

Figure SI 1. The RMSD plots and representative conformations of β-CD from MD simulations in vacuum in GAFF-CD and q4MD-CD. X-axes of RMSD plots are numbers of frames, and y-axes are RMSD against crystal structures in Å. Representative conformations are circled on the RMSD plots. Conformations A on the two plots have RMSD roughly equal to 1.7 Å. For GAFF-CD simulation, 80.5% conformations have RMSD smaller than 1.7 Å. For q4MD-CD simulations, the percentage is 95.12%.

3.   Details of Enthalpy Decomposition Calculations

The total enthalpy change (ΔH) is decomposed into changes of solute-solute interaction ($\Delta H_{Host\text{-}Guest}$) and desolvation energy ($\Delta H_{desolvation}$), where $\Delta H_{desolvation}$ includes interaction changes of host-water ($\Delta H_{Host\text{-}Water}$), guest-water ($\Delta H_{Guest\text{-}Water}$), and water-water ($\Delta H_{Water\text{-}Water}$). Because both conformational changes and intermolecular interactions contribute to $\Delta H_{Host\text{-}Guest}$, we further provide decomposition analysis of $\Delta H_{Host\text{-}Guest}$ into changes of host and guest conformational enthalpy $\Delta H_{Host\ Conf}$ and $\Delta H_{Guest\ Conf}$, respectively, and their interaction energy change ($\Delta H_{Solute\ Inter}$). To compute these decomposed energy terms, we also calculated average potential energies of water ($<E_{Water}>_{Complex}$), water+host ($<E_{Water+Host}>_{Complex}$), water+guest ($<E_{Water+Guest}>_{Complex}$), host+guest ($<E_{Host+Guest}>_{Complex}$), host ($<E_{Host}>_{Complex}$) and guest ($<E_{Guest}>_{Complex}$) in the complex trajectories, water ($<E_{Water}>_{Host}$), water+host ($<E_{Water+Host}>_{Host}$) and host ($<E_{Host}>_{Host}$) in the free β-CD trajectory, water ($<E_{Water}>_{Guest}$), water+guest ($<E_{Water+Guest}>_{Guest}$) and guest ($<E_{Guest}>_{Guest}$) in the free ligand trajectories, in addition to $<E>_{Complex}$, $<E>_{Water}$, $<E>_{Host}$ and $<E>_{Guest}$. Then we used eqs. 3.1 to 3.8 to calculate the decomposed energy terms.

$$\Delta H_{Host\text{-}Guest} = <E_{Host+Guest}>_{Complex} - <E_{Host}>_{Host} - <E_{Guest}>_{Guest} \qquad \text{eq. 3.1}$$

$$\Delta H_{Host\ Conf} = <E_{Host}>_{Complex} - <E_{Host}>_{Host} \qquad \text{eq. 3.2}$$

$$\Delta H_{Guest\ Conf} = <E_{Guest}>_{Complex} - <E_{Guest}>_{Guest} \qquad \text{eq. 3.3}$$

$$\Delta H_{Solute\ Inter} = <E_{Host+Guest}>_{Complex} - <E_{Host}>_{Complex} - <E_{Guest}>_{Complex} \qquad \text{eq. 3.4}$$

$$\Delta H_{Host\text{-}Water} = (<E_{Water+Host}>_{Complex} - <E_{Water}>_{Complex} - <E_{Host}>_{Complex}) -$$
$$(<E_{Water+Host}>_{Host} - <E_{Water}>_{Host} - <E_{Host}>_{Host}) \qquad \text{eq. 3.5}$$

$$\Delta H_{Guest\text{-}Water} = (<E_{Water+Guest}>_{Complex} - <E_{Water}>_{Complex} - <E_{Guest}>_{Complex}) -$$
$$(<E_{Water+Guest}>_{Guest} - <E_{Water}>_{Guest} - <E_{Guest}>_{Guest}) \qquad \text{eq. 3.6}$$

$$\Delta H_{Water\text{-}Water} = <E_{Water}>_{Complex} + <E>_{Water} - <E_{Water}>_{Host} - <E_{Water}>_{Guest} \qquad \text{eq. 3.7}$$

$$\Delta H_{Desolvation} = \Delta H_{Host\text{-}Water} + \Delta H_{Guest\text{-}Water} + \Delta H_{Water\text{-}Water} \qquad \text{eq. 3.8}$$

4.   Surface Area, Hydrogen Bond and Solvation Water

Figure SI 2. The hydrogen-bonding (H-bond) sites on β-CD. β-CD has totally 56 H-bonding sites, and on each of the seven glucopyranose units there are five H-bond acceptors (top) and three H-bond donors (bottom). The H-bonding sites are shown in spheres.

We evaluated the total solvent accessible surface area (SA_{Total}), polar solvent accessible surface area (SA_{Polar}) and non-polar solvent accessible surface area ($SA_{Non\ Polar}$) of β-CD. The solvent accessible surface area was evaluated by generating spherical meshes with radius set to the van der Waals radii of atoms plus 1.4 Å on the atoms and counting the exposed surface areas of the meshes. SA_{Total}, SA_{Polar} and $SA_{Non\ Polar}$ are evaluated by summing up the surface areas of all atoms, of oxygen atoms, and of carbon and hydrogen atoms, respectively. Values are averaged from all conformations of free β-CD and bound state conformations of complex trajectories.

To estimate the H-bonds with water molecules, we computed the average occupancy percentages of the 56 H-bonding sites (Figure SI 2) on β-CD. We only counted the H-bonds formed with water molecules on these H-bonding sites. For each H-bonding site, the number of conformations with water molecules forming H-bond to the H-bonding site was divided by the total number of conformations to compute the percentage of water H-bond occupancy percentage. The percentages of each H-bonding site were then averaged to compute the percentage of the conformation. Finally, the percentage of conformations were then average for all conformations of free β-CD or bound state conformations of complexes to compute the average H-bond occupancy percentage ($\Delta\%_{H\text{-}Bond}$).

The changes of total surface area (ΔSA_{Total}), polar surface area (ΔSA_{Polar}), non-polar surface area ($\Delta SA_{Non\ Polar}$), and average percentage of H-bond occupancy ($\Delta\%_{H\text{-}Bond}$) are calculated by subtracting values of free β-CD from values of complexes. The results are summarized in Table SI 2.

GAFF-CD								
Guest	SA_{Total}	SA_{Polar}	$SA_{Non Polar}$	%$_{H-Bond}$	ΔSA_{Total}	ΔSA_{Polar}	$\Delta SA_{Non Polar}$	Δ%$_{H-Bond}$
β-CD-1-Propanol	1019	216	803	34.25%	-37	-6	-31	-3.58%
β-CD-1-Butanol	1024	217	807	32.17%	-32	-5	-27	-5.66%
β-CD-Methyl Butyrate	998	209	789	32.31%	-58	-13	-45	-5.52%
β-CD-t-Butanol	1038	221	817	32.78%	-18	-1	-17	-5.04%
β-CD-1-Naphthyl Ethanol	994	211	783	29.64%	-62	-11	-52	-8.18%
β-CD-Aspirin	1002	212	790	29.86%	-54	-10	-44	-7.97%
β-CD-2-Naphthyl Ethanol	986	210	776	29.86%	-70	-12	-58	-7.97%
β-CD	1056	222	834	37.83%				
q4MD-CD								
Guest	SA_{Total}	SA_{Polar}	$SA_{Non Polar}$	%$_{H-Bond}$	ΔSA_{Total}	ΔSA_{Polar}	$\Delta SA_{Non Polar}$	Δ%$_{H-Bond}$
β-CD-1-Propanol	1058	228	829	28.74%	-66	-5	-61	-2.41%
β-CD-1-Butanol	1045	226	819	28.64%	-78	-7	-71	-2.51%
β-CD-Methyl Butyrate	1030	222	808	28.09%	-94	-12	-82	-3.05%
β-CD-t-Butanol	1055	231	824	28.44%	-68	-2	-66	-2.70%
β-CD-1-Naphthyl Ethanol	995	213	782	26.61%	-129	-21	-108	-4.54%
β-CD-Aspirin	1005	216	789	27.31%	-118	-17	-101	-3.83%
β-CD-2-Naphthyl Ethanol	993	212	781	27.18%	-130	-21	-109	-3.96%
β-CD	1123	234	890	31.14%				

Table SI 2. The solvent accessible surface area and H-bonds with water of β-CD in complexes and free state in GAFF-CD and q4MD-CD. The total surface area (SA_{Total}), polar surface area (SA_{Polar}) and non-polar surface area ($SA_{Non Polar}$) are in $Å^2$. The average H-bond occupancy percentage of 56 H-bond sites on β-CD (%$_{H-Bond}$) are in %. The corresponding changes (Δ) are calculated by subtracting values of free β-CD from values of complexes.

5.   Definition of Probability Distribution of Conformations of Solutes

Figure SI 3. A representative dihedral population histogram used to define the conformation of individual dihedral angles. The population distribution is in black. Four Gaussian functions (1 in Blue, 2 in cyan, 3in yellow and 4 in orange) were fitted onto the four population peaks and the averages (μ) and standard deviations (σ) are labeled.

We used the combinations of dihedral angles to define the conformations of β-CD and guest molecules, and used the probability distribution of conformations in the Gibbs entropy formula (eq. 1 in main text) to compute the internal entropy of β-CD and guest molecules. 14 dihedrals defined in Figure 1 were selected for β-CD while every dihedral except methyl and hydroxyl rotations were selected for the ligands. We obtained the population histograms of the dihedral angles from MD simulation. We defined the conformation of selected dihedral angles use the Gaussian functions illustrated in Figure SI 3. Given any dihedral angle, $\boldsymbol{\alpha}$, we define it in dihedral conformation $\boldsymbol{i}$ if,

$$\text{abs}\left(\frac{\text{Subtract}[\alpha,\mu(i)]}{\sigma(i)}\right) = \min\left\{\text{abs}\left(\frac{\text{Subtract}[\alpha,\mu(1)]}{\sigma(1)}\right), \text{abs}\left(\frac{\text{Subtract}[\alpha,\mu(2)]}{\sigma(2)}\right), \dots, \text{abs}\left(\frac{\text{Subtract}[\alpha,\mu(n)]}{\sigma(n)}\right)\right\} \quad \text{eq. 5.1}$$

where *abs(x)* is to take the absolute value of $\boldsymbol{x}$, $\mu(\boldsymbol{i})$ and $\sigma(\boldsymbol{i})$ indicate the average and standard deviation of Gaussian function $\boldsymbol{i}$, *min(…)* is to take the minimum value of given values, the $\boldsymbol{n}$ denotes the total number

of peaks of the selected dihedral, and *Subtract* is an operator defined as eq. 5.2 used to subtract two dihedral angles,

$$\text{Subtract}[a, b] = \text{atan2}[\sin(a)\cos(b) - \sin(b)\cos(a), \cos(a)\cos(b) + \sin(a)\sin(b)] \qquad \text{eq. 5.2}$$

where *atan2* is the standard arctangent function that convert from values of $\sin(\theta)$ and $\cos(\theta)$ to θ. We treated conformations that fall into each peak as one identical conformation of the corresponding dihedral angle, and neglected the width of the peaks shown in Figure SI 3.

With the definition of individual dihedral angles, we used the different combinations of individual dihedral conformations to define conformations of β-CD and ligands. Distinct combinations of dihedral angle conformations are considered as distinct conformations of β-CD and ligands. In β-CD case, its seven-fold symmetry was considered. The population distributions of β-CD and ligands were then calculated from the distinct conformations.

6. Solvation Entropy Calculation Using Water Molar Entropy

Using grid cell method [1-3], the water molar entropy was evaluated by separating it into vibrational entropy ($S_{Water\ Vib}$) and conformational entropy ($S_{Water\ Conf}$), as in eq. 6.1,

$$S_{Water} = S_{Water\ Vib} + S_{Water\ Conf} \qquad \text{eq. 6.1}$$

$S_{Water\ Vib}$ is further separated into translational entropy ($S_{Water\ Trans}$) and rotational entropy ($S_{Water\ Rot}$) as in eq. 6.2,

$$S_{Water\ Vib} = S_{Water\ Trans} + S_{Water\ Rot} \qquad \text{eq. 6.2}$$

$S_{Water\ Trans}$ and $S_{Water\ Rot}$ are computed from forces and torques on the three principal axes of the target water molecule as in eqs. 6.3 and 6.4,

$$S_{Water\ Trans} \text{ or } S_{Water\ Rot} = R\sum_{i=1}^{6}\left(\frac{\alpha_i}{\alpha_i-1} - \ln(1-e^{-\alpha_i})\right) \qquad \text{eq. 6.3}$$

$$\alpha_i = \begin{cases} \dfrac{F_i}{2k_BT}\dfrac{h}{(2\pi mk_BT)^{1/2}} & (i=1,2,3,\text{for trans}) \\[2ex] \dfrac{\tau_i}{2k_BT}\dfrac{h}{(2\pi I_i k_BT)^{1/2}} & (i=4,5,6,\text{for rot}) \end{cases} \qquad \text{eq. 6.4}$$

where, R is the gas constant, k_B is the Boltzmann constant, h is the Planck constant, T is the temperature, m is the mass of a water molecule, F_i are the force and τ_i are the torques on the three principal axes. The gas phase values of moments of inertia I_i about x, y, and z axes are used.

Figure SI 4. The definition of a loose H-bond. It is based on normal H-bond definition except that the complementary angle of X-H...Y (a) is less than 80° and the distance between H and Y (d) is less than 2.65Å.

Figure SI 5. The water conformational entropy definition. It is based on the Pauling's model. Around the target water molecule, the donor waters (DW) and acceptor waters (AW) are defined with the loose H-bond definition and any water molecule that is within 3.2 Å from the oxygen atom of the target water molecule but cannot form a loose H-bond is considered as a surrounding water (SW).

We derived model to evaluate water conformational entropy ($S_{Water\ conf}$) based on Pauling's model [2-4]. First a loose H-bond is defined as shown in Figure SI 4, where the complementary angle of X-H…Y (a) must be less than 80˚ and the H…Y distance (d) must be less than 2.65 Å. The purpose of the loose definition is to capture the forming and breaking H-bonds due to thermal fluctuation. Based on that, the conformational entropy is defined according to Figure SI 5. The water molecules around the target water molecule are classified into three groups, 1) acceptor waters (AW), which are the loose H-bond acceptors from the target water, 2) donor waters (DW), which are the loose H-bond donors to the target water, and 3) surrounding waters (SW), whose oxygen atoms are within 3.2 Å of the oxygen atom of the target water molecule but form no H-bond with it. The number of conformations (Ω_{Water}) of the target water is calculated using eq. 6.5,

$$\Omega_{Water} = C^2_{2+\#\ of\ DW+\#\ of\ SW} \cdot C^{\#\ of\ SW}_{\#\ of\ DW+\#\ of\ SW} \cdot \left(\frac{1}{2}\right)^{\#\ of\ DW+\#\ of\ SW} \qquad \text{eq. 6.5}$$

In the first term, both "2"s mean the two AW sites, regardless of the existence of water molecules there. This term can be interpreted as choosing two sites of the water molecules around as the acceptor sties. The second term means in the rest of the water around to choose # of SW of waters as the SW. The appearance of the last term is to compensate the double counting of water molecules around. For example, when there are exactly two AWs and two DWs, $\Omega_{Water} = C_4^2 \cdot C_2^0 \cdot \left(\frac{1}{2}\right)^2 = 6 \times \frac{1}{4} = 1.5$, which obeys the Pauling's model. When there are two AWs, two DWs and one SW, $\Omega_{Water} = C_5^2 \cdot C_3^1 \cdot \left(\frac{1}{2}\right)^3 = 6 \times \frac{1}{4} = 3.75$, which means an additional water molecule around increases the number of conformations of the target water. Finally, the conformational entropy of the water is calculated as eq. 6.6,

$$S_{Water\ Conf} = R\ln(\Omega_{Water}) \qquad\qquad \text{eq. 6.6}$$

The solvation entropy can be calculated using water molar entropy. The solvation energy of species X can be computed using eq. 6.7,

$$\Delta S_{Solv.} = N(water) \times (S_{Water}(X) - S_{Water}(0)) \qquad\qquad \text{eq. 6.7}$$

where *N(water)* is given as *# of Water* in Tables SI 3 to 5, $S_{Water}(X)$ is the molar water entropy with species *X*, and $S_{Water}(0)$ is the molar water entropy of bulk water. The computed water molar entropy and solvation entropy are summarized in Tables SI 3 to 5.

System	S$_{Water}$ (J/mol/K)					-TΔS$_{Solv.}$ (kcal/mol)					# of Water
	Trans	Rot	Vib	Conf	Total	Trans	Rot	Vib	Conf	Total	
β-CD-1-Propanol	47.69	20.83	68.52	5.01	73.53	4.62	0.59	5.21	2.48	7.69	403.01
β-CD-1-Butanol	47.7	20.85	68.54	5.01	73.55	4.44	0.23	4.66	2.53	7.19	401.58
β-CD-Methyl Butyrate	47.7	20.84	68.54	5.01	73.55	4.19	0.45	4.64	2.48	7.12	400.41
β-CD-t-Butanol	47.71	20.84	68.55	5.01	73.56	4.03	0.43	4.45	2.48	6.93	401.74
β-CD-1-Naphthyl Ethanol	47.72	20.85	68.56	5.01	73.57	3.74	0.21	3.95	2.46	6.41	397.53
β-CD-Aspirin	47.71	20.85	68.56	5.01	73.57	3.91	0.24	4.15	2.45	6.59	398.39
β-CD-2-Naphthyl Ethanol	47.72	20.85	68.57	5.01	73.58	3.69	0.13	3.82	2.47	6.29	397.49
β-CD	47.67	20.83	68.5	5.02	73.52	5.1	0.78	5.88	2.19	8.07	407.14

Table SI 3. The molar water entropy and solvation entropy of free β-CD and β-CD complexes in GAFF-CD. The molar water entropy (S$_{Water}$) is separated into vibrational (S$_{Water\ Vib}$), and conformational (S$_{Water\ Conf}$) terms. The vibrational (S$_{Water\ Vib}$) term is further separated into translational (S$_{Water\ Trans}$) and rotational (S$_{Water\ Rot}$) terms. The molar water entropy terms are in J/mol/K. The solvation entropy (-TΔS$_{Solv.}$) is separated into vibrational (-TΔS$_{Solv.\ Vib}$), and conformational (-TΔS$_{Solv.\ Conf}$) terms. The vibrational (-TΔS$_{Solv.\ Vib}$) term is further separated into translational (-TΔS$_{Solv.\ Trans}$) and rotational (-TΔS$_{Solv.\ Rot}$) terms. The solvation energy terms at 298K are in kcal/mol. The numbers of water molecules are corresponding to a volume of 13520 Å^3.

System	S_Water (J/mol/K)					-TΔS_Solv. (kcal/mol)					# of Water
	Trans	Rot	Vib	Conf	Total	Trans	Rot	Vib	Conf	Total	
β-CD-1-Propanol	47.71	20.85	68.56	5.01	73.57	4.11	0.09	4.20	2.35	6.55	401.93
β-CD-1-Butanol	47.71	20.85	68.57	5.01	73.58	3.89	-0.02	3.86	2.30	6.16	400.86
β-CD-Methyl Butyrate	47.72	20.86	68.58	5.01	73.59	3.71	-0.16	3.55	2.35	5.90	400.01
β-CD-t-Butanol	47.72	20.85	68.57	5.01	73.58	3.82	0.05	3.88	2.34	6.22	401.15
β-CD-1-Naphthyl Ethanol	47.72	20.87	68.59	5.01	73.60	3.57	-0.49	3.08	2.34	5.43	397.77
β-CD-Aspirin	47.72	20.86	68.58	5.01	73.59	3.68	-0.22	3.46	2.32	5.77	398.05
β-CD-2-Naphthyl Ethanol	47.73	20.87	68.60	5.01	73.61	3.44	-0.60	2.83	2.38	5.22	397.45
β-CD	47.69	20.84	68.54	5.02	73.56	4.50	0.27	4.77	2.20	6.98	406.33

Table SI 4. The molar water entropy and solvation entropy of free β-CD and β-CD complexes in q4MD-CD. Refer to Table SI 3 for meanings of symbols.

System	S_{Water} (J/mol/K)					$-T\Delta S_{Solv.}$ (kcal/mol)					# of Water
	Trans	Rot	Vib	Conf	Total	Trans	Rot	Vib	Conf	Total	
1-Propanol	47.80	20.83	68.63	5.07	73.71	1.54	0.71	2.26	0.68	2.93	441.36
1-Butanol	47.83	20.85	68.68	5.07	73.75	0.52	0.17	0.69	0.80	1.49	440.41
Methyl Butyrate	47.82	20.82	68.64	5.06	73.70	1.01	1.03	2.04	1.00	3.04	439.02
t-Butanol	47.85	20.85	68.70	5.07	73.76	0.09	0.11	0.20	0.87	1.08	440.53
1-Naphthyl Ethanol	47.86	20.86	68.72	5.05	73.77	-0.23	-0.39	-0.62	1.24	0.62	436.48
Aspirin	47.81	20.84	68.65	5.06	73.71	1.17	0.36	1.53	1.07	2.61	437.33
2-Naphthyl Ethanol	47.86	20.86	68.72	5.05	73.78	-0.26	-0.41	-0.67	1.22	0.55	436.41
Water	47.85	20.85	68.70	5.09	73.80	0.00	0.00	0.00	0.00	0.00	565.84

Table SI 5. The molar water entropy and solvation entropy of free ligands and water. Refer to Table SI 3 for meanings of symbols.

Alternatively, the solvation water entropy can be computed using forces ($F_{i,S}$), torques ($\tau_{i,S}$), and number of water conformation (Ω_S) directly using eqs. 6.8 to 6.11, instead of computing the water molar entropy first,

$$\Delta S_{Solv.Trans} = NRln\left(\prod_{i=1}^{3} \frac{F_{i,W}}{F_{i,S}}\right) \qquad \text{eq. 6.8}$$

$$\Delta S_{Solv.Rot} = NRln\left(\prod_{i=1}^{3} \frac{\tau_{i,W}}{\tau_{i,S}}\right) \qquad \text{eq. 6.9}$$

$$\Delta S_{Solv.\ Vib} = \Delta S_{Solv.\ Trans} + \Delta S_{Solv.\ Rot} \qquad \text{eq. 6.10}$$

$$\Delta S_{Solv.Conf} = NRln\left(\frac{\Omega_S}{\Omega_W}\right) \qquad \text{eq. 6.11}$$

where, R is the gas constant, $F_{i,W}$ and $F_{i,S}$ are the averages of forces on three principal axes without solute and with solute, $\tau_{i,W}$ and $\tau_{i,S}$ are the averages of torques on three principal axes without and with solute, Ω_W and Ω_S are the average of number of conformations of water molecules without and with solute, and N is the number of water molecules in the sub water box defined in Figure SI 6.

Figure SI 6. Illustration of water entropy calculation using free β-CD as representation. The original water box (deep blue) is align to the crystal structure of β-CD (deep green). One representative water molecule under calculation is shown in the sub water box (black lined box). Any periodic instances of atoms within the cutoff (purple circle) generated from original water box and periodic images of the original water box (light blue and light green) are used to calculate the forces and torques on the water molecule under calculation.

Practically, the complex and β-CD trajectories were aligned against the crystal structure of free β-CD. The ligand trajectories were aligned against the initial conformation of the ligand. Water molecules in a sub water box centered on the β-CD (in the case of free β-CD and complexes) and guest (in the case of free guests) with a size of 20Å ×26Å ×26Å were considered in the water entropy calculation. For the empty water box, a sub water box with size 22Å ×22Å ×22Å was used. A cutoff 50.0 Å was used for force and torque calculation in the vibrational term, and periodical atom instances generated from the full water box within this cutoff were used to calculate the forces and torques on the water molecules in the sub water box. The grid bin size was set to 1 Å in the grid cell method. The illustration of the water box for water entropy calculation is in Figure SI 6.

7.   Solvation Water

We also counted the numbers of water molecules in the first sovlation shells of free guest ($\#_{Guest}$), free β-CD ($\#_{Host}$) and the complex ($\#_{Complex}$) from the MD trajectory. We counted the number oxygen atoms in the water molecules within 3.5 Å of all atoms in the corresponding solute and took the average of the number of waters over the entire trajectory for free guest and free host and over the bound state conformations in trajectories for complexes. The changes of the numbers of water molecules in the first solvation shells upon ligand binding ($\Delta\#$) are calculated by $\#_{Guest} + \#_{Host} - \#_{Complex}$. The results are summarized in Table SI 6.

Guest	GAFF-CD				q4MD-CD			
	$\#_{Guest}$	$\#_{Host}$	$\#_{Complex}$	$\Delta\#$	$\#_{Guest}$	$\#_{Host}$	$\#_{Complex}$	$\Delta\#$
1-Propanol	15.21	77.92	80.46	12.67	15.21	81.43	80.21	16.43
1-Butanol	17.28	77.92	79.62	15.58	17.28	81.43	79.26	19.45
Methyl Butyrate	19.67	77.92	78.99	18.6	19.67	81.43	79.12	21.98
t-Butanol	16.66	77.92	79.75	14.83	16.66	81.43	79.5	18.59
1-Naphthyl Ethanol	23.53	77.92	78.47	22.98	23.53	81.43	78.53	26.43
Aspirin	22.02	77.92	78.83	21.11	22.02	81.43	78.92	24.53
2-Naphthyl Ethanol	24.02	77.92	79.5	22.43	24.02	81.43	79.18	26.27

Table SI 6. The number of water molecules in the first solvation shells of free guest ($\#_{Guest}$), free host ($\#_{Host}$), complex ($\#_{Complex}$) and number of water molecules released from the first solvation shell to the bulk ($\Delta\#$). Values are obtained by averaging the numbers of water molecules within 3.5 Å of atoms in the solute over the corresponding trajectories. The change is computed by $\#_{Guest} + \#_{Host} - \#_{Complex}$.

8.   Solute Concentrations Used in Association Rate Constant Calculations

We used this protocol to calculate the concentration used in association rate constant (k_{on}) calculation and the concentrations are summarized in Table SI 7. We obtained the cell sizes of free state conformations in the complex trajectories saved in original trajectories, and used eq. 8.1 to calculate the volume of the water box,

$$\text{Volume} = \frac{1}{N}\sum_{i=1}^{N}(\text{Cell Size X(i)} \times \text{Cell Size Y(i)} \times \text{Cell Size Z(i)}) \qquad \text{eq. 8.1}$$

where N is the number of free state conformations in the complex trajectory, and Cell Size X, Y, Z are the cell sizes of the water box in three dimensions.

Then we used eq. 8.2 to calculate the solute concentration ([solute]) of the free state in the complex trajectories,

$$[\text{Solute}] = \frac{1/N_A}{\text{Volume}} \qquad \text{eq. 8.2}$$

where N_A is the Avogadro constant, and "1" is the number of complex particles in the trajectory.

Guest	GAFF-CD	q4MD-CD
1-Propanol	0.03079	0.03079
1-Butanol	0.03078	0.03078
Methyl butyrate	0.03076	0.03076
t-Butanol	0.03078	0.03077
1-Naphthyl Ethanol	0.03070	0.03072
Aspirin	0.03072	0.03072
2-Naphthyl Ethanol	0.03070	0.03071
		Unit: M

Table SI 7. The solute concentrations of β-CD and guest molecules used in association rate constant (k_{on}) calculation in GAFF-CD and q4MD-CD. The concentrations correspond to free state conformations in the complex trajectories. All values are in mol/L (M).

9.   Diffusion Controlled Association Rate Constants Estimated from Diffusion Coefficient

System	D_{Expt} (m^2/s)	D_{TIP3P} (m^2/s)	D_{MD} (m^2/s)	$k_{on\_diffuse}$ (1/s·M)
1-Butanol	8.7×10^{-10}	2.4×10^{-9}	5.9×10^{-9}	4.45×10^{10}
Methyl Butyrate	8.1×10^{-10}	2.3×10^{-9}	5.2×10^{-9}	4.07×10^{10}
t-Butanol	8.7×10^{-10}	2.4×10^{-9}	5.7×10^{-9}	4.33×10^{10}
1-Propanol	9.4×10^{-10}	2.6×10^{-9}	6.6×10^{-9}	4.76×10^{10}
1-Naphthyl Ethanol	7.1×10^{-10}	2.0×10^{-9}	3.8×10^{-9}	3.26×10^{10}
Aspirin	7.3×10^{-10}	2.0×10^{-9}	3.7×10^{-9}	3.14×10^{10}
2-Naphthyl Ethanol	7.0×10^{-10}	2.0×10^{-9}	3.8×10^{-9}	3.28×10^{10}
β-CD	4.1×10^{-10}	1.1×10^{-9}	7.2×10^{-10}	/

Table SI 8. Diffusion coefficients of ligands and β-CD and diffusion controlled association rate constants ($k_{on\_diffuse}$) estimated from the diffusion coefficients. Diffusion coefficients (in m^2/s) are estimated by eq. 9.1 using water viscosities from experiments (D_{Expt}) and TIP3P water model (D_{TIP3P}), and directly measured from MD trajectories (D_{MD}). $k_{on\_diffuse}$ values are estimated using D_{MD} in eq. 9.2 for each β-CD-guest complex.

$$D = kT/6\pi\eta a \qquad\qquad \text{eq. 9.1}$$

where k is the Boltzmann constant, T is the temperature (298K), η is the viscosity of water, and a is the radius of solute.

$$k_{on\_diffuse} = 4\pi N_A DR \qquad\qquad \text{eq. 9.2}$$

where N_A is the Avogadro constant, D is the sum of the diffusion coefficient of β-CD and guest, and R is the sum of radii of β-CD and guest.

10.   Uncertainty Evaluation of Computed Properties

We evaluated the uncertainty of all computed properties. For potential energy terms ($<E>_{Complex}$, $<E>_{Water}$, $<E>_{Host}$, and $<E>_{Guest}$), we used standard deviation of mean to evaluate the uncertainty in the potential energy and it was calculated by using block analysis [5] with 2 ns in a block. The uncertainties of $<E>_{Complex}$, $<E>_{Water}$, $<E>_{Host}$, and $<E>_{Guest}$ were added in quadrature to get the uncertainty of binding enthalpy (ΔH). For properties *A* other than total binding enthalpy, we used bootstrap error analysis to get the uncertainties. We generated random subsets of the data, and calculated values of *A* from the random subsets, and finally calculated the standard deviations of the values of *A* from the random subsets, as the uncertainty of *A*. In principle, roughly 80% of the data were used to generate the random subsets. The number of random subsets is 100000 if non-repeated random subsets can be generated or otherwise the maximum number of possible subsets was used.

We computed the conformational population distribution of β-CD and used the Gibbs entropy formula (eq. 1) to evaluate $S_{Host\ Complex}$, $S_{Host\ Free}$, $S_{Guest\ Int\ Complex}$ and $S_{Guest\ Int\ Free}$. For $S_{Host\ Complex}$, we generated the random subsets of the conformations, and reconstructed the conformational population distributions for each subset, and calculated the standard deviation of $S_{Host\ Complex}$ obtained from each subset as the uncertainty of $S_{Host\ Complex}$. We did identical evaluations for uncertainty of $S_{Host\ Free}$, $S_{Guest\ Int\ Complex}$ and $S_{Guest\ Int\ Free}$. For example, for the free β-CD in GAFF-CD, we generated 100000 subsets with 200000 random conformations from the total 250000 conformations, and evaluated $S_{Host\ Free}$ using the strategy above. The uncertainty of ΔS_{Host} and $\Delta S_{Guest\ Int}$ were evaluated by summing uncertainties of $S_{Host\ Complex}$ and $S_{Host\ Free}$ and by summing uncertainties of $S_{Guest\ Int\ Complex}$ and $S_{Guest\ Int\ Free}$, respectively. We evaluated $S_{Guest\ Ext\ Complex}$ using eq. 2. Therefore, we used similar strategy as in internal entropy uncertainty evaluation to create the random subsets of conformations, and reconstructed the spatial and angular distribution histogram for evaluation of $S_{Guest\ Ext\ Complex}$ in each subset using eq. 2. The uncertainty was computed, again, as the standard deviation of $S_{Guest\ Ext\ Complex}$ obtained from the subsets. The uncertainty of solute entropy change (ΔS_{Solute}) was evaluated by summing uncertainties of ΔS_{Host}, $\Delta S_{Guest\ Int}$, and $S_{Guest\ Ext\ Complex}$. Note that $S_{Guest\ Ext\ Free}$ was computed using exact analytical equations so this term does not have uncertainty.

For uncertainties of $S_{Water\ Complex}$, $S_{Water\ Host}$ and $S_{Water\ Guest}$, the calculation is slightly different in the way the random subsets were created. We saved the water entropy values for every 10-ps-block (1000 frames) along the trajectories and obtained N water entropy values from each block. Then we generated subsets that contain random collections of roughly 0.8N water entropy values from the N values. The percentage

P% is changed for a suitable combination number $C_{N, P\%N}$ so that $C_{N, P\%N}$ is within 100000 to 4294967296 (the greatest number in unsigned integer or 2^{32}). The standard deviations of water entropy values averaged from the random subsets were reported as the uncertainties of the water entropy terms. For example, for β-CD-1-butanol complex in GAFF, we used 45 blocks from the total 49 blocks to generate 211876 ($C_{49, 45}$) subsets, and evaluated the uncertainty using first 100000 subsets. The uncertainties of water entropy change (ΔS_{Water}) were evaluated by summing up the uncertainties of $S_{Water\ Complex}$, $S_{Water\ Host}$ and $S_{Water\ Guest}$.

The uncertainty of binding entropy (ΔS) was evaluated by summing up the uncertainties of ΔS_{Solute} and ΔS_{Water}. The uncertainty of binding free energy from eq. 5 (ΔG_{Comp1}) was evaluated by summing up the uncertainties of ΔH and ΔS.

For association (k_{on}) and dissociation (k_{off}) rate constants, we created the random subsets of the time lengths of bound/free periods, and calculated the rate constants from these subsets, and finally computed the standard deviations of the rate constants from these subsets, as the uncertainties of k_{on} and k_{off}. The uncertainty of K_{eq} are evaluated based on subsets of random combinations of k_{on} and k_{off} values obtained from the random subsets used to evaluated uncertainty of k_{on} and k_{off}. For example, for β-CD-1-propanol complex in GAFF-CD, we used 34 bound state period lengths out of total 39 to generate 575757 ($C_{39, 34}$) random subsets, computed the uncertainty of k_{on} using the standard deviation of the k_{on} values from the first 100000 subsets.

11.   Experimental Data of ΔG, ΔH and –TΔS of β-CD-Alcohol Complexes Using ITC and UV

Ligand	Expt. Method	ΔG (kcal/mol)	ΔH (kcal/mol)	-TΔS (kcal/mol)
cyclohexanol	cal	-5.95	-1.67	-4.28
1,3-butanediol	cal	-5.57	-0.33	-5.23
1-pentanol	cal	-5.40	0.53	-5.93
1-hexanol	cal	-5.14	0.14	-5.28
phenol	cal	-4.54	-2.63	-2.15
trans-4-methylcyclohexanol	cal	-4.54	-2.17	-2.37
cis-4-methylcyclohexanol	cal	-4.33	-2.29	-2.03
1-methylcyclohexanol	cal	-4.18	-2.29	-1.89
(+-)-cis-2-methylcyclohexanol	cal	-4.08	-2.37	-1.72
4-iodophenol	cal	-4.06	-3.85	-0.22
3-methylcyclohexanol	cal	-3.98	-2.09	-1.90
(+-)-trans-2-methylcyclohexanol	cal	-3.91	-2.07	-1.85
2,2-dimethyl-1-propanol	uv	-3.70	-2.10	-1.51
cyclohexanol	uv	-3.66	-2.39	-1.22
3-nitrophenol	cal	-3.32	-2.89	-0.43
4-chlorophenol	uv	-3.27	-3.82	0.48
1-hexanol	uv	-3.18	0.10	-3.27
2-chlorophenol	uv	-3.13	-4.54	1.43
3-chlorophenol	uv	-3.13	-4.54	1.20
resorcinol	cal	-2.77	-4.35	1.55
1-pentanol	uv	-2.46	1.10	-3.56
5-methylresorcinol	cal	-2.34	-5.07	2.68
(R)-(-)-2-pentanol	cal	-2.08	0.88	-2.99
(S)-(+)-2-pentanol	cal	-2.06	0.98	-3.06
1-butanol	uv	-1.67	0.69	-2.34
2-propanol	cal	-0.57	2.65	-3.20

Table SI 9. Experimental data of ΔG, ΔH and –TΔS of β-CD-alcohol complexes using calorimetry (cal) and spectrophotometry (uv). The data are taken from ref [6].

Figure SI 7. Plot of experimental ΔH and –TΔS against experimental ΔG of β-CD-alcohol complexes using calorimetry and spectrophotometry. The plot uses values from Table SI 9. Enthalpy values are indicated by bordered rectangle, and entropy values are indicated by circles. The enthalpy and entropy of the same ligand are rendered in the same color. For ligands with ΔG ranging from 0 to -3 and -5 to -6 kcal/mol, –TΔS is the major contribution to ΔG; for ligands with ΔG ranging from -3 to -5 kcal/mol, both ΔH and –TΔS contributes to ΔG.

12.   Convergence of Enthalpy Calculations (<E>)

We investigated the convergence of enthalpy calculation by plotting average potential energy (<E> in kcal/mol) against number of frames used in the average in Figures SI 8 to 10. We calculate the standard deviation (SD) of the last region of the average potential energy to estimate the fluctuation in the calculated enthalpy. The regions used in the SD calculation are indicated on the figures. Note that the SD is different from the uncertainty reported in the manuscript.

Figure SI 8. The enthalpy convergence of ligand and water trajectories. The convergence is represented by average potential energy (<E>) plotted against the number of frames. X-axes are numbers of frames used in the average. Y-axes are average potential energies in kcal/mol. The Y value of the black line indicates the value used in reported enthalpy. The X range of the black line indicates the region used to calculate standard deviation (SD) of the averaged potential energy, and the SD (kcal/mol) is labeled on the figures.

Figure SI 9. The enthalpy convergence of free β-CD and β-CD complexes using GAFF-CD force field. Refer to Figure SI 8 for the interpretation of the figures.

Figure SI 10. The enthalpy convergence of free β-CD and β-CD complexes using q4MD-CD force field. Refer to Figure SI 8 for the interpretation of the figures.

13.   Movies of Association and Dissociation Pathways

In the movies, carbon atoms are in cyan and oxygen atoms are in red on β-CD. The ligands are rendered in green. Hydrogen atoms are omitted. Movies are listed as below.

From GAFF-CD force field:

β-CD-aspirin direct association: GAFF-CD-b-CD-aspirin_direct_ association.swf

β-CD-aspirin direct dissociation: GAFF-CD-b-CD-aspirin_direct_dissociation.swf

β-CD-aspirin sticky association: GAFF-CD-b-CD-aspirin_sticky_ association.swf

β-CD-aspirin sticky dissociation: GAFF-CD-b-CD-aspirin_ sticky_dissociation.swf

β-CD-t-butanol direct association: GAFF-CD-b-CD-t-butanol_direct_association.swf

β-CD-t-butanol direct dissociation: GAFF-CD-b-CD-t-butanol_direct_dissociation.swf

From q4MD-CD force field:

β-CD-aspirin direct association: q4MD-CD-b-CD-aspirin_direct_ association.swf

β-CD-aspirin direct dissociation: q4MD-CD-b-CD-aspirin_direct_dissociation.swf

β-CD-aspirin sticky association: q4MD-CD-b-CD-aspirin_sticky_ association.swf

β-CD-aspirin sticky dissociation: q4MD-CD-b-CD-aspirin_ sticky_dissociation.swf

β-CD-t-butanol direct association: q4MD-CD-b-CD-t-butanol_direct_association.swf

β-CD-t-butanol direct dissociation: q4MD-CD-b-CD-t-butanol_direct_dissociation.swf

14.   Ligand External Degrees of Freedom

1-Butanol   **1-Propanol**   **Methyl Butyrate**   **t-Butanol**

1-Naphthyl Ethanol   **Aspirin**   **2-Naphthyl Ethanol**

Figure SI 11. The binding poses of β-CD and ligand complex in GAFF-CD. For simplicity, the crystal structure of β-CD is shown in *VDW* representation. One representative conformation of ligands is rendered in green *VDW* representation. The volumes occupied by the bound state conformations are shown in cyan transparent *Surface* representation. The ligand conformations are obtained by superimposing β-CD coordinates in the trajectories to the crystal structure of β-CD. Similar binding volumes can be observed in q4MD-CD.

15.   Bound and Free Period Lengths of the Complexes

	1-Propanol		1-Butanol		Methyl butyrate				t-Butanol			
	Bound	Free	Bound	Free	Bound		Free		Bound		Free	
1	1801	5834	9769	899	5287	189	639	8354	8381	119	594	20517
2	167	1521	6455	8840	885	298	2516	2143	171	119	17392	7004
3	1090	482	353	77243	2931	111	1730	23327	418	704	28343	6095
4	731	241	105	33202	342	394	14352	41390	109	108	1257	1631
5	153	57826	218	2025	302	1305	2672	1141	179	6601	54931	8604
6	107	26676	104	10408	101	14187	11150	13848	181	347	5509	42418
7	102	31381	318	32669	203	103	13537	37917	104	364	345	1985
8	268	23494	129	1050	152	150	6029	9514	156	137	4972	220
9	199	2233	136	20238	153	152	34568	5714	105	153	8409	7350
10	184	17913	205	7329	252	17996	26941	4260	111	482	57093	512
11	741	13660	161	15286	198	506	13582	1550	103	193	20737	885
12	113	42695	156	9173	151	217	7508	6697	290	1669	25645	18557
13	145	28855	141	5079	109	183	722	348	118	146	2032	30213
14	103	4623	104	4886	205	102	163	16516	178		10998	
15	100	16875	404	27418	1604	112	8441	32830	19253		3640	
16	106	5690	737	27469	1044	153	1339	2681	164		4157	
17	148	2049	320	10703	1296	449	1372	188	18665		18096	
18	108	18646	132	4843	130	175	6512	128	141		241	
19	142	23696	105	4345	1719	282	5651	1263	546		22927	
20	226	26882	116	3582	111	1116	16376	3612	156		4234	
21	239	128	182	1894	420	228	5834	6019	2234		4107	
22	364	48217	121	38219	102	138	3798	3953	120		345	
23	159	18383	264	1773	2658	103	5771	1967	114		5678	
24	103	43352	201	24159	299	171	1467	119	130		8925	
25	128	12708	315	2688	695	134	8344	20276	172		2267	
26	401	13298	102	16344	465	189	18912	14028	128		5984	
27	153	23600	142	38714	104	109	2265	6755	133		3860	
28	201	787	110	1534	390	184	5454	2378	323		3144	
29	108	720	178	894	4840	324	1586	1427	108		214	
30	119	16617	130	1318	112	1490	11015	12289	431		12414	
31	105	23846	115	4496	123		2417		152		2287	
32	127	5159	282	1977	397		708		112		6350	
33	111	10813	113	16631	126		1523		147		3421	
34	162	1775	100	23787	238		1279		263		12449	
35	253	37611	196	25623	151		387		477		2439	
36	151	11333	7042	3528	196		16738		3728		3933	
37	174	2753	418	6781	600		3703		176		5920	
38	101	16619	5800	13087	224		21087		143		14056	
39	123	993	461	1300	266		9822		177		17870	
40			199	758	156		34084		117		8000	
41			188	62738	152		4415		640		458	
42			133	18110	100		5236		236		17375	
43					9922		1564		233		9313	
44					4159		635		108		21409	
45					12086		3829		120		878	
46					101		2604		101		46032	
47					252		337		135		10473	
48					826		13528		209		31839	
49					195		4446		771		28408	
Avg.	256.8	16409.8	880.0	14596.2	1250.4		8243.3		1171.1		11732.1	
# of Periods	39	39	42	42	79		79		52		52	

Continued.

	1-Naphthyl Ethanol		Aspirin						2-Naphthyl Ethanol	
	Bound	Free	Bound			Free			Bound	Free
1	60007	2437	14310	419	228	1315	1955	8016	79434	1094
2	292	2171	3850	122	368	673	2497	2020	15270	2190
3	181	767	29400	2178	269	3586	3234	3849	4633	1380
4	6478	3866	407	169	275	3655	675	3969	411	1117
5	7108	4005	1002	255	115	148	10442	2624	156	4645
6	287	853	1042	120	146	3279	1057	2389	10619	534
7	487	3852	126	104	266	155	11040	487	609	338
8	146	11887	119	153	376	1959	3667	4619	154	227
9	293800	9182	10190	247	176	1752	1603	2014	1785	1130
10	206	4737	264	312	370	1539	586	1991	363	1250
11	512	8456	1271	115	199	2940	6580	1221	125	1230
12	259	344	1315	254	105	1103	825	5897	288	1452
13	599630	206	46697	130	3301	2840	870	6453	4976	10239
14	111	1850	2729	417	238	293	1002	4969	823	5208
15	705	2377	5417	17434	125	4436	3449	10530	223	1495
16	72801		232	5570	343	1375	1431	4637	432	1638
17			14907	848	4538	799	430	4674	150	1935
18			247	217	474	210	640	736	771	8741
19			849	119	478	650	12050	7552	35439	5824
20			8436	119	118	5095	8876	5736	125	3210
21			233	150	289	391	1931	3842	489	722
22			448	271	112	1437	1344	189	165	776
23			1096	114	9699	2252	1266	1835	717	7637
24			197	117	7116	3407	2658	680	153	4746
25			362	130	204	1072	2082	1488	192	1610
26			16445	109	764	3851	8874	1440	198	3272
27			276	218	178	1648	229	2132	8415	3760
28			124	152	274	2465	640	432	3003	4103
29			167	483	149	1415	16437	1535	180	13227
30			243	109	184	2052	1347	6401	138	1061
31			829	108	105	2467	2244	3062	101	15234
32			360	247	205	6322	6913	4841	193	3048
33			103	143	148	2129	2248	2453	147	2372
34			395	713	5682	2254	7830	347	191	2997
35			141	140	208	1480	251	1220	154	7530
36			172	175		1441	11777		20316	1138
37			387	111		11984	1962		8508	737
38			530	246		1669	331		5201	199
39			346	165		820	8042		12334	3273
40			2540	207		1290	339		188001	537
41			199	234		1169	1093		214790	1692
42			152	118		703	3593		1340	1333
43			206	122		4479	2950		3312	4452
44			37326	124		215	236		217	2548
45			2111	288		1714	3461		95640	8830
46			1585	164		549	3077		9632	7124
47			270594	155		203	239		250	1809
48			2878	318		1078	6615		277	3692
49			121	245		1708	2957		204624	
Avg.	65188.1	3799.3	4183.3			2959.6			19095.2	3423.7
# of Periods	16	15	133			133			49	48

Table SI 10. The bound and free period lengths of the complexes in GAFF-CD. All lengths of periods are in 10 ps.

	1-Propanol						1-Butanol				Methyl Butyrate		t-Butanol	
	Bound			Free			Bound		Free		Bound	Free	Bound	Free
1	1066	101	1337	927	3992	3705	1669	311	5868	2015	4075	254	2079	4536
2	726	751	155	1994	3029	356	4436	1216	2529	1686	7411	560	200	2727
3	110	2499	247	3265	157	1527	3418	1606	691	1731	2199	1595	4888	2110
4	388	1001	815	9801	107	1917	675	4533	4797	415	4978	715	1518	581
5	909	463	4468	1944	1384	7095	839	1397	403	610	7610	10426	40660	3023
6	211	1323	581	5500	5008	3284	3525	219	4471	4280	7614	1374	17924	996
7	328	105	476	10021	292	2142	3339	721	1554	3039	2318	3102	19064	4732
8	642	1930	538	1772	4243	1017	1718	4044	1887	803	59426	1601	225	2162
9	3051	104	404	798	357	268	8429	2363	1076	1430	1513	2075	42030	5703
10	321	1181	455	4932	3457	4589	7477	2014	1874	2083	19404	1543	162	1199
11	823	1104	366	962	3717	4783	1597	1987	4394	898	21044	1334	109	1899
12	1468	613	2245	1553	2755	3235	123	12994	793	4889	22974	765	12323	1948
13	644	554	111	2321	10977	2270	130	3018	337	1162	6295	852	84946	8435
14	418	1459	295	986	3376	545	3550	2099	1987	101	5330	2041	11078	728
15	749	104	216	488	6438	2996	8262	4062	1588	428	9781	1596	9311	4121
16	1158	728	105	258	794	2695	12898	6707	220	2743	45096	3450	1798	5267
17	271	642	1766	1840	100	5162	5868	1271	711	171	16006	569	5173	332
18	411	156	2064	147	8925	5071	276	1564	3439	7825	1751	5507	7726	6462
19	2034	493	151	5741	1848	4353	375	4661	2854	759	15608	701	1875	3918
20	1274	1126	257	8362	2226	660	1082	553	2271	219	796	1086	2130	5128
21	227	1512	589	4757	2646	686	1014	679	810	3644	7949	1607	27334	1724
22	1112	1126	490	416	710	4381	211	7893	1368	321	22236	1400	16820	1317
23	1830	440	340	1205	1098	597	6426	911	1790	175	2675	1938	32273	2846
24	1121	571	633	826	2727	634	3235	2964	1530	1253	10181	487	22784	1384
25	921	1376	1172	4051	1468	1811	1562	371	4497	2381	2960	3447	141	458
26	2138	1502	154	4796	3993	1605	10609	156	2045	2131	5905	2755	558	839
27	634	413	323	1928	1560	768	670	175	990	4592	12412	1043	10296	
28	118	127	1134	1078	6956	687	5644	480	2284	2993	1748	370		
29	519	884	793	1290	1075	1204	4651	663	346	422	4834	4757		
30	1167	194	995	566	1807	519	240	8417	1629	1100	5901	262		
31	688	1792		575	3608		1358	6132	1072	1528	5626	466		
32	460	1625		1589	453		2035	111	5150	4204	1066	1872		
33	365	900		2779	6987		1284	416	7472	701	134	3888		
34	666	257		3514	1546		9153	4337	479	2745	4463	3745		
35	108	129		5846	7895		161	1744	317	1528	117	4174		
36	2889	183		5320	845		4359	2431	844	6045	2955	1884		
37	2551	379		196	5664		6786	440	2917	957	8285	1671		
38	510	900		1777	4104		8748	3703	2521	595	196	3873		
39	509	910		926	844		5470	4059	3535	6623	5821	1554		
40	909	115		2147	3121		208	2320	861	4446		968		
41	513	260		3317	6226		3263		485					
42	2092	600		608	1280		1238		2300					
43	251	715		6986	877		1517		4515					
44	199	664		209	1552		629		1418					
45	2411	604		955	3575		2902		720					
46	813	238		261	140		2728		1749					
47	777	1167		3399	5357		667		3268					
48	494	3235		4399	1890		1279		198					
Avg.	848.6			2722.8			2994.0		2119.6		9402.4	2082.7	13904.6	2868.3
# of Periods	128			128			89		89		39	40	27	26

Continued.

	1-Naphthyl Ethanol		Aspirin		2-Naphthyl Ethanol	
	Bound	Free	Bound	Free	Bound	Free
1	19239	890	29988	737	388957	1465
2	138	2805	3198	593	103585	203
3	77015	5636	9970	2614	105790	
4	167	1597	11443	400		
5	280005	996	94979	1252		
6	88794	4855	13411	436		
7	12888	2118	44896	672		
8	102857		46931	521		
9			4055	320		
10			112077	1570		
11			626	661		
12			31966	1974		
13			40488	1767		
14			64727	535		
15			14034	1247		
16			1007	1287		
17			426	717		
18			58475			
Avg.	17360.1	1184.2	22624.7	1831.1	23275.2	949.3
# of Periods	8	7	18	17	3	2

Table SI 11. The bound and free period lengths of the complexes in q4MD-CD. All lengths of periods are in 10 ps.

16. Comparison of ΔG_{Comp1} and ΔG_{Comp2}

Figure SI 12. The correlations between ΔH, $\Delta H_{Bound-unbound}$ and experimental values. $\Delta H_{Bound-unbound}$ was computed by using average potential energy of bound and unbound conformations in the trajectories of the complexes. The correlations are label correspondingly. Error bars indicate the uncertainties.

References

(1)   Henchman, R. H., *J. Chem. Phys.* **2007,** *126* (6).

(2)   Irudayam, S. J.; Henchman, R. H., *J. Phys. Condens. Matter* **2010,** *22* (28).

(3)   Gerogiokas, G.; Calabro, G.; Henchman, R. H.; Southey, M. W. Y.; Law, R. J.; Michel, J., *J. Chem. Theory Comput.* **2014,** *10* (1), 35-48.

(4)   Pauling, L., *J. Am. Chem. Soc.* **1935,** *57*, 2680-2684.

(5)   Flyvbjerg, H.; Petersen, H. G., *J. Chem. Phys.* **1989,** *91* (1), 461-466.

(6)   Rekharsky, M. V.; Inoue, Y., *Chem. Rev.* **1998,** *98* (5), 1875-1917.