The landscape of T cell epitope immunogenicity in sequence space

Authors: Masato Ogishi[1]* and Hiroshi Yotsuyanagi[1]

Affiliations:

[1]Division of Infectious Diseases and Applied Immunology, The Institute of Medical Sciences

Research Hospital, The University of Tokyo, Tokyo, 108-0071, Japan.

*Correspondence to: Masato Ogishi (oogishi-tky@umin.ac.jp)

Summary: The existence of population-wide T cell immunity is widely recognized for multiple

pathogen-derived immunodominant epitopes, despite the vast diversity and individualized nature

of T cell receptor (TCR) repertoire. We thus hypothesized that population-wide epitope

immunogenicity could be probabilistically defined by exploiting public TCR features. To gain a

proof-of-concept, here we describe a machine learning framework yielding probabilistic

estimates of immunogenicity, termed "immunogenicity scores," by utilizing features designed to

mimic thermodynamic interactions between peptides bound to major histocompatibility complex

(MHC) and TCR repertoire. Immunogenicity score dynamics among observed and

computationally simulated single amino acid mutants delineated the landscape of position- and

residue-specific mutational impacts, and even quantitatively estimated escaping potentials of

known epitopes with remarkable positional specificity. This study illustrates that the population-

wide aspect of adaptive immunity is predictable via non-individualized approach, possibly

indicating antigen-guided convergence of human T cell reactivity.

Introduction

T cell epitopes bound to major histocompatibility complex [MHC; also called the human leukocyte antigen (HLA) in humans] molecules activate T cells to initiate subsequent immunological orchestration (Garcia et al., 1999; Grakoui et al., 1999; Rudolph et al., 2006). MHC class I (MHC-I) molecules typically present 8- to 11-aa peptides generated through proteasomal cleavage of intracellular proteins to activate CD8[+] cytotoxic T lymphocytes (CTLs), whereas MHC class II (MHC-II) molecules with an open-ended binding groove accommodate peptides of more variable length derived from endocytosed proteins to activate CD4[+] T helper (Th) cells (Neefjes et al., 2011). Evidence suggests that not all peptides presented on MHC molecules are immunogenic, *i.e.*, trigger functional T cell activation (Bjerregaard et al., 2017; Newell et al., 2013; Sette et al., 1994). T cells recognize peptide-MHC (pMHC) complexes by their TCRs, most predominantly via complementarity determining region 3 (CDR3) loops (Borg et al., 2005; Tian et al., 2007; Zhong et al., 2013). TCR CDR3 sequences are exceedingly diversified (known as TCR repertoire) through gene segment rearrangement and in rare circumstances through somatic hypermutation (Cheynier et al., 1998). However, contrary to the general notion that TCR repertoire is highly stochastic and individualized, accumulating evidence suggests that the biased generation of TCR repertoire due to non-random rearrangement and positive/negative thymic selections is thought to result in an extensive utilization of a limited subset of possible sequences and larger-than-expected overlaps of repertoires between individuals knowns as public clonotypes (Britanova et al., 2014; Madi et al., 2014; Ndifon et al., 2012; Robins et al., 2010; Shugay et al., 2013). Moreover, even distinct TCR repertoires can recognize identical immunodominant epitopes (Chen et al., 2017). Finally, population-wide immunity to some pathogen-derived epitopes has been proven and even utilized clinically,

exemplified by the interferon-gamma release assay for tuberculosis (Horvati et al., 2016;

Udhayakumar et al., 1995). These observations led us to the hypothesis that population-wide

epitope immunogenicity could be probabilistically defined, and public TCR usage pattern might

serve as a probe. In line with this hypothesis, residue hydrophobicity at TCR contact residues has

been associated with immunogenicity, although the predictive capability was not satisfactory

(Chowell et al., 2015). In this context, here we propose that a computational framework

mimicking the intermolecular interactions between pMHC complexes and public human TCR

clonotypes, termed TCR-peptide contact potential profiling (CPP), generates probabilistic

estimates of immunogenicity, termed immunogenicity scores, which effectively recapitulate

essential characteristics of T cell immunity and enable quantitative assessment of potential

mutational impacts on the dynamics of immunogenicity, *i.e.*, escaping, in a position-specific

context.

Result

Concepts and analysis workflow

To gain a proof-of-concept, we first compiled epitope datasets comprising 21,162 8- to 11-aa HLA-I-restricted and 31,693 11- to 30-aa HLA-II-restricted peptide sequences from various sources (Figure 1, Figure S1 and Data S1) (Calis et al., 2013; Chowell et al., 2015; Fleri et al., 2017; Kuiken et al., 2005; Lata et al., 2009; Llano et al., 2013; Olsen et al., 2017; Reche et al., 2005; Tung et al., 2011). We defined positivity as the existence of at least one positive functional T cell assay result. A total of 6,957 and 16,642 HLA-I and HLA-II-restricted epitopes met the criteria. Next, we extracted 23,006,555 unique TCR β chain CDR3 (CDR3β) sequences from TCR repertoire datasets derived from healthy individuals using MiXCR software (Bolotin et al., 2015). We obtained 191,326 unique CDR3β sequences identified in at least 22 out of 206 different datasets to construct fragment libraries for the CPP feature computation.

The concept of CPP relies on the hypothesis that public TCR repertoire is biased toward immunodominant epitopes through evolutional adaptation and thymic negative selection, and therefore could be used to probe peptides with the highest immunogenic potentials (Figure 2A) (Blackman et al., 1990; Madi et al., 2017; Miles et al., 2011; Robins et al., 2010). However, because of the sequence-level and structural diversity of TCR and pMHC, it is unfeasible to comprehensibly characterize every single TCR-pMHC interaction experimentally. To circumvent this problem, we instead designed the CPP framework with the following simplifications. First, we focused on CDR3β loop because this region has the highest genetic variability and is primarily responsible for the interactions with peptides presented onto the MHC grooves, whereas more conserved CDR1 and CDR2 loops typically interact with MHC α-

helices (Bridgeman et al., 2012; Cole et al., 2014; Garcia et al., 1999; Rudolph et al., 2006).

Second, we assumed that initial intermolecular interaction requires only a small portion of the

CDR3β loop that could be approximated to linear structure, based on the low affinities ($K_D = 0.1$

~ >500 μM) compared to other immunoglobulin-like molecules including antibodies, the

relatively flat interfaces, and the substantial conformational changes induced upon recognition of

pMHC complexes (Boniface et al., 1999; Garcia et al., 1998; Hennecke and Wiley, 2001; Lee et

al., 2004). Indeed, analysis of known TCR-pMHC complex structures using PRODIGY

(Vangone and Bonvin, 2015) suggested that around 50% and more than 80% of the

intermolecular contacts are limited within the 3-aa and 5-aa ranges, respectively, in both peptides

and CDR3β regions (Figure 2C and Data S2). Finally, we approximated the "best-match"

problem between peptides and TCR repertoire to pairwise sequence alignment problem, where

the alignment scores are interpreted to reflect the contact energy potentials (Figure 2D).

TCR contact features essential for epitope immunogenicity prediction

We generated CPP features using public clonotype sequences, as well as other features

based on peptide physicochemical properties (Figure 1). Feature selection yielded the 32 and 27

most predictive features for MHC-I and MHC-II predictions, respectively (Figure S2 and Data

S3). Only nine of 32 and none of 27 features were derived from peptide physicochemical

descriptors, highlighting the indispensable contributions of CPP features on immunogenicity

prediction. Common parameter usage patterns for CPP features were observed in MHC-I and

MHC-II (Figure 3A-C). Notably, features derived from short fragments (*i.e.*, 3-aa and 4-aa) and

the longest fragment (*i.e.*, 8-aa and 11-aa in MHC-I and MHC-II, respectively) had higher

cumulative importance values (Figure 3A). Meanwhile, skewness- and kurtosis-derived features

showed markedly higher importance, indicating that distinct repertoire-wide contact potential

distributions are the hallmark of immunogenicity (Figure 3B). The inverted version of

BETM990101 (termed "BETM990101inv") (Betancourt and Thirumalai, 1999) and other six

AAIndex scales of the highest importance for both MHC-I and MHC-II (Figure 3C). To test the

hypothesis that selected AAIndex scales may reflect thermodynamic aspects of TCR-peptide

interactions, we analyzed 82 TCR-peptide complexes with experimentally determined affinities

collected from literature (Data S2) through a modified framework which we termed single-TCR

contact potential profiling (sCPP). We found that BETM990101inv and other six AAIndex

scales important for immunogenicity prediction also generated sCPP features correlating with

affinities (Figure 3D and E and Table S1). Sequence permutations significantly undermined the

observed correlations (Figure 3F). Collectively, these observations support the notion that our

CPP framework recapitulates the essential thermodynamic properties of TCR recognition of

pMHC complex which governs both MHC-I and MHC-II systems.

Probabilistic estimation of epitope immunogenicity

We applied machine learning techniques to convert these most predictive features into a

unidimensional scale of population-wide immunogenicity by averaging the probability estimates

from five times repeated five-fold cross-validations (CVs). We termed this scale as

"immunogenicity scores" (Data S4). The estimates were reasonably consistent across CVs; the

normalized standard deviations distributed with medians and interquartile ranges (IQRs) of 8.3%

(2.3% to 14.3%) and 5.0% (1.4% to 8.5%) in MHC-I and MHC-II, respectively. Addition of

MHC binding prediction results during machine learning only marginally improved the

predictive performance for MHC-I and had almost no effect for MHC-II. Both reduction of

features utilized during machine learning and removing peptides with high homology from the

dataset resulted in only a moderate decrease in predictive performance (Figure 4A and Figure

S3). Contrastingly, the MHC-I immunogenicity prediction tool (Calis et al., 2013) showed only

marginal predictive power on our dataset, with the area under the ROC curve (AUC) of 54%

(Figure S4), and there remains no existing tool for MHC-II to compare. Immunogenicity scores

stably differentiated epitopes from non-immunogenic binders across various subsets of the

datasets, including data obtained from autologous *ex vivo* assays (Figure 4B). Extrapolation of

human immunogenicity classifiers to non-human peptide datasets yielded estimates with slightly

lower predictive performances (Figure 4C and Figure S5), although the dearth of data prevents

us from drawing a solid conclusion on MHC-II-restricted primate epitopes.

We next asked the biological implications of the immunogenicity score framework. First,

thymically expressed self-epitopes identified previously (Adamopoulou et al., 2013) were more

likely annotated as non-immunogenic (Figure 4D). Moreover, even peptides annotated as

immunogenic had a score distribution lower than non-thymus immunogenic and comparable

with thymus non-immunogenic counterparts (Figure 4E). Immunogenic peptides from various

organisms generally had higher scores than non-immunogenic counterparts, with some

exceptions such as MHC-I-restricted peptides derived from human and *Mycobacterium*

tuberculosis. Meanwhile, immunogenicity scores were particularly well-suited for HIV and HCV

epitopes (Figures 4F and G), and this was also the case in the epitopes from the Los Alamos HIV

and HCV immune epitope databases (Figure 4H). Similarly, MHC-I-restricted neoepitopes

obtained from TANTIGEN database also exhibited higher scores than non-immunogenic

peptides from various origins (Figure 4I). Tumor-associated antigens yielding peptides of high

predicted immunogenicity included CTG1B and MAGB1 (Figure 4J).

The landscape of immunogenicity dynamics by single amino acid mutations

Single amino acid mutations can affect peptide immunogenicity bidirectionally, namely,

acquisition and loss of immunogenicity, in sequence space (Figure 5A). We termed the

discordance of annotated immunogenicity between neighbors, *i.e.*, single-aa mutants, as

"immune transition," and defined peptides with evidence of immune transition as "transitional."

Note that we cannot define "non-transitional" peptides from our datasets as the lack of evidence

of transition may simply result from the lack of assays. We identified 1,360 and 976 transitional

peptides for MHC-I and MHC-II, respectively. We noted that immunogenicity scores were less

applicable to transitional peptides (Figures 5B and C, and Figure S6), which may result from

suboptimal machine learning due to the dearth of transitional peptide data. We next asked

whether the integration of neighbor information improves classification. To test this, we

constructed neighbor networks for transitional peptides using all possible single-aa mutants

(N=232,723 and 292,437 for MHC-I and MHC-II, respectively) simulated computationally, the

process we termed "*in silico* mutagenesis." As expected, the averaged immunogenicity scores

classified the immunogenicity of transitional peptides more accurately (Figures 5B and C). To

further characterize the position- and residue-specific mutational impacts, we next focused on the

immunogenicity dynamics between neighbor pairs. We screened our epitope datasets to identify

6,179 and 3,076 MHC-I and MHC-II-restricted single-aa mutated peptide pairs. Immunogenicity

scores changed more dynamically in transitional pairs (Figure 5D). This observation was the case

even when focusing on neighbor pairs of common organism origins (Figure 5E). We also noted

that score changes between MHC-I neighbor pairs with mutations in their anchor residues were

significantly more eminent (Figure 5F). This is interesting as our framework does not utilize any

positionally defined features. To gain insights into the residue-specific mutational impacts, we

analyzed 2,580,890 and 3,101,092 neighbor pairs from all simulated single-aa mutants of

transitional MHC-I- and MHC-II-restricted peptides, respectively. We utilized simulated peptide

data because observed neighbor pairs were too sparsely distributed in terms of residue

combinations. Heatmap clustering analysis revealed residue-specific impacts on immunogenicity

that are partially interpretable with known physicochemical properties [*e.g.*,

hydrophobic/aliphatic (V, L, and I), negative (D and E), and aromatic (F and W)]. However,

some exceptions were also notable [*e.g.*, positive (K and R) in MHC-I]. Collectively, our

systematic characterization delineates the landscape of bidirectional effects of single-aa

mutations on immunogenicity in a position- and residue-dependent context.

Inference of escaping patterns from neighbor network architecture

Escaping mutations are the principal obstacle for vaccine development. Neither reactivity

of the specific T cell clones tested *in vitro* against target epitopes or immune activation after

administration *in vivo* does not guarantee a sustainable immune response in the real-world setting

because of the possibility of deleterious escaping mutations. We tackled this fundamental

problem by quantifying position-specific "escaping potentials" from neighbor network

architecture. Figure 6A shows two representative neighbor networks constructed from observed

neighbors of immunodominant influenza A virus epitopes GILGFVFTL (GIL) and

SRYWAIRTR (SRY). Apparently, GIL seems more robustly immunogenic than SRY because

the Cluster4 in the SRY network was enriched with non-immunogenic peptides, suggesting a

path for escaping. *In silico* mutagenesis followed by neighbor network clustering analysis

revealed position-specific dynamics of predicted immunogenicity (Figure 6B). P6 and P4 were

indicated to have the most substantial impact on escaping for GIL and SRY, respectively.

Intriguingly, P6 of GIL peptide has been shown to be involved in two hydrogen bonds with two

distinct TCR clones F50 and JM22 (Yang et al., 2017). Furthermore, five P4 mutants of SRY

peptide has been shown to result in an undetectable level of cytolysis by at least three CTL

clones (Bowness et al., 1994). Encouraged by these observations, we next examined two MHC-

II-restricted epitopes. GAGSLQPLALEGSLQKRG (GAG) is a self-epitope derived from insulin.

Our simulation indicated that P8-11 and P15-16 were required for its immunogenicity. Strikingly,

immunodominant epitope LALEGSLQK has been localized previously, showing a complete

match with the prediction (Congia et al., 1998). It is worth mentioning that this LALEGSLQK is

usually degraded proteolytically during the maturation of insulin molecule. As our framework

does not integrate factors affecting intracellular peptide processing and presentation, appropriate

peptide presentation onto MHC should be considered a prerequisite for performing

immunogenicity score-based analysis. PKGQTGEPGIAGFKGEQGPK (PKG) is another self-

epitope derived from collagen. Our simulation indicated that P10, P11, P13, and less evidently,

P14 and P16 were required for its immunogenicity. Indeed, mutations at P10, P13, and P16 have

been shown to almost diminish the T cell reactivity collected from pre-immunized transgenic

mice, and mutations at P14 partially impaired the response (Sakurai et al., 2006). To test if these

observations originate from overfitting to target peptides, we repeated the same analysis using

probabilistic estimates obtained from machine learning after removing both the target peptide

and all of its neighbors. The most prominent positions for every peptide, namely, P6 of GIL, P4

of SRY, P8-P14 of GAG, and P10/P13 of PKG, were reproducibly detected (Figure S7). In the

meantime, some of the positions such as P7 of SRY and P16 of PKG became statistically

insignificant, indicating that, as well as pan-epitope rules governing immunogenicity, epitope-

specific characteristics can also be incorporated when neighbor information is available.

It is of vast pragmatical significance to invent methodology to computationally prioritize

epitopes of high immunogenic potential and low escaping potential. To this end, we defined

"escaping potential" as the maximal reduction of cluster-mean immunogenicity score in the

neighbor network. Immunogenicity score-escaping potential (IS-EP) plots indicated that,

although there was a moderate correlation between IS and EP, considerable variation also existed

(Figures 7A and B). Of note, HIV and dengue virus (DGV)-derived epitopes had higher EP than

human epitopes, and this trend was not observed in non-immunogenic peptides (Figure 7C). Two

representative HIV-derived MHC-I epitopes GGKKKYKL and ITTESIVIW were shown in

Figure 7D. As indicated in our analysis, there are known naturally occurring escaping mutations

at P3, P5, and P7 of GGKKKYKL (Reid et al., 1996). Although we could not find reliable

information on escaping variants of ITTESIVIW, this epitope represents the most common

sequence found in the circulating HIV-1 clade B population in the Los Alamos HIV Sequence

Database (Du et al., 2016). Likewise, two representative MHC-II epitopes derived from *P.*

falciparum and *M. tuberculosis* were shown in Figures 7E and F. Indeed, the prevalence of T cell

reactivity to the low-EP peptide GKLLSTGLVQNFPNTIISK in Kenyan volunteers was reported

to be 25% (Udhayakumar et al., 1995), and, although not precisely the same, 29.4% of South

African patients with either latent or active tuberculosis infection were shown to be positive in

the ELISpot assay against the epitope VRAVAESHGVAAVLFAATAA (Horvati et al., 2016).

Collectively, these observations indicate that our framework enables prioritization of promising

epitopes based on their high immunogenic and low escaping potentials *in silico*.

Discussions

This study provides a proof-of-concept that population-wide aspect of T cell immunity is

predictable through probabilistic approach using features recapitulating physiological contacts

between MHC-presented peptides and TCR repertoire. It is of note that short (*i.e.*, 3-aa and 4-aa)

TCR fragments in combination with a restricted set of AAIndex scales generate features

contributing to both MHC-I and MHC-II immunogenicity prediction and also correlating with

individual TCR-pMHC affinities, because these commonly utilized parameters might represent

MHC-independent pan-epitope principles governing TCR-pMHC interactions. The utilization of

short fragment features is consistent with the induced fit model that only a limited

complimentary region of TCR is involved at least in initial engagement (Boniface et al., 1999;

Garcia et al., 1998; Lee et al., 2004). Meanwhile, the best AAIndex scale BETM990101 reflects

interaction energies between amino acid residues under physiological conditions (Betancourt and

Thirumalai, 1999), and hence the inverted version of BETM990101, giving larger values for

stronger interresidue contacts, is a reasonable scale for contact potential estimation. Indeed, a

positive skewness in the contact potential distribution, implying the coexistence of a small and a

large proportion of TCR fragments with high and low contact potentials, respectively, was

associated with stronger affinities in our analysis. TCR affinity has been correlated with T cell

activation (Sykulev et al., 1994), although there remains a debate as to whether other parameters

such as avidity, on- and off-rates are also essential (Stone et al., 2009). Further studies are

required to fully elucidate the biological relevance of simulated TCR-peptide contact features

quintessential for immunogenicity prediction to the thermodynamic parameters of TCR-pMHC

interactions and subsequent T cell activation.

Our findings bolster the hypothesis that population-level epitope immunogenicity could

indeed be probabilistically determined. Indeed, the concept of immunodominance bases this

hypothesis (Yewdell, 2006). For instance, only 0.03% of potential MHC-I-presented peptides

derived from vaccinia virus account for >90% of the CD8[+] T cell response in B6 mice (Moutaftsi

et al., 2006). Similarly, diverse TCR repertoires specific for two immunodominant viral epitopes

have been reported in human (Chen et al., 2017). Population-level immunity against various

immunodominant epitopes is another line of evidence (Horvati et al., 2016; Udhayakumar et al.,

1995). The fact that diverse TCRs can be utilized to achieve protective immunity against

immunodominant epitopes in multiple individuals indicates some underlying qualities of

epitopes guiding both genomic and thymic "convergent evolution" of human TCR repertoire. To

elucidate the hitherto unknown pan-epitope principles governing productive TCR-peptide

contacts, we took an inductive approach using a large set of *in vitro* T cell epitope assay results

which may vary depending on the specific T cell clones or TCR repertoires utilized. Our findings

indicate that a limited set of TCR-peptide contact features reproducibly serve as principal

determinants of T cell reactivity. Moreover, we show that the averaged probabilistic estimates,

termed immunogenicity scores, are consistently higher in epitopes with positive results from

various functional assays. Interestingly, peptides physiologically presented on the thymus have

lower immunogenicity scores than non-thymus counterparts, which is consistent with the biased

formation of human TCR repertoire owing to thymic negative selection (Adamopoulou et al.,

2013; Blackman et al., 1990). Practically speaking, the probabilistic scoring system would be

particularly useful for prioritizing peptide candidates to expedite the development of population-

level immunotherapies for various infectious diseases and cancer. However, it is notable that

immunogenicity scores do not distinguish immunogenic and non-immunogenic peptides from *M.*

tuberculosis. It has been shown that T cell epitopes of TB are evolutionary hyperconserved and

the bacteria seems to actually benefit from recognition by human T cells (Comas et al., 2010;

Coscolla et al., 2015). Further characterization of T cell reactivity for TB epitopes with high and

low immunogenicity scores may unveil critical insights into the uniqueness of anti-TB T cell

immunity.

287   One remarkable feature of our immunogenicity score system is that the score dynamics

between single-aa mutants are, to some extent, predictive of the mutational impacts on

immunogenicity. Moreover, the newly introduce metric termed "escaping potential" allows

researchers to quantitatively assess the "robustness" of immunogenicity for any epitope.

Practically, it could enable *in silico* epitope screening on the basis of their immunogenicity

scores and escaping potentials for developing off-the-shelf targeted immunotherapies available to

the general public. In the meantime, it could provide mechanistic insights into the co-

evolutionary trajectory of pathogens and general human ancestry. A caveat is that our current

model may be suboptimal because only a small fraction of possible sequence space has been

experimentally covered so far. Ideally, multiple assays using T cell clones of different

individuals should be performed for any peptide to evaluate inter-individual variations.

Nonetheless, it is encouraging that our current framework still infers escaping even without using

any of the peptide in question and its single-aa mutants during machine learning. Further

elucidation of the pan-epitope basis of population-level human T cell immunity has significant

clinical and biological implications.

Materials and Methods

Experimental design

The purpose of this retrospective study was to identify the essential biochemically relevant features predictive of T cell epitope immunogenicity and to develop a linear coordinate system reflecting the probabilities that those peptides are immunogenic. To this end, public MHC-I and MHC-II-restricted peptide sequence datasets were analyzed through a newly designed computational framework that mimics the molecular scanning of MHC-loaded peptides by human public TCR clonotypes. All analysis is exploratory; no predetermined protocols were applied before initiating the study. All data inclusion and exclusion criteria were described in the following sections. Accounting for missing data values including imputation is not applicable.

Computational analysis

All computational analyses were conducted using R ver. 3.5.0 (https://www.r-project.org/) (R Core Team, 2018). The latest versions of R packages were consistently used. Compiled datasets and essential in-house functions are available as the R package *Repitope* on GitHub (https://github.com/masato-ogishi/Repitope). Other scripts are available upon request.

Peptide sequence datasets

HLA-I-restricted peptide sequences of 8-aa to 11-aa lengths with T cell assay results were collected from public databases [Immune Epitope Database (IEDB, as of May 7th, 2018) (Fleri et al., 2017), the best-characterized CTL epitopes from Los Alamos National Laboratory (LANL) HIV Sequence Database (Llano et al., 2013), LANL HCV Sequence Database (Kuiken et al., 2005), EPIMHC (Reche et al., 2005), MHCBN (Lata et al., 2009), and TANTIGEN (Olsen et al., 2017)] and previous publications (Calis et al., 2013; Chowell et al., 2015; Tung et al., 2011). HLA-II-restricted peptide sequences of 11-aa to 30-aa lengths with T cell assay results were collected from the IEDB database (as of May 7th, 2018). As for peptides retrieved from the IEDB database, only those with evidence of functional T cell response were included. The list of evidence included is as follows: activation, antibody help, CCL2/MCP-1 release, CCL3/MIP-1a release, CCL4/MIP-1b release, CCL5/RANTES release, CXCL10/IP-10 release, CXCL9/MIG release, cytotoxicity, decreased disease, degranulation, disease exacerbation, GM-CSF release, granulysin release, granzyme A release, granzyme B release, IFNg release, IL-10 release, IL-12 release, IL-13 release, IL-17 release, IL-17A release, IL-1b release, IL-2 release, IL-21 release, IL-22 release, IL-23 release, IL-3 release, IL-4 release, IL-5 release, IL-6 release, IL-8 release, lymphotoxin A/TNFb release, pathogen burden after challenge, perforin release, proliferation, protection from challenge, survival from challenge, T cell- APC binding, T cell help, TGFb release, TNF release, TNFa release, tolerance, tumor burden after challenge, and type IV hypersensitivity (DTH). Peptides presented on non-human MHC molecules were discarded, whereas those presented on HLA molecules in non-human hosts (*e.g.*, transgenic mice) were included. In this manner, 21,162 and 31,693 HLA-I and HLA-II-restricted peptides were identified, respectively, of which 1,873 (8.9%) and 4,505 (14.2%) had contradicting annotations on immunogenicity. In such cases, peptides with at least one positive T cell assay result were considered immunogenic to maximize the breadth of population-wide immunity included in the analysis. Eventually, 6,957 (32.9%) and 16,642 (52.5%) HLA-I and HLA-II-restricted peptides were considered immunogenic, respectively. Collected peptide sequences and annotations are summarized in Data S1.

Sequences of peptides restricted on non-human MHC molecules were collected from the
IEDB database (as of as of May 7th, 2018) as described above. The following species were
considered primate: bonobo, chimpanzee, gorilla, marmoset, and rhesus macaque. Meanwhile,
mouse and rat were considered rodent. Eventually, 411 and 8,756 MHC-I-restricted peptides, and
76 and 8,445 MHC-II-restricted peptides were identified for primates and rodents, respectively.
Identified epitopes and annotated information can be found in Data S1.
355
TCR sequence datasets
TCR repertoire datasets were collected from NCBI Sequence Read Archive (SRA) and a
previous study led by Britanova *et al.* (Britanova et al., 2014). Repertoire datasets derived from
healthy donors were searched through SRA, and the following BioProjects were included:
PRJNA389805, PRJNA329041, PRJNA273698, PRJNA258001, PRJNA229070, PRJNA79707,
and PRJNA79435. Also, pooled sequence data from 39 healthy donors were retrieved from the
paper by Britanova *et al*. In this manner, a total of 206 datasets, of which only five were pooled
datasets, were collected. Pooled and individual datasets were derived from a total of 561 and 103
healthy donors, respectively. Fastq files were obtained using fastq-dump script with the
following options: --gzip --skip-technical --readids --read-filter pass --dumpbase --split-files --
clip --accession [SRA run number]. A total of 23,006,555 CDR3β sequences were extracted
using MiXCR software (Bolotin et al., 2015).
Public TCR clonotypes, the ones commonly observed in multiple datasets, were
identified from the pool of CDR3β sequences mentioned above. Britanova *et al.* (Britanova et
al., 2014) reported that 10,691 clonotypes were shared between at least six of 39 (15.4%) donor-
derived top 100,000 clonotype sets. Based on their findings, we decided to extract CDR3β
sequences identified in at least 22 out of 206 (10.8%) different datasets, and we subsequently
identified 191,326 (0.83%) public clonotypes. A randomly sampled 10,000 clonotypes were used
for TCR-peptide contact potential profiling.
375
TCR-peptide contact potential profiling (CPP)
T cell epitopes presented on the MHC molecules must be recognized by the TCRs of
CD8+ CTLs and CD4+ Th cells with sufficient affinity to trigger subsequent immunological
cascades. Undoubtedly, the vast majority of TCR repertoire is not involved in recognition of any
given peptide. Moreover, considering the substantial conformational flexibility observed upon
binding of TCRs to pMHC complexes, one can assume that only a subset of residues of the
peptide and the epitope-recognition domain, namely, the CDR3β loop, serve as a "seed" of
intermolecular docking during the early phase of molecular scanning (Boniface et al., 1999;
Bridgeman et al., 2012). With these in mind, identifying the best contact site between a given
pair of peptide and CDR3β sequences is conceptually similar to solving a local pairwise
sequence alignment problem. One caution is that higher scores must be given to more strongly
interacting residue pairs instead of more biochemically similar residue pairs as opposed to
ordinary alignments. For this purpose, amino acid pairwise contact potential (AACP) scales from
the AAIndex database (Kawashima et al., 2008)
(http://www.genome.jp/aaindex/AAindex/list_of_potentials) were adopted to generate custom
substitution matrices. As stronger interresidue interactions yield smaller free energy values
(negative numbers), and as the Smith-Waterman local alignment algorithm attempts to maximize
the alignment score of a given sequence (TCR fragment) against the whole target sequence
(peptide), the optimal pairwise alignment using a custom substitution matrix derived from the

sign-inverted version of a pairwise contact potential scale would in principle correspond to the best intermolecular contact. For comparative purposes, both inverted and non-inverted versions were tested. Values were rescaled to a range from zero to one for subsequent analyses. A set of non-inverted, non-rescaled AACP scales is provided as Data S6. For repertoire-wide CPP analysis, the 191,326 pooled public CDR3β sequences were randomly down-sampled to 10,000 sequences with the relative abundance ratios retained. The sequences were then fragmented by a sliding window strategy to generate a fragment library. Since the interacting orientations (either forward parallel or antiparallel) are unknown for most of the cases, reversed CDR3β sequences were also fragmented and combined. The sizes of fragments were from 3-aa to 8-aa and from 3-aa to 11-aa for MHC-I and MHC-II predictions, respectively. For single-TCR CPP (sCPP) analysis, every single TCR instead of public TCR repertoire was used to generate a fragment library. To perform sequence alignments, the *pairwiseAlignment* function implemented in the *Biostrings* package in *Bioconductor* (https://www.bioconductor.org/) (Gentleman et al., 2004) was utilized. This function seeks an optimal alignment which maximizes the overall alignment score defined as a sum of pairwise scores. Alignment type was set "global-local" to obtain an optimal alignment of a given set of TCR fragments against the consecutive subsequences of any given peptide. Gaps were not allowed. A set of alignment scores was summarized by calculating representative statistics. Following statistics were calculated using the functions implemented in the *psych* package: mean (Mean), standard deviation (SD), standard error of the mean (SE), median (Med), median absolute deviation (MAD), interquartile range (IQR), 10% quantile (Q10), 90% quantile (Q90), skewness (Skew), and kurtosis (Kurt). Predictive features were generated by combining the fragment length, the AACP scale, and the type of statistics. In this manner, 700 CPP features per one fragment length were generated for each of the peptides.

Peptide descriptors

Apart from CPP features, sequence-based estimates of physicochemical properties were also calculated. Each peptide sequence was converted into a set of consecutive fragments of a defined amino acid length, and peptide descriptors were calculated against each of the fragments using functions in the *Peptide* package. Following functions were utilized: *aIndex*, *blosumIndices*, *boman*, *charge*, *crucianProperties*, *fasgaiVectors*, *hmoment*, *hydrophobicity*, *instaIndex*, *kideraFactors*, *mswhimScores*, *pI*, *protFP*, *vhseScales*, and, *zScales*. The distributions of the values were summarized similarly to CPP features. Additionally, 20 categorical features indicating whether the peptide of interest is free from a specific amino acid residue were included. The peptide length was also included as a feature.

Feature selection

Preprocessing followed by importance-based feature selection was repeated ten times with different random seeds. The analysis workflow consisted of the following steps. First, the peptide set was randomly split in a ratio of 4:1 for the training and testing subdatasets. Second, using the training subdataset, a preprocessing function was defined; numerical features were centered and scaled; a peptide length and categorical features representing existence or absence of each residue remained unchanged. This preprocessing function was also applied to the testing subdataset. Third, highly correlated features were removed, with the threshold of correlation coefficient being 0.75. Fourth, features were filtered based on the importance values calculated using the *generateFilterValuesData* function with the random forest (RF) method (*randomForestSRC.rfsrc*) implemented in the *mlr* package (Bischl et al., 2016). The 100 most

important features were retained unless otherwise stated. In the first five repeats, the names of the peptide descriptors and the CPP features were ordered in ascending order. In the latter five repeats, on the other hand, the names of those features were ordered in descending order. In this manner, the inherent bias of feature selection owing to the lexicographical order of the names of features was avoided. Finally, features kept in all of the ten repeats were defined as the most predictive features and were utilized hereafter.

TCR-pMHC structure analysis

The structures of TCR-pMHC complexes with known experimentally determined affinities were collected manually from the literature using the ATLAS database (https://zlab.umassmed.edu/atlas/web/) (Borrman et al., 2017). A total of 95 structures were retrieved, of which 82 contained non-mutated (wildtype) TCRs. Contact sites were identified using the PRODIGY server (Vangone and Bonvin, 2015; Xue et al., 2016). TCR contact footprints were defined as the distributions of the numbers of contacts at each of the peptide positions.

For the analysis of correlation with affinities, sCPP features were computed as described in the above sections, and the single best feature was selected from a univariate analysis for each of the AAIndex scales. Multivariate regression with stepwise feature selection was conducted using the *stepAIC* function implemented in the *MASS* package. The variance inflation factors (VIFs) were computed using the *vif* function implemented in the *car* package.

Sequence permutation experiments were performed as follows. Peptide sequences were replaced with a set of random sequences of the same lengths. In contrast, because of the fragmentation strategy employed, a simple replacement would lead to incomplete disruption of hidden but essential motifs. Therefore, we instead replaced the TCR-derived fragment library with a randomly chosen set of random fragments that do not overlap with the original fragment library. Permutation experiments were iterated 1000 times.

HLA binding prediction

Predicted binding strength against twelve representative HLA-I alleles (A*01:01, A*02:01, A*03:01, A*24:02, A*26:01, B*07:02, B*08:01, B*27:05, B*39:01, B*40:01, B*58:01, and B*15:01) from NetMHC 4.0 (Andreatta and Nielsen, 2016) and six representative HLA-II alleles (DRB1*0101, DRB3*0101, DRB4*0101, DRB5*0101, DPA1*0103-DPB1*0101, and DQA1*0101-DQB1*0201) from NetMHCIIpan 3.2 (Jensen et al., 2018) for MHC-I and MHC-II immunogenicity prediction, respectively, were incorporated in the machine learning process for immunogenicity prediction because the stability of the peptide-MHC complex is a known correlate of immunogenicity (Rasmussen et al., 2016). Default parameters were used for prediction. Thresholds of percentile ranks for strong and weak binders were set at 0.5% and 2% in MHC-I, and 2% and 10% in MHC-II, respectively.

Peptide clustering by the IEDB Cluster Analysis Tool

The IEDB Epitope Cluster Analysis Tool (http://tools.iedb.org/cluster/) was utilized to test the effect of sequence-level peptide homology on immunogenicity prediction. Clustering was performed with the homology threshold of 80%. When clustering filter was applied, only a single peptide was randomly chosen from each of the clusters before feature preprocessing in the machine learning workflow.

Immunogenicity score

The most predictive peptide descriptors and CPP features, with or without predicted HLA binding-based features, were compressed into a linear coordinate system through machine learning. We chose extremely randomized trees (ERT) algorithm implemented in the *extraTrees* package (Geurts et al., 2006) from preliminary experiments. ERT is a derivative of RF and is more computationally efficient and at the same time less prone to overfitting. The model-specific parameter *mtry* was set to be 5. Other hyperparameters were set as defaults. Class weights were provided to compensate for imbalanced class distributions. Probabilistic estimates were computed by conducting five times repeated five-fold cross-validations (CVs). This strategy ensures that any peptide is subjected to five times repeated predictions by models trained from a set of peptides that do not contain the peptide in question. The averaged probability estimates from machine learning were termed "immunogenicity scores." Unless otherwise stated, we utilized immunogenicity scores from machine learning without predicted HLA binding-based features. For peptides not included in our epitope dataset, immunogenicity scores are defined as the averaged probability estimates of the 25 ERT models generated during CVs. Prediction variances were evaluated by computing coefficients of variance which are defined as standard deviations divided by means.

Immunogenicity prediction by the IEDB Class I Immunogenicity Tool

Immunogenicity was predicted on the HLA-I-restricted peptide set using the IEDB Class I Immunogenicity Tool (22) (http://tools.iedb.org/immunogenicity/) with default settings.

Neighbor network analysis

Single amino acid mutations, namely, substitutions, insertions, and deletions, have multifaceted effects on the immunogenicity of MHC-presented peptides. To systematically analyze their effects on immunogenicity, we adopted a network-style representation of the epitope data structures, termed a "neighbor network," where pairs of peptide sequences (nodes) with just one edit distance were defined as "neighbors" and regarded as edges. Each edge was directed from the peptide with a lower immunogenicity score to the peptide with a higher score. Edge weight was defined as the ratio of the lower score to the higher score so that less immunogenically similar peptides were mapped more distant from each other in sequence space due to the smaller edge weight. Clustering was performed using a walktrap algorithm implemented in the *igraph* package (Csárdi and Nepusz, 2006). For each cluster, sequences were aligned using the ClustalW algorithm implemented in the *msa* package (Bodenhofer et al., 2015). A consensus sequence was generated using the method implemented in the *Biostrings* package with an ambiguity threshold of 0.5. Gaps were treated equally to ambiguities. Gaps/ambiguities were expressed as "X."

In silico mutagenesis analysis

For a given "parental" peptide, the neighbor network was computationally expanded by simulating all possible single amino acid substituted mutant peptides. Insertions and deletions were not taken into consideration. These artificially introduced mutations could affect various aspects of T cell immunity, including aberrant proteolytic cleavage and altered MHC presentation, these possibilities were not rigorously examined since our primary interest is on exploring the net immunogenicity of mutated peptides if appropriately presented and scanned by

532 TCRs. Immunogenicity score computation and neighbor network analysis were conducted as
533 described.
534
<u>Immune transition</u>
Loss of immunogenicity due to mutation is also called escaping, evasion, and sometimes
immunoediting. In the present study, we refer to this phenomenon as escaping. In contrast, there
is no appropriate terminology for the mutation-driven acquisition of immunogenicity, and thus
we refer to this phenomenon as "epitope/neoepitope formation." For the sake of simplicity, we
do not take into consideration epitope-extrinsic mechanisms such as impaired intracellular
antigen processing, somatic loss of HLA heterozygosity (Chowell et al., 2018; McGranahan et
al., 2017), and checkpoint-mediated T cell exhaustion and apoptosis (Pardoll, 2012) in this work.
Escaping and epitope formation due to mutations can be understood as two sides of the same
coin in sequence space, and therefore, we propose a higher-order concept, "immune transition,"
which is defined as a change in immunogenicity between any MHC-presented peptide and its
single amino acid variant. Note that the synergistic effects of two or more mutations on the
changes in immunogenicity are beyond the scope of the present study.
To further characterize the theoretical boundary of immunogenic and non-immunogenic
peptides, we categorized peptides with evidence of immune transition found in our dataset as
"transitional" and analyzed separately. Note that we cannot certainly define "non-transitional"
peptides, as the lack of immune transition in our dataset may merely reflect the lack of
experimental evaluation. For transitional peptides, *in silico* mutagenesis followed by neighbor
network analysis was conducted. Mean immunogenicity scores of both all single amino acid
mutants and those within the same cluster to the parental peptide were computed.
555
<u>Immunogenicity score dynamics</u>
For quantitative comparison of immunogenicity for a given pair of peptides, either the
ratio of immunogenicity scores (divided by the lower score), termed "relative score change," or
the absolute difference of immunogenicity scores divided by the lower score, termed
"normalized score change," was used as a metric. To systematically explore the position-specific
mutational impacts for a given parental peptide, relative score change was computed for every
pair of computationally simulated mutants. For heatmap analysis, peptide pairs were grouped
based on their mutated residues, and median normalized score changes were calculated for each
group. Heatmaps were visualized using the *ComplexHeatmap* package (Gu et al., 2016).
565
<u>Escaping potential</u>
Escaping potential of a given peptide was defined as the difference between the mean
immunogenicity score of the cluster containing the parental peptide and the minimum mean
immunogenicity score of the other clusters in the neighbor network containing all possible
simulated mutants. A negative escaping potential indicates that the parental peptide resides in the
least immunogenic cluster.
572
<u>Statistical analysis</u>
Non-parametric hypothesis testing methods (Kruskal-Wallis test and Wilcoxon rank sum
test for continuous variables, and Fisher's exact test for categorical variables) were consistently
employed unless otherwise stated. In the case of multiple comparisons, *P*-values were FDR-
adjusted according to the method originally proposed by Benjamini and Hochberg (Benjamini

and Hochberg, 1995). When reporting measures of central tendency and dispersion, data were
presented as medians and interquartile ranges unless otherwise stated. AUC values were
calculated based on ten-fold CVs, and their 95% confidence intervals were inferred from
influence curves using the method implemented in the *cvAUC* package (Ledell et al., 2015). All
statistical analyses were conducted in R (R Core Team, 2018).
583

584   **Author contributions:** M.O. conceived the idea, designed the study, performed computational

585   analyses, and drafted the manuscript; M.O. and H.Y. wrote the manuscript.

586   **Acknowledgments:** We thank Ryo Ueno for helpful discussions. This work is not supported by

587   any external funding sources.

588   **Declaration of interest:** The authors declare no competing interests.

References

Adamopoulou, E., Tenzer, S., Hillen, N., Klug, P., Rota, I.A., Tietz, S., Gebhardt, M., Stevanovic, S., Schild, H., Tolosa, E., et al. (2013). Exploring the MHC-peptide matrix of central tolerance in the human thymus. Nat. Commun. *4*, 2039.

Andreatta, M., and Nielsen, M. (2016). Gapped sequence alignment using artificial neural networks: application to the MHC class I system. Bioinformatics *32*, 511–517.

Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. J. R. Stat. Soc. Ser. B *57*, 289–300.

Betancourt, M.R., and Thirumalai, D. (1999). Pair potentials for protein folding: choice of reference states and sensitivity of predicted native states to variations in the interaction schemes. Protein Sci. *8*, 361–369.

Bischl, B., Lang, M., Kotthoff, L., Schiffner, J., Richter, J., Studerus, E., Casalicchio, G., and Jones, Z.M. (2016). mlr: Machine Learning in R. J. Mach. Learn. Res. *17*, 1–5.

Bjerregaard, A.M., Nielsen, M., Jurtz, V., Barra, C.M., Hadrup, S.R., Szallasi, Z., and Eklund, A.C. (2017). An analysis of natural T cell responses to predicted tumor neoepitopes. Front. Immunol. *8*, 1–9.

Blackman, M., Kappler, J., and Marrack, P. (1990). The role of the T cell receptor in positive and negative selection of developing T cells. Science *248*, 1335–1341.

Bodenhofer, U., Bonatesta, E., Horejš-Kainrath, C., and Hochreiter, S. (2015). msa: an R package for multiple sequence alignment. Bioinformatics *31*, btv494.

Bolotin, D.A., Poslavsky, S., Mitrophanov, I., Shugay, M., Mamedov, I.Z., Putintseva, E. V, and

Chudakov, D.M. (2015). MiXCR: software for comprehensive adaptive immunity profiling. Nat.

Methods *12*, 380–381.

Boniface, J.J., Reich, Z., Lyons, D.S., and Davis, M.M. (1999). Thermodynamics of T cell

receptor binding to peptide-MHC: evidence for a general mechanism of molecular scanning.

Proc. Natl. Acad. Sci. U. S. A. *96*, 11446–11451.

Borg, N.A., Ely, L.K., Beddoe, T., Macdonald, W.A., Reid, H.H., Clements, C.S., Purcell, A.W.,

Kjer-Nielsen, L., Miles, J.J., Burrows, S.R., et al. (2005). The CDR3 regions of an

immunodominant T cell receptor dictate the "energetic landscape" of peptide-MHC recognition.

Nat. Immunol. *6*, 171–180.

Borrman, T., Cimons, J., Cosiano, M., Purcaro, M., Pierce, B.G., Baker, B.M., and Weng, Z.

(2017). ATLAS: A database linking binding affinities with structures for wild-type and mutant

TCR-pMHC complexes. Proteins Struct. Funct. Bioinforma. *85*, 908–916.

Bowness, P., Allen, R.L., and McMichael, A.J. (1994). Identification of T cell receptor

recognition residues for a viral peptide presented by HLA B27. Eur. J. Immunol. *24*, 2357–2363.

Bridgeman, J.S., Sewell, A.K., Miles, J.J., Price, D.A., and Cole, D.K. (2012). Structural and

biophysical determinants of αβ T-cell antigen recognition. Immunology *135*, 9–18.

Britanova, O. V., Putintseva, E. V., Shugay, M., Merzlyak, E.M., Turchaninova, M.A.,

Staroverov, D.B., Bolotin, D.A., Lukyanov, S., Bogdanova, E.A., Mamedov, I.Z., et al. (2014).

Age-Related Decrease in TCR Repertoire Diversity Measured with Deep and Normalized

Sequence Profiling. J. Immunol. *192*, 2689–2698.

Calis, J.J.A., Maybeno, M., Greenbaum, J.A., Weiskopf, D., De Silva, A.D., Sette, A., Keşmir, C., and Peters, B. (2013). Properties of MHC class I presented peptides that enhance immunogenicity. PLoS Comput. Biol. *9*, e1003266.

Chen, G., Yang, X., Ko, A., Sun, X., Gao, M., Zhang, Y., Shi, A., Mariuzza, R.A., and Weng, N. ping (2017). Sequence and Structural Analyses Reveal Distinct and Highly Diverse Human CD8+ TCR Repertoires to Immunodominant Viral Antigens. Cell Rep. *19*, 569–583.

Cheynier, R., Henrichwark, S., and Wain-Hobson, S. (1998). Somatic hypermutation of the T cell receptor Vβ gene in microdissected splenic white pulps from HIV-1-positive patients. Eur. J. Immunol. *28*, 1604–1610.

Chowell, D., Krishna, S., Becker, P.D., Cocita, C., Shu, J., Tan, X., Greenberg, P.D., Klavinskis, L.S., Blattman, J.N., and Anderson, K.S. (2015). TCR contact residue hydrophobicity is a hallmark of immunogenic CD8 + T cell epitopes. Proc. Natl. Acad. Sci. *112*, E1754–E1762.

Chowell, D., Morris, L.G.T.T., Grigg, C.M., Weber, J.K., Samstein, R.M., Makarov, V., Kuo, F., Kendall, S.M., Requena, D., Riaz, N., et al. (2018). Patient HLA class I genotype influences cancer response to checkpoint blockade immunotherapy. Science *359*, 582–587.

Cole, D.K., Miles, K.M., Madura, F., Holland, C.J., Schauenburg, A.J.A.A., Godkin, A.J., Bulek, A.M., Fuller, A., Akpovwa, H.J.E.E., Pymm, P.G., et al. (2014). T-cell Receptor (TCR)-peptide specificity overrides affinity-enhancing TCR-major histocompatibility complex interactions. J. Biol. Chem. *289*, 628–638.

Comas, I., Chakravartti, J., Small, P.M., Galagan, J., Niemann, S., Kremer, K., Ernst, J.D., and Gagneux, S. (2010). Human T cell epitopes of Mycobacterium tuberculosis are evolutionarily hyperconserved. Nat. Genet. *42*, 498–503.

Congia, M., Patel, S., Cope, A.P., De Virgiliis, S., and Sønderstrup, G. (1998). T cell epitopes of

insulin defined in HLA-DR4 transgenic mice are derived from preproinsulin and proinsulin.

Proc. Natl. Acad. Sci. U. S. A. *95*, 3833–3838.

Coscolla, M., Copin, R., Sutherland, J., Gehre, F., De Jong, B., Owolabi, O., Mbayo, G.,

Giardina, F., Ernst, J.D., and Gagneux, S. (2015). M. tuberculosis T cell epitope analysis reveals

paucity of antigenic variation and identifies rare variable TB antigens. Cell Host Microbe *18*,

538–548.

Csárdi, G., and Nepusz, T. (2006). The igraph software package for complex network research.

InterJournal *Complex Sy*, 1695.

Du, V.Y., Bansal, A., Carlson, J., Salazar-Gonzalez, J.F., Salazar, M.G., Ladell, K., Gras, S.,

Josephs, T.M., Heath, S.L., Price, D.A., et al. (2016). HIV-1–Specific CD8 T Cells Exhibit

Limited Cross-Reactivity during Acute Infection. J. Immunol. *196*, 3276–3286.

Fleri, W., Paul, S., Dhanda, S.K., Mahajan, S., Xu, X., Peters, B., and Sette, A. (2017). The

immune epitope database and analysis resource in epitope discovery and synthetic vaccine

design. Front. Immunol. *8*, 1–16.

Garcia, K.C., Degano, M., Pease, L.R., Huang, M., Peterson, P.A., Teyton, L., and Wilson, I.A.

(1998). Structural basis of plasticity in T cell receptor recognition of a self peptide-MHC antigen.

Science *279*, 1166–1172.

Garcia, K.C., Teyton, L., and Wilson, I.A. (1999). Structural basis of T cell recognition. Annu.

Rev. Immunol. *17*, 369–397.

Gentleman, R.C., Carey, V.J., Bates, D.M., Bolstad, B., Dettling, M., Dudoit, S., Ellis, B.,

Gautier, L., Ge, Y., Gentry, J., et al. (2004). Bioconductor: open software development for

computational biology and bioinformatics. Genome Biol. *5*, R80.

Geurts, P., Ernst, D., and Wehenkel, L. (2006). Extremely randomized trees. Mach. Learn. *63*, 3–

42.

Grakoui, A., Bromley, S.K., Sumen, C., Davis, M.M., Shaw, A.S., Allen, P.M., and Dustin, M.L.

(1999). The immunological synapse: a molecular machine controlling T cell activation. Science

285, 221–227.

Gu, Z., Eils, R., and Schlesner, M. (2016). Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. Bioinformatics *32*, 2847–2849.

Hennecke, J., and Wiley, D.C. (2001). T cell receptor-MHC interactions up close. Cell *104*, 1–4.

Horvati, K., Bősze, S., Gideon, H.P., Bacsa, B., Szabó, T.G., Goliath, R., Rangaka, M.X.,

Hudecz, F., Wilkinson, R.J., and Wilkinson, K.A. (2016). Population tailored modification of

tuberculosis specific interferon-gamma release assay. J. Infect. *72*, 179–188.

Jensen, K.K., Andreatta, M., Marcatili, P., Buus, S., Greenbaum, J.A., Yan, Z., Sette, A., Peters,

B., and Nielsen, M. (2018). Improved methods for predicting peptide binding affinity to MHC

class II molecules. Immunology *154*, 394–406.

Kawashima, S., Pokarowski, P., Pokarowska, M., Kolinski, A., Katayama, T., and Kanehisa, M.

(2008). AAindex: Amino acid index database, progress report 2008. Nucleic Acids Res. *36*, 202–

205.

Kuiken, C., Yusim, K., Boykin, L., and Richardson, R. (2005). The Los Alamos hepatitis C

sequence database. Bioinformatics *21*, 379–384.

Lata, S., Bhasin, M., and Raghava, G.P.S. (2009). MHCBN 4.0: A database of MHC/TAP

binding peptides and T-cell epitopes. BMC Res. Notes *2*, 61.

Ledell, E., Petersen, M., and Van Der Laan, M. (2015). Computationally efficient confidence

intervals for cross-validated area under the ROC curve estimates. Electron. J. Stat. *9*, 1583–1607.

Lee, J.K., Stewart-Jones, G., Dong, T., Harlos, K., Di Gleria, K., Dorrell, L., Douek, D.C., van

der Merwe, P.A., Jones, E.Y., and McMichael, A.J. (2004). T cell cross-reactivity and

conformational changes during TCR engagement. J. Exp. Med. *200*, 1455–1466.

Llano, A., Williams, A., Olvera, A., Silva-Arrieta, S., and Brander, C. (2013). Best-

Characterized HIV-1 CTL Epitopes: The 2013 Update. HIV Mol. Immunol. 2013 3–25.

Madi, A., Shifrut, E., Reich-Zeliger, S., Gal, H., Best, K., Ndifon, W., Chain, B., Cohen, I.R.,

and Friedman, N. (2014). T-cell receptor repertoires share a restricted set of public and abundant

CDR3 sequences that are associated with self-related immunity. Genome Res. *24*, 1603–1612.

Madi, A., Poran, A., Shifrut, E., Reich-Zeliger, S., Greenstein, E., Zaretsky, I., Arnon, T.,

Laethem, F. Van, Singer, A., Lu, J., et al. (2017). T cell receptor repertoires of mice and humans

are clustered in similarity networks around conserved public CDR3 sequences. Elife *6*, 1–17.

McGranahan, N., Rosenthal, R., Hiley, C.T., Rowan, A.J., Watkins, T.B.K., Wilson, G.A.,

Birkbak, N.J., Veeriah, S., Van Loo, P., Herrero, J., et al. (2017). Allele-Specific HLA Loss and

Immune Escape in Lung Cancer Evolution. Cell 1259–1271.

Miles, J.J., Douek, D.C., and Price, D.A. (2011). Bias in the αβ T-cell repertoire: implications for

disease pathogenesis and vaccination. Immunol. Cell Biol. *89*, 375–387.

Moutaftsi, M., Peters, B., Pasquetto, V., Tscharke, D.C., Sidney, J., Bui, H.-H., Grey, H., and

Sette, A. (2006). A consensus epitope prediction approach identifies the breadth of murine

TCD8+-cell responses to vaccinia virus. Nat. Biotechnol. *24*, 817–819.

Ndifon, W., Gal, H., Shifrut, E., Aharoni, R., Yissachar, N., Waysbort, N., Reich-Zeliger, S.,

Arnon, R., and Friedman, N. (2012). Chromatin conformation governs T-cell receptor J gene

segment usage. Proc. Natl. Acad. Sci. *109*, 15865–15870.

Neefjes, J., Jongsma, M.L.M., Paul, P., and Bakke, O. (2011). Towards a systems understanding

of MHC class I and MHC class II antigen presentation. Nat. Rev. Immunol. *11*, 823–836.

Newell, E.W., Sigal, N., Nair, N., Kidd, B.A., Greenberg, H.B., and Davis, M.M. (2013).

Combinatorial tetramer staining and mass cytometry analysis facilitate T-cell epitope mapping

and characterization. Nat. Biotechnol. *31*, 623–629.

Olsen, L.R., Tongchusak, S., Lin, H., Reinherz, E.L., Brusic, V., and Zhang, G.L. (2017).

TANTIGEN: a comprehensive database of tumor T cell antigens. Cancer Immunol. Immunother.

66, 731–735.

Pardoll, D.M. (2012). The blockade of immune checkpoints in cancer immunotherapy. Nat. Rev.

Cancer *12*, 252–264.

R Core Team (2018). R: A Language and Environment for Statistical Computing.

Rasmussen, M., Fenoy, E., Harndahl, M., Kristensen, A.B., Nielsen, I.K., Nielsen, M., and Buus,

S. (2016). Pan-Specific Prediction of Peptide–MHC Class I Complex Stability, a Correlate of T

Cell Immunogenicity. J. Immunol. *197*, 1517–1524.

Reche, P.A., Zhang, H., Glutting, J.P., and Reinherz, E.L. (2005). EPIMHC: A curated database

of MHC-binding peptides for customized computational vaccinology. Bioinformatics *21*, 2140–

2141.

Reid, S.W., McAdam, S., Smith, K.J., Klenerman, P., O'Callaghan, C.A., Harlos, K., Jakobsen,

B.K., McMichael, A.J., Bell, J.I., Stuart, D.I., et al. (1996). Antagonist HIV-1 Gag peptides

induce structural changes in HLA B8. J. Exp. Med. *184*, 2279–2286.

Robins, H.S., Srivastava, S.K., Campregher, P. V., Turtle, C.J., Andriesen, J., Riddell, S.R.,

Carlson, C.S., and Warren, E.H. (2010). Overlap and Effective Size of the Human CD8+ T Cell

Receptor Repertoire. Sci. Transl. Med. *2*, 47ra64.

Rudolph, M.G., Stanfield, R.L., and Wilson, I.A. (2006). How TCRs bind MHCs, peptides, and

coreceptors. Annu. Rev. Immunol. *24*, 419–466.

Sakurai, Y., Brand, D.D., Tang, B., Rosloniec, E.F., Stuart, J.M., Kang, A.H., and Myers, L.K.

(2006). Analog peptides of type II collagen can suppress arthritis in HLA-DR4 (DRB1*0401)

transgenic mice. Arthritis Res. Ther. *8*, R150.

Sette, A., Vitiello, A., Reherman, B., Fowler, P., Nayersina, R., Kast, W.M., Melief, C.J.,

Oseroff, C., Yuan, L., Ruppert, J., et al. (1994). The relationship between class I binding affinity

and immunogenicity of potential cytotoxic T cell epitopes. J. Immunol. *153*, 5586–5592.

Shugay, M., Bolotin, D.A., Putintseva, E. V., Pogorelyy, M. V., Mamedov, I.Z., and Chudakov,

D.M. (2013). Huge overlap of individual TCR beta repertoires. Front. Immunol. *4*, 9–11.

Stone, J.D., Chervin, A.S., and Kranz, D.M. (2009). T-cell receptor binding affinities and

kinetics: impact on T-cell activity and specificity. Immunology *126*, 165–176.

Sykulev, Y., Brunmark, A., Jackson, M., Cohen, R.J., Peterson, P.A., and Eisen, H.N. (1994).

Kinetics and affinity of reactions between an antigen-specific T cell receptor and peptide-MHC

complexes. Immunity *1*, 15–22.

Tian, S., Maile, R., Collins, E.J., and Frelinger, J.A. (2007). CD8+ T cell activation is governed

by TCR-peptide/MHC affinity, not dissociation rate. J. Immunol. *179*, 2952–2960.

Tung, C.W., Ziehm, M., Kamper, A., Kohlbacher, O., Ho, S.Y., Kämper, A., Kohlbacher, O.,

and Ho, S.Y. (2011). POPISK: T-cell reactivity prediction using support vector machines and

string kernels. BMC Bioinformatics *12*, 446.

Udhayakumar, V., Anyona, D., Kariuki, S., Shi, Y.P., Bloland, P.B., Branch, O.H., Weiss, W.,

Nahlen, B.L., Kaslow, D.C., and Lal, A.A. (1995). Identification of T and B cell epitopes

recognized by humans in the C-terminal 42-kDa domain of the Plasmodium falciparum

merozoite surface protein (MSP)-1. J. Immunol. *154*, 6022–6030.

Vangone, A., and Bonvin, A.M.J.J. (2015). Contacts-based prediction of binding affinity in

protein–protein complexes. Elife *4*, 1–15.

Xue, L.C., Rodrigues, J.P., Kastritis, P.L., Bonvin, A.M., and Vangone, A. (2016). PRODIGY: A

web server for predicting the binding affinity of protein-protein complexes. Bioinformatics *32*,

3676–3678.

Yang, X., Chen, G., Weng, N., and Mariuzza, R.A. (2017). Structural basis for clonal diversity of

the human T-cell response to a dominant influenza virus epitope. J. Biol. Chem. *292*, 18618–

18627.

Yewdell, J.W. (2006). Confronting Complexity: Real-World Immunodominance in Antiviral

CD8+T Cell Responses. Immunity *25*, 533–543.

Zhong, S., Malecek, K., Johnson, L.A., Yu, Z., Vega-Saenz de Miera, E., Darvishian, F.,

McGary, K., Huang, K., Boyer, J., Corse, E., et al. (2013). T-cell receptor affinity and avidity

defines antitumor response and autoimmunity in T-cell immunotherapy. Proc. Natl. Acad. Sci.

110, 6973–6978.

Peptide sequences

Dataset search
- Databases
  - IEDB, LANL (HCV & HIV), EPIMHC, MHCBN, and TANTIGEN
- Previous studies

Epitope metadatasets
- HLA-I-restricted (n=21,162)
  - Immunogenic (n=6,957; 32.9%)
  - Contradicting (n=1,873; 8.9%)
- HLA-II-restricted (n=31,693)
  - Immunogenic (n=16,642; 52.5%)
  - Contradicting (n=4,505; 14.2%)

TCR sequences

Dataset search
- TCR repertoire data from healthy subjects
  - Seven BioProjects + One non-BioProject study
  - Datasets (n=206)
    - Datasets from individual donors (n=201)
      - Donors (n=103)
    - Datasets from multiple donors pooled (n=5)
      - Donors pooled (n=589 in total)
  - Unique CDR3β sequences (n=23,006,555)

Public TCR clonotypes
- Clonotypes observed in multiple datasets
  - Publicity threshold: 22 datasets
  - Unique CDR3β sequences (n=191,326; 0.83%)

Feature computation
- Peptide length (n=1)
- Residue existence (n=20)
- Peptide descriptors (n=1,440 and 2,160 for HLA-I and -II)
- TCR-peptide contact potential profiling (CPP) (n=4,620 and 6,930 for HLA-I and -II)
  - Random down-sampling of TCR sequences (n=10,000)
  - Average of five computation with different random seeds

Feature selection
- Twice repeated 5-fold CV
  - Data splitting (train:test=4:1)
  - Preprocessing (centering, scaling, correlation-based filtering)
  - Importance-based selection (n=100)
- The most predictive features = Intersection of feature sets derived from all attempts combined (n=27 and 32 for HLA-I and –II)

Immunogenicity prediction
- Five times repeated 5-fold CV
  - Data splitting (train:test=4:1)
  - Machine learning on training sets using the most important features
  - Probability estimation on test sets
- Immunogenicity scores = Averaged probabilistic estimates

784   **Figure 1. Data analysis workflow.** The workflow for generation of the probabilistic estimates

785   of epitope immunogenicity, termed immunogenicity scores, is shown. All datasets were retrieved

786   from public sources. CV, cross-validation. See also Figures S1 and S2, Data S1, and Materials

787   and Methods.

Figure 2. TCR-peptide contact potential profiling. (A) Schematic of antigen-guided TCR

repertoire convergence hypothesis. We formulate the concept of TCR-peptide contact potential

profiling (CPP) to identify population-wide immunogenicity by exploiting public TCR repertoire

inherently biased toward common pathogen antigens but away from self-antigens due to

genome-level evolution and postnatal thymic selection. (B) Schematic of the linear contact

model. The underlying hypothesis is that only a small portion of the CDR3β loop structure

recognizes an epitope. The structure of HIV reverse-transcriptase epitope (TAFTIPSI) and

cognate TCR CDR3β chain is shown as an example (PDB ID: 4MJI). (C) Mean-centered contact

density distributions determined from 51 unique peptide-WT TCR structures. Inset numbers

indicate the proportions of contacts falling into the color-specified ranges. (D) Schematic of the

800   approximation of best-matching region problem to pairwise sequence alignment problem. See

801   also Data S2, and Materials and Methods.

Figure 3. Contact potential profiling parameters linked epitope immunogenicity and TCR

affinity. (A-C) Feature importance estimates from ten iterations. (A and B) Importance estimates

for all CPP features aggregated by (A) fragment length and (B) descriptive statistics. (C)

Importance estimates for the most predictive 21 and 26 CPP features for MHC-I and MHC-II,

respectively, aggregated by their source AAIndex scales. The seven most predictive AAIndex

scales selected both in MHC-I and MHC-II are shown. (D-F) Regression analysis against

individual TCR affinities. Single-TCR contact potential profiling (sCPP) was conducted for 82

wildtype (WT) TCR-peptide complexes. Thirteen mutant (MT) TCRs were excluded from

correlation analysis and only shown for visualization purposes (red points). Adjusted squared

Pearson's correlation coefficients are presented. (D) Representative univariate regression. (E)

Multivariate regression with five sCPP features derived from immunogenicity-predicting

AAIndex scales. (F) Permutation experiments with 1,000 iterations. Multivariate regression was

performed as in (E) using sCPP features computed from either randomly permuted peptide

sequences or CDR3β sequence fragments. The blue line represents the non-permuted regression

result. Dashed lines represent medians. See also Figure S2, Data S2 and S3, and Table S1.

Figure 4. Immunogenicity prediction. Averaged probabilistic estimates from iterative machine

learning were termed "immunogenicity scores." (A) Immunogenicity scores generating under

various conditions for machine learning processes. MHC binding: incorporation of MHC binding

prediction results in machine learning. R, percentile ranks; B, binding strength categories.

Feature: the cutoffs for feature selection. The numbers of features actually retained after feature

selection under the cutoff of 100, 75, 50, and 25 were 32, 26, 19, and 7 for MHC-I, and 27, 23,

21, and 12 for MHC-II, respectively. Clustering: Elimination of highly similar peptide using the

IEDB Epitope Cluster Analysis Tool (http://tools.iedb.org/cluster/) with either 100% (no

clustering) or 80% homology threshold. The numbers of peptides retained after 80% homology

clustering were 16,765 and 23,983 for MHC-I and MHC-II, respectively. (B) Immunogenicity

scores generated for various subsets of our epitope datasets. Machine learning was conducted

with 32 and 27 the most predictive features for MHC-I and MHC-II, respectively, but without

predicted MHC binding information. Evidence: IEDB-derived peptides with functional T cell

assay evidence (*i.e.*, without non-IEDB data); Human: epitopes tested in human (*i.e.*, without

transgenic animal hosts); Ex vivo: IEDB-derived peptides annotated from direct *ex vivo* assays

using autologous effector and antigen-presenting cells in human. The numbers of peptides

included after applying each filter were 21,162, 13,058, 11,084, and 6,048 for MHC-I, and

31,693, 31,693, 30,188, and 2,023 for MHC-II, respectively. (C) Immunogenicity scores of

human epitopes and extrapolated scores for primate and rodent epitopes (N=411 and 8,756 for

MHC-I and 76 and 8,445 for MHC-II, respectively). (D and E) Distributions of (D) annotated

immunogenicity and (E) immunogenicity scores for the human thymus MHC-I peptidome

(Adamopoulou et al., 2013). (F and G) Immunogenicity scores among IEDB-derived (F) MHC-I-

and (G) MHC-II-restricted peptides of various origins. HS, *homo sapiens*; HIV, human

immunodeficiency virus type I; HBV, hepatitis B virus; HCV, hepatitis C virus; IAV, influenza

A virus; DGV, dengue virus; PF, *Plasmodium falciparum*; TB, *Mycobacterium tuberculosis*. (H)

Immunogenicity scores among MHC-I-restricted peptides from various sources. HIV, Los

Alamos HIV Epitope Database; HCV, Los Alamos HCV Epitope Database. (I) Immunogenicity

scores of MHC-I-restricted neoepitopes. NE, neoepitopes from TANTIGEN database. (J)

Distributions of immunogenicity scores for neoepitopes grouped by their source antigen proteins.

Red points and bars represent median and interquartile ranges, respectively. In (A-I), orange and

blue represent immunogenic and non-immunogenic peptides, respectively. In (A-D) and (F-H),

bars represent medians. In (E-I), ns, not significant; *, $P < 0.05$; ***, $P < 0.001$; ****, $P <$

0.0001. See also Figure S2, S3, S4, and S5, Data S1, S3, and S4.

Figure 5. Systematic characterization of the impacts of single amino acid mutations on

immunogenicity in sequence space. (A) Schematic of the concept of immune transition in

sequence space. We termed the change in immunogenicity between neighbors, *i.e.*, single-aa

mutants, as "immune transition," and defined peptides with evidence of immune transition as

"transitional." (B and C) Immunogenicity scores among (B) MHC-I- and (C) MHC-II-restricted

transitional peptides, *i.e.*, peptides with at least one neighbor of opposite immunogenicity

annotation in our dataset. We identified 1,360 and 976 transitional peptides for MHC-I and

MHC-II, respectively. We expanded their neighbor networks by computing immunogenicity

scores for 232,723 and 292,437 all possible single-aa mutants of transitional MHC-I- and MHC-

II-restricted peptides, respectively. Original, original immunogenicity scores; Overall, mean

immunogenicity scores of all neighbors. Cluster, mean immunogenicity scores of neighbors

assigned to the cluster containing the parent peptide. Others, original immunogenicity scores for

peptides with no evidence of immune transition (shown for comparison purpose). Orange and

blue denote immunogenic and non-immunogenic peptides, respectively. (D-F) Immunogenicity

score dynamics among observed neighbor pairs. We identified 6,179 and 3,076 single-aa

mutated peptide pairs for MHC-I and MHC-II, respectively. (D) Score dynamics for all possible

peptide pairs, regardless of their origins. (E) Score dynamics for pairs of peptides derived from

the same organism. Blue and red denote non-transitional and transitional pairs, respectively. (F)

Score dynamics of MHC-I peptide pairs stratified by their mutated positions. Cter, C-terminal

residue. (G and H) Heatmap clustering analysis of immunogenicity score dynamics by mutating

residue pairs. We generated 2,580,890 and 3,101,092 neighbor pairs from 232,723 and 292,437

all possible single-aa mutants of 1,360 and 976 transitional MHC-I- and MHC-II-restricted

peptides, respectively. Color represents median normalized score changes. Grey means

unobserved residue pairs. In (D-F), ns, not significant; *, $P < 0.05$; **, $P < 0.01$; ****, $P <$

0.0001. See also Figure S6 and Data S5.

Figure 6. Identification of escape-prone positions from *in silico* mutagenesis and neighbor network analysis. (A) Example neighbor networks of MHC-I-restricted influenza A virus epitopes GILGFVFTL and SRYWAIRTR and their observed neighbors. Consensus sequences are indicated below the cluster IDs. (B and C) Relative score changes estimated from all possible simulated neighbors for (B) MHC-I-restricted influenza A virus epitopes GILGFVFTL and SRYWAIRTR and (C) MHC-II-restricted insulin-derived epitope GAGSLQPLALEGSLQKRG and collagen-derived epitope PKGQTGEPGIAGFKGEQGPK. Red characters indicate residues with known escaping mutations shown by *in vitro* T cell assays. Example peptides were chosen

based on their high immunogenicity scores with manual literature inspection. ns, not significant;

*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$. See also Figure S7.

Figure 7. Two-dimensional assessment of epitope quality using the immunogenicity score

and escaping potential. (A and B) Immunogenicity score-escaping potential (IS-EP) plots for

(A) MHC-I and (B) MHC-II peptides. Cluster-mean immunogenicity scores determined from

neighbor network analysis of all simulated neighbors were utilized. Labeled are representative

epitopes shown in Figures 6 and 7. (C) IS and EP distributions among pathogen-derived peptides.

ns, not significant; *, $P < 0.05$; **, $P < 0.01$. (D-F) Relative score changes estimated from all

possible simulated neighbors for (D) representative MHC-I-restricted HIV epitopes, and (E and

F) representative MHC-II-restricted epitopes derived from (E) *P. falciparum* and (F) *M.*

tuberculosis. In (D-F), representative peptides were chosen based on their high and low EPs with

manual literature inspection.

Supplementary Materials for

The landscape of T cell epitope immunogenicity in sequence space

Masato Ogishi and Hiroshi Yotsuyanagi

Correspondence to: Masato Ogishi (oogishi-tky@umin.ac.jp)

 Supplementary Figures

Figure S1. Base characteristics of the epitope dataset, Related to Figure 1. (A and B)
Relative abundance of (A) immunogenicity annotations and (B) conflicts in immunogenicity
annotations among peptides of different lengths. (C and D) Relative abundance of (C)

925 immunogenicity annotations and (D) conflicts in immunogenicity annotations among peptides of
926 different origins. HS, homo sapiens; HIV, human immunodeficiency virus type 1; HBV, hepatitis
927 B virus; HCV, hepatitis C virus; IAV, influenza A virus; DGV, dengue virus; PF, Plasmodium
928 falciparum; TB, Mycobacterium tuberculosis; Others, peptides derived from other organisms
929 (peptides with no source organism annotation were excluded).
930

A

B

932 **Figure S2. Correlograms of the minimal sets of features predictive of epitope**
933 **immunogenicity, Related to Figure 3.** (A) Features for MHC-I-restricted peptides (N=32). (B)
934 Features for MHC-II-restricted peptides (N=27). Features were reordered by the hierarchical
935 clustering algorithm.
936

Figure S3. Robustness analysis of immunogenicity predictions, Related to Figure 4.
Performances of probabilistic estimates computed through machine learning under various conditions were examined. (A and B) Addition of MHC binding prediction results as features for machine learning. Rank, percentile ranks; Bind, binding strength categories. (A) MHC-I-restricted peptides. (B) MHC-II-restricted peptides. (C and D) Machine learning with smaller cutoff values for feature selection. (C) MHC-I-restricted peptides. The actual numbers of features retained after feature selection were 32, 26, 19, and 7, respectively. (D) MHC-II-restricted peptides. The actual numbers of features retained were 27, 23, 21, and 12, respectively. (E and F) Peptide homology-based clustering before machine learning. The IEDB Epitope Cluster Analysis Tool (http://tools.iedb.org/cluster/) was used to cluster the peptides with the homology threshold of 80%. (E) MHC-I-restricted peptides. A total of 16,765 peptides were retained after clustering. (F) MHC-II-restricted peptides. A total of 23,983 peptides were retained after clustering. Non-filtered, all peptides. Filtered, single peptide randomly chosen from each of the clusters before preprocessing.

Figure S4. Immunogenicity prediction of HLA-I-restricted peptides by the IEDB Class I Immunogenicity Tool, Related to Figure 4. Classification performances were evaluated using the prediction scores (rescaled to 0-1 range) derived from the IEDB Class I Immunogenicity Tool (Calis et al., 2013) (http://tools.iedb.org/immunogenicity/) with default settings. Note that peptides used for training the model in the original paper were not excluded from prediction, which theoretically overestimates its predictive performance.

Figure S5. Extrapolation of the human immunogenicity prediction framework for primate and rodent epitopes, Related to Figure 4. (A and B) Overlaps between (A) MHC-I-restricted and (B) MHC-II-restricted peptides presented on MHC molecules of different species. (C and D) Predictive performances of the probabilities of immunogenicity for (C) MHC-I-restricted and (D) MHC-II-restricted peptides. Probabilities for non-human peptides were estimated by applying the classifiers trained on the human peptide datasets.

Figure S6. Improvement of predictive performances of immunogenicity scores for transitional peptides by integrating the neighbor network architectures in sequence space, Related to Figure 5. Predictive performances of the immunogenicity scores of (A) MHC-I- and (B) MHC-II-restricted transitional peptides, *i.e.*, peptides with at least one neighbor of opposite immunogenicity annotation in our dataset, were evaluated. We identified 1,360 and 976 transitional peptides for MHC-I and MHC-II, respectively. We expanded their neighbor networks by computing immunogenicity scores for 232,723 and 292,437 all possible single-aa mutants of transitional MHC-I- and MHC-II-restricted peptides, respectively. Original, original immunogenicity scores; Overall, mean immunogenicity scores of all neighbors. Cluster, mean immunogenicity scores of neighbors assigned to the cluster containing the parent peptide. Others, original immunogenicity scores for peptides with no evidence of immune transition (shown for comparison purpose).

A

B

Figure S7. Assessment of position-specific escaping potentials for representative peptides
using probabilistic estimates of immunogenicity inferred without target peptides and their
neighbors during machine learning, Related to Figure 6. Probabilities for the target and its
neighbors were computed by applying the classifiers trained without target peptides and their
neighbors. See the legend of Figure 6 for further information.

990 **Supplementary Tables**
991
992 **Table S1. Variance inflation factors (VIFs) of the features used in the multivariate**
993 **regression analysis against experimentally determined TCR affinities.**
994

Feature	VIF
sCPP_BETM990101inv_Skew_4	1.25
sCPP_KESO980102inv_Kurt_3	1.11
sCPP_KOLA930101_Skew_4	1.29
sCPP_SIMK990104_SD_3	1.14
sCPP_VENM980101inv_Min_3	1.06

995
996

Supplementary Datasets

Data S1. (separate file)
Human, primate, and rodent T cell epitope dataset.

Data S2. (separate file)
TCR-peptide affinity and structure dataset.

Data S3. (separate file)
Feature importance estimates.

Data S4. (separate file)
Immunogenicity scores.

Data S5. (separate file)
A list of single amino acid mutant pairs identified.

Data S6. (separate file)
A list of AAIndex amino acid pairwise contact potential scales.