Nuclease escape elements protect messenger RNA against cleavage by

multiple viral endonucleases

Mandy Muller[1,2]; Britt A. Glaunsinger[1,3,4*]

[1.] Department of Plant and Microbial Biology, University of California, Berkeley, California,
USA CA 94720-3370
[2.] Present address: Department of Microbiology, University of Massachusetts at Amherst,
Amherst, MA 01003 USA.
[3.] Department of Cell and Molecular Biology, University of California, Berkeley, California,
USA CA 94720-3370
[4.] Howard Hughes Medical Institute
12
13 * Corresponding author; glaunsinger@berkeley.edu

14 **Short Title**: Viral nuclease escape elements

ABSTRACT

During lytic Kaposi's sarcoma-associated herpesvirus (KSHV) infection, the viral endonu-clease SOX promotes widespread degradation of cytoplasmic messenger RNA (mRNA). However, select mRNAs, including the transcript encoding interleukin-6 (IL-6), escape SOX-induced cleavage. IL-6 escape is mediated through a 3' UTR RNA regulatory element that overrides the SOX targeting mechanism. Here, we reveal that this protective RNA element functions to broadly restrict cleavage by a range of homologous and non-homologous viral endonucleases. However, it does not impede cleavage by cellular endonucleases. The IL-6 protective sequence may be representative of a larger class of nuclease escape elements, as we identified a similar protective element in the GADD45B mRNA. The IL-6 and GADD45B-derived elements display similarities in their sequence, putative structure, and several associated RNA binding proteins. However, the overall composition of their ribonucleoprotein complexes appears distinct, leading to differences in the breadth of nucleases restricted. These findings highlight how RNA elements can selectively control transcript abundance in the background of widespread virus-induced mRNA degradation.

AUTHOR SUMMARY

The ability of viruses to control the host gene expression environment is crucial to promote

viral infection. Many viruses express factors that reduce host gene expression through

widespread mRNA decay. However, some mRNAs escape this fate, like the transcript

encoding the immunoregulatory cytokine IL-6 during KSHV infection. IL-6 escape relies on

an RNA regulatory element located in its 3'UTR and involves the recruitment of a

protective protein complex. Here, we show that this escape extends beyond KSHV to a

variety of related and unrelated viral endonucleases. However, the IL-6 element does not

protect against cellular endonucleases, revealing for the first time a virus-specific nuclease

escape element. We identified a related escape element in the GADD45B mRNA, which

displays several similarities with the IL-6 element. However, these elements assemble a

largely distinct complex of proteins, leading to differences in the breadth of their protective

capacity. Collectively, these findings reveal how a putative new class of RNA elements

function to control RNA fate in the background of widespread mRNA degradation by viral

endonucleases.

INTRODUCTION

A number of viruses restrict host gene expression to reduce competition for resources

and dampen immune responses. This 'host shutoff' phenotype can be triggered through a

range of mechanisms that operate at nearly every stage of the gene expression cascade.

Viruses whose host shutoff strategies involve the induction of widespread mRNA decay

include the alpha and gammaherpesviruses, vaccinia virus (VACV), influenza A virus (IAV),

and SARS coronavirus (SCoV) [1-5] In each of the above cases, mRNA degradation is

induced *via* one or more internal endonucleolytic cleavages in the target mRNA or, in the

case of VACV, direct removal of the mRNA 5' cap [5-9]. This is invariably followed by

exonucleolytic degradation of the cleaved fragment(s) by components of the mammalian

RNA decay machinery such as Xrn1 [1,10,11].

The viral strategy contrasts with basal mRNA degradation in eukaryotes, which is a

tightly regulated process that initiates with gradual shortening of the poly(A) tail, followed

by removal of the 5' cap prior to exonucleolytic degradation of the transcript body [12].

Although eukaryotes encode endonucleases, they are generally restricted to a highly

specific set of targets. For example, mRNAs containing premature stop codons are cleaved

by the Smg6 endonuclease during the translation-linked quality control process of

nonsense mediated decay (NMD) [13]. No-go decay is another form of quality control

activated in cases of ribosome stalling, although the specific endonuclease that cleaves the

mRNA remains unknown [14,15]. The use of endonucleases during quality control enables

more rapid removal of aberrant mRNAs from the translation pool, as their inactivation is

not reliant on the prior rate limiting steps of deadenylation and decapping. In this regard,

virus-induced mRNA decay resembles the cellular quality control mechanisms, but with

significantly expanded scope.

One of the well-studied viral endonucleases is the SOX protein encoded by ORF37 of

Kaposi's sarcoma-associated herpesvirus (KSHV). During lytic KSHV replication, SOX is

expressed with delayed early kinetics and its nuclease activity significantly reduces

cytoplasmic mRNA levels [16]. SOX is conserved throughout the herpesvirus family, but

only gammaherpesviral SOX homologs display ribonuclease activity in cells [16-18].

Perhaps surprisingly, its activity is not restricted to host mRNAs, and studies with the SOX

homolog from murine gammaherpesvirus 68 (MHV68) indicate that SOX activity helps fine

tune viral mRNA levels in a manner important for the *in vivo* viral lifecycle [19,20].

Although most mRNAs are subject to cleavage by SOX, a recent degradome-based

sequencing analysis together with studies on individual endogenous and reporter mRNAs

revealed that SOX cleavage sites are defined by a degenerate RNA motif [10,21]. The SOX-

targeting motif can be located anywhere within an mRNA and may be present multiple

times [10,21].

The observation that a targeting motif is present on SOX cleaved mRNAs suggests that

transcripts lacking this element should escape cleavage. Indeed, RNAseq analyses indicate

that approximately one-third of mRNAs are not depleted by SOX [22,23]. Studying these

'escapees' in aggregate is complicated, however, by the fact that multiple mechanisms can

promote apparent escape. These include lack of a targeting motif, indirect transcriptional

effects, and active evasion of cleavage [22,24-28]. This latter phenotype, termed dominant

escape, is particularly notable as it involves a specific RNA element whose presence in the

3' UTR of an mRNA protects against SOX cleavage, regardless of whether the RNA contains

a targeting motif. The one known example of dominant escape derives from the host

interleukin-6 (IL-6) transcript [26,27,29].

IL-6 expression is required for survival of B cells infected with KSHV, and the virus

engages a number of strategies to drive production of this cytokine [30-38]. The IL-6 mRNA

is directly refractory to SOX cleavage and thus remains robustly induced during host

shutoff due to the presence of a specific 'SOX resistance element' (SRE) [26,29]. Even in the

absence of infection, reporter mRNAs bearing the IL-6 SRE remain stable in SOX-expressing

cells, an observation that has helped delineate features of this novel RNA element required

for the protective phenotype [26,27]. The IL-6 SRE was fine mapped to a 200 nt sequence

within the 3' UTR, which was subsequently shown to assemble an 'escape complex' of at

least 8 cellular RNA binding proteins involved in protection [26,27]. How this complex

functions to restrict SOX recognition remains largely unknown, although nucleolin (NCL)

plays an essential role. NCL is partially relocalized from the nucleolus to the cytoplasm

during lytic KSHV infection, where it binds the SRE using its RNA recognition motif and

engages in protein-protein interactions related to escape with its carboxyl-terminal RGG

domain [27].

The IL-6 SRE is the first described ribonuclease escape element and much remains to

be learned about its function, as well as whether other related elements exist that protect

their associated mRNA. Here, we reveal that the IL-6 SRE is broadly protective against a

diverse group of viral endonucleases, suggesting an underlying commonality in the

mechanism by which these host shutoff factors recognize their mRNA targets. It is not

indiscriminately protective, however, as host quality control endonucleases are not

blocked by the IL-6 SRE. We then identify a second, novel SRE within the GADD45B mRNA,

which displays some physical and functional similarities to the IL-6 SRE. Collectively, these

findings suggest that a diversity of nuclease escape elements exist, and that their

characterization may lead to new insights into the control of mRNA fate in both infected

and uninfected cells.

RESULTS

The IL-6 derived resistance element broadly protects against cleavage by viral but

not cellular endonucleases

The 3' UTR of IL-6 contains a transferrable 200 nt SRE that protects its associated mRNA

from SOX-induced degradation [26,27]. The SRE also protects mRNA from cleavage by the

unrelated vhs endonuclease encoded by herpes simplex virus type 1 (HSV-1), hinting that

this RNA element may restrict endonuclease targeting in a broader capacity [27]. To test

this hypothesis, we assessed whether the presence of the SRE impacted the ability of a

panel of homologous and heterologous mRNA-specific viral endonucleases to degrade a

target mRNA. In addition to HSV-1 vhs, these included homologs of KSHV SOX from the

related gammaherpesviruses Epstein-Barr virus (EBV; BGLF5) and murine

gammaherpesvirus 68 (MHV68; muSOX), as well as the heterologous host shutoff

endonuclease from influenza A virus (IAV; PA-X) [3,17,39]. As anticipated, the control GFP

mRNA was readily degraded in 293T cells co-transfected with plasmids expressing each of

the viral endonucleases as measured by RT-qPCR (**Fig. 1A**). However, addition of the IL-6

derived SRE to the 3' UTR of GFP (GFP-IL-6-SRE) prevented each of the viral endonucleases

from degrading this mRNA (**Fig. 1B**). Fusion of a size matched segment of the IL-6 3' UTR

lacking the SRE to GFP (GFP-IL-6 ΔSRE) reinstated cleavage of the GFP reporter by the viral

endonucleases (**Fig. 1B**). These data confirm that the protection conferred by the SRE is

not specific to SOX-induced cleavage, but functions more broadly to restrict mRNA cleavage

by a diverse set of mammalian virus host shutoff endonucleases.

Cleavage sites for IAV PA-X have yet to be determined, but the other endonucleases

do not appear to target mRNA in the same location or using the same sequence features,

including the SOX homologs [1,9,21,40,41]. Thus, the IL-6 derived SRE is unlikely to

function through steric occlusion of a common cleavage site. We instead considered the

possibility that the SRE functions like an RNA 'zip code', directing its associated transcript

to a location in the cell inaccessible to endonucleases. To test this hypothesis, we used RNA

fluorescence *in situ* hybridization (FISH) to monitor how the presence of the IL-6 SRE

impacted the localization of its associated mRNA. Six phage MS2-derived stem-loops were

introduced upstream of the SRE or ΔSRE segment of the IL-6 3' UTR in the pcDNA3

luciferase reporter, enabling visualization of the RNA in transfected 293T cells using a Cy3-

labeled RNA probe directed against the MS2 sequences [42]. We verified that the presence

of the stem-loops did not prevent the escape of the SRE-containing reporter or degradation

of the ΔSRE reporter in SOX-expressing cells (**Fig. S1**). There was no distinguishable

difference in the localization of the SRE and ΔSRE containing mRNAs, both of which were

present relatively diffusely throughout the cell (**Fig. 1C**). The FISH signal was specific to

transfected cells, as we observed no fluorescence in neighboring untransfected cells nor

autofluorescence from transfected cells lacking the Cy3 probes (**Fig. 1C**). Although these

data to not exclude the possibility that the SRE-containing transcript became sequestered

into micro-aggregates or other structures not visible at this level of resolution, they do not

support relocalization as the driver of escape from viral endonuclease cleavage.

162
We next considered whether the inability to cleave an IL-6 SRE-containing

transcript was specific to viral endonucleases or similarly extended to host endonucleases.

Although basal cellular mRNA decay is carried out by exonucleases, host quality control

pathways involve endonucleases to promote rapid clearance of aberrant mRNA [43-45].

We applied two strategies to monitor the activity of the SRE against host endonucleases.

The first was to use a nonsense mediated decay (NMD) target containing a premature

termination codon (PTC) 100 amino acids into the body of an RFP reporter mRNA (dsRed2-

PTC) [10]. The NMD pathway detects PTCs during translation and directs cleavage of the

mRNA by the Smg6 endonuclease [46]. Depletion of Smg6 from 293T cells using siRNAs

restored dsRed2-PTC mRNA levels to those of the control dsRed2 transcript, confirming

that this was an NMD substrate degraded by Smg6 (**Fig. 2A-B**). We then fused the IL-6

derived SRE to the 3' UTR of dsRed2-PTC (dsRed2-PTC-SRE) or, as a control, the size

matched region from the IL-6 3' UTR lacking the SRE (dsRed2-PTC-ΔSRE) and measured

the levels of each mRNA by RT-qPCR in 293T cells (**Fig. 2C**). In contrast to the viral

endonucleases, the SRE did not impair Smg6-mediated cleavage of its target mRNA, as all

three PTC-containing transcripts were similarly degraded (**Fig. 2A & 2C**).

The second strategy to monitor host endonuclease activity was to express the nsp1

protein from SARS coronavirus. Nsp1 is not itself a nuclease, but it binds the 40S ribosome

and causes it to stall on the mRNA, thus activating cleavage of the mRNA by an as yet

unknown cellular endonuclease via a mechanism reminiscent of no-go decay [47,48].

Although the endonuclease involved in this pathway has not been established, it is known

not to be Smg6 and thus this enabled evaluation of a distinct host endonuclease [48]. Nsp1

was co-transfected with the GFP-SRE or GFP-ΔSRE reporter into 293T cells, and depletion

of the GFP transcript was measured by RT-qPCR. Similar to the NMD substrate, the SRE did

not prevent degradation of the GFP mRNA in nsp1-expressing cells (**Fig. 2D**). Collectively,

these results suggest that the IL-6 derived SRE confers broad protection against viral but

not cellular endonucleases. Given that the host endonucleases require ongoing translation

for target recognition, these data also confirm that the SRE does not pull its associated

mRNA out of the translation pool.

GADD45B mRNA is also resistant to SOX-induced degradation.

To determine whether this property of broad inhibition of viral endonucleases was

restricted to the IL-6 SRE, we sought to identify other SRE-bearing transcript(s). Based on

our previous finding that the IL-6 SRE required binding by a complex of cellular proteins

including NCL, we mined a published RNAseq dataset for transcripts that were not

depleted in SOX expressing 293T cells and were known to be bound by NCL [22,49-51]. A

transcript that fit these criteria was growth arrest and DNA damage-inducible 45 beta

(GADD45B).   Notably, the GADD45B mRNA was also previously shown to escape

degradation in HSV-1 vhs-expressing cells, further suggesting it might contain an SRE [52-

54]. We first examined whether the GADD45B mRNA was resistant to host shutoff upon

lytic reactivation of a KSHV-positive B cell line (TREX-BCBL1) and a renal carcinoma cell

line stably expressing the KSHV BAC16 (iSLK.219). Both TREX-BCBL1 and iSLK.219 cells

harbor a doxycycline (dox)-inducible version of the major viral lytic transactivator RTA

that promotes entry into the lytic cycle upon dox treatment [55,56]. Unlike the GAPDH

mRNA, which is degraded by SOX upon lytic reactivation, the GADD45B mRNA levels

remained unchanged both in reactivated TREX-BCBL1 and iSLK.219 cells as measured by

RT-qPCR (**Fig. 3A & 3B**). We also confirmed that, unlike the GAPDH mRNA, the endogenous

GADD45B mRNA was not depleted upon transfection of KSHV SOX or HSV-1 vhs into 293T

cells (**Fig. 3C**).   Finally, we noted that siRNA-based depletion of GADD45B from iSLK.219

cells resulted in decreased efficiency of lytic KSHV reactivation as well as reduced

expression of representative delayed early (ORF59) and late (K8.1) viral genes, suggesting

that GADD45B expression is important for the KSHV lytic cycle (**Fig. S2**).

Recently, we showed that KSHV SOX cleaves its targets at a specific but degenerate

RNA motif [21]. Thus, the failure of SOX (or perhaps vhs) to degrade the GADD45B mRNA

could either be due to the absence of such a targeting motif (e.g. passive escape), or to the

presence of a specific protective element like the IL-6 SRE (e.g. dominant escape). To

distinguish these possibilities, we constructed chimeras between GFP, which has a well-

characterized SOX cleavage element, and the GADD45B 5' UTR, 3' UTR, or coding region

(CDS), each of which were cloned downstream of the GFP coding region (**Fig. 3D**). Co-

transfection of the GFP-fused GADD45B 3' UTR construct with SOX into 293T cells did not

lead to degradation of this mRNA, whereas the GFP-GADD45B 5' UTR or CDS fusions were

readily degraded in SOX-expressing cells (**Fig. 3E**). Thus, similar to the IL-6 3' UTR, the

GADD45B 3' UTR contains a protective sequence that prevents SOX cleavage of an

established target mRNA.

Hairpin structures within the GADD45B and IL-6 SREs are important for conferring

resistance to SOX cleavage

To refine which GADD45B sequence encompassed the SRE, we initially looked for

similarities between the GADD45B 3' UTR and the IL-6 SRE using Clustal W alignment.

While there were no stretches of significant sequence identity between the two RNAs, the

last ~200nt of the GADD45B 3'UTR had the highest similarity (~46%) to the IL-6 SRE (**Fig.**

S3). We therefore fused this putative SRE segment of the GADD45B 3' UTR to GFP

(GADD45B-SRE), and found that it was sufficient to confer nearly the same level of

protection from SOX as the full GADD45B 3' UTR in transfected 293T cells (**Fig. 4A**). Thus,

similar to IL-6, the GADD45B 3' UTR contains a ~200 nt SRE (henceforth termed G-SRE).

We next sought to determine whether these two SREs could adopt a common

secondary structure. RNAfold-based predictions showed that the 3'-most segment of both

SREs form a long stem-loop protruding structure with at least one bulge near the middle of

the stem, whereas the other regions of the SREs did not fold into any similar high-

confidence structures (**Fig. 4B**). To validate this predicted SRE stem loop structure

experimentally, we applied in-line probing, an RNA cleavage assay in which base-paired or

structurally constrained nucleotides are protected from spontaneous phosphodiester bond

hydrolysis [57]. Results from the cleavage reaction (**Fig. S4**) largely confirmed the RNAfold

predictions, apart from a small variation in the IL-6 hairpin. We then tested whether this

structure is required for either IL-6 SRE or G-SRE function by changing two conserved TT

nucleotides located directly adjacent to the bulge in each hairpin structure to GG (SRE_GG;

mutated residues marked with asterisks Fig. 4B). We also separately mutated the AA

residues predicted to base pair with these nucleotides on the other side of the loop to CC

(SRE_CC). Both the SRE_GG and the SRE_CC mutations in the IL-6 or GADD45B GFP fusions

resulted in partial degradation of these mRNAs upon SOX expression in 293T cells,

suggesting that disruption of that portion of the SRE hairpin impaired the protective

capacity of each SRE **(Fig. 4C & 4D).** Notably, combining these mutations together

(SRE_GG+CC), which should restore the secondary structure of the stem-loop, rescued the

fully protective phenotype in SOX expressing cells (**Fig. 4C & 4D**). This suggests that, at

least for this region of each SRE, RNA structure rather than the specific sequence is

important for protection against SOX cleavage.

The GADD45B and IL-6 SREs assemble a partially overlapping ribonucleoprotein complex

Using an *in vitro* RNA-pulldown based strategy, we previously determined that the IL-6 SRE assembles a specific ribonucleoprotein (RNP) 'escape' complex that is critical for its ability to mediate protection from cleavage by SOX [26,27]. Given the partial similarities in length, sequence, and structure between the IL-6 SRE and the G-SRE, we hypothesized that they might also assemble a similar set of RNPs to mediate their protective function. Indeed, it had already been established that both transcripts bind NCL [27,49]. We therefore performed Comprehensive Identification of RNA binding Proteins by Mass Spectrometry (ChIRP-MS) to compare the set of proteins bound *in vivo* to the IL-6 and GADD45B 3'UTRs in transfected 293T cells (**Fig. 5A**). Briefly, ChIRP-MS involves purifying an RNA of interest along with its associated proteins from crosslinked, sonicated cells using specific RNA probe-based capture, then identifying the bound proteins by MS [58]. Control probes directed against GFP were included to identify nonspecific interactions, which were then filtered out of the dataset. Using ChIRP-MS, we identified 195 proteins associated with the GADD45B 3'UTR and 245 proteins associated with the IL-6 3'UTR, of which 124 were in common between the two sets (**Table S1 & Fig. S5**). Many of the proteins previously identified as bound to the IL-6 SRE (**Fig. 5B**, gray nodes; [27]) were also recovered using ChIRP-MS for the IL-6 3' UTR and were similarly associated with the GADD45B 3' UTR (**Table S2** & **Fig. 5B**, purple nodes). We sought to independently validate a subset of these common interactors by western blot following ChIRP (**Fig. 5C**). Again, we recovered NCL and HuR (also known as ELAVL1), two proteins that were previously identified as important for IL-6 escape from SOX degradation [26,27]. We also confirmed hnRNPU, a

protein that binds the IL-6 SRE but is not required for escape of the IL-6 SRE from SOX [27].

However, other IL-6 SRE-bound proteins required for its escape function were not detected

on the GADD45B 3' UTR by ChIRP-western blot, including IGF2BP1, STAU1, ZC3HAV1,

YTHDC2, NPM1, and hnRNPD (**Fig. 5C**). The fact that IGF2BP1 and NPM1 were recovered

in the GADD45B and IL-6 ChIRP-MS experiments but not in the ChIRP-western blots likely

reflects differences in sensitivity between MS and western blotting.

To further validate the interactions with NCL, HuR, and hnRNPU, we

immunoprecipited (IP) each endogenous protein from 293T cells and performed RT-qPCR

to measure the level of co-precipitating endogenous GADD45B RNA. We observed a >5-

fold enrichment of GADD45B mRNA over the mock (IgG) IP for both NCL and HuR, although

we were not able to detect an association with hnRNPU in this assay (data not shown) (**Fig.**

5D). We confirmed that the interaction occurs on the GADD45B 3'UTR by performing the

IPs from cells transfected with a GFP reporter fused to either the GADD45B 5' UTR or 3'

UTR (**Fig. 5E**). The structurally compromised mutant GADD45B 3'UTR_GG described in Fig.

4B failed to interact with NCL and HuR, confirming that the stem-loop structure in the SRE

is important for protein binding (**Fig. 5E**). Collectively, these data suggest that while there

is some overlap between the sequence, structure, and RNA binding proteins associated

with the GADD45B and IL-6 SREs, these elements likely assemble distinct RNP complexes

and thus may function in a related but non-identical manner.

302
The GADD45B and IL-6 SREs share a requirement for HuR but differ in the breadth of

their protective capacity

Depletion of either HuR or NCL impairs the ability of the IL-6 derived SRE to protect

its associated mRNA from degradation by SOX [26,27]. Given that both proteins are also

bound by the G-SRE, we evaluated whether they were similarly important for G-SRE-

mediated escape. We individually depleted each protein from 293T cells using siRNAs

targeting HuR or NCL (or control non-targeting siRNAs), then measured the ability of SOX

to degrade the GFP-GADD45B-3'UTR reporter by RT-qPCR (**Fig. 6A & 6B**). Similar to the IL-

6 SRE, depletion of HuR eliminated the protective capacity of the G-SRE, leading to

degradation of the GFP-GADD45B-3'UTR mRNA in SOX-expressing cells (**Fig. 6A**).

Surprisingly however, depletion of NCL did not impair the protective effect of the G-SRE in

SOX-expressing cells (**Fig. 6B**).

Finally, we used the GFP-GADD45B-3'UTR reporter to evaluate whether the G-SRE

conferred protection from cleavage by the panel of viral endonucleases described in Fig. 1.

Using the same experimental set up as was used for the IL-6 derived SRE reporter, we

observed that while the GADD45B 3' UTR protected against SOX and vhs, it was unable to

protect against degradation by muSOX, BGLF5, and PA-X (**Fig. 6C**). Furthermore, it did not

protect against cleavage by the nsp1-activated host endonuclease (**Fig. 6C**). Thus, although

the IL-6 and GADD45B derived nuclease escape elements exhibit a number of similarities,

they are distinct in both their RNP complex requirements and in the breadth of nucleases

they restrict.

DISCUSSION

Viruses extensively interface with the host gene expression machinery to promote

their own RNA and protein synthesis and to control the cellular response to infection. In

this regard, they have proven to be invaluable tools to dissect mechanisms of gene

regulation. Here, we reveal that a ~200 nt sequence present in the 3' UTR of the cellular IL-

6 mRNA functions as a broad-acting, virus-specific endonuclease escape element, and

identify a similar element in the GADD45B 3' UTR. Although these two SREs are not

identical in their protective capacity, they display some sequence and putative structural

similarities and are both functionally dependent on HuR binding. We hypothesize that

these may therefore be representative members of a new type of RNA regulatory element

engaged during viral endonuclease-triggered mRNA decay. Determining whether other

such elements exist in mammalian or viral mRNAs is an important future goal, as is

deciphering conditions under which such elements impact RNA fate in uninfected cells.

The IL-6 and GADD45B SREs display relatively limited sequence similarity but have

at least one structurally important stem-loop in common. Thus, RNA structure appears to

be a central component of an SRE and is likely to influence recruitment or arrangement of

proteins involved in escape. This is in line with the fact that RNP assembly can be more

heavily impacted by structural fidelity than primary sequence recognition [59]. We

hypothesize that there is a core set of proteins required for SRE function, such as HuR, but

that individual SREs recruit distinct accessory factors that dictate the conditions or

mechanism by which that SRE protects against nuclease targeting. In this regard, we have

not found significant overlap between known HuR mRNA targets and mRNAs that are not

downregulated by SOX [22]. This is in agreement with the fact that many of the SRE bound

proteins have a breadth of roles in controlling RNA fate beyond conferring nuclease escape

[60-64]. SRE activity may be further impacted by the cellular context and neighboring

regulatory features, particularly in the case of tightly controlled transcripts. The 3' UTR of

IL-6, for example, is targeted by several cellular endonucleases including MCPIP1 [65-67],

which has recently been reported to be inhibited during de novo KSHV infection [68]. The

above features underscore the importance of characterizing multiple SREs to parse out

specific protein requirements, but also present challenges for the identification of SREs, as

they cannot be easily predicted. Indeed, it has taken more than a decade from the first

report of IL-6 escaping SOX-induced host shutoff to identify a second SRE-containing mRNA

[29].

The observation that the IL-6 SRE in particular acts against diverse viral

endonucleases but does not impact host endonucleases has important implications for

understanding target recognition by these host shutoff factors. First, it suggests that there

are key commonalities to how the viral endonucleases are recruited to mRNAs, and that

these features are distinct from those involved in cellular endonuclease recruitment. The

viral endonucleases recognize translation competent mRNAs, but do not require ongoing

translation for cleavage [10]. While vhs is recruited to a 5' cap proximal location due to its

interaction with a eIF4H, the SOX homologs do not respond to particular location cues on

their target mRNA, and instead recognize a degenerate target motif that can be positioned

anywhere [10,21,69-71]. PA-X has been suggested to bind RNA processing factors in the

nucleus, but how it recognizes cytoplasmic mRNA targets remains unknown [72]. By

contrast, the cellular NMD and no-go decay quality control pathways show a clear

dependence on translation for target recognition, and mRNA cleavage occurs at the site of

the error or translation stall [13]. The observation that neither the IL-6 nor GADD45B SREs

impede these cellular endonucleases indicates that SREs are not inhibitory to translation.

Furthermore, our FISH data do not support mRNA relocalization or sequestration as a

driving feature of SRE-mediated escape. We instead hypothesize that the SRE somehow

occludes one or more factors required for recruitment of the viral endonucleases. This

could occur via long-range interactions with mRNA cap-associated proteins, which we

previously showed can take place for the IL-6 SRE [27]. The fact that the GADD45B SRE

restricts against a more limited set of host shutoff factors argues against a single factor

requirement. However, it is possible that there are multiple binding sites on one factor that

are more comprehensively blocked by the IL-6 SRE. Independent of the mechanisms

involved, nuclease escape elements could be developed as tools to broadly inhibit viral

endonucleases without disrupting normal RNA decay pathways.

How might viruses benefit from maintaining the expression of IL-6 and GADD45B

during infection? The requirement for IL-6 during KSHV infection is well documented [30-

38], but the role of GADD45B may be more nuanced. While we demonstrated that it

escapes host shutoff during lytic KSHV infection and appears important for viral

reactivation in the iSLK.219 model, GADD45B expression is repressed by viral miRNAs

during latency to avoid its cell cycle arrest and pro-apoptotic functions [73]. However,

GADD45B has a number of additional activities that could be necessary for the viral lytic

cycle. These include promoting DNA demethylation and Retinoblastoma (Rb) inactivation,

processes that are important for a number of DNA viruses, including KSHV [74-79]. If our

hypothesis that there are numerous SREs throughout the transcriptome is correct, it is

likely that individual viruses require only a subset of these host genes for replication. Other

mRNAs may collaterally escape simply because they contain SREs that functionally mimic

those present in the 'required' mRNAs. In this regard, the fact that a given mRNA escapes

degradation by multiple viral endonucleases does not necessarily indicate that expression

of that gene is broadly required for infection. Finally, it is important to bear in mind that

the SREs discovered thus far are cellular elements that assemble cellular proteins—and

thus presumably play host-directed roles in the regulation of their associated transcripts. It

may therefore be the case that some SRE-bearing mRNAs function in an antiviral capacity.

Exploring possible virus-host evolutionary interplay for SREs remains an exciting prospect

for future studies.

MATERIALS AND METHODS

Cells and transfections. The KSHV-positive B cell line bearing a doxycycline-inducible

version of the major lytic transactivator RTA (TREX-BCBL-1) [55] was maintained in RPMI

medium (Invitrogen) supplemented with 10% fetal bovine serum (FBS; Invitrogen), 200

μM L-glutamine (Invitrogen), 100 U/ml penicillin/streptomycin (Invitrogen), and 50 μg/ml

hygromycin B (Omega Scientific). Lytic reactivation was induced by treatment with 20

ng/ml 2-O-tetradecanoylphorbol-13-acetate (TPA; Sigma), 1 μg/ml doxycycline (BD

Biosciences), and 500 ng/ml ionomycin (Fisher Scientific) for 48h. 293T cells (ATCC) were

grown in DMEM (Invitrogen) supplemented with 10% FBS. The KHSV-infected renal

carcinoma cell line iSLK.219 bearing doxycycline-inducible RTA was grown in DMEM

supplemented with 10% FBS [56]. KSHV lytic reactivation of the iSLK.219 cells was

induced by the addition of 0.2 μg/ml doxycycline (BD Biosciences) and 110 μg/ml sodium

butyrate for 48 h.

For DNA transfections, cells were plated and transfected after 24h when 70%

confluent using linear PEI (polyethylenimine). For small interfering RNA (siRNA)

transfections, 293T cells were reverse transfected in 12-well plates by INTERFERin

(Polyplus-Transfection) with 10 μM of siRNAs. siRNAs were obtained from IDT as DsiRNA

(siRNA SMG6: hs.Ri.SMG6.13.1; siRNA NCL: hs.Ri.NCL.13.1; siRNA HuR ELAVL1#1:

hs.Ri.ELAVL1.13.2; siRNA ELAVL1#2: hs.Ri.ELAVL1.13.3). 48h following siRNA

transfection, the cells subjected to DNA transfection as indicated.

Plasmids. The GADD45B 5'UTR and CDS were obtained as G-blocks from IDT and cloned

into a pcDNA3.1 plasmid downstream of the GFP coding sequence. GADD45B 3'UTR was

cloned from a pDest-765 plasmid (kindly provided by J. Ziegelbauer) into a pcDNA3.1

plasmid downstream of the GFP coding sequence. The GFP-IL-6 3'UTR, SRE and ΔSRE

fusion constructs were described previously [27]. The dsRed2 and dsRed2-PTC reporters

were described elsewhere [10]. The SRE and ΔSRE were PCR amplified from the GFP

reporters and cloned downstream of the dsRed2 ORF.

Point mutations were introduced with the Quickchange site directed mutagenesis

protocol (Agilent) using the primers described in **Table S3**.

FiSH. Stellaris FiSH probes recognizing MS2 and GAPDH labeled with Quasar 570 Dye

(MS2: SMF-1063-5; GAPDH: SMF-2026-1) were hybridized in 293T cells following

manufacturer's instructions available online at www.biosearchtech.com/stellarisprotocols.

Briefly, 293T cells were grown on coverslips and transfected with either an empty vector

control or the MS2-SRE or MS2ΔSRE constructs. 24h later, cells were washed, fixed in 4%

formaldehyde and permeabilized in 70% ethanol. Probes (12.5μM) were then hybridized

for >5h at 37°C in Vanadyl ribonucleoside (10mM), formamide (10%), saline sodium citrate

(SSC), dextran (10%) and BSA (0.2%). DAPI was added for the last hour to stain cell nuclei.

Coverslips were washed in SSC and mounted in Vectashield mounting medium

(VectorLabs) before visualization by confocal microscopy on a Zeiss LSM 710 AxioObserver

microscope.

Comprehensive Analysis of RNA Binding Proteins (ChIRP). ChIRP was carried out

according to a protocol published previously [58] with minor modifications. Briefly, 293T

cells were transfected with the GFP-GADD45B 3'UTR plasmid and 24h later cells were

crosslinked in 3% formaldehyde for 30 minutes, quenched in 125mM Glycine, washed with

PBS and the pellet was flash frozen. Pellets were then lysed in fresh ChIRP Lysis Buffer

[Tris pH 7.4 50mM, EDTA 10mM, SDS 1%], sonicated, and cell debris was removed by

centrifugation at (14,000rpm for 10 min). Two mL of hybridization buffer was added to

each sample along with 100pmol of the indicated probes (see **Table S3** for sequences).

Hybridization was carried out for 4-12h. After adding streptavidin beads to the samples for

1h, samples were washed and reverse crosslinked at 65°C overnight. For western blotting,

beads were resuspended in 4X loading buffer, boiled for 30 minutes and resolved by SDS-

PAGE. For mass spectrometry, recovered proteins from two independent biological

replicates were gel extracted and subjected to LC-MS/MS at the UC Berkeley Vincent J.

Coates Proteomics/Mass Spectrometry Laboratory.

In-line probing. RNA was synthesized by Dharmacon (SRE-HP #CTM-299208, GADD45-

HP- #CTM-299209). Approximately 1ug of RNA was 5' end-labeled and purified on an 8M

urea gel. Samples were then ethanol precipitated in presence of glycogen, washed in 70%

ethanol and dissolved in 20 μL of DEPC-treated water. Reactions were carried out with

~1uL of purified RNA (~100,000cpm). In-line probing was performed following a standard

procedure [57]. Briefly, in-line probing assays were carried for 40h in 2X in-line buffer

[100 mM Tris·HCl, pH 8.3, 40 mM MgCl2, 200 mM KCl] and quenched by adding the same

volume of RNA loading Buffer [95% formamide, 10 mM EDTA and 0.025% xylene cyanol].

The no-reaction (NR) treatment, RNase T1 (T1) and partial base hydrolysis (–OH) ladders

were prepared as 20μL reactions and quenched with 20 μL RNA loading buffer. Dried gels

were exposed on a phosphorimager screen and scanned using a Typhoon laser-scanning

system (GE Healthcare).

RT-qPCR. Total RNA was harvested using Trizol following the manufacture's protocol.

cDNAs were synthesized from 1 μg of total RNA using AMV reverse transcriptase

(Promega), and used directly for quantitative PCR (qPCR) analysis with the DyNAmo

ColorFlash SYBR green qPCR kit (Thermo Scientific). Signals obtained by qPCR were

normalized to 18S.

RNA Immunoprecipitation. Cells were crosslinked in 1% formaldehyde for 10 minutes,

quenched in 125mM glycine and washed in PBS. Cells were then lysed in low-salt lysis

buffer [NaCl 150mM, NP-40 0.5%, Tris pH8 50mM, DTT 1mM, MgCl2 3mM containing

protease inhibitor cocktail and RNase inhibitor] and sonicated. After removal of cell debris,

specific antibodies were added as indicated overnight at 4°C. Magnetic G-coupled beads

were added for 1h, washed three times with lysis buffer and twice with high-salt lysis

buffer (low-salt lysis buffer except containing 400mM NaCl). Samples were separated into

two fractions. Beads containing the fraction used for western blotting were resuspended in

30μL lysis buffer. Beads containing the fraction used for RNA extraction were resuspended

in Proteinase K buffer (NaCl 100mM, Tris pH 7.4 10mM, EDTA 1mM, SDS 0.5%) containing

1μL of PK (Proteinase K). Samples were incubated overnight at 65°C to reverse

crosslinking. Samples to be analyzed by western blot were then supplemented with 10μL of

4X loading buffer before resolution by SDS-PAGE. RNA samples were resuspend in Trizol

and were processed as described above.

Western Blotting. Lysates were resolved by SDS-PAGE and western blotted with the

following antibodies at 1:1000 in TBST (Tris-buffered saline, 0.1% Tween 20): rabbit anti-

EST1A/SMG6 (Abcam), rabbit anti-NCL (Abcam), rabbit anti-HuR (Millipore), rabbit anti-

actin (Abcam), rabbit anti-IGF2BP1 (Abcam), rabbit anti-YTHDC2 (Abcam), rabbit anti-

hnRNPU (Abcam), rabbit anti-ZC3HAV1 (Abcam), rabbit anti-hnRNPD (Abcam), rabbit anti-

STAU1 (Pierce), mouse anti-NPM1 (Abcam), rabbit anti-GADD45 (Abcam), mouse anti-

GAPDH (Abcam), mouse anti-ORF59 (Adv biotechnologies), or rabbit anti-K8.1 (PRF&L,

inc). Primary antibody incubations were followed by HRP-conjugated goat anti-mouse or

goat anti-rabbit secondary antibodies (Southern Biotechnology, 1:5000).

Statistical analysis. All results are expressed as means ± S.E.M. of experiments

independently repeated at least three times. Unpaired Student's t test was used to evaluate

the statistical difference between samples. Significance was evaluated with P values as

follows: * p<0.1; ** p<0.05; *** p<0.01.

ACKNOWLEDGMENTS

We thank members of the Glaunsinger and Coscoy labs for helpful discussions. We also

thank Ming Hammond, Debojit Bose and Carolin Vogt for technical advice on in-line

probing.

FUNDING

This research was supported by NIH grants CA160556 and CA136367

(http://www.nih.gov/) and a Burroughs Wellcome Investigator in the Pathogenesis of

Infectious Diseases Award (http:// www.bwfund.org/grant-programs/infectious-diseases/

investigators-in-pathogenesis-of-infectious-disease) to BAG. B.A.G. is an investigator of the

Howard Hughes Medical Institute. Funding for open access charge: Howard Hughes

Medical Institute and MM is a HHMI Fellow of the Damon Runyon Cancer Research

Foundation (DRG- 2207-14) (http://www.damonrunyon.org/). This work used the Vincent

J. Proteomics/Mass Spectrometry Laboratory at UC Berkeley, supported in part by NIH S10

Instrumentation Grant S10RR025622.

REFERENCES

1. Gaglia MM, Covarrubias S, Wong W, Glaunsinger BA. A common strategy for host RNA degradation by divergent viruses. J Virol. 2012;86: 9527–9530.

2. Rivas HG, Schmaling SK, Gaglia MM. Shutoff of Host Gene Expression in Influenza A Virus and Herpesviruses: Similar Mechanisms and Common Themes. Viruses. 2016;8: 102.

3. Jagger BW, Wise HM, Kash JC, Walters K-A, Wills NM, Xiao Y-L, et al. An overlapping protein-coding region in influenza A virus segment 3 modulates the host response. Science. 2012;337: 199–204.

4. Kwong AD, Frenkel N. Herpes simplex virus-infected cells contain a function(s) that destabilizes both host and viral mRNAs. Proc Natl Acad Sci USA. 1987;84: 1926–1930.

5. Kamitani W, Huang C, Narayanan K, Lokugamage KG, Makino S. A two-pronged strategy to suppress host protein synthesis by SARS coronavirus Nsp1 protein. Nat Struct Mol Biol. 2009;16: 1134–1140.

6. Parrish S, Moss B. Characterization of a vaccinia virus mutant with a deletion of the D10R gene encoding a putative negative regulator of gene expression. J Virol. 2006;80: 553–561.

7. Parrish S, Moss B. Characterization of a second vaccinia virus mRNA-decapping enzyme conserved in poxviruses. J Virol. 2007;81: 12973–12978.

8. Parrish S, Resch W, Moss B. Vaccinia virus D10 protein has mRNA decapping activity, providing a mechanism for control of host and viral gene expression. Proc Natl Acad Sci USA. 2007;104: 2139–2144.

9. Elgadi MM, Hayes CE, Smiley JR. The herpes simplex virus vhs protein induces endoribonucleolytic cleavage of target RNAs in cell extracts. J Virol. 1999;73: 7153–7164.

10. Covarrubias S, Gaglia MM, Kumar GR, Wong W, Jackson AO, Glaunsinger BA. Coordinated destruction of cellular messages in translation complexes by the gammaherpesvirus host shutoff factor and the mammalian exonuclease Xrn1. PLoS Pathog. 2011 7(10): e1002339

11. Burgess HM, Mohr I. Cellular 5"-3" mRNA exonuclease Xrn1 controls double-stranded RNA accumulation and anti-viral responses. Cell Host Microbe. 2015;17:

332–344.

12.   Schoenberg DR, Maquat LE. Regulation of cytoplasmic mRNA decay. Nat Rev Genet.
2012;13: 246–259.

13.   Mühlemann O, Lykke-Andersen J. How and where are nonsense mRNAs degraded in
mammalian cells? RNA Biol. 2010;7: 28–32.

14.   Doma MK, Parker R. Endonucleolytic cleavage of eukaryotic mRNAs with stalls in
translation elongation. Nature. 2006;440: 561–564.

15.   Harigaya Y, Parker R. No-go decay: a quality control mechanism for RNA in
translation. Wiley Interdiscip Rev RNA. 2010;1: 132–141.

16.   Covarrubias S, Richner JM, Clyde K, Lee YJ, Glaunsinger BA. Host shutoff is a
conserved phenotype of gammaherpesvirus infection and is orchestrated exclusively
from the cytoplasm. J Virol. 2009;83: 9554–9566.

17.   Rowe M, Glaunsinger B, van Leeuwen D, Zuo J, Sweetman D, Ganem D, et al. Host
shutoff during productive Epstein-Barr virus infection is mediated by BGLF5 and
may contribute to immune evasion. Proc Natl Acad Sci USA. 2007;104: 3366–3371.

18.   Glaunsinger B, Ganem D. Lytic KSHV infection inhibits host gene expression by
accelerating global mRNA turnover. Mol Cell. 2004;13: 713–723.

19.   Abernathy E, Clyde K, Yeasmin R, Krug LT, Burlingame A, Coscoy L, et al.
Gammaherpesviral gene expression and virion composition are broadly controlled
by accelerated mRNA degradation. PLoS Pathog. 2014;10: e1003882.

20.   Richner JM, Clyde K, Pezda AC, Cheng BYH, Wang T, Kumar GR, et al. Global mRNA
degradation during lytic gammaherpesvirus infection contributes to establishment of
viral latency. PLoS Pathog. 2011;7: e1002150.

21.   Gaglia MM, Rycroft CH, Glaunsinger BA. Transcriptome-Wide Cleavage Site Mapping
on Cellular mRNAs Reveals Features Underlying Sequence-Specific Cleavage by the
Viral Ribonuclease SOX. PLoS Pathog. 2015;11: e1005305.

22.   Clyde K, Glaunsinger BA. Deep sequencing reveals direct targets of
gammaherpesvirus-induced mRNA decay and suggests that multiple mechanisms
govern cellular transcript escape. PLoS ONE. 2011;6: e19655.

23.   Chandriani S, Ganem D. Host transcript accumulation during lytic KSHV infection
reveals several classes of host responses. PLoS ONE. 2007;2: e811.

24.   Clyde K, Glaunsinger BA. Getting the message direct manipulation of host mRNA
accumulation during gammaherpesvirus lytic infection. Adv Virus Res. 2010;78: 1–
42.

25.   Lee YJ, Glaunsinger BA. Aberrant herpesvirus-induced polyadenylation correlates
599   with cellular messenger RNA destruction. PLoS Biol. 2009;7: e1000107.

600   26.   Hutin S, Lee Y, Glaunsinger BA. An RNA element in human interleukin 6 confers
601   escape from degradation by the gammaherpesvirus SOX protein. J Virol. 2013;87:
602   4672–4682.

603   27.   Muller M, Hutin S, Marigold O, Li KH, Burlingame A, Glaunsinger BA. A
604   ribonucleoprotein complex protects the interleukin-6 mRNA from degradation by
605   distinct herpesviral endonucleases. PLoS Pathog. 2015;11: e1004899.

606   28.   Abernathy E, Gilbertson S, Alla R, Glaunsinger B. Viral Nucleases Induce an mRNA
607   Degradation-Transcription Feedback Loop in Mammalian Cells. Cell Host Microbe.
608   2015;18: 243–253.

609   29.   Glaunsinger B, Ganem D. Highly selective escape from KSHV-mediated host mRNA
610   shutoff and its implications for viral pathogenesis. J Exp Med. 2004;200: 391–398.

611   30.   Sin S-H, Roy D, Wang L, Staudt MR, Fakhari FD, Patel DD, et al. Rapamycin is
612   efficacious against primary effusion lymphoma (PEL) cell lines in vivo by inhibiting
613   autocrine signaling. Blood. 2007;109: 2165–2173.

614   31.   Miles SA, Rezai AR, Salazar-González JF, Vander Meyden M, Stevens RH, Logan DM, et
615   al. AIDS Kaposi sarcoma-derived cells produce and respond to interleukin 6. Proc
616   Natl Acad Sci USA. 1990;87: 4068–4072.

617   32.   Screpanti I, Musiani P, Bellavia D, Cappelletti M, Aiello FB, Maroder M, et al.
618   Inactivation of the IL-6 gene prevents development of multicentric Castleman's
619   disease in C/EBP beta-deficient mice. J Exp Med. 1996;184: 1561–1566.

620   33.   Leger-Ravet MB, Peuchmaur M, Devergne O, Audouin J, Raphael M, Van Damme J, et
621   al. Interleukin-6 gene expression in Castleman's disease. Blood. 1991;78: 2923–2930.

622   34.   Xie J, Pan H, Yoo S, Gao S-J. Kaposi's sarcoma-associated herpesvirus induction of AP-
623   1 and interleukin 6 during primary infection mediated by multiple mitogen-activated
624   protein kinase pathways. J Virol. 2005;79: 15027–15037.

625   35.   An J, Sun Y, Sun R, Rettig MB. Kaposi's sarcoma-associated herpesvirus encoded
626   vFLIP induces cellular IL-6 expression: the role of the NF-kappaB and JNK/AP1
627   pathways. Oncogene. 2003;22: 3371–3385.

628   36.   Santarelli R, Gonnella R, Di Giovenale G, Cuomo L, Capobianchi A, Granato M, et al.
629   STAT3 activation by KSHV correlates with IL-10, IL-6 and IL-23 release and an
630   autophagic block in dendritic cells. Sci Rep. 2014;4: 4241.

631   37.   Deng H, Chu JT, Rettig MB, Martinez-Maza O, Sun R. Rta of the human herpesvirus
632   8/Kaposi sarcoma-associated herpesvirus up-regulates human interleukin-6 gene

633   expression. Blood. 2002;100: 1919–1921.

634   38.   McCormick C, Ganem D. The kaposin B protein of KSHV activates the p38/MK2
635   pathway and stabilizes cytokine mRNAs. Science. 2005;307: 739–741.

636   39.   Oishi K, Yamayoshi S, Kawaoka Y. Mapping of a Region of the PA-X Protein of
637   Influenza A Virus That Is Important for Its Shutoff Activity. J Virol. 2015;89: 8661–
638   8665.

639   40.   Smiley JR, Elgadi MM, Saffran HA. Herpes simplex virus vhs protein. Meth Enzymol.
640   2001;342: 440–451.

641   41.   Everly DN, Feng P, Mian IS, Read GS. mRNA degradation by the virion host shutoff
642   (Vhs) protein of herpes simplex virus: genetic and biochemical evidence that Vhs is a
643   nuclease. J Virol. 2002;76: 8560–8571.

644   42.   Aizer A, Kalo A, Kafri P, Shraga A, Ben-Yishay R, Jacob A, et al. Quantifying mRNA
645   targeting to P-bodies in living human cells reveals their dual role in mRNA decay and
646   storage. J Cell Sci. 2014;127: 4443–4456.

647   43.   Abernathy E, Glaunsinger B. Emerging roles for RNA degradation in viral replication
648   and antiviral defense. Virology. 2015;479-480: 600–608.

649   44.   Isken O, Maquat LE. Quality control of eukaryotic mRNA: safeguarding cells from
650   abnormal mRNA function. Genes Dev. 2007;21: 1833–1856.

651   45.   Schoenberg DR. Mechanisms of endonuclease-mediated mRNA decay. Wiley
652   Interdiscip Rev RNA. 2011;2: 582–600.

653   46.   Eberle AB, Lykke-Andersen S, Mühlemann O, Jensen TH. SMG6 promotes
654   endonucleolytic cleavage of nonsense mRNA in human cells. Nat Struct Mol Biol.
655   2009;16: 49–55.

656   47.   Narayanan K, Ramirez SI, Lokugamage KG, Makino S. Coronavirus nonstructural
657   protein 1: Common and distinct functions in the regulation of host and viral gene
658   expression. Virus Res. 2015;202: 89–100.

659   48.   Huang C, Lokugamage KG, Rozovics JM, Narayanan K, Semler BL, Makino S. SARS
660   coronavirus nsp1 protein induces template-dependent endonucleolytic cleavage of
661   mRNAs: viral mRNAs are resistant to nsp1-induced RNA cleavage. PLoS Pathog.
662   2011;7: e1002433.

663   49.   Zhang Y, Bhatia D, Xia H, Castranova V, Shi X, Chen F. Nucleolin links to arsenic-
664   induced stabilization of GADD45alpha mRNA. Nucleic Acids Res. 2006;34: 485–495.

665   50.   Zhang D, Liang Y, Xie Q, Gao G, Wei J, Huang H, et al. A novel post-translational
666   modification of nucleolin, SUMOylation at Lys-294, mediates arsenite-induced cell

667   death by regulating gadd45α mRNA stability. J Biol Chem. 2015;290: 4784–4800.

668   51.   Zheng X, Zhang Y, Chen Y-Q, Castranova V, Shi X, Chen F. Inhibition of NF-kappaB
669   stabilizes gadd45alpha mRNA. Biochem Biophys Res Commun. 2005;329: 95–99.

670   52.   Esclatine A, Taddeo B, Evans L, Roizman B. The herpes simplex virus 1 UL41 gene-
671   dependent destabilization of cellular RNAs is selective and may be sequence-specific.
672   Proc Natl Acad Sci USA. 2004;101: 3603–3608.

673   53.   Barzilai A, Zivony-Elbom I, Sarid R, Noah E, Frenkel N. The herpes simplex virus type
674   1 vhs-UL41 gene secures viral replication by temporarily evading apoptotic cellular
675   response to infection: Vhs-UL41 activity might require interactions with elements of
676   cellular mRNA degradation machinery. J Virol. 2006;80: 505–513.

677   54.   Khodarev NN, Advani SJ, Gupta N, Roizman B, Weichselbaum RR. Accumulation of
678   specific RNAs encoding transcriptional factors and stress response proteins against a
679   background of severe depletion of cellular RNAs in cells infected with herpes simplex
680   virus 1. Proc Natl Acad Sci USA. 1999;96: 12062–12067.

681   55.   Nakamura H, Lu M, Gwack Y, Souvlis J, Zeichner SL, Jung JU. Global changes in
682   Kaposi's sarcoma-associated virus gene expression patterns following expression of
683   a tetracycline-inducible Rta transactivator. J Virol. 2003;77: 4205–4220.

684   56.   Myoung J, Ganem D. Generation of a doxycycline-inducible KSHV producer cell line of
685   endothelial origin: maintenance of tight latency with efficient reactivation upon
686   induction. J Virol Methods. 2011;174: 12–21.

687   57.   Regulski EE, Breaker RR. In-line probing analysis of riboswitches. Methods Mol Biol.
688   2008;419: 53–67.

689   58.   Chu C, Zhang QC, da Rocha ST, Flynn RA, Bharadwaj M, Calabrese JM, et al. Systematic
690   discovery of Xist RNA binding proteins. Cell. 2015;161: 404–416.

691   59.   Bevilacqua PC, Ritchey LE, Su Z, Assmann SM. Genome-Wide Analysis of RNA
692   Secondary Structure. Annu Rev Genet. 2016;50: 235–266.

693   60.   Dean JLE, Sully G, Clark AR, Saklatvala J. The involvement of AU-rich element-binding
694   proteins in p38 mitogen-activated protein kinase pathway-mediated mRNA
695   stabilisation. Cell Signal. 2004;16: 1113–1121.

696   61.   Abdelmohsen K, Gorospe M. RNA-binding protein nucleolin in disease. RNA Biol.
697   2012;9: 799–808.

698   62.   Bell JL, Wächter K, Mühleck B, Pazaitis N, Köhn M, Lederer M, et al. Insulin-like
699   growth factor 2 mRNA-binding proteins (IGF2BPs): post-transcriptional drivers of
700   cancer progression? Cell Mol Life Sci. 2013;70: 2657–2675.

701   63.   White EJF, Brewer G, Wilson GM. Post-transcriptional control of gene expression by
702   AUF1: mechanisms, physiological targets, and regulation. Biochim Biophys Acta.
703   2013;1829: 680–688.

704   64.   Park E, Maquat LE. Staufen-mediated mRNA decay. Wiley Interdiscip Rev RNA.
705   2013;4: 423–435.

706   65.   Matsushita K, Takeuchi O, Standley DM, Kumagai Y, Kawagoe T, Miyake T, et al.
707   Zc3h12a is an RNase essential for controlling immune responses by regulating mRNA
708   decay. Nature. 2009;458: 1185–1190.

709   66.   Paschoud S, Dogar AM, Kuntz C, Grisoni-Neupert B, Richman L, Kühn LC.
710   Destabilization of interleukin-6 mRNA requires a putative RNA stem-loop structure,
711   an AU-rich element, and the RNA-binding protein AUF1. Mol Cell Biol. 2006;26:
712   8228–8241.

713   67.   Mino T, Murakawa Y, Fukao A, Vandenbon A, Wessels H-H, Ori D, et al. Regnase-1 and
714   Roquin Regulate a Common Element in Inflammatory mRNAs by Spatiotemporally
715   Distinct Mechanisms. Cell. 2015;161: 1058–1073.

716   68.   Happel C, Ramalingam D, Ziegelbauer JM. Virus-Mediated Alterations in miRNA
717   Factors and Degradation of Viral miRNAs by MCPIP1. PLoS Biol. 2016;14: e2000998.

718   69.   Feng P, Everly DN, Read GS. mRNA decay during herpes simplex virus (HSV)
719   infections: protein-protein interactions involving the HSV virion host shutoff protein
720   and translation factors eIF4H and eIF4A. J Virol. 2005;79: 9651–9664.

721   70.   Page HG, Read GS. The virion host shutoff endonuclease (UL41) of herpes simplex
722   virus interacts with the cellular cap-binding complex eIF4F. J Virol. 2010;84: 6886–
723   6890.

724   71.   Feng P, Everly DN, Read GS. mRNA decay during herpesvirus infections: interaction
725   between a putative viral nuclease and a cellular translation factor. J Virol. 2001;75:
726   10272–10280.

727   72.   Khaperskyy DA, Schmaling S, Larkins-Ford J, McCormick C, Gaglia MM. Selective
728   Degradation of Host RNA Polymerase II Transcripts by Influenza A Virus PA-X Host
729   Shutoff Protein. PLoS Pathog. 2016;12: e1005427.

730   73.   Liu X, Happel C, Ziegelbauer JM. Kaposi's Sarcoma-Associated Herpesvirus
731   MicroRNAs Target GADD45B To Protect Infected Cells from Cell Cycle Arrest and
732   Apoptosis. J Virol. 2017;91: e02045–16.

733   74.   Cho HJ, Park S-M, Hwang EM, Baek KE, Kim I-K, Nam I-K, et al. Gadd45b mediates
734   Fas-induced apoptosis by enhancing the interaction between p38 and
735   retinoblastoma tumor suppressor. J Biol Chem. 2010;285: 25500–25505.

736   75.   Hume AJ, Kalejta RF. Regulation of the retinoblastoma proteins by the human
737   herpesviruses. Cell Div. 2009;4: 1.

738   76.   Godden-Kent D, Talbot SJ, Boshoff C, Chang Y, Moore P, Weiss RA, et al. The cyclin
739   encoded by Kaposi's sarcoma-associated herpesvirus stimulates cdk6 to
740   phosphorylate the retinoblastoma protein and histone H1. J Virol. 1997;71: 4193–
741   4198.

742   77.   Chang Y, Moore PS, Talbot SJ, Boshoff CH, Zarkowska T, Godden-Kent, et al. Cyclin
743   encoded by KS herpesvirus. Nature. 1996;382: 410–410.

744   78.   Ma DK, Guo JU, Ming G-L, Song H. DNA excision repair proteins and Gadd45 as
745   molecular players for active DNA demethylation. Cell Cycle. 2009;8: 1526–1531.

746   79.   Purushothaman P, Uppal T, Verma SC. Molecular biology of KSHV lytic reactivation.
747   Viruses. 2015;7: 116–153.

 FIGURE LEGENDS
Figure 1: The IL-6-SRE broadly protects against multiple viral endonucleases. (A)
293T cells were transfected with empty vector (mock) or a plasmid expressing the
indicated endonucleases along with a GFP reporter. After 24 h, total RNA was harvested
and subjected RT-qPCR to measure GFP mRNA levels. Graphs here and afterwards display
individual replicates as dots, together with the mean values (±SEM). Statistical significance
was determined by the Student t test (* $p<0.1$; ** $p<0.05$; *** $p<0.01$). **(B)** 293T cells were
co-transfected with the indicated endonuclease-expressing plasmid expressing together
with a GFP-SRE or GFP-ΔSRE reporter. After 24 h, total RNA was harvested and subjected
RT-qPCR to measure GFP mRNA levels. **(C)** Top: diagram showing the structure of the
reporter mRNA containing MS2 repeats upstream of the SRE or ΔSRE fragment of the IL-6
3'UTR. Red lines denote the region detected by the MS2 probe. Bottom: 293T cells were
transfected with the indicated MS2 reporters. 24h later, cells were fixed and processed for
RNA FISH staining. Signals from the MS2 probes (red) and DAPI stained nuclei (blue) were
detected by confocal microscopy.
764
Figure 2: The IL-6-SRE does not protect against cellular endonucleases. (A) 293T cells
were transfected with the indicated dsRed2 reporters. After 24 h, total RNA was harvested
and subjected RT-qPCR to measure dsRed mRNA levels. **(B)** 293T cells were treated with
siRNAs targeting Smg6 or control non-targeting siRNAs. 48h later, cells were transfected
with the indicated dsRed reporters. After 24h, total RNA was harvested and subjected RT-
qPCR to measure dsRed mRNA levels (*left*). The efficiency of SMG6 knockdown was
measured by western blotting, with actin serving as a loading control (*right*). **(C)** 293T cells
were transfected with the indicated dsRed2 reporters. After 24 h, total RNA was harvested
and subjected to RT-qPCR to measure dsRed mRNA levels. **(D)** 293T cells were transfected
with an empty vector (mock) or a plasmid expressing SCoV nsp1 along with the indicated
GFP reporters. After 24 h, total RNA was harvested and subjected RT-qPCR to measure GFP
mRNA levels.
777
Figure 3: GADD45B mRNA is protected against SOX degradation. (A, B) Total RNA was
extracted from unreactivated or reactivated KSHV-positive TREX-BCBL-1 (A) cells or
iSLK.219 cells (B) and subjected to RT-qPCR to measure endogenous levels of the
GADD45B or GAPDH transcript. **(C)** 293T cells were transfected with an empty vector or a
plasmid expressing SOX or vhs. After 24 h, total RNA was harvested and subjected to RT-
qPCR to measure endogenous levels of GADD45B or GAPDH transcripts. **(D)** Diagram of the
reporter constructs where fragments of GADD45B transcript were fused to GFP. **(E)** 293T
cells were transfected with the indicated GFP-GADD45B fusion constructs in the presence
or absence of a plasmid expressing SOX. After 24 h, total RNA was harvested and subjected
to RT-qPCR to measure GFP levels.
788
Figure 4: The SREs contain a long hairpin required for protection against SOX. (A)
293T cells were transfected with the indicated GFP reporter along with a control empty
vector (mock) or a plasmid expressing SOX. After 24 h, total RNA was harvested and
subjected to RT-qPCR to measure GFP mRNA levels. **(B)** Diagram of the structure
prediction obtained with RNAfold for IL-6 and GADD45B SREs. The color scale represents

the confidence score of the structure as calculated by RNAfold, with red representing the
highest confidence. Asterisks denote the location of mutations that were introduced in the
structure for the following assays. The insets are RNAfold predictions of the stem loops of
interest in isolation. **(C)** 293T cells were transfected with the indicated GFP reporters
containing mutations within IL-6 SRE at the residues marked by a * in (B) along with a
control empty vector (mock) or a plasmid expressing SOX. After 24 h, total RNA was
harvested and subjected RT-qPCR to measure GFP mRNA levels. **(D)** 293T cells were
transfected with the indicated GFP reporters mutated within G- SRE at the residues marked
by a * in (B) along with a control empty vector (mock) or a plasmid expressing SOX. After
24 h, total RNA was harvested and subjected to RT-qPCR to measure GFP mRNA levels.

Figure 5: The IL-6 and GADD45B SRE bind a partially overlapping set of cellular
proteins. **(A)** Diagram outlining the ChIRP assay. **(B)** Cytoscape network showing the
proteins reported to interact with the IL-6 SRE from a previous screen [27] (gray nodes)
overlaid with the set of proteins that were also recovered by ChIRP-MS for the IL-6 and
GADD45B 3' UTRs (purple nodes) **(C)** 293T were transfected with the GFP-GADD45B
3'UTR reporter, then 24h later they were subjected to ChIRP analysis and protein samples
were western blotted. **(D)** Crosslinked lysates of 293T cells were subjected to RNA
immunoprecipitation (RIP) with control IgG, anti-NCL, or anti-HuR antibodies and the level
of co-purifying endogenous GADD45B mRNA was quantified by RT-qPCR. Bars represent
the fold enrichment over the mock IP. **(E)** 293T cells were transfected with either the GFP-
GADD45B 5'UTR, GFP-GADD45B 3'UTR, or GFP-GADD45B 3'UTR_GG reporter for 24 h.
Lysates were then subjected to RIP as described in (C). Bars represent the fold enrichment
over mock IP.
817
Figure 6: The GADD45B SRE function requires HuR, but is only protective against
SOX and vhs. (**A, B**) 293T cells were treated with siRNAs targeting HuR (A) or NCL (B) or
control non targeting siRNAs. 48h later, cells were transfected with the GFP-GADD45B-
3'UTR reporter along with a control empty vector or a plasmid expressing SOX. After 24 h,
total RNA was harvested and subjected to RT-qPCR to measure GFP mRNA levels. Protein
levels of HuR and NCL after siRNA-mediated depletion are shown under the bar graphs
(~30% and 45% of total protein left respectively) **(C)** 293T cells were transfected with the
GFP-GADD45B-3'UTR reporter along with a control empty vector or a plasmid expressing
the indicated viral endonucleases. After 24 h, total RNA was harvested and subjected to RT-
qPCR to measure GFP mRNA levels.
828
829
830
831 ## SUPPORTING INFORMATION
832
833 **Figure S1:** 293T cells were transfected with the indicated MS2 luciferase reporter
834 containing or lacking the IL-6 SRE +/- a KSHV SOX expression plasmid. After 24 h, total
835 RNA was harvested and subjected RT-qPCR to measure reporter mRNA levels.
836
Figure S2: (A) iSLK.219 cells were treated with siRNAs targeting GADD45B (or control
non-target siRNAs) for 48h. Cells were then reactivated with doxycycline and sodium

butyrate for 24 or 48h, fixed, and reactivation efficiency was monitored by expression of
red fluorescent protein, which is expressed from the viral genome under the control of the
lytic PAN promoter. **(B)** After siRNA treatment and reactivation as described in A, cell
lysates were subjected to western blotting to measure protein levels of GADD45B, the
KSHV proteins K8.1 and ORF59, and GAPDH (as a loading control).
844
845 **Figure S3:** Clustal W alignment of IL-6 SRE and GADD45B 3'UTR.
846
Figure S4: In-line probing of the IL-6 (left) and GADD45B (right) predicted stem-loop. ^{32}P-
labeled RNA (NR, no reaction) and products resulting from partial digestion with nuclease
T1 (T1; cuts after G residues), partial digestion with alkali (-OH), and spontaneous cleavage
during a 24h incubation are shown. Product bands corresponding to G residues (generated
by T1 digestion) are labeled with black arrows. Predicted paired or unpaired residues are
marked on the right of each gel and are shown on the RNA fold diagrams.
853
854 **Figure S5:** Network representation of the full set of proteins identified by ChIRP-MS for
855 either the IL-6 or GADD45B 3'UTR. Purple nodes represent proteins that were previously
856 identified using an *in vitro* pulldown/MS-based assay [27].
857
858
859 **Table S1:** List of proteins identified by ChIRP-MS. Proteins in bold were in common
860 between the IL-6 and GADD45B 3' UTR datasets. Background proteins found in the control
861 ChIRP-MS runs with probes were filtered out together with common contaminants.
862
863 **Table S2**: List of proteins identified by ChIRP-MS that are in common with previously
864 identified SRE-binding proteins [27].
865
866 **Table S3:** List of primers used in this study

A

Relative GFP mRNA levels

Mock SOX muSOX BGLF5 vhs PA-X

B

Relative GFP mRNA levels

Mock SOX vhs muSOX BGLF5 PAX   Mock SOX vhs muSOX BGLF5 PAX

GFP-SRE   GFP-ΔSRE

***

C

Cy3 MS2 probes

MS2 repeats   IL-6 SRE or ΔSRE

MS2-SRE   MS2-ΔSRE

MS2 probes   no probe