Title: A Bayesian Framework for Multiple Trait Colocalization from Summary Association Statistics

Claudia Giambartolomei[1], Jimmy Zhenli Liu[2], Wen Zhang[3], Mads Hauberg[3,4], Huwenbo Shi[5], James Boocock[1], Joe Pickrell[2], Andrew E. Jaffe[6], the CommonMind Consortium[#], Bogdan Pasaniuc*[1], Panos Roussos*[3,7,8]

[1]Department of Pathology and Laboratory Medicine, University of California, Los Angeles, Los Angeles, CA 90095, USA; Department of Human Genetics, University of California, Los Angeles, Los Angeles, CA 90095, United States of America.
[2] New York Genome Center, New York, New York, United States of America
[3]Department of Genetics and Genomic Science and Institute for Multiscale Biology, Icahn School of Medicine at Mount Sinai, New York, New York, 10029, United States of America.
[4]The Lundbeck Foundation Initiative of Integrative Psychiatric Research (iPSYCH), Aarhus University, Aarhus, 8000, Denmark.
[5]Bioinformatics Interdepartmental Program, University of California, Los Angeles, 90024
[6]Lieber Institute for Brain Development, Johns Hopkins Medical Campus; Departments of Mental Health and Biostatistics, Johns Hopkins Bloomberg School of Public Health Baltimore, MD, 21205, United States of America.
[7]Department of Psychiatry and Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, New York, 10029, United States of America.
[8]Mental Illness Research Education and Clinical Center (MIRECC), James J. Peters VA Medical Center, Bronx, New York, 10468, United States of America.

The CommonMind Consortium includes: Menachem Fromer, Panos Roussos, Solveig K Sieberts, Jessica S Johnson, Douglas M Ruderfer, Hardik R Shah, Lambertus L Klei, Kristen K Dang, Thanneer M Perumal, Benjamin A Logsdon, Milind C Mahajan, Lara M Mangravite, Hiroyoshi Toyoshiba, Raquel E Gur, Chang-Gyu Hahn, Eric Schadt, David A Lewis, Vahram Haroutunian, Mette A Peters, Barbara K Lipska, Joseph D Buxbaum, Keisuke Hirai, Enrico Domenici, Bernie Devlin, Pamela Sklar

Correspondence:
Dr. Panos Roussos
Icahn School of Medicine at Mount Sinai, Institute for Multiscale Biology
Department of Psychiatry and Department of Genetics and Genomic Science and
One Gustave L. Levy Place,
New York, NY, 10029, USA
panagiotis.roussos@mssm.edu

Dr. Claudia Giambartolomei
University of California, Los Angeles, Los Angeles ,
Department of Pathology and Laboratory Medicine, Los Angeles, CA 90095, USA
claudia.giambartolomei@gmail.com

ABSTRACT

Most genetic variants implicated in complex diseases by genome-wide association studies (GWAS) are non-coding, making it challenging to understand the causative genes involved in disease. Integrating external information such as quantitative trait locus (QTL) mapping of molecular traits (e.g., expression, methylation) is a powerful approach to identify the subset of GWAS signals explained by regulatory effects. In particular, expression QTLs (eQTLs) help pinpoint the responsible gene among the GWAS regions that harbor many genes, while methylation QTLs (mQTLs) help identify the epigenetic mechanisms that impact gene expression which in turn affect disease risk. In this work we propose **m**ultiple-trait-c**oloc** (*moloc*), a Bayesian statistical framework that integrates GWAS summary data with multiple molecular QTL data to identify regulatory effects at GWAS risk loci. We applied *moloc* to schizophrenia (SCZ) and eQTL/mQTL data derived from human brain tissue and identified 52 candidate genes that influence SCZ through methylation. Our method can be applied to any GWAS and relevant functional data to help prioritize disease associated genes.

AUTHOR SUMMARY

In this paper, we propose **m**ultiple-trait-c**oloc** (*moloc*), a statistical method that can use single variant summary statistics from multiple studies to test for colocalization. Using *moloc*, we integrate risk variants identified through genome-wide association studies (GWAS) in schizophrenia with quantitative trait loci that affect gene expression (eQTL) and DNA methylation (mQTL). Our method links non-coding risk variants with transcripts to identify possible biological mechanisms that affect the *cis* regulation of

67   gene expression. Overall, we find that integrating mQTL, eQTL and GWAS increases

68   discovery power as compared to standard pairwise colocalization. Our approach is

69   publically available as an R package from https://github.com/clagiamba/moloc.

INTRODUCTION

Genome-wide association studies (GWAS) have successfully identified thousands of genetic variants associated with complex diseases[1]. However, since the discovered associations point to non-coding regions, it is difficult to identify the causal genes and the mechanism by which risk variants mediate disease susceptibility. Advancement of high-throughput array and sequencing technology has enabled the identification of quantitative trait loci (QTLs), genetic variants that affect molecular phenotypes such as gene expression (expression QTL or eQTL) and DNA methylation (methylation QTL or mQTL). Integration of molecular QTL data has the potential to functionally characterize the GWAS results. Additionally, analyzing two datasets jointly has been a successful strategy to identify shared genetic variants that affect different molecular processes, in particular eQTL and GWAS[2–5] and mQTL and GWAS[6–9] integration. All these previous efforts have focused on pairwise trait integration (e.g. eQTL and GWAS or mQTL and GWAS). Analyzing multiple datasets jointly has the potential to increase power for functional characterization of non-coding, risk variants. In addition, it has the potential to link changes in methylation with specific transcripts, avoiding the issues of arbitrary annotating GWAS-mQTL associations to the nearby genes.

To our knowledge, a statistical approach to integrate multiple QTL datasets with GWAS is lacking. Therefore, we developed **m**ultiple-trait-**coloc** (*moloc*), a statistical method to quantify the evidence in support of a common causal variant at a particular risk region across multiple traits. We applied *moloc* to schizophrenia (SCZ), a complex polygenic psychiatric disorder, using summary statistics from the most recent and

largest GWAS by the Psychiatric Genomics Consortium[10], which reported association

for 108 independent genomic loci. eQTL data were derived from the CommonMind

Consortium[2], which generated the largest eQTL dataset in the dorsolateral prefrontal

cortex (DLPFC) from SCZ cases and control subjects (N=467). Finally, we leveraged

mQTL data that were previously generated in human DLPFC tissue (N=121) to

investigate epigenetic variation in SCZ[6]. Integration of multiple phenotypes helps better

characterize the genes predisposing to complex diseases such as SCZ.

RESULTS

Overview of the Method

We introduce an approach (***moloc***) to detect colocalization among any number of traits in a region in the genome. The input of the model is the set of summary statistics derived from three (or more) traits measured in distinct datasets of unrelated individuals. As a simplifying example, consider colocalization of three traits (e.g., GWAS, eQTL and mQTL); under a maximum of a single causal variant per trait, there can be up to three causal variants and 15 possible scenarios summarizing how the variants are shared among the traits (see Methods). For simplicity, we denote GWAS as G, eQTL as E, and mQTL as M. To illustrate our notation, consider a region with 8 SNP. Four examples of configurations are shown in **Figure 1**. The "." in the subscript denotes scenarios supporting different causal variants. For instance, GE summarizes the scenario for one causal variant shared between traits GWAS and eQTL (**Figure 1** - Right plot top panel); GE.M summarizes the scenario with one causal variant for traits GWAS and eQTL, and a different causal variant for trait mQTL (**Figure 1** - Left plot bottom panel). Our approach computes the posterior probability of any of the 15 configurations by weighting the likelihood of the data D given a configuration S, $P(D|S)$, by the prior probability of a configuration, $P(S)$ (see Methods). Although our approach estimates a posterior for each of the 15 configurations, in general, we are interested in the scenarios supporting a shared causal variant for two and three traits.

Sample Size Requirements

123   In a first set of simulations we explored false positive rates (**Tables S1, S2, S3**)

124 and the posterior probability under different sample sizes. **Figure S1** illustrate the

125 posterior probability distribution across all of the possible scenarios that includes three

126 traits: GWAS, eQTL and mQTL. With a GWAS sample size of 10,000 and eQTL and

127 mQTL sample sizes of 300, the method provides reliable evidence to detect a shared

128 causal variant behind the GWAS and another trait (median posterior probability of any

129 hypothesis >50%). Although in this paper we analyze GWAS, eQTL and mQTL, our

130 method can be applied to any combinations of traits, including 2 GWAS traits and an

131 eQTL dataset. We explored the minimum sample size required when analyzing two

132 GWAS datasets (GWAS1, GWAS2) and one eQTL (**Figure S2**). The method provides

133 reliable evidence for all hypotheses when the two GWAS sample sizes are 10,000 and

134 eQTL sample size reaches 300. False positive rates are below 0.05 even if a threshold

135 of 0.3 for posteriors is used, and where the causal variant is masked (**Tables S1**, **S2**,

136 **S3**). We then explored misclassification rates (**Tables S4, S5, S6**). When samples are

137 10,000 for GWAS and greater than 300 for eQTL and mQTL, misclassification rates for

138 detecting our hypotheses of interests at 80% threshold are all below 0.05 (**Table S4**).

139 Where the causal variant is masked, sample sizes also need to reach 10,000 for GWAS

140 and greater than 300 for eQTL and mQTL, for misclassification rates to be below 0.05

141 (**Table S5**). Given the small sample size for the mQTL data, the method has trouble

142 detecting a different causal variant for the mQTL dataset (**Table S6**). For example,

143 evidence pointing to two different causal variants between GWAS and eQTL could be

144 generated by the presence of three causal variants in reality, but the causal variant for

mQTL remains undetected. For this reason, we focused on cases with shared casual

variants between GWAS, eQTL, with or without mQTL.

It is instructive to observe where evidence for other hypotheses is distributed. **Figure 2**

illustrates the accuracy of our approach under different scenarios where two or three

causal variants are shared. For example, under simulations of one shared variant for

GWAS and eQTL and a second variant for mQTL (GE.M), on average 60% of the

evidence points to the simulated scenario, while 12% point to GE, 12% to G.E.M and

7.2% to GEM.

Choice of priors

The method requires the definition of prior probabilities at the SNP level for the

association with one (p1), two (p2), or three traits (p3). We set the prior probability that a

variant is associated with one trait as 1×10^{-4} for GWAS, eQTL and mQTL, assuming

that each genetic variant is equally likely a priori to affect gene expression or

methylation or disease. This estimate has been suggested in the literature for GWAS[12]

and used in similar methods[13]. We set the priors $p2 = 1 \times 10^{-6}$, $p3 = 1 \times 10^{-7}$ based on

sensitivity and exploratory analysis of genome-wide enrichment of GWAS risk variants

in eQTLs and mQTLs. In **Figure S3**, we find eQTLs and mQTLs to be similarly enriched

in GWAS, justifying our choice of the same prior probability of association across the

two traits. These values are also suggested by a crude approximation of *p2* and *p3* from

the common genome-wide significant SNPs across the three datasets.

We performed sensitivity analyses using different priors. Specifically, we fixed $p1$ to

1×10^{-4} and tested a range of priors for $p2$ and $p3$ from 1×10^{-5} to 1×10^{-8}, with increasing

difference between $p1$, $p2$ and $p3$. We used a form of internal empirical calibration to

compare our prior and posterior expectations. We considered a range of values for $p2$

and $p3$ while keeping $p1$ fixed, and find that the posterior expectation of colocalization

most closely resembled the prior expectation under our choice of priors (**Table S7**). We

note that our R package implementation allows users to specify a different set of priors.

Co-localization of eQTL, mQTL and risk for Schizophrenia

We applied our method to SCZ GWAS using eQTLs derived from 467 samples

and mQTL from 121 individuals. Our aim is to identify the genes important for disease

through colocalization of GWAS variants with changes in gene expression and DNA

methylation. We analyzed associations genome-wide, and report results both across

previously identified GWAS loci, and across potentially novel loci. While we consider all

15 possible scenarios of colocalization, here we focus on gene discovery due to higher

power in our eQTL dataset, by considering the combined probabilities of cases where

the same variant is shared across all three traits GWAS, eQTLs and mQTLs (GEM >

0.8) or scenarios where SCZ risk loci are shared with eQTL only (GE > 0.8 or GE.M >

0.8) (**Table 1**).   We identified 1,053 cis-regions/methylation pairs with posterior

probability above 0.8 that are associated with all three traits (GEM), or eQTLs alone

(GE or GE.M). These biologically relevant scenarios affect overall 84 unique genes and

include 39 genes that fall within the previously identified SCZ LD blocks (**Table S8**) and

45 potentially novel genes outside of these regions (**Table S9**). Fifty-two out of the 84

candidate genes influence SCZ, gene expression and methylation (GEM>=0.8). One

possible scenario is that the variants in these genes could be influencing the risk of SCZ

through methylation, although other potential interpretations such as pleiotropy should

be considered.

Addition of a third trait increases gene discovery

We examined whether moloc with 3 traits enhance power for GWAS and eQTL

colocalization compared to using 2 traits. In simulations to compare coloc and moloc

under one causal variant and our true sample sizes for all three datasets, we observe a

fold increase of 1.5 for gene discovery using moloc versus coloc. Moloc with three traits

recovers all the genes discovered using *coloc* with eQTL and mQTL, and additional

genes from the inclusion of the third layer. In our real data, colocalization analysis of

only GWAS and eQTL traits identified 45 genes with a posterior probability, PP4 in

coloc, of >= 0.8. The 39 additional genes that were found by adding methylation include

genes such as *CALN1,* a neuronal transcript associated with abnormalities in

sensorimotor gating in humans[14], that would have been missed by only GWAS and

eQTL colocalization.

Loci overlapping reported SCZ LD blocks

Psychiatric Genomics Consortium (PGC) identified 108 independent loci and

annotated LD blocks around these, 104 of which are within non-HLA, autosomal regions

of the genome[10]. In **Table 1** we report the number of identified gene-methylation pairs

and unique genes under each scenario that overlap the SCZ-associated LD blocks. Out

of the 78 SCZ-associated LD blocks we examined in our analysis, we found

colocalizations in 20 of them with an average gene density per block of 2.4 unique

genes. **Figure 3A** illustrates the average distribution of the posteriors across these

regions. Cumulatively, 12% of the evidence points to shared variation with an eQTL

(GE, GE.M and GEM). The majority of the evidence within these regions (64%) did not

reach support for shared variation across the three traits, with 20% not reaching

evidence for association with any traits, and 44% with only one of the three traits (36%

with GWAS; 6% with eQTL, 2% with mQTLs). The lack of evidence in these regions

could be addressed with greater sample sizes. **Figure 3B** shows the evidence for

colocalization of GWAS with eQTL or mQTL across the 39 candidate genes. We

provide illustrative examples of SCZ association with expression and DNA methylation

in the *FURIN* locus (**Figure 4** and **Figure S4A**).

Potentially novel SCZ loci

We found 45 unique genes that have a high posterior for SCZ and eQTL, but fall

in regions not previously identified to be associated with SCZ (at p-value of 5×10^{-8}).

All genes were far from a SCZ LD block (more than 150kb, **Table S9**), and contained

SNPs with p-values for association with SCZ ranging from 10^{-4} to 10^{-8}. These genes will

likely be identified using just the GWAS signal if the sample size is increased. *KCNN3* is

among these genes which encodes an integral membrane protein that forms a voltage-

independent calcium-activated channel. It regulates neuronal excitability by contributing

to the slow component of synaptic afterhyperpolarization[15]. A plot of the associations

with the three datasets within this locus is shown in **Figure S4B.**

Comparison with previous findings

We compare our gene discovery results to previous studies that assess GWAS-

eQTL[2–5] or GWAS-mQTL[5,7–9] colocalization using the same or similar datasets (**Table**

2). A substantial proportion of genes detected in our study (range 44%-85%) was

validated with four studies [2–5] that used eQTL and GWAS integration to prioritize genes

important for schizophrenia. Several studies have also linked methylation data with

schizophrenia [7–9]. Two recent studies[7,9] used blood mQTL data from 639 samples and

identified colocalization of SCZ loci with 32 and 200 methylation probes by applying

COLOC and SMR, respectively. A proportion of SCZ-mQTL colocalization was validated

in our study (COLOC: 46%; SMR: 18%, **Table 2**). Overlap between our analysis in brain

and these analyses point to shared mechanisms in blood and brain. Another study[8]

used mQTL data from 166 fetal brain samples and identified 297 methylation probes

important for schizophrenia. We analyzed 184 of those and found evidence for 13

probes. We note that our methylation data did not included fetal brain samples. Finally,

a recent study[5] identified 44 genes involved in schizophrenia through TWAS, followed

by integration with chromatin data in blood that resulted in 11 genes associated with

GWAS, eQTL and epigenome QTL. We analyzed 8 out of the 11 associations and

confirmed 6 of these genes that, in our study, influence SCZ through eQTL and mQTL.

Association of gene expression with methylation

We examined the association of DNA methylation and gene expression as a

function of distance from transcription start site. We explored direction of effects of

methylation and expression, for gene expression and DNA methylation that colocalize

(PPA.GEM + PPA.EM + PPA.G.EM >= 0.8). This approach has the advantage of linking

changes in methylation with specific transcripts, avoiding the issues of arbitrary

annotating CpG methylation sites to the nearby genes. Overall, we tested 1,947 DNA

methylation and gene expression pairwise interactions and found a significant negative

correlation between the effect sizes of methylation and expression in the proximity of

the transcription start site (**Figure 5**, p-value:<2.2e-16). **Table S10** provides a list of

methylation and gene expression pairwise interactions in human brain tissue for DNA

methylation probes that are proximal to the transcription start site (20kb upstream to 2kb

downstream of transcription start site).

DISCUSSION

In this paper, we propose a statistical method for integrating genetic data from molecular quantitative trait loci (QTL) mapping into genome-wide genetic association analysis of complex traits. The proposed approach requires only summary-level statistics and provides evidence of colocalization of their association signals. To our knowledge, a method integrating more than two traits is lacking. In contrast to other methods that attempt to estimate the true genetic correlation between traits such as LD score regression[16] and TWAS[17], *moloc* focuses on genes that are detectable from the datasets at hand. Thus, if the studies are underpowered, most of the evidence will lie in the null scenarios. We note that our model is the same as *gwas-pw* in Pickrell et al.[18] under specific settings. Precisely, *gwas-pw* averages over Bayes factors computed with W = 0.01, W = 0.1, and W = 0.5 (Methods). We provide this as an option that can be specified by the user. Additionally, *gwas-pw* estimates the prior parameters genome-wide using a maximization procedure. However, we note that, unlike *gwas-pw* that focuses on genome-wide estimation across pairs of traits, our approach focuses on one locus at a time with multiple traits.

Our method is complementary to methods quantifying local genetic covariance across traits. The aim of this method is to identify cases where the same causal variant is shared between the traits, and we do not capture cases where traits are driven by a different causal SNP. We argue that it is valuable to identify cases where the same signal is influencing multiple traits, for example when studying application to drug development and possible side effects of drugs to non-target biomarkers. Other

methods such as genetic covariance[19] can be used to identify genes shared across

traits even where the causal variants differ.

We expose one possible application of this approach in SCZ. In this application,

we focus on scenarios involving eQTLs and GWAS, alone or in combination with

mQTLs. Other scenarios are also biologically important. For example, colocalization of

GWAS and mQTL excluding eQTLs (GM.E scenario) could unveil important methylation

mechanisms affecting disease but not directly influencing gene expression in *cis*. We

report these and other scenarios in our web resource and encourage further

examination of these cases in future analyses. The GEM scenario provides evidence

that SCZ risk association is mediated through changes in DNA methylation and gene

expression. While our method does not detect causal relationships among the

associated traits, i.e. whether risk allele leads to changes in gene expression through

methylation changes or vice versa, there is evidence supporting the notion that risk

alleles might affect transcription factor binding and epigenome regulation that drives

downstream alterations in gene expression[20,21].

Our method helps to link a specific gene to a methylation probe. DNA

methylation is one the best studied epigenetic modifications. Methylation can alter gene

expression by disrupting transcription factor binding sites (with variable consequences

to expression depending on the transcription factor), or by attracting methyl-binding

proteins that initiate chromatin compaction and gene silencing. Therefore methylation

can be associated with both increased or decreased gene expression[20,22]. Increased

CpG methylation in promoter regions is usually associated with silencing of gene

expression[23]. However, in genome-wide expression and methylation studies, the

correlation of methylation and gene expression is low or the pattern of association is

mixed, even for CpG methylation within promoter regions[22]. We found a negative

correlation between the effect sizes of methylation and expression in the proximity of

the transcription start site. However, we note that the current assay (Illumina 450K)

cannot discriminate DNA methylation from hydroxymethylation sites. DNA methylation is

generally associated with the inhibition of gene expression, while DNA

hydroxymethylation, the oxidized form of methylation, has been associated with

enhancers and increased gene expression.

We provide posterior probabilities supporting respective hypotheses for each

gene-methylation pair analyzed, and the SNP for each trait with the highest probability

of colocalization with any other trait. For example, the SNP with the highest posterior

probability of GWAS colocalization with eQTL or mQTL will be computed from PPA of

GE + GE.M + GM + GM.E + GEM. However, the aim of this method is not fine-mapping

of SNPs and we encourage researchers to further analyze the identified local

associations with methods better suited for fine-mapping.

We assign a prior probability that a SNP is associated with one trait (1×10^{-4}), to

two (1×10^{-6}), and to three traits (1×10^{-7}). We find support for our choice of priors in the

data using two methods. The first uses stratified QQ plots (**Figure S3**). We find that

eQTL enrichment in GWAS has a similar enrichment to mQTL in GWAS. The second is

a form of empirical calibration as in Guo et al.[24]. We find that the prior and posterior

expectations of colocalisation matched more closely under our choice of priors (**Figure**

S6). However the choices for prior beliefs for each hypothesis are always arguable. One

could estimate priors for the different combinations of datasets. Pickrell et al.[18] proposed

estimation of enrichment parameters from genome-wide results maximizing *a posteriori*

estimates for two traits. For multiple traits, another possibility is using deterministic

approximation of posteriors[25]. We leave these explorations to future research.

The results of this analysis apply to a particular brain region (dorsolateral

prefrontal cortex), age period (adulthood), and ancestry (Caucasians only). We justify

using functional data from DLPFC to examine the colocalization with schizophrenia

variants as it is the most proximal to disease etiopathogenesis tissue [4,5]. In addition, due

to the tissue specificity of gene expression regulatory mechanisms, we matched eQTL

and mQTL data on the same brain region and age period. The genotypes and gene

expression data reported by the CommonMind Consortium (CMC) comprises the largest

existing postmortem brain genomic resource from individuals with psychiatric disorders

as well as unaffected controls. The mQTL dataset used here is the largest dataset in the

adult brain measured in the brain region and age period considered. Although this was

out of the scope of this paper, we encourage application of similar methods to functional

data derived from different tissues and age periods to elucidate the spatiotemporal

regulatory mechanisms of gene expression that are involved in the etiopathogenesis of

schizophrenia.

The method makes use of LD to weigh out the posteriors for the different models,

therefore the statistics (priors and posteriors of configurations) will depend on the

pattern of association (LD) and the number of SNPs in the region (Q). Specifically,

increased LD in a region with one causal variant decreases our chance to detect a

shared causal effect (**Figure S5A**). This is because in loci where the extent of LD is

strong (a large number of variants with the same strength of association), there is

uncertainty on whether we are dealing with a single variant causal for both traits, or two

independent variants in the same LD block. Similarly, the chance of the same SNP

being causal for all three traits decreases as Q increases (**Figure S5B**).

The Bayes Factors under the models of colocalization (for the case of a single

variant affecting more than one trait) are computed by multiplying the individual WABFs

together, and this assumes independence of the effect sizes for the traits.

Independence is appropriate in the case of analyzing different phenotypes, or eQTLs in

different tissues. On the other hand, this assumption may be inappropriate in other

cases, for example if we were looking at colocalization of a variant for the same trait in

two different studies, or the same eQTLs across different tissue type, or the same trait

in two different studies. A more general BF computation which includes cases where the

effects are correlated has been proposed in [26] and could be used to extend our

approach to these situations. Additionally, correlation among the SNP effect sizes may

result if the studies contain overlapping individuals. In this case we may overestimate

the evidence in favor of colocalization. We could adjust SNP Bayes factors to account

for this overlap as in [27]. The datasets we used in the current analysis do not contain

overlapping individuals, however we provide the code to use this approach.

We note that this approach can be extended to more than three traits. Since the

calculations are analytical and no recursive method is used, computation time for a

region with 1000 SNPs is less than one second. However, time increases exponentially

as number of traits increases. For four traits it is about 3 seconds, for five traits it is

greater than 22 minutes. Overall, owing to the increasing availability of summary

statistics from multiple datasets, the systematic application of this approach can provide

clues into the molecular mechanisms underlying GWAS signals and how regulatory

variants influence complex diseases.

MATERIALS AND METHODS

Ethics statement

This paper re-analyzes previously published datasets. All samples and patient data were handled in accordance with the policies and procedures of the participating organizations.

GWAS dataset

Summary statistics for genome-wide SNP association with Schizophrenia were obtained from the Psychiatric Genomics Consortium-Schizophrenia Workgroup (PGC-SCZ) primary meta-analysis (35,476 cases and 46,839 controls)[10].

Expression QTL (eQTL) analysis

For the eQTL associations, we used pre-QC'ed expression and imputed genotypes, on case-control of European-ancestry (N=467) collected by the CommonMind Consortium. Briefly, gene expression was assayed data from post-mortem DLPFC (Brodmann areas 9 and 46) with RNA-seq and mapped to human Ensembl gene reference (v70) using TopHat version 2.0.9 and Bowtie version 2.1.0. Genes with less than 1 CPM (counts per million) in at least 50% of the samples were discarded and RNA-seq data for a total of 16,423 Ensembl genes were considered for downstream analyses. After RNA sample QC, and retaining only genetically-identified European-ancestry individuals, a total of 467 samples were used for downstream analyses. These comprised 209 SCZ cases, 52 AFF (Bipolar, Major depressive disorder, or Mood disorder, unspecified) cases, and 206 controls. The expression data

was voom-adjusted[28] for both known covariates (RIN, library batch effect, institution,

diagnosis, post-mortem interval, age and sex) and hidden confounds identified based

on surrogate variable analysis.

Samples were genotyped at 958,178 markers using the Illumina Infinium

HumanOmniExpressExome array. Genotyped variants were QC'd, phased, and

imputed based on the Phase 1 reference panel from the 1,000 Genomes Project.

Variants with INFO < 0.8 or MAF < 0.05 were filtered out, and approximately 6.4 million

variants were included in further analysis. Further details on processing of raw data,

normalization, and quality control procedures, can be found in [2].

The R package MatrixEQTL [29] was used to fit an additive linear model between

the expression of 16,423 genes and SNP dosage data at 6.4 million markers within a 1

Mb window around the transcription start site for each gene, including covariates for

diagnosis, five ancestry covariates and known and hidden variables detected by

surrogate variable analysis.

Following eQTL identification, summary statistics on 72,441,005 SNPs and 15,791

autosomal genes were retained for downstream analyses. Overall, the model identified

2,154,331 significant cis-eQTL, (i.e., SNP–gene pairs within 1 Mb of a gene) at a false

discovery rate (FDR) ≤ 5%, for 13,137 (80%) genes.

Methylation QTL (mQTL) analysis

For the mQTL dataset, we used pre-QC'd DNA methylation data and imputed

genotypes on non-psychiatric adult Caucasian control donors (age > 13, N=121). We

briefly summarize here the procedures to process the raw data. More details on data

processing, normalization, and QC can be found in [6].

DNA methylation was assessed on postmortem tissue homogenates of the dorsolateral

prefrontal cortex (DLPFC, Brodmann areas 9 and 46) on 534 unique subjects using the

Illumina HumanMethylation450 ("450k") microarray (which measures CpG methylation

across 473,058 probes covering 99% of RefSeq gene promoters). The average intensity

was used to check for low quality samples, intensities from the sex chromosomes were

used to drop samples that had predicted sex which was different from its recorded

value, and across-sample quantile normalization was used. DNA for genotyping was

obtained from the cerebella of samples with either the Illumina Human Hap 650v3, 1M

Duo V3, or Omni 5M BeadArrays. Genotypes were called, QC'd, phased and imputed to

the 1000 Genomes Phase 3 variant set, separately by genotyping platform. Imputed

genotypes were then merged across the three platforms, and SNPs with MAF > 5% and

imputation quality above 90% were retained across the samples. Multidimensional

scaling (MDS) components were obtained for quantitative measures of ancestry.

For the mQTL association analysis, we only considered 121 adult Caucasian

individuals (age > 13). The analysis was conducted using the R package MatrixEQTL [29],

fitting an additive linear model up to 20kb distance between each SNP and CpG

analyzed, including as covariates the first five MDS components from the genetic data

and the first 8 PCs (based on the 100,000 most inter-individual variable probes for

computational efficiency). Summary statistics on 47,675,913 SNPs and 473,058

Illumina 450k probes were used for downstream analyses. Overall, the model identified

3,601,681 significant cis-mQTL, (i.e., SNP–probe pairs within 20kb distance of a probe)

at a false discovery rate (FDR) ≤ 5%, for 127,237 (27%) probes.

Method Description

Our approach is a multi-trait extension of the two-trait model described in

coloc[11]. We define a genomic region containing **Q** variants, for example a *cis* region

around expression or methylation probe. We are interested in a situation where

summary statistics (effect size estimates and standard errors) are available for all

datasets in the genomic region. We first derive our model using three traits, then

generalize to any number of traits. If we consider three traits, there are 15 possible

scenarios summarizing how the variants are shared among the traits. Each hypothesis

can be represented by a set of index sets according to which of the traits each SNP is

associated with: H_0- no association to any trait, H_1- association only for trait {1}, H_2-

association only for trait {2}, H_3- association only for trait {3}, H_4- association for traits {1

and 2}, H_5- association for traits {2 and 3}, H_6- association for traits {1 and 3}, H_7-

association for trait 1 and 2, but different causal variants for {1} and {2}, H_8- association

for traits 2 and 3, but different causal variants for {2} and {3}, H_9- association for traits 1

and 3, but different causal variants for {1} and {3}, H_{10}- association for traits 1, 2 and 3,

but different causal variants for {1} and {2,3}, H_{11}- association for traits 1, 2 and 3, but

different causal variants for {3} and {1,2}, H_{12}- association for traits 1, 2 and 3, but

different causal variants for {2} and {1,3}, H_{13}- association for traits 1, 2 and 3, but

different causal variants for {1}, for {2} and for {3}, and H_{14} SNP is associated with all 3

traits {1,2,3}.

Our aim is to find the evidence supporting 15 possible scenarios $(H_0 \cdots H_{14})$, of sharing

of SNPs among traits in the given genomic region. We compute the posterior probability

of a configuration by weighting the likelihood of the data *D* given a configuration *S*,

P(D|S), by the prior probability of a configuration, *P(S)*. We can reformulate the

posterior probability for each hypothesis as a ratio by dividing each by the baseline

likelihood supporting the first model of no association with any trait H_0. The probability of

the data for hypothesis h is then the sum over all configurations Sh, which are

consistent with the given hypothesis:

$$\frac{P(H_h|D)}{P(H_0|D)} = \sum_{S \in S_h} \frac{P(D \mid S)}{P(D \mid S_0)} \times \frac{P(S)}{P(S_0)} \qquad (1)$$

where, *P(D|S)/P(D|S₀)* is the Bayes Factor for each configuration compared to the

baseline configuration of no association with any trait *S₀*, *P(S)/P(S₀)* is the prior odds of

a configuration compared with the baseline configuration *S₀*, and the sum is over Sh,

the set of configurations supporting hypothesis H_0 to H_{14}. Similar to pairwise

colocalization[11] we then estimate the evidence in support of different scenarios in a

given genomic region using the posterior probability supporting hypothesis *h* among *H*

possible hypothesis, computed from:

$$PP_h = P(H_h|D) = \frac{P(H_h|D)}{\sum_{i=0}^{H} P(H_i)} = \frac{\frac{P(H_h|D)}{P(H_0|D)}}{1 + \sum_{i=1}^{H} \frac{P(H_i|D)}{P(H_0|D)}} \qquad (2)$$

Therefore, in our application, the method outputs 15 posterior probabilities. We are most

interested in the scenarios supporting a shared causal variant for two and three traits.

We make three important assumptions in *moloc*, the same that are made in our

previous *coloc* methodology. Firstly, that the causal variant is included in the set of *Q*

common variants, either directly typed or well imputed. If the causal SNP is not present,

the power to detect a common variant will be reduced depending on the linkage

disequilibrium (LD) between other SNPs included in the model and the causal SNP.

Secondly, we assume at most one causal variant is present for each trait. In the

presence of multiple causal variants per trait, this method is not able to identify

colocalization between additional association signals independent from the primary one.

Thirdly, as we do not explicitly model LD between SNPs, we assume the samples are

drawn from the same ethnic population and therefore have identical allele frequencies

and patterns of LD.

Bayes Factor of a SNP with one trait

We start by computing a Bayes Factor for each SNP and each trait (i.e GWAS, eQTL, mQTL). We assume a simple linear regression model to relate the phenotypes or a log-odds generalized linear model for the case-control dataset, and the genotypes. Using the Wakefield Approximate Bayes factors[30] (WABF), only the variance and effect estimates from regression analysis are needed, as shown below and previously described[11,18]:

$$WABF_i^j = \frac{1}{\sqrt{1-r}} \times exp\left[-\frac{Z_i^{2(j)}}{2} \times r\right] \quad (3)$$

where $Z_{ij} = \hat{\beta}/\sqrt{V}$ is the usual Z statistic and the shrinkage factor r is the ratio of the

variance of the prior and total variance ($r = W/(V + W)$).

The WABF requires specifying the variance W of the normal prior. In the moloc method

we set W to 0.15 for a continuous trait and 0.2 for the variance of the log-odds ratio

parameter, as previously described[11]. Another possibility is to average over Bayes

factors computed with W = 0.01, W = 0.1, and W = 0.5 [27]. We provide this as an option

that can be specified by the user.

If the variance of the estimated effect size V is not provided, it can be approximated

using the allele frequency of the variant f, the sample size N (and the case control ratio

s for binary outcome). For example, in the case-control setting, V is estimated using the

following equation[11]:

$$V_i = Var(\hat{\beta}_i) = \frac{1}{Ns_i(1 - s_i) \times 2f_i(1 - f_i)} \qquad (4)$$

Bayes factor of a SNP across more than one trait

To compute the BF where a SNP i associates with more than one trait, we use:

$$BF_{i,\mathbf{s}} = \prod_{j \in s} BF_i^j \qquad (5)$$

Where **s** is the set of trait indices for which SNP i is associated with.

Note that the computations under >1 trait multiply the individual Bayes Factors together.

This is equivalent to the Bayes Factor under the maximum heterogeneity model used in

Wen and Stephens[26]. Two key assumptions are necessary for the following

computations. Firstly, that the traits are measured in unrelated individuals, and secondly

that the effect sizes for the two traits are independent. We will consider when these

assumptions are appropriate in Discussion.

Prior probabilities that SNP i associates with traits indexed in s

The prior probability that SNP i associates with all traits indexed in a set in our

three trait model is: π_φ SNP i associates with no trait, with one trait, pairs or traits or all

traits $\pi_{\{1,2,3\}}$ such that they sum to 1: $\pi_\varphi + \pi_{\{1\}} + \pi_{\{2\}} + \pi_{\{3\}} + \pi_{\{1,2\}} + \pi_{\{1,3\}} + \pi_{\{2,3\}} +$

$\pi_{\{1,2,3\}} = 1$.

Simplified model under the assumption that all SNPs have the same prior

The probability of the data under each hypothesis can be computed by summing

the probability of all the causal configurations consistent with a particular hypothesis,

weighted by the prior probabilities (Equation 1).

In our model, *P(S)* is the prior probability under any one of the 15 hypotheses. We can

define these priors from the prior probability that a SNP i associates with traits indexed

in π_s (section above). Additionally, since the prior probability *P(S)* of any one

configuration in the different sets do not vary across SNPs that belongs to the same set

S_h, we can multiply the likelihoods by one common prior supporting the different

hypothesis[11]. In this framework, we can see that *P(S)* depends on a ratio of π_s and on

Q, the number of SNPs in the region (Text S1).

Therefore, across a set of j traits {1,2,3...,}, we compute the probability of the data

supporting hypothesis h, where one SNP is associated with j traits as:

$$\frac{P(H_h|D)}{P(H_0|D)} = \prod_{s\in h} \pi_s \sum_{i=1}^{Q} BF_{i,s} \quad 533 \qquad (6)$$

534   where $BF_{i,s}$ are the Bayes factor of a SNP across traits indexed in s (Equation 5), π are

535   the prior probabilities that SNP i is the causal SNP under a specific model.

The probability of the data where there are more than one independent associations

among the j traits (i.e. |h| > 1) can be derived from the pre-computed probability of the

data where there is one association among the j traits (i.e. |h| = 1, Text S1):

$$\frac{P(H_h|D)}{P(H_0|D)} = \prod_{s\in h} \pi_s \sum_{i=1}^{Q} BF_{i,s} - \frac{\prod_{s\in h} \pi_s}{\pi_t} \sum_{i=1}^{Q} \pi_t BF_{i,t} \qquad (7)$$

537   where t is the union of the index set in h.

Prior Probabilities of each hypothesis

540   In practice, we collapsed the prior probabilities to a smaller set for each kind of

541   configuration. We set the prior probability that a SNP is causal in each trait to be

542   identical (π{1} = π{2} = π{3}) and refer to this a p1. We also set the prior probability that

543   is associated with two traits to be identical (π{1,2},π{2,3},π{1,3}) and refer to this as p2.

544   We refer to the prior probability that SNP i the causal for all traits (π{1,2,3}) as p3.

Moloc Analysis

The GWAS, eQTL, mQTL datasets were filtered by minor allele frequency

greater than 5% and had individually been filtered by imputation quality (Methods).. The

Major Histocompatibility (MHC) region (chr 6: 25 Mb - 35 Mb) was excluded from all co-

localization analyses due to the extensive linkage disequilibrium. We applied a genic-

centric approach, defined *cis*-regions based on a 50kb upstream/downstream from the

start/end of each gene, since our goal is to link risk variants with changes in gene

expression. We evaluated all methylation probes overlapping the *cis*-region. The

number of cis-regions/methylation pairs is higher than the count of genes because, on

average, there are more than one methylation sites per gene. Common SNPs were

evaluated in the colocalization analysis for each gene, and each methylation probe, and

GWAS. In total, 12,003 *cis*-regions and 481,995 unique *cis*-regions/methylation probes

were tested. Genomic regions were analyzed only if greater than 50 SNPs were in

common between all the datasets. Across all of the analyses, a posterior probability

equal to, or greater than, 80% for each configuration was considered evidence of

colocalization.

In order to compare existing methods for colocalization of two trait analyses with

three traits, we applied *moloc* using the same region definitions across the same set of

SNPs, using a previously developed method (coloc[11]). Effect sizes and variances were

used as opposed to p-values, as this strategy achieves greater accuracy when working

with imputed data[11].

Simulations

We simulated genotypes from sampling with replacement among haplotypes of

SNPs with a minor allele frequency of at least 5% found in the phased 1000 Genomes

Project within 49 genomic regions that have been associated with type 1 diabetes (T1D)

susceptibility loci (excluding the major histocompatibility complex (MHC) as previously

described[31]. These represent a range of region sizes and genomic topography that

reflect typical GWAS hits in a complex trait. For each trait, two, or three "causal

variants" were selected at random. We have simulated continuous traits, and assume

that causal effects follow a multivariate Gaussian distribution, with each causal variant

explaining 0.01 variance of the trait in the GWAS data, and 0.1 in the eQTL and mQTL

datasets. Note that colocalisation testing may be applied equally to quantitative data

(using linear regression), and to case control data (using logistic regression). In our

application, one of the datasets is a case-control and we are directly using estimates,

and associated standard errors, from fitting a log-odds generalised linear model. This is

equivalent to applying *moloc* on the latent liability scores, which is a quantitative

phenotype. For the null scenario, the causal variants explain zero variance of the traits.

Each scenario was simulated 1,000 times. To quantify false positive rates on a large

number of tests, we simulated the null 500,000 times. We simulated the 15 possible

scenarios with different sharing patterns between the GWAS, eQTL, and mQTL

datasets. We used sample sizes of 82,315, 467, and 121 individuals to reflect our true

sample sizes. We also used different combinations of sample sizes to explore power to

detect the correct hypothesis.

We estimated the number of false positives within each simulated scenario, by

counting the proportion of simulations under the null that passed a posterior probability

supporting each of the 14 hypothesis at a particular threshold (PPA>=threshold). We

also report the false positives using the sum of the posteriors (PPA.ab + PPA.ab.c +

PPA.abc). The false positive rate is the number of false positives over 1,000

simulations. We repeated this procedure using 500,000 simulations under our true

sample sizes.

We next sought to compare the misclassification rates, and power to detect the

correct hypothesis. To compute the number of misclassified calls within each simulated

scenario, we counted the proportion of simulations that passed a posterior probability

supporting a different hypothesis from the one simulated at a particular threshold

(PPA>=threshold). We estimated power to distinguish a particular hypothesis from the

others by counting the proportion of correctly identified simulations at a particular

threshold (PPA(true)>=threshold). Since in most cases the causal variant will not be

included in the panel, we repeated simulations after removing the causal variant.

To explore the effect of linkage disequilibrium (LD) on estimated posterior

probability, we first computed an LD score for each SNP in the region, defined as the

sum of the squared correlation between a SNP and all the SNPs in the region. To

assess the degree of LD at a locus we took the average of these scores.

All analyses were conducted in R.

Web Resources

We developed a web site to visualize the colocalization results of SCZ GWAS,

eQTL, mQTLs under all possible scenarios (**icahn.mssm.edu/moloc**). The browser

allows searches by gene, methylation probe, and scenario of interest. The *moloc*

method is available as an R package from https://github.com/clagiamba/moloc.

ACKNOWLEDGMENTS

We would like to thank Chris Wallace at the Department of Medicine and MRC

Biostatistics Unit, Cambridge Biomedical Campus, University of Cambridge, Cambridge,

UK.

This work was supported by the National Institutes of Health (R01AG050986 Roussos

and R01MH109677 Roussos), Brain Behavior Research Foundation (20540 Roussos),

Alzheimer's Association (NIRG-340998 Roussos) and the Veterans Affairs (Merit grant

BX002395 Roussos). Additionally, this work was supported in part through the

computational resources and staff expertise provided by Scientific Computing at the

Icahn School of Medicine at Mount Sinai.

Data were generated as part of the CommonMind Consortium supported by funding

from Takeda Pharmaceuticals Company Limited, F. Hoffman-La Roche Ltd and NIH

grants   R01MH085542,   R01MH093725,   P50MH066392,   P50MH080405,

R01MH097276, RO1-MH-075916, P50M096891, P50MH084053S1, R37MH057881

and R37MH057881S1, HHSN271201300031C, AG02219, AG05138 and MH06692.

Brain tissue for the study was obtained from the following brain bank collections: the

Mount Sinai NIH Brain and Tissue Repository, the University of Pennsylvania

Alzheimer's Disease Core Center, the University of Pittsburgh NeuroBioBank and Brain

and Tissue Repositories and the NIMH Human Brain Collection Core. CMC Leadership:

Pamela Sklar, Joseph Buxbaum (Icahn School of Medicine at Mount Sinai), Bernie

Devlin, David Lewis (University of Pittsburgh), Raquel Gur, Chang-Gyu Hahn (University

of Pennsylvania), Keisuke Hirai, Hiroyoshi Toyoshiba (Takeda Pharmaceuticals

Company Limited), Enrico Domenici, Laurent Essioux (F. Hoffman-La Roche Ltd), Lara

Mangravite, Mette Peters (Sage Bionetworks), Thomas Lehner, Barbara Lipska (NIMH).

REFERENCES

1. Visscher, P.M., Brown, M.A., McCarthy, M.I., and Yang, J. (2012). Five Years of GWAS Discovery. Am. J. Hum. Genet. *90*, 7–24.

2. Fromer, M., Roussos, P., Sieberts, S.K., Johnson, J.S., Kavanagh, D.H., Perumal, T.M., Ruderfer, D.M., Oh, E.C., Topol, A., Shah, H.R., et al. (2016). Gene expression elucidates functional impact of polygenic risk for schizophrenia. Nat. Neurosci. *19*, 1442–1453.

3. Zhu, Z., Zhang, F., Hu, H., Bakshi, A., Robinson, M.R., Powell, J.E., Montgomery, G.W., Goddard, M.E., Wray, N.R., Visscher, P.M., et al. (2016). Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. Nat. Genet. *48*, 481–487.

4. Hauberg, M.E., Zhang, W., Giambartolomei, C., Franzén, O., Morris, D.L., Vyse, T.J., Ruusalepp, A., Sklar, P., Schadt, E.E., Björkegren, J.L.M., et al. (2017). Large-Scale Identification of Common Trait and Disease Variants Affecting Gene Expression. Am. J. Hum. Genet. *100*, 885–894.

5. Gusev, A., Mancuso, N., Finucane, H.K., Reshef, Y., Song, L., Safi, A., Oh, E., O'donovan, M.C., Katsanis, N., Crawford, G.E., et al. TITLE: Transcriptome-wide association study of schizophrenia and chromatin activity yields mechanistic disease insights.

6. Jaffe, A.E., Gao, Y., Deep-Soboslay, A., Tao, R., Hyde, T.M., Weinberger, D.R., and Kleinman, J.E. (2016). Mapping DNA methylation across development, genotype and schizophrenia in the human frontal cortex. Nat. Neurosci. *19*, 40–47.

7. Hannon, E., Dempster, E., Viana, J., Burrage, J., Smith, A.R., Macdonald, R., St

Clair, D., Mustard, C., Breen, G., Therman, S., et al. (2016). An integrated genetic-

epigenetic analysis of schizophrenia: evidence for co-localization of genetic

associations and differential DNA methylation. Genome Biol. *17*, 176.

8. Hannon, E., Spiers, H., Viana, J., Pidsley, R., Burrage, J., Murphy, T.M., Troakes, C.,

Turecki, G., O'Donovan, M.C., Schalkwyk, L.C., et al. (2016). Methylation QTLs in the

developing brain and their enrichment in schizophrenia risk loci. Nat. Neurosci. *19*, 48–

54.

9. Hannon, E., Weedon, M., Bray, N., O'Donovan, M., and Mill, J. (2017). Pleiotropic

Effects of Trait-Associated Genetic Variation on DNA Methylation: Utility for Refining

GWAS Loci. Am. J. Hum. Genet. *100*, 954–959.

10. of the Psychiatric Genomics Consortium, S.W.G. (2014). Biological insights from

108 schizophrenia-associated genetic loci. Nature *511*, 421–427.

11. Giambartolomei, C., Vukcevic, D., Schadt, E.E., Franke, L., Hingorani, A.D.,

Wallace, C., and Plagnol, V. (2014). Bayesian test for colocalisation between pairs of

genetic association studies using summary statistics. PLoS Genet. *10*, e1004383.

12. Stephens, M., and Balding, D.J. (2009). Bayesian statistical methods for genetic

association studies. Nat. Rev. Genet. *10*, 681–690.

13. Hormozdiari, F., van de Bunt, M., Segrè, A.V., Li, X., Joo, J.W.J., Bilow, M., Sul,

J.H., Sankararaman, S., Pasaniuc, B., and Eskin, E. (2016). Colocalization of GWAS

and eQTL Signals Detects Target Genes. Am. J. Hum. Genet. *99*, 1245–1260.

14. Roussos, P., Giakoumaki, S.G., Zouraraki, C., Fullard, J.F., Karagiorga, V.-E.,

Tsapakis, E.-M., Petraki, Z., Siever, L.J., Lencz, T., Malhotra, A., et al. (2016). The

Relationship of Common Risk Variants and Polygenic Risk for Schizophrenia to

Sensorimotor Gating. Biol. Psychiatry *79*, 988–996.

15. Deignan, J., Luján, R., Bond, C., Riegel, A., Watanabe, M., Williams, J.T., Maylie, J.,

and Adelman, J.P. (2012). SK2 and SK3 expression differentially affect firing frequency

and precision in dopamine neurons. Neuroscience *217*, 67–76.

16. Bulik-Sullivan, B., Finucane, H.K., Anttila, V., Gusev, A., Day, F.R., Loh, P.-R.,

Duncan, L., Perry, J.R.B., Patterson, N., Robinson, E.B., et al. (2015). An atlas of

genetic correlations across human diseases and traits. Nat. Genet. *47*, 1236–1241.

17. Gusev, A., Ko, A., Shi, H., Bhatia, G., Chung, W., Penninx, B.W.J.H., Jansen, R., de

Geus, E.J.C., Boomsma, D.I., Wright, F.A., et al. (2016). Integrative approaches for

large-scale transcriptome-wide association studies. Nat. Genet. *48*, 245–252.

18. Pickrell, J.K., Berisa, T., Liu, J.Z., Ségurel, L., Tung, J.Y., and Hinds, D.A. (2016).

Detection and interpretation of shared genetic influences on 42 human traits. Nat.

Genet. *48*, 709–717.

19. Shi, H., Mancuso, N., Spendlove, S., and Pasaniuc, B. (2016). Local genetic

correlation gives insights into the shared genetic architecture of complex traits. Doi.org

92668.

20. Tak, Y.G., and Farnham, P.J. (2015). Making sense of GWAS: using epigenomics

and genome engineering to understand the functional relevance of SNPs in non-coding

regions of the human genome. Epigenetics Chromatin *8*, 57.

21. Li, Y.I., Van De Geijn, B., Raj, A., Knowles, D.A., Petti, A.A., Golan, D., Gilad, Y.,

and Pritchard, J.K. RNA splicing is a primary link between genetic variation and

disease.

22. Wagner, J.R., Busche, S., Ge, B., Kwan, T., Pastinen, T., and Blanchette, M.

(2014). The relationship between DNA methylation, genetic and expression inter-

individual variation in untransformed human fibroblasts. Genome Biol. *15*, R37.

23. Du, X., Han, L., Guo, A.-Y., and Zhao, Z. (2012). Features of methylation and gene

expression in the promoter-associated CpG islands using human methylome data.

Comp. Funct. Genomics *2012*, 598987.

24. Guo, H., Fortune, M.D., Burren, O.S., Schofield, E., Todd, J. a, and Wallace, C.

(2015). Integration of disease association and eQTL data using a Bayesian

colocalisation approach highlights six candidate causal genes in immune-mediated

diseases. Hum. Mol. Genet. *24*, 3305–3313.

25. Wen, X., Pique-Regi, R., and Luca, F. (2017). Integrating molecular QTL data into

genome-wide genetic association analysis: Probabilistic assessment of enrichment and

colocalization. PLoS Genet. *13*,.

26. Wen, X., and Stephens, M. (2011). Bayesian methods for genetic association

analysis with heterogeneous subgroups: From meta-analyses to gene-environment

interactions.

27. Pickrell, J.K. (2014). Joint analysis of functional genomic data and genome-wide

association studies of 18 human traits. Am. J. Hum. Genet. *94*, 559–573.

28. Law, C.W., Chen, Y., Shi, W., and Smyth, G.K. (2014). voom: precision weights

unlock linear model analysis tools for RNA-seq read counts. Genome Biol. *15*, R29.

29. Shabalin, A.A. (2012). Matrix eQTL: ultra fast eQTL analysis via large matrix

operations. Bioinformatics *28*, 1353–1358.

30. Wakefield, J. (2009). Bayes factors for genome-wide association studies:

comparison with P-values. Genet. Epidemiol. *33*, 79–86.

31. Wallace, C. (2013). Statistical testing of shared genetic control for potentially related

traits. Genet. Epidemiol. *37*, 802–813.

FIGURES

Figure 1. Graphical representation of four possible configurations at a locus with 8

SNPs in common across three traits. The traits are labeled as G, E, M representing

GWAS (G), eQTL (E), and mQTL (M) datasets, respectively. Each plot represents one

possible configuration, which is a possible combination of 3 sets of binary vectors

indicating whether the variant is associated with the selected trait. Left plot top panel

(GEM scenario): points to one causal variant behind all of the associations; Right plot

top panel (GE scenario): represent the scenario with the same causal variant behind the

GE and no association or lack of power for the M association; Left plot bottom panel

(GE.M scenario): represents the case with two causal variants, one shared by the G

and E, and a different causal variant for M; Right plot bottom panel (G.E.M. scenario):

represents the case of three distinct causal variants behind each of the datasets

considered.

Figure 2. Results from simulations under colocalization/non-colocalization scenarios

using a sample size of 10,000 individuals for GWAS trait (denoted as G), 300 for eQTL

trait (denoted as E), and 300 for mQTL trait (denoted as M). X-axis shows all 15

simulated scenarios, e.g. G.E.M, three different causal variants for each of the three

traits; G.EM, 2 different causal variants, one for G and one shared between E and M;

GE, 1 shared causal variant for G and E; GE.M, 2 different causal variants, one shared

between G and E and one for M; GEM, one causal variant shared between all the three

traits. Y-axis shows the distribution of posterior probabilities under the simulated

scenario. The height of the bar represents the mean of the PPA for each configuration

across simulations.

A

B

Figure 3. Summary of genes identified using three-trait colocalization within the SCZ LD

blocks. **A**. Mean posterior probability for each hypotheses computed using the cis-

regions overlapping the SCZ LD blocks. Sections of the pie chart represent the 15

scenarios representing the possible combination of the three traits. The "." between the

traits denotes scenarios supporting different causal variants. The combined scenarios

GE, GE.M, GE account for 12.0%. **B**. Heatmap displaying the maximum posterior

probabilities reached by the 39 regions overlapping known SCZ LD blocks (gene,

number of methylation probes).

Figure 4. Illustration of one example of colocalization results with GWAS-eQTL-mQTL.

FURIN gene and cg24888049; Shown are Z-scores (regression coefficients/standard

errors) from association of expression (x-axis) and association of methylation (y-axis) at

the *FURIN* locus. Each point represents a SNP and the size of the dot represent the

level of association using Z-scores. The red point shows the SNP with the strongest

evidence for eQTL, mQTL, GWAS (rs4702).

Figure 5. Spearman correlation of eQTL and mQTL effect estimates by distance from

transcription start site of the gene. Intervals of methylation probe distance from TSS

were estimated based on 10 equal size bins.

784   **Table 1**. Number of genes with evidence of colocalization (PPA>=0.8) under each

785   scenario.

Scenarios	Sharing of variant	Unique gene-methylation pairs	Unique genes		
		Total PPA>=80%	Total PPA>=80%	Overlapping SCZ LD blocks	Number of LD blocks
Null	No associations	268569	9317	90	55
G	GWAS only	4331	215	146	63
E	eQTL only	97852	3597	14	12
M	mQTL only	21914	5588	42	28
G.E	GWAS not eQTL (2 causals)	1257	66	47	27
E.M	eQTL not mQTL (2 causals)	7542	1952	7	6
G.M	GWAS not mQTL (2 causals)	219	75	52	26
GE	GWAS,eQTL	359	30	18	14
EM	eQTL,mQTL	1494	769	3	3
GM	GWAS,mQTL	38	23	18	10
GM.E	GWAS,mQTL not eQTL (2 causals)	20	11	7	5
G.EM	eQTL,mQTL not GWAS (2 causals)	22	10	7	4
GE.M	GWAS,eQTL not mQTL (2 causals)	31	17	10	7
G.E.M	not GWAS not eQTL not mQTL (3 causals)	63	29	23	15
GEM	GWAS,eQTL,mQTL	123	52	25	11
GEM *or* GE.M *or* GE	**combined scenarios for GWAS,eQTL**	1053	84	39	20
total	total	481995	12003	273	78

786 **Table 2**. Summary of Previous Findings integrating SCZ GWAS, CMC eQTL and

787 methylation datasets.

Reference	Fromer et al. [2]	Zhu et al. [3]	Hauberg et al. [4]	Gusev at al. [5]	Hannon et al. [7]	Hannon et al. [8]	Hannon et al. [9]
Method Used	**Sherlock**	**SMR**	**SMR**	**TWAS**	**COLOC**	**COLOC**	**SMR**
Datasets	GWAS SCZ (N=82315); eQTL in brain (N=537)	GWAS SCZ (N=82315); eQTL in blood (N= 5311)	GWAS SCZ (N=82315); eQTL in brain (N=467)	GWAS SCZ (N=82315); eQTL in brain (N=621); chromatin QTL in blood (N=45)	GWAS SCZ (N=82315); mQTL in blood (N=639)	GWAS SCZ (N=82315); mQTL in fetal brain (N=166)	GWAS SCZ (N=82315); mQTL in blood (N=639)
Scenarios examined in our analysis	**22**	**9**	26	GWAS+eQTL: **35** GWAS+eQTL+mQTL: **8**	**15**	184	49
Validated scenarios (%) at PPA 0.8	**15 (68%)**	**4 (44.4%)**	**22 (85%)**	GWAS+eQTL: **21 (60%)** GWAS+eQTL+mQTL: **6 (75%)**	**7 (46%)**	**13 (7%)**	**9 (18%)**
Genes and methylation sites validated*	**GWAS+eQTL**: *SF3B1, C2orf47, CNTN4, CLCN3, ENSG00000253553, PPP1R13B, EFTUD1P1, ENSG00000225151, FURIN, INO80E, TOM1L2, DRG2, MAU2, GATAD2A, WBP2NL*	**GWAS+eQTL**: *SF3B1, PCCB, C17ORF39, IRF3*	**GWAS+eQTL**: *AL022476.2, ALMS1P, CLCN3, DOC2A, DRG2, EFTUD1P1, ELAC2, EMB, FAM86B3P, FURIN, GATAD2A, GOLGA2P7, INO80E, JRK, PCCB, PCDHA7, RBBP5, RP11-45P15.4, SF3B1, SLC9B1, SLCO4C1, VPS37A*	**GWAS+eQTL:** *ALMS1P, C2orf47, CPNE7, DOC2A, DRG2, ELOVL7, EMB, FURIN, GATAD2A, MAU2, MCHR1, NDUFA2, NT5C2, PCCB, PCDHA2, PRMT7, SEPT10, SF3B1, SLC45A1, TMEM81, ZMAT2* **GWAS+eQTL+mQTL:** *SLC45A1,PCCB,NDUFA2,PCDHA2,ZMAT2,PRMT7*	**GWAS+mQTL:** *cg00585072, cg01262667, cg02951883, cg08607108, cg08772003, cg19624444, cg26732615*	**GWAS+mQTL:** *cg08772003, cg19624444, cg08607108, cg02951883, cg00585072, cg26732615, cg02206767, cg26395211, cg25066665, cg04140906, cg04398451, cg10596483, cg24634471*	**GWAS+mQTL:** *cg25066665, cg02951883, cg07143303, cg08772003, cg14047540, cg04140906, cg12863693, cg04398451, cg26732615*

* Genes validated (GWAS+eQTL+mQTL and GWAS+eQTL) include the genes with

PPA.GEM+PPA.GE+PPA.GE.M>= 80%; methylation sites validated (GWAS+mQTL)

include PPA.GEM+PPA.GM+PPA.GM.E>= 80%.

SUPPORTING INFORMATION CAPTIONS

Figure S1. Simulations under different sample sizes for all scenarios in *moloc* of three

traits (GWAS, eQTL, mQTL). All cases include 10,000 individuals in the GWAS dataset.

The variance explained by the trait was set to 0.01 for GWAS (1%), and to 0.1 (10%) for

the eQTL and mQTL.

Figure S2. Simulations under different sample sizes for all scenarios in *moloc* of three

traits ("a" is GWAS 1, "b" is GWAS 2, "c" is eQTL). All cases include 10,000 individuals

in the biomarker dataset. The variance explained by the trait was set to 0.01 for GWAS

1 and GWAS 2, and to 1% (10%) for the eQTL.

Figure S3. Stratified Q-Q plots for eQTL and mQTLs in the SCZ GWAS. Enrichment of

each functional category (blue line: eQTL; green: mQTL; red: combined eQTL and

mQTL) illustrate the estimated increase in terms of the proportion of p-values expected

in each scenario and in the functionally unannotated variant (FUV) category (grey line).

The estimated enrichment for eQTL and mQTL is similar.

Figure S4. Illustration of two example of colocalization results with GWAS-eQTL-mQTL

in the *FURIN* locus and cg24888049 and *KCNN3* and cg09359103. −log10(p)

association p-values for GWAS (top panels), −log10(p) association p-values for eQTL

(center panels), -log10(p) association p-values for mQTL (bottom panels) with location

of genes at the FURIN locus, 200kb range. Highlighted is the region with the

methylation probe (15:91426667 +/- 20kb for cg24888049 in the FURIN locus, and

1:154839909 +/- 20kb for cg09359103 in the *KCNN3 locus*).

Figure S5. Simulation analysis with a shared causal variant between three studies

using a sample size of 10,000 individuals for GWAS trait, 300 for eQTL trait, and 300 for

mQTL trait. Y-axis is the posterior probability of a shared causal causal variant for all

three datasets (PPA.abc). **A.** X-axis is the mean LD in the region. **B**. X-axis is the

number of SNPs Q in the region. Number of simulations is 1,000.

Table S1. False positive rates under the null of no association with any traits and

different combination of sample sizes (N simulations = 1000). We compute the false

positives by counting the proportion of simulations under the null that passed a posterior

probability supporting the "Detected Hypothesis" at a particular threshold

(PPA>=threshold). PPA.comb is PPA.abc + PPA.ab + PPA.ab.c.

Table S2. False positive rates under the null of no association with any traits and when

the causal variant is masked (N simulations = 1000). We compute the false positives by

counting the proportion of simulations under the null that passed a posterior probability

supporting the "Detected Hypothesis" at a particular threshold (PPA>=threshold).

PPA.comb is PPA.abc + PPA.ab + PPA.ab.c.

Table S3. False positive rates using 1000 and 500,000 simulations and our true sample

sizes at a cut off of 80% (N simulations = 1000 and 500,000). We compute the false

positives by counting the proportion of simulations under the null that passed a posterior

probability supporting the "Detected Hypothesis" at a particular threshold

(PPA>=threshold). PPA.comb is PPA.abc + PPA.ab + PPA.ab.c.

Table S4. Misclassification rates and power to detect the correct hypothesis for

combination of sample sizes (N simulations = 1000). We simulated each hypothesis of

interest indicated under the column "TRUE", and compute the number of misclassified

calls under other 14 scenarios by counting the proportion of simulations that passed a

posterior probability supporting a different hypothesis from the one simulated at a

particular threshold (PPA>=cut). We estimated power to distinguish a particular

hypothesis from the others by counting the number of correctly identified cases out of

1000 simulations. PPA.comb is PPA.abc + PPA.ab + PPA.ab.c.

Table S5. Misclassification rates and power to detect the correct hypothesis for

combination of sample sizes when the causal SNP is masked (N simulations = 1000).

The column labels are the same as Table S4.

Table S6. Misclassification rates and power to detect the correct hypothesis (N

simulations = 1000) under the sample sizes of 82,315 for GWAS, 467 for eQTL, and

121 for mQTL. The column labels are the same as Table S4.

Table S7. Effect of prior probability of co-localization for hypotheses of interest (GEM,

GE, GE.M) on the prior and posterior support for colocalisation in regions where there is

evidence for association across the three dataests (PPA.GE + PPA.GE.M + PPA.GEM

+ PPA.G.E + PPA.G.EM + PPA.G.E.M). We repeated the moloc analysis with different

combinations of priors p2 and p3 while keeping p1 fixed to 1×10^{-4}. The posteriors with

support for colocalization (PPA.coloc = PA.GE + PPA.GE.M + PPA.GEM) are

contrasted to the ones with support against colocalization (PPA.nocoloc = PPA.G.M

+PPA.G.EM+PPA.G.E.M) for our hypothesis of interest. For each analysis in which

PPA.GE + PPA.GE.M + PPA.GEM + PPA.G.E + PPA.G.EM + PPA.G.E.M >= 0.8, we

report the mean relative posterior support for colocalization, defined by PPA.coloc /

(PPA.nocoloc + PPA.coloc), and the mean relative prior support, averaged across all

regions. We find that the combination of priors p1=1e-04, p2=1e-06, p3=1e-07 has the

lowest difference between the prior and posterior expectation.

Table S8. 39 genes falling within SCZ LD blocks with a high probability of influencing

SCZ through expression or methylation (GEM, GE, GE.M>=0.8). "CHR", "START",

"STOP" indicate the positions of the regions considered; "PGC2_LDclump_info"

indicates position of the previously identified SCZ LD block; "PPA.combined" shows

maximum posterior probability reached by a gene for evidence of association with

GWAS and eQTL (PPA.GE + PPA.GE.M + PPA.GE.M); "numbMethyl" indicates the

number of methylation probes overlapping this region.

Table S9. Potentially novel SCZ loci. 45 candidate genes outside of the SCZ LD blocks

with a high probability of influencing SCZ (GEM, GE, GE.M>=0.8).

Table S10. Methylation-gene expression pairs that colocalize (PPA.GEM + PPA.EM + PPA.G.EM >= 0.8) for DNA methylation probes that are proximal to the transcription start site and enhancer sequences (20kb upstream to 2kb downstream of transcription start site). "PPA.m" is PPA.GEM + PPA.EM + PPA.G.EM.

Text S1. Expanded methods, derivations and analyses.

1 Models

For three traits, the possible combinations of causal configurations (binary vectors with at most 1 entry equal to 1) can be assigned to one of the following 15 hypotheses listed below. Each hypothesis can be represented by a set of index set containing the traits each causal SNP associates with.

- $H_0 \longmapsto \{\phi\}$: No association in any trait

- $H_1 \longmapsto \{\{1\}\}$: One SNP is associated with trait 1 (a)

- $H_2 \longmapsto \{\{2\}\}$: One SNP is associated with trait 2 (b)

- $H_3 \longmapsto \{\{3\}\}$: One SNP is associated with trait 3 (c)

- $H_4 \longmapsto \{\{1,2\}\}$: One SNP is associated with trait 1 and 2 (ab)

- $H_5 \longmapsto \{\{2,3\}\}$: One SNP is associated with trait 2 and 3 (bc)

- $H_6 \longmapsto \{\{1,3\}\}$: One SNP is associated with trait 1 and 3 (ac)

- $H_7 \longmapsto \{\{1\},\{2\}\}$: One SNP is associated with trait 1, and a different SNP is associated with trait 2 (a.b)

- $H_8 \longmapsto \{\{2\},\{3\}\}$: One SNP is associated with trait 2, and a different SNP is associated with trait 3 (b.c)

- $H_9 \longmapsto \{\{1\},\{3\}\}$: One SNP is associated with trait 1, and a different SNP is associated with trait 3 (a.c)

- $H_{10} \longmapsto \{\{1\},\{2,3\}\}$: One SNP is associated with trait 1, and a different SNP is associated with trait 2 and 3 (a.bc)

- $H_{11} \longmapsto \{\{3\},\{1,2\}\}$: One SNP is associated with trait 3, and a different SNP is associated with trait 1 and 2 (c.ab)

- $H_{12} \longmapsto \{\{2\},\{1,3\}\}$: One SNP is associated with trait 2, and a different SNP is associated with trait 1 and 3 (b.ac)

- $H_{13} \longmapsto \{\{1\},\{2\},\{3\}\}$: One SNP is associated with trait 1, a different SNP is associated with trait 2, and a different SNP is associated with trait 3 (a.b.c)

- $H_{14} \longmapsto \{\{1,2,3\}\}$: One SNP is associated with trait 1 and 2 and 3 (abc)

To illustrate our notation, consider a region with 4 SNPs, then the set of combinations of causal configurations of each trait consistent with $H_4 = \{\{1,2\}\}$ (one SNP associates with trait 1 and 2) is the following set of combinations of binary vectors

$$\left\{ \begin{pmatrix} 1000 \\ 1000 \\ 0000 \end{pmatrix}, \begin{pmatrix} 0100 \\ 0100 \\ 0000 \end{pmatrix}, \begin{pmatrix} 0010 \\ 0010 \\ 0000 \end{pmatrix}, \begin{pmatrix} 0001 \\ 0001 \\ 0000 \end{pmatrix} \right\}. \tag{1}$$

This in turn can be represented by a set of tuples, indicating which SNP associates with which trait,

$$\{\{(1,1),(1,2)\},\{(2,1),(2,2)\},\{(3,1),(3,2)\},\{(4,1),(4,2)\}\}. \tag{2}$$

Formally, let S_h denote the set of configurations supporting hypothesis h, then S_h can be written as

$$S_h = \left\{ \cup_{k=1}^{|h|} \left\{ i_{|h|} \right\} \times h_k \mid \forall \, i_1 ... i_{|h|} \in \{1, ..., Q\}, i_1 \neq \cdots \neq i_{|h|} \right\}, \tag{3}$$

where h_k represents the k-th element (an index set) in the set representation of the hypothesis h.

Next, we aim to assess the support for each hypothesis. We can compute the posterior probabilities given the data for each of these hypothesis by summing the probability of all the causal configurations consistent with the hypothesis. We find the probability of the data under each hypothesis using the following equation,

$$\frac{P(H_h|D)}{P(H_0|D)} = \sum_{S \in S_h} \frac{P(D \mid S)}{P(D \mid S_0)} \times \frac{P(S)}{P(S_0)}, \tag{4}$$

where the first ratio in this equation is a Bayes Factor for each configuration, and the second ratio is the prior odds of a configuration compared with the baseline configuration S_0.

We can now assess the evidence supporting each hypothesis. To compute the likelihood of the data and the prior probabilities of each hypothesis, we first compute a BF for each SNP and each trait. To connect these approximate Bayes factors to the alternative models for the SNP, we find the likelihood of the data and the prior probabilities π_s under the model that SNP i associates with traits indexed in $\mathbf{s}$.

1.1   Bayes Factor of a SNP with traits indexed in s

First, a Bayes Factor for each SNP and each trait 1 and 2 and 3 is computed from summary statistics using the Asymptotic Bayes Factor derivation (Wakefield [2009], equation 3 in main text). This measures the relative support for a model in which the SNP is associated with the trait compared to the null model of no association.

Second, to assess the evidence for a SNP with j traits, let $\mathbf{s}$ be the set of trait indices for which SNP i is associated with. For example, $s = \{\phi\}$ means SNP i is associated with no trait; and $s = \{\{1, 2\}\}$ means SNP i is associated with trait 1 and 2. In this case, the BF for association of a SNP i with the two traits is:

$$BF_{i,\{1,2\}} = \sum_{i=1}^{Q} WABF_i^1 WABF_i^2 \tag{5}$$

In general, the strength of association of a SNP i with all traits indexed in $\mathbf{s}$ is:

$$BF_{i,\mathbf{s}} = \prod_{j \in s} BF_i^j \tag{6}$$

1.2   Prior Probabilities that SNP i associates with traits in s

In our three-trait model, the prior probability that SNP i associates with all traits indexed in a set is:

- π_ϕ: SNP i associates with no trait

- $\pi_{\{1\}}$: SNP i associates with trait 1

- $\pi_{\{2\}}$: SNP i associates with trait 2

- $\pi_{\{3\}}$: SNP i associates with trait 3

- $\pi_{\{1,2\}}$: SNP i associates with trait 1 and 2

- $\pi_{\{1,3\}}$: SNP i associates with trait 1 and 3

- $\pi_{\{2,3\}}$: SNP i associates with trait 2 and 3

- $\pi_{\{1,2,3\}}$: SNP i associates with trait 1 and 2 and 3

These are mutually exclusive, so we have:

$$\pi_\phi + \pi_{\{1\}} + \pi_{\{2\}} + \pi_{\{3\}} + \pi_{\{1,2\}} + \pi_{\{1,3\}} + \pi_{\{2,3\}} + \pi_{\{1,2,3\}} = 1 \tag{7}$$

1.3 Prior probabilities of each hypothesis

$$P(H_h \mid D) \propto \sum_{S \in S_h} P(D \mid S)P(S) \propto P(S \mid S \in S_h) \times \sum_{S \in S_h} P(D \mid S) \tag{8}$$

where $P(S)$ is the prior probability of a configuration, $P(D \mid S)$ is the probability of the observed data D given a configuration S, and the sum is over all configurations S which are consistent with a given hypothesis H_h, where $h = (1 \ldots 14)$. Since each binary vector that belongs to the same set S_h has the same prior probability, we can multiply likelihoods by one common prior probability for each hypothesis.

The prior probability of any one configuration in the different sets is:

- if $S \in S_0$, then $P(S) = \pi_\phi^Q$

- if $S \in S_1$, then $P(S) = Q \times \pi_0^{Q-1} \pi_{\{1\}}$

- if $S \in S_2$, then $P(S) = Q \times \pi_0^{Q-1} \pi_{\{2\}}$

- if $S \in S_3$, then $P(S) = Q \times \pi_0^{Q-1} \pi_{\{3\}}$

- if $S \in S_4$, then $P(S) = Q \times \pi_0^{Q-1} \pi_{\{1,2\}}$

- if $S \in S_5$, then $P(S) = Q \times \pi_0^{Q-1} \pi_{\{2,3\}}$

- if $S \in S_6$, then $P(S) = Q \times \pi_0^{Q-1} \pi_{\{1,3\}}$

- if $S \in S_7$, then $P(S) = Q(Q-1) \times \pi_0^{Q-2} \pi_{\{1\}} \pi_{\{2\}}$

- if $S \in S_8$, then $P(S) = Q(Q-1) \times \pi_0^{Q-2} \pi_{\{1\}} \pi_{\{3\}}$

- if $S \in S_9$, then $P(S) = Q(Q-1) \times \pi_0^{Q-2} \pi_{\{2\}} \pi_{\{2\}}$

- if $S \in S_{10}$, then $P(S) = Q(Q-1) \times \pi_0^{Q-2} \pi_{\{1\}} \pi_{\{2,3\}}$

- if $S \in S_{11}$, then $P(S) = Q(Q-1) \times \pi_0^{Q-2} \pi_{\{3\}} \pi_{\{1,2\}}$

- if $S \in S_{12}$, then $P(S) = Q(Q-1) \times \pi_0^{Q-2} \pi_{\{2\}} \pi_{\{1,3\}}$

- if $S \in S_{13}$, then $P(S) = Q(Q-1)(Q-2) \times \pi_0^{Q-3} \pi_{\{1\}} \pi_{\{2\}} \pi_{\{3\}}$

- if $S \in S_{14}$, then $P(S) = Q \times \pi_0^{Q-1} \pi_{\{1,2,3\}}$

1.4 Probability of the data under each hypothesis (the 3-trait special case)

Putting it all together. For three traits, the probability of the data under each hypothesis are:

- $\frac{P(H_0|D)}{P(H_0|D)} = 1$

- $\frac{P(H_1|D)}{P(H_0|D)} = \pi_{\{1\}} \times \sum_{i=1}^{Q} BF_{i,\{1\}}$

- $\frac{P(H_2|D)}{P(H_0|D)} = \pi_{\{2\}} \times \sum_{i=1}^{Q} BF_{i,\{2\}}$

- $\frac{P(H_3|D)}{P(H_0|D)} = \pi_{\{3\}} \times \sum_{i=1}^{Q} BF_{i,\{3\}}$

- $\frac{P(H_4|D)}{P(H_0|D)} = \pi_{\{1,2\}} \times \sum_{i=1}^{Q} BF_{i,\{1,2\}}$

- $\frac{P(H_5|D)}{P(H_0|D)} = \pi_{\{2,3\}} \times \sum_{i=1}^{Q} BF_{i,\{2,3\}}$

- $\frac{P(H_6|D)}{P(H_0|D)} = \pi_{\{1,3\}} \times \sum_{i=1}^{Q} BF_{i,\{1,3\}}$

- $\frac{P(H_7|D)}{P(H_0|D)} = \pi_{\{1\}} \times \pi_{\{2\}} \times \sum_{i=1}^{Q} \sum_{j=1}^{Q} BF_{i,\{1\}} BF_{j,\{2\}} I[i \neq j]$

- $\frac{P(H_8|D)}{P(H_0|D)} = \pi_{\{2\}} \times \pi_{\{3\}} \times \sum_{i=1}^{Q} \sum_{j=1}^{Q} BF_{i,\{2\}} BF_{j,\{3\}} I[i \neq j]$

- $\frac{P(H_9|D)}{P(H_0|D)} = \pi_{\{1\}} \times \pi_{\{3\}} \times \sum_{i=1}^{Q} \sum_{j=1}^{Q} BF_{i,\{1\}} BF_{j,\{3\}} I[i \neq j]$

- $\frac{P(H_{10}|D)}{P(H_0|D)} = \pi_{\{1\}} \times \pi_{\{2,3\}} \times \sum_{i=1}^{Q} \sum_{j=1}^{Q} BF_{i,\{1\}} BF_{j,\{2\}} BF_{j,\{3\}} I[i \neq j]$

- $\frac{P(H_{11}|D)}{P(H_0|D)} = \pi_{\{3\}} \times \pi_{\{1,2\}} \times \sum_{i=1}^{Q} \sum_{j=1}^{Q} BF_{i,\{3\}} BF_{j,\{1\}} BF_{j,\{2\}} I[i \neq j]$

- $\frac{P(H_{12}|D)}{P(H_0|D)} = \pi_{\{2\}} \times \pi_{\{1,3\}} \times \sum_{i=1}^{Q} \sum_{j=1}^{Q} BF_{i,\{2\}} BF_{j,\{1\}} BF_{j,\{3\}} I[i \neq j]$

- $\frac{P(H_{13}|D)}{P(H_0|D)} = \pi_{\{1\}} \times \pi_{\{2\}} \times \pi_{\{3\}} \times \sum_{i=1}^{Q} \sum_{j=1}^{Q} \sum_{k=1}^{Q} BF_{i,\{1\}} BF_{j,\{2\}} BF_{k,\{3\}} I[i \neq j \neq k]$

- $\frac{P(H_{14}|D)}{P(H_0|D)} = \pi_{\{1\}} \times \pi_{\{2\}} \times \pi_{\{3\}} \times \sum_{i=1}^{Q} BF_{i,\{1\}} BF_{i,\{2\}} BF_{i,\{3\}}$

where $BF_{i,\mathbf{s}}$ are computed from Wakefield's Asymptotic Bayes Factors for each SNP and each trait 1, 2 and 3 as outlined in eq. 3, $I[i \neq j \neq k]$ is an indicator that evaluates to 1 if SNP i and j and k are different and 0 otherwise, π are the prior probabilities that SNP i is the causal SNP under a specific model.

In practice, we collapsed the prior probabilities to a smaller set for each kind of configuration. We set the prior probability that a SNP is causal in each trait to be identical $(\pi_{\{1\}} = \pi_{\{2\}} = \pi_{\{3\}})$ and refer to this a p1. We also set the prior probability that is associated with two traits to be identical $(\pi_{\{1,2\}}, \pi_{\{2,3\}}, \pi_{\{1,3\}})$ and refer to this as p2. We refer to the prior probability that SNP i the causal for all traits $(\pi_{\{1,2,3\}})$ as p3.

1.5 Probability of the data under hypotheses involving more than one independent associations

The probability of the data under scenarios of more than one causal variants $[7 \ldots 13]$ can be derived from the probability of the data under one causal variant $[1 \ldots 6, 14]$. This generalization speeds up computation significantly.

- $\frac{P(H_7|D)}{P(H_0|D)} = \pi_{\{1\}} \times \pi_{\{2\}} \times \sum_{i=1}^{Q} BF_{i,\{1\}} \sum_{j=1}^{Q} BF_{j,\{2\}} - \left[\frac{\pi_{\{1\}} \times \pi_{\{2\}}}{\pi_{\{1,2\}}} \times \frac{P(H_4|D)}{P(H_0|D)} \right]$

- $\frac{P(H_8|D)}{P(H_0|D)} = \pi_{\{2\}} \times \pi_{\{3\}} \times \sum_{i=1}^{Q} BF_{i,\{2\}} \sum_{j=1}^{Q} BF_{j,\{3\}} - \left[\frac{\pi_{\{2\}} \times \pi_{\{3\}}}{\pi_{\{2,3\}}} \times \frac{P(H_5|D)}{P(H_0|D)} \right]$

- $\frac{P(H_9|D)}{P(H_0|D)} = \pi_{\{1\}} \times \pi_{\{3\}} \times \sum_{i=1}^{Q} BF_{i,\{1\}} \sum_{j=1}^{Q} BF_{j,\{3\}} - \left[\frac{\pi_{\{1\}} \times \pi_{\{3\}}}{\pi_{\{1,3\}}} \times \frac{P(H_6|D)}{P(H_0|D)} \right]$

- $\frac{P(H_{10}|D)}{P(H_0|D)} = \pi_{\{1\}} \times \pi_{\{2,3\}} \times \sum_{i=1}^{Q} BF_{i,\{1\}} \sum_{j=1}^{Q} BF_{j,\{2,3\}} - \left[\frac{\pi_{\{1\}} \times \pi_{\{2,3\}}}{\pi_{\{1,2,3\}}} \times \frac{P(H_{14}|D)}{P(H_0|D)} \right]$

- $\frac{P(H_{11}|D)}{P(H_0|D)} = \pi_{\{3\}} \times \pi_{\{1,2\}} \times \sum_{i=1}^{Q} BF_{i,\{3\}} \sum_{j=1}^{Q} BF_{j,\{1,2\}} - \left[\frac{\pi_{\{3\}} \times \pi_{\{1,2\}}}{\pi_{\{1,2,3\}}} \times \frac{P(H_{14}|D)}{P(H_0|D)} \right]$

- $\frac{P(H_{12}|D)}{P(H_0|D)} = \pi_{\{2\}} \times \pi_{\{1,3\}} \times \sum_{i=1}^{Q} BF_{i,\{2\}} \sum_{j=1}^{Q} BF_{j,\{1,3\}} - \left[\frac{\pi_{\{2\}} \times \pi_{\{1,3\}}}{\pi_{\{1,2,3\}}} \times \frac{P(H_{14}|D)}{P(H_0|D)} \right]$

In general, across a set of j traits $\{1, 2, 3..., j\}$, for a hypothesis h, where there are more than one independent associations among the traits (i.e. $|h| > 1$), the probability of the data supporting h can be written as

$$\frac{P(H_h|D)}{P(H_0|D)} = \prod_{s \in h} \pi_s \sum_{i=1}^{Q} BF_{i,s} - \frac{\prod_{s \in h} \pi_s}{\pi_t} \sum_{i=1}^{Q} \pi_t BF_{i,t}, \tag{9}$$

where t is the union of the index set in h, i.e. $t = \cup_{j=1}^{|h|} h_j$.

For example, let there be three traits $\{1, 2, 3\}$ with two independent associations. The first signal colocalizes with traits 1 and 2 while the second is an independent association with trait 3. Hence, $j = 3$, $h = \big\{ \{1, 2\}, \{3\} \big\}$ and:

$$\frac{P(H_h|D)}{P(H_0|D)} = \pi_{\{1,2\}} \pi_{\{3\}} \sum_{i=1}^{Q} BF_{i\{1,2\}} \sum_{j=1}^{Q} BF_{j\{3\}} - \frac{\pi_{\{1,2\}} \pi_{\{3\}}}{\pi_{\{1,2,3\}}} \sum_{i=1}^{Q} \pi_{\{1,2,3\}} BF_{i\{1,2,3\}} \tag{10}$$

Since we pre-compute all the single terms in the above equation, the computations are greatly simplified.

Similarly, let there be five traits $\{1, 2, 3, 4, 5\}$ with two independent associations. First one is common to traits 1, 2, the second one common to traits 4 and 5. Hence, $M = 5$, $h = \{\{1, 2\}, \{4, 5\}\}$ and:

$$\frac{P(H_h|D)}{P(H_0|D)} = \pi_{\{1,2\}} \pi_{\{4,5\}} \sum_{i=1}^{Q} BF_{i\{1,2\}} \sum_{j=1}^{Q} BF_{j\{4,5\}} - \frac{\pi_{\{1,2\}} \pi_{\{4,5\}}}{\pi_{\{1,2,4,5\}}} \sum_{i=1}^{Q} \pi_{\{1,2,4,5\}} BF_{i\{1,2,4,5\}}$$

$$\tag{11}$$

References

Claudia Giambartolomei, Damjan Vukcevic, Eric E Schadt, Lude Franke, Aroon D Hingorani, Chris Wallace, and Vincent Plagnol. Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS genetics*, 10:e1004383, May 2014. ISSN 1553-7404. doi: 10.1371/journal.pgen.1004383.

Jon Wakefield. Bayes factors for genome-wide association studies: comparison with p-values. *Genetic epidemiology*, 33:79–86, January 2009. ISSN 0741-0395. doi: 10.1002/gepi.20359.

Xiaoquan Wen and Matthew Stephens. Bayesian methods for genetic association analysis with heterogeneous subgroups: from meta-analyses to gene-environment interactions. *The annals of applied statistics*, 8(1):176, 2014.

Supplementary Figures

Figure S1. Simulations under different sample sizes for all scenarios in *moloc* of

three traits (GWAS, eQTL, mQTL). All cases include 10,000 individuals in the

GWAS dataset. The variance explained by the trait was set to 0.01 for GWAS (1%), and to 0.1 (10%) for the eQTL and mQTL.

Figure S2. Simulations under different sample sizes for all scenarios in *moloc* of three traits ("a" is GWAS 1, "b" is GWAS 2, "c" is eQTL). All cases include 10,000 individuals in the biomarker dataset. The variance explained by the trait was set to 0.01 for GWAS 1 and GWAS 2, and to 1% (10%) for the eQTL.

Figure S3. Stratified Q-Q plots for eQTL and mQTLs in the SCZ GWAS. Enrichment of each functional category (blue line: eQTL; green: mQTL; red: combined eQTL and mQTL) illustrate the estimated increase in terms of the proportion of p-values expected in each scenario and in the functionally unannotated variant (FUV) category (grey line). The estimated enrichment for eQTL and mQTL is similar.

Figure S4. Illustration of two example of colocalization results with GWAS-eQTL-mQTL in the *FURIN* locus and cg24888049 and *KCNN3* and cg09359103. –log10(p) association p-values for GWAS (top panels), –log10(p) association p-values for eQTL (center panels), -log10(p) association p-values for mQTL (bottom panels) with location of genes at the FURIN locus, 200kb range. Highlighted is the region with the methylation probe (15:91426667 +/- 20kb for cg24888049 in the FURIN locus, and 1:154839909 +/- 20kb for cg09359103 in the *KCNN3 locus*).

A.   B.

Figure S5. Simulation analysis with a shared causal variant between three studies using a sample size of 10,000 individuals for GWAS trait, 300 for eQTL trait, and 300 for mQTL trait. Y-axis is the posterior probability of a shared causal causal variant for all three datasets (PPA.abc). **A.** X-axis is the mean LD in the region. **B**. X-axis is the number of SNPs Q in the region. Number of simulations is 1,000.