Suppression of drug resistance reveals a genetic mechanism of metabolic

plasticity in malaria parasites

Running title: Mechanism of malaria metabolic plasticity

Ann M. Guggisberg[1]

Philip M. Frasse[1]

Andrew J. Jezewski[1]

Natasha M. Kafai[2]

Aakash Y. Gandhi[1]*

Samuel J. Erlinger[1]†

Audrey R. Odom John[1,3]#

[1] Department of Pediatrics, Washington University School of Medicine, St. Louis,

Missouri, USA

[2] Medical Scientist Training Program, Washington University School of Medicine, St.

Louis, Missouri, USA

[3] Department of Molecular Microbiology, Washington University School of Medicine, St.

Louis, Missouri, USA

* Current affiliation: Aakash Gandhi, Medical Scientist Training Program, UT

Southwestern Medical Center, Dallas, Texas, USA

† Current affiliation: Samuel Erlinger, Medical Scientist Training Program, Johns

Hopkins University, Baltimore, Maryland, USA

[#] Address correspondence to aodom@wustl.edu

ABSTRACT

In the malaria parasite *Plasmodium falciparum*, synthesis of isoprenoids from glycolytic

intermediates is essential for survival. The antimalarial fosmidomycin (FSM) inhibits

isoprenoid synthesis. In *P. falciparum*, we identify a loss-of-function mutation in *HAD2*

(PF3D7_1226300) as necessary for FSM resistance. Enzymatic characterization

reveals that HAD2, a member of the haloacid dehalogenase-like hydrolase (HAD)

superfamily, is a phosphatase. Harnessing a growth defect in resistant parasites, we

select for suppression of HAD2-mediated FSM resistance and uncover hypomorphic

suppressor mutations in the locus encoding the glycolytic enzyme phosphofructokinase

(PFK9). Metabolic profiling demonstrates that FSM resistance is achieved via increased

steady-state levels of MEP pathway and glycolytic intermediates and confirms reduced

PFK9 function in the suppressed strains. We identify HAD2 as a novel regulator of

malaria parasite metabolism and drug sensitivity and uncover PFK9 as a novel site of

genetic metabolic plasticity in the parasite. Our study informs the biological functions of

an evolutionarily conserved family of metabolic regulators and reveals a previously

undescribed strategy by which malaria parasites adapt to cellular metabolic

dysregulation.

IMPORTANCE

Unique and essential aspects of parasite metabolism are excellent targets for

development of new antimalarials. An improved understanding of parasite metabolism

and drug resistance mechanisms are urgently needed. The antibiotic fosmidomycin

targets the synthesis of essential isoprenoid compounds from glucose and is a

candidate for antimalarial development. Our study identifies a novel mechanism of drug

resistance and further describes a family of metabolic regulators in the parasite. Using a

novel forward genetic approach, we also uncover mutations that suppress drug

resistance in the glycolytic enzyme PFK9. Thus, we identify an unexpected genetic

mechanism of adaptation to metabolic insult that influences parasite fitness and

tolerance to antimalarials.

Keywords: Plasmodium, malaria, antimalarial agents, drug resistance, metabolism,

isoprenoids, fosmidomycin, suppressor, glycolysis, metabolic regulation

INTRODUCTION

Malaria remains a global health threat, infecting 216 million people per year and causing

nearly half a million deaths, mainly of pregnant women and young children (1).

Resistance to current therapies has limited control efforts for malaria (2, 3). New drugs

and a deeper understanding of drug resistance mechanisms are urgently needed.

Malaria is caused by infection with unicellular eukaryotic parasites of the genus

Plasmodium. The species *Plasmodium falciparum* is responsible for most life-

threatening malarial disease. As an obligate intracellular parasite of human

erythrocytes, *Plasmodium falciparum* has unique metabolic features that may be

exploited to discover new drug targets and develop new therapies. In the red blood cell

niche, *Plasmodium* parasites are highly dependent on glucose metabolism. Infection

with *Plasmodium* spp. results in a nearly 100-fold increase in glucose import in red

blood cells (4–6). Despite these energy requirements, the parasite demonstrates little

aerobic respiration via the TCA cycle. Instead, it relies on anaerobic glycolysis to

produce ATP (7–10).

Besides ATP production, glucose also has a number of anabolic fates in *P. falciparum*.

One such fate is the synthesis of isoprenoids. Isoprenoids are a large class of

hydrocarbons with extensive structural and functional diversity (11). In the malaria

parasite, isoprenoids perform several essential functions, including protein prenylation,

dolichylation, and synthesis of GPI anchors (12–14). Despite this diversity, all

isoprenoids are synthesized from a common five-carbon building block, isopentyl

pyrophosphate (IPP). Evolution has produced two distinct routes for IPP synthesis: the

mevalonate pathway, found in archaea, fungi, animals, and the cytoplasm of plants; and

the methylerythritol phosphate (MEP) pathway, found in most eubacteria, plant

chloroplasts, and apicomplexan parasites such as *P. falciparum* (15). Because it is both

essential for the parasite and absent from the human host, the MEP pathway is a

compelling target for antimalarial development. The antibiotic and antimalarial

fosmidomycin (FSM) is a competitive inhibitor of the first committed enzymatic step of

the MEP pathway, catalyzed by 1-deoxy-D-xylulose-5- phosphate reductoisomerase

(DXR, E.C. 1.1.1.267) (16–18). FSM has been validated as a specific inhibitor of the

MEP pathway in *P. falciparum* (19) and is a valuable chemical tool to study MEP

pathway biology and essential metabolism in the parasite. In this study, we find that

FSM is also a useful tool for probing glycolytic metabolism upstream of the essential

MEP pathway.

Parasites are likely to control the proportion of glucose used for energy production

versus production of secondary metabolites, such as isoprenoids. We previously used a

screen for FSM resistance to identify HAD1, a metabolic regulator whose loss results in

increased levels of MEP pathway intermediates and resistance to MEP pathway

inhibition. HAD1 is a cytoplasmic sugar phosphatase that dephosphorylates a number

of sugar phosphate intermediates upstream of the MEP pathway (20, 21). HAD1

belongs to the haloacid dehalogenase-like hydrolase (HAD) enzyme superfamily and

more specifically, the IIB and Cof-like hydrolase subfamilies (22). While HADs are found

in all kingdoms of life, HAD1 is most related to bacterial members of this superfamily

(20, 23), which have been implicated in metabolic regulation, stress response, and

phosphate homeostasis (24–28). However, most members of this superfamily remain

uncharacterized.

In this study, we describe the discovery of HAD2, a second HAD family member in *P.*

falciparum. We find that HAD2 is a cytosolic phosphatase required for metabolic

homeostasis. Loss of HAD2 dysregulates glycolysis and misroutes metabolites toward

the MEP pathway, conferring drug resistance. In our study, we harness a fitness defect

in *had2* parasite strains to employ an innovative screen for suppression of drug

resistance in the parasite. Selection for suppression of drug resistance identifies

mutations in *PFK9*, which encodes the canonical glycolytic regulatory enzyme,

phosphofructokinase. Reduction in PFK9 activity rescues the metabolic dysregulation in

our resistant mutants and restores FSM sensitivity. Our unique approach thus reveals

PFK9 as a site of exceptional metabolic plasticity in the parasite and uncovers a novel

genetic mechanism by which *P. falciparum* malaria parasites may adapt to metabolic

stress and drug selective pressure.

This article was submitted to an online preprint archive (29).

RESULTS

A FSM[R] strain possesses a nonsense allele of *HAD2*, homolog of the MEP

pathway regulator HAD1

The MEP pathway is responsible for the synthesis of the essential isoprenoid

precursors isopentenyl pyrophosphate (IPP) and dimethylallyl pyrophosphate (DMAPP).

This pathway is specifically inhibited by the antibiotic FSM (19, 30, 31). We previously

generated *P. falciparum* strains resistant to FSM. Mutations in *HAD1* (PF3D7_1033400)

cause the resistance phenotype in a majority of these strains (20). However, the genetic

and biochemical basis of FSM resistance in strain E2 remained unknown. As previously

reported, we find that E2 is less sensitive to FSM than its wild-type 3D7 parental line

(Fig. 1A) (20). Strain E2 has a FSM IC_{50} of 4.8 ± 1.2 μM, significantly greater than that

of its parent strain (0.9 ± 0.06 μM) (p≤0.01, unpaired Student's t-test).

We find that this resistance phenotype is not due to changes in expression of the genes

encoding the first two, rate-limiting, steps of the MEP pathway, *DXS* and *DXR* (32–35).

In addition, strain E2 does not have genetic changes in the known FSM resistance locus

and MEP pathway regulator, *HAD1*, nor are there changes in HAD1 expression by

immunoblot (Figs. S1A and S1B).

To identify new genetic changes that may result in FSM resistance, we performed whole

genome sequencing on strain E2 and identified an A469T mutation in PF3D7_1226300

(PlasmoDB ID), hereafter referred to as *HAD2* (36). Variant data for strains sequenced

in this study can be found in Dataset S1. Sanger sequencing of the *HAD2* locus in strain

E2 confirmed the presence of the A469T SNP. The A469T single nucleotide

polymorphism (SNP) yields a truncated (R157X) protein variant and therefore we expect

HAD2 function is lost in strain E2. Interestingly, HAD2 is a close homolog of the known

MEP pathway regulator, the sugar phosphatase HAD1 (20). Sequence homology places

both proteins in the haloacid dehalogenase-like hydrolase (HAD) superfamily and

further within the IIB and Cof-like hydrolase subfamilies (Interpro IPR006379 and

IPR000150, respectively) (22). While no structural information exists for *P. falciparum*

HAD2, the structure of the *Plasmodium vivax* HAD2 (PVX_123945, PvHAD2) has been

solved (PDB ID 2B30). PvHAD2 (93% identical and 98% similar to PfHAD2) contains

the common structural motifs found in other HADs, including a core and cap domain

(Fig. 1B). HAD2 possesses the four conserved sequence motifs found in HAD proteins

(Fig. 1C), which are involved in binding of the substrate, coordination of the phosphoryl

group and Mg^{2+} ion, and hydrolysis of the substrate phosphate (37–39). Overall, HAD2

and HAD1 protein sequences share ~29% sequence identity and ~53% sequence

similarity (Fig. 1C). We hypothesized that HAD2, like HAD1, regulates metabolism in *P.*

falciparum, and that loss of HAD2-mediated metabolic control was responsible for FSM-

resistance in malaria parasite strain E2.

HAD2 is a functional phosphometabolite phosphatase

We have previously established that *P. falciparum* HAD1 is a promiscuous sugar

phosphatase, with activity against a wide range of phosphometabolites. Similarly, *P.*

vivax HAD2 has been enzymatically characterized and found to possess phosphatase

activity against various monophosphorylated substrates, including glycerol 2-phosphate

(glc2P) and pyridoxal phosphate (PLP) (40). Recombinant PvHAD2 also utilizes

additional monophosphorylated substrates, such as adenosine 5'-monophosphate

(AMP) and glycerol 1-phosphate (glc1P), with moderate activity.

Based on the previous characterization of a close *Plasmodium* homolog, as well as

sequence homology to HAD1 and other HAD proteins, we predicted that PfHAD2 would

also function enzymatically as a phosphatase. We successfully purified recombinant

PfHAD2 in *E. coli* and confirmed the phosphatase activity of recombinant PfHAD2 using

para-nitrophenyl phosphate (*p*NPP), a promiscuous, chromogenic phospho-substrate

(Fig. 2A) (23, 41). Because *E. coli* expresses a number of HAD-like phosphatases (23),

we confirmed that the phosphatase activity was specific to purified PfHAD2 by

expression and purification of a catalytically inactive mutant (HAD2^{D26A}). The Asp26

residue was chosen for mutagenesis as the corresponding residue in PfHAD1 (Asp27)

has been previously shown to be required for catalysis (21).

We also established the activity of PfHAD2 against a panel of phosphorylated

substrates and determined that its substrate profile closely mirrors that of PvHAD2 (Fig.

2B). Overall, we find that PfHAD2 is a phosphatase with activity against small

phosphosubstrates, such as *p*NPP and glc2P. These data suggest that, like HAD1 and

related HADs in microbes and plants (23, 42–44), HAD2 is a promiscuous phosphatase

with a preference for a variety of monophosphorylated phosphometabolites.

In vitro evolution of mutations suppressing FSM resistance

During routine culturing of E2 FSM^{R} parasites, we observed that the E2 strain is growth-

attenuated compared to its parental parasite strain. Surprisingly, during prolonged

culture in the absence of FSM, this growth phenotype resolved, and improved growth

rates correlated with a return to FSM sensitivity (Fig. 3A). From these observations, we

hypothesized that $had2^{R157X}$-mediated FSM resistance led to a fitness cost in cultured

parasites. We sought to harness this fitness cost to drive *in vitro* evolution of a FSM-

sensitive (FSM^S) population, possessing additional, novel mutations that might suppress

FSM resistance in $had2^{R157X}$ parasite strains.

FSM-resistant strain E2 was cultured through multiple passages in the absence of FSM

selection. Through limiting dilution, we derived five E2-based clones in the absence of

drug pressure (Fig. 3B). Three of the five clones remained FSM^R (designated clones

E2-R1, -R2, and -R3), but two of these were found to be FSM^S (designated E2-S1 and -

S2) (Fig. 3B and 4A). To validate our novel suppressor screen approach, we

independently repeated this genetic selection with the three FSM^R E2 clones by again

culturing in the absence of FSM for >1 month (Fig. 3B). As before, these strains (E2-S3,

-S4, and -S5) also lost their FSM resistance phenotype (Fig. 4A).

Consistent with our initial observation that our $had2^{R157X}$ FSM resistant strain grows

poorly, we find that the FSM^R clone E2-R1 grows at a significantly reduced rate

compared to the parental strain, while two FSM^S clones (E2-S1 and E2-S3) have

restored growth rates, similar to that of wild-type (Fig. 4B).

Loss of FSM resistance might have occurred by reversion of the $had2^{R157X}$ mutation in

E2-derived strains. Instead, we find that all E2-SX clones maintained loss of HAD2 via

the $had2^{R157X}$ mutation. We hypothesized that the FSM^S E2 clones, driven by a fitness

advantage, had acquired new suppressor mutation(s) at an additional locus. We

performed whole genome sequencing on the original five E2 clones to identify any

genetic changes that segregated with FSM-sensitivity. Sequencing revealed that a new

mutation (C3617T) in the locus encoding phosphofructokinase-9 (*PFK9*,

PF3D7_0915400) is present in both suppressed (FSM^S) E2 clones and none of the

three FSM^R E2 clones (Fig. 3B, Dataset S1). The C3617T mutation results in a

PFK9^{T1206I} protein variant. *PFK9* contained the only SNP that segregated with the

change in FSM tolerance. Two other loci had indels that also segregated with our FSM

phenotype. These loci encode a tyrosine recombinase (MAL13P1.42) (45) and an

erythrocyte surface protein (PIESP1, PFC0435w). Given their predicted functions and

the presence of A/T indels in poly-A and poly-T tracts, we concluded that mutations in

these loci were unlikely to result in our suppressed phenotype and prioritized *PFK9* as

the likely locus of our suppressor mutation.

To verify whether mutations in *PFK9* were responsible for suppressing FSM resistance

in all of our suppressed strains, we investigated *HAD2* and *PFK9* in the E2-S3, -S4, and

-S5 strains, which were derived through independent evolution of the E2-R1, -R2, and -

R3 populations in the absence of FSM. By Sanger sequencing, we find that, as before,

all strains maintain the *had2*^{R157X} mutation and acquire new, independent *PFK9*

mutations (Fig. 4A). The independent acquisition of four different *PFK9* alleles during

selection, each of which was associated with both improved growth and loss of FSM

sensitivity, strongly indicates that loss of PFK9 function is responsible for these

phenotypes in strains lacking HAD2.

Loss of HAD2 is necessary for FSM-resistance in *had2^R157X* parasites

HAD2 was not the sole genetic change in FSM[R] strain E2. In addition, because

intraerythrocytic *P. falciparum* parasites are haploid, we cannot distinguish recessive

from dominant or gain-of-function mutations. Therefore, we sought to establish whether

restoring HAD2 expression in *trans* in a *had2^R157X* strain would restore FSM sensitivity.

Using a previously described expression system enabled by the piggyBac transposase

(20, 46), we expressed HAD2-GFP driven by the *heat shock protein 110* (*Hsp110*)

promoter (47). We confirmed that the transfected *had2^R157X* E2-R2 clone maintains the

had2^R157X allele at the endogenous locus and successfully expresses HAD2-GFP (Fig.

5A). Expression of HAD2-GFP in *had2^R157X* parasites results in restoration of FSM

sensitivity (Fig. 5B), confirming that loss of HAD2 is necessary for FSM resistance in

this strain. The resistant clone (E2-R2) has a FSM IC_{50} of 3.9 ± 0.2 μM, significantly

higher than that of the wild-type parent strain (0.9 ± 0.06 μM, p≤0.001, one-way ANOVA

and Sidak's post-test). Expression of HAD2-GFP in E2-R2 results in an IC_{50} of 0.6 ±

0.02 μM for FSM, significantly lower than the E2-R2 strain (p≤0.001) but not significantly

different than the parental strain (p>0.5).

Using fluorescence microscopy, we also investigated the localization of HAD2-GFP in

our E2-R2 Hsp110:HAD2-GFP strain. We observe that HAD2-GFP is diffusely present

throughout the cytoplasm in asexual *P. falciparum* trophozoites and schizonts but

excluded from the digestive food vacuole (Fig. S2). This finding is consistent with the

lack of a predicted signal sequence for HAD2 using SignalP, PlasmoAP, and PlasMit

algorithms (48–50).

PFK9 mutations in suppressed strains are hypomorphic

The *PFK9* locus encodes the enzyme phosphofructokinase (PFK, E.C. 2.7.11), which

catalyzes the first committed and canonically rate-limiting step of glycolysis, which is the

conversion of fructose 6-phosphate to fructose 1,6-bisphosphate. PFK9 is comprised of

two domains, alpha (α) and beta (β), which are typically encoded by independent genes

in non-apicomplexans (51) (Fig. 6A). While in other systems the α domain is regulatory,

previous work on *P. falciparum* PFK9 has demonstrated catalytic activity for both

domains (51–55).

Of the four PFK9 variants identified in this study, three variants map to the α domain,

while one variant (S335L) maps to the β domain (Fig. 6A). We projected our mutations

on a three-dimensional model of PfPFK9 to reveal a possible structural basis for altered

PFK function. Three variants (S335L, T1206I, and S1267L) align and model to currently

available crystal structures of PFK, while a fourth allele (N1359Y) does not. While three

out of four mutations map to the α domain of PfPFK9 (T1206I, S1267L, N1359Y), these

mutations do not appear to cluster in any particular region. All mutations affect amino

acid residues that are physically distant from the substrate-binding pocket of either

domain and are not predicted to impact binding or specific catalytic residues.

Consistent with a previous study on PfPFK9 (51), attempts to purify recombinant full-

length PFK9 were unsuccessful. To then understand the enzymatic impacts of our

PFK9 variants, we quantified the native PFK specific activity in *P. falciparum* (51, 56)

(Fig. 6B, Fig. S3). Lysates from strains possessing *PFK9* mutations (E2-SX strains)

have markedly reduced specific activity of PFK compared to the parental strain (Fig.

6B). Combined with the diverse mutation locations (Fig. 6A), the reduced lysate PFK

activity in E2-SX strains suggests that a variety of genetic strategies to alter PFK

function can lead to resistance suppression.

Metabolic profiling reveals mechanisms of resistance and suppression in HAD2

and PFK9 mutant parasites

Reduced activity of PFK9, which catalyzes the canonical rate-limiting step in glycolysis,

is associated with restored FSM sensitivity to *had2* mutant strains. Therefore, we

anticipated that metabolic changes might underlie both resistance and suppression in

our E2 clones. We performed targeted metabolic profiling on the parental parasite strain

as well as E2 clones R1-R3 and S1-S2 (Fig. 7A, Table S1). We find that levels of the

MEP pathway intermediate DOXP are significantly increased in FSM^R (*had2^{R157X}*,

PFK9^{WT}) strains (Fig. 7A, p≤0.05, one-way ANOVA and Sidak's post-test). FSM is

competitive with DOXP for inhibition of its target enzyme DXR. Therefore, our data

indicate that the FSM^R strains achieve resistance via increased levels of DOXP, which

outcompetes FSM. We also observe a significant increase in the downstream MEP

metabolite, MEcPP, in our FSM^R strains (p≤0.05).

To understand the role of PFK9 in conferring and suppressing FSM resistance, we

determined the steady-state levels of intermediates from glycolysis, metabolites of

which feed into the MEP pathway (Fig. 7A, Table S1). Hierarchical clustering indicates

that resistant clones are characterized by a metabolic signature of increased levels of

FBP, DOXP, and MEcPP (Fig. 7A). We observe that the abundance of DOXP and

MEcPP are tightly correlated with cellular levels of the PFK9 product, FBP (Fig. 7B,

$p<0.01$), but not with the other upstream glycolytic metabolites, such as glu6P/fru6P

(Fig. 7C, $p>0.2$).

Of note, the $pfk9^{T1206I}$ suppressor allele in strains S1 and S2 restores nearly parental

levels of FBP and downstream MEP pathway intermediates (Fig. 7A), consistent with

our finding that PFK activity is reduced in lysate from these strains (Fig. 6B).

DISCUSSION

Cells must control levels of critical metabolites in order to efficiently utilize carbon

sources for energy and biosynthesis of essential molecules. Cells may regulate their

metabolism via transcriptional, post-transcriptional, post-translational, allosteric, or

enzymatic mechanisms that are necessary for growth (57–60). In the glucose-rich red

blood cell niche, *Plasmodium* spp. malaria parasites display a unique dependence on

glycolysis for energy and biosynthesis.

Using resistance to a metabolic inhibitor, we identify a phosphatase member of the HAD

superfamily, HAD2, as a novel regulator of metabolism in *P. falciparum*. Importantly,

HAD2 controls substrate availability to the parasite-specific MEP pathway for isoprenoid

synthesis, which is a promising drug target for much needed new antimalarials. We find

that tolerance to inhibitors such as FSM is a robust and sensitive readout of metabolic

perturbation. HAD2 is necessary for metabolic homeostasis in malaria parasites. Cells

lacking HAD2 exhibit marked dysregulation of central carbon metabolism, including

altered steady-state levels of glycolytic intermediates and isoprenoid precursors. We

find that mutations in phosphofructokinase (*PFK9*) restore wild-type growth rates and

FSM sensitivity to our *had2* mutant strains. Our study thus genetically connects the

function of HAD2, a HAD superfamily member, to control of essential central carbon

metabolism. In addition, our work reveals a previously undescribed strategy by which

malaria parasites may respond to cellular metabolic dysregulation through mutation in

the gene encoding the rate-limiting glycolytic enzyme PFK9.

HAD2 is a member of the HAD superfamily and a homolog of the previously described

metabolic regulator HAD1. With our previous studies on HAD1 (20, 21), we define the

cellular role of these proteins in *P. falciparum* and contribute to the greater

understanding of the HADs, an evolutionarily conserved and widespread protein family.

Both enzymes belong to the IIB (IPR006379) and Cof-like hydrolase (IPR000150)

subfamilies (22). HAD enzymes display diverse substrate preferences (23, 27, 43, 44,

61–64), and their biological functions are largely unknown. Like other HAD homologs,

including PfHAD1 (20), HAD2 appears to be a non-specific, cytoplasmic phosphatase

with preference for small, monophosphorylated substrates. While the HAD superfamily

is thought to be highly evolvable pool of enzymes with broad substrate specificity (37,

42), our work strongly suggests that, like HAD1 and HAD2, other members of this

superfamily are likely to perform specific and biologically important cellular functions. As

P. falciparum has a smaller repertoire of HADs than bacteria, and *Plasmodium* HADs

influence easily quantified phenotypes (drug tolerance, growth, metabolite levels), the

malaria parasite may be an attractive system for study of the molecular mechanisms by

which HAD proteins control metabolic homeostasis and growth.

Metabolic profiling reveals that loss of HAD2 function leads to metabolic dysregulation,

which is centered around the canonical rate-limiting step of glycolysis, catalyzed by

PFK9. While the cellular abundance of the PFK9 product FBP is increased in *had2*

parasites, HAD2 does not directly utilize FBP as an enzymatic substrate, suggesting an

indirect mechanism of HAD2-mediated metabolic regulation. However, the distinct

metabolic signature of *had2* parasites, characterized by increased levels of the MEP

pathway metabolites DOXP and MEcPP and the key glycolytic metabolite FBP,

suggests that MEP pathway metabolism is precisely linked to FBP production. In other

microbial systems, FBP levels reflect metabolic flux and are cued to environmental

perturbations (65). FBP-centered metabolic regulation is also important to the related

apicomplexan *Toxoplasma gondii*, which constitutively expresses the fructose 1,6-

bisphosphatase (FBPase) to fine-tune glucose metabolism (57). While *P. falciparum*

does not appear to possess an FBPase, necessary for gluconeogenesis, the parasite

may possess alternative FBP-sensing mechanisms to tune metabolism, perhaps via

regulators such as HAD1 and HAD2.

The metabolic dysregulation we observe in the *had2* mutant strains appears to come

with a fitness disadvantage. Under FSM-selective pressure, the benefits of dysregulated

metabolism outweigh the costs. However, in the absence of FSM, *had2* parasites

achieve metabolic relief through secondary mutation in *PFK9*. The improved growth of

had2 pfk9 double mutant parasites, compared to parasites with a *had2* single mutation,

argues that both the growth and metabolic phenotypes are linked. However, our

complementation studies cannot strictly discern whether restoring HAD2 directly

improves the growth rate of the *had2* strain, as transfection and complementation of

HAD2 is inherently an additional selection for increased fitness. Of note, two recent

essentiality screens in *Plasmodium* spp. have found that HAD2 is dispensable for

growth, and loss of HAD2 was not found to be associated with any significant fitness

defect (66, 67). However, it is unknown whether the mutant strains generated in these

screens have also acquired additional suppressor mutations, such as polymorphisms in

PFK9, that have facilitated their growth.

Likewise, *PFK9* provides an additional case-study of the context dependence of gene

essentiality in *Plasmodium* spp. As the canonical rate-limiting step in glycolysis, *PFK9* is

strongly predicted to be essential for asexual growth of malaria parasites (66, 67). In the

context of a *had2* mutation, our strains readily develop mutations in *PFK9* that reduce

function but are nonetheless associated with increased fitness. Indeed, it is surprising

that parasites that are entirely dependent on glycolysis for ATP production tolerate such

a significant loss of activity in this enzyme. Because we identify mutations across the

length of *PFK9* in our suppressed strains, our studies do not appear to point to a

specific disrupted function, such as alterations in an allosteric binding pocket or a dimer

interface. The observed mutability of *PFK9* points to a remarkable and unexpected

metabolic plasticity in the parasite. That is, even though the parasite inhabits a highly

controlled intraerythrocytic niche, a wide range of metabolic states of *P. falciparum*

growth are still permissive for parasite growth. This previously undescribed metabolic

plasticity centered on PFK9 should be considered in future efforts to target essential

metabolism in *Plasmodium*.

Combined with the above study, our work highlights the central role of the glycolytic

enzyme PFK9. A recent kinetic model of parasite glycolysis confirms that PFK has a

high flux-control coefficient, is sensitive to competitive inhibition, and can effectively

reduce glycolytic flux (68, 69). Like HAD2, PFK9 is plant-like and evolutionarily

divergent from its mammalian homologs (51). These differences may be exploited for

PFK inhibitor design and may indicate that PFK9 can be specifically targeted for

antimalarial development. However, our work cautions that the parasite has a surprising

capacity to adapt to perturbations in central carbon metabolism, which may present

challenges when targeting these pathways.

Finally, our approach demonstrates the power of forward genetics to uncover novel

biology in a clinically relevant, non-model organism. We employ a previously described

screen (20) to uncover a novel resistance locus and employ a second selection for

parasite fitness to identify changes that suppress our resistance phenotype. Of the 19

strains in our original FSM resistance screen (20), we identify only one *had2* mutant,

likely due to the reduced fitness associated with resistance in this strain. While fitness

costs associated with antimalarial resistance are well known (70–73), this study

represents, to our knowledge, the first to harness this evolutionary trade-off to identify

suppressor mutations in a non-target locus. Additional methods to identify low fitness

resistant mutants have recently been recently described (73), and fitness assessment of

resistance mutations may allow for suppressor screening for other antimalarials or other

target pathways to reveal new aspects of biology and drug resistance in malaria

parasites.

MATERIALS AND METHODS

Parasite strains and culture

Unless otherwise indicated, parasites were maintained at 37 °C in 5% O_2/5% CO_2/90%

N_2 in a 2% suspension of human erythrocytes in RPMI medium (Sigma Aldrich)

modified with 27 mM $NaHCO_3$, 11 mM glucose, 5 mM HEPES, 0.01 mM thymidine, 1

mM sodium pyruvate, 0.37 mM hypoxanthine, 10 µg/mL gentamycin, and 5 g/L Albumax

(Thermo Fisher Scientific).

FSM[R] strain E2 was generated by selecting a clone of genome reference strain 3D7

(MRA-102, MR4, ATCC, Manassas, VA) under continuous treatment with 3 µM FSM, as

previously described (20). Clones of strain E2 were isolated by limiting dilution.

Quantification of FSM resistance

Opaque 96-well plates were seeded with asynchronous cultures at 0.5-1.0%

parasitemia (percent of infected red blood cells). After 3 days, media was removed and

parasitemia was measured via Picogreen fluorescence on a POLARStar Omega

spectrophotometer (BMG Labtech), as previously described (74). Half maximal

inhibitory concentration (IC_{50}) values were calculated using Graphpad Prism. Unless

otherwise indicated, all IC_{50} data are representative of means from at least ≥3

independent experiments performed with technical replicates.

HAD2 structural alignment

Structures were aligned using the TM-align algorithm in Lasergene Protean 3D software

(root mean square deviation (RMSD) of 1.9 Å).

Whole genome sequencing and variant analysis

Library preparation, sequencing, read alignment, and variant analyses were performed

by the Washington University Genome Technology Access Center. One microgram of

parasite genomic DNA was sheared, end repaired, and adapter ligated. Libraries were

sequenced on an Illumina HiSeq 2500 in Rapid Run mode to generate 101 bp paired

end reads. Reads were aligned to the *P. falciparum* 3D7 reference genome (PlasmoDB

v7.2) using Novoalign (V2.08.02). Duplicate reads were removed. SNPs were called

using samtools (quality score ≥20, read depth ≥5) and annotated using snpEff.

Background variants were removed using previously sequenced genomes from parental

and control strains (20). Mixed variant calls and variants in highly variable surface

antigen loci (75, 76) were removed.

Sanger sequencing

The *HAD2* (PlasmoDB PF3D7_1226300) A469T (R157X) SNP was amplified and

sequenced using the HAD2_R157X_F and HAD2_R157X_R primers. The *PFK9* locus

was amplified using the PFK9_F and PFK9_R primers. *PFK9* amplicons were

sequenced using the PFK9_seq (1-8) primers. Primer sequences can be found in Table

S2.

Generation of recombinant HAD2

The predicted coding sequence of *HAD2* was amplified using the HAD2_LIC_F and

HAD2_LIC_R primers (Table S2). A catalytic mutant (D26A) was also generated to use

as a negative control in activity assays. The *had2^{D26A}* allele was created using the

HAD2_D26A_F and HAD2_D26A_R site-directed mutagenesis primers (Table S2).

Ligation-independent cloning was used to clone *HAD2* and *had2^{D26A}* into vector BG1861

(77), which introduces an N-terminal 6xHis fusion to the expressed protein.

BG1861:6xHis-HAD2 was transformed into One Shot BL21(DE3)pLysS *Escherichia coli*

cells (Thermo Fisher Scientific). Protein expression was induced for 3 hours with 1 mM

isopropyl-β-D-thiogalactoside at mid-log phase (OD$_{600}$ 0.4-0.5). Cells were collected by

centrifugation and stored at -20°C.

Cells were lysed in buffer containing 1 mg/mL lysozyme, 20 mM imidazole, 1 mM

dithiothreitol, 1 mM MgCl$_2$ 10 mM Tris HCl (pH 7.5), 30 U benzonase (EMD Millipore),

and EDTA-free protease inhibitor tablets (Roche). 6xHis-HAD2 was bound to nickel

agarose beads (Gold Biotechnology), washed with 20 mM imidazole, 20 mM Tris HCl

(pH 7.5), and 150 mM NaCl and eluted in 300 mM imidazole, 20 mM Tris HCl (pH 7.5),

and 150 mM NaCl. This eluate was further purified by size-exclusion gel

chromatography using a HiLoad 16/600 Superdex 200 pg column (GE Healthcare)

equilibrated in 25 mM Tris HCl (pH 7.5), 250 mM NaCl, and 1 mM MgCl$_2$. The elution

fractions containing HAD2 were pooled and concentrated and glycerol was added to

10% (w/v). Protein solutions were immediately flash frozen and stored at -80°C.

HAD2 activity assays

The rate of *para*-nitrophenyl phosphate (*p*NPP; Sigma-Aldrich S0942) hydrolysis by

HAD2 was determined by continuous measurement of absorbance at 405 nm. Assays

were performed at 37°C in 50 µl with 50 mM Tris-HCl, pH 7.5, 5 mM MgCl$_2$, 10 mM

*p*NPP, and 1.2 µM enzyme.

Hydrolysis of other phosphorylated substrates by HAD2 was measured using the

EnzChek phosphate assay kit (Life Technologies). Reaction buffer was modified to

contain 50 mM Tris-HCl, 1 mM MgCl$_2$, pH 7.5, 0.1 mM sodium azide, 0.2 mM 2-amino-

6-mercapto-7-methylpurine riboside (MESG), and 1 U/mL purine nucleoside

phosphorylase (PNP). Reactions contained 5 mM substrate and 730 nM enzyme.

Activity was normalized to that obtained from catalytically inactive 6xHis-HAD2^{D26A}. All

data are means from ≥3 independent experiments performed with technical replicates.

P. falciparum **growth assays**

Asynchronous cultures were seeded at 1% parasitemia. Media (no drug) was

exchanged daily. Samples were taken at indicated timepoints and fixed in PBS + 4%

paraformaldehyde, 0.05% glutaraldehyde. Cells were stained with 0.01 mg/ml acridine

orange and parasitemia was determined on a BD Biosciences LSRII flow cytometer

(Thermo Fisher Scientific). All data are means from ≥3 independent experiments.

pTEOE110:HAD2 plasmid construction

The pTEOE110 construct contains the *heat shock protein 110* (PF3D7_0708800) 5'

UTR and a C-terminal GFP tag (20). Human dihydrofolate reductase (hDHFR) is

present as a selectable marker. Inverted terminal repeats are included for genome

integration by a co-transfected piggyBac transposase (pHTH, MRA912, MR4, ATCC,

Manassas, VA).

HAD2 was amplified with the HAD2_XhoI_F and HAD2_AvrII_R primers (Table S2) and

cloned into AvrII and XhoI sites in the pTEOE110 plasmid.

Parasite transfections

Transfections were performed as previously described (20). Briefly, 50-100 µg of

plasmid DNA was precipitated and resuspended in Cytomix (25 mM HEPES pH 7.6,

120 mM KCl, 0.15 mM $CaCl_2$, 2 mM EGTA, 5 mM $MgCl_2$, 10 mM K_2HPO_4).

A ring-stage *P. falciparum* culture was washed with Cytomix and resuspended in the

DNA/Cytomix solution. Cells were electroporated using a BioRad Gene Pulser II

electroporator at 950 µF and 0.31 kV. Electroporated cells were washed with media and

returned to normal culture conditions. Parasites expressing the construct were selected

by continuous treatment with 5 nM WR92210 (Jacobus Pharmaceuticals). Transfectants

were cloned by limiting dilution and presence of the HAD2-GFP construct was verified

by PCR using gene- and GFP-specific primers (HAD2_R157X_F and GFP_R, Table

S2). Maintenance of the endogenous *HAD2* and *PFK9* genotypes were verified by

Sanger sequencing.

Antiserum generation

Polyclonal anti-HAD2 antiserum was raised in rabbits against 6xHis-HAD2, with

Titermax as an adjuvant (Cocalico Biologicals). Antiserum specificity was confirmed by

immunoblot of lysate lacking HAD2. Polyclonal anti-HAD1 antiserum has been

previously described (MRA-1256, MR4, ATCC) (20).

Immunoblotting

Lysates were separated on a polyacrylamide gel and transferred to a polyvinylidene

difluoride membrane. Membranes were blocked in 5% non-fat dry milk, 0.1% Tween-20

in PBS. Rabbit polyclonal antisera were used at the following dilutions: 1:2,000-5,000

anti-HAD2 and 1:20,000 anti-HAD1 (20). For all blots, 1:20,000 HRP-conjugated goat

anti-rabbit IgG antibody was used as a secondary antibody (ThermoFisher 65-6120).

Blots were stripped with 200 mM glycine, 0.1% SDS, 1% Tween-20, pH 2.2 and re-

probed with 1:5,000 rabbit anti-heat shock protein 70 (Hsp70) (AS08 371, Agrisera

Antibodies) as a loading control. All blots shown are representative of ≥3 independent

experiments. Minimal adjustments were applied equally to all blot images.

PfPFK model construction

PfPFK subunits were searched against the HHpred server for protein remote homology

detection and 3D structure prediction using statistics as previously described (78–81).

The *Borellia burgdorferii* PFK structure (PDB 1KZH) (82) returned the highest similarity

for both PfPFK domains and was used to predict 3D structure for each domain using the

program MODELLER. PFK product orientation in the active site of the model was

predicted via the alignment tool, using PyMOL software against the *E. coli* PFK crystal

structure (PDB 1PFK) (83). The α domain model encompasses amino acids 779-1347

and the β domain model encompasses amino acids 110-638.

Assay of native PFK9 activity

Sorbitol-synchronized trophozites were isolated using 0.1% saponin. Cells were washed

in buffer containing 100 mM Tris-HCl (pH 7.5), 1 mM $MgCl_2$, 1 mM DTT, 10% glycerol,

and EDTA-free protease inhibitor tablets (Roche) and lysed by sonication at 4°C (Fisher

Scientific Model 550 Sonic Dismembrator, amplitude 3.5), followed by centrifugation at

4°C (10,000 x g, 10 min). An "RBC carryover" control was comprised of the trace

cellular material remaining after saponin lysis, centrifugation, and wash of uninfected

erythrocytes.

Lysate PFK9 activity was monitored by linking it to the oxidation of NADH, as previously

described (51, 56). Reactions contained 100 mM Tris-HCl (pH 7.5), 1 mM $MgCl_2$, 1 mM

DTT, 0.25 mM NADH, 1 mM ATP, 3 mM fructose 6-phosphate, and excess of linking

enzymes aldolase (7.5 U), triose-phosphate isomerase (3.8 U), and glycerol 3-

phosphate dehydrogenase (3.8 U). After adding fresh cell lysate (10-15 µg total protein),

absorbance at 340 nm was measured at 37°C for 40 min. Activity was determined by

linear regression using Graphpad Prism software. Unless otherwise indicated, data are

means from ≥3 independent experiments.

Metabolite profiling

Approximately ~1 x 10^9 sorbitol-synchronized early trophozites were isolated using

0.1% saponin, washed with ice-cold PBS + 2 g/L glucose, and frozen at -80 °C.

Samples were extracted in 600 µL of ice-cold extraction solvent [chloroform, methanol,

and acetonitrile (2:1:1, v/v/v)] using two liquid-nitrogen cooled 3.2 mm stainless steel

beads and homogenization in a Tissue-Lyser II instrument (Qiagen) at 20 Hz for 5

minutes in a cold sample rack. Ice-cold water was added and samples were

homogenized for 5 min at 20 Hz. Samples were centrifuged at 14,000 rcf at 4°C for 5

min. The polar phase was lyophilized and re-dissolved in 100 µL water and analyzed by

LC-MS/MS. LC-MS/MS was performed on a 4000QTRAP system (AB Sciex) in multiple-

reaction monitoring mode using negative ioniziation and 10 mM tributylammonium

acetate (pH 5.1-5.5) as the ion pair reagent. The specific parameters used for analysis

of MEP pathway metabolites have been previously described (19). Liquid

chromatography separation was performed using ion pair reverse-phase

chromatography (84) with the following modifications: (1) RP-hydro 100 mm × 2.0 mm,

2.5 μm high performance liquid chromatography column (Phenomenex), (2) flow rate of

0.14 mL/min, (3) solvent A of 10 mM tributylammonium acetate in 5% methanol, (4)

binary LC gradient (20% solvent B (100% methanol) from 0 to 2.5 min, 30% B for 12.5

min, 80% B for 5 min, and column equilibration at for 5 minutes), and (5) 20 μL

autosampler injection volume.

Accession numbers

All genome data have been deposited in the NCBI BioProject database (PRJNA222697)

and Sequence Read Archive (SRP038937).

ACKNOWLEDGEMENTS

We thank the Genome Technology Access Center in the Department of Genetics at

Washington University School of Medicine for help with genomic analysis. The Center is

partially supported by NCI Cancer Center Support Grant #P30 CA91842 to the

Siteman Cancer Center and by ICTS/CTSA Grant# UL1TR002345 from the National

Center for Research Resources (NCRR), a component of the National Institutes of

Health (NIH), and NIH Roadmap for Medical Research. This publication is solely

the responsibility of the authors and does not necessarily represent the official view

of NCRR or NIH. We thank Sophie Alvarez and the Proteomics and Mass Spectrometry

Facility at the Donald Danforth Plant Science Center for performing the LC-MS/MS

sample prep and analysis. This work was supported by the National Science

Foundation under grant number DBI-0521250 for acquisition of the QTRAP LC-MS/MS.

We thank Daniel Goldberg (Washington University) for supplying the pTEOE110

plasmid. We thank the Malaria Research Reference and Reagent Resource (MR4) for

providing reagents contributed by D.J. Carucci (MRA-102) and John Adams (MRA-912).

We thank Katherine Andrews for sharing unpublished data relevant to this study.

REFERENCES

1.   World Health Organization. 2017. World Malaria Report 2017. Geneva,

Switzerland.

2.   Dondorp AM, Nosten F, Yi P, Das D, Phyo AP, Tarning J, Lwin KM, Ariey F,

Hanpithakpong W, Lee SJ, Ringwald P, Silamut K, Imwong M, Chotivanich K, Lim

P, Herdman T, An SS, Yeung S, Singhasivanon P, Day NP, Lindegardh N,

Socheat D WN. 2009. Artemisinin resistance in *Plasmodium falciparum* malaria. N

Engl J Med 455–467.

3.   Ashley EA, Dhorda M, Fairhurst RM, Amaratunga C, Lim P, Suon S, Sreng S,

Anderson JM, Mao S, Sam B, Sopha C, Chuor CM, Nguon C, Sovannaroth S,

Pukrittayakamee S, Jittamala P, Chotivanich K, Chutasmit K, Suchatsoonthorn C,

Runcharoen R, Hien TT, Thuy-Nhien NT, Thanh NV, Phu NH, Htut Y, Han K-T,

Aye KH, Mokuolu OA, Olaosebikan RR, Folaranmi OO, Mayxay M, Khanthavong

M, Hongvanthong B, Newton PN, Onyamboko MA, Fanello CI, Tshefu AK, Mishra

N, Valecha N, Phyo AP, Nosten F, Yi P, Tripura R, Borrmann S, Bashraheil M,

Peshu J, Faiz MA, Ghose A, Hossain MA, Samad R, Rahman MR, Hasan MM,

Islam A, Miotto O, Amato R, MacInnis B, Stalker J, Kwiatkowski DP, Bozdech Z,

Jeeyapant A, Cheah PY, Sakulthaew T, Chalk J, Intharabut B, Silamut K, Lee SJ,

Vihokhern B, Kunasol C, Imwong M, Tarning J, Taylor WJ, Yeung S, Woodrow

CJ, Flegg JA, Das D, Smith J, Venkatesan M, Plowe C V, Stepniewska K, Guerin

PJ, Dondorp AM, Day NP, White NJ. 2014. Spread of artemisinin resistance in

Plasmodium falciparum malaria. N Engl J Med 371:411–23.

4.   Mehta M, Sonawat HM, Sharma S. 2006. Glycolysis in *Plasmodium falciparum*

results in modulation of host enzyme activities. J Vector Borne Dis 43:95–103.

5.   Roth E. 1990. *Plasmodium falciparum* carbohydrate metabolism: a connection

between host cell and parasite. Blood Cells 16:453–66.

6.   Roth EF. 1987. Malarial parasite hexokinase and hexokinase-dependent

glutathione reduction in the *Plasmodium falciparum*-infected human erythrocyte. J

Biol Chem 262:15678–82.

7.   Bowman IB, Grant PT, Kermack WO, Ogston D. 1961. The metabolism of

Plasmodium berghei, the malaria parasite of rodents. 2. An effect of mepacrine on

the metabolism of glucose by the parasite separated from its host cell. Biochem J

78:472–8.

8.   Krungkrai J, Burat D, Kudan S, Krungkrai S, Prapunwattana P. 1999.

Mitochondrial oxygen consumption in asexual and sexual blood stages of the

human malarial parasite, *Plasmodium falciparum*. Southeast Asian J Trop Med

Public Health 30:636–42.

9.   MacRae JI, Dixon MW, Dearnley MK, Chua HH, Chambers JM, Kenny S, Bottova

I, Tilley L, McConville MJ. 2013. Mitochondrial metabolism of sexual and asexual

blood stages of the malaria parasite *Plasmodium falciparum*. BMC Biol 11:67.

10.   Scheibel LW, Miller J. 1969. Glycolytic and cytochrome oxidase activity in

Plasmodia. Mil Med 134:1074–80.

11.   Gershenzon J, Dudareva N. 2007. The function of terpene natural products in the

natural world. Nat Chem Biol 3:408–14.

12.   Guggisberg AM, Amthor RE, Odom AR. 2014. Isoprenoid biosynthesis in

Plasmodium falciparum. Eukaryot Cell 13:1348–59.

13.   Imlay L, Odom AR. 2014. Isoprenoid metabolism in apicomplexan parasites. Curr

Clin Microbiol Reports 1:37–50.

14.   Jordão FM, Kimura EA, Katzin AM. 2011. Isoprenoid biosynthesis in the

erythrocytic stages of *Plasmodium falciparum*. Mem Inst Oswaldo Cruz 106:134–

141.

15.   Lange BM, Rujan T, Martin W, Croteau R. 2000. Isoprenoid biosynthesis: the

evolution of two ancient and distinct pathways across genomes. Proc Natl Acad

Sci U S A 97:13172–7.

16.   Koppisch AT, Fox DT, Blagg BSJ, Poulter CD. 2002. *E. coli* MEP synthase:

steady-state kinetic analysis and substrate binding. Biochemistry 41:236–43.

17.   Kuzuyama T, Shimizu T, Takahashi S, Seto H. 1998. Fosmidomycin, a specific

inhibitor of 1-deoxy-d-xylulose 5-phosphate reductoisomerase in the

nonmevalonate pathway for terpenoid biosynthesis. Tetrahedron Lett 39:7913–6.

18.   Steinbacher S, Kaiser J, Eisenreich W, Huber R, Bacher A, Rohdich F. 2003.

Structural basis of fosmidomycin action revealed by the complex with 2-C-methyl-

D-erythritol 4-phosphate synthase (IspC). Implications for the catalytic mechanism

and anti-malaria drug development. J Biol Chem 278:18401–7.

19.   Zhang B, Watts KM, Hodge D, Kemp LM, Hunstad DA, Hicks LM, Odom AR.

2011. A second target of the antimalarial and antibacterial agent fosmidomycin

revealed by cellular metabolic profiling. Biochemistry 50:3570–7.

20.   Guggisberg AM, Park J, Edwards RL, Kelly ML, Hodge DM, Tolia NH, Odom AR.

2014. A sugar phosphatase regulates the methylerythritol phosphate (MEP)

pathway in malaria parasites. Nat Commun 5:4467.

21.   Park J, Guggisberg AM, Odom AR, Tolia NH. 2015. Cap-domain closure enables

diverse substrate recognition by the C2-type haloacid dehalogenase-like sugar

phosphatase *Plasmodium falciparum* HAD1. Acta Crystallogr D Biol Crystallogr

71:1824–34.

22.   Hunter S, Jones P, Mitchell A, Apweiler R, Attwood TK, Bateman A, Bernard T,

Binns D, Bork P, Burge S, de Castro E, Coggill P, Corbett M, Das U, Daugherty L,

Duquenne L, Finn RD, Fraser M, Gough J, Haft D, Hulo N, Kahn D, Kelly E,

Letunic I, Lonsdale D, Lopez R, Madera M, Maslen J, McAnulla C, McDowall J,

McMenamin C, Mi H, Mutowo-Muellenet P, Mulder N, Natale D, Orengo C,

Pesseat S, Punta M, Quinn AF, Rivoire C, Sangrador-Vegas A, Selengut JD,

Sigrist CJA, Scheremetjew M, Tate J, Thimmajanarthanan M, Thomas PD, Wu

CH, Yeats C, Yong S-Y. 2012. InterPro in 2011: new developments in the family

and domain prediction database. Nucleic Acids Res 40:D306-12.

23. Kuznetsova E, Proudfoot M, Gonzalez CF, Brown G, Omelchenko M V, Borozan

I, Carmel L, Wolf YI, Mori H, Savchenko A V, Arrowsmith CH, Koonin E V,

Edwards AM, Yakunin AF. 2006. Genome-wide analysis of substrate specificities

of the *Escherichia coli* haloacid dehalogenase-like phosphatase family. J Biol

Chem 281:36149–61.

24. Kang MJ, Lee YM, Yoon SH, Kim JH, Ock SW, Jung KH, Shin YC, Keasling JD,

Kim SW. 2005. Identification of genes affecting lycopene accumulation in

Escherichia coli using a shot-gun method. Biotechnol Bioeng 91:636–42.

25. Kang Y, Weber KD, Qiu Y, Kiley PJ, Blattner FR. 2005. Genome-wide expression

analysis indicates that FNR of *Escherichia coli* K-12 regulates a large number of

genes of unknown function. J Bacteriol 187:1135–1160.

26. Pratish G, Radhakrishnan M. 2014. Engineering *Escherichia coli* for D-ribose

production from glucose-xylose mixtures. Ind Biotechnol 10:106–114.

27. Roberts A, Lee S-Y, McCullagh E, Silversmith RE, Wemmer DE. 2005. YbiV from

Escherichia coli K12 is a HAD phosphatase. Proteins 58:790–801.

28. Sun Y, Vanderpool CK. 2013. Physiological consequences of multiple-target

regulation by the small RNA SgrS in *Escherichia coli*. J Bacteriol 195:4804–15.

29. Guggisberg AM, Gandhi AY, Erlinger SJ, Merino EF, Cassera MB, John ARO.

2017. Suppressor screen connects HAD protein function to metabolic control in

Plasmodium falciparum. bioRxiv 155523.

30. Yeh E, DeRisi JL. 2011. Chemical rescue of malaria parasites lacking an

apicoplast defines organelle function in blood-stage *Plasmodium falciparum*.

PLoS Biol 9:e1001138.

31. Jomaa H, Wiesner J, Sanderbrand S, Altincicek B, Weidemeyer C, Hintz M,

Türbachova I, Eberl M, Zeidler J, Lichtenthaler HK, Soldati D, Beck E. 1999.

Inhibitors of the nonmevalonate pathway of isoprenoid biosynthesis as

antimalarial drugs. Science 285:1573–6.

32. Estévez JM, Cantero A, Reindl A, Reichler S, León P. 2001. 1-Deoxy-D-xylulose-

5-phosphate synthase, a limiting enzyme for plastidic isoprenoid biosynthesis in

plants. J Biol Chem 276:22901–9.

33. Mahmoud SS, Croteau RB. 2001. Metabolic engineering of essential oil yield and

composition in mint by altering expression of deoxyxylulose phosphate

reductoisomerase and menthofuran synthase. Proc Natl Acad Sci U S A 98:8915–

20.

34. Carretero-Paulet L, Cairó A, Botella-Pavía P, Besumbes O, Campos N, Boronat

A, Rodríguez-Concepción M. 2006. Enhanced flux through the methylerythritol 4-

phosphate pathway in *Arabidopsis* plants overexpressing deoxyxylulose 5-

phosphate reductoisomerase. Plant Mol Biol 62:683–95.

35. Morrone D, Lowry L, Determan MK, Hershey DM, Xu M, Peters RJ. 2010.

Increasing diterpene yield with a modular metabolic engineering system in *E. coli*:

comparison of MEV and MEP isoprenoid precursor pathway engineering. Appl

Microbiol Biotechnol 85:1893–1906.

36. Aurrecoechea C, Brestelli J, Brunk BP, Dommer J, Fischer S, Gajria B, Gao X,

Gingle A, Grant G, Harb OS, Heiges M, Innamorato F, Iodice J, Kissinger JC,

Kraemer E, Li W, Miller JA, Nayak V, Pennington C, Pinney DF, Roos DS, Ross

C, Stoeckert CJ, Treatman C, Wang H. 2009. PlasmoDB: a functional genomic

database for malaria parasites. Nucleic Acids Res 37:D539-43.

37. Burroughs AM, Allen KN, Dunaway-Mariano D, Aravind L. 2006. Evolutionary

genomics of the HAD superfamily: understanding the structural adaptations and

catalytic diversity in a superfamily of phosphoesterases and allied enzymes. J Mol

Biol 361:1003–34.

38. Koonin E V., Tatusov RL. 1994. Computer analysis of bacterial haloacid

dehalogenases defines a large superfamily of hydrolases with diverse specificity.

Application of an iterative approach to database search. J Mol Biol 244:125–32.

39. Aravind L, Galperin MY, Koonin E V. 1998. The catalytic domain of the P-type

ATPase has the haloacid dehalogenase fold. Trends Biochem Sci 23:127–9.

40. Srinivasan B, Kempaiah Nagappa L, Shukla A, Balaram H. 2015. Prediction of

substrate specificity and preliminary kinetic characterization of the hypothetical

protein PVX_123945 from *Plasmodium vivax*. Exp Parasitol 151:56–63.

41. Kuznetsova E, Proudfoot M, Sanders SA, Reinking J, Savchenko A, Arrowsmith

CH, Edwards AM, Yakunin AF. 2005. Enzyme genomics: Application of general

enzymatic screens to discover new enzymes. FEMS Microbiol Rev 29:263–79.

42. Kuznetsova E, Nocek B, Brown G, Makarova KS, Flick R, Wolf YI, Khusnutdinova

A, Evdokimova E, Jin K, Tan K, Hanson AD, Hasnain G, Zallot R, de Crécy-

Lagard V, Babu M, Savchenko A, Joachimiak A, Edwards AM, Koonin E V,

Yakunin AF. 2015. Functional diversity of haloacid dehalogenase superfamily

phosphatases from *Saccharomyces cerevisiae*: biochemical, structural, and

evolutionary insights. J Biol Chem 290:18678–98.

43.   Titz B, Häuser R, Engelbrecher A, Uetz P, Alatossava T, Jutte H, Kuhn A,

Kellenberger E, Aravind L, Koonin EV, Baba T, Ara T, Hasegawa M, Takai Y,

Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL, Mori H, Bacon DF,

Treffers HP, Bateman A, Coin L, Durbin R, Baykov AA, Evtushenko OA, Avaeva

SM, Bessman MJ, Frick DN, O'Handley SF, Cohen SS, Flaks JG, Barner HD,

Loeb MR, Lichtenstein J, Cox EC, Yanofsky C, Datsenko KA, Wanner BL,

Galperin MY, Moroz OV, Wilson KS, Murzin AG, Guzman LM, Belin D, Carson

MJ, Beckwith J, Harbers E, Chaudhuri NK, Heidelberger C, Herrick DJ, Major PP,

Kufe DW, Kamiya H, Koonin EV, Tatusov RL, Kufe DW, Major PP, Kuznetsova E,

Proudfoot M, Gonzalez CF, Lorkiewicz Z, Szybalski W, Lusk JE, Williams RJ,

Kennedy EP, Major PP, Egan E, Herrick D, Kufe DW, Maki H, Sekiguchi M,

Maliekal P, Vertommen D, Delpierre G, Schaftingen E Van, Nakanishi T, Sekimizu

K, O'Brien KP, Remm M, Sonnhammer EL, O'Donovan GA, Neuhard J, Okazaki

R, Kornberg A, Outten CE, O'Halloran TV, Pratt S, Shepard RL, Kandasamy RA,

Johnston PA, Perry W, Dantzig AH, Proudfoot M, Kuznetsova E, Brown G, Rao

NN, Kitagawa M, Mori H, Savchenko A, Yakunin AF, Rosner A, Yagil E, Santi DV,

McHenry CS, Sommer H, Thompson J, Treffers HP, Spinelli V, Belser NO,

Uchiyama I, Wilkinson DS, Crumley J, Zhang J, Inouye M. 2007. The *Escherichia*

coli protein YjjG is a house-cleaning nucleotidase in vivo. FEMS Microbiol Lett

270:49–57.

44.   Caparrós-Martín JA, McCarthy-Suárez I, Culiáñez-Macià FA. 2013. HAD

810  hydrolase function unveiled by substrate screening: enzymatic characterization of

811  *Arabidopsis thaliana* subclass I phosphosugar phosphatase AtSgpp. Planta

812  237:943–54.

813 45. Ghorbal M, Scheidig-Benatar C, Bouizem S, Thomas C, Paisley G, Faltermeier C,

814  Liu M, Scherf A, Lopez-Rubio J-J, Gopaul DN. 2012. Initial characterization of the

815  Pf-Int recombinase from the malaria parasite *Plasmodium falciparum*. PLoS One

816  7:e46507.

817 46. Balu B, Shoue DA, Fraser MJ, Adams JH. 2005. High-efficiency transformation of

818  *Plasmodium falciparum* by the lepidopteran transposable element piggyBac. Proc

819  Natl Acad Sci U S A 102:16391–6.

820 47. Muralidharan V, Oksman A, Pal P, Lindquist S, Goldberg DE. 2012. *Plasmodium*

821  *falciparum* heat shock protein 110 stabilizes the asparagine repeat-rich parasite

822  proteome during malarial fevers. Nat Commun 3:1310.

823 48. Foth BJ, Ralph SA, Tonkin CJ, Struck NS, Fraunholz M, Roos DS, Cowman AF,

824  McFadden GI. 2003. Dissecting apicoplast targeting in the malaria parasite

825  *Plasmodium falciparum*. Science 299:705–8.

826 49. Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0:

827  discriminating signal peptides from transmembrane regions. Nat Methods 8:785–

828  6.

829 50. Zuegge J, Ralph S, Schmuker M, McFadden GI, Schneider G. 2001. Deciphering

830  apicoplast targeting signals--feature extraction from nuclear-encoded precursors

831  of *Plasmodium falciparum* apicoplast proteins. Gene 280:19–26.

832 51. Mony BM, Mehta M, Jarori GK, Sharma S. 2009. Plant-like phosphofructokinase

from *Plasmodium falciparum* belongs to a novel class of ATP-dependent

enzymes. Int J Parasitol 39:1441–53.

52. Bär J, Golbik R, Hübner G, Kopperschläger G. 2000. Denaturation of

phosphofructokinase-1 from *Saccharomyces cerevisiae* by guanidinium chloride

and reconstitution of the unfolded subunits to their catalytically active form.

Biochemistry 39:6960–8.

53. Kemp RG, Gunasekera D. 2002. Evolution of the allosteric ligand sites of

mammalian phosphofructo-1-kinase. Biochemistry 41:9426–30.

54. Klinder A, Kirchberger J, Edelmann A, Kopperschläger G. 1998. Assembly of

phosphofructokinase-1 from *Saccharomyces cerevisiae* in extracts of single-

deletion mutants. Yeast 14:323–34.

55. Theodorou ME, Cornel FA, Duff SM, Plaxton WC. 1992. Phosphate starvation-

inducible synthesis of the alpha-subunit of the pyrophosphate-dependent

phosphofructokinase in black mustard suspension cells. J Biol Chem 267:21901–

5.

56. Beutler E. 1984. Red Cell Metabolism: A Manual of Biochemical Methods. Grune

& Stratton, Philadelphia, PA.

57. Blume M, Nitzsche R, Sternberg U, Gerlic M, Masters SL, Gupta N, McConville

MJ. 2015. A *Toxoplasma gondii* gluconeogenic enzyme contributes to robust

central carbon metabolism and is essential for replication and virulence. Cell Host

Microbe 18:210–20.

58. Brown AJP, Brown GD, Netea MG, Gow NAR. 2014. Metabolism impacts upon

Candida immunogenicity and pathogenicity at multiple levels. Trends Microbiol

22:614–622.

59.   Jurica MS, Mesecar A, Heath PJ, Shi W, Nowak T, Stoddard BL. 1998. The

allosteric regulation of pyruvate kinase by fructose-1,6-bisphosphate. Structure

6:195–210.

60.   Kronstad J, Saikia S, Nielson ED, Kretschmer M, Jung W, Hu G, Geddes JMH,

Griffiths EJ, Choi J, Cadieux B, Caza M, Attarian R. 2012. Adaptation of

Cryptococcus neoformans to mammalian hosts: integrated regulation of

metabolism and virulence. Eukaryot Cell 11:109–18.

61.   Cabello-Díaz JM, Gálvez-Valdivieso G, Caballo C, Lambert R, Quiles FA, Pineda

M, Piedras P. 2015. Identification and characterization of a gene encoding for a

nucleotidase from *Phaseolus vulgaris*. J Plant Physiol 185:44–51.

62.   Caparrós-Martín JA, Reiland S, Köchert K, Cutanda MC, Culiáñez-Macià FA.

2007. *Arabidopsis thaliana* AtGppl and AtGpp2: two novel low molecular weight

phosphatases involved in plant glycerol metabolism. Plant Mol Biol 63:505–17.

63.   Huang H, Pandya C, Liu C, Al-Obaidi NF, Wang M, Zheng L, Toews Keating S,

Aono M, Love JD, Evans B, Seidel RD, Hillerich BS, Garforth SJ, Almo SC,

Mariano PS, Dunaway-Mariano D, Allen KN, Farelli JD. 2015. Panoramic view of

a superfamily of phosphatases through substrate profiling. Proc Natl Acad Sci U S

A 112:E1974-83.

64.   Liu J, Zhou W, Liu G, Yang C, Sun Y, Wu W, Cao S, Wang C, Hai G, Wang Z,

Bock R, Huang J, Cheng Y. 2015. The conserved endoribonuclease YbeY is

required for chloroplast ribosomal RNA processing in *Arabidopsis*. Plant Physiol

168:205–221.

65.   Kochanowski K, Volkmer B, Gerosa L, Haverkorn van Rijsewijk BR, Schmidt A,

880   Heinemann M. 2012. Functioning of a metabolic flux sensor in *Escherichia coli*.

881   Proc Natl Acad Sci 110:1130–1135.

882   66.   Bushell E, Gomes AR, Sanderson T, Anar B, Girling G, Herd C, Metcalf T,

883   Modrzynska K, Schwach F, Martin RE, Mather MW, McFadden GI, Parts L,

884   Rutledge GG, Vaidya AB, Wengelnik K, Rayner JC, Billker O. 2017. Functional

885   profiling of a *Plasmodium* genome reveals an abundance of essential genes. Cell

886   170:260–272.e8.

887   67.   Zhang M, Wang C, Otto TD, Oberstaller J, Liao X, Adapa SR, Udenze K, Bronner

888   IF, Casandra D, Mayho M, Brown J, Li S, Swanson J, Rayner JC, Jiang RHY,

889   Adams JH. 2018. Uncovering the essential genes of the human malaria parasite

890   Plasmodium falciparum by saturation mutagenesis. Science 360:eaap7847.

891   68.   Penkler G, du Toit F, Adams W, Rautenbach M, Palm DC, van Niekerk DD,

892   Snoep JL. 2015. Construction and validation of a detailed kinetic model of

893   glycolysis in Plasmodium falciparum. FEBS J 282:1481–511.

894   69.   van Niekerk DD, Penkler GP, du Toit F, Snoep JL. 2016. Targeting glycolysis in

895   the malaria parasite *Plasmodium falciparum*. FEBS J 283:634–46.

896   70.   Gabryszewski SJ, Dhingra SK, Combrinck JM, Lewis IA, Callaghan PS, Hassett

897   MR, Siriwardana A, Henrich PP, Lee AH, Gnädig NF, Musset L, Llinás M, Egan

898   TJ, Roepe PD, Fidock DA. 2016. Evolution of fitness cost-neutral mutant PfCRT

899   conferring *P. falciparum* 4-aminoquinoline drug resistance is accompanied by

900   altered parasite metabolism and digestive vacuole physiology. PLOS Pathog

901   12:e1005976.

71.   Hastings IM, Donnelly MJ. 2005. The impact of antimalarial drug resistance

mutations on parasite fitness, and its implications for the evolution of resistance.

Drug Resist Updat 8:43–50.

72.   Rosenthal PJ. 2013. The interplay between drug resistance and fitness in malaria

parasites. Mol Microbiol 89:1025–1038.

73.   Lim MY-X, LaMonte G, Lee MCS, Reimer C, Tan BH, Corey V, Tjahjadi BF, Chua

A, Nachon M, Wintjens R, Gedeck P, Malleret B, Renia L, Bonamy GMC, Ho PC-

L, Yeung BKS, Chow ED, Lim L, Fidock DA, Diagana TT, Winzeler EA, Bifani P.

2016. UDP-galactose and acetyl-CoA transporters as *Plasmodium* multidrug

resistance genes. Nat Microbiol 1:16166.

74.   Corbett Y, Herrera L, Gonzalez J, Cubilla L, Capson TL, Coley PD, Kursar TA,

Romero LI, Ortega-Barria E. 2004. A novel DNA-based microfluorimetric method

to evaluate antimalarial drug activity. Am J Trop Med Hyg 70:119–24.

75.   Su XZ, Heatwole VM, Wertheimer SP, Guinet F, Herrfeldt JA, Peterson DS,

Ravetch JA, Wellems TE. 1995. The large diverse gene family var encodes

proteins involved in cytoadherence and antigenic variation of *Plasmodium*

falciparum-infected erythrocytes. Cell 82:89–100.

76.   Kyes SA. 1999. Rifins: A second family of clonally variant proteins expressed on

the surface of red cells infected with *Plasmodium falciparum*. Proc Natl Acad Sci

U S A 96:9333–9338.

77.   Alexandrov A, Vignali M, LaCount DJ, Quartley E, de Vries C, De Rosa D,

Babulski J, Mitchell SF, Schoenfeld LW, Fields S, Hol WG, Dumont ME, Phizicky

EM, Grayhack EJ. 2004. A facile method for high-throughput co-expression of

protein pairs. Mol Cell Proteomics 3:934–8.

78.   Zimmermann L, Stephens A, Nam S-Z, Rau D, Kübler J, Lozajic M, Gabler F,

Söding J, Lupas AN, Alva V. 2017. A completely reimplemented MPI

bioinformatics toolkit with a new HHpred server at its core. J Mol Biol.

79.   Söding J. 2005. Protein homology detection by HMM-HMM comparison.

Bioinformatics 21:951–60.

80.   Hildebrand A, Remmert M, Biegert A, Söding J. 2009. Fast and accurate

automatic structure prediction with HHpred. Proteins 77 Suppl 9:128–32.

81.   Meier A, Söding J. 2015. Automatic prediction of protein 3D structures by

probabilistic multi-template homology modeling. PLoS Comput Biol 11:e1004343.

82.   Moore SA, Ronimus RS, Roberson RS, Morgan HW. 2002. The structure of a

pyrophosphate-dependent phosphofructokinase from the Lyme disease

spirochete *Borrelia burgdorferi*. Structure 10:659–671.

83.   Shirakihara Y, Evans PR. 1988. Crystal structure of the complex of

phosphofructokinase from *Escherichia coli* with its reaction products. J Mol Biol

204:973–94.

84.   Luo B, Groenke K, Takors R, Wandrey C, Oldiges M. 2007. Simultaneous

determination of multiple intracellular metabolites in glycolysis, pentose

phosphate pathway and tricarboxylic acid cycle by liquid chromatography-mass

spectrometry. J Chromatogr 1147:153–64.

Figure 1. FSM^R strain E2 possesses a mutation in HAD2, a homolog of the MEP

pathway regulator HAD1. (A) Representative FSM dose-response of the parental

strain and strain E2. **(B)** *P. vivax* HAD2 (teal, PDB 2B30) is structurally similar to

PfHAD1 (grey, PDB 4QJB). Ions (Mg^{2+}, Ca^{2+}, Cl^-) are shown in yellow. **(C)** HAD2 is a

homolog of HAD1 (29% identity and 53% similarity) and possesses all conserved HAD

sequence motifs required for catalysis (37).

Figure 2. PfHAD2 is a phosphatase. (A) HAD2 is an active phosphatase, and

HAD2^{D26A} is a catalytic mutant (cat. mut.) that can be used as a negative control for

HAD2 specific activity. "No enz" represents a no enzyme control. Data shown are the

enzyme activities using the synthetic phosphatase substrate *p*NPP. Error bars represent

S.E.M. (**** = p≤0.0001, unpaired t-test). **(B)** Activity of HAD2, normalized to the activity

of the catalytic mutant (HAD2^{D26A}), for variety of substrates (2-GlcP, 2-glycerol-

phosphate; M6P, mannose-6-phosphate; FBP, fructose-2,6-bisphosphate; dAMP,

deoxy-adenosine monophosphate; AMP, adenosine monophosphate). Error bars

represent S.E.M.

Figure 3. Leveraging a resistance-associated growth attenuation to identify

genetic changes that modulate FSM sensitivity.

(A) Prolonged culture results in loss of FSM resistance in strain E2. Shown are single

FSM dose-responses of the strain E2 before (9 days) and after (79 days) prolonged

culture without FSM. Nine days after thawing the resistant E2, we observe a FSM IC_{50}

of 4.9 µM, while after 79 days of culture without FSM, E2 has an IC_{50} of 1.3 µM. Each

data point is representative of the mean from two technical replicates. Error bars

represent S.E.M. **(B)** Parasites are colored by their FSM phenotype (teal, FSM^S; purple,

FSM^R). Cloned strains are named by FSM phenotype (E2-SX, sensitive; E2-RX,

resistant). A FSM^S parental strain was selected under FSM pressure to enrich for FSM^R

strain E2 ($had2^{R157X}$). After relief of FSM pressure, a fitness advantage selects for

spontaneous suppressor mutations in *PFK9* ($pfk9^{mut}$, yellow star) that result in FSM

sensitivity. FSM^R clones are grown without FSM pressure and a fitness advantage

again selects for suppressor mutations in *PFK9* that result in increased growth rate and

loss of FSM resistance.

Figure 4. Suppressor strains with *PFK9* mutations display changes in FSM

tolerance and growth.

(A) Suppressed clones have significantly lower FSM IC_{50}s (**** = p≤0.0001). Error bars

represent S.E.M. *HAD2* and *PFK9* genotypes for each strain are indicated. **(B)** FSM

resistance results in a fitness cost. A representative FSM^R clone with the $had2^{R157X}$

allele (R1, purple line) has a reduced growth rate compared to the wild-type parental

strain (black) (** = p≤0.01). The growth defect is rescued in two representative clones

with mutations in *PFK9* (S1 and S3, teal lines). Growth is normalized to parasitemia on

day 0. Error bars represent S.E.M. from independent growth experiments.

Figure 5. Loss of HAD2 is necessary for FSM resistance. (A) Successful expression

of pTEOE110:HAD2-GFP in strain R2 (*had2*^{R157X}, *PFK9*) was confirmed by immunoblot.

Marker units are kilodaltons (kDa). The top blot was probed with anti-HAD2 antiserum

(expected masses: HAD, 33 kDa; HAD2-GFP, 60 kDa). The bottom blot was probed

with anti-heat shock protein 70 (Hsp70) antiserum as a loading control. **(B)**

Representative FSM dose-response demonstrating expression of HAD2-GFP in strain

R2 (*had2*^{R157X}, *PFK9*) results in restored sensitivity to FSM. Strain R2 has an elevated

FSM IC$_{50}$ compared to the parental strain. When HAD2 expression is restored in strain

R2, the resulting strain has an IC$_{50}$ near that of the parent strain.

1003

1004

1005

Figure 6. *PFK9* alleles in suppressed strains are hypomorphic. (A) Schematic of

suppressor mutations found in *PFK9*. Resulting amino acid changes and strain names

are indicated. Three of the four mutations are found on the structural model of

PfPFK. The parts of the protein represented by the model are notated by the teal arrows

under the α and β domains. Total protein length is 1418 amino acids. N1359Y fits

outside of the model. The other mutations are represented by their stick model structure

with the resulting change shown in magenta. Orientations of the close-up

representations of the mutations are indicated when they differ from the main model. **(B)**

Measurement of PFK activity of *P. falciparum* lysate indicates that E2-SX clones with

PFK9 suppressor mutations have significantly reduced PFK activity (**** = p≤0.0001,

ANOVA, Sidak's post-test). Error bars represent S.E.M. Assay is linear with respect to

protein content and specific for PfPFK9 activity (S3 Fig).

1018

Figure 7. *HAD2* and *PFK9* alleles alter FSM resistance and metabolite levels in *P. falciparum*. (A) Metabolic profiling and clustering of parental and E2 clone strains demonstrates a metabolic signature of resistance, which includes increased levels of MEP pathway intermediates DOXP and MEcPP and the glycolytic metabolite and PFK product FBP. Glu6P/fru6P and DHAP/gly3P are isomer pairs that cannot be confidently distinguished. Clustering performed using the heatmap function in R. Data are also summarized in S1 Table. FSM IC_{50}s are shown for reference. **(B)** DOXP levels are highly correlated to levels of the upstream glycolytic metabolic FBP (Pearson's r=0.95). **(C)** In contrast, DOXP levels are not correlated to the glycolytic metabolites glu6P/fru6P (Pearson's r=0.57).

1031 **SUPPLEMENTAL MATERIAL LEGENDS**

1032

Supplemental Figure 1. FSM^R strain E2 does not have increased expression of

DXS, DXR,* or *HAD1.

Supplemental Figure 2. HAD2 is cytosolic.

Supplemental Figure 3. Assay of PFK activity from *P. falciparum* lysate is linear

and specific.

Supplemental Table 1. Relative metabolite levels in strains described in this

study.

Supplementary Table 2. Primers used in this study.