Sharing of heteroplasmies between human liver lobes varies across the mtDNA genome

Alexander Hübner[1,§,*], Manja Wachsmuth[1,§], Roland Schröder[1], Mingkun Li[2], Anna Maria Eis-Hübinger[3], Burkhard Madea[4] and Mark Stoneking[1,*]

[1] Department of Evolutionary Genetics, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, D-04103 Leipzig, Germany

[2]Key Laboratory of Genomic and Precision Medicine, Beijing Institute of Genomics, Chinese Academy of Sciences, 100101, Beijing, China

[3]Institut für Virologie, Universitätsklinikum Bonn, Sigmund-Freud-Str. 25, D-53105 Bonn, Germany

[4]Institut für Rechtsmedizin, Universitätsklinikum Bonn, Stiftsplatz 12, D-53111 Bonn, Germany

* To whom correspondence should be addressed. Tel: +49 (0) 341 3550-504; Email: stonekg@eva.mpg.de; alexander_huebner@eva.mpg.de

§ These authors contributed equally to the manuscript.

Keywords: mtDNA, heteroplasmy, liver, sharing

ABSTRACT

Mitochondrial DNA (mtDNA) heteroplasmy (intra-individual variation) varies among different human tissues and increases with age, suggesting that the majority of mtDNA heteroplasmies are acquired, rather than inherited. However, the extent to which heteroplasmic sites are shared across a tissue remains an open question. We therefore investigated heteroplasmy in two liver samples (one from each primary lobe) from 83 Europeans, sampled at autopsy. Minor allele frequencies (MAF) at heteroplasmic sites were significantly correlated between the two liver samples from an individual, with significantly more sharing of heteroplasmic sites in the control region than in the coding region. We show that this increased sharing for the control region cannot be explained by recent mutations at just a few specific heteroplasmic sites or by the possible presence of 7S DNA. Moreover, we carried out simulations to show that there is significantly more sharing than would be predicted from random genetic drift from a common progenitor cell. We also observe a significant excess of non-synonymous vs. synonymous heteroplasmies in the coding region, but significantly more sharing of synonymous heteroplasmies. These contrasting patterns for the control vs. the coding region, and for non-synonymous vs. synonymous heteroplasmies, suggest that selection plays a role in heteroplasmy sharing.

INTRODUCTION

The mitochondrial genome is present in many copies in a single cell, and inter-individual variation in the mitochondrial genome of an individual is called mtDNA heteroplasmy (Larsson 2010). In humans, it has been shown that detrimental mtDNA mutations are usually present in a heteroplasmic state at low frequencies, with high frequencies of the deleterious allele leading to functional defects and a disease phenotype (Larsson 2010; Wallace and Chalkia 2013; Stewart and Chinnery 2015). In addition, heteroplasmy is a general phenomenon in aging individuals, where the minor allele is present at rather low frequencies (often below 4%) and many of the affected sites are part of the control region (Stewart and Chinnery 2015). The

total number of heteroplasmic sites strongly correlates with age and several studies have

shown that heteroplasmic sites are tissue specific (Michikawa et al. 1999; Calloway et al. 2000;

Wang et al. 2001; Samuels et al. 2013; Li et al. 2015; Naue et al. 2015), i.e. sites which are

frequently heteroplasmic in one tissue are homoplasmic in all other tissues of the same

individual.

The tissue specificity of heteroplasmic sites and the association between the number of

heteroplasmies and age would suggest that the majority of heteroplasmies are not inherited

from the previous generation but are acquired during the lifetime of an individual in a tissue-

dependent manner. Therefore, the question arises, whether cells from the same tissue share

a similar profile of heteroplasmic sites despite being separated for many cell divisions? Studies

on mouse embryos containing two different mtDNA haplotypes have shown that mtDNA

segregation occurs rapidly between generations and the distribution of mtDNA haplotypes in

the F1 generation resembles a pattern expected under random genetic drift (Jenuth et al.

1996). This result is expected when the underlying heteroplasmies are evolving neutrally.

However, it has been shown that mtDNA variants do not behave fully neutrally (Nachman et

al. 1996; Jenuth et al. 1997) and more recent studies on human cells indicated that the

observed variance in heteroplasmic levels between cells is less stochastic than expected by

random genetic drift (Raap et al. 2012; Jayaprakash et al. 2015). This suggests that there

might be further population genetic forces, e.g. within-cell mtDNA population structure (Kowald

and Kirkwood 2011) or selection, that shape the distribution of heteroplasmies within as well

as between tissues.

While most age-related heteroplasmies occur in the control region, human liver tissue is

unusual in showing an excess of heteroplasmies involving non-synonymous mutations in the

mtDNA protein-coding genes (Li et al. 2015). This result is remarkable because these coding

region mutations are likely to have a functional effect (Li et al. 2015), and coding region

mutations are strongly selected against during transmission from one generation to another

(Stewart et al. 2008). Thus, it seems that mtDNA in human liver tissue exhibits a relaxation of

purifying selection and age-related positive selection for somatic mutations that decrease mitochondrial function (Li et al. 2015). This makes liver a good candidate tissue to analyze the sharing of heteroplasmic sites with respect to different evolutionary forces.

While some studies have investigated the amount of variation in levels of heteroplasmy in cells that arose from a single ancestor cell in cell culture (Raap et al. 2012; Jayaprakash et al. 2015), to date there has been no such investigation comparing heteroplasmy across a tissue. We therefore obtained one blood sample and two liver samples (one from each primary lobe) from 83 Europeans, sampled at autopsy. MtDNA heteroplasmy was evaluated by capture-enrichment sequencing (Li et al. 2010; Maricic et al. 2010; Li and Stoneking 2012; Li et al. 2015), and we analyzed sharing of mtDNA heteroplasmy between the liver lobes for different regions of the mitochondrial genome. We find a high correlation in the minor allele frequency (MAF) at heteroplasmic sites in the control region between the two liver samples, but a much weaker correlation in the coding region.

RESULTS

Heteroplasmy sharing within the liver

We investigated mitochondrial DNA heteroplasmy in liver and blood tissue samples of 83 individuals. For liver, two samples taken from different lobes were analyzed in order to compare the heteroplasmic pattern in different parts of the tissue. The samples were capture-enriched for mtDNA and sequenced to an average sequencing depth of 1,175-fold for blood samples and 2,640-fold for liver samples. Applying a threshold of 2.5 % MAF, we detected 541 heteroplasmic sites at 343 different positions (Supplementary Table S1). More heteroplasmic sites were observed in the coding region for liver (280 sites) compared to blood (64 sites), but the most abundant heteroplasmic sites in liver were in the control region (site 72: 67 individuals, site 60: 26 individuals, site 94: 20 individuals), which were only rarely observed in blood (site 72: 1 individual, site 60: 1 individual, site 94: no individuals). These

data are in accordance with results from a previous study ((Li et al. 2015), Supplementary

Figure S1), indicating that heteroplasmy is tissue specific, with different individuals exhibiting

similar heteroplasmic patterns.

Virological tests revealed that three individuals had active hepatitis B virus infection, one had

active hepatitis C virus infection and one individual was HIV positive, with low viral load

(Supplementary Table S2). Those individuals were kept in all downstream analyses, as the

number of positive cases was too low to analyze separately. There was no effect of liver fat

content on either the total number or the MAFs of heteroplasmic sites (Supplementary Figure

S2, p>0.05) and the mitochondrial DNA copy numbers, estimated for each liver sample as

described before (Wachsmuth et al. 2016), were highly correlated between corresponding liver

samples of an individual, suggesting no functional differences between the liver lobes

(Supplementary Figure S3, r=0.81, p<0.001).

Figure 1: Correlation of MAFs at heteroplasmic sites in liver lobes. Each dot is one

heteroplasmic site in one individual. Pearson's correlation coefficient r is given.

Heteroplasmic sites are compared in **A** liver sample 1 and 2, **B** the control region of liver

sample 1 and 2, **C** the non-control region of liver sample 1 and 2. Heteroplasmic sites

shared between liver sample 1 and 2 are in dark grey, non-shared sites in light grey.

We next asked whether heteroplasmy was correlated between the three different samples

from an individual. For blood and each liver sample, there is a low but significant correlation

(r=0.35 and 0.38, p<0.001, Supplementary Figure S4) with only 7.5% of the heteroplasmies

shared between the tissues. However, the correlation between the two liver samples was

higher (Figure 1A, r=0.90, p<0.001). While 355 sites were heteroplasmic in only one of the

two liver samples of an individual, 136 sites (28%) were heteroplasmic in both liver samples

and these exhibited similar MAFs (Figure 1A). Moreover, there were more shared

heteroplasmies from the control region than from the rest of the genome (Table 1, p<0.001):

55% of control region heteroplasmies were shared, vs. 7.5% of heteroplasmies outside the

control region. The high correlation between the MAFs in the two liver samples was mainly

driven by the control region (Figure 1B, r=0.90, p<0.001), while there was a lower, but still

significant correlation outside the control region (Figure 1C, r=0.29, p<0.001). The difference

in correlation coefficients is significant, based on random partitions of all of the heteroplasmic

sites into two sets with the same number of sites as observed for the control region and the

non-control region (p<0.001, Supplementary Figure S5). Thus, not only are more

heteroplasmies shared in the control region than outside the control region, but control region

heteroplasmies also have more similar MAFs.

Table 1: Heteroplasmic mutations in the control region and the non-control region and shared

135   and non-shared heteroplasmies. p<0.001 (two-sided Fisher's exact test).

heteroplasmies in liver1 and liver2	control region	non-control region	total
shared	115	21	136
not shared	95	260	355
total	210	281	

137   **Potential mechanisms underlying the more frequent sharing in the control region**

Figure 2: Heteroplasmy sharing in liver samples and correlation with single sites. MAFs

per site in liver sample 1 and 2 for: **A** sites 60, 72 and 94; and **B** for all other

heteroplasmic sites in the control region. Each dot is one heteroplasmic site in one

individual. Pearson's correlation coefficient r is given. **C** distribution of r-values for

correlation of liver sample 1 and 2 MAFs per site for random subsamples of

heteroplasmies in the control region (same number as the sum of sites 60, 72 and 94).

The r-value for sites 60, 72 and 94 only is shown as a red bar.

Frequent heteroplasmic sites. There are three sites that are frequently heteroplasmic in liver

(Supplementary Table S1) and all are in the control region (sites 60, 72, and 94). To determine

if these three sites are driving the higher correlation in MAF in the control region, we analyzed

them separately. While these three sites did indeed show a high correlation in MAF between

the two liver samples (Figure 2A, r = 0.91, p<0.001), the remaining sites in the control region

still showed a significant correlation (Figure 2B, r=0.82, p<0.001) that is higher than the

correlation for the non-control region (Figure 1C). To determine if the difference in correlations

for the three sites vs. the remaining sites in the control region was statistically significant, we

performed a subsampling test. As there were 113 occurrences of heteroplasmy at these three

sites, we sampled 113 heteroplasmies at random from the control region, calculated the

correlation in MAF between the two liver samples, and repeated this 1000 times. The r-value

for the three sites was not significantly higher than the r-values for the random subsamples

(Figure 2C), indicating that the higher correlation between MAF for the control region than for

the rest of the genome is not driven solely by these three sites.

7S DNA. The control region includes the D-loop region, in which a third strand, the 7S DNA,

displaces the heavy strand and binds to the light strand. Inferred heteroplasmies in the D-loop

region might therefore reflect mutations in the 7S DNA rather than mutations in the mtDNA

itself. To see if sequences from 7S DNA were likely to be present in the sequencing libraries,

we estimated the relative mtDNA copy number from the capture-enrichment sequencing

coverage of the D-loop region and the rest of the mtDNA molecule separately, as described

before (Wachsmuth et al. 2016). As the 7S DNA has several starting and end points, we used

the outer limits reported in the literature, namely from site 16,097 to site 191 (Roberti et al.

1998; Nicholls and Minczuk 2014). The D-loop did not exhibit a higher copy number than the

other parts of the mtDNA genome, indicating that 7S DNA is unlikely to be present in the DNA

libraries (p=0.30 and 0.41 for blood and liver, respectively, Supplementary Figure S6A).

Furthermore, the correlation between the MAF for the two liver samples is almost as high for

the D-loop region as it for the rest of the control region (Supplementary Figure S6 B,C: r=0.90

vs. r = 0.89). Hence, the significant correlation in MAFs in the control region is likely a

phenomenon of the entire control region.

Higher mutation rate. We also tested if heteroplasmic sites showed a higher mutation rate than

non- heteroplasmic sites by comparing the number of heteroplasmies to the inferred mutation

rate at each site, based on observed polymorphism data (Soares et al. 2009). Heteroplasmic

sites had significantly higher mutation rates than sites that were not heteroplasmic (p<0.001,

Supplementary Figure S7) with the mutation rate being higher in the control region than

outside the control region. However, mutation rates did not differ between shared vs. non-

shared heteroplasmies (p>0.05, Supplementary Figure S7), suggesting that the mutation rate

does not increase the probability of heteroplasmies to be shared.

Figure 3: Expected difference in minor allele frequency (MAF) assuming random genetic drift. A schematic scheme of the two-step mtDNA replication model used for the simulations: first, all mtDNA molecules are segregated equally into two daughter cells followed by replication step to re-gain the same number of mtDNA copies as in the

progenitor cell. After each generation n_i, the MAF frequency of one random cell on each

site of the pedigree is determined and the MAF difference between the two cells

calculated. **B** expected difference in MAF after 10, 20, 50, 100, and 250 generations

since the split of the two cells. Different mtDNA copy numbers (1,000, 2,500, and 5,000;

kept constant throughout simulation) and initial MAFs (5%, 10%, 20%, and 50%) were

used to simulate random genetic drift between two cells. For each combination of mtDNA

copy number and initial MAF, 1000 replicates were simulated. The black, dotted line

indicates the observed MAF difference between the shared liver heteroplasmies in the

data set. The asterisk highlights the simulation with the parameters closest to the ones

observed in the data set. **C** the mtDNA copy number distribution estimated from the

capture-enriched sequencing data of the liver samples. The values were corrected using

a correction "ratio" of 1/150 (Wachsmuth et al. 2016) to convert the relative to absolute

mtDNA copy numbers. **D** the distribution of MAF in the liver heteroplasmies.

Random genetic drift. If control region heteroplasmies are more likely to have arisen in a

common progenitor cell of the cells sampled in the two liver lobes, then the higher correlation

in MAF might reflect this common ancestry. We therefore tested if the correlation in MAF is

compatible with expected amount of random genetic drift from a common progenitor cell by

simulating mtDNA as a two-step model (Figure 3A) following (Jayaprakash et al. 2015). We

started with different MAFs at the initial generation n_0 (5%, 10%, 20%, 50%) and different

constant mitochondrial DNA copy numbers (1,000, 2,500, and 5,000) and sampled the

difference in MAF 10, 20, 50, 100, and 250 generations after the initial split of the cells (Figure

3B). For all simulations, we observed a higher MAF difference with an increasing number of

cell replications. This effect was more pronounced for higher MAF at generation n_0 and lower

mtDNA copy number. Based on the observed mtDNA copy number (Figure 3C; mean=2,528)

and MAF of shared heteroplasmies with the same consensus allele (Figure 3D; mean=10.8%),

the most similar corresponding simulation is with a copy number of 2,500 and a MAF at

generation n_0 of 10%; for this simulation, the variance in observed MAF differences was

smaller than would be expected after 50 generations of random genetic drift (one-sided F-test

for the equality of two variances, p<0.001). With an increasing number of cell replications, the

simulated MAF difference further diverged from the observed values, which suggests that

random genetic drift cannot explain the observed sharing.

221 Differences in MAF between corresponding liver samples are not correlated with age

We next investigated the influence of age on heteroplasmy sharing between different liver

regions of an individual. Overall, the total number of different heteroplasmic sites in an

individual increased with age for both the control region and the coding region (Supplementary

Figure S8A, r=0.42 and adjusted p<0.001 both within and outside of the control region).

However, the MAF at heteroplasmic sites did not increase with age (Supplementary Figure

S8B, r=0.09 for the control region, -0.01 for outside of the control region, adjusted p>0.05).

Although some heteroplasmies exhibit high MAFs only at ages above 50, many sites remain

at low frequencies even at older ages (Supplementary Figure S8B). The hypothesis that

random genetic drift has an effect on the difference in MAF of two corresponding liver samples

would suggest that with increasing age the difference in MAF would increase, too. Yet, when

testing for a correlation between the difference in MAF between two corresponding liver

samples and age of the individual, we did not observe a significant correlation either within or

outside the control region (Figure 4, r=0.06 within control region, r=-0.01 outside the control

region, both p>0.05). In order to test whether, in contrast, specific sites have a significant

correlation between the MAF difference and age, we separately calculated the correlation

between MAF difference and age for the three most frequent heteroplasmy sites (sites 60, 72,

and 94), for which at least 20 individuals were heteroplasmic (Supplementary Figure S9). The

correlation between MAF and age was not significant for any of these three sites (adjusted

p>0.05), which further supports the hypothesis that the difference in MAF does not increase

with age, contrary to what would be expected from random genetic drift.

Figure 4: Minor allele frequency difference and age. The difference in minor allele frequency between liver sample 1 and 2 versus age plotted separately for sites within and outside the control region. Dark grey dots indicate sites that were shared (i.e. heteroplasmic in both liver samples from an individual), light grey dots indicate sites that were not shared.

Synonymous heteroplasmies are more often shared than non-synonymous ones

Previous studies showed that liver has a significant excess of non-synonymous heteroplasmies which are predicted to have an impact on function (Li et al. 2015). We investigated this by calculating the ratio of non-synonymous heteroplasmies per non-synonymous site vs. synonymous heteroplasmies per synonymous site (hN/hS). In the absence of any selection this ratio has an expected value of 1; purifying selection results in values less than 1 and positive selection results in values greater than 1. As found previously (Li et al. 2015), the hN/hS ratio is significantly greater than 1 (adjusted $p < 0.05$, Supplementary Figure S10A), indicating positive selection for non-synonymous heteroplasmies rather than relaxation of functional constraints (Li et al. 2015). Additionally, we observed a strong increase in the number of heteroplasmies occurring the coding region compared to the control region in individuals older than 60 years (Supplementary Figure S8A). We, therefore, investigated whether the excess in non-synonymous heteroplasmies is age-dependent (Supplementary

Figure S10B). While both synonymous and non-synonymous heteroplasmies increased in

number with increasing age, only non-synonymous heteroplasmies showed a significant

difference between the age groups below 60 years and above 60 years (two-sided Mann-

Whitney U test, adjusted p<0.05). We then asked if either synonymous or non-synonymous

heteroplasmic sites were more likely to be shared between different liver samples. Although

there were more than twice as many non-synonymous than synonymous heteroplasmies in

the data set (163 vs. 62), there were significantly more synonymous sites shared than non-

synonymous sites (11 vs. 6, Table 2). Accordingly, the median MAF difference between

corresponding liver samples was on average higher for non-synonymous heteroplasmies than

synonymous ones (median MAF difference of 4.4% to 3.8%), albeit not significantly so (p>0.05,

Supplementary Figure S11A). These results suggest that non-synonymous heteroplasmies

more often arise independently in different cells. However, neither non-synonymous nor

synonymous heteroplasmies showed a significant correlation for the difference in MAF

between corresponding liver samples with age (p>0.5, Supplementary Figure S11B).

Table 2: Non-synonymous and synonymous heteroplasmic mutations in the coding region and

shared and non-shared heteroplasmies. p<0.001 (two-sided Fisher's exact test)

heteroplasmies in liver1 and liver2	non-synonymous	synonymous	total
shared	6	11	17
not shared	157	51	208
total	163	62	

DISCUSSION

Heteroplasmies are shared across liver lobes

In this study, we compared mtDNA heteroplasmy patterns in samples collected from the two

primary liver lobes of 83 individuals sampled at autopsy and found a significant correlation in

heteroplasmy MAF between the two liver samples from an individual. Moreover, if sites are

heteroplasmic in both liver samples of an individual, the MAFs at those sites tend to be similar

independent of age. Previous studies found similar correlations in MAF between cell colonies

derived from single cells and grown in culture (Raap et al. 2012; Jayaprakash et al. 2015), so

our results extend these findings to different parts of the same organ within living individuals.

In addition, we analyzed patterns of heteroplasmy sharing and MAF correlation across

different parts of the mtDNA genome, which provide further information concerning the

potential mechanism(s) behind these observations. In particular, we found significantly more

sharing of heteroplasmic sites and higher MAF correlations for the control region than for the

rest of the genome. We showed that these differences are not due to the presence of 7S DNA,

nor are they driven by just a few sites. Moreover, within the coding region, we found

significantly more sharing of heteroplasmies and higher MAF correlations at synonymous sites

than at nonsynonymous sites.

These observations allow us to evaluate several potential explanations for this heteroplasmy

sharing within liver. First, a trivial potential explanation is that a heteroplasmy could appear to

be shared between liver lobes when it is actually present in blood. About 1 liter of blood per

minute flows through the blood vessels of the liver (Wynne et al. 1989), so the DNA from the

liver samples also contains DNA arising from blood cells. Heteroplasmies present in the blood

cells could therefore be detected as heteroplasmies in the liver samples and hence seem to

be shared across lobes. There are $\sim 10^{10}$ liver cells in 10 g of liver tissue and 5×10^{10} blood

cells in 10 ml of blood (Sender et al. 2016), and in liver autopsy samples there is approximately

10 ml blood per 100 g of tissue (Greenway and Stark 1971). The ratio of liver to blood cells in

a liver autopsy sample is therefore about 1.8. As we additionally observed for our samples

that the average mitochondrial copy number is five times higher in liver than in blood samples

(Supplementary Figure S6a), the overall ratio of mitochondrial genomes from liver cells to

those from blood cells is approximately 9 to 1. Therefore, a heteroplasmy in blood would need

a MAF of at least 22.5% to be called a heteroplasmy in liver (minimum MAF = 2.5%) when it

was actually totally absent in the liver samples. Only 8 out of the 541 heteroplasmies in blood

have a MAF of ≥ 22.5%, of which four are shared across blood and both liver lobes

(Supplementary Table S1). Moreover, the heteroplasmies that are commonly observed in liver (at positions 60, 72, and 94) are practically absent from blood; these results rule out experimental contamination with blood DNA as the primary driver for heteroplasmy sharing across liver lobes.

A second reason for heteroplasmy sharing could be a high mutational pressure for some sites, with de novo mutations at the same site occurring independently throughout the tissue. If this was the case, one would expect to see a higher correlation of MAFs for common heteroplasmic sites, as those would be under high mutational pressure. We showed that this is not the case for the most common heteroplasmic sites 60, 72, and 94 in the data set (Figure 2), and hence this explanation is unlikely. While we further observed a higher mutation rate for heteroplasmic sites than for non-heteroplasmic sites (Supplementary Figure S7), there was no difference in mutation rate between shared and non-shared heteroplasmies, suggesting that the mutation rate has little impact on heteroplasmy sharing.

Third, it has been shown that colonic stem cells with non-synonymous mtDNA mutations can expand clonally from a few cells and spread throughout a tissue by crypt fission (Greaves et al. 2006). During this process they retain the same level of mutant DNA on a cellular level. While a similar mechanism in liver could explain the results presented here, this process would require a cellular turnover on a huge scale throughout the entire liver, starting from just a few stem cells. This does not seem very likely, as clonal expansion was shown for single intestinal crypts only and is supposed to be rather slow (Greaves et al. 2006). Moreover, it is not clear to what extent liver regeneration is driven by stem cells vs. mature hepatocytes (Grompe 2014).

Fourth, heteroplasmy sharing could also derive from a pre-existing, inherited heteroplasmy (Guo et al. 2013; Payne et al. 2013; Rebolledo-Jaramillo et al. 2014; Li et al. 2015) that remains at similar frequencies across the tissue because drift (random changes in MAF) is limited. We tested this possibility, using a simulation scheme from a previous study (Jayaprakash et al. 2015), with different initial heteroplasmy frequencies, and mtDNA copy numbers that were kept constant (Berk and Clayton 1974). As expected, we observed a larger

variance in MAF between daughter cells with increasing starting MAF and decreasing mtDNA copy number (Figure 3B). For the simulation results closest to our observed average mtDNA copy number and average heteroplasmy MAF (2,500 and 10%, Figure 3C-D), the observed difference in MAF between shared heteroplasmies was smaller than would be expected after random genetic drift acting on 50 mtDNA replication steps. However, more than these 50 mtDNA replication steps are expected to have taken place during the life of an average individual in our study. Assuming 3.61×10^{11} liver cells in an adult liver (Bianconi et al. 2013), a total of 39 hepatocyte cell replications (including mtDNA replication) are needed to obtain a full-size, adult liver from a single hepatocyte cell. After the development stage, the post-mitotic cells continue to replicate their mtDNA independently of cell replication ("relaxed" replication (Poovathingal et al. 2009)). The estimated half-life of mtDNA in post-mitotic cells ranges between 2-10 days (Miwa et al. 2008) and 30-300 days (Poovathingal et al. 2012). Assuming an mtDNA half-life of 30 days, there would be complete replacement of all mtDNA molecules of a cell within a year, given a mtDNA copy number of 2,500 per cell, so approximately one "relaxed" replication cycle occurs within the liver of an adult for every year of age. Thus, for the age range in our data set of 24 to 94 years, the liver samples would have gone through about 62 - 132 cell replications prior to sampling, and so the observed difference in MAF in the liver samples is significantly smaller than expected. Moreover, liver samples accumulated significantly more heteroplasmies with age (Supplementary Figure S8A), further arguing that shared heteroplasmies do not reflect pre-existing, inherited heteroplasmies. Also, there was no significant correlation of MAF difference with age, as would be expected with random genetic drift. In sum, our results extend to liver tissues the previous observations (Raap et al. 2012; Jayaprakash et al. 2015) that random genetic drift alone cannot explain heteroplasmy sharing between cells in culture.

Finally, an equilibrium of heteroplasmy across an entire tissue could be explained by an exchange of genetic material from mitochondria between cells. Cells can donate whole mitochondria to adjacent cells through nanotubes, but this has been suggested for distances

up to 100 μM only and the exchange is often triggered by functional impairments in the

acceptor cell (Rogers and Bhattacharya 2013). An additional way for cells to exchange DNA

material could be the uptake of extracellular DNA material that is either secreted by healthy

cells or is present as the remains of apoptotic cells (van der Vaart and Pretorius 2008). While

the uptake and integration of cell-free nuclear DNA material has been shown (Basak et al.

2016), it is unclear whether cells would also accept mitochondrial DNA. However, studies of

heteroplasmy at the single cell level (reviewed in (Yao et al. 2015)) do suggest the possibility

of transfer between cells. Experiments with cell culture mixes of fluorescently labelled cell lines

suggested the exchange of mtDNA between co-cultured partner cell lines, although the

specific mechanism, either transfer of mitochondrial organelles or transfer of free mtDNA,

could not be identified (Jayaprakash et al. 2015). Overall, such intercellular DNA exchange,

followed by incorporation of mtDNA fragments into the mtDNA of the recipient cells, could

account for the significant correlation we observe in MAF between liver lobes.

However, other aspects of our data are incompatible with the hypothesis of intercellular DNA

exchange. In particular, intercellular exchange cannot explain the significantly higher number

of shared heteroplasmies and correlation in MAF for the control region vs. the rest of the

genome, unless one postulates that mtDNA fragments arising from the control region are

either exchanged or incorporated between cells more frequently than mtDNA fragments

arising from the rest of the genome. But even then, intercellular exchange cannot explain the

significantly higher number of shared heteroplasmies and correlation in MAF for synonymous

vs. non-synonymous heteroplasmies, as both should be exchanged at the same rate between

cells.

Instead, our data suggest that even if intercellular exchange is occurring, selection must be

involved in the sharing of heteroplasmies and correlation in MAF between liver lobes. Several

aspects of the data suggest that selection influences heteroplasmies. In the coding region of

the mtDNA, we observed a significant excess of non-synonymous vs. synonymous

heteroplasmies (Table 2), more so than can be explained by relaxation of functional

constraints on non-synonymous mutations. Moreover, the number of non-synonymous

heteroplasmies increased significantly in individuals above 60 years. Overall, these results

strongly suggest positive selection for nonsynonymous heteroplasmies in liver, as found

previously (Li et al. 2015), and possibly reflecting the hypothesis of the "survival of the slowest"

(deGrey 1997), which postulates that mitochondria with reduced respiratory function due to

increasing mutations suffer less degradation from the production of reactive oxygen species

(ROS). Hence, mitochondria that lack these mutations suffer ROS-related damage and are

removed from cells, thereby resulting in an increase in frequency in mtDNAs with non-

synonymous mutations that decrease mitochondrial function.

However, our results indicate a more complex role for selection in the different patterns of

heteroplasmy sharing and MAF correlation across different regions of the mtDNA genome, in

keeping with evidence from other studies. In mice and humans, significantly more synonymous

than non-synonymous heteroplasmies are transmitted to the next generation (Stewart et al.

2008; Rebolledo-Jaramillo et al. 2014; Floros et al. 2018), suggesting selection against non-

synonymous heteroplasmies during transmission. The notable tissue-specificity and allele-

specificity of particular heteroplasmic sites in the control region also suggests a role for

positive selection on heteroplasmies during aging (Samuels et al. 2013; Li et al. 2015). The

increasing evidence for both purifying and positive selection acting on heteroplasmic variants

warrants further investigation, particularly into the potential health-related consequences.

MATERIAL AND METHODS

Tissue collection and DNA extraction

Blood and liver were sampled at autopsy from 94 individuals (57 males, 37 females, age range:

24-94, mean: 63, median: 63). Two samples were taken from each liver, one from the right

lobe and one from the left lobe. DNA was extracted as previously described (Li et al. 2015).

The collection of samples and the experimental procedures were approved by the Ethics

Commissions of the Rheinische Friedrich Wilhelms University Medical Faculty (Lfd. Nr. 097/15)

and the University of Leipzig Medical Faculty (Az. 305-15-24082015).

Virological assays and histological investigation

Human immunodeficiency virus (HIV) RNA and Hepatitis C virus (HCV) RNA concentration in

blood was determined by using the Abbott RealTi*me*® HIV-1 and HCV systems and the

m2000sp/m2000rt instruments according to the instructions of the manufacturer. For detection

of Hepatitis B virus (HBV) DNA the Abbott RealTi*me*® HBV system was used. The 95% limit

of detection (LOD_{95}) of the HIV, HCV and HBV assay was 40 copies/mL, 12 IU/ mL, and 10

IU/ mL, respectively. If inhibitory effects on enzymatic reactions were present (detected via

co-amplification of control RNA or DNA sequences), blood samples were re-tested at dilutions

of 1/5, 1/10, and 1/15 (13 samples (13%) for HIV, 91 (93%) for HCV, and 8 (7%) for HBV load).

For dilution, a plasma donation from a blood donor negative for HIV, HCV, and HBV was used.

Of the diluted samples tested for HIV, one sample was diluted 1/5, nine had to be diluted 1/10,

and two had to be diluted 1/15, lowering the LOD_{95} of the assay to 200, 400, and 600,

respectively. In one sample, inhibition could not be eliminated by sample dilution. Of the diluted

samples tested for HCV, one, 78, and two samples were diluted 1/5, 1/10, and 1/15,

respectively, lowering the LOD_{95} of the assay to 60, 120, 180 IU/ml, respectively. In 10

additional samples no result could be achieved, even after dilution. Of the diluted samples

tested for HBV load, 3, 2, and 2 samples were diluted 1/5, 1/10, and 1/15, respectively,

lowering the LOD_{95} of the assay to 50, 100, and 150 IU/mL, respectively. All other samples

were tested without any dilution.

Fat content of the liver was determined by histological investigations and Sudan staining

(Mulisch M 2015). Tissues with <10% hepatocytes including fat droplets were considered low,

10-30% were medium, 31-50% were high fat and >50% were considered adipohepatic.

Illumina library preparation and sequencing

Double-barcoded DNA libraries for sequencing were prepared and capture-enriched for mtDNA as previously described (Li et al. 2015). DNA was sequenced on the Illumina HiSeq platform in rapid mode with 95 bp paired-end reads. Bases were called with FreeIbis (Renaud et al. 2013) and reads were subsequently trimmed and merged using leeHom (Renaud et al. 2014).

Heteroplasmy detection

Heteroplasmy was detected according to the DREEP pipeline (Li et al. 2010; Li and Stoneking 2012). First, heteroplasmies were called if: the minor allele frequency (MAF) for the most frequent minor allele was at least 2.5% on both the forward and reverse strand; the sequencing depth was at least 500-fold at a candidate site; and there were at least 10 reads supporting the minor allele on each strand. Additionally, we required a minimum heteroplasmic quality score of 10 on each strand. In order to discriminate a true heteroplasmy from sequencing error, the DREEP pipeline compares the minor allele pattern of any inferred heteroplasmic site to a database that comprises the minor allele patterns at this site from all other individuals in the study. When the majority of the samples are from a single tissue like liver in this study (two liver samples and one blood sample per individual), DREEP is prone to considered commonly heteroplasmic sites as elevated sequencing error and therefore under-estimates the heteroplasmic quality score for these. Thus, we used the information about commonly heteroplasmic sites from (Li et al. 2015) to flag these for both blood and liver tissue in this study, respectively. A site was considered commonly heteroplasmic in a tissue when at least five individuals were heteroplasmic for it. Based on this criterion, we ignored the heteroplasmic quality for sites 60, 72, 94, 185, 189, 203, 11,126, 16,093, and 16,126 for liver and 12,705 for blood and considered a site heteroplasmic if all other criteria were fulfilled. The following regions were excluded for heteroplasmy analysis: 302-316, 513-526, 566-573, and 16,181-16,194. We confirmed that all heteroplasmies were within a coverage between 20 % and 200 % of the average coverage of the sample. In addition, all samples that could have been contaminated with other samples during library preparation/extraction were removed. To

detect such contamination, pairwise comparisons of all liver samples with each other as well

as all blood samples with each other were performed. All three samples of an individual were

removed if all of the following criteria were fulfilled for any pairwise comparison between

individuals across all sites of the mitochondrial genome: 1) for at least 80% of the sites, for

which two samples had different consensus alleles, the minor allele in the recipient sample

was identical to the major allele in the donor sample; 2) the average MAF across these sites

was at least 1%; and 3) at least 60% of all sites in the recipient sample, for which a minor

allele was observed, were identified as heteroplasmies by the DREEP pipeline (Li et al. 2010;

Li and Stoneking 2012). Furthermore, an additional filter for potential contamination was

applied, in which the heteroplasmic sites for each sample were checked to see if five or more

sites could be explained by contamination from another haplogroup. In total, ten individuals

were removed in these contamination filter steps. Finally, we removed a single individual

because its samples had more than 2% of the MT genome below a sequencing depth of 500-

fold, the cut-off for being considered a heteroplasmy, and thus would have had a higher false

negative rate than the other samples. Overall, we retained 83 individuals for the subsequent

analyses. Minor allele frequencies were calculated with respect to the major allele in blood.

Correlation analysis

Statistical analysis was performed using R (https://www.R-project.org), with analyses

performed for the entire mtDNA genome and separately for the coding region (577-16,023),

the control region (16,024-576) and the D-loop region (16,097-191). For correlation of MAFs,

we selected only sites that were identified as heteroplasmies and passed our quality filters in

at least one of the tissues. We then compared the MAFs of these heteroplasmies to the MAF

in the other tissues of an individual, even if the site was not detected as a heteroplasmy in the

other tissues. Pearson correlation coefficients were calculated for correlations between MAFs

among samples as well as for correlations with age assuming a two-sided alternative

hypothesis; the significance of the correlation was tested by randomly permuting the data.

Permutation tests were also used to assess the association between specific sites or regions

and minor allele sharing between liver lobes; all permutations were carried out 1000 times.

Whenever categorical data were compared (e.g. synonymous vs. non-synonymous sites),

two-sided Mann-Whitney U tests were used to test for significant differences. Fisher's exact

test was used to test for an association of sharing of heteroplasmic sites between liver lobes

for the control region vs. non-control region and for synonymous vs. non-synonymous

heteroplasmies using a two-sided alternative hypothesis. P-values were adjusted for multiple

testing using Benjamini-Hochberg correction (Benjamini and Hochberg 1995) and highlighted

in the text.

Coverage across the mitochondrial genome

505 We used per-site coverage determined by the *filter_and_summary.pl* script of the DREEP

506 pipeline (Li and Stoneking 2012) for each sample and calculated the average coverage across

507 the D-loop region and across the rest of the mtDNA genome.

Non-synonymous heteroplasmies

The hN/hS ratio (Li et al. 2015) was calculated by calculating the Ka/Ks ratio using

KaKs_Calculator 2.0 (Wang et al. 2010) between the revised Cambridge Reference Sequence

(rCRS; Andrews, 1999 doi:10.1038/13779) and a mock sequence created by introducing all

minor alleles of heteroplasmies into the coding region of the rCRS. A significance test was

performed by randomly introducing the same substitutions as observed for heteroplasmies at

any site of the coding region of rCRS and calculating the hN/hS ratio in comparison to the non-

altered rCRS.

The potential functional impact of non-synonymous heteroplasmies was analyzed by

overlapping the position of the heteroplasmy and its minor allele with the MitImpact database

(Castellana et al. 2015) and comparing the *MutationAssessor* (Reva et al. 2011) score.

DATA ACCESS

The raw sequencing data have been deposited with the European Nucleotide Archive under

accession number PRJEB27731. The scripts for heteroplasmy detection and filtering can be

found at http://dmcrop.sourceforge.net/ and https://github.com/alexhbnr/liverlobes

_heteroplasmy.

ACKNOWLEDGEMENTS

We thank Anne Haubner, Antje Weihmann, Barbara Höber and Melanie Grabmüller for

technical support. We thank Enrico Macholdt, Michael Dannemann, and Martin Petr for

valuable discussion. The research was funded by the Max Planck Society

AUTHOR CONTRIBUTIONS

MS, BM, ML, AH and MW designed the study. BM collected the samples. RS performed DNA

isolation and sequencing library preparation, AMEH performed virological assays and BM

performed histological analyses. ML and AH wrote pipeline and executed heteroplasmy

detection, AH and MW performed statistical analyses. AH, MW, AMEH and MS wrote the

manuscript. All authors read and approved the final manuscript.

DISCLOSURE DECLARATION

The authors declare that they have no competing interests.

REFERENCES

Basak R, Nair NK, Mittra I. 2016. Evidence for cell-free nucleic acids as continuously arising

endogenous DNA mutagens. *Mutation Research-Fundamental and Molecular*

Mechanisms of Mutagenesis **793**: 15-21.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate - a practical and powerful

approach to multiple testing. *Journal of the Royal Statistical Society Series B-*

Methodological **57**: 289-300.

Berk AJ, Clayton DA. 1974. Mechanism of mitochondrial DNA replication in mouse L-cells:

asynchronous replication of strands, segregation of circular daughter molecules,

aspects of topology and turnover of an initiation sequence. *Journal of molecular biology*
86: 801-824.

Bianconi E, Piovesan A, Facchin F, Beraudi A, Casadei R, Frabetti F, Vitale L, Pelleri MC,
Tassani S, Piva F et al. 2013. An estimation of the number of cells in the human body.
Annals of human biology **40**: 463-471.

Calloway CD, Reynolds RL, Herrin GL, Anderson WW. 2000. The frequency of heteroplasmy
in the HVII region of mtDNA differs across tissue types and increases with age.
American Journal of Human Genetics **66**: 1384-1397.

Castellana S, Ronai J, Mazza T. 2015. MitImpact: an exhaustive collection of pre-computed
pathogenicity predictions of human mitochondrial non-synonymous variants. *Hum*
Mutat **36**: E2413-2422.

deGrey A. 1997. A proposed refinement of the mitochondrial free radical theory of aging.
Bioessays **19**: 161-166.

Floros VI, Pyle A, Dietmann S, Wei W, Tang WWC, Irie N, Payne B, Capalbo A, Noli L,
Coxhead J et al. 2018. Segregation of mitochondrial DNA heteroplasmy through a
developmental genetic bottleneck in human embryos. *Nature cell biology* **20**: 144-144.

Greaves LC, Preston SL, Tadrous PJ, Taylor RW, Barron MJ, Oukrif D, Leedham SJ,
Deheragoda M, Sasieni P, Novelli MR et al. 2006. Mitochondrial DNA mutations are
established in human colonic stem cells, and mutated clones expand by crypt fission.
Proceedings of the National Academy of Sciences of the United States of America **103**:
714-719.

Greenway C, Stark R. 1971. Hepatic vascular bed. *Physiological reviews* **51**: 23-65.

Grompe M. 2014. Liver stem cells, where art thou? *Cell stem cell* **15**: 257-258.

Guo Y, Li CI, Sheng QH, Winther JF, Cai QY, Boice JD, Shyr Y. 2013. Very Low-Level
Heteroplasmy mtDNA Variations Are Inherited in Humans. *Journal of Genetics and*
Genomics **40**: 607-615.

Jayaprakash AD, Benson EK, Gone S, Liang R, Shim J, Lambertini L, Toloue MM, Wigler M,
Aaronson SA, Sachidanandam R. 2015. Stable heteroplasmy at the single-cell level is
facilitated by intercellular exchange of mtDNA. *Nucleic Acids Research* **43**: 2177-2187.

Jenuth JP, Peterson AC, Fu K, Shoubridge EA. 1996. Random genetic drift in the female
germline explains the rapid segregation of mammalian mitochondrial DNA. *Nature*
genetics **14**: 146-146.

Jenuth JP, Peterson AC, Shoubridge EA. 1997. Tissue-specific selection for different mtDNA
genotypes in heteroplasmic mice. *Nature genetics* **16**: 93-93.

Kowald A, Kirkwood TBL. 2011. Evolution of the mitochondrial fusion--fission cycle and its role
in aging. *Proceedings of the National Academy of Sciences* **108**: 10237-10242.

Larsson NG. 2010. Somatic mitochondrial DNA mutations in mammalian aging. *Annual review*
of biochemistry **79**: 683-706.

Li M, Schoenberg A, Schaeferd M, Schroeder R, Nasidze I, Stoneking M. 2010. Detecting
Heteroplasmy from High-Throughput Sequencing of Complete Human Mitochondrial
DNA Genomes. *American Journal of Human Genetics* **87**: 237-249.

Li M, Schroeder R, Ni S, Madea B, Stoneking M. 2015. Extensive tissue-related and allele-
related mtDNA heteroplasmy suggests positive selection for somatic mutations.
Proceedings of the National Academy of Sciences of the United States of America **112**:
2491-2496.

Li M, Stoneking M. 2012. A new approach for detecting low-level mutations in next-generation
sequence data. *Genome Biology* **13**: R34.

Maricic T, Whitten M, Paeaebo S. 2010. Multiplexed DNA Sequence Capture of Mitochondrial
Genomes Using PCR Products. *Plos One* **5**: e14004.

Michikawa Y, Mazzucchelli F, Bresolin N, Scarlato G, Attardi G. 1999. Aging-dependent large
accumulation of point mutations in the human mtDNA control region for replication.
Science **286**: 774-779.

Miwa S, Lawless C, Von Zglinicki T. 2008. Mitochondrial turnover in liver is fast in vivo and is
accelerated by dietary restriction: application of a simple dynamic model. *Aging cell* **7**:
920-923.

Mulisch M WU. 2015. *Romeis - Mikroskopische Technik*. Springer Spektrum, Berlin
Heidelberg.

Nachman MW, Brown WM, Stoneking M, Aquadro CF. 1996. Nonneutral mitochondrial DNA
variation in humans and chimpanzees. *Genetics* **142**: 953-963.

Naue J, Horer S, Sanger T, Strobl C, Hatzer-Grubwieser P, Parson W, Lutz-Bonengel S. 2015.
Evidence for frequent and tissue-specific sequence heteroplasmy in human
mitochondrial DNA. *Mitochondrion* **20**: 82-94.

Nicholls TJ, Minczuk M. 2014. In D-loop: 40 years of mitochondrial 7S DNA. *Experimental*
Gerontology **56**: 175-181.

Payne BAI, Wilson IJ, Yu-Wai-Man P, Coxhead J, Deehan D, Horvath R, Taylor RW, Samuels
DC, Santibanez-Koref M, Chinnery PF. 2013. Universal heteroplasmy of human
mitochondrial DNA. *Human Molecular Genetics* **22**: 384-390.

Poovathingal SK, Gruber J, Halliwell B, Gunawan R. 2009. Stochastic drift in mitochondrial
DNA point mutations: a novel perspective ex silico. *PLoS computational biology* **5**:
e1000572.

Poovathingal SK, Gruber J, Lakshmanan L, Halliwell B, Gunawan R. 2012. Is mitochondrial
DNA turnover slower than commonly assumed? *Biogerontology* **13**: 557-564.

Raap AK, Tafrechi RSJ, van de Rijke FM, Pyle A, Wählby C, Szuhai K, Ravelli RBG, de Coo
RFM, Rajasimha HK, Nilsson M et al. 2012. Non-random mtDNA segregation patterns
indicate a metastable heteroplasmic segregation unit in m. 3243A> G cybrid cells.
PLoS One **7**: e52080-e52080.

Rebolledo-Jaramillo B, Su MS-W, Stoler N, McElhoe JA, Dickins B, Blankenberg D,
Korneliussen TS, Chiaromonte F, Nielsen R, Holland MM et al. 2014. Maternal age
effect and severe germ-line bottleneck in the inheritance of human mitochondrial DNA.
Proceedings of the National Academy of Sciences of the United States of America **111**:
15474-15479.

Renaud G, Kircher M, Stenzel U, Kelso J. 2013. freeIbis: an efficient basecaller with calibrated
quality scores for Illumina sequencers. *Bioinformatics* **29**: 1208-1209.

Renaud G, Stenzel U, Kelso J. 2014. leeHom: adaptor trimming and merging for Illumina
sequencing reads. *Nucleic Acids Research* **42**: e141.

Reva B, Antipin Y, Sander C. 2011. Predicting the functional impact of protein mutations:
application to cancer genomics. *Nucleic Acids Res* **39**: e118.

Roberti M, Musicco C, Polosa PL, Milella F, Gadaleta MN, Cantatore P. 1998. Multiple protein-
binding sites in the TAS-region of human and rat mitochondrial DNA. *Biochemical and
Biophysical Research Communications* **243**: 36-40.

Rogers RS, Bhattacharya J. 2013. When Cells Become Organelle Donors. *Physiology* **28**:
414-422.

Samuels DC, Li C, Li B, Song Z, Torstenson E, Clay HB, Rokas A, Thornton-Wells TA, Moore
JH, Hughes TM et al. 2013. Recurrent Tissue-Specific mtDNA Mutations Are Common
in Humans. *Plos Genetics* **9**: e1003929.

Sender R, Fuchs S, Milo R. 2016. Revised estimates for the number of human and bacteria
cells in the body. *PLoS biology* **14**: e1002533.

Soares P, Ermini L, Thomson N, Mormina M, Rito T, Rohl A, Salas A, Oppenheimer S,
Macaulay V, Richards MB. 2009. Correcting for Purifying Selection: An Improved
Human Mitochondrial Molecular Clock. *American Journal of Human Genetics* **84**: 740-
759.

Stewart JB, Chinnery PF. 2015. The dynamics of mitochondrial DNA heteroplasmy:
implications for human health and disease. *Nat Rev Genet* **16**: 530-542.

Stewart JB, Freyer C, Elson JL, Wredenberg A, Cansu Z, Trifunovic A, Larsson N-G. 2008.
Strong purifying selection in transmission of mammalian mitochondrial DNA. *PLoS
biology* **6**: e10-e10.

van der Vaart M, Pretorius PJ. 2008. Circulating DNA - Its origin and fluctuation. In *Circulating
Nucleic Acids in Plasma and Serum V*, Vol 1137 (ed. PB Gahan, R Swaminathan), pp.
18-26.

Wachsmuth M, Huebner A, Li M, Madea B, Stoneking M. 2016. Age-Related and
Heteroplasmy-Related Variation in Human mtDNA Copy Number. *Plos Genetics* **12**:
e1005939.

Wallace DC, Chalkia D. 2013. Mitochondrial DNA Genetics and the Heteroplasmy Conundrum
in Evolution and Disease. *Cold Spring Harbor Perspectives in Biology* **5**: a021220.

Wang D, Zhang Y, Zhang Z, Zhu J, Yu J. 2010. KaKs_Calculator 2.0: a toolkit incorporating
gamma-series methods and sliding window strategies. *Genomics, proteomics &*
bioinformatics **8**: 77-80.

Wang Y, Michikawa Y, Mallidis C, Bai Y, Woodhouse L, Yarasheski KE, Miller CA, Askanas
V, Engel WK, Bhasin S et al. 2001. Muscle-specific mutations accumulate with aging
in critical human mtDNA control sites for replication. *Proceedings of the National*
Academy of Sciences of the United States of America **98**: 4022-4027.

Wynne HA, Cope LH, Mutch E, Rawlins MD, Woodhouse KW, James OF. 1989. The effect of
age upon liver volume and apparent liver blood flow in healthy man. *Hepatology* **9**:
297-301.

Yao YG, Kajigaya S, Young NS. 2015. Mitochondrial DNA mutations in single human blood
cells. *Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis*
779: 68-77.