1

Intestine-to-neuronal signaling alters risk-taking behaviors in food-deprived

Caenorhabditis elegans

Molly A. Matty[1†], Hiu E. Lau[1,2†], Anupama Singh[1,3], Jessica A. Haley[1,4], Ahana Chakraborty[1],

Karina Kono[1], Kirthi C. Reddy[1], Malene Hansen[3], and Sreekanth H. Chalasani[1]*

[1] Molecular Neurobiology Laboratory, The Salk Institute for Biological Studies, La Jolla, CA

92037, USA.

[2] Division of Biological Sciences, University of California, San Diego, La Jolla, CA 92093,

USA.

[3] Development, Aging and Regeneration Program, Sanford Burnham Prebys Medical Discovery

Institute, La Jolla, CA 92037, USA.

[4] Neurosciences Graduate Program, University of California, San Diego, La Jolla, CA 92093,

USA.

*Author for correspondence: schalasani@salk.edu (S.H.C.).

[†] *These authors contributed equally to this project*

Long Title: Intestine-to-neuronal signaling alters risk-taking behaviors in food-deprived

Caenorhabditis elegans

Short Title: Food deprivation alters behavior

Author Summary

We have all experienced behavioral changes when we are hungry - the pang in our stomach can

cause us to behave erratically. In particular, hungry animals, including humans, are known to

pursue behaviors that involve higher risk compared to when they are well-fed. Here we explore

the molecular details of this behavior in the invertebrate animal model C. elegans. This behavior,

termed sensory integration, shows that C. elegans display reduced copper sensitivity when

hungry. Copper is toxic and repellant to C. elegans; reduced avoidance indicates that these

animals use riskier food search behaviors when they are hungry. Luckily, like us, this behavioral

change is reversible upon re-feeding. This hunger-induced behavioral change is not due to

increased attraction to food or depletion of fat stores, but rather insulin signaling between the

intestine and specific neurons. We use genetic tools, microscopy, and behavioral tests to

determine that this risky behavior involves sensation of "lack of food" in the intestine, release of

signaling molecules, and engagement with sensory neurons. Our work highlights new and

potentially evolutionarily conserved ways in which intestinal cells and neurons communicate

leading to largescale behavioral change, providing further support for the importance of the gut-

brain-axis.

Abstract

Animals integrate changes in external and internal environments to generate behavior. While

neural circuits detecting external cues have been mapped, less is known about how internal states

like hunger are integrated into behavioral outputs. We use the nematode *C. elegans* to decode

how changes in internal nutritional status affects chemosensory behaviors. We show that acute

food deprivation leads to a reversible decline in repellent, but not attractant, sensitivity. This

behavioral change requires two conserved transcription factors MML-1 (Mondo A) and HLH-30

(TFEB), both of which translocate from the intestinal nuclei to the cytoplasm upon food

deprivation. Next, we identify insulin-like peptides INS-23 and INS-31 as candidate ligands

relaying food-status signals from the intestine to other tissues. Furthermore, we show that ASI

chemosensory neurons use the DAF-2 insulin receptor, PI-3 Kinase, and the mTOR complex to

integrate these intestine-released peptides. Together, our study shows how internal food status

signals are integrated by transcription factors and intestine-neuron signaling to generate flexible

behaviors.

Keywords

Behavioral plasticity; sensory integration; food-deprivation, MML-1, HLH-30, insulin-like

peptides, DAF-2 receptors, ASI neurons, gut-brain axis.

Introduction

Animals evaluate their environment, integrating prior experiences and internal state

information to optimize their behaviors in order to maximize rewards and avoid threats [1].

Moreover, changes in internal states play a critical role in adjusting the animal's responses to

external stimuli [2, 3]. One critical internal state is hunger, which has a profound effect on

animal survival and elicits dramatic changes in food-seeking behaviors [2, 4]. Multiple species,

including humans, have been shown to alter their chemosensory behavior during periods of

starvation [5-10]. Despite this, less is known about how the nervous system integrates

information about hunger status.

The nematode *Caenorhabditis elegans*, with just 302 neurons [11], and 20 cells in its

intestine [12], provides a unique opportunity for a high-resolution analysis of how the nervous

system integrates internal signals. Previous studies have shown that *C. elegans*, similar to

mammals, exhibits a number of behavioral, physiological, and metabolic changes in response to

altered nutritional status. *C. elegans* hermaphrodites retain eggs [13], are unlikely to mate with

males [14], initiate altered foraging behaviors [15-17], and change their responses to

environmental CO_2 [18], salt [19], and pheromones [20] upon food deprivation. Moreover, many

molecules that signal hunger are conserved between *C. elegans* and vertebrates. For example,

neuropeptide Y (NPY) signaling influences feeding behaviors in nematodes and mammals [21-

23]. Similar effects are also seen with insulin and dopamine signaling, which seem to act via

modifying chemosensory activity and behavior in nematodes [24, 25]; and on mammalian neural

circuits [26-28] to modify feeding behavior. While neuronal pathways responding to food-

deprivation on the multiple-minute timescales have been mapped [17, 29], those integrating these

signals on the multiple hour timescales are poorly understood.

Here we use *C. elegans* to dissect the machinery required to integrate internal food

signals and modify behaviors. We combined food deprivation over multiple hours with a

behavioral assay that quantifies the animal's ability to integrate both toxic and food-related

signals, mimicking a simplified ecologically relevant scenario. In this sensory integration assay,

animals cross a toxic copper barrier (repellent) and chemotax towards a point source of a volatile

food-associated odor, diacetyl (attractant) [30]. We show that animals' food-deprived for

multiple hours have reduced sensitivity to the repellent and cross the copper barrier more readily

than well-fed animals. Next, we show that two transcription factors translocate from the

intestinal nuclei to the cytoplasm upon multiple hours of food deprivation. We confirm a role for

these transcription factors and identify the downstream peptides released by the intestine to relay

"the lack of food" signal to other tissues. Finally, we show that ASI chemosensory neurons

integrate these intestine-released peptides. This allows animals to reduce their avoidance to

repellents and undertake a higher risk strategy in their search for food.

Results

Acute food deprivation specifically alters repellent-driven behaviors

Animals simultaneously integrate both attractant and repellent signals from their

environment to generate appropriate behavioral readouts. To mimic these interactions, animals

are exposed to a copper repellent barrier ($CuSO_4$) and a gradient of a volatile attractant, diacetyl

[30]. The proportion of animals that cross the copper barrier are counted and expressed as a

chemotactic index (**Figure 1A**). We analyzed the behavior of well-fed, wild-type animals and

found that ~30% crossed the copper barrier and locomote towards the spot of diacetyl (black

bars, **Figure 1B and Movie S1**). In contrast, food-deprived animals were more likely to cross the

copper barrier (blue bars, **Figure 1B and Movie S2**). We also found that animals needed to be

food deprived for at least 1 hr before they significantly altered their behavior with a maximal

effect at 3 hrs (**Figure 1B**). Next, we tested whether the food-deprivation effect was reversible.

We food-deprived animals for 3 hrs and then returned them to food for different durations and

analyzed animal behavior after the food experience. We found that 3-hr food-deprived animals

that had been returned to food for at least 3 hrs reverted to the "well-fed" state (**Figure 1B**).

Taken together, these results indicate that food deprivation reversibly modifies sensory

integration behavior.

We then tested whether this food deprivation-evoked change in sensory integration

behavior was specific to the copper repellent and diacetyl attractant used in the assay. We

observed that food-deprived animals did not cross the repellent barrier when diacetyl was paired

with other repellents like fructose (except one intermediate concentration), sodium chloride, or

quinine (**Figure 1C**). In contrast when copper was paired other attractants like benzaldehyde and

isoamyl alcohol, food deprived animals continued to cross the copper barrier more readily than

well-fed animals (**Figure 1D**). Next, we tested responses of these animals to varying

concentrations of copper or diacetyl alone. We found that food-deprived animals crossed the

copper barrier more readily than well-fed animals, suggesting that their responsiveness to copper

is reduced even in the absence of an attractant (**Figure 1E**). In contrast, food-deprived animals

did not discernably alter their attraction to diacetyl in the absence of the copper repellant (**Figure**

1F). Given the small number of well-fed animals that cross the copper barrier alone (**Figure 1E**),

we continued to pair copper with the diacetyl attractant for further analysis. We also tested

whether altering the concentrations of the copper barrier has an effect on food-deprived animals

and confirmed these concentrations using a copper indicator (**Supplementary Figure 1**). We

found that food-deprived animals showed significant increase in their ability to cross the

repellent barrier above a threshold of 5 mM copper concentration (**Figure 1G**). To gain further

confirmation of this copper-specific change, we tested food-deprived animals in a single animal

copper drop assay (**Supplementary Figure 2A**). In this assay, the response of a single animal to

a drop of 1.5 mM CuSO$_4$ solution placed in its path was monitored. Most repellents can be tested

in this assay with animals generating a robust avoidance response [31]. We found that food-

deprived animals showed a significant deficit in their copper avoidance response

(**Supplementary Figure 2B**). Collectively, these data show that food-deprived animals display

reduced sensitivity to copper, which we dissected further using genetic methods and tracking

software.

Dynamics of risky search strategies in food-deprived animals

To analyze how food deprivation modifies animal behavior, we recorded and tracked

populations of animals over 45 minutes in the sensory integration assay. Individual animal

trajectories were identified and used for analysis (see Methods and **Supplementary Figure 3**).

We found that fewer well-fed animals cross the repellent copper barrier (**Figure 2A**) as

compared to food-deprived animals (**Figure 2B**) during the entire 45 min assay (example tracks

for all groups in **Supplementary Figure 3 E-H**). To quantify this difference, we plotted the

fraction of tracks that crossed the copper barrier as a function of time (**Figure 2C**). We found

that food-deprived, wild-type animals were more likely to cross the barrier at all time points (15,

30, and 45 mins) compared to well-fed animals. Thus, the differences between well-fed and

food-deprived animals were not limited to specific time windows in the assay. To further assess

the increased likelihood of food-deprived animals crossing the repellent barrier, we compared the

probability of animal tracks being located at given distances from the barrier (**Figure 2D,**

methods described in **Supplementary Figure 3A-C**)). We found that food-deprived worms are

nearly twice as likely to reside within +/- 0.5 cm from the copper barrier while well-fed animals

are more likely to be found 2.1 cm from the barrier, not far from where the animals were placed

on the assay plate (**Figure 2D**, statistics summarized in **Supplementary Table 2**). These data

suggest that well-fed animals reorient upon detection of the copper thereby increasing the

likelihood of animals being located in regions well before the barrier. In contrast, food-deprived

animals cross the barrier more frequently. To further dissect these behavioral differences, we

quantified the mean velocity of worm tracks (**Figure 2E**). We find that well-fed animals move

more slowly when physically closer to the copper barrier. Food-deprived worms are significantly

slower at distances far from the copper barrier (2-3.5 cm), but then accelerate to speeds matching

well-fed behavior as they approach the barrier before slowing down as they reach the barrier

consistent with well-fed animals (**Figure 2E, Supplementary Table 2**).

When this assay is run in the absence of the attractant diacetyl, there is no significant

difference in the probability of crossing the copper barrier between well-fed and food-deprived

worms (**Figure 2F**), but this may be related to the low numbers of experiments we analyzed in

this group. However, this is consistent with an increased chemotactic index of food-deprived

worms across the copper barrier in the absence of an attractant at all time points

(**Supplementary Figure 3D**). The probability of food-deprived animals locomoting close to the

copper barrier is higher for 0.3 and 0.5 cm before the barrier **(Figure 2G, Supplementary Table**

2) while the velocity of these worms as a function of distance to the barrier is distributed

similarly to worms assayed with copper and diacetyl (**Figure 2H, Supplementary Table 2).**

These data suggest that the increased likelihood of food-deprived animals to cross the copper

barrier cannot be explained by a deficiency in copper sensation alone. Further, these data suggest

that food-deprived animals display similar locomotion dynamics to well-fed animals in response

to copper in the absence of diacetyl.

In the absence of copper, food-deprived and well-fed animals behave similarly with no

significant difference in the fraction of "barrier" (no copper) crossings toward the diacetyl or

their localization on the assay plate (**Figure 2I, 2J, Supplementary Table 2**). Further, food-

deprived worms display a decreased velocity on average (**Figure 2K, Supplementary Table 2**),

consistent with previous studies [32]. Collectively, these data suggest that increase in food-

deprived animals crossing the copper barrier is not due increased mobility, but may rather be a

result of these animals pursuing navigation that is unfavorable (copper is toxic to *C. elegans*

[33]).

Lack of food and not changes in fat drives the food-deprivation induced behavioral change

Given that the change in sensory integration behavior requires multiple hours of food-

deprivation, we hypothesized that metabolic signals like changes in fat content might play a

crucial role. Additionally, previous studies have shown that prolonged starvation can deplete fat

stores in *C. elegans*, which in turn can affect behavior [34, 35]. We tested whether 3 hrs of food

deprivation alters the fat content of animals. Oil-Red O (ORO), a fat-soluble dye that stains

triglycerides and lipoproteins has been used to label and quantify fat stores in *C. elegans* (**Figure**

3A-B) [36]. We used this dye and found that 3 hrs of food-deprivation did not alter the ORO

signal or the area of the animal labeled by this stain (**Figure 3C-D**). In contrast, we observed a

significant change in the both the intensity of the signal and area of animal stained in 6-hr food-

deprived animals, consistent with previous studies [37]. These data suggest that changes in

sensory integration behavior, which occurs after 3 hrs of food-deprivation is likely to be

independent of fat metabolism.

Next, we sought to identify the relevant aspects of the bacterial experience contribute to

the food deprivation-triggered behavioral change. *C. elegans* has been shown to evaluate

multiple aspects of the food experience, including changes in food distribution, oxygen and

carbon dioxide concentrations, small molecule metabolites and others [38-40]. To uncouple the

tactile and chemosensory input of the bacteria from the nutritional value of ingesting bacteria, we

analyzed the effect of using Sephadex gel beads on animal behavior. Animals exposed to gel

beads experience the tactile input, but are not exposed to the nutritional value of food (**Figure**

3E) [15]. Notably, we found that animals exposed o Sephadex beads in the absence of *E. coli*

OP50 for 3 hrs behaved the same as food-deprived animals in the sensory integration assay

(**Figure 3F**). Together, these results show that the lack of food in the *C. elegans* intestine, but not

the absence of chemosensory cues, reduces the animal's sensitivity to copper.

Transcription factors mediate food deprivation-induced behavioral change

Our studies indicated that the lack of food inside the animal was responsible for the transient

reduction in copper sensitivity. To gain insights into the underlying molecular machinery, we

investigated the role of nutritional-responsive transcription factors in the sensory integration

assay. In mammalian cells, glucose is rapidly converted to glucose-6-phostphate, whose levels

are sensed by a two basic-helix-loop-helix-leucine zipper transcription factors, MondoA and

ChREBP (Carbohydrate Response Element Binding Protein). In well-fed conditions, MondoA

binds the excess glucose-6-phosphate and Mlx (Max-like protein X) and translocates to the

nucleus where it activates transcription of glucose-responsive genes. In the absence of glucose,

MondoA remains in the cytoplasm [41, 42] (**Figure 4A**). *C. elegans* orthologs of MondoA and

Mlx have been identified as MML-1 and MXL-2, respectively [43]. Furthermore, MML-

1/MondoA has also been shown to translocate into the intestinal nuclei under well-fed conditions

(**Figure 4A**) [44]. We predicted that *mml-1* mutants would be unable to sense the lack of food

and thereby unable to reduce copper sensitivity after food deprivation. Consistently, we found

that *mml-1*, but not *mxl-2* mutants were defective in their integration responses after food

deprivation (**Figure 4B**). We then tested whether food deprivation alters the sub-cellular

localization of the MML-1 protein. We monitored the GFP fluorescence under well-fed and

food-deprived conditions in a *mml-1* knockout transgenic animal expressing GFP fused to the

full-length coding sequence of MML-1/MondoA under well-fed and food-deprived conditions.

We found that 3 hrs of food-deprivation resulted in an increased translocalization of MML-

1/MondoA from the nucleus to the cytoplasm of the intestinal cells (**Figure 4C, 4D**). We suggest

that this cytosolic MML-1/MondoA reduces copper sensitivity by modifying signaling between

tissues.

Previous studies have shown that MML-1 regulates the activity and nuclear localization

of a second bHLH transcription factor HLH-30 (*C. elegans* TFEB, **Figure 4A**) [45]. In multiple

animal models, HLH-30 functions as a key regulator of longevity pathways by promoting

autophagy and lysosome biogenesis [46-49]. We tested whether HLH-30/TFEB was also

required for food deprivation-evoked change in sensory integration. We found that, unlike wild-

type animals, *hlh-30* null mutants did not show a change in their behavior after food-deprivation

in the sensory integration assay (**Figure 4E**). We then tested whether the subcellular localization

of HLH-30/TFEB was also affected by food deprivation. We observed an initial decrease in

cytosolic GFP fluorescence at 1 hr of food-deprivation in HLH-30::GFP transgenic animals

(Lapierre et al., 2011). Subsequently, at 3 hr of food-deprivation we found a robust increase in

cytosolic HLH-30::GFP fluorescence (**Figure 4F, 4G**). Collectively, these data show that both

MML-1 and HLH-30 accumulate in the intestinal cytoplasm upon 3 hr of food-deprivation and

are required for the consequent behavioral change in sensory integration.

Intestine-to-neuron signaling involves insulin signaling

Previous studies have shown that the *C. elegans* intestine is a major site for the transcriptional

regulation of insulin-like peptide genes in response to starvation [50]. In addition, HLH-

30/TFEB has been shown to act upstream of the insulin-signaling pathway in regulating the

expression of neuronal chemoreceptor genes [51]. The *C. elegans* genome encodes about 40

insulin-like peptides [52] and all of these ligands are thought to bind and signal via a single

tyrosine kinase DAF-2 receptor [53]. We hypothesized that insulin-like peptides might also act

downstream of HLH-30/TFEB in relaying food status signals from the intestine to other tissues.

Consistent with our hypothesis, multiple insulin-like peptides including INS-3, INS-4, INS-6,

INS-10, INS-17, INS-18, INS-23, and INS-31 contain HLH-30/TFEB binding sites in their

promoters [51]. In addition, INS-7, INS-8 and INS-37 have been shown to affect the subcellular

localization of HLH-30/TFEB in the *C. elegans* intestine after mating (**Figure 5A**). [54]. We

tested mutants in these insulin-like peptide genes for their ability to alter their sensory integration

behavior after food deprivation. We similarly tested animals with a semi-dominant mutation in

daf-28(*sa191*), since this allele has been shown to the prevent other insulin-like peptides from

binding the common DAF-2 insulin receptor [55-57]. We found that multiple insulin-like

peptides including INS-23 (*tm1875*), INS-31 (*tm3543*) and DAF-28 (*sa191*) were unable to

respond to food deprivation. Specifically, these mutant animals did not display an increased

ability to cross the repellent copper barrier when food deprived (**Figure 5B**), implying that these

might be candidate signals relaying food status signals. In contrast, *ins-7* (*tm2001*), *ins-8*

(*tm4144*)*,* and *ins-37* (*tm6061*) mutants were similar to wild-type animals in their ability to cross

the copper barrier in both well-fed and food-deprived conditions (**Figure 5C**). Taken together,

these data suggest that the intestine might release INS-23, INS-31 and DAF-28 (or other insulin-

like peptides whose binding to the DAF-2 insulin receptor is blocked by DAF-28) to relay

hunger information to other tissues.

Next, we probed the role of the insulin receptor, DAF-2, in affecting 3 hr-food

deprivation evoked changes in sensory integration. Consistent with our analysis of mutants in

various insulin-like peptide genes, we found that mutants in the insulin receptor, DAF-2, were

also defective in their response to food deprivation (**Figure 5D**). To localize the site of DAF-2

action, we analyzed the effect of rescuing this receptor in different tissues. We found that

expressing *daf-2* under neuronal, but not intestine or pharyngeal muscle promoters [57] restored

normal behavior to the *daf-2* mutants (**Figure 5D**). Taken together, these results suggest that

neuronally expressed DAF-2 receptors might detect INS-23, INS-31, and other insulin-like

peptides released from the intestine, particularly those hindered by DAF-28 binding.

We then sought to test whether the upstream MML-1/Mondo A and the downstream

DAF-2 insulin receptor function in the same pathway. We generated an *daf-2,mml-1* double

mutant, which did not show any additional defects in the 3 hr food-deprivation evoked change in

sensory integration when compared to either *mml-1* or *daf-2* single mutant (**Figure 5E**),

suggesting that these two might function in the same pathway. We also found that expressing

MML-1 in the intestine alone was not sufficient to restore wild-type behavior to the double

mutant. In contrast, restoring MML-1 to the intestine and DAF-2 in ASI sensory neurons

restored normal integration response after food deprivation (**Figure 5E**). Together, these data

show that while MML-1 is required in the intestine, DAF- 2 is required in ASI neurons and these

genes act in the same pathway to alter food deprivation-modulated integration behavior (**Figure**

5F).

ASI chemosensory neurons use insulin-signaling pathways to integrate intestine-released

peptide signals

We then sought to identify components of the DAF-2 signaling pathway (**Figure 6A**) in ASI

chemosensory neurons that were required to alter food-deprivation evoked change in sensory

integration. We observed that mutants in the insulin-signaling pathway components including the

FOXO family transcription factor *daf-16*, serine/threonine kinases AKT-1, AKT-2 (*akt-1*, *akt-2*),

3-phosphoinositide-dependent kinase 1 (*pdk-1*) and lipid phosphatase (*daf-18*, PTEN suppressor)

performed normally in the sensory integration assay after food deprivation (**Figure 6B**) [58, 59].

In contrast, mutants in the phosphoinositide 3-kinase (PI3K, *age-1*) and Rictor (*rict-1)* [a key

component of the mTORC2 complex [60, 61]] were defective in their copper sensitivity after

food deprivation (**Figure 6B and 6C**). Similar to our *daf-2* rescue experiments, we found that

Rictor was also required in ASI neurons to restore normal food-deprivation behavior to *rict-1*

mutants (**Figure 6C**). These results suggest that PI-3 Kinase and Rictor might function in the

same pathway downstream of DAF-2 receptors in ASI neurons. Previously, Rictor has been

shown to act in the intestine to mediate an intestine-to-neuron signaling to affect dauer formation

[62]. Our results show that Rictor can also act in ASI neurons as part of an intestine-to-neuron

signal to alter sensory integration behavior. Collectively, we suggest that food deprivation

engages DAF-2 signaling in ASI chemosensory neurons to alter the animal's copper sensitivity

allowing it to cross the copper barrier more readily.

Discussion

We used food deprivation in *C. elegans* as a model to understand how food deprivation modifies

behavior. We show that food-deprived animals reversibly alter their behavior by reducing their

repellent responsiveness, allowing them to traverse potentially toxic environments in their search

for food. The *C. elegans* intestine sense the lack of food leading to cytosolic MML-1 and HLH-

30, which in turn promotes the release of multiple insulins (INS-23, INS-31 and potentially

others). These intestine-released peptides bind DAF-2 receptors and are processed by

downstream PI-3 Kinase and RICT-1 in ASI and other neurons to reduce copper sensitivity and

alter behavior (**Figure 6D**).

Multicellular animals' sense and regulate glucose homeostasis at several levels. While

insulin and glucagon maintain constant levels of circulating glucose, the Myc-family

transcription factors are used within cells. Glucose uses cell membrane-localized transporters to

enter cells, where it is rapidly converted into glucose-6-phosphate [63]. This intermediate

metabolite is sensed by the Myc-Max complex, which binds glucose-6-phosphate and

translocates to the nucleus where it regulates the transcription of glucose-responsive genes [41].

While the role of ChREBP/MondoA-Mlx-glucose-6-phosphate complex in regulating

transcription is well studied [42, 64, 65], the role of these proteins in the cytoplasm remains

poorly understood. We show a specific role for MML-1 (MondoA homolog), but not MXL-2

(Mlx homolog) in the intestine in reducing copper sensitivity after food deprivation.

Additionally, we show that HLH-30, an ortholog of TFEB, is also required for attenuating

copper sensitivity after food deprivation. Intriguingly, MML-1/MondoA and HLH-30/TFEB are

both basic helix-loop-helix transcription factors and have been shown to act in concert to modify

signaling networks and affecting global states like reproduction or survival [45]. Like MML-

1/MondoA, a role for HLH-30/TFEB in the cytoplasm has also not been defined. We suggest

that MML-1/MondoA and HLH-30/TFEB accumulation in the cytoplasm (in food deprived

animals) enables the intestine to release peptide(s) relaying a "lack of glucose" signal to other

tissues.

Helix-loop-helix transcription factors in *C. elegans* like MML-1/MondoA and HLH-

30/TEFB are known to bind similar E-box elements (CACGTG) and have large overlap in their

target gene expression [43, 45]. Additionally, previous studies have identified multiple insulin-

like peptide genes whose expression is regulated by HLH-30 and other insulin-like peptide

genes, which can affect the subcellular localization of HLH-30 [51, 54]. We screened this subset

of insulin-like peptide genes to identify multiple candidates relaying food status signals from the

intestine to other tissues. The *C. elegans* intestine has been previously shown to be a key tissue

where the transcription of insulin peptide genes is regulated [50]. While we have not directly

demonstrated that our candidate insulin-like peptides are released from the intestine, we

speculate that food deprivation promotes their release relaying the "lack of food" signal.

We show that the ASI chemosensory neurons use the tyrosine kinase insulin receptor

(DAF-2) to integrate these signals. Three lines of evidence suggest that the intestine is releasing

multiple insulin-like peptide(s) - first, mutants in *ins-23*, *ins-31* and *daf-28* are defective in their

food-deprivation evoked change in behavior, second, the insulin receptor (DAF-2) integrates

these signals and third, MML-1 is required in the intestine and acts in the same pathway as the

DAF-2 receptor, which acts in ASI neurons. We also define additional insulin signaling pathway

components in ASI neurons. While FOXO (DAF-16), AKT kinase -1 and -2, PDK-1 and PTEN

(DAF-18) are not required, we show that AGE-1 (PI-3 Kinase) and Rictor (a component of the

mTORC2 complex) are required to integrate intestine-released peptide signals. While PI-3 kinase

has been shown to act via AKT kinase to activate the Rictor [60, 61], our result hint at an AKT

kinase-independent mechanism for PI-3 Kinase to signal to Rictor and the mTORC2 complex.

Multiple studies have also highlighted the role of insulin signaling in relaying starvation-

related signals to various neurons. Starvation has been shown to decrease the secretion of INS-18

from the intestine, which antagonizes DAF-2 receptor in ADL neurons and modifies pheromone-

mediated behaviors [20]. Also, starvation has been shown to be associated with increased

octopamine signaling, which transforms CO_2 attraction to repulsion in starved animals [18].

Moreover, starvation has also been shown to recruit ASG neurons to cooperate with ASE

neurons and drive avoidance to high salt [19]. We speculate that our intestine-to-neuron insulin

signaling pathway leads to altered ASI function and altered copper sensitivity. Consistently,

starvation has been shown to increase ASI neural activity in response to food-stimuli [66]. These

data are also consistent with previous studies showing that ASI neurons playing a crucial role in

modifying behavior after 6 hours of food deprivation [17, 67]. Taken together, we speculate that

food deprivation leads to an increase in insulin signaling from the intestine to ASI neurons,

which alters neuronal activity and reduces the animal's sensitivity to copper, allowing it to cross

the barrier more readily. More broadly, these studies link transcription factors and insulin

signaling from the intestine to neurons to modify sensory behavior, a mechanism likely

conserved across species.

Methods

Strains

C. elegans strains were grown and maintained under standard conditions [68]. All strains used

are listed in **Supplementary Table 1**.

Behavior Assays

All animals were grown to adulthood on regular nematode growth medium (NGM) plates seeded

with OP50 ($OD_{600} \sim 0.2$) before they were washed and transferred to new food (standard NGM

plates seeded with OP50) or food-free plates (standard NGM plates) respectively for the

indicated duration. Sephadex beads (G-200) were added to both the empty NGM plate and the

OP50 lawn in experiments for Figure 3E-F. Sensory integration assays were performed on 2%

agar plates containing 5 mM potassium phosphate (pH 6), 1 mM $CaCl_2$ and 1 mM $MgSO_4$, made

the day before the experiment. Repellent gradients (including $CuSO_4$ (Copper (II) sulfate

pentahydrate, Sigma 209198), glycerol (, NaCl (, fructose (D-(-) Fructose Sigma F0127), and

quinine (Sigma 22620)) were established by dripping 25 µl of solution, dissolved in water,

across the midline of the plate [30]. This solution was allowed to dry overnight. Prior to the

assay, the animals were washed from the food or food-free plates into Eppendorf tubes. Each

treatment group was serially washed once with M9+MgSO$_4$ and 3 times with Chemotaxis buffer

(5 mM potassium phosphate (pH 6, Fisher BP362 monobasic and Fisher BP363, dibasic), 1 mM

CaCl$_2$ (Sigma C1016) and 1 mM MgSO$_4$ (Sigma M7506)) before being transferred to the assay

plates. Glass Pasteur pipets were used to prevent loss of animals due to sticking in plastic pipette

tips. Immediately after plating 100-200 animals in a small drop of chemotaxis buffer, 1 μL of

attractant with 1 μL of 1M sodium azide in water (Sigma 71289) was placed on the opposite side

of the chemotaxis plate. Attractants used were diacetyl (2,3-Butanedione Sigma 11038), Isoamyl

alcohol (3-methyl-1-butanol, Sigma 77664), and Benzaldehyde (Sigma 418099) diluted in

Ethanol. If necessary, the small drop of animals was dabbed gently with the edge of a Kim wipe

and the lid was immediately replaced. After 45 minutes or at indicated times, the integration

index was computed as the number of worms in the odor half of the plate divided by the total

number of animals on the plate. For each experiment, at least two plates were tested each day

with experiments performed on at least three different days. Unless otherwise noted, the repellant

is a dried stripe of 25 μL 50 mM CuSO$_4$ (Copper (II) sulfate pentahydrate, Sigma 209198) in

water and the attractant is 1 μL 0.2% diacetyl (2,3-Butanedione Sigma 11038) diluted 1:500 in

100% ethanol.

Statistics

For sensory integration, experiments were performed at least 3 times with at least 2 plates per

genotype/condition (unless otherwise noted). For strains with extrachromosomal arrays, only

animals expressing the co-injection markers were counted. Every condition was performed with

N2 (wild-type) controls at the same time. Two-way ANOVAs with post-hoc Bonferroni-

corrected multiple comparisons were performed across WF/FD conditions, only if the factor was

significant. For all figures, p values are represented by: * $p<0.05$, ** $p<0.01$, *** $p<0.001$, ****

$p<0.0001$.

Single animal avoidance assay: Copper drop test

Experiments were performed as previously described [31]. Animals are moved from a food to a

food-free assay plate. A capillary tube is used to deliver a drop of test compound (1.5 mM

$CuSo_4$) 0.5 - 1 mm away from the head of the animal and its responses scored. Positive

avoidance indicates an animal executing a large reversal and omega bend within 3 seconds of

sensing the test compound. Five animals are tested per condition and with each animal exposed

to 10 drops and the percent avoidance is plotted. Assay is replicated at least three times by an

investigator who is blind to the conditions being tested.

Tracking

Sensory integration behavior assays using 50 mM $CuSO_4$ (Copper (II) sulfate pentahydrate,

Sigma 209198) in water and 1 μL 0.2% diacetyl (2,3-Butanedione Sigma 11038) attractant

(1:500 in 100% Ethanol) were performed with well-fed and food-deprived animals. Animal

behavior was recorded for 45 minutes using a Pixelink camera (1024x1024 pixels at 3 frames per

second). The imaging field of view was approximately 47 mm x 47 mm. WormLab software

(MBF Bioscience) was used to identify and track the midpoints of worms in each video. Custom

MATLAB software (https://github.com/shreklab/Matty-et-al-2021) was used to further clean the

data (i.e. remove putative tracks that did not correspond to animal behavior) and analyze

individual tracks. Tracks were excluded if they met any of the following criteria: 1) overlapped

with shadows or markings; 2) lasted less than 10 seconds; 3) travelled fewer than 30 pixels^2; or

4) traveled less than 10 pixels in any direction. Valid animal tracks were then plotted (**Figure**

2A, 2B) and analyzed as described below.

Tracking analysis

The number of animals in each experiment is estimated from the maximum number of

simultaneous tracks identified in a single frame. An average of 33.2 ± 10.4 animals were assayed

across all conditions. Because the field of view does not encompass the entire plate, the number

of tracks identified in each frame decreases over time as animals crawl to other regions of the

plate. To quantify the number of animals crossing the copper barrier as a function of time, the

number of unique tracks that started past the copper barrier was divided by the number of unique

tracks in the entire field-of-view. This fraction of cumulative unique tracks that crossed the

copper barrier was calculated for 15, 30, and 45 minutes (**Figure 2C, 2F, 2I; Supplementary**

Figure 3A). To better understand animals' avoidance of copper and attraction to diacetyl, the

probability of an animal residing at a particular distance from the barrier was calculated for 1

mm bins. The total number of tracked midpoints at each time point located in each 1 mm bin was

summed and divided by the total number of tracked midpoints across all bins (**Figure 2D, 2G,**

2J; Supplementary Figure 3C). Additionally, animal velocity was calculated by computing the

Euclidean distance of a track over a 2 second window. In each video, the average velocity of all

tracks was computed as a function of distance from the copper barrier in 1 mm bins (**Figure 2E,**

2H, 2K; Supplementary Figure 3B).

Visualizing Copper Gradients

Copper sulfate gradients were visualized using 1-(2-Pyridylazo)-2-naphthol (PAN, Sigma 101036). Plates with 25 μL of 5 mM, 25 mM, 50 mM and 100 mM $CuSO_4$ dripped down the midline were dried overnight. 1 mL of 0.01% PAN indicator was added to plates the next day and allowed to dry. The plates with PAN indicator were incubated overnight and imaged the following day to allow for saturation of the signal. Images and quantification of the copper barrier is shown in **Supplementary Figure 1**.

Fat quantification

Oil red O staining was conducted as previously described [36]. Briefly, 10-20 N2 adults were allowed to lay eggs for 1 hour on NGM plates seeded with OP50. The adults were removed and eggs were allowed to develop for 3 days. These day-1 adult animals were either removed from food and placed on an empty NGM plate for 3 hours or 6 hours or placed on a new plate with OP50 food. 5 mg/mL Oil Red O (Sigma, O9755) in 100% isopropanol was prepared as a working solution and diluted 3:2 in 60% isopropanol on the day before use. Mixture was kept from the light and filtered using a 0.2 μm cellulose acetate syringe filter and allowed to mix on a rocker overnight. Animals were washed off plates with PBST (PBS + 0.01% Triton X-100 (Sigma, X100)at the appropriate times and washed once. Animals were fixed in 40% isopropanol and shaken at room temperature for 3 minutes. Isopropanol was removed and 600 μL of the Oil Red O diluted solution was added to each tube. Each tube was nutated for 2 hours at room temperature, away from light. Animals were washed once with PBST and nutated for another 30 minutes. Animals were washed once more and prepared for imaging. Approximately 20 worms from each treatment group were pipetted onto a microscope slide and covered with a coverslip.

Images were collected on upright Zeiss Axio Imager.M2 at 10X using an AxioCam 506 Color

camera. Images were quantified using color deconvolution in ImageJ, normalized to background

and an unstained region of an animal. Within each experiment, the same thresholds were used

across treatments. Approximately 20 animals were quantified within each condition on each

experimental day, performed across three different days.

Imaging

Transgenic animals (HLH-30::GFP and MML-1::GFP) were grown to day 1 adulthood (3 days

post hatching) via a one-hour hatch off on standard NGM plates seeded with OP50. Animals

were picked onto empty NGM plates for 1, 2, and 3 hours for food deprivation or placed on a

new NGM plates with OP50. Animals were picked onto thin agar pads on microscope slides and

anesthetized with 100 µM tetramisole hydrochloride (Sigma-Aldrich L9756) immediately prior

to imaging. Animals were imaged at 10X using an upright Zeiss Axio Imager .M2. At least 12

animals per group on three different days were imaged and qualitatively analyzed for localization

to primarily cytoplasmic, nuclear, or both in intestinal cells, with the investigator blind to food

deprivation status.

Molecular Biology and Transgenics

The following primers were used for amplifying full-length cDNAs:

mml-1

forward 5 'TATTTAGCTAGCATGTCGCGCGGGCAGATTATACACAG

reverse 5'CGGGGTACCGAGCAGTTCAAAATGGATTTTTGAGTTGTTGC

rict-1

forward 5'TATTTAGCTAGCATGGACACTCGTCGAAAAGTGTATCAC

reverse 5'CGGGGTACCTAAAAGATTTGCTGCAGGAATGCTCTCG

daf-2

forward 5' TATTTAGCTAGCAATGAATATTGTCAGATGTCGGAGACGA 3'

reverse 5' CGGGGTACCTCAGACAAGTGGATGATGCTCATTATC 3'

cDNAs corresponding to the entire coding sequences of *mml-1, rict-1,* and *daf-2* genomic region

were amplified by PCR using primers above and expressed under tissue or cell selective

promoters.

Tissue specific expression was achieved with *Prgef-1* for neurons, *Pges-1* and *Pgly-19* for the

intestine, and *Pmyo-2* for pharynx, [69-74]. Cell-specific expression used *Pstr-3* was used for

cell-specific expression in ASI neurons [75]. For all experiments, a splice leader (SL2) fused to

mCherry or *gfp* transgene was used to confirm expression of the gene of interest in either specific

cells or tissues. Germline transformations were performed by microinjection of plasmids [76] at

concentrations between 50 and 100 ng/µl with 10 ng/µl of *elt-2::gfp* as a co-injection marker.

Author Contributions

M.A.M. and H.E.L. conceived and conducted the experiments, interpreted the data, and co-wrote

the paper. A.S., A.C., and K.K., conducted behavioral assays; A.S.' experiments were conducted

in M.H.'s laboratory. J.A.H. analyzed tracking data. S.H.C. conceived the experiments,

interpreted the data and co-wrote the paper. All co-authors provided feedback on the manuscript.

Acknowledgements

We thank A. Dillin, T. Ishihara, S. Lockery, A. Samuelson, E. Troemel, M. Zhen, the National

BioResource Project (NBRP, Japan) and Caenorhabditis Genetics Center (CGC) for strains; C.

Bargmann, E. Hallem, M. Hilliard, A. van der Linden, P. McGrath, D. Pilgrim and P. Sengupta

for constructs; S. Srinivasan and lab members for RNAi clones and help with Oil-Red O

staining; and M. Tamés, Z. Liu, and C. Yang for technical help with behavioral and imaging

studies. We are also grateful to Jing Wang, and members of the Chalasani lab for critical

comments, advice, and insights. This work was funded by grants from The Rita Allen

Foundation, The W.M. Keck Foundation and NIH R01MH096881 to S.H.C., NSF Postdoctoral

Research Fellowships in Biology Program Under Grant No. 2011023 (M.A.M), Socrates

Program (H.E.L. Award #NSF-742551) and Graduate Research Fellowship from NSF (H.E.L.,

and J.A.H.).

References

1.   Dugatkin LA. Principles of animal behavior: University of Chicago Press; 2020.

2.   Sternson SM, Nicholas Betley J, Cao ZF. Neural circuits and motivational processes for

hunger. Curr Opin Neurobiol. 2013;23(3):353-60. doi: 10.1016/j.conb.2013.04.006. PubMed

PMID: 23648085; PubMed Central PMCID: PMC3948161.

3.   Taghert PH, Nitabach MN. Peptide neuromodulation in invertebrate model systems.

Neuron. 2012;76(1):82-97. doi: 10.1016/j.neuron.2012.08.035. PubMed PMID: 23040808;

PubMed Central PMCID: PMC3466441.

4.   Atasoy D, Betley JN, Su HH, Sternson SM. Deconstruction of a neural circuit for hunger.

Nature. 2012;488(7410):172-7. doi: 10.1038/nature11270. PubMed PMID: 22801496; PubMed

Central PMCID: PMC3416931.

5.   Gillette R, Huang RC, Hatcher N, Moroz LL. Cost-benefit analysis potential in feeding

behavior of a predatory snail by integration of hunger, taste, and pain. Proc Natl Acad Sci U S A.

2000;97(7):3585-90. PubMed PMID: 10737805; PubMed Central PMCID: PMC16283.

6.   Kawai K, Sugimoto K, Nakashima K, Miura H, Ninomiya Y. Leptin as a modulator of

sweet taste sensitivities in mice. Proc Natl Acad Sci U S A. 2000;97(20):11044-9. doi:

10.1073/pnas.190066697. PubMed PMID: 10995460; PubMed Central PMCID: PMC27145.

7.   Sengupta P. The belly rules the nose: feeding state-dependent modulation of peripheral

chemosensory responses. Curr Opin Neurobiol. 2013;23(1):68-75. doi:

10.1016/j.conb.2012.08.001. PubMed PMID: 22939570; PubMed Central PMCID:

PMC3524363.

8.   Inagaki HK, Panse KM, Anderson DJ. Independent, reciprocal neuromodulatory control

of sweet and bitter taste sensitivity during starvation in Drosophila. Neuron. 2014;84(4):806-20.

doi: 10.1016/j.neuron.2014.09.032. PubMed PMID: 25451195; PubMed Central PMCID:

PMC4365050.

9.   Smeets PA, de Graaf C, Stafleu A, van Osch MJ, Nievelstein RA, van der Grond J. Effect

of satiety on brain activation during chocolate tasting in men and women. The American journal

of clinical nutrition. 2006;83(6):1297-305.

10.   O'doherty J, Rolls ET, Francis S, Bowtell R, McGlone F, Kobal G, et al. Sensory-specific

satiety-related olfactory activation of the human orbitofrontal cortex. Neuroreport.

2000;11(2):399-403.

11.   White JG, Southgate E, Thomson JN, Brenner S. The structure of the nervous system of

the nematode Caenorhabditis elegans. Philos Trans R Soc Lond B Biol Sci. 1986;314(1165):1-

340. PubMed PMID: 22462104.

12.   McGhee JD. The C. elegans intestine. WormBook. 2007:1-36. Epub 2007/12/01. doi:

10.1895/wormbook.1.133.1. PubMed PMID: 18050495.

13.   Trent C, Tsuing N, Horvitz HR. Egg-laying defective mutants of the nematode

Caenorhabditis elegans. Genetics. 1983;104(4):619-47. PubMed PMID: 11813735.

14.   Lipton J, Kleemann G, Ghosh R, Lints R, Emmons SW. Mate searching in

Caenorhabditis elegans: a genetic model for sex drive in a simple invertebrate. J Neurosci.

2004;24(34):7427-34. PubMed PMID: 15329389.

15.   Sawin ER, Ranganathan R, Horvitz HR. C. elegans locomotory rate is modulated by the

environment through a dopaminergic pathway and by experience through a serotonergic

pathway. Neuron. 2000;26(3):619-31. PubMed PMID: 10896158.

16.   Hills T, Brockie PJ, Maricq AV. Dopamine and glutamate control area-restricted search

behavior in Caenorhabditis elegans. J Neurosci. 2004;24(5):1217-25. Epub 2004/02/06. doi:

10.1523/JNEUROSCI.1569-03.2004

24/5/1217 [pii]. PubMed PMID: 14762140.

17.   Gray JM, Hill JJ, Bargmann CI. A circuit for navigation in Caenorhabditis elegans. Proc

Natl Acad Sci U S A. 2005;102(9):3184-91. PubMed PMID: 15689400.

18.   Rengarajan S, Yankura KA, Guillermin ML, Fung W, Hallem EA. Feeding state sculpts a

circuit for sensory valence in Caenorhabditis elegans. Proc Natl Acad Sci U S A.

2019;116(5):1776-81. Epub 2019/01/18. doi: 10.1073/pnas.1807454116. PubMed PMID:

30651312; PubMed Central PMCID: PMCPMC6358703.

19.   Jang MS, Toyoshima Y, Tomioka M, Kunitomo H, Iino Y. Multiple sensory neurons

mediate starvation-dependent aversive navigation in Caenorhabditis elegans. Proc Natl Acad Sci

U S A. 2019;116(37):18673-83. Epub 2019/08/29. doi: 10.1073/pnas.1821716116. PubMed

PMID: 31455735; PubMed Central PMCID: PMCPMC6744849.

20.   Ryu L, Cheon Y, Huh YH, Pyo S, Chinta S, Choi H, et al. Feeding state regulates

pheromone-mediated avoidance behavior via the insulin signaling pathway in Caenorhabditis

elegans. EMBO J. 2018;37(15). Epub 2018/06/22. doi: 10.15252/embj.201798402. PubMed

PMID: 29925517; PubMed Central PMCID: PMCPMC6068425.

21.   Mercer RE, Chee MJ, Colmers WF. The role of NPY in hypothalamic mediated food

intake. Front Neuroendocrinol. 2011;32(4):398-415. doi: 10.1016/j.yfrne.2011.06.001. PubMed

PMID: 21726573.

22.   Nassel DR, Wegener C. A comparative review of short and long neuropeptide F signaling

in invertebrates: Any similarities to vertebrate neuropeptide Y signaling? Peptides.

2011;32(6):1335-55. doi: 10.1016/j.peptides.2011.03.013. PubMed PMID: 21440021.

23.   de Bono M, Bargmann CI. Natural variation in a neuropeptide Y receptor homolog

modifies social behavior and food response in C. elegans. Cell. 1998;94(5):679-89. PubMed

PMID: 9741632.

24.   Chalasani SH, Kato S, Albrecht DR, Nakagawa T, Abbott LF, Bargmann CI.

Neuropeptide feedback modifies odor-evoked dynamics in Caenorhabditis elegans olfactory

neurons. Nat Neurosci. 2010;13(5):615-21. Epub 2010/04/07. doi: nn.2526 [pii]

10.1038/nn.2526. PubMed PMID: 20364145; PubMed Central PMCID: PMC2937567.

25.   Ezcurra M, Tanizawa Y, Swoboda P, Schafer WR. Food sensitizes C. elegans avoidance

behaviours through acute dopamine signalling. Embo J. 2011;30(6):1110-22. Epub 2011/02/10.

doi: emboj201122 [pii]

10.1038/emboj.2011.22. PubMed PMID: 21304491; PubMed Central PMCID: PMC3061029.

26. Air EL, Benoit SC, Clegg DJ, Seeley RJ, Woods SC. Insulin and leptin combine

additively to reduce food intake and body weight in rats. Endocrinology. 2002;143(6):2449-52.

doi: 10.1210/endo.143.6.8948. PubMed PMID: 12021212.

27. Berthoud HR. Metabolic and hedonic drives in the neural control of appetite: who is the

boss? Curr Opin Neurobiol. 2011;21(6):888-96. doi: 10.1016/j.conb.2011.09.004. PubMed

PMID: 21981809; PubMed Central PMCID: PMC3254791.

28. Figlewicz DP, Sipols AJ. Energy regulatory signals and food reward. Pharmacol

Biochem Behav. 2010;97(1):15-24. doi: 10.1016/j.pbb.2010.03.002. PubMed PMID: 20230849;

PubMed Central PMCID: PMC2897918.

29. Chalasani SH, Chronis N, Tsunozaki M, Gray JM, Ramot D, Goodman MB, et al.

Dissecting a circuit for olfactory behaviour in Caenorhabditis elegans. Nature.

2007;450(7166):63-70. Epub 2007/11/02. doi: nature06292 [pii]

10.1038/nature06292. PubMed PMID: 17972877.

30. Ishihara T, Iino Y, Mohri A, Mori I, Gengyo-Ando K, Mitani S, et al. HEN-1, a secretory

protein with an LDL receptor motif, regulates sensory integration and learning in Caenorhabditis

elegans. Cell. 2002;109(5):639-49. Epub 2002/06/14. doi: S0092867402007481 [pii]. PubMed

PMID: 12062106.

31. Hilliard MA, Bargmann CI, Bazzicalupo P. C. elegans responds to chemical repellents by

integrating sensory inputs from the head and the tail. Curr Biol. 2002;12(9):730-4. PubMed

PMID: 12007416.

32. Lüersen K, Faust U, Gottschling D-C, Döring F. Gait-specific adaptation of locomotor

activity in response to dietary restriction in Caenorhabditis elegans. Journal of Experimental

Biology. 2014;217(14):2480-8. doi: 10.1242/jeb.099382.

33.   Williams PL, Dusenbery DB. A promising indicator of neurobehavioral toxicity using the

nematode Caenorhabditis elegans and computer tracking. Toxicol Ind Health. 1990;6(3-4):425-

40. PubMed PMID: 2237928.

34.   Hyun M, Davis K, Lee I, Kim J, Dumur C, You Y-J. Fat metabolism regulates satiety

behavior in C. elegans. Scientific reports. 2016;6(1):1-9.

35.   Lemieux GA, Ashrafi K. Investigating connections between metabolism, longevity, and

behavior in Caenorhabditis elegans. Trends in Endocrinology & Metabolism. 2016;27(8):586-96.

36.   Escorcia W, Ruter DL, Nhan J, Curran SP. Quantification of lipid abundance and

evaluation of lipid distribution in Caenorhabditis elegans by nile red and oil red O staining.

Journal of visualized experiments: JoVE. 2018;(133).

37.   Srinivasan S, Sadegh L, Elle IC, Christensen AG, Faergeman NJ, Ashrafi K. Serotonin

regulates C. elegans fat and feeding through independent molecular mechanisms. Cell Metab.

2008;7(6):533-44. Epub 2008/06/05. doi: S1550-4131(08)00144-7 [pii]

10.1016/j.cmet.2008.04.012.   PubMed   PMID: 18522834;   PubMed   Central   PMCID:

PMC2495008.

38.   Calhoun AJ, Tong A, Pokala N, Fitzpatrick JA, Sharpee TO, Chalasani SH. Neural

Mechanisms for Evaluating Environmental Variability in Caenorhabditis elegans. Neuron.

2015;86(2):428-41.   doi: 10.1016/j.neuron.2015.03.026.   PubMed   PMID: 25864633;   PubMed

Central PMCID: PMC4409562.

39.   Carrillo MA, Hallem EA. Gas sensing in nematodes. Mol Neurobiol. 2015;51(3):919-31.

doi: 10.1007/s12035-014-8748-z. PubMed PMID: 24906953.

40.   Ludewig AH, Schroeder FC. Ascaroside signaling in C. elegans. WormBook. 2013:1-22.

Epub 2013/01/29. doi: 10.1895/wormbook.1.155.1. PubMed PMID: 23355522; PubMed Central

PMCID: PMC3758900.

41.   Havula E, Hietakangas V. Glucose sensing by ChREBP/MondoA-Mlx transcription

factors. Semin Cell Dev Biol. 2012;23(6):640-7. doi: 10.1016/j.semcdb.2012.02.007. PubMed

PMID: 22406740.

42.   Stoltzman CA, Peterson CW, Breen KT, Muoio DM, Billin AN, Ayer DE. Glucose

sensing by MondoA:Mlx complexes: a role for hexokinases and direct regulation of thioredoxin-

interacting protein expression. Proc Natl Acad Sci U S A. 2008;105(19):6912-7. doi:

10.1073/pnas.0712199105. PubMed PMID: 18458340; PubMed Central PMCID: PMC2383952.

43.   Grove CA, De Masi F, Barrasa MI, Newburger DE, Alkema MJ, Bulyk ML, et al. A

multiparameter network reveals extensive divergence between C. elegans bHLH transcription

factors. Cell. 2009;138(2):314-27. doi: 10.1016/j.cell.2009.04.058. PubMed PMID: 19632181;

PubMed Central PMCID: PMC2774807.

44.   Johnson DW, Llop JR, Farrell SF, Yuan J, Stolzenburg LR, Samuelson AV. The

Caenorhabditis elegans Myc-Mondo/Mad complexes integrate diverse longevity signals. PLoS

Genet. 2014;10(4):e1004278. doi: 10.1371/journal.pgen.1004278. PubMed PMID: 24699255;

PubMed Central PMCID: PMC3974684.

45.   Nakamura S, Karalay Ö, Jäger PS, Horikawa M, Klein C, Nakamura K, et al. Mondo

complexes regulate TFEB via TOR inhibition to promote longevity in response to gonadal

signals. Nature communications. 2016;7(1):1-15.

46.   Lapierre LR, Gelino S, Melendez A, Hansen M. Autophagy and lipid metabolism

coordinately modulate life span in germline-less C. elegans. Curr Biol. 2011;21(18):1507-14.

doi: 10.1016/j.cub.2011.07.042. PubMed PMID: 21906946; PubMed Central PMCID:

PMC3191188.

47.   Pyo J-O, Yoo S-M, Ahn H-H, Nah J, Hong S-H, Kam T-I, et al. Overexpression of Atg5

in mice activates autophagy and extends lifespan. Nature communications. 2013;4(1):1-9.

48.   Settembre C, Di Malta C, Polito VA, Arencibia MG, Vetrini F, Erdin S, et al. TFEB links

autophagy to lysosomal biogenesis. science. 2011;332(6036):1429-33.

49.   O'Rourke EJ, Ruvkun G. MXL-3 and HLH-30 transcriptionally link lipolysis and

autophagy to nutrient availability. Nature cell biology. 2013;15(6):668-76.

50.   Chen Y, Baugh LR. Ins-4 and daf-28 function redundantly to regulate C. elegans L1

arrest. Developmental biology. 2014;394(2):314-26.

51.   Gruner M, Grubbs J, McDonagh A, Valdes D, Winbush A, van der Linden AM. Cell-

Autonomous and Non-Cell-Autonomous Regulation of a Feeding State-Dependent

Chemoreceptor Gene via MEF-2 and bHLH Transcription Factors. PLoS Genet.

2016;12(8):e1006237. Epub 2016/08/04. doi: 10.1371/journal.pgen.1006237. PubMed PMID:

27487365; PubMed Central PMCID: PMCPMC4972359.

52.   Hobert O. The neuronal genome of Caenorhabditis elegans. WormBook. 2013:1-106. doi:

10.1895/wormbook.1.161.1. PubMed PMID: 24081909.

53.   Pierce SB, Costa M, Wisotzkey R, Devadhar S, Homburger SA, Buchman AR, et al.

Regulation of DAF-2 receptor signaling by human insulin and ins-1, a member of the unusually

large and diverse C. elegans insulin gene family. Genes Dev. 2001;15(6):672-86. PubMed

PMID: 11274053.

54.   Shi C, Booth LN, Murphy CT. Insulin-like peptides and the mTOR-TFEB pathway

protect Caenorhabditis elegans hermaphrodites from mating-induced death. eLife. 2019;8. Epub

2019/07/10. doi: 10.7554/eLife.46413. PubMed PMID: 31282862; PubMed Central PMCID:

PMCPMC6697448.

55.   Li W, Kennedy SG, Ruvkun G. daf-28 encodes a C. elegans insulin superfamily member

that is regulated by environmental cues and acts in the DAF-2 signaling pathway. Genes Dev.

2003;17(7):844-58. PubMed PMID: 12654727.

56.   Gruner M, Nelson D, Winbush A, Hintz R, Ryu L, Chung SH, et al. Feeding state, insulin

and NPR-1 modulate chemoreceptor gene expression via integration of sensory and circuit

inputs. PLoS genetics. 2014;10(10):e1004707-e. doi: 10.1371/journal.pgen.1004707. PubMed

PMID: 25357003.

57.   Hung WL, Wang Y, Chitturi J, Zhen M. A Caenorhabditis elegans developmental

decision requires insulin signaling-mediated neuron-intestine communication. Development.

2014;141(8):1767-79. doi: 10.1242/dev.103846. PubMed PMID: 24671950; PubMed Central

PMCID: PMC3978837.

58.   Lapierre LR, Hansen M. Lessons from C. elegans: signaling pathways for longevity.

Trends Endocrinol Metab. 2012;23(12):637-44. doi: 10.1016/j.tem.2012.07.007. PubMed PMID:

22939742; PubMed Central PMCID: PMC3502657.

59.   Murphy CT, Hu PJ. Insulin/insulin-like growth factor signaling in C. elegans.

WormBook. 2013:1-43. doi: 10.1895/wormbook.1.164.1. PubMed PMID: 24395814.

60.   Soukas AA, Kane EA, Carr CE, Melo JA, Ruvkun G. Rictor/TORC2 regulates fat

metabolism, feeding, growth, and life span in Caenorhabditis elegans. Genes Dev.

2009;23(4):496-511. Epub 2009/02/26. doi: 23/4/496 [pii]

10.1101/gad.1775409. PubMed PMID: 19240135; PubMed Central PMCID: PMC2648650.

61.   Jones KT, Greer ER, Pearce D, Ashrafi K. Rictor/TORC2 regulates Caenorhabditis

elegans fat storage, body size, and development through sgk-1. PLoS Biol. 2009;7(3):e60. Epub

2009/03/06. doi: 08-PLBI-RA-3327 [pii]

10.1371/journal.pbio.1000060.   PubMed   PMID: 19260765;   PubMed   Central   PMCID:

PMC2650726.

62.   O'Donnell MP, Chao P-H, Kammenga JE, Sengupta P. Rictor/TORC2 mediates gut-to-

brain signaling in the regulation of phenotypic plasticity in C. elegans. PLoS genetics.

2018;14(2):e1007213.

63.   Jordan SD, Konner AC, Bruning JC. Sensing the fuels: glucose and lipid signaling in the

CNS controlling energy homeostasis. Cell Mol Life Sci. 2010;67(19):3255-73. Epub 2010/06/16.

doi: 10.1007/s00018-010-0414-7.   PubMed   PMID: 20549539;   PubMed   Central   PMCID:

PMC2933848.

64.   Stoeckman AK, Ma L, Towle HC. Mlx is the functional heteromeric partner of the

carbohydrate response element-binding protein in glucose regulation of lipogenic enzyme genes.

J Biol Chem. 2004;279(15):15662-9. doi: 10.1074/jbc.M311301200. PubMed PMID: 14742444.

65.   Li MV, Chen W, Harmancey RN, Nuotio-Antar AM, Imamura M, Saha P, et al. Glucose-

6-phosphate mediates activation of the carbohydrate responsive binding protein (ChREBP).

Biochem   Biophys   Res   Commun.   2010;395(3):395-400.   doi: 10.1016/j.bbrc.2010.04.028.

PubMed PMID: 20382127; PubMed Central PMCID: PMC2874883.

66.   Davis KC, Choi Y-I, Kim J, You Y-J. Satiety behavior is regulated by ASI/ASH

reciprocal antagonism. Scientific reports. 2018;8(1):1-7.

67.   Shimizu K, Ashida K, Hotta K, Oka K. Food deprivation changes chemotaxis behavior in

Caenorhabditis elegans. Biophysics and physicobiology. 2019;16:167-72.

68.   Brenner S. The genetics of Caenorhabditis elegans. Genetics. 1974;77(1):71-94. PubMed

PMID: 4366476.

69.   Shioi G, Shoji M, Nakamura M, Ishihara T, Katsura I, Fujisawa H, et al. Mutations

affecting nerve attachment of Caenorhabditis elegans. Genetics. 2001;157(4):1611-22. PubMed

PMID: 11290717; PubMed Central PMCID: PMC1461592.

70.   Okkema PG, Harrison SW, Plunger V, Aryana A, Fire A. Sequence requirements for

myosin gene expression and regulation in Caenorhabditis elegans. Genetics. 1993;135(2):385-

404. PubMed PMID: 8244003; PubMed Central PMCID: PMC1205644.

71.   Warren CE, Krizus A, Dennis JW. Complementary expression patterns of six

nonessential Caenorhabditis elegans core 2/I N-acetylglucosaminyltransferase homologues.

Glycobiology. 2001;11(11):979-88. PubMed PMID: 11744632.

72.   Hilliard MA, Bargmann CI. Wnt signals and frizzled activity orient anterior-posterior

axon outgrowth in C. elegans. Dev Cell. 2006;10(3):379-90. PubMed PMID: 16516840.

73.   Maduro M, Pilgrim D. Identification and cloning of unc-119, a gene expressed in the

Caenorhabditis elegans nervous system. Genetics. 1995;141(3):977-88. PubMed PMID:

8582641.

74.   Egan CR, Chung MA, Allen FL, Heschl MFP, Van Buskirk CL, McGhee JD. A Gut-to-

Pharynx/Tail Switch in Embryonic Expression of the Caenorhabditis elegans ges-1 Gene Centers

on   Two   GATA   Sequences.   Developmental   Biology.   1995;170(2):397-419.   doi:

https://doi.org/10.1006/dbio.1995.1225.

75.   Troemel ER, Chou JH, Dwyer ND, Colbert HA, Bargmann CI. Divergent seven

transmembrane receptors are candidate chemosensory receptors in C. elegans. Cell.

1995;83(2):207-18. PubMed PMID: 7585938.

76.   Mello C, Fire A. DNA transformation. Methods Cell Biol. 1995;48:451-82.

Main Figure Legends

Figure 1: Starvation reduces copper avoidance

A) Schematic of the sensory integration assay. ~100-200 day 1 adult animals (n) are placed in

the black rectangle. Blue barrier represents copper barrier (or other repellant) and star represents

diacetyl or other attractant. Chemotactic Index is the number of animals that have crossed the

barrier (odor side) divided by the total number of animals on the plate (odor + origin sides).

Experiments with well-fed (WF) animals will appear with black dots and those with food-

deprived (FD) animals will be indicated with blue dots. Unless otherwise noted, FD is 3 hours

with no food. Each dot represents a single plate (N) of animals (n).

B) Animals are deprived of food for increasing periods of time. Animals that have been starved

for 3 hours are allowed to recover for 1, 3, or 5 hours on OP50. Well-fed matched partners are

kept on OP50 plates for the entire length of the experiment. Animals are exposed to 50 mM

$CuSO_4$ repellant and 1:500 (0.2%) diacetyl attractant. $N \geq 6$.

C) Animals are exposed to increasing concentrations of other repellants (Fructose, NaCl,

Quinine) with the attractant 0.05% diacetyl (1:2000) in each condition $N \geq 7$.

D) Animals are exposed to decreasing concentrations of diacetyl (0.2%, 0.1% and 0.05%, or

1:200, 1:1000, and 1:2000, respectively) and other volatile attractants 0.1% Benzaldehyde (BZ)

and 0.05% Isoamyl Alcohol (IAA). 50 mM $CuSO_4$ is the repellant in each condition $N \geq 6$.

E) Animals are exposed to $CuSO_4$ in increasing concentrations (5 mM, 25 mM, 50 mM, 100

mM) without any attractant $N \geq 8$.

F) Animals are exposed to diacetyl alone in decreasing concentrations (0.2%, 0.1%, 0.05%). Full

assay (0.2% diacetyl and 50 mM $CuSO_4$) is included as a control N≥7.

G) Animals are exposed to 1:500 diacetyl and increasing concentrations of $CuSO_4$ (5 mM, 25

mM, 50 mM, 100 mM) N≥6.

All graphs are analyzed using a two-way ANOVA, determined to have significant differences

across well-fed and food-deprived conditions. WF/FD comparisons were then performed as

pairwise comparisons within each genotype or treatment as t-tests with Bonferroni corrections

for multiple comparisons. * $p<0.5$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$, ns $p>0.05$. Error

bars are S.D.

Figure 2: Riskier search strategies in starved worms

(A) Worm tracks (n = 32) are plotted for a representative sensory integration assay of well-fed

worms behaving in the presence of 50mM $CuSO_4$ (blue stripe) and 1 μL 0.2% diacetyl (1:500)

(location not shown). Regions of the plate that were not able to be tracked are in gray with the

edge of the plate indicated in black. Tracks are plotted and color coded for time (0 – 45 minutes).

(B) Worm tracks (n = 31) are plotted for a representative sensory integration assay of 3 hour

food-deprived worms. Conditions and plotting the same as in A.

(C, F, I) The mean cumulative sum of worm tracks that cross the barrier as a fraction of the total

worm tracks is plotted at three time points (15, 30, and 45 minutes). Well-fed (WF) animals

appear with black dots and food-deprived (FD) animals are indicated with blue dots. Each dot

represents a single plate of animals. C) 50 mM $CuSO_4$ and 0.2% diacetyl F) 50 mM $CuSO_4$, no

diacetyl I) No copper, 0.2% (1:500) diacetyl. Graphs are analyzed using a two-way ANOVA to

determine significant differences across well-fed and food-deprived conditions. WF/FD

comparisons were then performed as pairwise comparisons within each time period as t-tests

with Bonferroni corrections for multiple comparisons. * $p<0.5$, ** $p<0.01$, *** $p<0.001$, ****

$p<0.0001$, ns $p>0.05$.

(D, G, J) The probability of an animal being located at 1 mm binned distances from the barrier is

plotted for well-fed (black) and food-deprived animals (blue). The dark line represents the mean

probability of residence with the shaded areas representing the standard error of the mean. D) 50

mM $CuSO_4$ and 0.2% diacetyl G) 50 mM $CuSO_4$, no diacetyl J) No copper, 0.2% diacetyl. For

each graph, multiple unpaired t-tests with Welch's correction were performed with correction for

multiple comparisons with Holm-Šídák post-hoc test. Corrected p values <0.05 are indicated by

yellow shading. A comprehensive list of the statistics can be found in Supplementary Table 2.

(E, H, K) The mean velocity of worms as a function of distance from the barrier is plotted for

well-fed (black) and food-deprived animals (blue). Conditions, plotting, and statistics are the

same as in D, G, and J.

Figure 3: Lack of food, not fat or physical interactions, drive behavioral changes

(A) Schematic of Oil Red O experiments. Animals are raised together to day 1 of adulthood and

separated into three groups: well-fed (on food), 3 hour food-deprived, and 6 hour food-deprived.

Animals are stained using Oil Red O and then imaged using a color camera. (B) Representative

images of well-fed (WF, black), 3 hour food-deprived (3hr FD, blue), and 6 hour food-deprived

(6hr FD, green). Inset images are shown, highlighting the regions where there is the most

difference in staining. Black arrows highlight regions of no Oil Red O stain in 6hr FD. (C) Graph

showing the percent change in Oil Red O staining when compared to the average of the area of

Oil Red O signal above a threshold value in the well-fed group within each independent

experiment. N=3, n>20 within each experimental treatment group. (D) Graph showing the

percent of the animals' area that contains Oil Red O signal above threshold N=3, n>20 within

each experimental treatment group. Same data as in C, shown as non-normalized values. (E) A

schematic representing the experiment in F, in which populations of animals are either well-fed

or food-deprived in the presence or absence of Sephadex beads before performing the sensory

integration assay. (F) Prior to the sensory integration assay, animals are exposed to either

standard OP50 ("no beads WF") or empty plates ("no beads FD"), or Sephadex gel beads as

chemosensory input. Alternatively, animals were exposed to beads and no food ("beads FD") or

OP50 with Sephadex beads on top ("beads WF") for 3 hours. Animals were then exposed to

standard Sensory Integration Assay set-up with 50 mM $CuSO_4$ and 1 μL of 0.2% diacetyl. N≥18.

C and D were analyzed using Welch's ANOVA test with Dunnett's multiple comparisons test. *

$p<0.5$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$, ns $p>0.05$. F was analyzed using a full model

two-way ANOVA, determined to have significant differences across well-fed and food-deprived

conditions but no difference between "bead"/"no bead" groups. Those comparisons are shown to

indicate no difference between "beads" and "no beads". Pairwise comparisons within each

treatment were performed as t-tests with Tukey's multiple comparisons test. Error bars are S.D.

Figure 4: *mml-1* and *hlh-30* are required for sensory integration shift upon food

deprivation, correlated with shifts in intestinal localization.

(A) Schematic showing the 20 intestinal cells in a day 1 adult *C. elegans*. Our findings for

MML-1:GFP and HLH-30::GFP are shown in the dotted box, while previously published

paradigms are within the solid line box. Addition of glucose has been shown to induce nuclear

localization of MondoA. Autophagy has been shown to increase nuclear localization of HLH-30.

(B) Standard sensory integration assay with *mml-1(ok849)* and *mxl-2(tm1516* and wildtype

controls. N=20.

(C) Representative images of MML-1::GFP localization in day 1 adult animals (data quantified

in D). All images were collected with the same exposure time and laser power.

(D) Intestinal MML-1::GFP expression in animals during static timepoints food deprivation.

Only intestinal expression was characterized as "nuclear", "nuclear/cytoplasmic", or

"cytoplasmic". Each dot represents the proportion of animals within an experiment with the

phenotype. N=6, n=296.

(E) Standard sensory integration assay with *hlh-30(tm1978)* mutant animals and wildtype

controls. N=9.

(F) Representative images of HLH-30::GFP localization in day 1 adult animals (data quantified

in G). All images were collected with the same exposure time and laser power.

(G) Intestinal HLH-30::GFP expression in animals during static timepoints of food deprivation.

Only intestinal expression was characterized as "nuclear", "nuclear/cytoplasmic", or

"cytoplasmic". Each dot represents the proportion of animals within an experiment with the

phenotype. N=3, n=149.

B and E were analyzed using two-way ANOVA, determined to have significant differences

across well-fed and food-deprived conditions. WF/FD comparisons were then performed as

pairwise comparisons within each genotype or treatment as t-tests with Bonferroni's multiple

comparisons test. D and G were analyzed using Two-Way ANOVA, determined to have

significant differences across localization and an interaction between time of food deprivation

and localization. Within each localization group, pairwise comparisons were performed across

each time point and tested for significance using Tukey's multiple comparisons test. * p<0.5, **

p<0.01, *** p<0.001, **** p<0.0001, ns p>0.05. Error bars are S.D.

Figure 5: Sensory integration changes require HLH-30-regulated insulins and *daf-2* is

required in neurons with *mml-1* in the intestine.

(A) HLH-30 interacts with *C. elegans* insulin peptides. Of the 40 insulin-like peptides encoded in

the *C. elegans* genome, 22% have an HLH-30 binding motif (CANNTG E-box, blue) in the 5'

UTR (< 300bp upstream of start site) [51]. 7% of insulins have been shown to regulate the

localization of HLH-30 but do not contain an E-box (orange, "HLH-30 modifiers"). An

illustration of a representative insulin peptide with two yellow exons and an upstream E-box with

HLH-30 initiating transcription.

(B) All insulins known to contain an HLH-30 binding motif in the 5' UTR were tested using the

standard sensory integration assay. When available, more than one allele was tested (N≥8) for

each insulin, with wild-type (N2) animals tested with each mutant.

(C) Insulins previously shown to regulate HLH-30 localization (*ins-7, ins-8, ins-37*) were tested

using the standard sensory integration assay alongside wildtype (N2) control. N≥7.

(D) *daf-2* mutants and tissue-specific rescues are tested in the standard sensory integration assay

N ≥ 9 for each strain tested alongside wild-type N2. *daf-2* is rescued in neurons, intestines, and

pharynx using tissue-specific promoters.

(E) *daf-2, mml-1,* and *daf-2 mml-1* mutants were tested in standard sensory integration assays.

mml-1 was rescued in a tissue-specific manner in intestinal cells (*Pgly-19)* and *daf-2* was rescued

in the ASI neurons (P*str-3*), alongside N2 controls. N≥10.

(F) Schematic showing requirement of *mml-1* in the intestine, insulin-like peptides, and *daf-2* in

ASI neurons. CI phenotype means Chemotactic Index phenotype, where wildtype animals

displace a chemotactic index of WF < FD.

All graphs were analyzed using a two-way ANOVA, determined to have significant differences

across well-fed and food-deprived conditions. WF/FD comparisons were then performed as

pairwise comparisons within each genotype or treatment as t-tests with Bonferroni's multiple

comparisons test. * $p < 0.5$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, ns $p > 0.05$.

Figure 6: ASI chemosensory neurons use insulin-signaling pathway to integrate intestine-

released peptide signals

(A) Schematic of an ASI neuron's *daf-2*-mediated canonical and non-canonical insulin signaling.

Summary of the findings in B and C.

(B) Standard sensory integration assay performed with mutants in the canonical insulin signaling

pathway (*daf-16, age-1, daf-18, pdk-1, akt-1,* and *akt-2*), alongside wild-type N2 N≥7.

(C) Standard sensory integration assay performed with wildtype N2 (*rict-1 +*), *rict-1* mutants,

and *rict-1* mutants with *rict-1* rescued in ASI (*Pstr-3*). N ≥9.

(D) Summary of data and proposed model through which food deprivation alters animal

behavior.

(B-C) were analyzed using two-way ANOVA, determined to have significant differences across

well-fed and food-deprived conditions. WF/FD comparisons were then performed as pairwise

comparisons within each genotype or treatment as t-tests with Bonferroni's multiple comparisons

test. * $p < 0.5$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, ns $p > 0.05$.

Supplementary Material Legends

Supplementary Figure 1 (to accompany Figure 1): Spread of copper sulfate CuSO₄ on agar plates visualized using 1-(2-pyridylazo)-2- naphthol

(A-E) 25 µl of (A) water as control, (B) 5 mM $CuSO_4$, (C) 25 mM $CuSO_4$, (D) 50 mM $CuSO_4$, and (E) 100 mM CuSO4 was dripped and dried overnight along the midline of the plate to form a copper gradient. PAN indicator (1-(2-pyridylazo)-2-naphthol) distributed over the entire plate shows a gradient of orange-red upon chelation with copper ions. (F) Measured width of colored area with each data point representing the average width, error bars indicate SEM. N = 9.

Supplementary Figure 2 (to accompany Figure 1): Food-deprived animals fail to avoid copper in single animal drop test.

(A) Schematic for dry drop test shown in B. ~300 nL of 1.5 mM CuSO4 is dropped ~1 mm away from the animal's forward motion. Turning away or backing up is considered "avoidance" and given a score of 1. Heading toward the dried drop is considered "no avoidance" and given a score of 0. (B) Quantification of the dry drop test. Food-deprived (FD) animals were starved for 3 hours. Each dot represents the average of ten trials (drops) for a single animal, N=15. Analyzed with an unpaired t-test * $p<0.5$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$, ns $p>0.05$. Error bars are S.D.

Supplementary Figure 3 (to accompany Figure 2): Description of measurements to define

tracking dynamics and additional treatment groups

(A) Measuring Barrier Crossings. Worm tracks (n = 31) are plotted for the first 15 minutes of a

representative sensory integration assay. 188 tracks are plotted in a unique color with the start of

each track labelled with a numbered, circular marker. The number of unique, continuous tracks

that started past the copper barrier was divided by the number of unique tracks in the entire field-

of-view to obtain a measure of barrier crossing. In the example experiment shown, 11 unique

tracks crossed the barrier out of 188 total tracks, resulting in a Barrier Crossings score of 0.0585

for this experiment after 15 minutes of recording.

(B) Measuring Velocity. 10 seconds (i.e. 30 frames) of a single example worm track is plotted.

The midpoint positions of the worm at each frame as identified by WormLab are plotted as filled

circles connected by lines. For each time t, the velocity was calculated by computing the

Euclidean distance of the track from time $t – 1$ second to time $t + 1$ second and dividing by the

length of time, 2 seconds. Because these videos lack the special resolution necessary to

accurately estimate absolute path length (and thus body bends), Euclidean distance is used. In the

example given, the Euclidean distance of the 2 second time window centered at time t was 425.7

μm resulting in an instantaneous velocity of 212.9 μm/s. Velocity was calculated for every time

point in this way.

(C) Measuring Probability of Location. 9 unique worms tracks are plotted in a 3 mm x 3 mm

field-of-view, a 9 mm^2 inset of a 45-minute example experiment. The midpoint positions of the

worms at each frame are plotted as circles connected by lines. Midpoints located in Bin X (1 mm

wide) are represented by filled circles while midpoints located in the neighboring bins (Bin X-1

and Bin X+1, each 1 mm wide) are represented by open circles. The probability of a worm being

located in Bin X is calculated by dividing the number of tracked midpoints in Bin X by the total

number of tracked points in all bins. In the small example area shown, there are 139 points in Bin

X and a total of 345 points across all 3 bins resulting in a p(Location) score of 0.4029. In the

entire field of view there are 45 bins, yielding an average p(Location) score of 0.0222. This

analysis was used in Figure 2D, 2G, 2J.

(D) Graph of the chemotactic index (# animals on odor side / total # of animals) over time (15,

30, 45 minute bins). Well-fed (WF) animals appear with black dots and food-deprived (FD)

animals are indicated with blue dots. Each dot represents a single plate of animals, with each

plate measured at each time point (matched). Analyzed using a Two-Way ANOVA, determined

to have significant differences across well-fed and food-deprived conditions. WF/FD

comparisons were then performed as pairwise comparisons within each time period as t-tests

with Bonferroni's correction for multiple comparisons. * $p < 0.5$, ** $p < 0.01$, *** $p < 0.001$, ****

$p < 0.0001$, ns $p > 0.05$.

(E) Worm tracks (n = 36) are plotted for a representative sensory integration assay of well-fed

worms behaving in the presence of 50 mM $CuSO_4$ in water (blue stripe) with no attractant

(location not shown). Regions of the plate that were not able to be tracked are in gray with the

edge of the plate indicated in black. Tracks are plotted and color coded for time.

(F) Worm tracks (n = 40) are plotted for a representative sensory integration assay of 3 hour

food-deprived worms. Conditions and plotting the same as in E.

(G) Worm tracks (n = 33) are plotted for a representative sensory integration assay of well-fed

worms behaving in the presence of no barrier (blue stripe) with attractant is 1 μL 0.2% diacetyl

(1:500) in 100% ethanol (location not shown). Regions of the plate that were not able to be

tracked are in gray with the edge of the plate indicated in black. Tracks are plotted and color

coded for time.

(H) Worm tracks (n = 31) are plotted for a representative sensory integration assay of 3 hour

food-deprived worms. Conditions and plotting the same as in G.

1024

1025 **Supplementary Table 1: All worm strains used in the experiments.** Strain ID,

1026 genotype/allele, and how it is referenced in the paper is provided. If the strain is first described

1027 here (all IV strains), the method of creation is provided.

1028

1029 **Supplementary Table 2: All p-values for figures 2D, 2G, 2J and 2E, 2H, 2K.** The p-values

1030 shown are the result of multiple unpaired t-tests with Welch's correction with Holm-Šídák post-

1031 hoc tests correction for multiple comparisons. Adjusted p-values are shown, with yellow shading

1032 for adjusted p-values <0.05, same shading as in figures 2D, 2G, 2J and 2E, 2H, 2K.

1033

1034 **Supplemental Movie S1. Sensory integration behavior of well-fed animals.** ~150 Well-fed

1035 wild-type animals are placed in the standard sensory integration assay. Bracket indicates origin

1036 where animals are placed, spot shows position of 1:500 diacetyl odor, midline indicates repellent

1037 $CuSO_4$ barrier.

1038

1039 **Supplemental Movie S2. Sensory integration behavior of food-deprived animals.** ~150

1040 Wild-type animals' food-deprived for three hours are placed in sensory integration behavior

1041 assay. Bracket indicates origin where animals are placed, spot shows position of 1:500 diacetyl

1042 odor, midline indicates $CuSO_4$ barrier.

Figure 1: Starvation reduces copper avoidance

Chemotactic Index = # worms odor half / (# worms origin half + odor half)

Figure 3

A

WF

3 hr FD

6 hr FD

B

WF

3 hr FD

6 hr FD

20 µm

100 µm

C

% change in Oil Red O staining from Well-Fed

****

ns

****

WF   3 hr FD   6 hr FD

D

% Area of worm ORO+

****

ns

****

WF   3 hr FD   6hr FD

E

No beads
(standard assay)

WF   FD

Beads
(+chemosensory input)

WF   FD

F

Chemotactic Index

ns

ns

****

****

No Beads   Beads

● WF
● FD

Figure 4

A

B

C

MML-1::GFP nuclear

MML-1::GFP n/c

MML-1::GFP cytoplasmic

D

- nuclear
- nuclear/cytoplasmic
- cytoplasmic

E

F

hlh-30::GFP nuclear   hlh-30::GFP n/c   hlh-30::GFP cytoplasmic

G

- nuclear
- nuclear/cytoplasmic
- cytoplasmic

Figure 5

mml-1;daf-2   CI phenotype

no rescue   WF ~ FD

Pgly-19::mml-1   WF ~ FD

Pstr-3::daf-2   WF ~ FD

Pgly-19::mml-1
Pstr-3::daf-2   WF < FD

⬠ INS-23
⬛ INS-31
▲ DAF-28
⬠ ?

Figure 6

A

B

A

$$\text{Barrier Crossings} = \frac{\text{\# Unique Tracks Across the Barrier}}{\text{\# Unique Tracks on Plate}}$$

$$e.g. \quad = \frac{11}{188} = 0.0585$$

B

$$\text{Velocity} = \frac{\text{Euclidean Distance (µm)}}{2 \text{ s}}$$

$$e.g. \quad = \frac{425.7 \text{ µm}}{2 \text{ s}} = 212.9 \text{ µm/s}$$

C

$$p(\text{Location}) = \frac{\text{\# Tracked Points in Bin X}}{\text{\# Total Tracked Points in All Bins}}$$

$$e.g. \quad = \frac{139}{345} = 0.4029$$

D 50 mM CuSO₄+ 1:500 Diacetyl

E 50 mM CuSO₄ Only

F 50 mM CuSO₄ Only

G 1:500 Diacetyl Only

H 1:500 Diacetyl Only