Habitat preference of an herbivore shapes the habitat distribution of its host plant

Nicolas M. Alexandre[1,2,3,†], Parris T. Humphrey[2,3,4,†], Joseph Frazier[2], Andrew D. Gloss[2,3,5],

Jimmy Lee[2], Henry A. Affeldt III[2], Noah K. Whiteman[1,2,3*]

[1] Department of Integrative Biology, University of California, Berkeley, CA 94720.

[2] Rocky Mountain Biological Laboratory, Crested Butte, CO 81224.

[3] Department of Ecology & Evolutionary Biology, University of Arizona, Tucson, AZ 85721.

[4] Department of Organismic & Evolutionary Biology, Harvard University, Cambridge, MA

02138.

[5] Department of Ecology & Evolution, University of Chicago, Chicago, IL 60637.

† *equal contribution.*

*Author for correspondence: NKW, whiteman@berkeley.edu, phone: +1 520-696-3950

Running title: Herbivore habitat choice shapes herbivory

Article type: Note

Author contributions: NMA, ADG, JL, HAA, and NKW designed the experiments; NMA,

ADG, JL, HAA, and JF collected the data; PTH and NMA analyzed the data; NMA, PTH and

NKW wrote the paper. All authors contributed to editing the submission.

Data accessibility: All data will be deposited in the Dryad data repository (doi pending).

Abstract

Habitat distributions of plants are often driven by abiotic factors, but growing evidence suggests

an important role for consumers. A textbook example of a plant whose habitat distribution is

shaped by consumers is bittercress (*Cardamine cordifolia*). Bittercress is more abundant in shade

than in sun habitats, and this is thought to arise because herbivore pressure is lower in the shade.

The bittercress case study remains incomplete, as we still do not understand why herbivory is

lower in the shade. Herbivores may avoid shaded bittercress because the plants are lower quality,

or because herbivores simply prefer brighter, warmer habitats. We tested these alternative

hypotheses through a series of herbivore choice experiments. *Scaptomyza nigrita*, a locally

abundant specialist and dominant herbivore of bittercress, strongly preferred feeding and laying

eggs on bittercress we collected from shade versus sun habitats. Thus, shaded bittercress are

more, not less, palatable to these herbivores. Separately, *S. nigrita* strongly preferred feeding and

laying eggs on leaves held in treatments that simulated sun rather than shade habitats—

regardless of whether leaves came from sun or shade habitats originally. The underlying

mechanism for a consumer-driven plant distribution appears to be a simple behavioral preference

of herbivores for brighter, warmer habitats.

Keywords: *Brassicaceae*, bittercress, *Drosophilidae,* leaf miner, *Scaptomyza*, herbivory.

Introduction

Habitat distributions of plants are thought to be shaped primarily by abiotic environmental

gradients (Whittaker 1967), but there is growing evidence that consumers can have a major

impact as well (Maron and Crone 2006). A series of studies in the 1980–90s on bittercress

(Brassicaceae: *Cardamine cordifolia*) in the Elk Mountains of Colorado were among the first to

explore the role of herbivory in shaping the fine-scale distribution of a plant species (Collinge

and Louda 1988, 1990; Louda 1984; Louda and Rodman 1983, 1996), and this system serves as

a textbook example of biotic drivers of plant distributions (Ricklefs and Miller 2000). Adult

females of *Scaptomyza nigrita* flies (Drosophilidae) are the primary herbivores of bittercress and

create feeding punctures in leaves with serrated ovipositors (Collinge and Louda 1988). Their

larvae, which form leaf-mines, can defoliate up to 75% of leaf area from bittercress in sun

habitats (Collinge and Louda 1988). Louda and Rodman (1996) argued that fitness effects of

high herbivory in sun habitats were strong enough to drive bittercress into shade habitats, where

herbivory was low. But nearly 30 years later, we still do not understand the causes of differential

herbivory across this sun–shade ecotone. Louda and Rodman (1996) proposed two mechanisms

to explain this pattern. Plants in sun habitats could be less resistant to herbivores, likely due to

water stress, than those in shade habitats. Herbivores were also more abundant and active in sun

habitats (Louda and Rodman 1996), leading them to propose as an alternative that these

herbivores prefer warmer, brighter habitats (Louda and Rodman 1996). We addressed these

alternative hypotheses in order to better understand the factors that drive a textbook case of

consumer-driven habitat limitation for a native plant.

Because previous herbivory data were collected nearly 30 years ago (Louda 1988), we

first verified that *S. nigrita* herbivory remained higher in sun relative to shade habitats. Second,

we tested the plant quality hypothesis by offering *S. nigrita* females a choice between sun- and

shade-derived bittercress under laboratory conditions. Finally, we conducted choice trials in

laboratory and field settings in which we manipulated the light and temperature environment of

plants within choice trials to measure how these abiotic variables impacted *S. nigrita* foraging

patterns, using plants collected from both sun and shade habitats. We found that *S. nigrita* prefer

bittercress from the shade, yet their strong attraction to plants held in bright and warm habitats is

sufficient to override this preference for shade-derived leaves. Our data lead us to reject the plant

quality hypothesis and instead conclude that the habitat preference of this herbivore for brighter,

warmer habitats underlies the higher rates of herbivory in sun habitats. Thus, the habitat

distribution of bittercress arises from the habitat preferences of its dominant herbivore.

Materials and Methods

Herbivory surveys. All experiments were conducted between 2010 and 2015 at the Rocky

Mountain Biological Laboratory (RMBL) in Gothic, CO, USA. In 2011, we conducted field

surveys of herbivore damage on bittercress in nine sun habitats (no tree canopy) and nine shade

habitats (dense evergreen tree canopy present) (Appendix S1: Fig. S1). Shade and sun sites were

interspersed geographically and in elevation and systematically differed in photosynthetically

active radiation, % shade cover, and nearby canopy tree size (Appendix S1: Table S1). We

recorded adult *S. nigrita* feeding punctures (stipples) and larval mines in two basal leaves from

each of ten ramets from the same bittercress patch (*n*=180 observations per habitat type).

We modeled stipple and mine counts using zero-inflated negative binomial (ZINB)

generalized linear models (GLMs). We chose this model because a zero count can arise because

local herbivore abundance is too low ('false' zeros, Zuur et al. 2009). These are distinct from the

expected proportion of zeros arising from non-truncated count distributions such as Poisson or

NB ('true' zeros, Zuur at al. 2009). While all zeros in our dataset are meaningful, we modeled

whether a leaf belongs to the putatively un-sampled (i.e. 'false zero') class with probability π_0, as

a function of the fixed effects of source habitat (sun vs. shade) and leaf area (mm^2), using the

canonical logit link function in a binomial GLM. The count distribution containing the 'true'

zero class [with probability $(1-\pi_0)$] was simultaneously fit under a NB distribution with a log

link function, with habitat (sun vs. shade), leaf area, and an arbitrary leaf ID (two-levels)

modeled as fixed effects. Coefficients were estimated under maximum likelihood using **R** v3.3.3

(R Core Team 2017) package *pscl* v1.4.9. (Jackman 2015). We further describe our statistical

approach in Appendix S3.

Host choice experiment I: Sun versus shade-derived bittercress. In 2010 we tested

whether *S. nigrita* adult females prefer feeding on individual bittercress derived from sun or

shade habitats. We transplanted bittercress plants from the field into soil within plastic pots and

placed in the laboratory under fluorescent lighting (16:8 light:dark) for < 24 h. In each of eight

replicates, we randomly assigned two shade-derived and two sun-derived bittercress plants to the

four corners of a mesh 35.5 x 35.5 x 61 cm cage (livemonarch.com). All leaves were un-mined,

and we subtracted pre-existing stipple damage from final counts. In each cage we placed two

petri dishes (100 mm diameter) containing 100% recycled paper towels: one moistened with a

5% sucrose solution, and one with tap water. Four, field-collected adult female flies were

introduced into each cage and allowed to feed for 24 h, after which stipples and eggs were

counted using a dissecting microscope. See Appendix S2: Fig. S1A for a schematic.

To control for differences in plant architecture between sun- and shade-derived

bittercress, we conducted a detached leaf assay using cauline leaves clipped from the first or

second position from sun or shade habitats. For each of 15 replicate trials, two leaves each from

sun and shade plants were inserted by their petioles into a half liter-sized plastic container filled

with 1.5 cm of 2% Phytoblend (Caisson Laboratories, Logan, UT). Leaves were randomly

assigned to positions for each assay container, which was closed with a mesh lid. We introduced

one field-caught adult female fly into each container and allowed it to forage for 24 h, after

which we counted stipples and eggs as above. No flies were used for multiple trials. See

Appendix S2: Fig. S1B for a schematic.

For both assays, we modeled stipple and egg counts using NB GLMs with plant habitat

(sun vs. shade), number of cauline leaves (for whole-plants), and leaf width (for detached leaves;

mm) as fixed effects, and cage ID (i.e. replicate assay) as a random intercept. Coefficients were

estimated via maximum likelihood using **R** package *lme4* v1.1-13 (Bates et al. 2015). For

comparisons to Poisson and zero-inflated models see Appendix S3.

Host choice experiment II: Effects of light and temperature. In 2014 and 2015 we

conducted choice experiments to decouple the effects of light and temperature on *S. nigrita*

foraging behavior. In 2014, multiple sets of trials were conducted in a temperature-controlled

laboratory setting as well as in a field setting. In both settings, we manipulated light levels at one

end of large mesh cages (35.5 x 35.5 x 185 cm) and performed choice assays under warmer and

cooler air temperatures. For the field trials, we placed two mesh cages lengthwise on the ground

in adjacent sun and shade plots at a site at RMBL (Appendix S1: Fig. S1). Each cage was placed

under a large wooden picnic table wrapped in reflective Mylar to protect the surface of each

mesh cage from sun exposure. Shade cloth (70% opacity) was then used to wrap each cage until

ambient light was similar to that of shade habitats as determined by a light meter. Within each

cage, we established a light gradient by affixing two LED lights (18–20 lumens, 7000K lights,

LX-8058, Gemini, USA) via hooks at 14 and 21 cm from the bottom of one of two 1.25 cm thick

plywood boards placed vertically against both far sides of each cage. Both plywood boards had a

20 x 30 cm sheet of aluminum foil affixed to them, which focuses the light beam on detached

leaves on the light treatment side. Adult female flies, when placed into the middle of the cage at

the start of the experiments, were not within line of sight of the light source. Two data-loggers

were mounted on each board (Thermocron iButton DS1921G, Maxim, USA) to continuously

measure temperature in the cages. See Appendix S2: Fig. S3 for a schematic.

At two-day intervals, we conducted six trials using both sun-warmed and shade-cooled

cages at once. One side of each cage was randomized to receive the LED light. Ten un-damaged

bittercress plants were collected near RMBL from along the Copper Creek Drainage (Appendix

S1: Fig. S1) and maintained in pots in the laboratory for ≤ 4 d prior to each trial. Four leaves

from each of the ten plants were detached at the petiole and randomized to each of four

experimental conditions (two cage-level temperature treatments X two light environments per

cage). Each group of ten leaves was then placed into petri dishes (100 mm) with petioles

wrapped in a moistened 100% recycled paper towel. The two sides of a cage contained ten petri

dishes, each with a leaf (Appendix S2: Fig S2). Ten *S. nigrita* adult females were collected near

RMBL along the Copper Creek drainage (Appendix S1: Fig. S1) and released into the middle of

each cage. Flies foraged for 24 hours starting at 1100 h.

For the six 2014 laboratory choice trials, the same two cages were placed without Mylar-

table overheads into temperature-controlled environmental chambers. One cage was placed into

each chamber, which was either cooled or held at ambient temperature (~16°C and ~21°C,

respectively; Appendix S4: Fig. S1). Plants, leaves, and flies were collected and utilized as

above, except that flies were allowed to feed for 8 hours (1100–1900 h) during each trial. LED

and data-logger placement in cages were the same for each cage as in the field trials.

We carried out similar trials in 2015 but in a single environmental chamber at two-day

intervals, alternating between two temperatures (approximately 20 °C and 24 °C; Appendix S4:

Fig. S1). Leaves were obtained from plants in sun and shade habitats along the Copper Creek

Drainage near RMBL (Appendix S1: Fig. S1) and were randomized with equal representation of

sun and shade-derived leaves across treatments. Baseline temperatures in 2015 were elevated by

4 °C relative to 2014 (Appendix S4: Fig. S2). In addition to stipples, we counted eggs deposited

by foraging *S. nigrita*, which were not counted in 2014 because our experiment began later in the

season when adult females were less likely to be gravid.

For all 2014 trials, we modeled stipple and egg counts using NB generalized linear mixed

models (GLMMs) with the following fixed effects: leaf width (mm^2), leaf position along stem

from which it was removed ('position'), light environment (light vs. dark), temperature (warm

vs. cool), as well a fixed interaction term between temperature and light environment, which

estimates how the effect of light differs depending on temperature. We modeled between-trial,

between-room, between-cage, and between-side-of-cage effects as a series of nested random

intercept terms. For both years, and for both stipple and egg intensity in 2015, we modeled

counts with NB GLMMs using **R** package *lme4*. For 2015 trials, we included plant source habitat

(sun vs. shade) as a fixed effect. For all of the statistical models, statistical significance of fixed

effects was assessed at the *p*≤0.05 level via asymptotic Wald tests. See Appendix S3 for details.

Results

Herbivory surveys. Bittercress plants in shade habitats had a lower prevalence of stippling (odds

ratio ['OR'] = 0.36 [0.16–0.82 c.i.]) and were less than one tenth as likely to have leaf mines

(OR = 0.08 [0.02–0.42 c.i.]) than bittercress in sun habitats (Table 1; Binomial model). Average

stippling intensity was over four times higher in sun (rate ratio ['RR'] = 4.45 [3.6–5.5 c.i.]), and

leaf miner damage was over 10 times in sun (RR = 11.5 [3.3–39.3 c.i.]; Table 1, 'NB model')

than in shade habitats (Fig. 1A, Table 1). Larger leaves were more likely to be damaged and had

a higher average stippling and mining intensity (Table 1). Leaf size substantially overlapped

between habitats but was slightly larger on average for shade habitats (Appendix S1: Table S2).

Host choice experiment I: Sun versus shade-derived bittercress. In laboratory choice

tests, *S. nigrita* female flies strongly preferred feeding and laying eggs on shade-derived

bittercress. Sun-derived plants received only a third as much stippling (RR = 0.33 [0.22–0.51

c.i.]), and a quarter as many eggs (RR = 0.25 [0.11–0.55 c.i.]) compared to shade-derived plants

(Fig. 1B, Table 1). Results were similar in the detached leaf assay (Appendix S3: Fig. S2, Table

1). Leaf area was not a significant predictor of herbivory in this assay (Table 1).

Host choice experiment II: Effects of light and temperature. In the 2014 field trials,

stippling intensity was eight times higher on plants under lights compared to those not under

lights (RR = 7.9 [2.6–24.5 c.i.]; Fig. 2A, Table 2), and stippling intensity in warmer cages was

three times higher than in cooler cages (RR = 3.1 [1.00–9.54 c.i.]; Fig. 2A, Table 2). We detected

no effect of a light-by-temperature interaction on stippling intensity (Table 2).

S. nigrita exhibited a similarly strong light preference in both 2014 and 2015 laboratory

choice trials. Stippling intensity was six times higher in 2014 (RR = 6.0 [1.16–31.06 c.i.]) and

eight times higher in 2015 (RR = 8.7 [3.2–23.8 c.i.]) on leaves under lights compared to those

not under lights (Fig. 2A, Table 2). No effect of cage temperature was detected for stippling

(Table 2). In all models of stippling intensity, simpler models without an interaction term

between light and temperature were favored via Akaike's Information Criterion (AIC; Appendix

S3: Tables S5–S6). When 2014 and 2015 were pooled, the results were qualitatively unchanged

(Appendix S3: Tables S4,S7).

Egg deposition intensity (measured in 2015 only) was over 30 times higher in leaves

under lights than those not under lights (RR = 36.5 [12.5–104.6 c.i.]) and over seven times

higher in warmer cages (RR = 7.1 [1.99–25.4 c.i.]) than in cooler cages (Fig. 2B, Table 2).

Overall, the model estimated similar egg laying intensities on leaves under lights at both

temperatures (Fig. 2B, Table 2), and the light-by-temperature interaction term reflects that leaves

not under lights received more eggs in the warmed cages than in the cooler cages (Fig. 2B). Leaf

size positively impacted stippling intensity in 2014, and egg intensity in 2015 (Table 2). Plant

source habitat (sun vs. shade) did not significantly impact stippling or egg intensity (Table 2).

Discussion

We found that herbivory was far higher in bittercress found in sun than in shade habitats (Fig.

1A, Table 1), which is the same pattern found nearly 30 years ago (Louda and Rodman 1996). To

address the two alternate hypotheses proposed by Louda and Rodman (1996) to explain this

pattern—one plant-centric (plant quality) and one herbivore-centric (herbivore habitat

preference)—we conducted a series of choice experiments allowing us to dissect the variables

which may be driving this pattern. We found that when given a choice between sun- and shade-

derived bittercress in the laboratory, female flies actually preferred bittercress from the shade

(Fig. 1B, Table 1). Thus, we did not find support for the hypothesis that herbivory is higher in

sun habitats because of higher plant palatability. Given this result, we test the hypothesis that

herbivores prefer brighter, warmer sun habitats. Choice tests in the field in which light levels and

air temperature were experimentally varied within cages allowed us to disentangle the effects of

each variable on herbivore preferences. We consistently found that female flies strongly

preferred foraging on leaves under lights compared to those not under lights when allowed to

move across light environments within cages (Fig. 2A, Table 2). In addition, stippling intensity

was higher in the warmer habitat cages in the field, and egg intensity was higher at warmer

temperatures in 2015 laboratory trials (Fig. 2A, Table 2). The fewest eggs were laid on plants

away from lights in cooler cages (Fig. 2B, Table 2). Thus, warmer temperatures and high light

levels combine to promote herbivory in sun habitats, releasing bittercress in shade habitats from

herbivory.

Lower plant secondary compounds may be one reason *S. nigrita* preferred shade-derived

bittercress in our pairwise tests under constant light (Fig. 2A, Table 2; Humphrey et al. 2016).

But the preference of *S. nigrita* for sun habitats appears stronger than their preference for shade-

derived bittercress. We included leaves sampled from both sun and shade sites in our 2015

habitat choice trials. If host quality were a major driver of habitat specificity of *S. nigrita*

damage, we would expect to have observed higher damage levels on shade-derived than sun-

derived leaves placed under either light treatment. We detected no such effect (Table 2).

The proximal mechanism behind herbivore habitat choice could be a simple positive

phototaxis or thermotaxis. However, phototactic behavior is known to vary among individuals

and strains of *Drosophila melanogaster* and may be under active neuronal control (Gorostiza et

al. 2016). Given that attraction to light is a genetically labile trait yet persists in *S. nigrita,* we

hypothesize that there may be benefits to feeding in warm, sunlit habitats that outweigh any

advantages to feeding on the more palatable plants in the shade. However, it is completely

unknown if the observed preference of *S. nigrita* for sun habitats is adaptive. Habitat preference

could be adaptive due to phenological differences in host plant availability, lower parasitism in

the shade, or extended developmental times of larvae in the shade. Insects, as ectotherms, are

highly sensitive to the temperature of their environment (Sinclair et al. 2012). Cool temperatures

restrict the ability of insects to oviposit on available host plants, even when they are abundant,

because the temperature in such areas is too low for flight (Kingsolver 1989). This may explain

why insects are often restricted to sunny habitats (Huffaker and Kennett 1959; Kaufman 1968),

areas experiencing sunny weather (Whitman 1987), or areas within a plant exposed to the sun,

regardless of plant quality (Casey 1992). It could also be that *S. nigrita* uses visual cues to find

bittercress plants—its only known host—and as a result, has a reduced ability to find bittercress

growing under shade cover (Wallace 1958,Vernon and Gillepsie 1990). Bittercress plants in the

shade are less clumped than those in the sun in distribution and often occur as single plants,

potentially reducing encounters between *S. nigrita* and their hosts (Landa and Rabinowitz 1983,

Finch and Collier 1994). Further studies are required to determine if the light and temperature

preferences of *S. nigrita* are adaptive or represent a constraint that limits the habitat distribution

of this herbivore. Release of shade bittercress from herbivory could also promote divergence in

plant defense strategies and responses to light between bittercress populations in sunlit versus

shaded habitats, especially if they differ in flowering time; these questions await future study.

Our study adds to the evidence that herbivory can have a major impact on fine-scale

habitat distributions of plants. But even in the best-studied systems (e.g., Bruelheide and

Schiedel 1999), the mechanisms driving differential herbivory have been difficult to ascertain.

Here, we found a simple habitat preference of a specialist herbivore species for sunnier, warmer

habitats in the sub-alpine environment of the Rocky Mountains. Regardless of the adaptive value

for the flies, because of the high defoliation potential of larval leaf miners, this herbivore habitat

preference may be sufficient to drive bittercress into the shade.

Acknowledgments

We thank Ian Billick (RMBL), Jennifer Reithel (RMBL), Kailen Mooney (UC-Irvine), Carol

Boggs (University of South Carolina), Nick Waser (RMBL), Mary Price (RMBL), Brian Enquist

(University of Arizona), and Tom Miller (Rice University) for feedback. Financial support was

Literature cited

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models using

lme4. Journal of Statistical Software, 67:1–48.

Bruelheide, H., and U. Scheidel. 1999. Slug herbivory as a limiting factor for the geographical

range of *Arnica montana*. Journal of Ecology 87:839-848.

Casey, T.M. 1993. Effects of temperature on foraging of caterpillars. *Caterpillars: Ecological*

and Evolutionary Constraints on Foraging. Chapman and Hall, New York.

Collinge, S.K., and S.M. Louda. 1988. Patterns of resource use by a drosophilid (Diptera) leaf

miner on a native crucifer. Annals of the Entomological Society of America 81:733-741.

Collinge, S.K., and S.M. Louda 1989. *Scaptomyza nigrita* Wheeler (Diptera: Drosophilidae), a

leaf miner of the native crucifer, *Cardamine cordifolia* A. Gray (Bittercress). Journal of the

Kansas Entomological Society 62:1-10.

Finch, S., and R. Collier. 2000. Host-plant selection by insects–a theory based on

'appropriate/inappropriate landings' by pest insects of cruciferous plants. Entomologia

Experimentalis et Applicata 96:91-102.

Gorostiza, E.A., J. Colomb, and B. Brembs. 2016. A decision underlies phototaxis in an insect.

Open Biology 6:160229.

Huffaker, C.B. and C.E. Kennett. 1959. A ten-year study of vegetational changes associated with

biological control of Klamath weed. Journal of Range Management 12:69–82.

Humphrey, P.T., A.D. Gloss, N.M. Alexandre, M.M. Villalobos, M.R. Fremgen, S.C. Groen,

L.N. Meihls, G. Jander, and N.K. Whiteman. 2016. Aversion and attraction to harmful plant

secondary compounds jointly shape the foraging ecology of a specialist herbivore. Ecology

& Evolution 6:3256-3268.

Jackman, S. 2015. *pscl*: Classes and methods for R developed in the political science

computational laboratory, Stanford University. Department of Political Science, Stanford

University, Stanford, CA. R package version 1.4.9.

Kingsolver, J.G. 1989. Weather and the population dynamics of insects: integrating

physiological and population ecology. Physiological Zoology 2:314.

Landa, K., and D. Rabinowitz. 1983. Relative preference of *Arphia sulphurea* for sparse and

common prairie grasses. Ecology 64:392-395.

Louda, S.M., and J. Rodman. 1983. Ecological patterns in the glucosinolate content of a native

mustard, *Cardamine cordifolia*, in the Rocky Mountains insect-plant interactions, chemical

defense, Colorado. Journal of Chemical Ecology 3:397-422.

Louda, S.M. 1984. Herbivore effect on stature, fruiting, and leaf dynamics of a native crucifer.

Ecology 65:1379-1386.

Louda, S.M., and J.E. Rodman. 1996. Insect herbivory as a major factor in the shade distribution

of a native crucifer (*Cardamine cordifolia* A. Gray, bittercress). Journal of Ecology 84:229-

237.

Louda, S. M., P.M. Dixon, and N.J. Huntly.1987. Herbivory in sun versus shade at a natural

meadow-woodland ecotone in the Rocky Mountains. Plant Ecology 72:141-149.

Louda, S.M. 1990. Predation in the dynamics of seed regeneration. In, *Ecology of Soil Seed*

Banks 124: 105-113. Eds. Leck, M.A., T.V. Parker, and R.L. Simpson, Academic Press, Inc.

Maron, J.L., and E. Crone. 2006. Herbivory: effects on plant abundance, distribution and

population growth. Proceedings of the Royal Society, London, Series B: Biological Sciences

273:2575-2584.

R Core Development Team 2017. R: A language and environment for statistical computing. R

Foundation for Statistical Computing.

Sinclair, B. J., C. M. Williams, and J.S. Terblanche. 2012. Variation in thermal performance

among insect populations. Physiological and Biochemical Zoology: Ecological and

Evolutionary Approaches 6:594-606.

Vernon, R.S., and D.R. Gillespie.1990. Spectral responsiveness of *Frankliniella occidentalis*

(Thysanoptera: Thripidae) determined by trap catches in greenhouses. Environmental

Entomology 19:1229-1241.

Wallace, G.K. 1958. Some experiments on form perception in the nymphs of the desert locust,

Schistocerca gregaria. Journal of Experimental Biology 35:765-775.

Whittaker, R.H. 1967. Gradient analysis of vegetation. Biological Reviews 49:207-264.

Whitman, D.W. 1987. Thermoregulation and daily activity patterns in a black desert

grasshopper, *Taeniopoda eques*. Animal Behavior 35:1814-1826.

Zuur, A., E.N. Ieno, N. Walker, A.A. Saveliev, and G.M. Smith. 2009. Mixed Effects Models

and Extensions in Ecology with R. Statistics for Biology and Health. Springer

Science+Business Media. ISBN 978-0-387-87458-6.

Table 1: Model coefficient estimates for herbivory field survey and lab choice assay of plants from sun and shade habitats.

			Herbivory type	
Dataset	Model type	Coefficient	Stipples	Mines
Herbivory survey	Binomial[†]	Constant	−0.655 (0.486)	1.152 (0.884)
	(π_0)	Site type [sun]	−1.035** (0.424)	−2.491*** (0.782)
		Leaf area (mm^2)	−0.027* (0.015)	–
	NB[‡]	Constant	1.223*** (0.129)	−1.165* (0.648)
	(count)	Site type [sun]	1.494*** (0.107)	2.438*** (0.629)
		Leaf area (mm^2)	0.004* (0.002)	0.009*** (0.003)
		Leaf ID	0.051 (0.098)	−0.332** (0.152)
			Stipples	Eggs
Choice assay	NB[‡]	Constant	1.777*** (0.393)	1.079* (0.476)
(whole plants)	(count)	Leaf position	−0.179*** (0.035)	−0.432*** (0.075)
		Source type [sun]	−1.158** (0.404)	−1.484** (0.470)
Choice assay	NB[‡]	Constant	2.177*** (0.761)	0.856 (0.811)
(detached leaves)	(count)	Leaf area (mm^2)	0.034 (0.048)	0.019 (0.053)
		Source type [sun]	−1.743*** (0.286)	−0.567* (0.304)

*$p < 0.1$; **$p < 0.05$; ***$p < 0.01$

[†]Coefficients are log odds for Constant; log odds ratios for others.

[‡]Coefficients are log rate for Constant; log rate ratios for others.

Table 2: Coefficient estimates for models of habitat choice assays.

Year (setting)	Stipples			Eggs
	2014 (field)	2014 (lab)	2015 (lab)	2015 (lab)
Constant	−2.038***	−2.514***	0.521	−4.870***
	(0.603)	(0.870)	(0.885)	(0.824)
Leaf width (mm)	0.068***	0.063**	−0.003	0.054***
	(0.012)	(0.026)	(0.022)	(0.019)
Plant source [sun]	–	–	−0.291	−0.002
	–	–	(0.702)	(0.19)
Light [light]	2.072***	1.792**	2.145***	3.60***
	(0.574)	(0.838)	(0.240)	(0.540)
Temp. [warm]	1.129**	0.399	−0.125	1.960***
	(0.575)	(0.885)	(0.501)	(0.650)
Light × Temp. [light:warm]	−1.263	1.123	0.325	−1.99***
	(0.788)	(1.172)	(0.870)	(0.630)

$^{*}p < 0.1$; $^{**}p < 0.05$; $^{***}p < 0.01$

Note: Coefficients are log rate for Constant; log rate ratios for all others.

    Figure captions

Fig. 1. Herbivory is higher on bittercress in sun versus shade habitats, but female *S. nigrita*

prefer shade-grown bittercress when given a choice. (**A**) Herbivory field survey results from

sun and shade habitats show higher stipples and mines on bittercress in sun habitats. Raw data

points for both stipples and leaf mines are shown in dark gray and are jittered for visual clarity;

medians are depicted as black bars, while light gray kernel-smoothed density underlay depicts

the distribution of the data. (**B**) Adult female *S. nigrita* stippled and laid more eggs in bittercress

derived from shade versus sun habitats in laboratory choice trials. Plotted are raw leaf-level

counts of stipples and eggs on leaves along bittercress stems. Statistical results are presented in

Table 1.

Fig. 2. Female *S. nigrita* stippled (A) and laid more eggs (B) in bittercress leaves in

simulated sun compared to shade habitats in field and laboratory choice trials. The field

and laboratory choice trials between light and dark sides of assay cages were conducted at two

temperatures (see Appendix S4: Fig. S1 for full temperature profiles), which are indicated below

each sub-plot. Eggs were counted only for laboratory trials conducted in 2015 (see Materials &

Methods). Plot features are depicted as in Fig. 1. Statistical results are presented in Table 2.

Figures

370   **Fig. 1**

373   **Fig. 2**

Appendix S1. Characteristics of source locations for bittercress herbivory survey.

At each site, we recorded leaf area of all sampled leaves, photosynthetically active radiation (PAR) using a light meter (Spectrum Technologies, Inc.), percent canopy cover using a densiometer, diameter at breast height (*dbh*) of the four largest trees within four meters, and latitude, longitude, and elevation using a GPS unit (Garmin) (Table S1). Environmental variables at each site were compared using one-way ANOVAs. Sun habitats had higher average PAR and % open canopy than shade habitats (both $p < 0.001$) and did not systematically differ in elevation ($p > 0.8$, Table S2).

Fig. S1. Map of source sites used in the herbivory surveys in the East River Valley and Copper Creek drainages, near the RMBL in Gothic, CO. A. Base map showing all sites within region (1:48,000). **B–D**. Maps showing detail of site locations (all same scale, 1:7500).

Fig. S2. Distribution of bittercress leaf area observed across all shade and sun habitats.

Table S1. Attributes of sites used for herbivory survey.

Site #	Date sampled	Lat.	Long.	Elevation (m)	Soil Moisture	Light Environment	PAR (μmol•s•m-1)[1]	% Canopy Open	Average DBH (cm)
1	10-Jul-11	39.0050602776535	-106.943317166893	3461	Very Wet	Sun	1980	99.84	0
2	10-Jul-11	38.9885304491745	-106.945774321867	3218	Moist Loamy	Shade	300	6.76	10
3	11-Jul-11	38.9900162385746	-106.944210201638	3240	Moist Loamy	Shade	130	8.06	121.75
4	11-Jul-11	38.9900829890862	-106.943450029132	3253	Very Wet	Sun	2030	82.94	0
5	11-Jul-11	38.98969822732227	-106.9432894133	3261	Very Wet	Sun	2034	95.42	0
6	11-Jul-11	38.9897805369006	-106.94429611421	3244	Moist Loamy	Shade	130	4.16	83.25
7	11-Jul-11	38.9890385630266	-106.944483622637	3220	Very Wet	Sun	2134	99.84	0
8	11-Jul-11	38.9876108211294	-106.944190554921	3250	Moist Loamy	Shade	28	3.38	81.75
9	12-Jul-11	38.9607056556777	-106.973679741745	3023	Very Wet	Sun	2026	88.14	0
10	12-Jul-11	38.9600423351315	-106.973476684125	3013	Wet Loamy	Shade	35	5.72	72.25
11	12-Jul-11	38.9609507737634	-106.973571138078	3994	Very Wet	Sun	1933	86.84	0
12	12-Jul-11	38.9655594715881	-106.968516958823	3994	Very Wet	Shade	32	3.9	101
13	13-Jul-11	38.9828665210329	-106.989822099201	4058	Very Wet	Sun	1780	96.2	0
14	13-Jul-11	38.9774145794955	-106.98986568487	4082	Very Wet	Sun	1870	88.4	0
15	13-Jul-11	38.9771831850899	-106.989697600334	4057	Very Wet	Sun	1850	99.84	0
16	13-Jul-11	38.976356744414	-106.990078474138	4052	Very Wet	Shade	54	4.68	98
17	13-Jul-11	38.975757817399	-106.990327194402	4046	Moist Loamy	Shade	60	4.16	140.5
18	13-Jul-11	38.974195183581	-106.989498527449	4051	Very Wet	Shade	40	7.28	78.75

[1] Light meter was positioned above the center of each source collection plot, at times without cloud cover between 1:00 pm and 3:00 pm during the week of July 17, 2011.

Table S2. Environmental attributes of sun and shade sites where herbivory survey was conducted

Site attributes	Sun Sites (N = 9)		Shade sites (N = 9)		ANOVA	
	μ	se	μ	se	F	P
PAR ($\mu mol{\cdot}s^{-1}{\cdot}m^{-2}$)	1959.7	37.0	89.0	29.4	1564	$<10^{-10}$
% Canopy Open	93.1	2.2	5.3	0.6	1527	$<10^{-10}$
DBH (*cm*)	0.00	0.00	87.5	12.2	51.77	$<10^{-10}$
Elevation (*m*)	3601	146	3568	150	0.025	0.875

Sample attributes	Sun leaves (N = 157)		Shade leaves (N = 140)		ANOVA	
	μ	se	μ	se	F	P
Leaf size (*mm*)	4.98	0.15	5.47	0.13	5.95	0.015

Appendix S2: Design schematics for choice experiments.

A. Whole-plant choice assay.

B. Detached leaf choice assay.

Fig. S1. Schematic of experimental design for Host choice experiment I: Sun versus shade-derived bittercress. (**A**) Whole-plant assay depicted (eight replicate trials were conducted; see Materials & Methods, main text). (**B**) Detached leaf assay (fifteen replicate trials were conducted; see Materials & Methods, main text).

1 meter

Fig. S2. Schematic of experimental design for Host choice experiment II: Effects of light and temperature.

Appendix 3: Statistical Supplement

Contents

1   Herbivory survey

Herbivore damage from *Scaptomyza nigrita* on its host plant, *Cardamine cordifolia* arises from a sequence of contingent events: (i) the arrival by an herbivore, usually a female, at a host plant (or leaf), and (ii) the damage that results from acceptance of such a host by an adult female or a larva. The former process can be modeled as the probability that an herbivore arrives at a plant (prevalence), while the latter captures the probability that the resulting damage is of a particular extent (intensity). We model each of these steps explicitly by considering herbivore damage count data as a mixture of these two processes in a way that is directly compatible with standard generalized linear regression

approaches. Specifically, herbivore damage (in this case, counts of feeding punctures made by adult females ['stipples'], leaf mines made by larvae, and eggs laid by adult females) is always recorded as positive integer counts, and such count data typically exhibit under-dispersion (i.e. 'zero-inflation'), over-dispersion ('excess variance'), or both, with respect to expectations of Poisson or Poisson–Gamma mixture (i.e. negative binomial, 'NB', models.

Our understanding of the foraging ecology of *S. nigrita* leads us to expect that dispersion of both of these types is plausible. Female flies are choosy, often visiting several leaves before making feeding punctures; females may also avoid leaves entirely, either actively or due to stochasticity in the host sampling process, or to differences in local abundances of foraging *S. nigrita*. Once a host has been preliminarily accepted, we expect that variation in the intensity of feeding damage arises from factors perceived subsequent to the initiation of damage. We thus assume that separate (but potentially related) biological processes govern the host acceptance vs. the host damage stages of herbivory as measured in our study.

Figure 1 shows the distribution of stipple and mine counts from our herbivory survey, broken down by habitat type.

1.1 Modeling the host selection process

Practically, our herbivore survey data contain both an excess of zeros as well as an over-dispersed count distribution relative to expectations of a Poisson error model This makes zero-inflated count models a natural choice [10], which are mixture models composed of a binomial component that models the zero-inflation, and a count component (typically Poisson or negative binomial). In our case, a simplest example of the binomial component of such a mixture model assumes that the number of damaged leaves y_i in group i of size n_i, each with probability p_i, is a realization of a binomially distributed random variable Y_i:

$$Y_i \sim B\left(n_i, p_i\right)$$

In the context of a zero-inflated GLM, it is not p_i that is estimated but rather $1 - p_i$, or the probability of zero (i.e. failure), which we define as π_0. Thus, our expression for the binomial probability distribution of y zeros, dropping the i subscript, is

$$Pr\left\{Y(0) = y\right\} = \binom{n}{y}\pi_0^y(1 - \pi)^{n-y} \tag{1}$$

To describe the influence of habitat type and leaf area on $\pi_{0,i}$ we construct a binomial GLM which estimates the linear effects of habitat and leaf area on the logit of $\pi_{0,i}$ (i.e. the log odds), which we call η_i:

$$\text{logit}(\pi_{0,i}) = \log\left(\frac{\pi_{0,i}}{1 - \pi_{0,i}}\right) = \eta_i$$

Figure 1: Distribution of (**A**) stippling and (**B**) leaf miner damage on sun and shade-grown bitter-cress.

Our binomial GLM for stipple abundance takes on the following linear expression for η_i:

$$\eta_i = \begin{cases} \alpha_0 + \beta x_i, & \text{if habitat = shade.} \\ \alpha_0 + \alpha_1 + \beta x_i, & \text{if habitat = sun.} \end{cases} \tag{2}$$

where α_0 is the coefficient for the shade habitat, taken as the reference level (i.e. the 'Constant' or intercept), to which the effect of the sun habitat (α_1) is added; β captures the effect of the area of each leaf (x_i).

In Table 1 of the main text, we report the values of the coefficients α_0 as log odds (the Constant), and α_1, and β as log odds ratios for the factor levels listed in the table. In the Results section of the main text, for ease of interpretation, we report the exponentiated coefficients (e.g. $\exp[\alpha_0]$) to communicate risk of herbivory in terms of odds ($\frac{\pi_i}{1-\pi_i}$) and odds ratios (ORs); we also discuss differences between habitats in terms of the probability of zero counts, π, calculated as:

$$\pi_i = \text{logit}^{-1}(\eta_i) = \frac{e^{\eta_i}}{1 + e^{\eta_i}}$$

Linear estimators ($\hat{\eta}_i$) were generated by maximum likelihood fits of coefficients, as well as the standard error of each coefficient estimate, using **R** [7] package *pscl* [3, 9, 5] using function `zeroinfl`, which jointly estimates these parameters along with those for the count process (see below). Model code in **R** syntax is presented in section 1.3 below.

1.2 Modeling the damage intensity process

We model counts of herbivore damage (stipples and leaf mines) as a Poisson random variable (Y_i) where the Poisson mean (Θ_i) is itself a random variable, which gives rise to the following expression for the conditional probability of Y_i given Θ_i:

$$Pr\{Y_i = y_i | \Theta_i = \theta_i\} = \begin{cases} \pi_{0,i} + e^{-\theta}, & \text{if } y_i = 0. \\ (1 - \pi_{0,i}) \cdot \dfrac{\theta_i^{y_i}}{y_i!} e^{-\theta_i}, & \text{if } y_i > 0. \end{cases} \tag{3}$$

Conceptually, this means that the variance of our random variable Y includes the Poisson variance associated with each Poisson mean Θ as well as additional variance in the distribution of θ itself. We adopt the conventional probability density function for θ as a Gamma distribution with scale parameter α and rate parameter β [10]:

$$g(\theta) = \frac{\alpha^\beta}{\Gamma(\beta)} \theta^{\beta-1} \cdot e^{-\alpha\theta}$$

We recover the the unconditional probability of Y_i by integrating out the θ (i.e. the variable Poisson mean) to recover the parameterization of the negative binomial (NB) that we will use [10]:

$$Pr\{Y_i = y_i\} = \int_0^\infty Pr\{Y_i = y_i | \Theta_i = \theta_i\} \cdot g(\theta) \cdot d\theta$$

$$= \frac{\Gamma(\alpha + y_i)}{y_i! \Gamma(\alpha)} \cdot \left(\frac{\mu_i}{\mu_i + \beta}\right)^{y_i} \left(\frac{\beta}{\beta + \mu_i}\right)^\beta \tag{4}$$

Note that the mean of $g(\theta)$ is $\frac{\alpha}{\beta}$ and its variance is $\frac{\alpha}{\beta^2}$. Setting $\alpha = \beta = r$ allows us to re-write the mean as equal to 1 and the variance σ^2 as $\frac{1}{r}$. This gives the expectation of Y_i equal to μ_i and the variance equal to $\mu + \sigma^2 \mu^2$. Thus, the 'dispersion parameter' of the NB, σ^2 (or alternatively, $\frac{1}{r}$), can be interpreted as the variance of the Gamma distribution from which the Poisson means (θ_i) are drawn. When $\sigma^2 \to 0$ the negative binomial collapses to the Poisson, where $E[Y_i] = \text{Var}[Y_i] = \mu_i$. Note that the μ_i of the NB is not equivalent to the $\frac{\alpha}{\beta}$ from the Gamma distribution.

When combined with the binomial probability of zero (π_i), the full expression for the unconditional probability of Y_i becomes (using the $\sigma^2 = \frac{1}{r}$ parameterization to be consistent with **R**):

$$Pr\{Y_i = y_i\} = \begin{cases} \pi_{0,i} + (1 - \pi_{0,i}) \cdot \left(\dfrac{r}{r + \mu_i}\right)^r, & \text{if } y_i = 0. \\[2ex] (1 - \pi_{0,i}) \cdot \dfrac{\Gamma(r + y_i)}{y_i! \Gamma(r)} \cdot \left(\dfrac{\mu_i}{\mu_i + r}\right)^{y_i} \left(\dfrac{r}{r + \mu_i}\right)^r, & \text{if } y_i > 0. \end{cases} \tag{5}$$

Thus, $Pr\{Y_i\}$ depends on the three parameters, π_0, μ, and r. In our zero-inflated negative binomial (ZINB) GLM framework, the mean of the NB μ_i for each group i is modeled, via the log link function, as a linear function of our predictor variables. Thus, we model $\log(\mu_i)$ as a function of bittercress habitat, leaf area, as well as an additional 'structural' fixed effect, the arbitrary leaf ID (levels A or B):

$$\log(\mu_i) = \begin{cases} \alpha_0 + \beta x_i + \gamma_i, & \text{if habitat = shade.} \\ \alpha_0 + \alpha_1 + \beta x_i + \gamma_i, & \text{if habitat = sun.} \end{cases} \tag{6}$$

Here, α_1 indicates the fixed effect of the sun habitat type; γ_i represents leaf ID, which has two levels (arbitrarily 'A' and 'B'). We set γ_1 to 0 because the reference level of this factor is embedded in the Constant (α_0); β captures the effect of leaf position along the stem (x_i). In the main text, we report model coefficient estimates corresponding to each level of α_i and the single β term for the stipple model. The residual error term is implied.

1.3 Model comparisons

To justify our use of a ZINB model for the herbivory survey data, we constructed ZI-Poisson models and the non-ZI versions of both NB and Poisson models and compared model fits. For stipple data,

all GLMs contained the same terms (Eq. 2, both with logit link functions; Eq. 6 for count models, both with log link function). The same was true for the leaf mine models, but we removed the leaf area term (β) from the binomial component because its inclusion increased overall model AIC by 1.95. In contrast, for the ZINB stipple models, adding leaf area to the binomial component reduced AIC by > 2.5 points and was thus retained. In table S2 below and in Table 1 in the main text, we report the leaf mine model results without the leaf area term.

We compared non-nested models (Poisson, NB-only, ZIP, ZINB) using an analog of the likelihood ratio test (Vuong test) as implemented in **R** package `pscl` [5]. This test calculates $p_i = \hat{Pr}(y_i|M_1)$, the predicted probabilities of each data point from model 1, evaluated under the maximum likelihood estimates of the coefficients, as well as q_i, the corresponding probabilities from model 2. The Vuong (Z) statistic is $Z = \sqrt{N}\bar{m}/\sigma_m$ where $m_i = log(p_i) - log(q_i)$ and σ_m is the sample standard deviation of m_i. P-value of the Vuong test is taken as the $Z < \alpha$, $Z > (1 - \alpha)$ quantile of the standard Normal distribution; here, and throughout this study, we use $\alpha = 0.05$. We report a version of Z which scales $\bar{m}$ according to an AICc finite sample size correction [5]. Tables 1 and 2 show coefficient estimates for stipples and leaf mines (respectively) for each of the four candidate models, as well as their log likelihoods and AIC scores.

Table S1: GLM comparisons for **stipple** counts from herbivory survey.

Model type	Coefficient	Poisson	NB	ZIP	ZINB
Count model (μ_i)	Constant (α_0)	1.00***	0.92***	1.36***	1.22***
		(0.06)	(0.14)	(0.06)	(0.13)
	Leaf area (β)	0.005***	0.007*	0.003***	0.004
		(0.001)	(0.003)	(0.001)	(0.002)
	Habitat [sun] (α_1)	1.60***	1.63***	1.39***	1.49***
		(0.05)	(0.12)	(0.05)	(0.11)
Binomial model (π_i)	Constant (α_0)			−0.37	−0.66
				(0.36)	(0.49)
	Leaf area (mm^2) (β)			−0.02*	−0.03
				(0.01)	(0.02)
	Habitat [sun] (α_1)			−1.35***	−1.04*
				(0.35)	(0.42)
	AIC	2999.90	1847.35	2490.04	1820.66
	Log Likelihood	-1495.95	-918.68	-1238.02	-902.33

***$p < 0.001$, **$p < 0.01$, *$p < 0.05$

Notice that the coefficient estimates are largely similar for both ZINB and NB-only models, while both Poisson family models are far worse fits to the data, indicated by the very large ΔAIC between these and the NB models (Tables S1-S2). Overall, the ZINB model has the lowest AIC (ΔAIC > −20 between ZINB and NB-only models). Comparing the ZINB model to the NB-only model gave an AIC-corrected Vuong statistic of $Z = 2.45$ ($p < 0.01$) for stipples and $Z = 1.41$ ($p = 0.07$) for leaf mines; Z values were well above 5 when ZINB models were compared to ZIP or Poisson models (all $p < 0.001$). Full **R** code for all figures, models, and calculations can be found in the Dryad data repository (doi pending).

Table S2: GLM comparisons for **larval leaf mine** counts from herbivory survey.

Model type	Coefficient	Poisson	NB	ZIP	ZINB
		\multicolumn{4}{c}{Model}			
Count model (μ_i)	Constant (α_0)	-2.60^{***}	-2.65^{***}	-0.82	-1.19
		(0.30)	(0.34)	(0.56)	(0.67)
	Leaf area (mm^2) (β)	0.006^{***}	0.009^{*}	0.008^{***}	0.009^{**}
		(0.002)	(0.004)	(0.002)	(0.0003)
	Habitat [sun] (α_1)	3.61^{***}	3.64^{***}	2.19^{***}	2.46^{***}
		(0.29)	(0.32)	(0.56)	(0.65)
Binomial model (π_i)	Constant (α_0)			1.65^{*}	1.24
				(0.71)	(0.75)
	Habitat [sun] (α_1)			-2.57^{***}	-2.44^{**}
				(0.67)	(0.63)
	AIC	1003.99	802.94	839.12	792.92
	Log Likelihood	-497.99	-396.47	-412.56	-389.46

$^{***}p < 0.001$, $^{**}p < 0.01$, $^{*}p < 0.05$

2 Herbivore choice tests I: Sun versus shade derived bittercress

2.1 Models for choice experiments

Using NB generalized linear mixed models (GLMMs), we model variation in stipple and egg counts on plants arising from source habitat (sun versus shade) and leaf attributes (as fixed effects), as well as 'structural' random effects of plant ID nested within cage ID (i.e. replicate) to capture experimental design constraints that determine the level of independence among datapoints. Our mixed model for stipple and eggs counts, for both the whole-plant and detached leaf assays, takes the following form:

$$\log(\mu_{ijk}) = \begin{cases} \alpha_0 + \beta x_i + \gamma_{(jk)} + \gamma_k, & \text{if habitat = shade.} \\ \alpha_0 + \alpha_1 + \beta x_i + \gamma_{(jk)} + \gamma_k, & \text{if habitat = sun.} \end{cases} \tag{7}$$

In this model, μ_{ijk} is the estimate for each leaf i, α_0 and α_1 are as in Eqn. 6, and β is leaf position along stem (low to high) for the whole-plant model or leaf area (mm^2) for the detached leaf assay; finally, γ_k represents a random effect of cage ID (k) and $\gamma_{(jk)}$ represents a random effect for plant ID j nested within each level of cage ID k; $\gamma_j \sim N(0, \sigma_1^2)$ and $\gamma_{(jk)} \sim N(0, \sigma_2^2)$.

In our assay cages, we assume that adult females *S. nigrita* flies had sufficient time to potentially visit all available plant tissue; thus, *a priori* we favor using NB-only compared to ZINB models; we do not consider Poisson models further in this analysis on the basis of far worse model fits observed for stipple and leaf mine data, above. We directly compared NB-only models to a ZINB version to justify this approach. We found that adding a ZI term did not improved model fit for stipples ($\Delta AIC = 1.24$) nor for eggs ($\Delta AIC = 1.94$). For the detached leaf assay, the ZINB model was also

a marginally worse fit for both stipples ($\Delta AIC = 2$) and eggs ($\Delta AIC = 1.77$) as well (all ΔAIC are ZINB $-$ NB; Table S3). *P*-values for Vuong tests between NB and ZINB models were all > 0.1.

ZINB and NB mixed models were fit using **R** package `glmmADMB` with the NB parameterization described in section 1 [2, 8]. `glmmADMB` handles ZI mixed models by fitting a single constant π_0 term across all groups, rather than a $\pi_{0,i}$ for each designated group, making it less flexible than the ZINB implementation in package `pscl`; however, `glmmADMB` handles NB and ZINB mixed models, making it appropriate for fitting models with random effects. Coefficient estimates for NB and ZINB models for stipple and egg counts were very similar for both assay types (Table S3); in the main text, we report the NB-only model results.

Table S3: Model estimates for plant source choice experiments.

	stipples		eggs	
Whole-plant assay	NB	ZINB	NB	ZINB
Constant (α_0)	1.777 (0.393)***	1.927 (0.412)***	1.079 (0.476)*	1.285 (0.802)
Leaf position (β)	−0.179 (0.035)***	−0.184 (0.036)***	−0.432 (0.075)***	−0.430 (0.075)***
Plant source (α_1)	−1.158 (0.404)**	−1.127 (0.406)**	−1.484 (0.470)**	−1.469 (0.469)**
Cage (σ_1^2)	0.323	0.331	0.0004	0
Plant:Cage (σ_2^2)	0.919	0.938	0.473	0.466
AIC	1143.024	1144.268	386.976	388.920
Log Likelihood	−565.512	−565.134	−187.488	−187.460
Num. obs.	356	356	356	356
Zero inflation: parameter		0.138		0.200
Zero inflation: SD		0.121		0.589
Detached leaf assay				
Constant (α_0)	2.177 (0.761)**	2.273 (0.770)**	0.856 (0.811)	1.433 (0.718)*
Leaf position (β)	0.034 (0.048)	0.034 (0.049)	0.019 (0.053)	0.009 (0.047)
Plant source (α_1)	−1.743 (0.286)***	−1.770 (0.291)***	−0.567 (0.304)	−0.368 (0.305)
Cage (σ_1^2)	1.35	1.135	0.561	0
AIC	395.818	397.554	280.105	281.934
Log Likelihood	−192.909	−192.777	−135.052	−134.967
Num. obs.	60	60	60	60
Num. groups: cage	15	15	15	15
Zero inflation: parameter		0.000		0.217
Zero inflation: SD		0.000		0.103

***$p < 0.001$, **$p < 0.01$, *$p < 0.05$

3 Herbivore choice tests II: Effects of light and temperature.

The NB model structures for this set of choice experiments were similar to those described in above except for an expanded random effects structure. All models described below are NB-only GLMMs; ZINB models will not be evaluated further in this section. The choice trials in 2014 and 2015 were

conducted slightly differently, which means that each year's data calls for slightly different random effects. We present the analyses of each dataset separately, then jointly.

3.1   2014 Trials

The 2014 field and lab trials were conducted in two temperature environments simultaneously, with one cage held in each. This gives a structure of temperature environment ($\gamma_{(lk)}$, $n = 2$) nested within trial (γ_l, $n = 6$ each for the 2014 field and lab assays). Nested within temperature environment is cage, but since we have only a single level of cage for each temperature environment per trial, this level is irrelevant for the 2014 dataset. However, we include side-of-cage ($\gamma_{(kj)}$, $n = 2$; left or right, arbitrarily) to control for pseudo-replication at the level of the main treatment effect (i.e. Light environment, light versus dark), which was applied with randomization to the sides of each cage. Each random effect is $\sim N(0, \sigma_i^2)$. The full model is given below using the above notation for the random effects. The number of independent data points in the 2014 field and lab trials is 24 each (2 sides per cage, 1 cage per trial, 2 temperature settings per trial, and 6 trials).

$$
\log(\mu_{ijkl}) = \begin{cases} \alpha_0 + \alpha_{1,i} + \alpha_{2,i} + \alpha_{12} + \beta x_i + \gamma_l + \gamma_{(lk)} + \gamma_{(kj)}, & \text{if Light = light \& Temp = warm} \\ \alpha_0 + \alpha_{1,i} + \alpha_{2,i} + \beta x_i + + \gamma_l + \gamma_{(lk)} + \gamma_{(kj)}, & \text{otherwise.} \end{cases} \tag{8}
$$

We set the first levels of coefficients for the fixed effects of Light ($\alpha_{1,1}$ = dark) and Temperature ($\alpha_{2,1}$ = cool) equal to zero since these two states are incorporated into the Constant (α_0, i.e. the reference level of the model). Our experiment was designed to allow us to fit an interaction term α_{12} which estimates how much the effect of Light is impacted by the Temperature environment (thus α_{12} has a single level). NB models were estimated using **R** package *lme4* [1]. We fit models with the same structure for field and laboratory trials.

3.2   2015 Trials

In the 2015 trials, we used bittercress leaves collected from both sun and shade habitats, and our randomization scheme ensured equal representation of sun- and shade-derived leaves across all treatments. We did this to test whether *S. nigrita* would exhibit preference for shade-derived bittercress in the context of our temperature and light manipulations. In our analysis, we included leaf source (sun v. shade) as an additional fixed factor. Our model structure was thus

$$
\log(\mu_{ijkl}) = \begin{cases} \alpha_0 + \alpha_{1,i} + \alpha_{2,i} + \alpha_{12} + \alpha_{3,i} + \beta x_i + \gamma_l + \gamma_{(lk)} + \gamma_{(kj)}, & \text{if Light = light \& Temp = warm} \\ \alpha_0 + \alpha_{1,i} + \alpha_{2,i} + \alpha_{3,i} + \beta x_i + \gamma_l + \gamma_{(lk)} + \gamma_{(kj)}, & \text{otherwise.} \end{cases}
$$

$$\tag{9}$$

All coefficients are as in the 2014 trials above, except that the term $\alpha_{3,i}$ is for source habitat, and the first level ($\alpha_{3,1}$ = shade) was set to 0 and was thus incorporated into the intercept term α_0. Thus, this model estimates only a single coefficient for the effect of habitat.

The experimental procedures also differed slightly in 2015. We placed two cages in one environmental chamber which was set to a single temperature per trial, and individual trials were conducted at different temperatures sequentially. This implies a structure of cage (γ_{kl}, $n = 2$) nested within trial (γ_l, $n = 5$), along with the side-of-cage nested within cage ($\gamma_{(kj)}$, $n = 2$). The number of independent data points in the 2015 trials is thus 40 (2 sides per cage, 2 cages per room, 10 trials [5 in each temperature regime]). While the model structure is the same as in Eqn. 8, the meaning of the random effects (and their coding in the design matrix) are slightly different. Below we discuss how we reconciled the random effects to analyze both years' data together.

3.3 Combined 2014-2015 Analysis

We reconcile the slight distinctions between 2014 and 2015 datasets by including trial-year as a composite random effect, now re-coded to reflect each experiment conducted in a given room (compared to a trial containing two separate temperature settings, as in the 2014-only analysis). Nested within the new trial factor is cage, which has $n = 1$ for 2014 and $n = 2$ for 2015; side-of-cage is modeled in the same way as above. This model structure is now identical to Eqn. 8.

For the combined analysis, we drop the term for plant source habitat (α_3) since the 2014 trials were not designed to examine plant source habtat. Additionally, we model both a continuous and discrete versions of the Temperature factor: the discrete model is the same as Eqn. 8, while in the continuous form, $\alpha_{2,j}$ is replaced with $\beta_2 y_j$, where y_j is the temperature measured at the level of cage for each trial separately (Appendix B, Fig. 2C). Additionally, the fixed interaction term α_{12} is now replaced with an interaction modeled by the expression $\beta_3 y_j(\alpha_{1,j})$, which captures how the effect of light ($\alpha_{1,j}$) changes as a function of cage temp (y_i); since $\alpha_{1,i}$ has one level (the first level is set to 0), estimating this interaction term β_3 adds only a single parameter to the model. The full model is thus:

$$\log(\mu_{ijkl}) = \alpha_0 + \alpha_{1,i} + \alpha_{2,i} + \beta_1 x_i + (\beta_2 + \beta_3 \alpha_{1,i})y_j + \gamma_l + \gamma_{(lk)} + \gamma_{(kj)} \tag{10}$$

Because the full model with the continuous temperature interaction term (β_3) exhibited such a poor fit to the data as judged by AIC (see below), we report the coefficient estimates for the nested model containing only the additive effect of the continuous temparture term (β_2; Table S4).

3.4 Model Selection

In the main text, we report the results from the NB GLMMs fit to the 2014 and 2015 datasets (separately) using **R** package `lme4`. The full results for the combined analysis is displayed below (Table S4). For the 2014 and 2015 datasets, we also analyzed a series of nested models to evaluate the relative performance of simpler nested models (Table S5-S7). We calculated AIC and also performed likelihood ratio tests (not shown, but can be calculated from reported log likelihoods), which agreed with ΔAIC results. Additionally, we evaluated overall model fit for the fixed effects-only portion with $R^{2(M)}_{GLMM}$ and for all model terms combined with $R^{2(C)}_{GLMM}$ using which were calculated as derived in [6] using **R** package `piecewiseSEM` [4]. The models reported in the main text are not typically the best model as judged by AIC. Nonetheless, we report the more complex models in the main text

Table S4: Coefficient estimates for all light–temp choice experiments.

	Stipples					Eggs
	Field (2014)	Lab (2014)	Lab (2015)	Both years (1)	Both years (2)	Lab (2015)
Fixed effects						
α_0 (Constant)	−2.038***	−2.514**	0.364	−1.147	−4.123	−4.870***
	(0.603)	(0.870)	(0.885)	(0.661)	(2.162)	(0.812)
β_1 leaf width (*mm*)	0.068***	0.063*	−0.003	0.033	0.031	0.054**
	(0.012)	(0.026)	(0.022)	(0.018)	(0.017)	(0.019)
α_1, [light]		1.792*	2.160***	2.019***	2.321***	3.598***
		(0.838)	(0.515)	(0.470)	(0.339)	(0.537)
α_2 [warm]		0.399	−0.100	0.109		1.960**
		(0.885)	(0.880)	(0.667)		(0.650)
α_{12} [light:warm]		1.123	0.293	0.636		−1.994**
		(1.172)	(0.718)	(0.654)		(0.628)
α_3 [sun]			−0.291			−0.002
			(0.701)			(0.19)
β_2 [temp]					0.149	
					(0.102)	
Random effects						
σ^2_{kj} [trial/cage/side]	0.559	1.397	0.720	1.102	1.129	0.152
σ^2_{lk} [trial/cage]	0.000	0.099	2.451	1.687	1.587	0.170
σ^2_{l} [trial]	0.204	0.000	0.000	0.000	0.063	0.163
σ^2_{m} [year]				0.036	0.000	
AIC	1152.4	874.4	2125.7	2995.8	2993.4	861.2
Log Likelihood	−567.2	−428.2	−1053.9	−1487.9	−1487.7	−421.6
Num. obs.	240	240	398	638	638	398

***$p < 0.001$, **$p < 0.01$, *$p < 0.05$

so that the coefficient estimates for all terms are available to the reader; the terms retained in the models with lowest AIC in Tables S5-S7 correspond to those with statistically significant ($p < 0.05$) coefficient estimates in the main text tables.

Table S5: Model comparisons for different fixed effect combinations: 2014.

Dataset	Fixed effects	$R^{2(M)}_{GLMM}$	$R^{2(C)}_{GLMM}$	AIC	ΔAIC	log Lik
2014 Field	α_0	0.00	0.85	1190.05	37.98	-590.03
(stipples)	α_0, β	0.31	0.91	1158.50	6.43	-573.25
	$\alpha_0, \beta, \alpha_1$	0.52	0.91	1152.07	**0.00**	-569.03
	$\alpha_0, \beta, \alpha_1, \alpha_2$	0.54	0.91	1152.81	0.75	-568.41
$\longrightarrow$	$\alpha_0, \beta, \alpha_1, \alpha_2, \alpha_{12}$	0.58	0.90	1152.41	0.34	-567.21
2014 Lab	α_0	0.00	0.86	888.57	15.23	-439.29
(stipples)	α_0, β	0.06	0.87	883.03	9.69	-435.52
	$\alpha_0, \beta, \alpha_1$	0.41	0.86	873.84	0.50	-429.92
	$\alpha_0, \beta, \alpha_2$	0.12	0.87	883.41	10.08	-434.71
	$\alpha_0, \beta, \alpha_1, \alpha_2$	0.47	0.85	873.34	**0.00**	-428.67
$\longrightarrow$	$\alpha_0, \beta, \alpha_1, \alpha_2, \alpha_{12}$	0.48	0.85	874.43	1.09	-428.22

Notes: ΔAIC is each model minus model with lowest AIC.
                       $\longrightarrow$ indicates model reported in main text, Table 3.

Table S6: Model comparisons for different fixed effect combinations: 2015.

Dataset	Fixed effects	$R^{2(M)}_{GLMM}$	$R^{2(C)}_{GLMM}$	AIC	ΔAIC	log Lik
2015 Lab	α_0	0.00	0.95	2140.95	19.06	-1065.48
(stipples)	α_0, β	0.00	0.95	2142.79	20.90	-1065.40
	$\alpha_0, \beta, \alpha_1$	0.28	0.95	2121.89	**0.00**	-1053.94
	$\alpha_0, \beta, \alpha_2$	0.00	0.95	2144.78	22.89	-1065.39
	$\alpha_0, \beta, \alpha_1, \alpha_2$	0.28	0.95	2123.88	1.99	-1053.94
	$\alpha_0, \beta, \alpha_1, \alpha_2, \alpha_{12}$	0.29	0.95	2124.47	2.59	-1053.24
	$\alpha_0, \beta, \alpha_1, \alpha_2, \alpha_3$	0.29	0.95	2126.26	4.37	-1053.13
$\longrightarrow$	$\alpha_0, \beta, \alpha_1, \alpha_2, \alpha_{12}, \alpha_3$	0.28	0.95	2125.72	3.83	-1053.86
(eggs)	α_0	0.00	0.64	908.39	47.22	-449.19
	α_0, β	0.02	0.64	902.03	40.86	-445.01
	$\alpha_0, \beta, \alpha_1$	0.43	0.64	867.95	6.78	-426.97
	$\alpha_0, \beta, \alpha_2$	0.05	0.65	902.43	41.26	-444.21
	$\alpha_0, \beta, \alpha_1, \alpha_2$	0.46	0.65	867.88	6.71	-425.94
	$\alpha_0, \beta, \alpha_1, \alpha_2, \alpha_{12}$	0.56	0.68	861.17	**0.00**	-421.58
	$\alpha_0, \beta, \alpha_1, \alpha_2, \alpha_3$	0.46	0.65	869.87	8.71	-425.94
$\longrightarrow$	$\alpha_0, \beta, \alpha_1, \alpha_2, \alpha_{12}, \alpha_3$	0.56	0.68	863.17	2.00	-421.58

Notes: ΔAIC is each model minus model with lowest AIC.
                       $\longrightarrow$ indicates model reported in main text, Table 3.

Table S7: Model comparisons for different fixed effect combinations: 2014–2015 combined

Dataset	Fixed effects	$R^{2(M)}_{GLMM}$	$R^{2(C)}_{GLMM}$	AIC	ΔAIC	log Lik
Combined Lab	α_0	0.00	0.93	3025.93	32.55	-1506.97
(Stipples)	α_0, β_1	0.02	0.93	3022.43	29.04	-1504.21
	$\alpha_0, \beta_1, \alpha_1$	0.31	0.92	2993.43	0.04	-1488.71
	$\alpha_0, \beta_1, \alpha_2$	0.02	0.93	3023.84	30.45	-1503.92
	$\alpha_0, \beta_1, \alpha_1, \alpha_2$	0.32	0.93	2994.78	1.40	-1488.39
	$\alpha_0, \beta_1, \alpha_1, \alpha_2, \alpha_{12}$	0.32	0.93	2995.84	2.46	-1487.92
	$\alpha_0, \beta_1, \alpha_1, \beta_2$	0.35	0.92	2993.38	**0.00**	-1487.69
	$\alpha_0, \beta_1, \alpha_1, \beta_2, \beta_3$	0.35	0.92	2994.27	0.89	-1487.14

Notes: ΔAIC is each model minus model with lowest AIC.

References

[1] Douglas Bates, Martin Mächler, Ben Bolker, and Steve Walker. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1):1–48, 2015.

[2] D. A. Fournier, H. J. Skaug, J. Ancheta, J. Ianelli, A. Magnusson, M.N. Maunder, A. Nielsen, and J. Sibert. AD Model Builder: using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. *Optim. Methods Softw.*, 27:233–249, 2012.

[3] Simon Jackman. *pscl: Classes and Methods for R Developed in the Political Science Computational Laboratory, Stanford University*. Department of Political Science, Stanford University, Stanford, California, 2015.

[4] Jonathan S. Lefcheck. piecewisesem: Piecewise structural equation modelling in r for ecology, evolution, and systematics. *Methods in Ecology and Evolution*, 7(5):573–579, 2016.

[5] Tom Loeys, Beatrijs Moerkerke, Olivia De Smet, and Ann Buysse. The analysis of zero-inflated count data: beyond zero-inflated poisson regression. *Br J Math Stat Psychol*, 65(1):163–80, Feb 2012.

[6] Shinichi Nakagawa and Holger Schielzeth. A general and simple method for obtaining r2 from generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 4(2):133–142, 2013.

[7] R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2017.

[8] Hans Skaug, Dave Fournier, Ben Bolker, Arni Magnusson, and Anders Nielsen. *Generalized Linear Mixed Models using 'AD Model Builder'*, 2016-01-19.

[9] Achim Zeileis, Christian Kleiber, and Simon Jackman. Regression models for count data in r. *Journal of Statistical Software, Articles*, 27(8):1–25, 2008.

[10] A. Zuur, E.N. Ieno, N. Walker, A.A. Saveliev, and G.M. Smith. *Mixed Effects Models and Extensions in Ecology with R*. Statistics for Biology and Health. Springer New York, 2009.

Appendix S4. Temperature profiles for field and laboratory habitat preference choice tests.

Fig. S1. Temperature profiles for field and laboratory habitat preference choice tests. (**A**) Temperature profiles for field cages (from 2014). Both 'Sun' and 'Shade' cages were equally masked from natural sunlight but were either sun-exposed or canopy shaded in order to confer different temperature profiles. We collected a full 24 h of temperature data during each trial (ordered in time and labeled A–F), which took place for 24 h beginning at 1100 h. 0 h represents midnight. (**B**) Differences in temperature between sun-exposed and canopy-shaded assay cages (sun − shade), showing a maximal difference of 5 °C in mid-afternoon during each trial. (**C**) Average temperature for each laboratory trial for 2014 and 2015 (left; ordered in time and labeled 1–6), and the mean (right; ± 1 standard deviation) over all trials for each year. 2014 trials used two environmental chambers, while 2015 trials alternated between warm and cool trials at two-day intervals (see Methods in the main text for details).