Evolution with a seed bank: the population genetic consequences of microbial dormancy

WR Shoemaker and JT Lennon

Indiana University, Department of Biology

corresponding authors: wrshoema@indiana.edu and lennonj@indiana.edu

ABSTRACT

Dormancy is a common bet-hedging strategy that allows organisms to persist through

26   biotic and abiotic conditions that are sub-optimal for growth and reproduction by reducing their

metabolic activity. The ability to enter and exit a dormant state allows a population to maintain a

28   reservoir of genetic and phenotypic diversity that can contribute to the long-term survival of the

population (i.e., a seed bank), a potentially adaptive strategy has long been of interest to

30   ecologists and evolutionary biologists. However, we know comparatively little about how

dormancy influences the fundamental evolutionary forces of genetic drift, mutation, selection,

32   recombination, and gene flow. Here, we discuss how seed banks affect the fundamental forces

that underpin evolution by reviewing existing population genetic models, implementing novel

34   simulations, and determining how and when dormancy can influence evolution as a population

genetic process. Our simulations show that dormancy can only influence selection under certain

36   demographic conditions, suggesting that the active portion of a population can continue to

undergo adaptive evolution even in the presence of a large seed bank. We then extend our

38   analysis to examine how seed banks can alter macroevolutionary dynamics and determine

whether metabolically inactive microbial populations are engaging in dormancy as a life history

40   strategy. We propose that by examining the population genetic and macroevolutionary

implications of dormancy as a life history strategy we can understand the extant that seed banks

42   influence microbial evolutionary dynamics.

KEYWORDS: microorganisms, evolution, population genetics, macroevolution, seed banks,

46   dormancy, bacteria, bet-hedging

INTRODUCTION

48   Nature is rarely predictable. Resource availability, disease pressure, and temperature are

just a few of the abiotic and biotic factors that fluctuate over space and time. Variation in these,

50   and other, biotic and abiotic factors have important consequences for the growth, survival, and

reproduction of individuals. Many taxa respond to variability in their environment by entering a

52   reversible state of reduced metabolic activity, a phenomenon known as dormancy (Lennon &

Jones, 2011). Dormancy is an adaptive trait that has independently evolved multiple times across

54   the tree of life (Guppy & Withers, 1999). By entering a dormant state individuals can endure

conditions that are sub-optimal for growth and reproduction, which results in the net increase of

56   a population's long-term geometric fitness (Cohen, 1966). However, dormancy comes at a cost.

This cost not only reflects the missed opportunities for reproduction, but also the energetic

58   investment required to produce resting structures (Bradshaw et al., 1998; Cáceres & Tessier,

2004; van Bodegom, 2007). Despite these costs, dormant individuals accumulate in many

60   systems and form a seed bank (Locey et al., 2017), which serves as a reservoir of genetic and

phenotypic diversity (Fig. 1). Seed banks have important implications for a range of ecological

62   phenomena, perhaps most central is the maintenance of biodiversity. Specifically, dormancy

reduces interspecific competition allowing for coexistence, a mechanism known as the storage

64   effect (Chesson & Warner, 1981; Chesson, 1994). Seed bank mediated diversity has

consequences for other ecological processes including patterns of succession (Marks, 1974;

66   Bazzaz, 1979; Lennon & Jones, 2011), community responses to perturbations (Kalamees &

Zobel, 2002), and ecosystem processes (Wang et al., 2014).

68   Dormancy also has important consequences for evolution. For example, rates of

phenotypic evolution are reduced in freshwater zooplankton that persist in a dormant state

70   (Hairston & De Stasio, 1988). Similarly, dormancy has also been found to be associated with a

higher rate of lineage diversification in plants (Willis et al., 2014). However, to fully understand

72   how dormancy influences evolution, it is necessary to understand how it affects the relative

strength of the fundamental forces that govern the rate and direction that allele frequencies

74   change over multiple generations. Evolution is a population genetic process governed by both

deterministic and stochastic forces, the relative strengths of which dictate how genetic diversity

76   is generated, lost, and maintained. Because dormancy as a life history strategy assumes trade-offs

that affect fitness, it is subject to the deterministic force of selection if it is sufficiently strong

78   (Cohen, 1966; Templeton & Levin, 1979; Brown & Venable, 1986). In addition, dormancy can

trade-off with dispersal as an alternative life history strategy (Venable & Lawlor, 1980; Venable

80   & Brown, 1988; Olivieri, 2001; de Casas et al., 2015), affecting the evolutionary force of gene

flow (Vitalis et al., 2004). However, far less progress has been made towards understanding how

82   dormancy influences the stochastic forces of evolution that operate within a single population

(e.g., genetic drift, mutation, and recombination). To gain a complete understanding of how

84   dormancy influences evolutionary dynamics it is necessary to consider how it affects the

stochastic and deterministic forces that underpin evolutionary biology.

86   While dormancy can influence eco-evolutionary dynamics across the tree of life

(Hairston et al., 1999; Willis et al., 2014), it could have a particularly large effect on

88   microorganisms. Dormancy is prevalent across ecosystems, ranging from the human gut to

engineered ecosystems to the global ocean (Lennon & Jones, 2011). For example, > 90% of

90   microbial biomass is soils is metabolically inactive (Alvarez et al., 1998; Lennon & Jones, 2011;

Blagodatskaya & Kuzyakov, 2013). In addition, microorganisms can alter their metabolically

92   activity in a matter of hours, suggesting that they can quickly transition between active and

dormant states (Fig. 2; Walker & Winslow, 1932; Kaprelyants & Kell, 1993; Votyakova et al.,

94   1994; Ishiguro et al., 2015). Although metabolic transition may occur stochastically (Epstein,

2009), dormancy can also be regulated by environmental cues, such as changes in temperature

96   (Oliver et al., 1995), pH (Keynan et al., 1964), water (Aadnerud et al., 2015), and resource

supply (Dworkin & Shah, 2010). The amount of time that microorganisms spend in a dormant

98   state is variable. Some microorganisms succumb to environmental stress within days (Bale et al.,

1993), while others can persist in a dormant state for prolonged periods of time. For example,

100   viable microorganisms have been retrieved from ancient materials (e.g., permafrost, amber,

halite crystals) that, in some cases, are hundreds of millions of years old (e.g., Johnson et al.,

102   2007). As a consequence, dormant microorganisms can survive for periods of time that far

exceed the average generation time of actively reproducing individuals (Cano & Borucki, 1995;

104   Vreeland et al., 2000). This means that active individuals can continue to evolve while

individuals in the seed bank remains relatively static. Because a considerable fraction of

106   phylogenetically diverse microbial taxa are able to enter a dormant state across disparate

environments, it is likely that dormancy has influenced the evolutionary history of

108   microorganisms. This influence has the potential to extend across evolutionary scales, ranging

from the population genetic processes that underlie the evolutionary dynamics of populations

110   (i.e., genetic drift, mutation, selection, recombination, and gene flow) to the rates that microbial

lineages diverge and go extinct.

112   In this paper, we focus on how dormancy affects microbial evolution. We do this by

synthesizing existing population genetic theory and developing novel simulations that provide

114   insight into how seed banks can affect the fundamental forces that govern the rate and direction

of evolution. We then examine how dormancy influences evolutionary dynamics at longer time

116   scales, including macroevolutionary processes such as speciation and extinction. In addition, we

examine environments where organisms are often found in a dormant state to determine whether

118   dormancy is an evolutionarily viable bet-hedging strategy or simply a metabolic response to

conditions that are sub-optimal for growth and reproduction (Box 1). Environments do not

120   always improve and if organisms never have the opportunity to reproduce, the ability to enter a

dormant state will merely increase the amount of time it takes for a population go extinct.

122   Throughout this paper, we emphasize the importance of using population genetic theory to

understand how dormancy as a life history strategy can alter evolutionary dynamics (Box 2).

124   While we focus on dormancy and the evolution of microbial populations, we believe that the

framework and conclusions apply to organisms across the tree of life

POPULATION GENETIC CONSEQENCES OF DORMANCY

128   Seed banks preserve genetic and phenotypic diversity by stratifying the population. In a

population with a seed bank, individuals can effectively be grouped into active and dormant sub-

130   populations, where they enter and exit a dormant state in a manner analogous to migration

between sub-populations (Lennon & Jones, 2011; Blath et al., 2015) (Fig. 1). Seed banks are

132   sometimes viewed simply as an evolutionary buffer (Koopmann et al., 2016). Dormancy

preserves existing genetic diversity by decreasing the rate that genetic diversity is removed from

134   the population (Hairston & De Stasio, 1988; Vitalis et al., 2004; Koopmann et al., 2016) and

increasing in the effective size of the population (Nunney, 2002). However, when individuals

136   remain dormant for extremely long periods of time, dormancy can have complex and non-

intuitive effects on evolutionary dynamics (Blath et al., 2015). In the following sections, we use

138   population genetic theory to examine how seed banks affect each of the fundamental forces of

        evolution and expand on that theory through the use of novel simulations.

        Genetic drift — The amount of genetic diversity that can be maintained in an ideal population of

142   finite size is determined by the number of individuals in the population (N) and the per-

        generation mutation rate (μ). When the rate that genetic diversity is acquired by mutation is

144   equal to the rate that it is lost by drift (i.e., mutation-drift equilibrium), our expectation for the

        maximum amount of genetic diversity that can be maintained in a population of haploid

146   individuals is $\theta = 2N\mu$. This equation can conveniently be interpreted as the ratio of the rate that

        genetic diversity is acquired by mutation (2μ) and lost by drift ($1/N$) each generation (Kimura,

148   1969). However, our expectation needs to be modified when a population contains dormant

        individuals. Because dormant individuals do not reproduce they are not subject to random

150   genetic drift, which increases the maximum amount of genetic diversity that can be maintained

        in a finite population. Thus, all else being equal, dormancy should inflate estimates of genetic

152   diversity (Levin, 1990).

        Two classes of theoretical population genetic models have been used to explore how

154   genetic diversity scales with dormancy. The first class of models examines what is known as the

        weak seed bank effect. It assumes that individuals can spend a maximum number of generations

156   in the seed bank that is smaller than the number of active individuals in a population, meaning

        that individuals enter and exit the seed bank before the genetic diversity in the active portion of

158   the population turns over due to the joint effects of mutation and drift (Kaj et al., 2001). The

        weak seed bank model predicts that dormant individuals increase genetic diversity, but do not

160   change the pattern of ancestry among individuals in the population (i.e., the shape of the

population's genealogy). In contrast, the strong seed bank model has no constraint on the

162   maximum number of generations that an individual can remain in the dormant state (González-

Casanova et al., 2014; Blath et al., 2015; 2016). This lack of constraint means that the length of

164   time that an individual spends in the seed bank (where time is scaled as generations) can be

longer than the number of active individuals in the population. Under this scenario genetic

166   diversity within the active portion of the population effectively turns over before an individual in

the seed bank has the opportunity to reproduce. The result of the strong seed bank effect is that

168   dormancy drastically alters the pattern of ancestry among individuals in the total population in

addition to increasing the maximum amount of genetic diversity that can be maintained in the

170   population (Fig. 3a). Here the expected amount of genetic diversity ($\mathbf{E}[\pi]$) in a population with

M dormant individuals is $\mathbf{E}[\pi] = \theta + \theta(\frac{M}{N})$, which can be interpreted as the amount of genetic

172   diversity expected in a population with N actively reproducing individuals ($\theta = 2N\mu$) plus the

seed bank's contribution towards reducing the rate of genetic drift scaled by the relative excess

174   of dormant individuals $(\frac{M}{N})$.

In addition to increasing genetic diversity in a population, our simulations suggest that

176   the average length of time required for a population to reach mutation-drift equilibrium

drastically increases under the assumption of a strong seed bank (Fig. 3b). However, while the

178   weak and strong seed bank models present viable hypotheses about how dormancy can increase

genetic diversity, little work has been done to determine if inferred patterns of ancestry in actual

180   populations capable of entering a dormant state resemble predictions from the strong or weak

seed bank effect. Determining whether natural populations capable of entering a dormant state

182   exhibit the strong or weak seed bank effect will require comparing inferred patterns of ancestry

to the predictions of each model.

Mutation — Mutations are the ultimate source of genetic diversity. In populations composed entirely of reproducing (i.e., active) individuals, the majority of newly acquired genetic variation is due to genome replication errors acquired during cell division (Kunkel, 2004). Given that dormant individuals do not reproduce, it is reasonable to expect that all observed genetic diversity was acquired at some point in the past when the population was active, an assumption that is often made in population genetic models that include dormancy (Vitalis et al., 2004; Blath et al., 2015). However, mutations can still arise in populations that experience sub-optimal conditions. For example, the rate of mutation can be effectively decoupled from the rate of replication in *Mycobacterium tuberculosis* due to oxidative DNA damage that occurs when individuals spend a prolonged period of time in a dormant state (Ford et al., 2011; 2013). This reduced correlation between the rate of replication and the rate of mutation via oxidation-induced DNA damage likely applies to other types of mutagens (e.g., high intracellular concentrations of a chemical or physical mutagen). If acquiring mutations during dormancy is common among microorganisms, then it is necessary to consider the rate of mutation that occurs between replication events as well as the mutation rate during replication to understand how genetic diversity is acquired in populations.

The extant that mutation can be decoupled from replication depends on the mechanism used to enter dormancy. For example, DNA is protected during dormancy in spore-forming taxa like *Bacillus* by the extremely low water content in the spore coat and small, acid-soluble proteins that bind to the DNA (Setlow, 1992; 2006; Moeller et al., 2009; 2014), as well as the up-regulation of DNA repair during the early phases of resuscitation (Lenhart et al., 2012; Ramírez-Guadianaa et al., 2012). These mechanisms protect spores from the loss of guanine due to

depuriation as well as UV radiation-induced pyrimidine dimers (Setlow, 1992), suggesting that

208   spores acquire mutations at a reduced rate. However, many taxa enter and exit dormancy without

producing specialized resting structures (e.g., spores, cysts, akinetes). In contrast, non-

210   sporulating taxa can likely only rely on mismatch repair mechanisms that are up-regulated during

resuscitation (Mizrahi & Anderson, 1998; Rittershaus et al., 2013) as well as on DNA repair

212   mechanisms that can be maintained under an energy-limited state (Gong et al., 2005; Dieser et

al., 2013; Rittershaus, 2013). For non-sporulating taxa that spend extended periods of time in an

214   energy-limited state it is likely that the long-term mutation rate is primarily determined by the

fidelity of DNA repair mechanisms that operate during dormancy, rather than that of

216   polymerases used during periods of rapid growth. Maintained DNA repair may be a better

mechanism for long-term cell survival rather than an absolute lack of metabolic activity, as

218   evidence of DNA repair has been found in bacteria isolated from half-million-year-old

permafrost samples (Johnson et al., 2007). While the exact cost of maintaining DNA repair

220   mechanisms relative to the total energy budget of energy-limited cells is not known, it is small

enough that it has been ignored in estimates of the cost of cell maintenance under conditions of

222   homeostasis (Lynch & Marinov, 2015) and extreme energy limitation (Kempes et al., 2017),

suggesting that a certain level of mismatch repair can be maintained over long periods of

224   dormancy without draining the total cellular budget.

        Attempting to reproduce under sub-optimal conditions, rather than going dormant, can

226   elevate the rate of mutation in a population. Microbial mutation rates are often elevated under

stressful conditions due to the up-regulation of error-prone machinery used for DNA replication

228   and repair (Witkin, 1976; Foster, 2007). It has been argued that this error-prone machinery may

be beneficial under times of stress, as an increased mutation rate should increase the number of

230 beneficial mutations acquired over a given length of time (Rosenberg, 2001; Foster, 2007;

Galhardo et al., 2007). For example, DNA polymerases used under times of stress can confer a

232 competitive advantage to starved populations of *Escherichia coli*, known as the Growth

Advantage at Stationary Phase (GASP) phenotype (Finkel, 2006). However, this temporary

234 fitness advantage can have long-term consequences for the survival of the population. The

majority of mutations are deleterious (Lynch et al., 1999) and there is no reason to expect that an

236 increased mutation rate will change the distribution of mutational fitness effects. Thus, any

increase in the mutation rate will likely be accompanied by a proportional increase in the number

238 of deleterious mutations acquired per unit time, contributing towards the long-term deterioration

of the genome (Gerrish et al., 2013; Lynch et al., 2016). Populations that continue to grow in a

240 starved state likely constitute an evolutionary dead end, as the accumulated deleterious mutations

would result in a decrease in fitness once the environment improves. Rather, if there is sufficient

242 temporal variation in the environment then it is possible that the long-term geometric fitness of a

population would be maximized by persisting in a dormant state rather than attempting to

244 reproduce and incurring the negative effects of an elevated mutation rate.

246 *Selection* — Because dormancy can act as a buffer against the stochastic forces of mutation and

genetic drift, it is reasonable to expect that the presence of a seed bank will reduce the rate that

248 the directional force of natural selection removes genetic diversity from the population. Natural

selection is less efficient in models that assume a weak seed-bank effect, where the average

250 amount of time required for selection to drive a beneficial allele to fixation (T_{fix}) increases

quadratically with the average number of generations that an individual spends in a dormant state

252 (Koopmann et al., 2016). However, little work has been done to examine selection under a strong

seed-bank effect or how selection affects T_{fix} differently for active and dormant portions of the

254 population. Intuitively we would expect that T_{fix} would increase, since individuals can spend an

even longer period of time in a dormant state. Consequently, beneficial alleles should take a

256 longer period of time to go to fixation because individuals must enter and exit the seed bank to

spread the beneficial allele throughout the whole population. To test this hypothesis, we

258 simulated the trajectory of a beneficial allele in a population subject to the strong seed bank

effect and examined the dormant and active portions of the population separately (see

260 Supplemental Materials: Selection simulation). As expected, we found that the average amount

of time required for a beneficial allele to reach fixation increases with the average number of

262 generations that an individual spends in the seed bank. Our simulations are supported by

previously developed theory that found a similar increase in T_{fix} as the average number of

264 generations that an individual spends in a dormant state increases (Fig. 4a,b) (Koopmann et al.,

2016). However, because the seed bank stratifies the population into active and dormant

266 components and a fitness advantage only increases the frequency of a beneficial allele if its

carrier is reproducing, it is necessary to examine the active sub-population separately from the

268 dormant sub-population. In our model, the beneficial allele initially rose in frequency until it was

effectively fixed (a.k.a., quasi-fixation), where it then fluctuated below a frequency of one as

270 individuals continued to exit and enter the seed bank (analogous to back-mutation in a single-

locus model). By examining the active portion of a population with a seed bank in a simulation

272 of single locus selection, we found that the average number of generations required for a

beneficial allele to reach quasi-fixation is similar to that for the dormant portion up until the

274 average time in the seed bank approaches 100 generations, which is the threshold between the

weak and strong seed bank effects (Fig. 4b). Surprisingly, the average length of time until a

276 beneficial allele is quasi-fixed among actively reproducing individuals decreases after the strong

seed bank threshold, approaching the expected T_{fix} for a population without a seed bank. This

278 decrease in T_{fix} is likely because the input of genetic diversity via resuscitation is low enough

that it does not increase the amount of time required for a beneficial allele to reach fixation, a

280 novel result that has not previously been reported. Contrary to expectations, our simulations

suggest that if the average length of time that an individual spends in a dormant state (scaled as

282 generations) is longer than the number of active individuals, the presence of a seed bank will not

interfere with selection in the active portion of the population.

284 The decline of T_{fix} for the active portion of a population subject to the strong seed bank

effect has important implications for the long-term evolutionary dynamics of microbial

286 populations. Because the majority of a given microbial genome is likely clonal, the rate of

evolution in a population undergoing adaptive evolution can be sufficiently described by the

288 beneficial mutation rate (U) and the strength of selection operating on a population of constant

size (N) (Desai et al., 2007). Given that the average time it takes for a beneficial mutation to arise

290 (T_{mut}) is much greater than T_{fix}, the population will evolve at a rate (v), which is proportional to

the rate that beneficial mutations of small effect arise and fix within a population (i.e., $v \propto$

292 $s^2 NU$) (Park et al., 2010). If this condition is violated, multiple beneficial mutations can arise on

different genetic backgrounds and compete in the population, increasing T_{fix} through the process

294 of clonal interference (Gerrish & Lenski, 1998). As no one has examined clonal interference in

an asexual population with a seed bank, there is no theory to draw on that would allow one to

296 determine whether dormancy increases or decreases the rate of evolution in the multiple

mutation regime. Our results suggest that the rate of evolution in the active portion of a microbial

298 population with a strong seed bank can likely be approximated as a series of mutations fixing in

the population. However, T_{fix} will increase if individuals on average spend fewer generations in

300   a dormant state than the number of active individuals (Fig. 4b). If the presence of a seed bank

increases T_{fix} to the point that it is not much larger than T_{mut}, then multiple beneficial mutations

302   will segregate simultaneously, moving the population into the clonal interference regime. To

fully describe the dynamics of clonal interference in a population with a seed bank it will be

304   necessary to extend the single beneficial mutation dynamics described here to the multiple

mutation regime using simulations and population genetic theory.

306   The average amount of genetic diversity among all individuals in a population with a

seed bank will be preserved for longer periods of time in the face of selection. This preservation

308   of genetic diversity has important implications for microbial evolution, as it is often thought that

microbial populations contain low levels of diversity across their genomes due to linked positive

310   selection (i.e., genetic draft) (Gillepsie, 2000; Lynch, 2007). Genetic draft can dominate genetic

drift as the primary stochastic force in large microbial populations, constraining the maximum

312   amount of genetic diversity that can be maintained in a population (Smith & Haigh, 1974;

Gillespie, 2000). Evolution experiments have confirmed that linked selection regularly reduces

314   genetic diversity in microbial populations and that it operates on the relatively fast time scale of

hundreds to thousands of generations (Maddamsetti et al., 2015; Tenaillon et al., 2016).

316   However, such experiments are commonly carried out in constant environments that are near-

optimal for growth and reproduction, potentially providing an inaccurate representation of how

318   selection operates in naturally occurring populations where individuals can enter a dormant state.

While it is difficult to examine the genetic diversity of natural microbial populations over time, a

320   recent study found evidence of only a single case of genome-wide linked selection in 30

freshwater microbial populations monitored over nine years (Bendall et al., 2015). Given that

322 approximately half of all bacterial cells in freshwater lakes may be dormant (Lennon & Jones,

2011), the small number of selective sweeps observed could be explained by the presence of a

324 seed bank that increased the length of time required for selection to drive beneficial alleles to

fixation among both active and dormant individuals. Accurately inferring the extant that

326 selection can act on a population that capable of entering a dormant state requires accounting for

the seed bank.

Recombination — Dormancy can likely preserve the diversity of genes as well as the diversity

330 of alleles generated by recombination. The ability for microorganisms to acquire and exchange

DNA by horizontal gene transfer (HGT) is a major evolutionary feature that can promote the

332 spread of beneficial genes within a population (Medini et al., 2005; Lapierre & Gogarten, 2009;

Dixit et al., 2016). The stochastic processes of gene gain and loss via HGT leads to the uneven

334 distribution of genes among genomes in a population (Berg & Kurland, 2002; Baumdicker et al.,

2012). This uneven distribution is often summarized as the total number of genes found within

336 the population (i.e., the pangenome) and the number of genes shared among all individuals

within the population (i.e., the core genome). Similar to how the presence of a seed bank

338 increases the maximum amount of genetic diversity that can be maintained in a finite population

by reducing the effect of genetic drift, it is possible that the presence of a large seed bank would

340 increase the size of the pangenome. In addition, a strong seed bank could alter the distribution of

gene frequencies within a population, analogous to how the strong seed bank effect alters the

342 distribution of allele frequencies within a population (Blath et al., 2015). Given that closely

related microorganisms can often share very little gene content, it is possible that the ability for

344 microorganisms to persist in a dormant state can preserve genic diversity acquired by

recombination.

Gene flow — Seed banks may reduce the effect that migration has on altering allele frequencies

348 within a population (i.e., gene flow). The ability to enter a dormant state can increase the chance

that any two individuals share a common ancestor within a population (i.e., the probability of

350 identity by descent) and reduce the amount of estimated differentiation among populations (i.e.,

the fixation index, F_{st}) (Vitalis et al., 2004; Živković & Tellier, 2012; Hollander & Pederzani,

352 2017). This increase in genetic similarity due to the presence of a seed bank has been found in

natural populations of plants (Tellier et al., 2011; Živković & Tellier, 2012). However, dormancy

354 and migration are not always independent. For many organisms, dormancy can increase the

probability of successfully migrating and establishing in a new population. For example, the

356 spatial distribution of endospore-forming thermophilic bacteria in marine sediments closely

reflects global ocean currents (Müller et al., 2014). However, dormancy and migration can

358 represent a trade-off if the ability to enter a dormant state involves investing energy into a

different mechanism than the one used to migrate (Olivieri, 2001). While the potential for

360 dormancy and migration to be either positively or negatively correlated is well known in life

history theory, little work has been made towards accounting for this relationship in population

362 genetic theory. To understand the evolutionary consequences of dormancy and migration it is

necessary to examine how these life history strategies interact to shape patterns of shared genetic

364 diversity between populations.

366 **MACROEVOLUTIONARY CONSEQUENCES OF DORMANCY**

In the previous section, we examined how seed banks influence the rate that allele

368   frequencies change over relatively short periods of time within a lineage (i.e., microevolution).

However, if seed banks buffer the aggregated effects of multiple evolutionary forces over long

370   periods of time, then dormancy may have important implications for macroevolutionary

phenomena. In the following section, we examine how dormancy influences processes and

372   patterns such as the rate of molecular evolution, shared ancestry among lineages, speciation, and

extinction.

The rate of molecular evolution — If dormancy can affect the rate of evolution in microbial

376   populations, we would likely observe it in bacteria that are uniquely capable to persist for

extended periods of time in an inactive state via their capacity to form resistant, long-lived

378   spores. Many bacteria in the phylum *Firmicutes* possess the ability to form endospores, which

are thought to be one of the most metabolically inert forms of life on the planet (Setlow, 2014).

380   Endospore formation is triggered by low resource availability, where the endospore is built

within the cell, lyses, and is released into the environment (Tan & Ramamurthi, 2013). These

382   endospores are persistent and in some extreme instances have reportedly been found to survive

for hundreds of millions of years (Vreland, 2000). Endospores can endure exposure to extreme

384   conditions such as high doses of gamma and UV radiation (Nicholson et al., 2000; 2005),

desiccation (Nicholson et al., 2000), and the vacuum of space (Horneck et al., 1994; 2012).

386   Given that spore-forming bacteria can persist for long periods of time without reproducing, one

might expect non-spore-forming relatives to have more rapid rates of molecular evolution. This

388   hypothesis was tested by analyzing a large collection of *Firmicutes* genomes. Some of these

genomes came from isolates that had the ability to form endospores while other genomes came

390   from isolates that had lost the ability to form endospores (Weller & Wu, 2015). The rate of

       amino acid and synonymous substitutions for non-spore forming isolates were significantly

392   elevated relative to spore-forming taxa and the phylogenetic branch length declined as the

       number of spore-forming genes within a genome increased (Fig. 5a; Weller & Wu, 2015). These

394   results suggest that the rate of evolution increases when a lineage loses the ability to enter

       dormancy via sporulation. To further evaluate this effect of dormancy on the rate of evolution,

396   we ran a simulation for the simple case of a neutrally evolving population, where we

       manipulated the average number of generations that an individual spends in the seed bank and

398   calculated the substitution rate after 10,000 generations (see Supplementary Methods for more

       detail). For the entire population (i.e., both active and dormant individuals) we found that

400   substitution rate declined as average time in the seed bank increased, a finding that should apply

       to microbial evolution irrespective of the mechanisms controlling transitions into and out of a

402   dormant state (Fig. 5b).

             While theoretical and empirical evidence suggests that seed banks can reduce the rate of

404   evolution within a lineage, this effect may be short lived if the ability to enter and exit a dormant

       state cannot be maintained. For example, the ability to form endospores has been lost multiple

406   times within the *Firmicutes* phylogeny (Weller & Wu, 2015). The repeated loss of this seemingly

       beneficial trait across lineages can be explained by adaptive and non-adaptive evolutionary

408   processes operating within a lineage. Organisms often need to invest in morphological or

       physiological structures that allow individuals to enter and exit a dormant state in response to

410   fluctuating environmental conditions (Ayati & Klapper, 2012; Akiyama et al., 2017). The

       selective advantage of maintaining the ability to enter a dormant state depends on both this

412   energetic cost as well as the long-term fitness contribution of dormancy in a fluctuating

environment, where the strength and direction of natural selection can change over time (Simons,

414 2009). In relatively stable environments, the mechanism necessary to enter a dormant state may

be lost from a lineage, either due to relaxed selection leading to the accumulation of mutations at

416 dormancy-encoding loci or selection against the energetic cost of the mechanism (Dawes &

Thornley, 1970). For example, in long-term evolution experiments, five populations of *Bacillus*

418 *subtilis* lost the ability to form endospores when medium was continuously supplied so that

individuals did not experience starvation (Maughan et al., 2007). The accumulation of mutations

420 due to relaxed selection was responsible for the loss of endospore formation in four of the five

populations, while the loss of sporulation actually increased fitness in the other population. Spore

422 formation may be lost because it is a trait that is encoded by more than 200 genes (Galperin et

al., 2012), making it a large target for mutation. Alternatively, it is possible that the underlying

424 genes may be lost because the energetic cost of even a single nucleotide is visible to natural

selection in bacteria (Lynch & Marinov, 2015). In contrast to endospore-formation, other

426 mechanisms regulating dormancy appear to require fewer genes than what is reported in spore-

forming bacteria. For example, resuscitation promoting factor (Rpf) is an enzyme encoded by a

428 single gene that terminates dormancy for many bacteria belonging to the *Actinobacteria* phylum

(Mukamolova et al., 1998; Keep et al., 2006). Rpf cleaves the 1-4 linkage between the amino

430 sugars N-acetylglucosamine and N-acetylmuramic acid in peptidoglycan, a major cell-wall

component of nearly all bacteria. The gene encoding Rpf likely has a lower probability of being

432 lost owing to its small size and lower energetic cost. Ultimately, whether a mechanism used to

enter a dormant state is retained in a lineage long enough to affect the rate of evolution will

434 depend on the amount of environmental variation the lineage experiences over extended periods

of time, the energetic cost of the mechanism, the mutation rate of the lineage, and the number of

436   nucleotides that encode the mechanism.

438   ***Shared ancestry among lineages*** — Dormancy may provide novel insight into confusing

patterns of ancestry among microbial lineages. Many microbial genomes harbor genes that have

440   no detectable homologues in closely related lineages (i.e., ORFans) (Daubin & Ochman, 2004;

Mira et al., 2004; Zhaxybayevaa et al., 2009). One explanation for the existence of ORFans is

442   horizontal gene transfer (HGT). HGT is a reasonable hypothesis given that genic and genomic

content can vary among and within microbial lineages due to the exchange of genetic material

444   (Ochman et al., 2000). However, certain ORFans are essential for the process of cell division,

even though they are not present in neighboring lineages (e.g., Mira et al., 2004; Zhaxybayevaa

446   et al., 2009). Therefore, it is unclear to what extant essential genes can be lost and acquired

without resulting in the death of the cell. Another explanation for existence of ORFans is seed

448   banks (González-Casanova et al., 2014). If some fraction of individuals can remain dormant for

extremely long periods of time (González-Casanova et al., 2014), then the active portion of the

450   population may continue to accumulate substitutions to the point that certain genes in the active

pool no longer share homology with members in the dormant pool. To test this, a set of ORFan

452   genes from the nitrogen-fixing soil bacterium *Azotobacter vinelandii* and neighboring lineages

were examined (González-Casanova et al., 2014). These ORFan genes had the characteristic

454   codon-usage and GC-content biases of *A. vinelandii*, but showed little homology to closely

related members of the genus *Pseudomonas*, a result that was interpreted as support for the

456   strong seed bank hypothesis.

While the strong seed bank hypothesis may explain the long-term evolutionary

458   consequences of remaining in a dormant state, an alternative hypothesis can explain the existence

of ORFans. First, additional copies of essential genes could be acquired by HGT, leading to

460   relaxed selection on each gene copy. The original copy could then be physically lost due to gene

deletion or functionally lost due to accumulated mutations (Lynch, 2007). This new copy would

462   then gradually acquire the same GC-content as the rest of the genes within the lineage through a

combination of lineage-specific mutation bias and the rate of GC-biased gene conversion

464   (Lynch, 2007; Lassalle et al., 2015). In addition, lineage-specific selection on translational

efficiency would shape the codon composition of the gene (Drummond & Wilke 2008; Plotkin &

466   Kudla, 2011), particularly if the lineage has a large effective population size (Sharp & Li, 1987;

Sharp et al., 2005). These directional pressures suggest that horizontally acquired genes that are

468   retained in the genome may lose their signal of homology while becoming similar in composition

to vertically acquired genes. While the theoretical models necessary to describe the effect of the

470   seed bank on the genetic composition of microbial lineages exist (González-Casanova et al.,

2014), further study is needed to determine whether the strong seed bank effect applies to gene-

472   specific patterns of ancestry.

474   *Speciation* — While not theoretically well-developed, it is possible that dormancy can affect

rates of speciation. Population genetic theory suggests that the presence of a seed bank can

476   preserve genetic similarity among separate populations (Vitalis et al., 2004). The ability to exit a

dormant state effectively acts like gene flow in physically separated populations, preserving

478   genetic diversity and similarity that might otherwise be lost due to selection acting on newly

arisen alleles within a population. The preserved genetic similarity due to the seed bank would

480   likely slow the rate of evolutionary divergence between lineages due to geographic divergence,

reducing the rate of allopatric speciation. However, if a portion of the population remains in a

482   dormant state while the rest of the population diverges to the extent that genetic material can no

longer be exchanged, then dormant and active individuals may ultimately diverge into separate

484   lineages. This process of divergence is an extension of the strong seed bank effect previously

discussed (González-Casanova et al., 2014), where the focus is on how evolutionary divergence

486   between dormant and active individuals can prevent the exchange of genetic material, rather than

on patterns of ancestry. While seed banks may influence the rate of speciation, very little is

488   known about how microbial lineages split or the rate that they split, regardless of the microbial

species definition. While the rates of speciation are unclear, recent research on the net difference

490   of speciation and extinction rates (i.e., diversification) in microorganisms suggests that microbial

taxa split at a constant rate through time (Marin et al., 2017). If so, then we would expect that

492   any decrease in the rate of speciation due to the presence of a seed bank would be associated

with a similar increase in the rate of extinction, or vice-versa. The exact evolutionary

494   mechanisms that are responsible for this constant rate of lineage diversification are currently

unknown. Determining to what extant the ability to persist in a dormant state reduces the rate of

496   speciation requires further research.

498   ***Extinction*** — Because seed banks can maintain preserve the size of the population, they can

likely buffer populations from extinction (Kalisz & McPeek, 1993). However, while next to

500   nothing is known about extinction rates in microorganisms (Weinbauer & Rassoulzadegan,

2007), we can extend evolutionary models of microbial extinction to examine how the ability to

502   enter a dormant state might decrease the rate of extinction. The primarily clonal structure of

microorganisms means that most mutations cannot be removed by recombination once they

504   become fixed. It is often argued that the rate of extinction in a microbial lineage is determined by

the fixation rate of deleterious mutations, since once a deleterious mutation is fixed it can only be

506   removed by the fixation of a reverse mutation at the same site; a process known as Muller's

Ratchet (Muller, 1964; Felsenstein, 1974). The negative fitness effect of deleterious mutations

508   accumulated over time can lead to the extinction of a lineage. In this case, extinction happens

because reproductive output is reduced due to fixed deleterious mutations, increasing the

510   strength of genetic drift relative to natural selection, leading to a positive feedback loop where

deleterious mutations continue to accumulate and reproductive output declines over time. This

512   feedback loop eventually results in the extinction of the lineage, a phenomenon known as a

mutational meltdown (Gabriel et al., 1993; Lynch et al., 1993). Because lineages persisting in a

514   dormant state likely fix mutations at a far lower rate (Fig. 5), extinction via mutational meltdown

is less likely to occur within lineages capable of forming a seed bank. Given that dormant

516   individuals can persist on time scales upwards of millions of years (Cano et al., 1995; Greenblatt

et al., 2004), it is worth determining whether there is a relationship between the age of a

518   microbial lineage and its ability to persist in a dormant state.

520   **CONCLUSION**

The ability to enter a dormant state is a common life history strategy that is found across

522   the tree of life. Historically dormancy has been viewed as an adaptive trait that preserves genetic

and taxonomic diversity, with comparatively little attention given to its effects on the

524   fundamental forces that underlie evolution. Using recent developments in theoretical population

genetics and novel simulations we examined how the ability to enter a dormant state effects the

526   population genetic forces that underlie evolutionary biology as well as common estimates of

genetic diversity. Specifically, we identified the demographic scenarios where dormancy can

528   affect natural selection. We then extended these results to determine how dormancy can

influence the evolutionary dynamics of microbial lineages and in what environmental systems

530   the ability to enter a dormant state is a viable life history strategy. We conclude that while

dormancy can influence evolutionary dynamics and common estimates of genetic diversity, it is

532   necessary to consider whether dormant organisms within a system are engaging in a life history

strategy or simply reducing their metabolic rate out of physiological necessity. By considering

534   the short-term population genetic and long-term macroevolutionary implications of dormancy as

an adaptive trait researchers can examine the evolutionary importance of dormancy. Determining

536   the extant that dormancy influences microbial evolution will require empirical research on

whether dormant microorganisms in the environment are engaging in a life history strategy as

538   well as continued development of the population genetic theory that accounts for the

evolutionary effects of dormancy.

Box 1: Dormancy in low-energy environments

550   In nature, most organisms experience conditions that are suboptimal for growth and

reproduction. This is true for microorganisms, just as it is for plants and animals. For example,

552   approximately 30-50% of the prokaryotic (i.e. Bacteria and Archaea) cells on Earth live under

continents and on the ocean floor in a marine subsurface habitat known as the "deep biosphere".

554   In this environment, microorganisms are fueled by remnants of organic matter produced in the

well-lit, productive surface waters that sink to the bottom of the ocean. As a result,

556   microorganisms in the deep biosphere rest on the thermodynamic edge of life and death

(Kallmeyer et al., 2012; Parkes et al., 2014; Jørgensen & Marshall, 2016; Starnawski et al.,

558   2017). The deep biosphere likely constitutes the largest seed bank on the planet, where spore-

forming bacteria are one of the most dominant groups of microorganisms (Lomstein et al., 2012).

560   Furthermore, microorganisms in the deep biosphere spend an extremely long time in the seed

bank. Based on rates of amino acid racemization, it is estimated that the microbial biomass pool

562   may only turnover once every thousand years (Lomstein et al., 2012; Hoehler & Jørgensen,

2013; Jørgensen & Marshall, 2016). In a recent study where four microbial lineages within the

564   deep biosphere were sequenced, all lineages showed little-to-no signature of divergence or

difference in standing genetic variation from populations at the surface. In addition, the ratio of

566   synonymous to non-synonymous polymorphisms did not change with sediment depth for three

out of four lineages, suggesting that populations within the deep biosphere are not undergoing

568   adaptive evolution as they sink deeper into the sediment (Starnawski et al., 2017). These findings

are consistent with expectations of how dormancy should affect evolution as a buffer, suggesting

570   that the lack of divergence, adaptive evolution, and similar estimates of standing genetic

variation across sediment depth could be explained by the presence of a large microbial seed

572   bank. However, certain areas of the deep biosphere are thought to be remarkably stable for long

periods of time, perhaps millions of years (Jørgensen, 2012), a length of time that is likely

574   several orders of magnitude longer than the average generation time. However, the slow turnover

rate raises the question of whether dormancy is a viable life history strategy for microorganisms

576   in the deep biosphere. Without a change in environmental conditions which would occasionally

favor growth and reproduction, the deep biosphere may simply reflect a very large collection of

578   microorganisms that are on a slow march to death. Without the possibility of resuscitation,

remaining in a dormant state is ultimately an evolutionary dead end. These stable environmental

580   conditions suggest that dormancy is unlikely to be a viable life history strategy (Malik & Smith,

2008). Rather, in the deep biosphere dormancy may simply be a metabolic necessity due to the

582   energetic constraints of the environment.

In contrast, dormancy is likely a viable life history strategy in certain regions of the

584   permafrost, a temperature and energy-limited environment covering 25% of Earth's terrestrial

surface (Graham et al., 2012). Upper layers of the permafrost that are not covered with ice can go

586   through annual freeze-thaw cycles, providing the temporal environmental variation necessary to

favor dormancy as a life history strategy (Cohen, 1966; Malik & Smith, 2008). These freeze-

588   thaw cycles could produce boom-and-bust periods of population growth that would leave a

generation-time effect on the rate of molecular evolution, where lineages closer to the poles

590   spend more time on average in a dormant state and show a reduced rate of molecular evolution.

If so, this might result in latitudinal patterns of diversification, which has been a long-standing

592   pattern of diversity in a range of biological systems. However, many microorganisms within the

permafrost may not be completely dormant. For example, psychrophilic taxa isolated from the

594   permafrost are capable of genome replication at temperatures as low as -20 °C (Tuorto et al.,

2014) and evidence of DNA repair has been found in taxa that have persisted in the permafrost

596   for hundreds of thousands to millions of years (Johnson et al., 2007; Dieser et al., 2013). Unlike

the deep biosphere, an appreciable fraction of the microorganisms within the permafrost have

598   likely adapted to grow and reproduce in their environment. However, even with reproduction in

the permafrost, a generation-time effect would still be possible if generation time varied

600   seasonally. In addition, evidence of DNA repair does not mean that microorganisms within the

permafrost maintain a high level of metabolic activity. The energetic cost of DNA repair is low

602   (see section on mutation), suggesting that it is possible for cells within the permafrost to maintain

genome integrity in an otherwise metabolically inactive state. However, as the average global

604   temperature increases due to human-induced climate change, the range of the permafrost will

recede (Seneviratne et al., 2012). This rapid directional shift in Earth's climate will likely impose

606   a strong selective pressure against dormancy as a life history strategy and favoring sustained

growth and reproduction as seasonal environmental variation declines.

 Box 2: Dormancy and the evolution of infectious diseases

Pathogens exist in environments that fluctuate rapidly owing to the immune response of the host and the processes of infection and transmission. Microorganisms often experience

periods of high-growth during the early stages of infection and low-growth in the later stages of infection and when they disperse outside of the host environment (Oliver et al., 1995). These

temporal fluctuations in growth conditions suggest that dormancy would be favored by selection as a life history strategy. For example, *Yersinia pestis*, the causative agent of the plague, often

enters a dormant stage that allows it to persist between infections (Pawlowski et al., 2011). This ability to persist is likely an adaptation, as *Y. pestis* contains several highly conserved genes that

can contribute towards its survival long after the death of the host (Easterday et al., 2012). There is also evidence that dormancy can contribute towards the survival of a pathogen within the host.

Approximately one-third of the world's population is infected with "latent" (i.e., dormant) strains of *Mycobacterium tuberculosis* (Dye et al., 1999), the causative agent of tuberculosis that kills

over 1 million people per year (GBD, 2016). In addition to preventing population extinction, the ability to enter a dormant state would decrease the temporal variation in the size of the

population, maintaining standing genetic variation that could contribute towards adaptive evolution in a new host. If dormancy is as common as these examples suggest, then the ability to

enter and exit a dormant state could be a major feature in the evolutionary dynamics of pathogenic microorganisms.

Dormancy may also contribute to the evolution of antibiotic resistant pathogens. The evolution of disease-causing bacteria that are resistant to the most commonly used antibiotics is a

major global health care concern (WHO, 2014). Methicillin-resistant *Staphylococcus aureus*

infections alone kill more than 11,000 Americans each year (Gross, 2013; Golkar et al., 2014).

642   Antibiotic resistant infections in general are estimated to kill 10 million people by 2050 (O'Neill,

2014). Dormancy can contribute towards the severity of these infections, as bacterial cells can

644   often survive antibiotic treatment by remaining in a dormant state, contributing towards chronic

infections within hosts (Wood et al., 2013; Zhang, 2014). Natural selection can optimize the

646   length of time that bacteria remain dormant under antibiotic treatment (Fridman et al., 2014),

suggesting that dormancy contributes towards long-term geometric fitness as a life history

648   strategy. In addition to preventing cell death, bacteria that survive antibiotic treatment can evolve

antibiotic resistance, simply because they are alive long enough to acquire a beneficial mutation

650   (Levin-Reisman et al., 2017). As the annual number of deaths due to the persistent infection of

antibiotic resistant bacteria continues to increase (WHO, 2014), it will be necessary to develop

652   treatments and strategies to remove reservoirs of dormant bacteria. One potential strategy for

targeting pathogens while minimizing the risk of antibiotic resistance is to resuscitate dormant

654   cells alongside a course of antibiotics. The development of treatments designed to resuscitate

dormant pathogenic bacteria is in the early stages, but holds promise. For example, in sputum

656   samples of individuals infected with *M. tuberculosis*, 80–99.99% of cells were only detectable

after being resuscitated with Rpf (Mukamolova et al., 2010). However, because a large fraction

658   of infections are acquired in hospitals (Klevens et al., 2007), it will be necessary to develop

strategies to remove dormant disease-causing bacteria in buildings (i.e., the built environment) as

660   well as in infected hosts. Recent work on characterizing the built environment suggests that the

ability to persist is the primary trait that shapes the composition of indoor microbial communities

662   (Gibbons et al., 2015; Gibbons, 2016), a trait found in many disease-causing bacteria. Combating

the emergence of antibiotic resistant bacteria and the role of dormancy as an adaptive trait will

664   require re-examining the architecture and materials used to construct the buildings where

        infection is treated as well as infection treatment itself.

ACKNOWLEDGMENTS

688   We acknowledge MG Behringer, J Davis, V Kuo, KJ Locey, RW Moger-Reischer, and

NI Wisnoski for feedback on earlier versions of the manuscript. BK Lehmkuhl and E Polezhaeva

690   provided technical assistance. This work was supported by National Science Foundation

Dimensions of Biodiversity Grant 1442246 (JTL) and US Army Research Office Grant

692   W911NF-14-1-0411 (JTL). The data and code for the simulations used in this study can be found

in a public GitHub repository (https://github.com/LennonLab/EvoDorm).

AUTHOR CONTRIBUTIONS

696   WRS and JTL conceived of the ideas for the paper; WRS created and executed the

simulations; WRS and JTL wrote the paper.

700   **Acknowledgements**

We acknowledge MG Behringer, J Davis, and NI Wisnoski for feedback on

702   earlier versions of the manuscript This work was supported by National Science Foundation

Dimensions of Biodiversity Grant 1442246 (JTL) and US Army Research Office Grant

704   W911NF-14-1-0411 (JTL). The data and code for the simulations used in this study can be found

in a public GitHub repository (https://github.com/LennonLab/EvoDorm).

708   **REFERENCES**

Aanderud, Z. T., Jones, S. E., Fierer, N., & Lennon, J. T. (2015). Resuscitation of the rare

710   biosphere contributes to pulses of ecosystem activity. *Frontiers in Microbiology*, 6, 1–11.

Akiyama, T., Williamson, K. S., Schaefer, R., Pratt, S., Chang, C. B., & Franklin, M. J.

712   (2017). Resuscitation of *Pseudomonas aeruginosa* from dormancy requires hibernation

promoting factor (PA4463) for ribosome preservation. *Proceedings of the National*

714   *Academy of Sciences, USA* 114, 3204–3209.

Alvarez, C. R., Alvarez. R., Grigera, S., Lavado. R. S. (1998). Associations between organic

716   matter fractions and the active soil microbial biomass. *Soil Biology. Biochemistry*, 30,

767–773.

718   Ayati, B.P., & Klapper, I. (2012). Models of microbial dormancy in biofilms and

planktonic cultures. *Communications in Mathematical Sciences,* 10, 493–511

720   Bale, M. J., Bennett, P. M., Beringer, J. E., and Hinton, M. (1993). The survival of bacteria

exposed to desiccation on surfaces associated with farm buildings. Jo*urnal of Applied*

722   *Bacteriology*, 75, 519-528.

Baumdicker, F., Hess, H. R., & Pfaffelhuber, P. (2012). The Infinitely Many Genes

724   Model for the Distributed Genome of Bacteria. *Genome Biology and Evolution,* 4, 443–

456.

726   Bazzaz, F. A. (1979). Physiological ecology of plant succession. *Annual Review of Ecology,*

Evolution, and Systematics, 10, 351–371.

728   Bendall, M. L., Stevens, S. L., Chan, L. K., Malfatti, S., Schwientek, P., Tremblay, J., …

Malmstrom, R. R. (2016). Genome-wide selective sweeps and gene-specific sweeps in

730   natural bacterial populations. *The ISME Journal,* 10, 1589–1601.

Berg, O. G., & Kurland., C. G. (2002). Evolution of Microbial Genomes: Sequence

732   Acquisition and Loss. *Molecular Biology and Evolution,* 19, 2265–2276.

Blagodatskaya, E., & Kuzyakov, Y. (2013). Active microorganisms in soil: critical review of

734   estimation criteria and approaches. *Soil Biology Biochemistry*, 67, 192–211.

Blath, J., González-Casanova, A., Eldon, B., Kurt, N., & Wilke-Berenguer, M. (2015).

736   Genetic Variability Under the Seedbank Coalescent. *Genetics,* 200, 921–934.

Blath, J., González-Casanova, A., Kurt, N., & Wilke-Berenguer, M. (2016). A new

738   coalescent for seed-bank models. *Annals of Applied Probability,* 26, 857-891

Bradshaw, W. E., Armbruster, P. A., & Holzapfel, C. M. (1998). Fitness consequences of

740   hibernal diapause in the pitcher-plant mosquito, *Wyeomyia smithii. Ecology,* 79, 1458–

1462.

742   Brown, J. S., & Venable, D. L. (1986). Evolutionary ecology of seed-bank annuals in

temporally varying environments. *The American Naturalist,* 127, 31–47.

744   Cáceres, C. E., & Tessier, A. J. (2004). To sink or swim: Variable diapause strategies

among *Daphnia* species. *Limnology and Oceanography,* 49, 1333–1340.

746   Cano, R. J., & Borucki, M. K. (1995). Revival and identification of bacterial spores in

25- to 40-million-year-old Dominican amber. *Science,* 268, 1060-1064.

748   Chesson, P. (1994). Multispecies competition in variable environments. *Theoretical*

population biology, 45, 227–276.

750   Chesson, P. L., & Warner, R. R. (1981). Environmental variability promotes coexistence

in lottery competitive systems. *The American Naturalist,* 117, 923–943.

752   Cohen, D. (1966). Optimizing reproduction in a randomly varying environment. *Journal*

        of Theoretical Biology, 12, 119–129.

754   Dawes, I. W., & Thornley, J. H. M. (1970). Sporulation in *Bacillus subtilis*. Theoretical

        and experimental studies in continuous culture systems. *Journal of General*

756   *Microbiology,* 62, 49–66.

        Daubin, V., & Ochman, H. (2004). Bacterial genomes as new gene homes: The genealogy of

758   ORFans in *E. coli*. *Genome Research,* 14, 1036–1042.

        de Casas, R. R., Donohue, K., Venable, D. L., & Cheptou, P. (2015). Gene-flow through

760   space and time: dispersal, dormancy and adaptation to changing environments.

        Evolutionary Ecology, 29, 813–831.

762   Desai, M. M., Fisher, D. S., & Murray, A. W. (2007). The speed of evolution and

        maintenance of variation in asexual populations. *Current Biology,* 17, 385–394.

764   Dieser, M., Battista, J. E., & Christner, B. C. (2013). DNA double-strand break repair at −

        15 C. *Applied and Environmental Microbiology,* 79. 7662–7668.

766   Dixit, P., Pang, T. Y., & Maslov, S. (2016). Recombination-driven genome evolution and

        stability of bacterial species. doi:10.1101/067942

768   Drummond, D. A., & Wilke, C. O. (2008). Mistranslation-induced protein misfolding as a

        dominant constraint on coding-sequence evolution. *Cell*, 134, 341-352.

770   Dworkin, J. & Shah, I. M. (2010). Exit from dormancy in microbial organisms. Nature Reviews

        Microbiology, 8, 890–896.

772   Dye, C., Scheele, S., Dolin, P., Pathania, V., Raviglione, M. C. (1999). Consensus statement.

Global burden of tuberculosis: estimated incidence, prevalence, and mortality by country.

774   WHO global surveillance and monitoring project. *The Journal of the American Medical*

Association, 282, 677–686.

776 Easterday, W. R., Kausrud, K. L., Star, B., Heier, L., Haley, B. J., Ageyev, V., …

Stenseth, N. C. (2012). An additional step in the transmission of *Yersinia pestis*? *The*

778   *ISME Journal,* 6, 231–236.

Epstein, S. S. (2009). Microbial awakenings. *Nature*, 457, 1083–1083.

780 Eriksson, A., Fernstrom, P., Mehlig, B., and Sagitov, S. (2008). An Accurate Model for

Genetic Hitchhiking. *Genetics,* 178, 439–451.

782 Felsenstein, J. (1974). The evolutionary advantage of recombination. *Genetics,* 78, 737–

756.

784 Finkel, S. E. (2006). Long-term survival during stationary phase: evolution and the GASP

phenotype. *Nature Reviews Microbiology,* 4, 113-120.

786 Ford, C. B., Shah, R. R., Maeda, M. K., Gagneux, S., Murray, M.B., Cohen, T., …

Fortune, S. M. (2013). *Mycobacterium tuberculosis* mutation rate estimates from

788   different lineages predict substantial differences in the emergence of drug-resistant

tuberculosis. *Nature Genetics,* 45, 784–790.

790 Ford, C. B., Lin, P. L., Chase, M. R., Shah, R. R., Iartchouk, O., Galagan, J., … Fortune,

S. M. (2011). Use of whole genome sequencing to estimate the mutation rate of

792   *Mycobacterium tuberculosis* during latent infection. *Nature Genetics,* 43, 482–486.

Foster, P. L. (2007). Stress-induced mutagenesis in bacteria. *Critical Reviews in*

794   *Biochemistry and Molecular Biology,* 42, 373-397.

Fridman, O., Goldberg, A., Ronin, I., Shoresh, N., Balaban, N. Q. (2014). Optimization of lag

796 time underlies antibiotic tolerance in evolved bacterial populations. *Nature,* 513, 418–

421.

798 Gabriel, W., Lynch, M., & Burger, R. (1993). Muller's Ratchet and mutational

meltdowns. *Evolution,* 47, 1744-1757.

800 Galhardo R. S., Hastings, P. J., & Rosenberg, S. M. (2007). Mutation as a stress response

and the regulation of evolvability. C*ritical Reviews in Biochemistry and*

802 *Molecular Biology*, 42, 399-435.

Galperin, M. Y., Mekhedov, S. L. , Puigbo, P., Smirnov, S. , Wolf, Y. I. , & Rigden, D. J.

804 (2012). Genomic determinants of sporulation in *Bacilli* and *Clostridia*: towards the

minimal set of sporulation-specific genes. *Environmental Microbiology,* 14, 2870–2890.

806 GBD 2015 Mortality and Causes of Death Collaborators (2016). Global, regional, and national

life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death,

808 1980-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet,*

388, 1459–1544.

810 Gerrish, P. J., Colato, A. & Sniegowski, P. D. (2013). Genomic mutation rates that

neutralize adaptive evolution and natural selection. *Journal of The Royal Society*

812 *Interface,* 10, 20130329.

Gerrish, P. J., & Lenski, R. E. (1998). The fate of competing beneficial mutations in an

814 asexual population. *Genetica,* 102/103, 127–144.

Gibbons, S. M. (2016). The built environment is a microbial wasteland. *mSystems,* 1,

816 e00033-16.

Gibbons S. M., Schwartz, T., Fouquier, J., Mitchell, M., Sangwan, N., Gilbert, J. A., &

818 Kelley, S. T. (2015). Ecological succession and viability of human-associated

microbiota on restroom surfaces. *Applied Environmental Microbiology,* 81, 765–

820   773.

Gillespie, J. H. (2000). Genetic drift in an infinite population. The pseudohitchhiking

822   model. *Genetics,* 155, 909-919.

Golkar Z, Bagazra, O., Pace, D. G. (2014). Bacteriophage therapy: a potential solution for the

824   antibiotic resistance crisis. *Journal of infection in developing countries*, 8, 129–136.

Gong, C., Bongiorno, P., Martins, A., Stephanou, N. C., Zhu, H., Shuman, S., &

826   Glickman, M. S. (2005). Mechanism of nonhomologous end-joining in mycobacteria: a

low-fidelity repair system driven by Ku, ligase D and ligase C. *Nature Structural &*

828   *Molecular Biology,* 12, 304–312.

González-Casanova, A., Aguirre-von-Wobeser, E., Espín, G., Servín-González, L., Kurt,

830   N., Spanò, D., Blath, J., & Soberón-Chávez, G. (2014). Strong seed-bank effects

in bacterial evolution. *Journal of Theoretical Biology,* 356, 62–70.

832   Graham, D. E., Wallenstein, W. D., Vishnivetskaya, T. A., Waldrop, M. P., Phelps, T. J.,

Pfiffner, S. M., … Jansson, J. K. (2012). Microbes in thawing permafrost: the

834   unknown variable in the climate change equation. *The ISME Journal,* 6, 709–712.

Greenblatt, C. L., Baum, J., Klein, B. Y., Nachshon, S., Koltunov, V., & Cano, R. J.

836   (2004). *Micrococcus luteus* - Survival in amber. *Microbial Ecology,* 48, 120-127.

Gross, M. (2013). Antibiotics in crisis. *Current Biology*, 23, 1063–1065.

838   Guppy, M., & Withers, P. (1999). Metabolic depression in animals: physiological

perspectives and biochemical generalizations. *Biological reviews of the*

840   *Cambridge Philosophical Society,* 74, 1–40.

Hairston Jr., N. G. & De Stasio Jr., B. T. (1988). Rate of evolution slowed by a dormant

842   propagule pool. *Nature,* 336, 239–242.

Hairston, N. G., Lampert, W., Cáceres, C. E., Holtmeier, C. L., Weider, L. J., Gaedke, U.,

844   Fischer, J. M. … Post, D. M. (1999). Lake ecosystems: Rapid evolution revealed by

dormant eggs. *Nature,* 401, 446–446.

846 Hoehler, T. M., & Jørgensen, B. B. (2013). Microbial life under extreme energy

limitation. *Nature Reviews Microbiology,* 11, 83–94.

848 Hollander, F. den., & Pederzani, G. (2017). Multi-colony Wright-Fisher with seed-bank.

Indagationes Mathematicae, doi:10.1016/j.indag.2017.02.002

850 Horneck, G., Bucker, H., & Reitz, G. (1994). Long-term survival of bacterial spores in

space. *Advances in Space Research,* 14, 41–45.

852 Horneck, G., Moeller, R., Cadet, J., Douki, T., Mancinelli, R. L., Nicholson, W. L., …

Venkateswaran, K. J. (2012). Resistance of bacterial endospores to outer space for

854   planetary protection purposes—Experiment PROTECT of the EXPOSE-E

mission. *Astrobiology,* 12, 445–456.

856 Ishiguro, K., Washio, J., Sasaki, K., & Takahashi, N. (2015). Real-time monitoring of the

metabolic activity of periodontopathic bacteria. *Journal of Microbiological Methods*,

858   115, 22–26.

Johnson, S. S., Hebsgaard, M. B., Christensen, T. R., Mastepanov, M., Nielsen, R.,

860   , K., Brand, T., … Willerslev, E. (2007). Ancient bacteria show evidence of DNA

repair. *Proceedings of the National Academy of Sciences, USA,* 104, 4401-14405.

862 Jones, S. E. & Lennon, J. T. (2010). Dormancy contributes to the maintenance of

microbial diversity. *Proceedings of the National Academy of Sciences, USA,* 107,

864   5881–5886.

Jørgensen, B. B. (2012). Shrinking majority of the deep biosphere. *Proceedings of the*

866   *National Academy of Sciences, USA,* 109, 15976–15977.

Jørgensen, B. B., & Marshall, I. P. G. (2016). Slow Microbial Life in the Seabed. *Annual*

868   *Review of Marine Science,* 8:311–332.

Kaj, I., Krone, S. M., & Lascoux, M. (2001). Coalescent theory for seed bank models.

870   *Journal of Applied Probability,* 38, 285–300.

Kalamees, R. & Zobel, M. (2002). The role of the seed bank in gap regeneration in a calcareous

872   grassland community. *Ecology*, 83, 1017–1025.

Kalisz, S. & McPeek, M. A. (1993). Extinction dynamics, population growth and seed banks.

874   *Oecologia*, 95, 314-320.

Kallmeyer, J., Pockalny, R., Adhikari, R. R., Smith, D. C. & D'Hondt, S. (2012). Global

876   distribution of microbial abundance and biomass in subseafloor sediment.

           Proceedings of the National Academy of Sciences, USA, 109, 16213–16216.

878   Kaprelyants, A. S., & Kell, D. B. (1993). Dormancy in Stationary-Phase Cultures of

           Micrococcus luteus: Flow Cytometric Analysis of Starvation and Resuscitation. *Applied*

880   *and Environmental Microbiology*, 59, 3187–96.

Keep, N. H., Ward, J. M., Cohen-Gonsaud, M., & Henderson, B. (2006). Wake up!

882   Peptidoglycan lysis and bacterial non-growth states. *Trends in Microbiology,* 14, 271–

           276.

884   Kempes, C. P., van Bodegom, P. M., Wolpert, D., Libby, E., Amend, J., & Hoehler, T.

           (2017). Drivers of Bacterial Maintenance and Minimal Energy Requirements.

886   *Frontiers in Microbiology,* 8, 1–10.

Keynan, A., Evanchik, Z., Halvorson, H. O., & Hastings, J. W. (1964). Activation of bacterial

888   endospores. *Journal of Bacteriology,* 88, 313–8.

Kimura, M. (1969). The number of heterozygous nucleotide sites maintained in a finite

890   population due to steady flux of mutations *Genetics,* 61, 893-903.

Klevens, R. M., Edwards, J. R., Richards Jr., C. L., Horan, T. C., Gaynes, R. P., Pollock, D. A.,

892   & Cardo, D. M. (2007). Estimating health care-associated infections and deaths in U.S.

                 hospitals, 2002. *Public Health Reports*, 122, 160–166.

894   Koopmann, B., Müller, J., Tellier, A., & Živković, D. (2017). Fisher–Wright model with

                 deterministic seed bank and selection. *Theoretical Population Biology,* 114, 29–39.

896   Kunkel, T. A. (2004). DNA replication fidelity. *The Journal of Biological Chemistry,*

                 279, 16895–16898.

898   Lapierre, P., & Gogarten, J. P. (2009). Estimating the size of the bacterial pan-genome.

                 Trends in Genetics, 25, 107–110.

900   Lassalle, F., Périan, S., Bataillon, T., Nesme, X., Duret, L. & Daubin, V. (2015). GC-

                 content evolution in bacterial genomes: The biased gene conversion hypothesis expands.

902   *PLOS Genetics,* 11, e1004941.

Lenhart, J.S., Schroeder, J. W., Walsh, B. W., & Simmons, L. A. (2012). DNA repair and

904   genome maintenance in *Bacillus subtilis*. *Microbiology and Molecular Biology Reviews,*

                 76, 530–564.

906   Lennon, J. T., & Jones, S. E. (2011). Microbial seed banks: the ecological and

                 evolutionary implications of dormancy. *Nature Reviews Microbiology,* 9, 119–30.

908   Levin, D. A. (1990). The seed bank as a source of genetic novelty in plants. *The*

                 American Naturalist, 135, 563–572.

910   Levin-reisman, I., Ronin, I., Gefen, O., Braniss, I., Shoresh, N., & Balaban, N. Q. (2017).

Antibiotic tolerance facilitates the evolution of resistance. *Science*, 355, 1–10.

912   Locey, K. J., Fisk, M. C., & Lennon, J. T. (2017). Microscale insight into microbial seed banks.

Frontiers in Microbiology, 7, 2040

914   Lomstein, B. A., Langerhuus, A. T., D'Hondt, S., Jørgensen, B. B., & Spivack, A. J.

(2012). Endospore abundance, microbial growth and necromass turnover in deep sub-

916   seafloor sediment. *Nature,* 484, 101–104.

Lynch, M. (2007). The Origins of Genome Architecture. Sunderland, MA: Sinauer

918   Associates.

Lynch, M., Ackerman, M. S., Gout, J.-F., Long, H., Sung, W., Thomas, W. K., & Foster,

920   P. L. (2016). Genetic drift, selection and the evolution of the mutation rate.

Nature Reviews Genetics, 17, 704–714.

922   Lynch, M., Blanchard, J., Houle, D., Kibota, T., Schultz, S., Vassilieva, L., & Willis, J. (1999).

Perspective: Spontaneous Deleterious Mutation. *Evolution*, 53, 645-663.

924   Lynch, M., Bürger, R., Butcher, D., & Gabriel, W. (1993). The mutational meltdown in

asexual populations. *The Journal of Heredity,* 84, 339–344.

926   Lynch, M., & Marinov, G. K. (2015). The bioenergetic costs of a gene. *Proceedings of*

the National Academy of Sciences, USA, 112, 15690-15695.

928   Maddamsetti, R., Hatcher, P. J., Green, A. G., Williams, B. L., Marks, D. S., & Lenski,

R. E. (2017). Core genes evolve rapidly in the long-term evolution experiment

930   with *Escherichia coli. Genome Biology and Evolution,* 9, 1072-1083..

Malik, T. & Smith, H. L (2008). Does dormancy increase fitness of bacterial populations

932   in time-varying environments? *Bulletin of Mathematical Biology*, 70, 1140–1162.

Marin, J., Battistuzzi, F. U., Brown, A. C., & Hedges, S. B. (2017). The timetree of

934   prokaryotes: New insights into their evolution and speciation. *Molecular Biology and Evolution,* 34, 437-446.

936   Marks, P. L. (1974). The Role of Pin Cherry (Prunus pensylvanica L.) in the Maintenance of Stability in Northern Hardwood Ecosystems. *Ecological Monographs*, 44, 73–88.

938   Maughan, H., Masel, J., Birky, C. W., & Nicholson, W. L. (2007). The roles of mutation accumulation and selection in loss of sporulation in experimental populations of *Bacillus*

940   *subtilis. Genetics,* 177, 937–948.

Medini, D., Donati, C., Tettelin, H., Masignani, V., & Rappuoli, R. (2005). The microbial

942   pan-genome. *Current Opinion in Genetics & Development,* 15, 589–594.

Mira, A., Pushker, R., Legault, B. A., Moreira, D., & Rodríguez-Valera, F. (2004).

Evolutionary relationships of *Fusobacterium nucleatum* based on phylogenetic analysis and comparative genomics. *BMC Evolutionary Biology,* 4, 50.

Mizrahi, V., & Andersen, S. J. (1998). DNA repair in *Mycobacterium tuberculosis*. What have we learnt from the genome sequence? *Molecular Microbiology,* 29, 1331–1339.

Moeller, R., Setlow, P., Reitz, G., & Nicholson, W. L. (2009). Roles of small, acid-soluble spore proteins and core water content in survival of *Bacillus subtilis*

spores exposed to environmental solar UV radiation. *Applied and Environmental Microbiology*, 75, 5202–5208.

Moeller, R., Raguse, M., Reitz, G., Okayasu, R., Li, Z., Klein, S. … Nicholson, W. L. (2014). Resistance of *Bacillus subtilis* spore DNA to lethal ionizing radiation damage

relies primarily on spore core components and DNA repair, with minor effects of oxygen radical detoxification. *Applied and Environmental Microbiology,* 80, 104–109.

Mukamolova, G. V., Kaprelyants, A. S., Young, D. I., Young, M. & Kell, D. B. A.

(1998). A bacterial cytokine. *Proceedings of the National Academy of Sciences, USA,* 95,

958   8916–8921.

Mukamolova, G. V., Turapov, O., Malkin, J., Woltmann, G., & Barer, M. R. (2010).

960   Resuscitation-promoting factors reveal an occult population of tubercle Bacilli in

Sputum. *American Journal of Respiratory and Critical Care Medicine*, 181, 174–80.

962 Muller, H. J. (1964). The relation of recombination to mutational advance. *Mutation*

Research, 106, 2–9.

964 Müller, A. L., de Rezende, J. R., Hubert, C. R. J., Kjeldsen, K. U., Lagkouvardos, I.,

Berry, D., … Loy, A. (2014). Endospores of thermophilic bacteria as tracers of

966   microbial dispersal by ocean currents. *The ISME Journal,* 8, 1153–1165.

Nicholson, W.L., Munakata, N., Horneck, G., Melosh, H. J., & Setlow, P. (2000).

968   Resistance of *Bacillus* endospores to extreme terrestrial and extraterrestrial environments.

Microbiology and Molecular Biology Reviews, 64, 548–572.

970 Nicholson, W.L., Schuerger, A. C., & Setlow, P. (2005). The solar UV environment and

bacterial spore UV resistance: considerations for Earth-to-Mars transport by natural

972   processes and human spaceflight. *Mutation Research,* 571, 249–264.

Nunney, L. (2002). The effective size of annual plant populations: The interaction of a

974   seed bank with fluctuating population size in maintaining genetic variation. *The*

American Naturalist, 160, 195–204.

976 Ochman, H., Lawrence, J. G., & Groisman, E. A. (2000). Lateral gene transfer and the

nature of bacterial innovation. *Nature*, 405, 299–304.

978 Oliver, J. D. (1995). The viable but non-culturable state in the human pathogen *Vibrio*

vulnificus. *FEMS Microbiology Letters,* 133, 203–208.

980   Olivieri, I. (2001). The evolution of seed heteromorphism in a metapopulation:

interactions between dispersal and dormancy. *In* "Integrating Ecology and

982   Evolution in a Spatial Context" (J. Silvertown and J. Antonovics eds.), pp. 245-268.

Oxford: Cambridge University Press.

984   O'Neill J. (2014). Review on Antimicrobial Resistance Antimicrobial Resistance: Tackling a

crisis for the health and wealth of nations. London: Review on Antimicrobial Resistance.

986   Park, S.-C., Simon, D., & Krug, J. (2010). The speed of evolution in large asexual

populations. *Journal of Statistical Physics,* 138, 381-410.

988   Pawlowski, D. R., Metzger, D. J., Raslawsky, A., Howlett, A., Siebert, G., Karalus, R. J.

… Whitehouse, C. A. (2011). Entry of *Yersinia pestis* into the viable but nonculturable

990   state in a low-temperature tap water microcosm. *PLoS ONE,* 6, e17585.

Plotkin, J. B., & Kudla, G. (2011). Synonymous but not the same: the causes and

992   consequences of codon bias. *Nature Reviews Genetics,* 12, 32–42.

Ramirez-Guadiana, F. H., Barraza-Salas, M., Ramirez-Ramirez, N., Ortiz-Cortes, M.,

994   Setlow, P., & Pedraza-Reyes, M. (2012). Alternative excision repair of ultraviolet B- and

C-induced DNA damage in dormant and developing spores of *Bacillus subtilis*. *Journal*

996   *of Bacteriology,* 194, 6096–6104.

Rittershaus, E. S. C., Baek, S., & Sassetti, C. M. (2013). The normalcy of dormancy:

998   common themes in microbial quiescence. *Cell Host & Microbe* 13, 643–651.

Rosenberg, S. M. (2001). Evolving responsively: adaptive mutation. *Nature Reviews*

1000   *Genetics,* 2, 504-515.

Seneviratne, S. I. Nicholls, N., Easterling, D., Goodess, C. M., Kanae, S., … Zhang, X.

1002   (2012). Changes in climate extremes and their impacts on the natural physical

environment: an overview of the IPCC SREX report. *EGU General Assembly Conference*

1004   *Abstracts,* 14, 12566.

Setlow, P. (1992). I will survive: protecting and repairing spore DNA. *Journal of*

1006   *Bacteriology,* 174, 2737–2741.

Setlow, P. (2006). Spores of *Bacillus subtilis*: their resistance to and killing by radiation,

1008   heat and chemicals. *Journal of Applied Microbiology,* 101, 514–525.

Sharp, P. M., Bailes, E., Grocock, R. J., Peden, J. F., & Sockett, R. E. (2005). Variation in the

1010   strength of selected codon usage bias among bacteria. *Nucleic Acids Research*. 33, 1141-

1153.

1012   Sharp, P. M., & Li, W.-H. (1987). The rate of synonymous substitution in enterobacterial genes

is inversely related to codon usage bias. *Molecular Biology and Evolution*, 4, 222-230.

1014   Simons, A. M. (2009). Fluctuating natural selection accounts for the evolution of

diversification bet hedging. *Proceedings of the Royal Society B*: *Biological Sciences*, 276,

1016   1987–1992.

Smith, J. M., & Haigh, J. (1974). The hitch-hiking effect of a favourable gene. *Genetical*

1018   *Research*, 23, 23–35.

Starnawski, P., Bataillon, T., Ettema, T. J. Jochum, L. M., Schreiber, L., Chen, X. …

1020   Kjeldsen, K. U. (2017). Microbial community assembly and evolution in

subseafloor sediment. *Proceedings of the National Academy of Sciences, USA,* 114,

1022   2940–2945.

Tan, I. S., & Ramamurthi, K. S. (2014). Spore formation in *Bacillus subtilis*. *Environmental*

1024   *Microbiology Reports*, 6, 212–225.

Tellier, A., Laurent, S. J. Y., Lainer, H., Pavlidis, P., & Stephan, W. (2011). Inference of

1026   seed bank parameters in two wild tomato species using ecological and genetic

data. *Proceedings of the National Academy of Sciences, USA,* 108, 17052–17057.

1028 Templeton, A. R., & Levin, D. A. (1979). Evolutionary consequences of seed pools. *The*

American Naturalist, 114, 232–249.

1030 Tenaillon, O., Barrick, J. E., Ribeck, N., Deatherage, D. E., Blanchard, J. L., Dasgupta.

A., … Lenski, R. E. (2016). Tempo and mode of genome evolution in a 50,000-

1032 generation experiment. *Nature,* 536, 165–170.

Tuorto, S. J., Darias, P., McGuinness, L. R., Panikov, N., Zhang, T., Häggblom, M. M.,

1034 & Kerkhof, L. J. (2014). Bacterial genome replication at subzero temperatures in

permafrost. *The ISME Journal,* 8, 139–149.

1036 van Bodegom, P. (2007). Microbial maintenance: a critical review on its quantification.

Microbial Ecology, 53, 513–523.

1038 Venable, D. L., & Brown, J. S. (1988). The selective interactions of dispersal, dormancy,

and seed size as adaptations for reducing risk in variable environments. *The American*

1040 *Naturalist,* 131, 360–384.

Venable, D. L., & Lawlor, L. (1980). Delayed germination and dispersal in desert

1042 annuals: Escape in space and time. *Oecologia*, 46, 272–282.

Vitalis, R., Glémin, S., & Olivieri, I. (2004). When genes go to sleep: the population

1044 genetic consequences of seed dormancy and monocarpic perenniality. *The*

American Naturalist, 163, 295–311.

1046 Votyakova, T., Kaprelyants, A., & Kell, D. B. (1994). Influence of viable cells on the

resuscitation of dormant cells in *Micrococcus luteus* cultures held in an extended

1048   stationary phase: the population effect. *Applied and Environmental Microbiology*, 60,

3284–3291.

1050   Vreeland, R. H., Rosenzweig, W. D., & Powers, D. W. (2000). Isolation of a 250 million-

year-old halotolerant bacterium from a primary salt crystal. *Nature,* 407, 897–900.

1052   Walker, H. H., & Winslow, C.-E. A. (1932). Metabolic Activity of the Bacterial Cell at Various

Phases of the Population Cycle. *Journal of Bacteriology*, 24, 209–241.

1054   Wang, G. S., Mayes, M. A., Gu, L. H., and Schadt, C. W. (2014). Representation of dormant and

active microbial dynamics for ecosystem modeling. *PLoS ONE*, 9, 10.

1056   Weinbauer, M., & Rassoulzadegan, F. (2007). Extinction of microbes: evidence and potential

consequences. *Endangered Species Research*, 3, 205–215.

1058   Weller, C., & Wu, M. (2015). A generation-time effect on the rate of molecular evolution

in bacteria. *Evolution,* 69, 643–652.

1060   Willis, C. G., Baskin, C. C., Baskin, J. M., Auld, J. R., Venable, D. L., Cavender-Bares,

J., Donohue, K., … NESCent Germination Working Group. (2014). The evolution of

1062   seed dormancy: environmental cues, evolutionary hubs, and diversification of the seed

plants. *New Phytologist,* 203, 300–309.

1064   Witkin, E. M. (1976). Ultraviolet mutagenesis and inducible DNA repair in *Escherichia*

coli. Bacteriological Reviews, 40, 869-907.

1066   Wood, T. K., Knabel, S. J., & Kwan, B. W. (2013). Bacterial persister cell formation and

dormancy. *Applied and Environmental Microbiology,* 79, 7116–7121.

1068   World Health Organization. (2014). Antimicrobial resistance: 2014 global report on surveillance.

World Health Organization. Chicago

1070   Zhang, Y. (2014). Persisters, persistent infections and the Yin–Yang model. *Emerging*

Microbes & Infections, 3, e3.

1072 Zhaxybayeva, O., Swithers, K. S., Lapierre, P., Fournier, G. P., Bickhart, D. M., DeBoy,

    R. T. … Noll, K. M. (2009). On the chimeric nature, thermophilic origin, and

1074   phylogenetic placement of the *Thermotogales*. *Proceedings of the National Academy of*

    Sciences, USA, 106, 5865–5870.

1076 Živković, D., & Tellier, A. (2012). Germ banks affect the inference of past demographic

    events. *Molecular Ecology,* 21, 5434–5446.

1078

1080

1082

1084

1086

1088   **Figure 1:** Transitioning in and out of a seed bank is analogous to the migration of individuals

between populations. In **a)** we see a population with a seed bank where individuals switch

1090   between active and dormant states. In **b)** we see two populations where individuals can migrate

between each population. An important difference between **a)** and **b)** is that individuals must

1092   originate in the active pool, because dormant individuals cannot reproduce. While dormant

individuals can die in this model, their death rate is much lower than that of active individuals.

1094   This conceptual model was first presented in an ecological context (Lennon & Jones, 2011) and

later used for the seed bank coalescent (Blath et al., 2015).

1096

Fig. 2: The single cell metabolic activity distribution in a bacterial population at different time-

1098   points. The bacterium used is a strain of *Janthinobacterium*, an aerobic, Gram-negative, soil-

dwelling β-proteobacteria. The blue, yellow, and red histograms represent samples taken at 3, 9,

1100   and 200 hours after the culture was inoculated, respectively. The distribution of metabolic

activity depends on the growth state of the population, where the mode decreases as well as the

1102   shape of the distribution as the population enters a dormant state. During the process of

resuscitation the mode increases corresponding to an increase in growth rate (see Supplemental

1104   Materials).

1106   **Figure 3: a)** The expected level of nucleotide diversity ($\mathbf{E}[\pi]$) for a sample of active individuals

increases with the number of generations that an individual on average spends in the seed bank

1108   for a population at mutation-drift equilibrium with a given population-scaled mutation rate ($\theta =$

$2N\mu$). Here N is the number of active individuals, M is the number of dormant individuals, and

1110 μ is the mutation rate for active individuals, where diversity is estimated as $\mathbf{E}[\pi] = \theta + \theta(\frac{M}{N})$

(Blath et al., 2015, eq. 21). The average number of generations that an individual spends in the

1112   seed bank can be modeled as a geometric distribution, where the probability of exiting the seed

bank each generation is $\frac{c}{M}$ and c is the number of individuals that exit the seed bank each

1114   generation (Blath et al., 2015; 2016). The vertical grey-dashed line represents the point where the

average number of generations that an individual spends in the seed bank is greater than the

1116   number of active individuals in the populations (i.e., the strong seed bank threshold). **b)** The

length of time required to reach mutation-drift equilibrium increases with the average number of

1118   generations that an individual spends in the seed bank. These results are from a simulated

population evolving over time that is subject only to mutation and drift with a constant seed bank

1120   size. In both **a)** and **b)** We assume that dormant individuals do not acquire mutations. See

Supplementary Information for more detail on implementation of model.

1122

Figure 4: a) The average length of time that an individual spends in a dormant state influences

1124   the trajectory of a beneficial allele destined to go to fixation. The horizontal grey line indicates

an allele frequency of 0.5, showing how the beneficial allele first becomes fixed in the active

1126   portion of the population and requires an extended period of time to become fixed in the seed

bank as the average length of time spent in a dormant state increases. **b)** The length of time until

1128   an allele is effectively fixed (T_{fix}) for both active and dormant individuals scales with the

average number of generations that an individual spends in a dormant state. However, T_{fix} for

1130   the active portion of the population decreases in the strong seed bank regime and eventually

approaches the expected time to fixation for an ideal population with the same number of active

1132   individuals and strength of selection (horizontal grey line) (Eriksson et al., 2008, eq. 37). We

believe that this is because the beneficial allele can reach fixation before dormant individuals

1134   with the alternative allele resuscitate, preventing the seed bank from reducing the efficiency of

selection. These results for **a)** and **b)** were obtained from a simulated Moran model of a selective

1136   sweep with a seed bank (see Supplementary Materials for details).

1138   **Figure 5: a)** The evolutionary distance (calculated as the root-to-tip sum on the branch of the

phylogeny) declines as the number of sporulation-associated genes increases among *Firmicutes*

1140   taxa, a common group of bacteria found in soils and hosts. The publically available data

presented here is from the *Firmicutes* phylogeny of conserved genes (Weller & Wu, 2015). **b)**

1142   The evolutionary distance (calculated as the JC69 corrected distance) decreases as the average

length of time that an individual spends in the seed bank increases (see Supplementary

1144   Materials). The lines in both **a)** and **b)** are the slopes from simple linear regression.

1146

1148

1150

1152

1154