1

TITLE: Increased prefrontal activity with aging reflects nonspecific neural responses rather
than compensation

ABBREVIATED TITLE: Nonspecific prefrontal activity in aging

Alexa M. Morcom[1]

Richard N. A. Henson[2], for Cambridge Centre for Ageing and Neuroscience[3]

[1]Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, United
Kingdom

[2]MRC Cognition and Brain Sciences Unit, Cambridge CB2 3EB, United Kingdom

[3]Cambridge Centre for Ageing and Neuroscience (Cam-CAN), University of Cambridge and
MRC Cognition and Brain Sciences Unit, Cambridge CB2 3EB, United Kingdom

Alexa M. Morcom can be contacted at: alexa.morcom@ed.ac.uk

No. of Figures = 3

No. of Tables = 5

No multimedia/ extended data

Abstract

Elevated prefrontal cortex activity is often observed in healthy older adults despite declines in their memory and other cognitive functions. According to one view, this activity reflects a compensatory functional posterior-to-anterior shift, which contributes to maintenance of cognitive performance when posterior cortical function is impaired. Alternatively, the increased prefrontal activity may be less efficient or less specific, owing to structural and neurochemical changes accompanying aging. These accounts are difficult to distinguish on the basis of average activity levels within brain regions. Instead, we used a novel model-based multivariate analysis technique, applied to two independent functional magnetic resonance imaging datasets from an adult-lifespan human sample (N=123 and N=115; approximately half female). Standard analysis replicated the age-related increase in average prefrontal activation, but multivariate tests revealed that this activity did not carry additional information. The results contradict the posterior-to-anterior shift hypothesis, suggesting reduced efficiency or specificity, rather than compensation.

Significance statement

Functional brain imaging studies have often shown increased activity in prefrontal brain regions in older adults. This has been proposed to reflect a compensatory shift to greater reliance on prefrontal cortex, helping to maintain cognitive function. Alternatively, activity may become less specific as people age. This is a key question in the neuroscience of aging. In this study, we used novel tests of how different brain regions contribute to long-and short-term memory. We found increased activity in prefrontal cortex in older adults, but this activity carried less information about memory outcomes than activity in visual regions. These findings are relevant for understanding why cognitive abilities decline with age, suggesting that optimal function depends on successful brain maintenance rather than compensation.

Introduction

It is well established that healthy aging is associated with a decline in cognitive processes like

memory, but mechanistic explanation of this decline is impeded by difficulties in interpreting

the underlying brain changes. Functional magnetic resonance imaging (fMRI) of such

cognitive processes shows striking increases, as well as decreases, in brain activity of older

relative to younger adults. One leading theory – the Posterior-to-Anterior Shift in Aging (PASA)

– states that recruitment of anterior regions like prefrontal cortex (PFC) contributes to

maintenance of cognitive performance when posterior cortical function is impaired (Davis,

Dennis, Daselaar, Fleck, & Cabeza, 2008; Grady, 2012; Park & Reuter-Lorenz, 2009).

Alternatively, age-related increases in PFC activity may reflect reduced efficiency or specificity

of neuronal responses, reflecting primarily age-related functional impairment within PFC

(West, 1996; Glisky et al., 2001; Raz and Rodrigue, 2006; Nyberg, Lövdén, Riklund,

Lindenberger, & Bäckman, 2012; Park et al., 2004). It is difficult to adjudicate between these

theories based on average activity levels within brain regions (Morcom and Johnson, 2015).

We used a novel multivariate approach to directly test predictions of the PASA theory.

With multivariate methods that examine distributed patterns of brain activity over many voxels,

one can ask whether increased anterior activity provides additional information, and whether

this information goes beyond that provided by posterior cortical regions. Such increases in the

information represented by PFC activity would support theories that attribute additional PFC

recruitment to compensatory mechanisms. We used a model-based decoding approach called

multivariate Bayes (MVB) (Friston et al., 2008; Morcom and Friston, 2012; Chadwick et al.,

2014), which estimates the patterns of activity that best predict a target cognitive outcome.

Importantly, MVB allows formal comparison of models comprising different brain regions, such

as PFC, posterior cortex, or their combination.

In this study, we applied MVB to fMRI data from two different paradigms in population-derived,

adult-lifespan samples (N=123 and N=115, 19-88 years; Shafto et al. 2014). In the first, long-

term memory (LTM) experiment, participants were scanned while encoding new memories of

unique pairings of objects and background scenes, and the target cognitive outcome was

whether or not these associations were subsequently remembered (Figure 1a). A previous

behavioral study in an independent sample showed a strong decline in such associative

memory across the adult lifespan (Henson et al., 2016). To test whether findings generalized

across tasks and cognitive domains, as PASA predicts (Davis et al., 2008), we replicated our

findings in a second, visual short-term memory (STM) experiment. In the STM experiment, a

separate sample of participants was scanned while maintaining visual dot patterns online for

a few seconds in the presence of distraction, and the target cognitive outcome was the

increase in the number of patterns to be maintained (i.e., load; Figure 1b). Increases in PFC
activity have frequently been reported in older adults in similar tasks (Grady et al., 1998;
Cabeza et al., 2004), although sometimes activity reductions are found at higher loads
(Cappell et al., 2010).

We defined two regions-of-interest (ROIs): posterior visual cortex (PVC), comprising lateral
occipital and fusiform cortex, and PFC, comprising ventrolateral, dorsolateral, superior and
anterior regions (Fig 2a). These ROIs were based on previous fMRI studies of memory tasks,
and those cited in the context of the PASA theory (Davis et al., 2008; Maillet and Rajah, 2014).

Figure 1. Memory tasks. a and b, long-term memory (LTM) task and c, short-term memory (STM) task.
a, Associative encoding. In the scanned Study phase of the LTM task, participants were asked to make
up a story that linked each object with its background scene (120 trials total). A scene with positive
valence is illustrated. On each trial, the scene was presented for 2 sec, then the object superimposed
for 7.5 sec, finally the screen was blanked for 0.5 sec before the next trial. b, Associative retrieval. At
Test (out of the scanner), each object was presented again, and after a measure of priming, item
memory and background valence memory, participants were asked to verbally describe the scene with
which it was paired at Study. The latter verbal recall was scored as correct or incorrect, which was then
used to classify the trials at Study into "remembered" and "forgotten" (see text for details). The example
illustrates encoding and retrieval of a trial with neutral valence. c, An example trial of STM task with
memory load of 2 items. Trials began with fixation dot for 7 sec. On each trial, three dot displays were
displayed in red, yellow and blue for 500 msec each (250 msec gap). To vary load, the dots in either
one, two or three of the dot displays moved in a consistent direction (the to-be-ignored displays rotated).
After the last display, the screen was blanked for an 8 sec maintenance period. Then the probe display
presented a colored circle to indicate which dot display to recall (red, yellow or blue). Participants had
up to 5 sec to adjust the pointer until the direction matched that of the to-be-remembered display.

Materials and Methods

Experiment 1: Long-term Memory Encoding

Participants

A healthy, population-derived adult lifespan human sample (N=123; 19-88 years; 66 female)
was collected as part of the Cam-CAN study (Shafto et al., 2014). Participants were fluent
English speakers in good physical and mental health. Exclusion criteria included a low Mini
Mental State Examination (MMSE) score (<=24), serious current medical or psychiatric
problems, or poor hearing or vision, as well as standard MRI safety criteria. Two participants
were excluded from fMRI analysis as subsequent memory could not successfully be decoded
from either region of interest (see Multivariate Bayesian decoding). Two further participants
were excluded because of statistical outlier values in the analysis of univariate subsequent
memory effects (see Statistics for criteria). The experiment used a within-participant design,
so all participants received all the task conditions. Therefore, randomization and blinding were
not required. The study was approved by the Cambridgeshire 2 (now East of England—
Cambridge Central) Research Ethics Committee. Participants gave informed written consent.

Materials

Stimuli were 160 pictures of everyday emotionally-neutral objects taken from Smith et al.
(2004). For the study phase, objects were presented within a square yellow background on
one of 120 scenes from the IAPS emotional pictures database (Lang et al., 1997). Scenes
were grouped into 40 per valence (positive, neutral, negative), selected based on a pilot study,
with the same randomized trial order for each valence condition for all participants. To control
for stimulus effects, the 160 objects were divided randomly into 4 sets, and the allocation of
object sets to scene valence rotated across participants in 4 different counterbalances (see
Henson et al., 2016, for further details).

Behavioral procedure

The paradigm is summarized in Figure 1a. The scanned study phase comprised 120 trials,
presented in two 10 min blocks separated by a short break. On each study trial, a background
scene was first presented for 2 sec, and an object then superimposed for 7.5 sec, slightly
above center and either to the left or right. Participants were asked to create a story that linked
the object to the scene, to press a button when they had made the story. In order to equate
the amount of time spent processing the story, participants were asked to continue to
elaborate it until the scene and object disappeared. A blank screen of 0.5 sec was then
presented prior to the next trial. Participants were informed that the task would include some

pleasant and unpleasant scenes. They were not told that their memory would be tested later.
A practice session of 6 study trials was given just beforehand.

The test phase took place outside the scanner, following a short break of approximately 10
min involving refreshment and conversation with the experimenter. The 120 objects from the
study phase were presented again, randomly intermixed with 40 new objects, and divided into
4 blocks lasting approximately 20 min each. The first stage of each test trial involved tests of
priming, item memory and memory for the picture valence (see Henson et al., 2016 for details)
However, for the present fMRI analysis, we focused on the fourth question in each trial.
Participants were asked to verbally recall the scene that had been paired with the test object
at study. Trials at study in which scenes that were correctly recalled at test, in terms of detail
or gist, were scored as "remembered". Remaining trials were split according to whether the
scenes could not be recalled ("associative misses"), or for which an incorrect scene was
described instead ("associative intrusions"), or for which the object was not recognized ("item
misses"). Initial analyses showed no evidence that valence interacted with age, so trials were
collapsed across valence (see Behavioral results). Table 1 summarizes the trial numbers per
condition split by age tertile.

	Younger (19-45 years) n=38	Middle-Age (46-64 years) n=43	Older (65-88 years) n=42
Remembered	55 (25)	44 (24)	23 (15)
Forgotten			
Associative Miss	31 (13)	38 (17)	46 (18)
Associative Intrusion	11 (8)	13 (10)	10 (8)
Item Miss	22 (15)	25 (17)	42 (24)

Table 1. Trial numbers divided by condition. Remembered = trials with correct object recognition and
scene recall; Associative Miss = trials with correct object recognition but no scene recall; Associative
Intrusion = trials with correct object recognition but recall of an incorrect scene; Item Miss = trials with
misclassification of the object as unstudied. Data are split by age tertile. Means are given with
standard deviations (SD).

For the main imaging analyses, we combined the three types of forgotten trial in order to
maximise power. However, in case processes that lead to subsequent memory for associative
memory versus item memory differ (e.g., Dennis et al., 2008; Mattson et al., 2014), we ran a
subsidiary imaging analyses with item misses excluded.

Imaging data acquisition and preprocessing

The MRI data were collected using a Siemens 3 T TIM TRIO system (Siemens, Erlangen,
Germany). MR data preprocessing and univariate analysis used the SPM12 software
(Wellcome Department of Imaging Neuroscience, London, UK, www.fil.ion.ucl.ac.uk/spm),
release 4537, implemented in the AA 4.0 pipeline
(https://github.com/rhodricusack/automaticanalysis). The functional images were acquired
using T2*-weighted data from a Gradient-Echo Echo-Planar Imaging (EPI) sequence. A total
of 320 volumes were acquired in each of the 2 Study sessions, each containing 32 axial slices
(acquired in descending order), slice thickness of 3.7 mm with an interslice gap of 20% (for
whole brain coverage including cerebellum; TR =1970 msec; TE = 30 msec; flip angle =78
degrees; field of view (FOV) =192 mm × 192 mm; voxel-size = 3 mm × 3 mm × 4.44 mm). A
structural image was also acquired with a T1-weighted 3D Magnetization Prepared RApid
Gradient Echo (MPRAGE) sequence (repetition time (TR) 2250ms, echo time (TE) 2.98 ms,
inversion time (TI) 900 msec, 190 Hz per pixel; flip angle 9 deg; FOV 256 x 240 x 192 mm;
GRAPPA acceleration factor 2).

The structural images were rigid-body registered with an MNI template brain, bias-corrected,
segmented and warped to match a gray-matter template created from the whole CamCAN
Stage 3 sample (N=272) using DARTEL (Ashburner, 2007) (see Taylor *et al.*, 2015) for more
details). This template was subsequently affine-transformed to standard Montreal
Neurological Institute (MNI) space. The functional images were then spatially realigned,
interpolated in time to correct for the different slice acquisition times, rigid-body coregistered
to the structural image and then transformed to MNI space using the warps and affine
transforms from the structural image, and resliced to 3x3x3mm voxels.

Univariate imaging analysis

For each participant, a General Linear Model (GLM) was constructed, comprising three neural
components per trial: 1) a delta function at onset of the background scene, 2) an epoch of 7.5
sec which onset with the appearance of the object (2 sec after onset of scene) and offset when
both object and scene disappeared, and 3) a delta function for each keypress. Each neural
component was convolved with a canonical haemodynamic response function (HRF) to create
a regressor in the GLM. The scene onset events were split into 3 types (i.e, 3 regressors)
according to the valence of the scene on each trial, while the keypress events were modelled

by the same regressor for all trials (together, these four regressors served to model trial-locked
responses that were not of interest). The responses of interest were captured by the epoch
neural component, during which participants were actively relating the scene and object (see
Behavioral Procedure). The duration of this component did not depend on response time, as
participants were instructed to continue to link the object and scene mentally for the full
duration of the display.

For the principal GLMs, the epoch component was split into 6 types (regressors) according to
the 3 scene valences and 2 types of subsequent memory, i.e., study trials for which the scenes
were correctly recalled ("remembered"), and those for which the scenes could not be recalled,
an incorrect scene was described instead, or the object was not recognized ("forgotten").
When comparing remembered and forgotten trials, we averaged across the three valences
because 1) there was no behavioral evidence of an interaction between age and valence on
subsequent memory, 2) there was no imaging evidence of an interaction between age and
valence on subsequent memory, and 3) there would have been insufficient numbers of each
trial-type to examine each valence separately. Thus the main target contrast for the univariate
and multivariate analyses were remembered versus forgotten trials.

As noted above, given that encoding of associative (source) information versus item
information may differ with regard to additional recruitment and (potentially) to compensation
(e.g., Dennis et al., 2008; Mattson et al., 2014), we ran a subsidiary analysis in which the
"forgotten" category excluded item misses. In these GLMs, study trials were modelled using 9
regressors according to the 3 scene valences and 3 (rather than 2) types of subsequent
memory: trials on which the object was recognized but the scene forgotten or incorrectly
recalled ("association forgotten") and trials on which the object was not recognized ("item
forgotten"). Participants for whom one of the sessions did not contain at least one trial of each
type were removed, leaving n=109 (note this involved removal of more participants in the
oldest age tertile: 0 removed aged 19-35, 2 aged 46-64, and 12 aged 65-88 years). One
remaining outlier (>5 SD) on the univariate measures was removed, giving n=108. As reported
in the Results section, this subsidiary analysis corroborated the findings of the main analysis.

Six additional regressors representing the 3 rigid body translations and rotations estimated in
the realignment stage were included in each GLM to capture residual movement-related
artifacts. Finally the data were scaled to a grand mean of 100 over all voxels and scans within
a session, and the model was fitted to the data in each voxel. The autocorrelation of the error
was estimated using an AR(1)-plus-white-noise model, together with a set of cosines that
functioned to highpass the model and data to 1/128 Hz, fit using Restricted Maximum
Likelihood (ReML). The estimated error autocorrelation was then used to "prewhiten" the

model and data, and ordinary least squares used to estimate the model parameters. To
compute subsequent memory effects, the parameter estimates for the 6 epoch components
were averaged across the two sessions and the three valences (weighted by number of trials
per session/valence), and contrasted directly as remembered minus forgotten (Morcom et al.,
2003; Maillet and Rajah, 2014). Univariate statistical analyses were conducted on the mean
subsequent memory effect across all voxels in the MVB analysis, in each ROI for each
participant (see next section).

Experiment 2: Visual STM

Participants

Participants were a separate subset (N=115; 25-86 years; 54 female) of those recruited to the
Cam-CAN study (see Experiment 1, Participants, for details). Nineteen participants were
excluded from the current analysis as the contrast of interest could not successfully be
decoded from either region of interest (see Multivariate Bayesian decoding). None were
excluded because of statistical outlier values on the measures used (see Statistics for criteria).
The experiment used a within-participant design, so all participants received all the task
conditions.

Materials

The task was adapted from Emrich et al. (2013). Stimuli were three patches of coloured dots,
one red, one yellow, and one blue. Dots were 0.26 degrees of visual angle (dva) in diameter,
at a density of 0.7 per square degree, and viewed though a circular aperture of diameter 11
dva. As a manipulation of set size, one, two, or three of the dot displays moved (at 2 dva/ sec)
in a single direction which had to be remembered. The other, distractor, displays rotated
around a central axis, and were be ignored. On 90% of trials the probed movements were in
one of three directions (7, 127, or 247 degrees). Other directions were selected at random, to
avoid subjects learning the target directions. Order of presentation of the 3 display colors was
randomized trial by trial, as was memory load. Rotation direction alternated across trials of a
given load.

Behavioral procedure

Each trial began with a grey fixation dot in the middle of a black screen for 5 sec, which then
turned white for 2 sec. Participants then saw the 3 dot displays for 500 msec each, with 250
msec in-between. After the third display, an 8 sec blank fixation delay was presented, followed
by the probe display. The probe display showed a colored circle to indicate which dot display
to recall (red, yellow, or blue), with a pointer. Participants had up to 5 sec to adjust the pointer
using 2 buttons until it matched the direction of motion of the remembered target dot display.

After responding, a third button cleared the probe display. Participants completed 3 runs of 30

trials per run (10 for each load). The direction of the target, the sequential position of the target,

and the set size were counterbalanced within each run, and presented in random order. Colour

and position of target were also counterbalanced using a Greco-Latin square design.

Imaging data acquisition and preprocessing.

Imaging data were acquired on the same scanner as Experiment 1. Functional T2*-weighted

data were acquired using a Multi-Echo Gradient-Echo Echo-Planar Imaging (EPI) sequence.

Approximately 300 volumes were acquired in each of the 3 VSTM task sessions (duration

depending on response times). Each volume had 34 axial slices (acquired in descending

order), slice thickness of 2.9 mm with an interslice gap of 20% (FOV = 224 mm × 224 mm, TR

= 2000 msec; TE = 12 msec, 25 msec, 38 msec; flip angle = 78 degrees; voxel-size = 3.5 mm

× 3.5 mm × 3.48 mm). Structural image sequences were the same as in Experiment 1. The

multiple echos were combined by computing their average, weighted by their estimated T2*

contrast. The functional images were spatially realigned and interpolated in time to correct for

different slice acquisition times. Spatial normalisation used the 'new segment' protocol in

SPM12 (Ashburner and Friston, 2005). Participants' structural scans were coregistered to their

mean functional image, then segmented into 6 tissue classes. Functional images were rigid-

body coregistered to the structural image then transformed to MNI space using the warps and

affine transforms estimated from the structural image, and resliced to 2x2x2mm voxels.

Univariate imaging analysis

The GLM for each participant comprised three neural components per trial: 1) encoding,

modelled as an epoch of 1 sec duration at onset of the first moving dot pattern, 2)

maintenance, modelled as an epoch of 4 sec at offset of the last moving dot pattern (2.25 sec

after onset of 1), and 3) probe, a delta function at the time of the participant's response. These

components were each split into 3 types (regressors) according to the 3 STM load levels. As

in Experiment 1, 6 additional regressors were added representing the motion parameters

estimated in the realignment stage. Finally the data were scaled to a grand mean of 100 over

all voxels and scans within a session. To confirm that this dataset was suitable as a replication

of Experiment 1's multivariate results, we first checked that at least one significant cluster

within the PFC region of interest (ROI) showed increased univariate activity in older people.

This was done using a standard analysis of effects of a linear contrast of increasing VSTM

load on activity during the delay period, whole-brain corrected for multiple comparisons at p <

.05 (voxel threshold), and a linear contrast of age. Details of this analysis are not reported and

it did not contribute to ROI selection, which was the same as for Experiment 1. Note that the

continuous nature of the judgment in the VSTM task precludes definition of individual trials as

correct or incorrect (rather, performance is used to estimate continuous summary measures
for each participant, as in Emrich et al, 2013). Therefore all trials were included in the fMRI
analysis, and the main target contrast for the univariate and multivariate analyses was the
linear effect of load from 1-3 during the delay period.

Regions of interest

ROIs were defined using WFU PickAtlas (http://fmri.wfubmc.edu/, version 3.0.5) with AAL and
Talairach atlases (Lancaster et al., 2000; Tzourio-Mazoyer et al., 2002; Maldjian et al., 2003).
The posterior visual cortex (PVC) mask comprised bilateral lateral occipital cortex and fusiform
cortex (from AAL, fusiform and middle occipital gyri), and the PFC mask comprised bilateral
ventrolateral, dorsolateral, superior and anterior regions: from AAL, the inferior frontal gyrus
(IFG), both pars triangularis and pars orbitalis; middle frontal gyrus, lateral part (MFG);
superior frontal gyrus (SFG); and from Talairach, Brodmann Area 10 (BA10), dilation factor =
1. In the subregion analyses, separate masks were created for BA10, IFG, MFG and SFG
(regions included in the BA10 mask were excluded from the other masks).

Multivariate Bayesian decoding

A series of MVB decoding models were fit to assess the information about subsequent
memory carried by individual ROIs or combinations of ROIs. Each MVB decoding model is
based on the same design matrix of experimental variables used in the univariate GLM, but
the mapping is reversed: many physiological data features (derived from fMRI activity in
multiple voxels) are used to predict a psychological target variable (Friston et al., 2008). This
target (outcome) variable is specified as a contrast. In Experiment 1 (LTM) the outcome was
subsequent memory, and in Experiment 2 (STM) it was the linear increase in STM load during
maintenance periods. Modelled confounds in the design (all covariates apart from those
involved in the target contrast) are removed from both target and predictor variables.

Each MVB model is fit using hierarchal parametric empirical Bayes, specifying empirical priors
on the data features (voxel-wise activity) in terms of patterns over voxel features and the
variances of the pattern weights. Since decoding models operating on multiple voxels (relative
to scans) are ill-posed, these spatial priors on the patterns of voxel weights act as constraints
in the second level of the hierarchical model. MVB also uses an overall sparsity (hyper) prior
in pattern space which embodies the expectation that a few patterns make a substantial
contribution to the decoding and most make a small contribution. The pattern weights
specifying the mapping of data features to the target variable are optimised with a greedy
search algorithm using a standard variational scheme which iterates until the optimum set size
is reached (Friston et al., 2007). This is done by maximizing the free energy, which provides

an upper bound on the Bayesian log-evidence (the marginal probability of the data given that

model). The evidence for different models predicting the same psychological variable can then

be compared by computing the difference in their log-evidences, giving the log (marginal)

likelihood ratio test (Bayes factor) (see Friston et al., 2007; Chadwick et al., 2012; Morcom

and Friston, 2012). In this work, the main outcome measures were the log-evidence for each

model and the set of fitted weights for all patterns (voxels) in the ROI, which can be examined

to assess their distribution over voxels and the contributions of different combinations of

voxels. These analyses were implemented in SPM12 v6486 and custom MATLAB scripts.

Features (voxels) for MVB analysis were selected using an orthogonal contrast and a leave-

one-participant-out scheme. For each participant and ROI, these were the 1000 voxels with

the strongest responses to the task: in Experiment 1 (LTM), the 6 epoch regressors modelling

object onsets in the GLM, and in Experiment 2 (STM), the 3 epoch regressors modelling

maintenance periods in the GLM (defined using an F contrast in all other participants testing

variance explained by these regressors, regardless of valence or subsequent memory). We

used a sparse spatial prior, in which each pattern is an individual voxel (Morcom and Friston,

2012; Chadwick et al., 2014; Hulme et al., 2014; Maass et al., 2014). We first checked that

the target memory variables could reliably be decoded from the selected features by

contrasting the evidence for each model with the evidence for models in which the design

matrix (and therefore the target variable) had been randomly phase-shuffled, taking the mean

over 20 repetitions, and comparing log-evidence for real versus phase-shuffled models. One-

tailed t-tests compared the difference in real versus shuffled model evidences to a

hypothesized population mean difference of 3 which would reflect good Bayesian evidence for

the real over the shuffled models. These showed that the difference in log-evidence was

robustly greater than this in both PVC, $t(118) = 6.08$, p < .0001, $r^2_{adj} = .225$, mean difference =

9.72; and PFC, $t(118) = 7.70$, p < .0001, $r^2_{adj} = .323$, mean difference = 11.8. The same applied

to Experiment 2: for PVC, $t(95) = 8.42$, p < .0001, $r^2_{adj} = .415$, mean difference = 23.0, and

PFC, $t(95) = 11.4$, p < .0001, $r^2_{adj} = .569$, mean difference = 35.0. To confirm that the sparse

prior represented the best spatial model, we then compared the log-evidence with that for

models with smooth spatial priors, in which each pattern is a local weighted mean of voxels

(Gaussian FWHM = 8). For Experiment 1 (LTM): log evidence was substantially greater for

the sparse priors in both ROIs: in PVC, $t(118) = 18.4$, p < .0001, $r^2_{adj} = .737$, and PFC, $t(118)$

= 18.0, $r^2_{adj} = .728$, p < .0001, two-tailed tests. The same was true for Experiment 2 (STM): for

PVC, $t(95) = 10.3$, $r^2_{adj} = .464$, p < .0001, and PFC, $t(95) = 14.9$, $r^2_{adj} = .650$, p < .0001.

Unlike univariate activation measures such as subsequent memory effects, but like other

pattern-information methods, MVB finds the best non-directional model of activity predicting

the target variable, so positive and negative pattern weights are equally important. Therefore,

the principle MVB measure of interest for each ROI was the spread (standard deviation) of the
weights over voxels, reflecting the degree to which multiple voxels carried substantial
information about subsequent memory. To test whether PFC activity was compensatory, we
also constructed a novel measure of the contribution of prefrontal cortex to subsequent
memory. This used Bayesian model comparison within participants to assess whether a joint
PVC-PFC model boosted prediction of subsequent memory relative to a PVC-only model. The
PASA hypothesis, in which PFC is engaged to a greater degree in older age and this
contributes to cognitive outcomes, predicts that a boost will be more often observed with
increasing age. The initial dependent measure was the log model evidence, coded
categorically for each participant to indicate the outcome of the model comparison. The 3
possible outcomes were: a boost to model evidence for PVC-PFC relative to PVC models, i.e.,
better prediction of subsequent memory (difference in log evidence > 3), equivalent evidence
for the two models (-3 < difference in log evidence < 3), or a reduction in prediction of
subsequent memory for PFC-PVC relative to PVC (difference in log evidence < -3).

Lastly, given that the relative contribution of anterior versus posterior regions could change
with age, even if the absolute amount of pattern information decreased with age in both
regions, we computed a second novel measure: we estimated the PFC contribution to
cognitive outcome in terms of the proportion of top-weighted voxels in the joint PVC-PFC
model that were located in PFC, as opposed to PVC, derived from joint PVC-PFC models. In
each participant, the voxels making the strongest contribution to the cognitive outcome,
defined as those with absolute voxel weight values greater than 2 standard deviations from
the mean, were split according to their anterior versus posterior location. The dependent
measure was the proportion of these top voxels located in PFC.

Experimental design and statistical analysis

Sample size was determined by the initial considerations of Stage 3 of the CamCAN study –
see Shafto et al.(Shafto et al., 2014) for details. For the LTM experiment, a sensitivity analysis
indicated that with N=123, we would have 80% power to detect a small to medium effect
explaining 6.5% of the variance on a two tailed test for a model with 2 predictors (r^2 = .0658).
For the STM experiment with N = 115, the corresponding minimal effect size for 80% power
was 6.9% of the variance (r^2 = .0694). In our previous report of aging and successful memory
encoding, an *a priori* test of a between-region difference in subsequent memory effects
according to age showed a large effect (r^2 = .257) (Morcom et al., 2003).

Age effects on continuous multivariate or univariate dependent measures were tested using
robust second-order polynomial regression with "rlm" in the package MASS for R (Venables
et al., 2002); MASS version 7.3-45; R version 3.3.1) with standardized linear and quadratic

age predictors. For analysis of covariance for behavioral data we used JASP version 0.8.3.1.

Analysis of outcomes of the between-region MVB model comparison (PVC and PFC combined

versus PVC, see Fig 2 and main text) used ordinal regression with "polr" in MASS.

Distributions were also trimmed to remove extreme outliers (> 5 SD above or below the mean).

In Experiment 1 (LTM), the two participants (aged 72 and 80) with outlier values for univariate

effects were also removed from the MVB analyses so the samples examined were

comparable. We excluded two further participants (aged 68 and 83) in whom subsequent

memory could not be decoded from at least one of the two ROIs (log model evidence <= 3),

giving n=119. In Experiment 2 (STM), we excluded nineteen participants in whom VSTM load

could not be decoded during maintenance, giving n=96. All tests were two-tailed and used an

alpha level of .05.

Where it was important to test for evidence for the null hypothesis over an alternative

hypothesis, we supplemented null-hypothesis significance tests with Bayes Factors

(Wagenmakers, 2007; Rouder et al., 2009). The Bayes Factors were estimated using Dienes'

online calculator (Dienes, 2014) which operationalizes directional hypotheses such as PASA

in terms of a half-normal distribution. Here, we assumed an effect size of 1 SD and therefore

defined the half-normal distribution with mean=0 and SD=1. All statistics and p values are

reported to 3 significant figures, except where $p < .0001$.

Results

Experiment 1: Long-term Memory (LTM) Encoding

Behavioral results

We examined age effects on the number of trials in each remembered and forgotten condition

(see Table 2). For remembered trials, there was a significant linear decrease with age (t(118)

= -7.30, p < .0001, r^2_{adj} = .299), with no significant quadratic component (t(118) = -0.104, p =

.917; alpha = .0125). As a consequence, the number of forgotten trials increased with age,

and this was true for both associative misses (linear t(118) = 4.82, p < .0001, r^2_{adj} = .150;

quadratic, t(118) = 0.630, p = .532) and item misses (linear t(118) = 5.43, p < .0001, r^2_{adj} =

.186; quadratic, t(118) = 1.57, p = .120), although not for associative intrusions (linear t(118)

= -1.38, p = .163; quadratic, t(118) = -2.29, p = .0221). Analysis of covariance with the factor

of valence (Positive, Neutral, Negative) showed no interaction between age and valence on

the number of remembered items (for linear effect of age, F(2,231) < 1, p = .486; quadratic,

F(2,231) = 1.59, p = .206).

Testing compensation

Standard univariate activation analyses assessed mean activity in each ROI across all voxels included in the multivariate analysis (see Materials and Methods). Consistent with the PASA account, the increase in activity associated with subsequent memory became more pronounced with age, particularly in later years (linear effect of age, t(118) = 2.43, p = .0166; quadratic effect of age, t(118) = 2.58, p = .0111) (Fig 2a, Table 2). Age effects in PVC were not significant (see Table 2). The age effects in PFC were also present after removal of the older participant with the largest SM effect (although they did not meet our criterion for an outlier; see Fig 2a; linear t(117) = 2.14, p = ; quadratic F(117) = 2.31, p = .033). In both ROIs results were very similar for the models excluding forgotten trials for which the items themselves were forgotten (see Methods; PFC: linear t(107) = 2.22, p = .0316; quadratic t(107) = 2.91, p = .00527; PVC: linear, t(107) = 1.10, p= .288; quadratic, t(107) = 1.24, p = .233).

Figure 2. Relationship between age and univariate and multivariate effects within ROIs. **(a)**. Univariate subsequent memory effects (mean activity for remembered - forgotten), showing increased activity with age in PFC but not PVC. **(b).** Spread of multivariate responses predicting subsequent memory (standard deviation of fitted MVB voxel weights), showing reduced spread of responses with age in both ROIs. **(c).** Univariate effects of load (positive linear contrast) during STM maintenance, showing increased activity with age in both ROIs. **(e).** Spread of multivariate responses during STM maintenance predicting increasing load, showing reduced spread of responses with age in both ROIs. Red and blue lines are robust-fitted second-order polynomial regression lines and shaded areas show 95%

461 confidence intervals. **(e).** PVC (blue) and PFC (red) ROIs overlaid on sagittal section (x=+36) of a

462 canonical T1 weighted brain image. Note that *y*-axis scales are not comparable across tasks.

ROI/ measure	Model		Linear			Quadratic		
	F	p	t	r^2_{adj}	p	T	r^2_{adj}	p
Mean univariate SM activation								
PFC	5.49	.00525	2.43	.0312	0.0166	2.58	.0371	.0111
PVC	0.426	.654	0.728	--	0.480	0.703	--	.495
PFC-PVC	0.837	.436	0.883	--	0.388	1.084	--	0.293
Multivariate spread (SD) of SM activity								
PFC	6.36	.00240	-3.33	.0701	.000998	-1.44	--	.151
PVC	11.3	< .0001	-3.49	.0780	.000650	-3.50	.0784	.000621
PFC-PVC	2.02	.109	4.16	--	.0437	.398	--	.690

Table 2. Age effects on mean univariate SM effects and spread of multivariate SM effects in the LTM

task. PFC-PVC refers to analyses where the dependent variable was the difference in each measure

between PFC and PVC. SD = standard deviation. r^2_{adj} = the unbiased estimate of the amount of variance

explained in the population. n = 119.

If the increasing PFC activation with age reflected compensation, we expected the multivariate

analyses to show that this increased activity carried additional information about subsequent

memory. However, the data revealed a different pattern. In MVB models, like other linear

models with multiple predictors, each voxel within an ROI has a weight that captures the

unique information it contributes (in this case, for predicting subsequent memory). Because

both positive and negative weights carry information, we summarised the MVB results by the

spread (standard deviation) of weights over voxels (see Materials and Methods).

In both ROIs, this spread was markedly reduced during later life (PVC: linear t(118) = -3.49,

p= .000650; quadratic t(118) = -3.50, p = .000621; PFC: linear t(118) = -3.33, p = .000998;

quadratic t(118) = -1.44, p = .151); see Fig 2b and Table 2. This means that, contrary to a

compensatory PASA shift, PFC showed fewer, rather than more, voxels with large positive or

negative weights with increasing age. Again, the results were similar for the subsidiary models

excluding item misses (PVC: linear t(107) = -1.41, p= .158; quadratic t(107) = -2.81, p =

.000544; PFC: linear t(107) = -2.21, p = .0280; quadratic t(107) = -0.566, p = .570). By contrast,

the spread of univariate activities across voxels increased with age in both ROIs (for PVC,

linear effect of age, t(118) = 5.91, p < .0001, r^2_{adj} = .215; quadratic effect of age, t(118) = 1.72,

p = .0881; for PFC, linear effect of age, $t(118)$ = 5.64, p < .0001, r^2_{adj} = .199; quadratic $t(118)$

= -2.31, p = .0226, r^2_{adj} = .0268).

To provide a more direct test for a compensatory posterior-to-anterior shift, we assessed the

specific contribution of PFC to subsequent memory, over and above that of PVC. We fitted a

joint MVB model that included both posterior and anterior ROIs, and contrasted this with a

model including PVC only, using Bayesian model comparison. Thus we could ask, for each

participant, whether or not adding PFC to the model "boosted" prediction of subsequent

memory (see Methods). Contrary to the PASA theory of a compensatory shift towards greater

reliance on PFC, a Bayes Factor comparing these two models revealed strong evidence for

the null hypothesis of no boost (BF_{01} = 11.1); indeed, the probability of a boost to model

evidence for PVC-PFC compared to PVC-only actually *decreased* with age numerically (Fig

3a; linear $t(118)$ = -1.54, p = .126). Excluding item misses from the model enhanced this

pattern (for linear age effect $t(107)$ = -2.34, p = .0211, BF_{01} = 14.3).

Figure 3. Evidence against a compensatory posterior-to-anterior shift from MVB comparisons between

ROIs. Ordinal regression of Bayesian model comparison of combined PVC+PFC model versus PVC-

only model using age to predict outcomes of model comparison: adding PFC to the model boosts

prediction of the cognitive outcome (difference in log-evidence > 3), or there is no boost (-3 < difference

< 3), or a reduction in log-evidence (difference < -3). (a) LTM, for subsequent memory effects, a boost

was no more frequent with increasing age (b) STM, for load effects, a boost was less frequent with

increasing age.

Testing sub-regions within PFC

We also explored whether the pattern of results was similar across subregions within PFC.

The PASA theory does not specify which areas are involved in a compensatory shift, but aging

does not affect all subregions equally (Raz and Rodrigue, 2006). In functional studies,
univariate SM effects in ventrolateral and dorsolateral PFC tend to be age-invariant while
anterior and superior prefrontal regions show age-related increases (Morcom et al., 2003;
Maillet and Rajah, 2014). In contrast, Davis et al.'s (2008) PASA proposal was based on
increased activation in older people in anterior ventrolateral PFC and anterior cingulate during
visual perception and episodic retrieval. In the present episodic encoding task, there were
significant age-related increases in univariate activation in anterior PFC (BA10) and lateral
inferior frontal gyrus (IFG), and significant decreases in the spread of multivariate voxel
weights in BA10 and superior (medial) frontal gyrus (SFG), as well as numerical decreases
also in IFG and in lateral middle frontal gyrus including dorsolateral PFC (MFG) (Table 3; see
Materials and Methods, Regions of Interest for region definition). Thus the overall age-related
increase in activation was mainly driven by BA10 and IFG, but no subregion showed a
decrease in activation with age. The reduction in multivariate information and evidence against
a functional boost were relatively uniform over subregions (Table 3). Direct model comparison
showed no evidence that PFC activity was compensatory in older age, even in the two
subregions with strong increases in activation: Bayes Factors weighed against any boost to
prediction of subsequent memory for joint PFC-PVC models relative to PVC-only models (BF_{01}
favoring the null over positive linear effect of age = 11.1 for BA10, 12.5 for MFG, 5.00 for IFG
and 14.2 for SFG).

ROI/ measure	Model			Linear			Quadratic		
	F	p	t	r^2_{adj}	p	t	r^2_{adj}	p
BA10								
Mean univariate SM	5.24	.00660	1.33	--	.180	2.36	--	.0189
MVB spread (SD)	8.28	.000433	-3.75	.0911	.0003249	-1.86	--	.0623
PFC boost	--	--	-1.44	--		-0.321	--	
IFG								
Mean univariate SM	5.39	.00572	2.48	.204	.0152	2.72	.227	.00824
MVB spread (SD)	3.30	.0405	-2.50	--	.0127	-0.653	--	.514
PFC boost	--	--	0.00665	--		-0.210	--	
MFG								
Mean univariate SM	1.34	.266	1.56	--	.119	1.38	--	.169
MVB spread (SD)	4.54	.0126	-2.86	.0487	.00466	-1.10	--	.271
PFC boost	--	--	-1.45	--		-.132	--	
SFG								
Mean univariate SM	1.72	.184	1.33	--	.181	1.34	--	.179
MVB spread (SD)	4.39	.0145	-2.83	.0474	.00533	-1.09	--	.275

PFC boost	--	--	-1.68	--	1.03	--

Table 3. Age effects for PFC subregions in the LTM task. The table lists mean univariate SM effects, the spread (SD) of multivariate Bayesian (MVB) voxel weights predicting SM, and results of the between-region tests of 'boost' to model evidence for PFC plus PVC models compared to PVC-only. See text for details. Alpha = .0125. SD = standard deviation. SM = subsequent memory. n=119.

Testing posterior-anterior shift

The foregoing analyses provide strong evidence that the increase in (univariate) PFC activity observed in this task did not reflect compensation. Nonetheless, the PASA theory is more general, describing a shift in relative reliance on posterior and anterior regions with age, which need not be compensatory, but could reflect differential age effects in posterior and anterior cortices. In other words, the relative involvement of anterior versus posterior regions could increase with age, even if the absolute involvement of both decreased with age. Direct comparison of the mean univariate activation between ROIs did not reveal any evidence for such relative differences in age effects, with strong Bayesian evidence against the predicted greater age-related increase in PFC (BF_{01} for null hypothesis = 25; Table 2). We next tested for a shift in the relative multivariate information between regions. In the separate MVB models, the age-related reduction in spread of weights was numerically *greater* in PFC than PVC (linear age effect on PFC-PVC difference, $t(118) = 4.16$, p = .0437); see Table 2. We also measured the proportion of top-weighted voxels (> 2 standard deviations above the mean) that were located in PFC as opposed to PVC in the joint PVC-PFC model. This proportion decreased significantly with age (overall model, $F(2,116) = 3.27$, p = .0415; linear $t(118) = -2.55$, $r^2_{adj} = .359$, p = .0119; quadratic $t(118) = -.106$, p = .915), with mean 52.4% of top voxels located in PFC in the younger tertile (SD = 9.09; 18-45 years) and 47.1% in the older tertile (SD = 8.57; 65-88 years), although this was no longer significant when item misses were excluded (overall model, $F(2,105) = 2.60$, p = .0799, linear t(107) = -1.86, p = .0638). Thus, there was no evidence that in older age there is a general shift in the areas contributing to subsequent memory from posterior to anterior (though see Experiment 2 below).

Experiment 2: Short-term Memory (STM) Maintenance

Behavioral results

For the visual STM task, analysis of the effects of increasing load on performance showed a strong age-related increase in the effect of load on accuracy, measured using the root mean squared error of the estimated dot direction relative to the actual dot direction in degrees (Fig 1b). As expected, older people showed a larger increase in error at load = 3 compared to load

= 1 (linear contrast) than younger people (for linear age-by-load interaction, $t(95) = 5.53$, p <

.0001, $r^2_{adj} = .192$; quadratic $t(95) = =1.27$, p = .203), although some age-related decrement in

accuracy was present even at load = 1 (for linear effect of age, $t(95) = 2.607$, p = 0110, $r^2_{adj} =$

.0382; quadratic $t(95) = 0.388$, p = .699).

Testing compensation

For STM, standard univariate activation analyses showed that increasing load elicited activity

increases during the maintenance period which varied according to age in both ROIs. As in

the LTM experiment, and consistent with the PASA account, PFC activation increased with

age, particularly in later years (linear t(95) = 3.01, p = .003; quadratic $t(95) = -0.505$, p = .615)

(Fig 2c, Table 4). Unlike for LTM encoding, there was also a significant increase in load-related

PVC activation over the lifespan (linear $t(95) = 4.28$, p < .0001; quadratic $t(93) = -0.988$, p =

.324; see Table 4.

ROI/ measure	Model		Linear			Quadratic		
	F	p	t	r^2_{adj}	p	T	r^2_{adj}	p
Mean univariate STM activation								
PFC	4.57	.0128	3.01	.0553	.00336	-0.505	--	.615
PVC	9.43	.000187	4.28	.119	< .0001	-0.988	--	.324
PFC-PVC	.606	.548	-0.587	--	.559	-0.878	--	.380
Multivariate spread (SD) of STM activity								
PFC	13.4	< .0001	.662	--	.507	-5.03	.162	< .0001
PVC	9.30	.000210	-1.01	--	.308	-4.07	.108	< .0001
PFC-PFC	3.00	.0547	.674	--	.497	2.26	.0250	0244

Table 4. Age effects on mean univariate SM effects and spread of multivariate SM effects in the STM

task. PFC-PVC refers to analyses where the dependent variable was the difference in each measure

between PFC and PVC. SD = standard deviation. n = 96. R^2_{adj} = the unbiased estimate of the amount

of variance explained in the population.

Separate MVB analysis in each ROI showed a similar pattern of age effects to the LTM task.

In both PFC and PFC, the spread (SD) of individual voxel weights predicting increased STM

load was particularly reduced during later life, with a significant quadratic component (PVC:

linear $t(95) = -1.01$, p = .308; quadratic $t(95) = -4.07$, p < .0001; PFC: linear $t(95) = 0.662$, p =

.507; quadratic $t(95) = -5.03$, p < .0001) (see Fig 2d and Table 4). The result for PVC was

unchanged by removing a subset of subjects with very low values (i.e., SD weights < .0005;
for quadratic age effect $t(69) = -5.32$, p = .0012). As found for LTM encoding, the direction of
the effect in PFC was contrary to a compensatory PASA shift, i.e., PFC voxels contributed
less to the cognitive task in old age. Again, the spread of univariate effects did not show the
same effects of age (in PVC linear $t(95) = 1.52$, p = .134; quadratic $t(95) = 0.831$, p = .406; in
PFC, linear $t(95) = -0.471$, p = .641; quadratic $t(95) = 1.70$, p = .0912).

As for LTM encoding, we used model comparison of a joint PVC-PFC model with a PVC-only
model to directly evaluate the compensatory PASA hypothesis. The results were similar to the
LTM experiment: The "boost" to prediction of the cognitive variable obtained by adding PFC
to the model showed a significant age-related *reduction* in the probability of a boost for STM
load (in an ordinal regression, $t(95) = -2.00$, p = .0479). The Bayes Factor provided strong
evidence against the compensatory hypothesis of an increased boost (for unidirectional
hypothesis, BF = 10.2), although evidence was only anecdotal for the presence of an age-
related reduction in boost (for bidirectional hypothesis, BF = 1.81). Thus, like for the LTM
experiment, there was clear evidence against a compensatory increase in prefrontal
contribution to the task with age.

Testing sub-regions within PFC

Again, we examined the four prefrontal subregions separately to assess whether the findings
were driven by a specific part or parts of the large ROI (Table 5). For this experiment, the age-
related increase in univariate activation was not separately significant in any subregion, which
may have reflected relatively distributed effects and the more inclusive selection of 'active'
voxels. As for the LTM experiment however, overall age effects on the spread of multivariate
voxel weights were significant in three subregions, and those in IFG were similar in magnitude
and form, suggesting reductions in spread across PFC, with no major between-subregion
differences. Moreover, all ROIs showed Bayes Factors of at least 6 against a boost to model
evidence from adding PFC to the posterior-only models predicting increasing STM load, again
consistent with the overall results.

ROI/ measure	Model		Linear			Quadratic		
	F	P	t	r^2_{adj}	p	t	r^2_{adj}	p
BA10								
Mean univariate	1.90	.155	1.79	--	.0787	-0.937	--	.352
MVB spread (SD)	12.7	< .0001	-1.41	--	.158	-4.69	.171	< .0001
PFC boost t	--	--	-1.48	--	.142	-1.00	--	.320
PFC boost BF_{01}	--	--	10.0	--	--	--	--	--
IFG								

Mean univariate	2.64	.0767	1.99	--	.0512	0.997	--	.323
MVB spread (SD)	6.26	.00282	-0.787	--	.427	-3.35	.0864	.00109
PFC boost t	--	--	-1.11	--	.270	-1.10	--	.274
PFC boost BF_{01}	--	--	7.69	--	--	--	--	--
MFG								
Mean univariate	2.08	.131	2.03	--	.0459	-0.447	--	.654
MVB spread (SD)	11.0	< .0001	0.300	--	.762	-4.60	.165	< .0001
PFC boost t	--	--	-1.38	--	.171	-0.788	--	.433
PFC boost BF_{01}	--	--	6.25	--	--	--	--	--
SFG								
Mean univariate	2.64	.0770	2.27	--	.0264	0.241	--	.812
MVB spread (SD)	7.37	.00106	-1.57	--	.116	-3.34	.0858	.00111
PFC boost t	--	--	-2.73	.0528	.00755	-0.720	--	.473
PFC boost BF_{01}	--	--	14.3	--	--	--	--	--

Table 5. Age effects for PFC subregions in the visual short-term memory task. The table lists mean univariate activation during maintenance in response to increasing VSTM load, the spread (SD) of MVB voxel weights predicting linearly increasing VSTM load, and results of the between-region tests of 'boost' to model evidence for PFC plus PVC models compared to PVC-only. See text for details. Alpha = .0125. SD = standard deviation. n=96.

Testing posterior-anterior shift

Finally, even if age increased activity and decreased multivariate information in both PFC and PVC, it is possible that the PFC:PVC ratio of activity and/or multivariate information increases with age, consistent with the general PASA claim. As for LTM encoding, there was no evidence that age effects on (univariate) activation in the two ROIs differed, i.e. the increase in activation in older people was similar in magnitude (Table 4; BF_{01} for null hypothesis over prediction of a greater age-related increase in PFC = 33.3). However, multivariate analysis revealed a picture different from that in the LTM task. In the separate MVB models, the age-related reduction in spread of weights showed a stronger quadratic component in PFC than PVC (for PFC-PFC, quadratic $t(95) = 2.26$, p = .0244; Table 4). More clearly, when examining the location of top-weighted voxels from the joint PFC+PVC model, a higher proportion were located in PFC in older age (for model, $F(2,93) = 22.4$, p < .0001, linear $t(95) = 3.72$, p < .001, quadratic $t(95) = 5.20$, p < .0001), with mean 69.1% of top voxels located in PFC in the younger tertile (SD = 14.1; 25-43 years) but 81.0% in the older tertile (SD = 13.3; 66-85 years). Thus while the STM experiment, like the LTM experiment, found decreases in absolute PFC (and PVC) involvement in old age, the relative involvement of PFC versus PVC voxels (at least in

terms of those with high weights in the joint model) did increase with age, unlike in the LTM
experiment. This provides some support for a PASA pattern in this task, even though there
was no evidence that this greater PFC involvement was compensatory.

Discussion

This study investigated the proposal that there is a posterior-to-anterior shift in task-related
brain activity during aging, with the greater reliance on prefrontal cortex in older age reflecting
compensatory mechanisms. We tested the predictions of this PASA theory with data from two
memory tasks that were conducted on independent and relatively large population-derived
adult lifespan samples. Using novel model-based multivariate analyses, we provide direct
evidence against a compensatory posterior-to-anterior shift. Instead, our data suggest that the
increased prefrontal activation reflects less specific or less efficient activity, rather than
compensation.

The results of our standard univariate activation analyses are consistent with previous studies
showing age-related increases in activation in prefrontal cortex, which form the basis of the
PASA theory (Grady et al., 1994; Davis et al., 2008). Many studies have found such increases
in PFC activation in different cognitive tasks, although regional reductions in activation are
also found (e.g. see Rajah and D'Esposito, 2005; Spreng et al., 2010; Li et al., 2015) We found
such age-related increases in both the PFC activation associated with trials that were later
remembered many minutes later (in the LTM experiment) and the activation associated with
maintaining increasing numbers of items for a few seconds (in the STM experiment). We also
further generalized previous findings across PFC sub-regions, in that the increased activation
was reliable across lateral, anterior and superior prefrontal areas (although in the LTM
experiment, it was mainly driven by inferolateral and anterior PFC).

Importantly, despite this increased univariate activity, multivariate analysis of both
experiments showed that with increasing age, PFC possessed less, rather than more,
information about the cognitive outcome. This reduced pattern information was evident both
in terms of the spread of voxel weights (Figure 2) and the lack of a meaningful boost to model
evidence when adding PFC voxels to the model (Figure 3). The latter type of inference was
made possible by our novel use of multivariate Bayesian (MVB) classification.

If the increased prefrontal activation with age is not compensatory, then what does it reflect?
One possibility is that neural function is less efficient, such that a greater BOLD signal is
required for the same level of computation, i.e, less "bang for the buck" for the same level of
neural activity (Grady, 2008; see also Rypma and Esposito, 2000; Morcom et al., 2007;
Reuter-Lorenz and Campbell, 2008; Nyberg et al., 2014). This could reflect the proposed

greater sensitivity of prefrontal cortices than other brain regions to aging (West, 1996; Glisky

et al., 2001; Raz and Rodrigue, 2006). Another possibility is that PFC activity becomes less

specific with age, as might be expected by theories of age-related dedifferentiation, particularly

in complex cognitive functions (Li et al., 2001; Park et al., 2004; Carp et al., 2011;

Abdulrahman et al., 2014). Partial support for the latter comes from the LTM experiment,

where the negative effect of age on the spread of multivariate weights across voxels was

accompanied by a positive effect of age on the spread (as well as mean) of univariate activity.

This suggests that, while more voxels showed substantial (positive or negative) activity related

to subsequent memory in older people, these additional responses were redundant, with fewer

voxels contributing uniquely to memory encoding, as expected if the increased prefrontal

activity is less specific. In the STM experiment, on the other hand, the spread of univariate

responses was age-invariant, suggesting a more spatially uniform increase in response to

load with age, although the MVB results suggested that – just as in the LTM task – this

increased response carried less information. Whether the present results reflect reductions in

efficiency or reductions in specificity, they are more consistent with the general idea of brain

maintenance (Nyberg et al., 2014) – that cognitive function in older age is determined by the

ability to maintain a youth-like brain – than with the idea associated with PASA of functional

compensation by anterior brain regions.

Despite age-related decreases in overall multivariate information in both PFC and PVC, it is

possible that the *relative* contribution of anterior regions to cognitive tasks could increase with

age. There is some evidence for such a shift from studies showing crossover effects in which

age-related decreases in posterior cortical activity occur alongside age-related increases in

PFC (e.g., Grady et al., 1994; Davis et al., 2008; see also recent meta-analysis by Maillet and

Rajah, 2014). However, our univariate activation analyses showed little evidence of such a

relative posterior-to-anterior shift: despite increased prefrontal activation, age effects on

univariate activation in PFC and PVC did not differ significantly in either experiment. In terms

of multivariate information, the LTM experiment actually showed, if anything, a decrease rather

than increase in the contribution of PFC relative to PVC. The only comparison that provided

some support for a relative increase in anterior contribution was for multivariate information

about load in the STM experiment. Thus the posterior-to-anterior shift claimed by PASA seems

to be task-dependent (see also Ford and Kensinger, 2017). This is consistent with other meta-

analyses, which have found age-related decreases as well as increases in activation,

depending on the task (Spreng, Wojtowicz, & Grady, 2010; Li et al., 2015). Moreover, most

studies have not made the direct statistical comparisons needed to test for anterior-posterior

differences in the absence of crossover effects (see Morcom & Johnson, 2015). A strength of

our approach is that our analyses encompassed large ROIs in both anterior and posterior
cortices, as well as direct comparisons between the two.

In summary, our data replicate an increase in PFC activity over the adult lifespan, but do not
support the idea that this reflects a compensatory posterior-to-anterior shift, at least in the
context of the two memory tasks considered here. Our results are inconsistent both with the
proposal that the increased activity is compensatory, and with a generalized shift with age to
greater relative reliance on prefrontal cortex. The data are most parsimoniously explained by
reduced efficiency or specificity of neural responses, reflecting primary age-related deleterious
changes in posterior as well as prefrontal cortex. Our results therefore help to adjudicate
between competing accounts of neurocognitive aging, while also illustrating the more general
ability of MVB to compare models that comprise different sets of voxels, thereby offering an
exciting new general way to test the relative contributions of brain regions to cognitive
outcomes.

Acknowledgments

A.M.M. is a member of the University of Edinburgh Centre for Cognitive Ageing and Cognitive
Epidemiology (CCACE), part of the UK cross-council Lifelong Health and Wellbeing Initiative,
Grant number G0700704/84698. The Cambridge Centre for Ageing and Neuroscience (Cam-
CAN) research was supported by the Biotechnology and Biological Sciences Research
Council (grant number BB/H008217/1). The full Cam-CAN author list can be found here:
http://www.cam-can.org/index.php?content=corpauth. R.N.A.H. is funded by the Medical
Research Council (SUAG/010 RG91365) with additional support by the European Union's
Horizon 2020 research and innovation programme (grant agreement No 732592).

Author contributions

A.M.M. designed research; A.M.M., R.N.A.H. and Cam-CAN performed research; A.M.M. and
R.N.A.H. analyzed data; A.M.M. and R.N.A.H. wrote the paper.

References

Abdulrahman H, Fletcher PC, Bullmore E, Morcom AM (2014) Dopamine and memory
dedifferentiation in aging. Neuroimage.

Ashburner J (2007) A fast diffeomorphic image registration algorithm. 38:95–113.

Ashburner J, Friston KJ (2005) Unified segmentation. Neuroimage 26:839–851.

Cabeza R, Daselaar SM, Dolcos F, Prince SE, Budde M, Nyberg L (2004) Task-independent
and Task-specific Age Effects on Brain Activity during Working Memory, Visual Attention

and Episodic Retrieval. Cereb Cortex 14:364–375.

Cappell KA, Gmeindl L, Reuter-lorenz PA (2010) Age differences in prefontal recruitment
during verbal working memory maintenance depend on memory load. CORTEX 46:462–
473 Available at: http://dx.doi.org/10.1016/j.cortex.2009.11.009.

Carp J, Park J, Polk TA, Park DC (2011) Age differences in neural distinctiveness revealed by
multi-voxel pattern analysis. Neuroimage 56:736–743 Available at:
http://dx.doi.org/10.1016/j.neuroimage.2010.04.267.

Chadwick MJ, Bonnici HM, Maguire EA (2012) Decoding information in the human
hippocampus: A user's guide. Neuropsychologia 50:3107–3121 Available at:
http://dx.doi.org/10.1016/j.neuropsychologia.2012.07.007.

Chadwick MJ, Bonnici HM, Maguire E a. (2014) CA3 size predicts the precision of memory
recall. Proc Natl Acad Sci 111:10720–10725 Available at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4115494&tool=pmcentrez&re
ndertype=abstract.

Davis SW, Dennis NA, Daselaar SM, Fleck MS, Cabeza R (2008) Que PASA? the posterior-
anterior shift in aging. Cereb Cortex 18:1201–1209.

Dienes Z (2014) Using Bayes to get the most out of non-significant results. Front Psychol 5:1–
17.

Emrich SM, Riggall AC, LaRocque JJ, Postle BR (2013) Distributed Patterns of Activity in
Sensory Cortex Reflect the Precision of Multiple Items Maintained in Visual Short-Term
Memory. J Neurosci 33:6516–6523 Available at:
http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.5732-12.2013.

Ford JH, Kensinger EA (2017) Age-Related Reversals in Neural Recruitment across Memory
Retrieval Phases. J Neurosci:0521–17 Available at:
http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.0521-17.2017.

Friston K, Chu C, Mour??o-Miranda J, Hulme O, Rees G, Penny W, Ashburner J (2008)
Bayesian decoding of brain images. Neuroimage 39:181–205.

Friston K, Mattout J, Trujillo-Barreto N, Ashburner J, Penny W (2007) Variational free energy
and the Laplace approximation. Neuroimage 34:220–234.

Glisky EL, Rubin SR, Davidson PSR (2001) Source Memory in Older Adults : An Encoding or
Retrieval Problem ? 27:1131–1146.

Grady C (2012) The cognitive neuroscience of ageing. Nat Rev Neurosci 13:491–505
Available at: http://www.ncbi.nlm.nih.gov/pubmed/22714020.

Grady CL, Maisog JM, Horwitz B, Ungerleider LG, Mentis MJ, Salerno JA, Pietrini P, Wagner
E, Haxby J V (1994) Age-related changes in cortical blood flow activation during visual
processing of faces and location. J Neurosci 14:1450–1462.

Grady CL, Mcintosh AR, Bookstein F, Horwitz B, Rapoport SI, Haxby J V (1998) Age-Related
Changes in Regional Cerebral Blood Flow during Working Memory for Faces. 425:409–
425.

Henson RN et al. (2016) Multiple determinants of lifespan memory differences. Sci Rep
6:32527 Available at: http://www.nature.com/articles/srep32527.

Hulme OJ, Skov M, Chadwick MJ, Siebner HR, Rams??y TZ (2014) Sparse encoding of
automatic visual association in hippocampal networks. Neuroimage 102:458–464
Available at: http://dx.doi.org/10.1016/j.neuroimage.2014.07.020.

Lancaster JL, Woldorff MG, Parsons LM, Liotti M, Freitas CS, Rainey L, Kochunov P V.,
Nickerson D, Mikiten SA, Fox PT (2000) Automated Talairach Atlas labels for functional
brain mapping. Hum Brain Mapp 10:120–131.

Lang PJ, Bradley MM, Cuthbert BN (1997) International Affective Picture System (IAPS):
Technical Manual and Affective Ratings. NIMH Cent Study Emot Atten:39–58 Available
at:
http://www.unifesp.br/dpsicobio/adap/instructions.pdf%5Cnhttp://econtent.hogrefe.com/
doi/abs/10.1027/0269-8803/a000147.

Li H-J, Hou X-H, Liu H-H, Yue C-L, Lu G-M, Zuo X-N (2015) Putting age-related task activation
into large-scale brain networks: A meta-analysis of 114 fMRI studies on healthy aging.
Neurosci Biobehav Rev 57:156–174.

Li S-C, Li S-C, Lindenberger U, Lindenberger U, Sikstrom S, Sikstrom S (2001) Aging
cognittion: From neuromodulation to representation. 5:479–486.

Maass A, Schütze H, Speck O, Yonelinas A, Tempelmann C, Heinze H-J, Berron D,
Cardenas-Blanco A, Brodersen KH, Enno Stephan K, Düzel E (2014) Laminar activity in
the hippocampus and entorhinal cortex related to novelty and episodic encoding. Nat
Commun   5:5547   Available   at:
http://www.nature.com/doifinder/10.1038/ncomms6547%5Cnhttp://www.pubmedcentral.
nih.gov/articlerender.fcgi?artid=4263140&tool=pmcentrez&rendertype=abstract.

Maillet D, Rajah MN (2014) Age-related differences in brain activity in the subsequent memory
paradigm: A meta-analysis. Neurosci Biobehav Rev 45:246–257 Available at:
http://dx.doi.org/10.1016/j.neubiorev.2014.06.006.

Maldjian JA, Laurienti PJ, Kraft RA, Burdette JH (2003) An automated method for
neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets.
Neuroimage 19:1233–1239.

Morcom AM, Friston KJ (2012) Decoding episodic memory in ageing: A Bayesian analysis of
activity patterns predicting memory. Neuroimage 59.

Morcom AM, Good CD, Frackowiak RSJ, Rugg MD (2003) Age effects on the neural correlates
of successful memory encoding. Brain 126.

Morcom AM, Johnson W (2015) Neural reorganization and compensation in aging. J Cogn
Neurosci 27.

Morcom AM, Li J, Rugg MD (2007) Age effects on the neural correlates of episodic retrieval:
Increased cortical recruitment with matched performance. Cereb Cortex 17.

Nyberg L, Andersson M, Kauppi K, Lundquist A, Persson J, Pudas S, Nilsson LG (2014) Age-
related and genetic modulation of frontal cortex efficiency. J Cogn Neurosci 26:746–764.

Nyberg L, Lövdén M, Riklund K, Lindenberger U, Bäckman L (2012) Memory aging and brain
maintenance. Trends Cogn Sci 16:292–305.

Park DC, Polk TA, Park R, Minear M, Savage A, Smith MR (2004) Aging reduces neural
specialization in ventral visual cortex. 101:13091–13095.

Park DC, Reuter-Lorenz P (2009) The adaptive brain: aging and neurocognitive scaffolding.
Annu   Rev   Psychol   60:173–196   Available   at:
http://www.ncbi.nlm.nih.gov/pubmed/19035823.

Rajah MN, D'Esposito M (2005) Region-specific changes in prefrontal function with age: A
review of PET and fMRI studies on working and episodic memory. Brain 128:1964–1983.

Raz N, Rodrigue KM (2006) Differential aging of the brain: Patterns, cognitive correlates and
modifiers. Neurosci Biobehav Rev 30:730–748.

Reuter-Lorenz PA, Campbell KA (2008) Neurocognitive ageing and the Compensation
Hypothesis. Curr Dir Psychol Sci 17:177–182.

Rouder JN, Speckman PL, Sun D, Morey RD, Iverson G (2009) Bayesian t tests for accepting

and rejecting the null hypothesis. Psychon Bull Rev 16:225–237.

Rypma B, Esposito MD (2000) Isolating the neural mechanisms of age-related changes in
human working memory. 3:509–515.

Shafto M a, Tyler LK, Dixon M, Taylor JR, Rowe JB, Cusack R, Calder AJ, Marslen-Wilson
WD, Duncan J, Dalgleish T, Henson RN, Brayne C, Matthews FE (2014) The Cambridge
Centre for Ageing and Neuroscience (Cam-CAN) study protocol: a cross-sectional,
lifespan, multidisciplinary examination of healthy cognitive ageing. BMC Neurol 14:204
Available   at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4219118&tool=pmcentrez&re
ndertype=abstract.

Smith APR, Henson RNA, Dolan RJ, Rugg MD (2004) fMRI correlates of the episodic retrieval
of emotional contexts. Neuroimage 22:868–878.

Spreng RN, Wojtowicz M, Grady CL (2010) Reliable differences in brain activity between
young and old adults: A quantitative meta-analysis across multiple cognitive domains.
Neurosci Biobehav Rev 34:1178–1194 Available at:
http://dx.doi.org/10.1016/j.neubiorev.2010.01.009.

Taylor JR, Williams N, Cusack R, Auer T, Shafto MA, Dixon M, Tyler LK, Cam-CAN X, Henson
RN (2015) The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) data
repository: Structural and functional MRI, MEG, and cognitive data from a cross-sectional
adult lifespan sample. Neuroimage 144:262–269 Available at:
http://dx.doi.org/10.1016/j.neuroimage.2015.09.018.

Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, Mazoyer
B, Joliot M (2002) Automated anatomical labeling of activations in SPM using a
macroscopic anatomical parcellation of the MNI MRI single-subject brain. Neuroimage
15:273–289.

Venables WN (William N., Ripley BD, Venables WN (William N). (2002) Modern applied
statistics with S.

Wagenmakers E-J (2007) A practical solution to the pervasive problems of p values. 14:779–
804.

West RL (1996) An application of prefrontal cortex function theory to cognitive aging. Psychol
Bull 120:272–292.