Nanopore-based single molecule sequencing of the D4Z4 array

responsible for facioscapulohumeral muscular dystrophy

Satomi Mitsuhashi[1,2], So Nakagawa[1,3], Mahoko Ueda[1,3], Tadashi Imanishi[1],

Hiroaki Mitsuhashi[4]

1.   Biomedical Informatics Laboratory, Department of Molecular Life Science,

Tokai University School of Medicine, Isehara, Kanagawa 259-1193, Japan

2.   Department of Human Genetics, Yokohama City University

Graduate School of Medicine, Yokohama, Kanagawa 236-0004, Japan

3.   Micro/Nano Technology Center, Tokai University, Hiratsuka, Kanagawa,

Japan

4.   Department of Applied Biochemistry, School of Engineering, Tokai

University, Hiratsuka, Kanagawa, Japan

Corresponding author

Satomi Mitsuhashi

Department of Human Genetics Yokohama City University Graduate School of

Medicine, Yokohama, Kanagawa 236-0004, Japan

Tel. +81-45-787-2606

Fax. +81-45-786-5219

E-mail: satomits@yokohama-cu.ac.jp

Abstract

Subtelomeric macrosatellite repeats are difficult to sequence using conventional

sequencing methods owing to the high similarity among repeat units and high

GC content. Sequencing these repetitive regions is challenging, even with

recent improvements in sequencing technologies. Among these repeats, a

haplotype of the telomeric sequence and shortening of the D4Z4 array on

human chromosome 4q35 causes one of the most prevalent forms of muscular

dystrophy with autosomal-dominant inheritance, facioscapulohumeral muscular

dystrophy (FSHD). Here, we applied a nanopore-based ultra-long read

sequencer to sequence a BAC clone containing 13 D4Z4 repeats and flanking

regions. We successfully obtained the whole D4Z4 repeat sequence, including

the pathogenic gene *DUX4* in the last D4Z4 repeat. The estimated sequence

accuracy of the total repeat region was 99.7% based on a comparison with the

reference sequence. Errors were typically observed between purine or between

pyrimidine bases. Further, we analyzed the D4Z4 sequence from publicly

available ultra-long whole human genome sequencing data obtained by

nanopore sequencing. This technology may become a new standard for the

molecular diagnosis of FSHD in the future and has the potential to widen our

understanding of subtelomeric regions.

Introduction

Facioscapulohumeral muscular dystrophy (FSHD) is one of the most

prevalent adult-onset muscular dystrophies. The genomes of most patients with

FSHD have a common feature, i.e., a contracted subtelomeric macrosatellite

repeat array called D4Z4 on chromosome 4q35. The D4Z4 array consists of a

highly similar 3.3-kb single repeat unit. Normally, the D4Z4 array is highly

methylated and forms heterochromatin. Patients with FSHD have less than 11

D4Z4 repeats (1-3). In Japan, the majority of patients with FSHD have less than

7 repeats (4). Shortening of the D4Z4 array causes the de-repression of the

flanking genes as well as *DUX4*, located in the last D4Z4 repeat. The ectopic

expression of *DUX4* is toxic in muscle tissues and is thought to be a causal

factor for FSHD (5-9). In addition to the repeat number, the haplotype of the last

D4Z4 repeat is important for the development of FSHD (1, 2). The telomeric

flanking region of D4Z4 contains the 3′ UTR of *DUX4* and is called the pLAM

region. The presence of a polyadenylation signal in this region allows *DUX4*

expression and disease manifestation (10). In contrast, individuals without

polyadenylation signals do not manifest the disease (2).

Molecular diagnosis of FSHD is commonly made by Southern blotting

of genomic DNA after restriction enzyme digestion to measure the D4Z4 array

length and estimate the number of repeats. Haplotype analysis requires a

different probe (1). Sequencing of this D4Z4 array using Sanger sequencing or

short-read sequencers (up to 600 bp for Illumina and IonTorrent) is technically

difficult owing to the high similarity and the high GC content of the repeats. The

Oxford Nanopore Technologies MinION (Oxford, UK) is a single-molecule

sequencer that can produce long reads exceeding 100 kbp (11). Therefore,

MinION sequencing may enable the determination of pathogenicity by

sequencing the complete D4Z4 array.

Results

Nanopore-based D4Z4 sequencing using a BAC clone

The D4Z4 array on 4q35 has *Eco*RI sites in its flanking region. We

took advantage of this restriction enzyme to excise the full-length D4Z4 repeats

with flanking sequences, for a total of 49,877 bp. Both sides of the *EcoRI*-

digested DNA fragment had unique sequences that are not found in the D4Z4

repeats (4,823 bp on centromeric side and 865 bp on the telomeric side). RP11-

242C23 contained multiple *Eco*RI sites. pBACe3.6 vector-derived DNA was

digested, yielding fragments of less than 10 kb (Figure 1a). We were able to

easily separate the D4Z4-containing DNA fragment (49877 bp) from vector-

derived DNA by agarose gel electrophoresis and gel extraction (Figure 1b). We

extracted the D4Z4 array-containing DNA and subjected it to MinION 1D

sequencing (Oxford Nanopore Technologies, Oxford, UK). Base-calling was

initially performed using MinKNOW ver. 1.5.12 and fastq conversion was

performed using poretools to obtain 20,761 reads (12). Base-calling was not

possible for 87,410 reads using real-time MinKNOW basecaller probably due to

out of computer memory; we used Albacore (v.1.1.0) to obtain the fastq

sequences in these cases. A total of 128,171 reads were obtained, with an

average read length of 7,577 bp (Supplemental Table 1).

We mapped the reads to the reference BAC clone sequence

(GenBank accession number CT476828.7) using LAST (13). Visualization of

mapped reads using IGV showed coverage of the whole D4Z4 array (Figure 2).

The longest read mapped to the D4Z4 repeat was 29,060 bp. The consensus

sequence had an accuracy of 99.72%. We also used BWA-MEM for mapping

and found that the consensus sequence had a lower accuracy (99.18%). Thus,

we used LAST for subsequent analyses.

The haplotype of the telomeric flanking region of the final D4Z4 repeat

known as pLAM is important for disease manifestation. There are two equally

common haplotypes, A and B. Haplotype A has an added polyadenylation signal

at the 3′ UTR of the *DUX4* gene(10). This polyadenylation signal allows the

ectopic expression of *DUX4*, which is toxic in muscle cells of patients with

FSHD with the contracted D4Z4 array (14). Haplotype B lacks homologous

sequence to pLAM. Individuals with haplotype B do not manifest the disease,

despite having the contracted D4Z4 allele. Thus, it is important to identify the

pLAM sequence for the molecular diagnosis of FSHD. Using MinION, we

successfully sequenced the whole pLAM region with an accuracy of 100%

(Figure 3). In total, 135 bases were different from the reference genome

sequence among the whole D4Z4 array sequence of 49,877 bp (0.27%)

(Supplemental Figure 1a). Among 135 bases, 115 (85.2%) substitutions were

between purines or between pyrimidines (Supplemental Figure 1b). Most of

these errors were repeatedly detected at the same position in the repeats

(indicated by asterisks in Supplemental Figure 1a). Interestingly, 16 out of 18

recurrent errors were seen in the CCXGG sequence at the X position.

We also compared the nanopore-sequenced *DUX4* open-reading

frame (ORF) to the reference and the Sanger sequencing results for the

subcloned *DUX4* ORF. The accuracy of the *DUX4* ORF sequence was 99.9%

(Supplemental Figure 2) and there were 3 errors. These errors were also

located in the X position in the CCXGG sequence.

D4Z4 detection using nanopore-based whole human genome sequencing

We tested whether we can identify the D4Z4 array from whole genome

sequencing data obtained from the MinION sequencer. We used the publicly

available human reference standard genome NA12878 with R9.4 chemistry

(11). This project contains two sets of data. The rel3 dataset had approximately

26× coverage with an N50 length of 10.6 kb. Rel4 had 5× coverage of ultra-long

reads with an N50 of 99.7 kb, indicating that rel4 contained reads that possibly

cover the whole D4Z4 region. We performed a blastn similarity search against

rel4 reads using the pLAM region as a query and obtained 18 hits (e-value =

0.0). These reads were aligned to the D4Z4 repeat reference sequence with the

pLAM region. The consensus sequence identity to the last D4Z4 repeat and the

pLAM region was 96.3% (Figure 4). Among 18 reads, 2 had homologous

sequences to both centromeric and telomeric flanking regions of D4Z4 (Figure

2, Supplemental Table 2). These two reads are expected to cover the whole

D4Z4 array, which usually includes more than 16 D4Z4 repeats. Although, we

could not determine the exact number of D4Z4 repeats owing to the high rate of

deletion errors, this range of read fragment is capable of detecting contracted

D4Z4 array which is seen in the most of the FSHD patients.

Discussion

Sequencing a highly repetitive subtelomeric region is extremely

challenging. There is variation in the number of repeats among individuals and

sometimes within individuals, i.e., somatic mosaicism. It has been reported that

subtelomeric regions form heterochromatin, functioning as an insulator or

repressor of near-by genes or preventing telomere shortening (15, 16). It is

important to determine the relationship between phenotypic differences and

either sequence or structural variation in repeats not only to decipher the

pathomechanisms of the disease, but also to obtain a deeper understanding of

human genomes. Here, we applied a nanopore-based sequencer to investigate

the subtelomeric repeat array associated with FSHD for the first time. In the

near future, it will be feasible to search for these sequence-difficult regions to

find causal relationships between the human genome and genetic diseases;

even given the prevailing use of high-throughput sequencing of coding regions,

the genetic causes of many diseases remain unsolved.

The disease locus of FSHD was identified at 4q35 more than 20 years

ago; however, the mechanism underlying the disease has been a mystery for

years and the causative genes have not been identified until recently, when

accumulating evidence has shown that the misexpression of *DUX4* is

associated with the disease. Further, it is still unclear whether there is any

sequence polymorphism in the *DUX4* gene or flanking regions, as it is difficult to

sequence the gene or the *DUX4* transcript, which is expressed at the very low

levels even in the muscle tissues of patients (14, 17). Since therapeutic

approaches including nucleic acid drugs targeting *DUX4* mRNA are being

studied (18, 19), it may be useful to determine the exact *DUX4* sequence of

patients for the development of effective therapies as well as an integrative

diagnostic method.

Currently, the number of D4Z4 repeats is usually determined by

Southern blotting using a probe that hybridizes to the centromeric flanking

sequence, p13E-11 (1). If the patient has a deletion at this probe site, it is not

possible to detect the D4Z4 repeat by Southern blotting. The Southern blotting

technique is complicated and time-consuming. Alternative methods have been

investigated, but are not widely used (4, 20).

Morioka et al. sequenced D4Z4 using the PacBio sequencer (21) and

analyzed random fragments from the BAC clone. The advantage of the

nanopore sequencer over the PacBio sequencer is the ultra-long read capability

(11). It has the potential to obtain reads of more than 100 kbp, the approximate

mean size of D4Z4 in healthy individuals. Currently, we could only obtained two

reads that potentially cover all D4Z4 repeats from human genome data with 5×

coverage using 14 flow-cells (11). Our estimate of the number of D4Z4 repeats

was more than 16, the normal size observed in healthy individuals. As the

NA12878 standard DNA originated from healthy individuals without FSHD, this

repeat number is reasonable. As FSHD patients have D4Z4 number less than

11, we think this ultra-long read sequencing may be usable to detect the

disease-causing contracted D4Z4 array. If the data output for the MinION

sequencer improves, it will be possible to obtain sequence data with better

resolution. This approach is potentially applicable to subtelomeric regions of

other chromosomes or even to centromere sequences.

In nanopore-based sequencing, changes in electric current are

detected as nucleotides pass through the pore. We observed that the errors

tend to occur between purines or between pyrimidines, probably because they

have similar chemical structures (11). In addition, we also observed that

substitution errors tend to occur at the same nucleotide position across repeats

(Supplemental Figure 1, asterisk). This may reflect the fact that the nanopore

detects combinations of nucleotides and the specific combination CCXGG was

prone to be misread. We anticipate further improvements of the base-calling

algorithm, which will make MinION more beneficial for medical applications.

Sequencing technologies are continuously developed. During the

preparation of this manuscript, the new chemistry R9.5 with the new flow-cell

FLO-MIN107 was released. Considering the rapid improvements in this

technique, it may not be very long before this sequencing technology is used for

D4Z4 repeat analyses for patients with FSHD.

Conclusions

Using MinION with a R9.4 flow-cell and 1D sequencing chemistry, we

successfully sequenced the complete *EcoRI*-digested D4Z4 array from a BAC

clone that contained the D4Z4 repeat region of human chromosome 4. Our

deep sequencing results had an accuracy of 99.8% for the whole D4Z4 array

and flanking region. This includes the pLAM region, with an accuracy of 100%,

and the whole ORF of the pathogenic gene *DUX4,* with the accuracy of 99.9%,

which are important regions for determining the pathogenesis. This short report

may provide a basis for the future use of nanopore sequencing to deepen our

understanding of highly heterogenous subtelomeric regions that may contribute

to human disease.

Materials and Methods

BAC clone

The RP11-242C23 human BAC clone was obtained from BAC PAC Resources

Center (https://bacpaacresources.org). This BAC clone was sequenced and

deposited at GenBank under accession number CT476828.7 by the Wellcome

Trust Sanger Institute. It contained 13 3306-bp D4Z4 repeats.

Preparation of D4Z4 repeats from the BAC clone

RP11-242C23 was digested using EcoRI and treated with Klenow Fragment

DNA Polymerase (Takara, Shiga, Japan) at 37°C for 20 min. DNA was

subjected to electrophoresis on a 0.5% agarose gel. Bands larger than the 10-

kb marker (GeneRuler 1kb DNA Ladder; Thermo Fisher Scientific, Waltham,

MA, USA) were excised using a razor under ultraviolet light. The DNA

fragments larger than 1 kb were subjected to phenol-chloroform DNA

preparation. Agarose gels were soaked in phenol and incubated for 30 min at -

80°C. Then, the aqueous phase was collected and phenol-chloroform DNA

preparation was performed. The EcoRI-digested whole D4Z4 repeat was

enriched in the DNA sample.

MinION 1D sequencing

Library preparation was performed using a SQK-LSK108 Sequencing Kit R9.4

version (Oxford Nanopore Technologies, Oxford, UK) using 500 ng of DNA.

MinION sequencing was performed using one FLO-MIN106 (R9.4) flow cell with

the MinION MK1b sequencer (Oxford Nanopore Technologies). Base-calling

and fastq conversion were performed with MinKNOW ver. 1.5.12 followed by

poretools or Albacore.

Sequence alignment by LAST and BWA-MEM

Sequence reads were aligned to the EcoRI-digested D4Z4 repeat reference

(Figure 1b, Supplemental material) using the LAST aligner (13). Sequences

were also mapped to the reference genome hg19 using BWA-mem with default

settings (22). Consensus sequences were obtained and sequence identity was

calculated using UGENE (23). Mapped reads were visualized using IGV

software (24).

Subcloning of the last D4Z4 repeat

An *Escherichia coli* transformant with the RP11-242C23 human BAC clone was

cultured in LB medium containing 12.5 μg/ml chloramphenicol at 37°C The

human BAC clone DNA was purified using the QIAGEN Plasmid Midi Kit

(Hilden, Germany) according to the "User-Developed Protocol (QP01)." Briefly,

bacterial lysate from a 100-ml scale culture was passed through a QIAGEN-tip

100 column. The BAC clone DNA was eluted with buffer QF prewarmed to 65°C

and concentrated by isopropanol precipitation.

To obtain the DNA clone containing the last D4Z4 repeat with the pLAM region,

50 ng of the purified BAC clone was used as a template for PCR with the

forward primer 5′-cgcgtccgtccgtgaaattcc-3′ and the reverse primer 5′-

caggggatattgtgacatatctctgcac-3′. PCR was performed with PrimeSTAR GXL

DNA Polymerase (Takara) with the following cycling conditions: 98°C for 2 min

and 30 cycles of 98°C for 10 s, 60°C for 15 s, and 68°C for 30 min. PCR

products were gel-purified and cloned into a pCR blunt vector (ThermoFisher

Scientific) with the Mighty Mix DNA Ligation Kit (Takara). The sequence of the

resulting plasmid was confirmed by Sanger sequencing with M13 forward and

M13 reverse primers.

D4Z4 sequence analysis using the ultra-long human whole genome

sequence

The human whole genome sequenced by MinION sequencers was downloaded

(https://github.com/nanopore-wgs-consortium/NA12878) (11). A Blastn search

was performed against the ultra-long read dataset, rel4, using the pLAM

sequence as a query. A total of 18 sequence reads containing pLAM hits were

extracted (e-value = 0.0). These extracted reads were mapped to the D4Z4

reference sequence using LAST and consensus sequences for the last D4Z4

and pLAM regions were obtained as described above.

References

1.   Wijmenga C, Hewitt JE, Sandkuijl LA, Clark LN, Wright TJ, Dauwerse HG,

et al. Chromosome 4q DNA rearrangements associated with

facioscapulohumeral muscular dystrophy. Nat Genet. 1992;2(1):26-30.

2.   Lemmers RJ, de Kievit P, Sandkuijl L, Padberg GW, van Ommen GJ,

Frants RR, et al. Facioscapulohumeral muscular dystrophy is uniquely

associated with one of the two variants of the 4q subtelomere. Nat Genet.

2002;32(2):235-6.

3.   van Deutekom JC, Wijmenga C, van Tienhoven EA, Gruter AM, Hewitt JE,

Padberg GW, et al. FSHD associated DNA rearrangements are due to

deletions of integral copies of a 3.2 kb tandemly repeated unit. Hum Mol

Genet. 1993;2(12):2037-42.

4.   Goto K, Nishino I, Hayashi YK. Rapid and accurate diagnosis of

facioscapulohumeral muscular dystrophy. Neuromuscul Disord.

2006;16(4):256-61.

5.   Mitsuhashi H, Mitsuhashi S, Lynn-Jones T, Kawahara G, Kunkel LM.

Expression of DUX4 in zebrafish development recapitulates

facioscapulohumeral muscular dystrophy. Hum Mol Genet. 2013;22(3):568-

77.

6.   Kowaljow V, Marcowycz A, Ansseau E, Conde CB, Sauvage S, Matteotti C,

et al. The DUX4 gene at the FSHD1A locus encodes a pro-apoptotic

protein. Neuromuscul Disord. 2007;17(8):611-23.

7.   Snider L, Asawachaicharn A, Tyler AE, Geng LN, Petek LM, Maves L, et al.

RNA transcripts, miRNA-sized fragments and proteins produced from D4Z4

units: new candidates for the pathophysiology of facioscapulohumeral

dystrophy. Hum Mol Genet. 2009;18(13):2414-30.

8.   Wallace LM, Garwick SE, Mei W, Belayew A, Coppee F, Ladner KJ, et al.

DUX4, a candidate gene for facioscapulohumeral muscular dystrophy,

causes p53-dependent myopathy in vivo. Ann Neurol. 2011;69(3):540-52.

9.   Wuebbles RD, Long SW, Hanel ML, Jones PL. Testing the effects of FSHD

candidate gene expression in vertebrate muscle development. Int J Clin

Exp Pathol. 2010;3(4):386-400.

10.   Lemmers RJ, van der Vliet PJ, Klooster R, Sacconi S, Camano P,

Dauwerse JG, et al. A unifying genetic model for facioscapulohumeral

muscular dystrophy. Science. 2010;329(5999):1650-3.

11.   Jain M, Koren S, Quick J, Rand AC, Sasani TA, Tyson JR, et al. Nanopore

sequencing and assembly of a human genome with ultra-long reads.

BioRxiv. 2017. **doi:** https://doi.org/10.1101/128835

12.   Loman NJ, Quinlan AR. Poretools: a toolkit for analyzing nanopore

sequence data. Bioinformatics. 2014;30(23):3399-401.

13.   Kielbasa SM, Wan R, Sato K, Horton P, Frith MC. Adaptive seeds tame

genomic sequence comparison. Genome Res. 2011;21(3):487-93.

14.   Snider L, Geng LN, Lemmers RJ, Kyba M, Ware CB, Nelson AM, et al.

Facioscapulohumeral dystrophy: incomplete suppression of a

retrotransposed gene. PLoS Genet. 2010;6(10):e1001181.

15.   Ottaviani A, Rival-Gervier S, Boussouar A, Foerster AM, Rondier D,

Sacconi S, et al. The D4Z4 macrosatellite repeat acts as a CTCF and A-

type lamins-dependent insulator in facio-scapulo-humeral dystrophy. PLoS

Genet. 2009;5(2):e1000394.

333 16. Stadler G, Rahimov F, King OD, Chen JC, Robin JD, Wagner KR, et al.

334   Telomere position effect regulates DUX4 in human facioscapulohumeral

335   muscular dystrophy. Nat Struct Mol Biol. 2013;20(6):671-8.

336 17. Jones TI, Chen JC, Rahimov F, Homma S, Arashiro P, Beermann ML, et al.

337   Facioscapulohumeral muscular dystrophy family studies of DUX4

338   expression: evidence for disease modifiers and a quantitative model of

339   pathogenesis. Hum Mol Genet. 2012;21(20):4419-30.

340 18. Ansseau E, Vanderplanck C, Wauters A, Harper SQ, Coppee F, Belayew A.

341   Antisense oligonucleotides used to target the DUX4 mRNA as therapeutic

342   approaches in faciosscapulohumeral muscular dystrophy (FSHD). Genes.

343   2017;8(3).

344 19. Wallace LM, Liu J, Domire JS, Garwick-Coppens SE, Guckes SM, Mendell

345   JR, et al. RNA interference inhibits DUX4-induced muscle toxicity in vivo:

346   implications for a targeted FSHD therapy. Mol Ther. 2012;20(7):1417-23.

347 20. Vasale J, Boyar F, Jocson M, Sulcova V, Chan P, Liaquat K, et al. Molecular

348   combing compared to Southern blot for measuring D4Z4 contractions in

349   FSHD. Neuromuscul Disord. 2015;25(12):945-51.

350 21. Morioka MS, Kitazume M, Osaki K, Wood J, Tanaka Y. Filling in the gap of

351   human chromosome 4: single molecule real time sequencing of

352   macrosatellite repeats in the facioscapulohumeral muscular dystrophy

353   locus. PLoS One. 2016;11(3):e0151963.

354 22. Li H. Aligning sequence reads, clone sequences and assembly

355 23. contigs with BWA-MEM. arXiv. 2013;arXiv:1303.3997.

24. Okonechnikov K, Golosova O, Fursov M, UGENE team. Unipro UGENE: a

unified bioinformatics toolkit. Bioinformatics. 2012;28(8):1166-7.

25. Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz

G, et al. Integrative genomics viewer. Nat Biotechnol. 2011;29(1):24-6.

Author Contributions

SM and HM designed the study and collected experimental materials.

SM, SN, MU, HM, and TI analyzed and interpreted the data. SM drafted the

original manuscript.

Competing interests

The authors report no disclosures relevant to the manuscript.

Web Resources

LAST: http://last.cbrc.jp

BWA: http://bio-bwa.sourceforge.net

Ape: http://biologylabs.utah.edu/jorgensen/wayned/ape/

UGENE: http://ugene.net

The URL for the human whole genome sequence data for NA12878 used in this

study is as follows:

https://github.com/nanopore-wgs-consortium/NA12878

Funding

This study was supported by MEXT-Supported Program for the Strategic

Research Foundation at Private Universities (to S.N. and M.T.U.). This work

was supported by JSPS KAKENHI Grant Number JP15K19477 (to H.M).

Figure legends

Figure 1

(a) Vector map of RP11-242C23 generated using Ape software. EcoRI sites are

shown. The D4Z4 array with 13 repeats and flanking regions was excised using

EcoRI digestion, yielding a 49877-bp product. (b) Agarose gel electrophoresis

of the EcoRI-digested vector DNA. Arrow shows the band of the 49877-bp D4Z4

array.

Figure 2

Mapped reads were visualized using IGV software. Coverage of reads is shown

on the upper part of the IGV image. Scheme shows the 13 D4Z4 repeats with

flanking sequences. The bottom scheme shows the enlarged last D4Z4 repeat

with the pLAM region (haplotype A). This region encodes pathogenic *DUX4*.

Figure 3

Nanopore sequence of the pLAM region. Exon 3 of *DUX4*, the 3′ UTR of the

gene, and polyA signal were determined with an accuracy of 100%. The upper

sequence is the reference and the bottom shows the nanopore sequence.

405   Figure 4

406   The consensus sequence of the last D4Z4 repeat and pLAM region obtained by

407   whole human genome nanopore sequencing. The upper sequence is the

408   reference and the lower sequence is the publicly available rel4 data set.

Figure 1

a

b

Figure 2

Figure 3

Figure 4

The last D4Z4

pLAM