1
7 **Variation in neuronal activity state, axonal projection target, and position principally define**
8 **the transcriptional identity of individual neocortical projection neurons**
9
10

11 *Maxime Chevée[1,2], Johanna D. Robertson[3], Gabrielle H. Cannon[4]*
12 *Solange P. Brown[2,*] and Loyal A. Goff[2,4,*]*

Affiliations:
[1]Biochemistry, Cellular and Molecular Biology Graduate Program, Johns Hopkins University
School of Medicine, Baltimore, Maryland, 21205, USA.

[2]Solomon H. Snyder Department of Neuroscience, Johns Hopkins University School of Medicine,
Baltimore, Maryland, 21205, USA.

[3]Human Genetics Training Program, McKusick-Nathans Institute for Genetic Medicine, Johns
Hopkins University School of Medicine, Baltimore, Maryland, 21205, USA.

[4]McKusick-Nathans Institute for Genetic Medicine, Johns Hopkins University School of Medicine,
Baltimore, Maryland, 21205, USA.

[*]Corresponding Authors: spbrown@jhmi.edu (SPB) and loyalgoff@jhmi.edu (LAG)

ABSTRACT

Single-cell RNA sequencing technologies have generated the first catalogs of transcriptionally

defined neuronal subtypes of the brain. However, the biologically informative cellular

processes that contribute to neuronal subtype specification and transcriptional heterogeneity

remain unclear. By comparing the gene expression profiles of single layer 6 corticothalamic

neurons in somatosensory cortex, we show that transcriptional subtypes primarily reflect

axonal projection pattern, laminar position within the cortex, and neuronal activity state.

Pseudotemporal ordering of 1023 cellular responses to manipulation of sensory input

demonstrates that changes in expression of activity-induced genes both reinforced cell-type

identity and contributed to increased transcriptional heterogeneity within each cell type. This

is due to cell-type specific biases in the choice of transcriptional states following manipulation

of neuronal activity. These results reveal that axonal projection pattern, laminar position, and

activity state define significant axes of variation that contribute both to the transcriptional

identity of individual neurons and to the transcriptional heterogeneity within each neuronal

subtype.

INTRODUCTION

Single-cell RNA sequencing (scRNA-seq) approaches have been harnessed to reveal

previously hidden levels of complexity in cell types and states within a given tissue[1, 2]. Nowhere

is this more relevant than in the mammalian central nervous system, an organ system

dependent on a remarkable diversity of cell types and cell states for its function. The neocortex

contains a wide variety of neuronal cell types organized into the circuits that direct higher brain

functions such as perception, memory and cognition. Furthermore, cortical neurons are highly

dynamic, undergoing significant changes in gene expression during development and

throughout adulthood in response to activity and experience[3-8]. Recent studies have generated a

more comprehensive 'parts list' of discrete cell types within the neocortex than previously

available[9-16]. While such surveys yield insights into the diversity of cortical cell types, the sources

of transcriptional variation both within and across cell types remain poorly understood[17-19].

We compared expression profiles of layer 6 corticothalamic neurons (L6CThNs), a

heterogeneous population of related cortical projection neurons defined in anatomical,

functional, and gene expression studies, making them ideally suited for investigating the

relationships between transcriptional subtypes and other cellular properties[12, 16, 20-29]. By

combining scRNA-seq with an enrichment strategy that preserved axonal target information, we

identified two transcriptionally distinct L6CThN subtypes that each demonstrates a reproducible

bias for long-range projection targets. The transcriptional profiles of L6CThNs also reflect their

laminar position within L6. These two L6CThN subtypes exhibit divergent signatures of neuronal

activity both in their baseline expression patterns as well as in their responses following

manipulation of sensory input. The subtype biases in the choice of response following sensory

manipulation increased transcriptional heterogeneity both within and between cell types and

reinforced the transcriptional identities of the two L6CThN subtypes. These results demonstrate

that scRNA-seq resolves relationships between gene expression and features such as axonal

projection pattern, spatial organization, and cell state, and identifies the independent

contributions of multiple biological signals that together determine transcriptional

heterogeneity within and across neuronal populations.

RESULTS

Transcriptional profiling of layer 6 corticothalamic neurons reveals two subtypes which reflect

axonal projection bias

Studies of primary sensory cortex demonstrate that L6CThNs are heterogeneous[20-29]. In rat

barrel cortex, L6CThNs in upper layer 6 project to the ventral posterior medial nucleus (VPM) of

the thalamus, while L6CThNs in lower layer 6 project to both VPM and the posterior medial

nucleus (POm)[21, 25]. To distinguish between these two projection classes in mouse, we first

validated Cre recombinase expression as a reliable marker for L6CThNs in the barrel cortex of

Neurotensin receptor 1-Cre mice[30-32] (Fig. 1a-e; *Ntsr1-Cre*, Gensat 220). Next, we showed that a

subset of Cre-expressing, VPM-projecting L6CThNs in lower layer 6 also projects to POm (Fig. 1f-

j).

87   To determine whether L6CThNs subclasses projecting to VPM only or to both VPM and

POm are distinguished by their gene expression profiles, we labeled the two subclasses in

adolescent mice, microdissected barrel cortex containing retrogradely labeled cells, dissociated

the tissue into a single-cell suspension, sorted the differentially labeled L6CThNs using

Fluorescence Assisted Cell Sorting (FACS; Supplementary Fig. 1a-b), and collected enriched

populations of each subclass for bulk RNA sequencing. We identified 69 genes with significant

differential expression (DE) between the two sorted populations (Fig. 1k, Supplementary File 1,

Supplementary Table 1; Cuffdiff2; 10% false discovery rate (FDR)). These 69 DE genes

demonstrate that subclasses of L6CThNs distinguished by their long-range axonal projection

patterns are differentiated by their gene expression profiles.

This bulk analysis was predicated on prior knowledge of existing morphological

subclasses and may have obscured underlying transcriptional subtypes that comprise each

projection class of L6CThNs. To resolve transcriptionally defined subtypes of L6CThNs, we next

evaluated the gene expression landscape of single L6CThNs using an unbiased classification

approach. We first sorted individual, labeled L6CThNs (Supplementary Fig. 1c) and collected 96

VPM-only and 96 VPM/POm L6CThNs from each of two replicate mice, totaling 384 single

L6CThNs. Individual cell lysates were subjected to a modified Smart-Seq2 library preparation

and scRNA-seq analysis. In total, 346 single L6CThNs passed quality control filters and were used

for further analysis (Supplementary Fig. 1d-i; Supplementary File 2). We confirmed the fidelity of

our enrichment by assessing each cell for neuronal and non-neuronal markers (Supplementary

Fig. 1j).

To identify transcriptional subtypes of L6CThNs, we selected genes with the greatest

likelihood of contributing to differences across the ensemble of single L6CThNs by identifying

genes with high residuals to a mean-variance model fit independently in each of the two

replicates and selecting the intersection of these two sets (Supplementary Fig. 2a). These 261

genes with high dispersion were therefore depleted for genes associated with technical

variation between replicates[33, 34] (Supplementary Table 2). Weights on the first three significant

principal components across all cells using this gene set[35] (Supplementary Fig. 2b; permutation

parallel analysis; p < 0.001) were used for a tSNE dimensionality reduction followed by k-means

clustering (Fig. 2a). Single-cell transcriptional profiles of the 346 L6CThNs clustered into at least

two distinct subtypes. Fitting the data to three subtypes (k=3) or more did not significantly

improve the performance of the clustering (Supplementary Fig. 2c). We compared our

classification approach to several recently described single-cell clustering utilities[36, 37] and found

a high-degree of agreement (Supplementary Fig. 2d; SC3: 93.77% agreement; CIDR: 90%

agreement). These results indicate that independent, unbiased clustering approaches based on

genes with higher than expected variance across the L6CThN population identify two major

subtypes of L6CThN.

To determine the relationship between transcriptional identity and morphological

subtypes, we next compared the distribution of VPM-only and VPM/POm projection labels

across the two transcriptionally defined subtypes (Fig. 2b). The majority of neurons in subtype 2

were labeled VPM-only (79%, 103 of 130 cells, $p < 3.768 \times 10^{-17}$, hypergeometric test) while most

neurons in subtype 1 were VPM/POm (65%, 141 of 216 cells, $p < 3.768 \times 10^{-17}$, hypergeometric

test), a distribution significantly different from that expected by chance. Furthermore, the

axonal projection bias was reproducible across replicates (Fig. 2c) even though the segregation

of projection-defined subclasses was incomplete for both transcriptionally defined subtypes.

Together our results indicate that each transcriptionally defined subtype of L6CThN is enriched

for neurons targeting specific sets of thalamic nuclei.

Transcriptional differences between subtypes of layer 6 corticothalamic neurons

To assess the transcriptional differences between the two identified L6CThN subtypes, all

expressed genes were subjected to the Monocle2 differential test, using model formulas that

accounted for both batch effects and the number of genes detected in each cell, a proxy

measure for efficiency of RNA capture and library synthesis. We identified 286 genes that were

significantly differentially expressed between the two subtypes (Fig. 2d; Monocle likelihood

ratio test, 0.1% FDR; Supplementary File 2, Supplementary Table 3), only six of which overlapped

with the 69 DE genes observed in our bulk RNA-seq analysis of sorted L6CThNs despite the high

correlation between our bulk RNA-seq and scRNA-seq data (Supplementary Fig. 2e-g). This

result, in conjunction with our previous observation of incomplete label segregation across

L6CThN cell types, suggests that this discrepancy is primarily due to significant sample

heterogeneity arising from retrograde label inefficiencies. Importantly, parameters such as total

mapped fragments (mass), total estimated mRNAs per cell, number of genes detectably

expressed per cell, and replicate did not result in biased clustering across the DE gene list (Fig.

2d), suggesting a minimal influence of technical variation on the list of DE genes. The significant

differential expression of genes, including *Fxyd6* and *Lamp*5, between the two subtypes was

consistent with expression patterns seen in the Allen Mouse Brain Atlas[38] (Fig. 2e;

http://mouse.brain-map.org, Fxyd6-RP_051017_01_E10-coronal, Lamp5-RP_050725_01_B03-

coronal). The two subtypes also shared some transcriptional similarities with two recently

defined subtypes in primary visual cortex[12] (Supplementary Fig. 3). Several long non-coding

RNAs (lncRNAs) were also specifically enriched in each subtype. For example, linc-Tmem20[8] was

significantly enriched in subtype 1 L6CThNs (Fig. 2f) and was preferentially expressed in lower

layer 6 (Fig. 2g). Conversely, the lncRNA *Pantr1* was the gene with the greatest predictive power

for neurons in subtype 2 (Fig. 2f, AUC = 0.876, power = 0.752, ROC analysis). Using a mouse line

in which *LacZ* was knocked into the *Pantr1* locus[39], we confirmed greater *LacZ* expression in

VPM-only L6CThNs relative to VPM/POm L6CThNs (Fig. 2h,i). Furthermore, in contrast to our

analysis of L6CThNs by anatomical labeling, we found that the total number of genes with

detectable expression in each transcriptional subtype was significantly greater for subtype 1

relative to subtype 2, a finding consistent across replicates (Fig. 2j), and that the mean pairwise

Euclidean distance between cells within each subtype was also greater in subtype 2 than in

subtype 1, indicating greater cell-cell variation across subtype 2 (Fig. 2k; subtype 1 μ=72.16,

σ=5.80; subtype 2 μ=80.59, σ=7.07; $p < 2.2 \times 10^{-16}$, Welch's two-sample t-test). Importantly, we

found only two genes, *Lypd1* and *Calm2*, with a significant combinatorial effect of subtype and

label, suggesting that label does not distinguish subpopulations within each subtype. Together,

these data identify two L6CThN subtypes and confirm a relationship between transcriptionally

defined neuronal subtypes and the projection targets of these neurons.

To identify cellular processes that differentiate the two transcriptional subtypes of

L6CThNs, we queried the DE gene list for enrichment of annotated gene sets from public

databases (Supplementary Fig. 4a-d). Significant gene sets included Gene Ontology and

Reactome terms related to general features of neurons such as "Neuronal part" and "Synaptic

transmission" ($p < 0.01$; hypergeometric test, Benjamini-Hochberg corrected), highlighting the

limited resolution of gene sets in currently available public databases for generating biological

insights among neuronal subtypes. To identify more informative biological processes that shape

differences in the response properties of L6CThNs, we compared the expression of voltage-

gated ion channels, neurotransmitter receptors and neuropeptides, a number of which were

differentially expressed (Supplementary Fig. 4e-h). Several G protein-coupled receptors and

neuropeptides were differentially expressed. For example, *Adcyap1* (PACAP) and a gene

encoding a peptide for processing PACAP, *Pam*, were preferentially expressed in subtype 2

(Supplementary Fig. 4h-i). Interestingly, receptors for PACAP are found in primary sensory

thalamic nuclei[40] and modulate thalamocortical interactions[41], consistent with the projection

pattern bias of subtype 2 L6CThNs. Our data indicate that a focused analysis identifies gene

expression differences that reflect relevant functional features.

Distinct cellular processes are coordinately regulated within subtypes of L6CThN

To identify cellular processes that contribute to the heterogeneity of gene expression across the

transcriptomes of all L6CThNs analyzed, we performed a weighted gene co-expression network

analysis (WGCNA) on all genes expressed in the 346 L6CThNs[42]. This analysis yielded 22 modules

of co-regulated genes, which we classified using hierarchical clustering (Fig. 3a, Supplementary

Fig. 5a). We tested for significant correlations between subtype identity for each cell and the

eigenvalues of each module. Three modules significantly correlated with subtype 1 (Black,

Turquoise, and Cyan; Fig. 3b, Supplementary Fig. 5b; $p < 0.01$; Pearson's product moment

correlation test) and four with subtype 2 (Red, Purple, Blue and Midnight Blue; Fig. 3b,

Supplementary Fig. 5b; $p < 0.01$; Pearson's product moment correlation test). When the same

analysis was performed based on cell label, VPM-only or VPM/POm, correlation coefficients and

confidence measures were all weakened (Fig. 3b, Supplementary Fig. 5c). These seven modules

also showed significant correlation with the first principal component of the PCA on the high-

variance gene set (Fig. 3c, Supplementary Fig. 3h), suggesting that subtype identity explains a

significant amount of variation in gene expression across these neurons. Five of these modules

were enriched for genes identified as significantly differentially expressed between subtype 1

and subtype 2 (Fig. 3d, Supplementary Fig. 3d). No module was correlated with replicate or

other potentially confounding technical parameters (Supplementary Fig. 5f,g), confirming that

the variations we observe are driven by biological differences among neurons rather than

technical variation. Taken together, these data demonstrate that the greatest source of

variation in the transcriptomes of L6CThNs is the difference between subtypes and reveal

several discrete modules of gene expression contributing to this difference.

Projection-dependent and position-dependent gene expression differences contribute to the

transcriptional identity of layer 6 neurons

Although the transcriptional signature of L6CThNs reflects projection target bias, axonal

projection pattern and sublaminar position are confounded among L6CThNs. VPM/POm

neurons are predominantly located in lower layer 6 while VPM-only neurons are biased towards

upper layer 6 (Fig. 1h). Thus the gene expression differences we identified may represent

differences in sublaminar position within layer 6 rather than axonal projection pattern *per se*. To

test this hypothesis, we took advantage of the fact that L6CThNs represent only approximately

half of the neurons in layer 6[29, 31]. If a gene's expression reflects axonal projection pattern, its

expression should be restricted to L6CThNs in upper or lower layer 6 only. On the other hand, if

a gene's expression reflects laminar position, we predict that its expression would be restricted

to neurons in either upper or lower layer 6, regardless of their projection pattern.

To select target genes to evaluate using single molecule fluorescence *in situ*

hybridization (smFISH), we performed a PCA on the mean-centered expression estimates of the

high-variance genes across all 346 L6CThNs and used the rotations from this analysis to project

all expressed genes into this PCA space to rank order candidate genes (Supplementary Fig. 6a-d).

We quantified the expression of target gene mRNAs in individual tdTomato-positive, NeuN-

positive neurons identified as L6CThNs and in tdTomato-negative, NeuN-positive neurons in

slices of barrel cortex from Ntsr1-Cre;tdTomato mice. These data were then fit to a generalized

additive model to test the independent contributions of laminar position and expression of

tdTomato (Fig. 4). We found that *Lamp5*, *Serpini1* and *Gabra5* were selectively expressed in a

subtype of L6CThNs (Fig. 4b,e-f). In contrast, *Pantr1* varied with laminar position within layer 6

in both L6CThNs and non-L6CThNs (Fig. 4c). Our findings reveal that information about position

within layer 6 and long range axonal projection pattern is contained in the gene expression

differences between the two transcriptional subtypes of L6CThNs.

Neural activity significantly contributes to transcriptionally defined subtype identityNeuronal

activity is known to strongly influence gene expression[4-6]. We therefore hypothesized that

neuronal activity state influences the transcriptional profiles of the L6CThNs and may contribute

to their transcriptional identity. To test whether any modules reflect activity state, we assessed

genes assigned to each module for enrichment of a curated gene set representing the ensemble

of genes regulated after induction of neural activity compiled from recently published studies[43-

46] (Fig. 3e, Supplementary Fig. 5, Supplementary Table 4). Four of the seven modules correlated

with transcriptional subtype (Black, Green, Purple, and Midnight Blue) were significantly

enriched for genes induced by neuronal activity[43-46] (Fig. 3e, Supplementary Fig. 5, $p < 0.01$,

hypergeometric test). Each of these four modules was also significantly correlated with the first

and second principal components of the PCA on the high-variance gene set, suggesting that

differences in neuronal activity state explain a significant amount of transcriptional variation

across L6CThNs as well. Taken together, our results show that the long range axonal projection

pattern, laminar position within layer 6, and the activity state of each neuron are all reflected in

the transcriptional profiles of individual L6CThNs and are principal contributors to the identity of

the two L6CThN subtypes.

253   Among the four activity-associated modules (Black, Green, Purple, and Midnight Blue),

Black was specifically correlated with subtype 1, and Purple and Midnight Blue with subtype 2,

suggesting subtype-specific engagement of activity-induced genes in the steady state. The

Green module was correlated with PC1-PC3, but demonstrated no significant bias for either cell

type. To assess whether these signatures of neuronal activity represent a fundamental aspect of

subtype identity, we re-evaluated our classification workflow after regressing out the

eigenvalues for each activity-associated module and observed a reduction in the separation of

the two L6CThN subtypes in each case, suggesting that steady state differences in neuronal

activity genes are a defining characteristic of these two L6CThN subtypes (Supplementary Fig.

6e).

To further test the influence of neuronal activity on transcriptional identity, we

evaluated the enrichment of activity genes along the first two principal components of a gene-

centric PCA of the mean-centered expression estimates of high variance genes across all 346

L6CThNs. All expressed genes were projected in this PCA space and ranked using their weights

along PC1 and PC2 (Supplementary Fig. 6a-d). Both PC1 and PC2 were significantly enriched for

genes drawn from our curated list of activity-induced genes[43-46] (p < 0.01; Kolmogorov-Smirnov

test, Pre-ranked GSEA). The expression levels of the most heavily weighted PC1 genes varied

predominantly between subtypes (Fig. 5b-c), while the expression levels of heavily weighted PC2

genes additionally varied within each subtype in a pattern similar to classical activity-induced

genes like *Fos* and *Bdnf* (Fig. 5d-g), suggesting that activity contributes both to subtype identity

and to transcriptional heterogeneity within each subtype. The expression patterns of genes such

as *Cdh13*, *Igfbp6*, and *Sla* that strongly contributed to both PC1 and PC2 provide further

evidence that axonal projection pattern, laminar location, and activity state are not fully

orthogonal axes contributing to cellular identity (orange asterisks, Fig. 5c, Supplementary Fig.

6c,d). These data indicate that neuronal activity accounts for significant variation in the gene

expression of L6CThNs, contributing to transcriptional heterogeneity both within and between

L6CThN subtypes.

Modulation of neuronal activity influences the transcriptional state and identity of L6CThNs

Our data demonstrate that the gene expression profiles of L6CThNs reflect an integration of

multiple basis vectors corresponding to discrete, but potentially dependent sources of variation

dominated by axonal projection pattern, sublaminar position within layer 6 and neuronal

activity. Alterations in the molecular cascades engaged by different patterns of neural activity

therefore have the potential to modulate the transcriptional identity of L6CThNs. To test this

hypothesis, we unilaterally removed whiskers in a chessboard pattern, a pattern of sensory

deprivation shown to engage plasticity mechanisms in the barrel cortex[47, 48], from Ntsr1-

Cre;tdTomato mice which were bilaterally injected with a retrograde tracer in POm to label

VPM-only and VPM/POm L6CThNs as in our baseline experiments (Fig. 6a). At one and seven

days following this sensory manipulation, we collected and sequenced single L6CThNs from the

barrel cortex both contralateral and ipsilateral to the manipulation. After preprocessing and

quality control, we obtained high-quality transcriptional profiles for 133 of 151 sequenced

L6CThNs from 2 replicates one day following whisker removal (Day 1) and for 550 of 563

sequenced L6CThNs from 2 replicates seven days following whisker removal (Day 7). When

combined with L6CThNs collected under baseline conditions (Day 0), we generated a data set

comprised of 1023 sequenced L6CThNs collected from two replicates at each of three time

points (Supplementary Fig. 7a-d, Supplementary File 3).

To assign each newly acquired L6CThN to its transcriptional subtype, gene expression

profiles were first pre-processed and normalized as described for the baseline data set (Day 0).

We then performed a PCA using the 286 genes we identified as differentially expressed between

transcriptional L6CThN subtypes under baseline conditions and a tSNE analysis on the 1023

L6CThNs (Fig. 6b). Neurons were clustered using a Gaussian mixture model[49, 50] (MClust), and

cluster assignments were used to designate a subtype identity to each newly profiled neuron.

This approach largely recapitulated our original subtype assignments, as only ten neurons

classified under baseline conditions were assigned to a different subtype (Fig. 6c). The significant

axonal projection bias of the two transcriptional subtypes was maintained as subtype 1 neurons

were predominantly labeled VPM/POm (70.7%, 408 of 585 cells, $p < 2.97 \times 10^{-71}$, hypergeometric

test) and subtype 2 VPM-only (84.2%, 369 of 438 cells, $p < 2.97 \times 10^{-71}$, hypergeometric test),

highlighting the reproducibility of our method and the robustness of our subtype assignment

(Fig. 6d).

We identified 1134 genes with significant differential expression across the three time

points sampled, independent of stable, baseline differences in expression between the two

transcriptional subtypes (Fig. 6e, q ≤ 0.001; Monocle2 test). A k-means clustering analysis of

mean expression profiles identified 16 clusters of genes with different temporal expression

following deprivation, but failed to identify any clusters with significant subtype-specific effects

over time (Fig. 6e). However, the transcriptional changes induced in response to altered sensory

input are unlikely to be synchronous across all neurons collected at each time point. To describe

the cellular responses to altered sensory input without the confounding effects of neurons in

diverse states intermixed at each time point, we established a pseudotemporal ordering for the

1023 L6CThNs derived from the 1134 genes with significant differential expression across time

points (Fig. 7a). Briefly, using the Monocle2 DDRTree algorithm, cells were arranged in an

embedded graph representation in a reduced dimensional space. In this manner, cells with

similar expression profiles across the 1134 target genes were positioned next to each other, and

a traversal through the graph revealed the sequence of transcriptional changes that reflect

progression through a given biological process. Through this projection, we captured a smooth

representation of transcriptional modulation to altered sensory input. As expected, the

distribution of L6CThNs along pseudotime generally followed the temporal order of collection

following whisker removal (Fig. 7a,b). Nonetheless, neurons from each time point were found

throughout pseudotime, confirming that the transcriptional response to sensory manipulation is

asynchronous across the population of L6CThNs. Interestingly, we observed no bias in the

distribution of L6CThNs across pseudotime when the neurons were grouped by hemisphere ipsi-

or contralateral to whisker removal, suggesting that longer term transcriptional responses in

L6CThNs from both hemispheres are similar (Supplementary Fig. 7e).

We identified 1507 genes that were differentially expressed across pseudotime at a

much higher stringency than our aggregate analysis independent of transcriptional subtype

designation (q ≤ 0.0000001; Monocle2), indicating that a significantly greater fraction of the

transcriptome was identified as modulated by sensory manipulation using the increased

resolution of pseudotemporal ordering. To identify cellular processes modulated along

pseudotime, we clustered the normalized response curves of the differentially expressed genes

(Fig. 7c, Supplementary Table 5) and queried each cluster for enrichment of annotated gene sets

from public databases and our curated list of activity-induced genes. The cluster with the

earliest changes in expression (Cluster 2) corresponded to significant downregulation of the

activity-associated genes (p < 5.65 x 10^{-7}, hypergeometric test) that were enriched in the four

activity-associated modules contributing to cellular identity. In contrast, the cluster representing

the late response (Cluster 3) corresponded to upregulation of genes associated with chromatin

remodeling and reorganization as well as upregulation of lncRNAs, suggesting a slower

epigenetic response to sensory manipulation. Interestingly, genes associated with long-term

potentiation (LTP) from Cluster 2 (*Calm1*, *Calm2*, *Gria1*, *Gria2*, *Plcb1*, *Plcb4*, *Ppp1ca*, and *Ppp3r1*)

were downregulated early in the response while LTP-associated genes in Cluster 3 (*Crebbp*,

Adcy1, *Grin1*, *Prkcb*, *Ppp3ca*, and *Ppp3cb*) were upregulated towards the end of pseudotime,

suggesting that non-overlapping subsets of genes in this single gene set are regulated at distinct

phases of the L6CThN response to sensory manipulation. We also identified 75 transcription

factors (TFs) with significant differential regulation after sensory manipulation including activity-

associated TFs[5, 6] such as *Arc*, *Fos*, *Ier2*, *Junb*, *Mef2c*, and *Nr4a1,* which were expressed early and

downregulated over the course of pseudotime. Several TFs involved in neural development[3]

including *Neurod6*, *Fezf2*, *Mef2c*, and *Foxp2* were transiently expressed during the response,

suggesting a regulatory relationship between activity-dependent plasticity and neural

development.

Activity induced changes in gene expression enhance the distinction between subtypes

Our data demonstrate that a transcriptional signature of activity significantly contributes to

L6CThN subtype identity under baseline conditions, and that altered sensory inputs result in

dramatic transcriptional responses in L6CThNs. These results suggest that the response to

sensory manipulation may alter the transcriptional relationship among neurons of the same

subtype as well as the distinction between the two transcriptionally defined L6CThN subtypes.

To test these hypotheses, we assessed the distribution of pairwise Euclidean distances of the

variance stabilized gene expression estimates across all expressed genes for L6CThNs within

each subtype for each day following sensory manipulation. In both subtypes, we observed that

the response to altered sensory input resulted in an increased mean distance among cells across

days (Fig. 8a; $p < 2.2 \times 10^{-16}$, Welch's two-sample t-test). We also identified a significant increase

in the variance of the intra-subtype distances across days ($p < 3.59 \times 10^{-5}$, F-test) for all adjacent

time points except for subtype 2 Day 1 versus Day 7, indicating that the cell-to-cell variation

within both L6CThN subtypes increases in response to altered sensory input. Second, we

assessed the distribution of the inter-subtype pairwise Euclidean distances between subtype 1

and subtype 2. We found that the mean distance significantly increased from Day 0 to Day 1 as

well as from Day 1 to Day 7 (Fig. 8b; $p < 2.2 \times 10^{-16}$, Welch's two-sample t-test) as did the

variance of inter-subtype distances ($p < 4.67 \times 10^{-8}$, F-test). These results demonstrate that

modulation of neuronal activity increases both the transcriptional heterogeneity within each

L6CThN subtype and the relative transcriptional differences between L6CThN subtypes,

confirming a dependent relationship between activity level and transcriptional identity.

Subtype-biased responses contribute to transcriptional heterogeneity and subtype identity

These results raise the possibility that subtype-specific responses to sensory manipulation drive

the increase in transcriptional heterogeneity and enhance the distinction between L6CThN

subtypes. We found that subtype 1 and subtype 2 were differentially distributed along

pseudotime at each time point assessed following sensory manipulation (Fig. 8c). Furthermore,

pseudotemporal ordering identified a single major branch point in the transcriptional responses,

with each of the two subsequent branches exhibiting a significant bias for a specific L6CThN

subtype (Branch A and Branch B; Fig. 8d). Consistent with the proportion of the two subtypes in

the baseline data, neither subtype 1 nor subtype 2 was significantly enriched in the root state

(grey arrow; Subtype 1: p < 0.0715; Subtype 2: p < 0.91, hypergeometric test), indicating that

both transcriptional subtypes share a similar early response to sensory manipulation. In

contrast, Branch A and Branch B exhibited significant subtype bias: Branch A (red arrow) was

biased for subtype 1 (VPM/POM; 80.4%, 148/184 neurons; $p < 7.03 \times 10^{-14}$, hypergeometric test)

and Branch B (blue arrow) for subtype 2 (VPM-only; 58.4%, 202/346 neurons; $p < 2.04 \times 10^{-13}$,

hypergeometric test). Similar subtype-specific biases were seen at a subsequent branch point

(Supplementary Fig. 8). These results indicate that although either cell type may engage the

discrete processes represented by each branch in response to sensory manipulation, the cellular

decisions to engage a particular response are biased with respect to L6CThN subtypes.

Using the Monocle2 branch expression analysis modeling (BEAM) test, we identified

1392 genes with significant branch-dependent differential expression (Supplementary Table 6; q

≤ 0.0001; Monocle2 BEAM test); 926 of these overlapped with the 1507 genes with pseudotime-

dependent expression suggesting that discrete cellular responses independently contribute to

the aggregate transcriptional response to altered sensory manipulation. The branch-dependent

genes were organized into seven distinct clusters using hierarchical clustering of the Monocle2

branched model fits (Fig. 8e). Hypergeometric testing demonstrated that neurons that progress

along Branch A (VPM/POM enriched) were enriched for genes associated with engagement of

the proteasome complex, a process involved in synaptic remodeling[51], while Branch B cells

(VPM-only enriched) were significantly enriched for genes related to the organization of the

postsynaptic density as well as to the regulation of LTP. We identified 79 TFs expressed in a

branch-dependent manner, including 31 TFs not identified in our initial analysis of the aggregate

response (Fig. 8f), suggesting that much of the aggregate L6CThN response induced by altered

sensory inputs is confounded across these two alternative cellular responses. The remaining 27

pseudotime-dependent TFs may regulate a uniform response independent of these two

subtype-biased responses. Additional genes regulated in a branch-specific manner included

thirty-two presynaptic genes (Fig. 8g). Furthermore, neurons committing to a Branch A response

demonstrated enhanced expression of GABA receptor subunits (*Gabra5*, *Gabrb1* and *Gabrb2*)

while neurons along Branch B induced ionotropic glutamate receptors (Fig. 8h; *Grin1*, *Grin2b*,

Grik5, and *Grin2a*). Taken together, these subtype-specific biases in transcriptional responses

induced by sensory manipulation likely underlie the overall effect we observed on L6CThN

identity, enhancing the distinctions between subtype 1 and subtype 2, and significantly

increasing both the inter- and intra-subtype heterogeneity of L6CThNs.

DISCUSSION

Using scRNA-seq, we identified two transcriptionally defined subtypes of L6CThNs, each

exhibiting significant bias for particular long-range axonal projection targets. Analysis of gene

expression differences among the profiled neurons determined that variation in axonal

projection pattern, laminar position within the cortex, and neuronal activity state all significantly

contribute to transcriptional identity, and that these properties are at least partially dependent.

Manipulating the activity states of L6CThNs by altering tactile input further showed that each

subtype was biased for particular cellular responses to sensory manipulation, revealing

functional differences among these neurons. These subtype-biased transcriptional responses

not only increased cell-to-cell transcriptional heterogeneity within each subtype, but also

enhanced the transcriptional differences between the two subtypes. Together, these data

identify the most significant influences on the transcriptional identity of individual cortical

projection neurons, and suggest that discrete cellular properties and responses affect the

population-level variation and identity of neuronal subtypes.

Although it is possible that projection target and transcriptional identity are decoupled

for a minority of neurons, the incomplete segregation of retrograde label across subtypes

observed here likely represents mislabeling of a subset of L6CThNs, as expected due to the close

proximity of VPM and POm and the difficulty of retrogradely labeling all neurons projecting to

adjacent targets. Interestingly, the subtypes identified here share some transcriptional

similarities with two recently defined subtypes in primary visual cortex[12] (Supplementary Fig. 3),

suggesting a conservation of corticothalamic neuron transcriptional identity and axonal

projection target bias across functional regions of the cortex[20]. Future studies will be required to

determine whether VPM-only and LGN-only projecting neurons as well as VPM/POM and

LGN/LP neurons are transcriptionally analogous, and to identify sources of cellular variation

across cortical areas. Although the population of L6CThNs we analyzed is comprised of two

subtypes, profiling a greater number of neurons or profiling at greater depth may reveal

additional rare subtypes of L6CThNs or may show that subtle variations in axonal projection

pattern identified in some anatomical studies are not apparent in the expression profiles of

cortical neurons[20, 21].

Pseudotemporal ordering of the states induced in L6CThN transcriptomes by altered

sensory input assessed over seven days revealed that L6CThNs engage at least two molecularly

distinct responses in a subtype-biased, but not subtype-specific, manner. Although the distinct

transcriptional responses were dominated by neurons collected one and seven days following

whisker manipulation, neurons collected in the baseline state were found throughout

pseudotime, suggesting that individual cortical neurons may engage these plasticity responses in

the non-manipulated state. L6CThNs in the initial phase of cellular response to altered sensory

input exhibited similar gene expression changes, indicating that both subtypes engaged a

common initial transcriptional response. These results are consistent with previous studies of

responses induced by neural activity measured over hours across brain regions and between

inhibitory and excitatory neurons which found that common early transcriptional responses lead

to cell-type specific late responses[52, 53].

The differential response to sensory manipulation resulted in an increase in the

transcriptional variation across L6CThNs within each subtype as well as a significant

enhancement of the distinction between the subtypes, demonstrating a non-dissociable

relationship between neuronal identity and neuronal activity. Because single L6CThNs have the

potential to engage either transcriptional response regardless of subtype, our data suggest that

extrinsic factors such as distinct activation patterns, due to differences in the circuits in which

each subtype is embedded, induce neurons from a given subtype preferentially toward a similar

response rather than solely cell-autonomous properties. Interestingly, we found that the

transcriptional responses for neurons ipsilateral and contralateral to the sensory manipulation

were similar, suggesting that shared cellular mechanisms underlie the previously described

functional effects ipsilateral and contralateral to unilateral manipulation at timescales similar to

those assessed here[48, 54, 55]. Furthermore, expression of genes that strongly contributed to

subtype identity such as *Lamp5* was altered in response to sensory manipulation. These data

indicate that factors that change cell state such as plasticity or injury affect our ability to

accurately define discrete, stable transcriptional subtypes.

Our results indicate that the transcriptional profiles of cortical neurons reflect specific

features of these cells, but also that the transcriptional variation across individual neurons is a

principal feature of subtype identity with significant functional consequences for individual

neuronal responses and subtype function. Subtype 2 L6CThNs were more transcriptionally

heterogeneous than subtype 1 neurons under steady-state conditions in part because of

baseline differences in gene expression associated with neural activity. In addition, the intra-cell

type variation across subtype 2 at day 1 and at day 7 was greater than the inter-cell type

variation at the same time points, suggesting that changes in cell-to-cell variation, rather than

subtype specific differences, dominate the transcriptional responses to experience dependent

plasticity. Together, these data indicate that the contribution of neuronal activity to gene

expression differs across distinct neuronal subtypes and that the transcriptional variation due to

differences in neuronal activity state plays a central role in defining the identity of cortical

projection neurons.

 METHODS

Mice
All procedures were approved by the Johns Hopkins Animal Care and Use Committee and
followed the guidelines of the Society for Neuroscience and the National Institutes of Health.
Animals used for RNA-sequencing ranged from postnatal day 23 (P23) to P28; animals used for
immunohistochemistry and *in situ* hybridization ranged from P23 to P32. Both males and
females were used in this study (gender indicated for each replicate below). All animals were
kept on a 12 h light/dark cycle, housed with at least two mice per cage and provided with
unlimited water and food. The following mouse lines were used:
Neurotensin receptor-1 Cre recombinase line[30] (Ntsr1-Cre, GENSAT 220), loxP-STOP-loxP-
tdTomato Cre reporter lines[56] (Ai9 and Ai14, Allen Institute for Brain Science), loxP-STOP-loxP-
eYFP Cre reporter line[56] (Ai3, Allen Institute for Brain Science), and Pantr1-LacZ[39]. Mice were
maintained on mixed backgrounds including C57BL/6 and CD-1.
507
Stereotactic injections
To identify layer 6 corticothalamic neurons (L6CThNs), mice P18 to P23 were anesthetized with
ketamine (50 mg/kg), dexmedetomidine (25 μg/kg) and the inhalation anesthetic, isoflurane.
Animals were placed in a custom-built stereotactic frame and anesthesia was maintained with
isoflurane (1-2.5%). A small craniotomy was performed, and a glass pipette (10–25 μm tip
diameter) was lowered into the thalamus. To identify L6CThNs that project to the ventral
posterior medial nucleus (VPM) of the thalamus, Ntsr1-Cre;tdTomato mice were injected in VPM
(1.1 mm posterior, 1.7 mm lateral , 1.4 mm ventral from bregma) with either Alexa 488 Cholera
toxin B (CTB488, ThermoFisher) or green Retrobeads IX (Lumafluor), and Ntsr1-Cre;YFP mice
were injected with Alexa 555 Cholera toxin B (CTB555, ThermoFisher). To identify L6CThNs that
project to both VPM and to the posterior medial nucleus of the thalamus (POm), Ntsr1-
Cre;tdTomato mice were injected in POm (1.35 mm posterior, 1.23 mm lateral , 3.3 mm ventral
from bregma) with green Retrobeads IX, and Ntsr1-Cre;YFP mice were injected with either
CTB555 or red Retrobeads (Lumafluor). Between 30 and 100 nl of tracer were pressure-injected,
and the pipette was kept in position for 5-10 minutes before removal. Following the injection,
the incision was sutured and buprenorphine (0.05 mg/kg) was administered to all animals
postoperatively. Quantification of the colocalization between retrogradely labeled neurons and
Cre expression in Ntsr1-Cre mice (Fig. 1e,j) was performed on mice injected with either CTB or
Retrobeads. Preliminary experiments indicated that all tracers used generated similar labeling.
However, Retrobeads resulted in increased discrimination of labeled neurons during
Fluorescence Assisted Cell Sorting (FACS) and were therefore used for all FACS experiments.
529
Immunohistochemistry
Brains from animals injected with neuronal tracers were removed and placed in a solution of 4%
paraformaldehyde (PFA) in 0.01 M phosphate buffered saline (PBS) for 3 h. Coronal sections
were cut on a vibratome (50 μm, VT-1000S, Leica). Mice expressing YFP were subjected to
immunohistochemistry (1:1000, chicken anti-GFP, GFP-1020, Aves, RRID:AB_10000240, and
1:300 AlexaFluor 488-conjugated goat anti-chicken, Life Technologies, A11039,
RRID:AB_10563770). Sections were then mounted using Aqua Poly/Mount mounting medium
(Polysciences, Inc) and visualized on a confocal microscope (LSM 510, Zeiss) using 10x (0.3 NA),
25x (0.8 NA) or 40x (1.3 NA) objectives or on an epifluorescence microscope (AxioObserver.Z1,

Zeiss) using a 5x (0.16 NA) objective. Colocalization of tdTomato, YFP and retrograde tracer was
quantified using single-plane confocal images and the Cell Counter plugin in Fiji[57]. For all
anatomical comparisons, upper layer 6 and lower layer 6 were defined as the top 50% and
bottom 50% of the layer defined by Cre-expressing neurons in Ntsr1-Cre mice.
543
Cell isolation and enrichment
For bulk sequencing experiments (bulk RNA-seq), POm was injected with green Retrobeads in
three Ntsr1-cre;tdTomato mice. For single-cell sequencing experiments (scRNA-seq), POm was
injected with red or green Retrobeads in Ntsr1-Cre;YFP mice or in Ntsr1-Cre;tdTomato mice,
respectively. Mice were sacrificed 5-7 days after surgery (Bulk RNA-seq: Replicate 1: P23 female;
Replicate 2: P23 female; Replicate 3: P26 male; scRNA-seq baseline: Replicate 1: P27 male,
Replicate 2: P28 female; scRNA-seq Day 1: Replicate 1: P24 female, Replicate 2: P24 female;
scRNA-seq Day 7: Replicate 1: P29 female, Replicate 2: P27 female). Brains were rapidly
removed and 300 µm thick somatosensory thalamocortical[58] or coronal slices were sectioned
(VT-1200s, Leica) in ice-cold sucrose solution containing the following (in mM): 76 NaCl, 25
NaHCO$_3$, 25 glucose, 75 sucrose, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 0.5 CaCl$_2$, 7 MgSO$_4$, pH 7.3, 310 mOsm
continuously bubbled with 95% O$_2$/5% CO$_2$. The slices were then placed in a submersion
chamber on an upright microscope in ice-cold sucrose solution continuously bubbled with
95% O$_2$/5% CO$_2$. The barrel cortex was identified using differential interference contrast (DIC)
and epifluorescence microscopy (Axioskop2 FsPLus, Zeiss) using 4x (0.1 NA) and 40x (water
immersion, 0.75 NA) objectives. The location of the retrograde tracer in the lower half of layer 6
was visually confirmed, and then regions of cortex below barrels with appropriate labeling were
microdissected and immediately dropped into 2.6 mL equilibrated Papain DNase-I solution. For
all experiments, at least three consecutive slices confirmed as clearly labeled were
microdissected.
The dissociation protocol was adapted from the trehalose-enhanced neuronal isolation
protocol described in Saxena et al, 2012[59] which is based on the Worthington Papain
Dissociation System (Worthington Biochemical Corporation). Briefly, tissue pieces were dropped
into a 5 ml tube containing 2.6 mL of a trehalose-Papain-DNase1 solution which included RNase
inhibitor, incubated 30-45 min at 37°C with gentle trituration at the half way point using a p1000
pipetter. The reaction was stopped by adding a protease inhibitor solution (for solution
compositions, see Saxena et al, 2012[59]. The tissue was then further dissociated and washed by
sequential pipetting and low-speed centrifugations. The final pellet was resuspended in 200 µL
of media (DMEM, 5% trehalose (w/v), 25 µM AP-V, 0.4 mM kynurenic acid, 6 µL of 40 U/µl
RNase inhibitor) at room temperature and introduced into a FACS machine (Beckman Coulter
MoFlo Cell Sorter). Neurons were sorted based on their fluorescence (either tdTomato-positive
(red) and Retrobead-positive (green) versus tdTomato-positive only or YFP-positive (green) and
Retrobead-positive (red) versus YFP-positive only) directly into lysis buffer either in eppendorf
tubes for bulk sequencing (350 µL lysis buffer, Qiagen RNeasy Micro Kit; Replicate 1: tdTomato-
only: 1943 cells, tdTomato and tracer: 2287 cells; Replicate 2: tdTomato-only: 386, tdTomato
and tracer: 782; Replicate 3: tdTomato-only: 604, tdTomato and tracer: 934) or into individual
wells of 96-well plates for single-cell sequencing (2 µL Smart-Seq2 lysis buffer + RNAase
inhibitor, 1 µL oligo-dT primer, and 1 µL dNTPs according to Picelli et al, 2014[60]. Upon
completion of a sort, the plates were briefly spun in a tabletop microcentrifuge and immediately
placed on dry ice. Single-cell lysates were subsequently kept at -80°C until cDNA conversion.
584

Library preparation and amplification

585
For bulk samples, RNA was extracted with a Qiagen RNeasy Micro Kit. RNA quality was assessed
with the Bioanalyzer Pico RNA kit and samples had RIN scores of 8.2-8.8. Library preparation and
amplification were performed according to the Smart-Seq2 protocol[60] with some modifications.
Principally, all primer and adapter sequences were synthesized with a 5'-biotin to minimize
primer dimerization. Briefly, 200 pg of RNA per sample were used as input for template
switching cDNA synthesis. Full-length cDNA was amplified by KAPA HiFi mediated PCR for 20
cycles using a 5'-biotinylated ISPCR primer. Then, 500 pg of Ampure XP bead cleaned (1:1) cDNA
was used as input for a standard Nextera XT tagmentation reaction, and amplification of
adapter-ligated fragments was carried out for 12 cycles during which individual index sequences
were added to each distinct sample. Library concentration was assessed with Qubit and library
fragment size distribution was assessed on the Agilent Bioanalyzer. Pooled, indexed bulk RNA-
seq samples were initially sequenced on the Illumina MiSeq 500 platform, and subsequently the
same pool was sequenced on one lane of the Illumina HiSeq 2500 platform to produce 50 bp
paired-end reads. Libraries were sequenced to an average depth of 4.2×10^7 fragments each.
Reads from both runs were aggregated by index prior to mapping.
Library preparation and amplification of single-cell samples was performed using a
modified version of the Smart-Seq2 protocol. Briefly, 96-well plates of single-cell lysates were
thawed to 4°C, heated to 72°C for 3 minutes, and then immediately placed on ice. Template
switching first-strand cDNA synthesis was performed as described above using a 5'-biotinylated
TSO oligo. cDNAs were amplified using 18 cycles of KAPA HiFi PCR and 5'-biotinylated ISPCR
primer. Amplified cDNA was cleaned with a 1:1 ratio of Ampure XP beads and 300 pg was used
for a one-half standard-sized Nextera XT tagmentation reaction. Tagmented fragments were
amplified for 12 cycles and dual indexes were added to each well to uniquely label each library.
Concentrations were assessed with Quant-iT PicoGreen dsDNA Reagent (Invitrogen) and
samples were diluted to ~2 nM and pooled. Pooled libraries were sequenced on the Illumina
HiSeq 2500 platform to a target mean depth of ~8.0×10^5 50 bp paired end fragments per cell[61]
at the Hopkins Genetics Research Core Facility to generate 50 bp paired-end reads.
613
***In situ* hybridization and immunohistochemistry**
Immunohistochemistry (IHC) for tdTomato and *in situ* hybridization (ISH) for *linc-Tmem20* were
combined to show co-localization of the lncRNA transcript and tdTomato protein in L6CThNs.
Ntsr1-Cre;tdTomato mice (P23 to P30) were transcardially perfused with 0.1 M PBS followed by
a 4% PFA solution, and brains were post-fixed for 3 h at room temperature in 4% PFA. Then, 30
µm sections were cut on a vibratome (VT-1000S, Leica) and first processed for IHC (1:1000,
Rabbit anti-DsRed polyclonal antibody, Living Colors, Clontech Laboratories, RRID: AB_10013483
and 1:300 AlexaFluor 568-conjugated goat anti-rabbit, Life Technologies, A11011, RRID:
AB_143157). Next, the sections were processed for ISH following the protocol described in
Blackshaw, 2013[62] without proteinase K treatment. Sections were incubated overnight at 65°C
with 5 ng/µL of an anti-sense *linc-Tmem20* DIG labeled probe (DIG RNA labeling Kit (SP6/T7),
Roche). Detection of the hybridized DIG probes was performed using an alkaline phosphatase
(AP) conjugated antibody (1:1000, Anti-Digoxigenin [21H8] Alkaline Phosphatase, Abcam, Cat.
No.: ab119345, RRID:AB_10901703) followed by application of the AP substrate BCIP/NBT
(SigmaFast BCIP/NBT, Sigma-Aldrich). The enzymatic reaction was carried out until the desired
signal to noise ratio was observed. Images were then taken on an AxioObserver.Z1 (Zeiss) using

a combination of epifluorescence and brightfield imaging (10x objective, NA: 0.3). Brightness
and contrast were adjusted using Adobe Photoshop (Adobe Systems Incorporated).
For validation of *Pantr1* expression, transgenic mice expressing LacZ driven by the
endogenous *Pantr1* promoter locus[39, 63] were injected with CTB488 in VPM and CTB555 in POm.
IHC was performed on 50 μm-thick sections (rabbit anti-β galactosidase antibody 1:5000; a kind
gift from Dr. Joshua R. Sanes, Harvard University, Cambridge, MA and 1:300 AlexaFluor 647-
conjugated donkey anti-rabbit, Life Technologies, A31573, RRID: AB_2536183). Imaging was
performed on a confocal microscope (LSM 510, Zeiss) using 10x (0.3 NA), 25x (0.8 NA) or 40x
(1.3 NA) objectives. LacZ-positive and retrogradely labeled neurons were manually counted on
single-plane confocal images using the Cell Counter plugin in Fiji[57]. Brightness and contrast were
adjusted using Adobe Photoshop (Adobe Systems Incorporated).
For detection of single mRNA molecules, we used the commercially available system
RNAscope (ACDbio). Ntsr1-Cre;tdTomato mice (P23 to P30) were transcardially perfused with
0.1 M PBS followed by a 4% PFA solution, and brains were post-fixed for 2 h at room
temperature in 4% PFA. Fixed brains were then washed in PBS and allowed to equilibrate in a
30% sucrose-PBS solution for 36 h before being embedded and frozen in OCT compound
(Scigen, Tissue-Plus #4583). Frozen brains were kept at -80°C for up to 2 months. 20 μm sections
were collected on a cryostat (Leica CM3050) and processed according to instructions provided
by RNAscope. The following probes were used: tdTomato-C2 (317041-C2), tdTomato-C3 (
317041-C3), Mm-Rbfox3 (313311), Rbfox3-C2 (313311-C2), Mm-Lamp5 (451071), Mm-Pantr1
(483711), Mm-Serpini1 (501441), Gabra5-C3 (319481-C3). Imaging was performed on a confocal
microscope (LSM 510, Zeiss) using a 63x objective (NA 1.4). Sections from three animals were
used for *Pantr1*, *Lamp5* and *Gabra5* and from four animals for *Serpini1*. For quantification of the
number of mRNAs per neuron obtained using RNAscope, single plane images were first
processed with the open-source software CellProfiler (Broad Institute) to automatically identify
individual cells based on DAPI signal and to generate outline files. These outlines were then used
as input, in combination with the corresponding confocal images, into FISH-quant[64] (Matlab) for
quantification of transcript number per cell. Cells with no detectable NeuN (*Rbfox3*) transcript
were discounted as non-neuronal cells. A threshold of 20 *tdTomato* puncta was selected as the
cutoff for distinguishing tdTomato-positive from tdTomato-negative neurons. This resulted in
164 tdTomato-positive and 188 tdTomato-negative neurons for *Serpini1*, 118 tdTomato-positive
and 150 tdTomato-negative neurons for *Lamp5*, 111 tdTomato-positive and 198 tdTomato-
negative neurons for *Pantr1*, and 135 tdTomato-positive and 160 tdTomato-negative
neurons for *Gabra5*. Individual images were stitched to reconstitute an image spanning layer 6
vertically, and the distance of each neuron from the bottom of the layer was measured manually
using Fiji[57] and normalized to the height of layer 6. Statistical testing (Likelihood ratio test) and
curve fitting (Loess) was performed in R/Bioconductor.
667
Sensory manipulation
Prior to sensory manipulation, animals were bilaterally injected in POm with green Retrobeads
IX (Lumafluor) as described above. To induce experience dependent plasticity, mice (P18 to P23)
were anesthetized with the inhalation anesthetic, isoflurane. Once an adequate level of
anesthesia was achieved, the animals were placed under a surgical microscope and, using
tweezers, individual vibrissae were gently pulled by applying slow, steady tension to the base of
the whisker[65]. To produce a chessboard pattern, vibrissae alpha, gamma, A2, A4, B1, B3, C2, C4,
C6, D1, D3, D5, D7, E1, E3, E5 and E7 were pulled unilaterally while the rest remained

untouched. Following the procedure, the animals were returned to their home cage and housed
with at least one other animal until tissue collection. We checked for regrowing vibrissae every
two days and removed them if visible.
679
Read pre-processing
For both bulk and single-cell libraries, paired-end reads were aligned to the mouse reference
genome (mm10) using Hisat2[66] with the default parameters except: -p 8. Aligned reads from
individual samples were quantified against a reference transcriptome[61] (GENCODE vM8)
supplemented with additional lncRNA genes described previously[8] (Supplementary File S1) with
cuffquant[67] using default parameters with the following exceptions: --no-update-check –p 8. For
bulk RNA-Seq, pre-processed expression estimates were used as input for cuffdiff using default
parameters except: -p 16, and the output of cuffdiff was analyzed using the cummeRbund
R/Bioconductor package[68]. For single-cell experiments, normalized expression estimates across
all samples were obtained using cuffnorm with default parameters. For comparison to results
from primary visual cortex[12], raw reads were downloaded from the short read archive
(GSE71585), processed as described above for single-cell samples, and normalized expression
estimates were obtained from combining both sources of single-cell reads together via
cuffnorm.
694
Single-cell analysis
The normalized FPKM matrix from cuffnorm was used as input for the Monocle2 single-cell RNA-
seq framework[69] in R/Bioconductor[70]. Relative FPKM values for each cell were converted to
estimates of absolute mRNA counts per cell (RPC) using the Monocle2 census utility. A total of
346 individual cells under baseline conditions passed quality filters including: a) total mRNAs
between 2000 and 200,000 per cell, and b) >10,000 fragments mapped (total_mass). 12,537
genes were identified as expressed in at least 15 cells across all conditions.
To identify high-variance genes, a generalized additive model[71] (MGCV R package) was
fit to the $\log_2$ mean RPC expression versus a cubic spline fit to the $\log_2$ coefficient of
variation(BCV) for each replicate dissociation independently (Supplementary Fig. 2a and Fig. 7d).
The intersection of genes with residuals to this fit greater than 1.0 from each replicate were
chosen as 'high-variance' genes and the $\log_2$ expression estimates (with a pseudocount of 1) of
these selected genes were used as input for PCA analysis and t-stochastic nearest neighbor
embedding (tSNE) clustering of individual cells[72].
To cluster individual cells, we employed a spectral tSNE approach[73], combining non-
linear dimensional reduction and density clustering. Briefly, PCA analysis was conducted on
high-variance genes and then a subset of components (n = 3, selected by permutation parallel
analysis, Supplementary Fig. 2b) were used as input for the tSNE visualization algorithm[72]
(perplexity = 30). K-means clustering was used to partition cells using the two reduced tSNE
dimensions into two discrete subtypes. The choice of k was made using 1000 bootstrapped tSNE
estimates and evaluating k = 2 through k = 8 for each round. The average silhouette coefficient
for each tSNE*k was determined and the mean and standard deviation of these estimates was
estimated for each k (Supplementary Fig. 2c). k = 2 was chosen as the k value with the highest
average silhouette coefficient and the lowest value of k for which there was no significant
improvement in mean silhouette score.
After cluster assignment, differential expression testing was performed across all
expressed genes using the Monocle2 VGAM model comparison test[69]. To test for differentially

expressed genes between transcriptionally-defined cell types, the following full model was fit to
each expressed gene: ~replicate+num_genes_expressed+celltype, and compared to a reduced
model in which celltype was removed. The number of genes expressed in each cell was included
as an explanatory variable as a proxy for the sensitivity of scRNA-seq in each individual cell. To
test for differentially expressed genes between anatomically labeled cells, the following full
model was fit to each expressed gene: ~replicate+num_genes_expressed+label, and compared
to a reduced model in which label was removed. In both cases, genes were selected as
significantly differentially expressed with a 0.1% false discovery rate (Monocle2 test; Benjamini-
Hochberg corrected). DE gene lists were tested for gene set enrichment against Gene
Ontology and Reactome genesets[74, 75] using the clusterProfiler Bioconductor package[76]. The
heatmap of differentially expressed genes was composed using the pheatmap R package[77] and
dendrograms were generated from the hierarchical clustering of the Euclidean distances
between points.
To identify gene sets contributing to transcriptional identity in our single-cell dataset,
gene-centric PCA was performed on a mean-centered matrix of variance-stabilized expression
estimates for high-variance genes across all cells. The resulting rotations were used to project
all expressed genes into the same PCA space to identify their weights. These weights were used
to rank-order all expressed genes and this ordering was used as input for a pre-ranked GSEA
analysis[78]. Gene sets for the GSEA analysis were derived from the Monocle2 differential gene
tests described above or from our curated list of neuronal activity genes[43-46] using their adjusted
p-value cutoff of $p < 0.01$ (Supplementary Table 4).
743
WGCNA
To identify modules of correlated gene expression we used the Weighted Gene Correlation
Network Analysis package WGCNA[42]. Normalized expression data were first batch-corrected
with respect to replicates using limma, and the resulting matrix of 12,537 detectably expressed
genes and 346 cells was used as input for WGCNA. Cells were hierarchically clustered using the
Jensen-Shannon distance with average linkage. A soft threshold power of 11 was used to create
a signed Topological overlap matrix directly from the normalized and batch-corrected
expression estimates. Modules were then learned with a minimum module size of 30. Modules
were merged below a distance threshold of 0.5. Module eigenvalues were estimated and
correlated with cellular traits using the Pearson product moment correlation test. Module gene
membership was determined in a similar manner in which individual gene expression estimates
were correlated with module eigenvalues and tested for significance. To test the effect of each
module on the segregation of L6CThN cell types, each module eigenvalue was separately
regressed out of the expression matrix using limma, and subsequent values were used as input
for a tSNE using identical parameters to the original assay.
759
Pseudotime and BEAM analyses
Pseudotemporal ordering was performed using the prescribed Monocle2 workflow. Briefly, all
1023 L6CThNs passing quality controls were used as input. To reconstruct a trajectory that
reflected cellular progression in response to altered sensory input, we first performed a
differential test to identify genes whose expression changed as a function of collection day,
independent of baseline differences between celltype (fullModelFormulaStr =
"~num_genes_expressed+Total_mRNAs+sex+celltype*Day, reducedModelFormulaStr = "
~num_genes_expressed+Total_mRNAs+sex+celltype", $q \leq 0.01$). These 1134 genes were used as

a filtering set for the DDRTree dimensionality reduction. Cells were tested for differential
expression along pseudotime using the following model comparisons with q ≤ 0.0000001:
fullModelFormulaStr = "~num_genes_expressed+celltype+sm.ns(Pseudotime,df=3)",
reducedModelFormulaStr = "~num_genes_expressed+celltype". Branch dependent expression
was determined using the Monocle2 BEAM test (fullModelFormulaStr =
"~num_genes_expressed+celltype+sm.ns(Pseudotime, df =
3)*Branch",reducedModelFormulaStr = "~num_genes_expressed+celltype+sm.ns(Pseudotime,
df = 3)", q ≤ 0.0001). All significant gene lists were tested for gene set enrichment using the
hypergeometric test.
777
778 **Data and software availability**
779 All primary data are archived on the SRA and Gene Expression Omnibus and are available for
780 direct download on our companion website
781 (http://rstudio.gofflab.org:8383/apps/L6CthPN_dash/). Source code and software tools are
782 available upon request.
783

784 **Author Contributions**: MC, SPB and LAG conceived the experiments. MC, JDR, and GHC
785 performed all experiments. MC, SPB and LAG analyzed the data and wrote the paper with input
786 from all authors.
787
788 **Conflict of Interest**: The authors declare no competing financial interests.
789
Acknowledgments: We thank Hao Zhang and the Johns Hopkins Bloomberg School of Public
Health Flow Cytometry and Cell Sorting Core Facility. Sequencing service was provided by the
Johns Hopkins Genetic Resources Core Facility. We thank Dr. Joshua R. Sanes, Harvard
University, for the anti-LacZ antibody and Dr. Jesse Gray, Harvard Medical School, for the
curated list of neuronal activity-induced genes. This work was supported by a Klingenstein-
Simons Fellowship (SPB), a Johns Hopkins Science of Learning Grant (SPB, LAG), National Science
Foundation Grant 1656592 (SPB, LAG), NIH training grant T32 GM07814 (JDR), a Boerhinger-
Ingelheim Fonds Fellowship (MC), and the NINDS (NS050274).

803 **REFERENCES**

 FIGURE LEGENDS

Figure 1: Layer 6 corticothalamic neuron subclasses distinguished by their axonal projections

have distinct gene expression profiles. (**a**, **f**) The labeling schemes for two classes of layer 6

corticothalamic neurons (L6CThNs; VPM: Ventral posterior medial nucleus; POm: Posterior

medial nucleus). (**b**) Injection of a retrograde tracer (red, Alexa 555-Cholera toxin B) into VPM in

an Ntsr1-Cre;YFP mouse. Low (**c**) and high (**d**) magnification images of layer 6 (L6) of barrel

cortex (BC) showing colocalization of the retrograde tracer (red) and YFP (green). (**e**)

Quantification of the colocalization (mean±SE; n=4 mice). (**g**) Injection of a retrograde tracer

(red) in POm of an Ntsr1-Cre;YFP mouse. Low (**h**) and high (**i**) magnification images of BC

showing the colocalization of the tracer to CThNs in lower L6. (**j**) Quantification of the

colocalization (mean±SE; n=3 mice). (**k**) Matrix showing the 69 genes differentially expressed

between pools of VPM/POm and VPM-only L6CThNs from three replicates. Scale bars, (**b,g**) 500

μm; (**c,h**) 50 μm; and (**d,i**) 10 μm.

Figure 2: Unbiased clustering of single transcriptomes of layer 6 corticothalamic neurons

defines two subtypes with strong axonal projection bias. (**a**) tSNE plot showing two subtypes of

layer 6 corticothalamic neurons (L6CThNs) identified through unbiased clustering of single

L6CThN transcriptomes. (**b**) Same tSNE plot as in (**a**), with each L6CThN color-coded by its axonal

projection label. (**c**) Fraction of VPM-only (green) and VPM/POm (red) L6CThNs in each

transcriptionally defined subtype for each replicate. (**d**) Matrix showing the hierarchical

clustering of the 346 individual L6CThNs (x-axis) and the 286 genes differentially expressed (DE)

between the two subtypes (y-axis, 0.1 % FDR). (**e**) tSNE plots showing the normalized expression

levels of two differentially expressed genes enriched in subtype 1 (*Fxyd6*, left) or subtype 2

(*Lamp5*, right). (**f**) tSNE plots showing the normalized expression levels of two differentially

expressed long-noncoding RNAs enriched in subtype 1 (*linc-Tmem20*, left) or subtype 2 (*Pantr1*,

right). (**g**) Low magnification image of *linc-Tmem20* (red) in barrel cortex of an Ntsr1-

Cre;tdTomato (green) mouse using a combination of *in situ* hybridization (*linc-Tmem20*) and

immunohistochemistry (tdTomato). Insets show higher expression of *linc-Tmem20* in L6CThNs in

lower layer 6 (L6, Inset 2) relative to upper L6 (Inset 1). (**h**) LacZ expression in barrel cortex of a

heterozygous *Pantr1-LacZ* mouse following injections of green retrograde tracer in VPM and red

tracer in POm. Insets show puncta of LacZ in VPM-only L6CThNs (Column 1, green) and not in

VPM/POm L6CThNs (Column 2, red and green L6CThNs). (**i**) Barplot showing the fraction of

VPM-only and VPM/POm L6CThNs expressing LacZ (mean±SE; n=3 mice). (**j**) Boxplot of the

median number of genes detected across all cells for each subtype by replicate pair. (Replicate

1: Median number of genes detected: Subtype 1: 5582 ± 526.3 (sd), Subtype 2: 5080 ± 650.0

(sd), $p < 2.169 \times 10^{-10}$, Mann-Whitney test; Replicate 2: Median number of genes detected:

Subtype 1: 6950 ± 545.4 (sd), Subtype 2: 6569 ± 478.7 (sd), $p < 7.071 \times 10^{-7}$, Mann-Whitney test).

(**k**) Cumulative probability distribution of the pairwise Euclidean distances among cells in

subtype 1 (gold) and subtype 2 (blue) ($p < 2.2 \times 10^{-16}$, Welch's two-sample t-test). The black line

represents the pairwise distances among a random sample of 100 cells drawn from the 346

cells. 95% confidence interval is shown in light grey (Dvoretzky-Kiefer-Wolfowitz inequality).

Scale bars, (**g**) 100 μm, 20 μm; (**h**) 20 μm, 5 μm.

Figure 3: Coordinately regulated gene sets contribute to the transcriptional identities of layer

6 corticothalamic neuron subtypes. (**a**) Weighted gene co-expression network analysis

(WGCNA) on variance-stabilized gene expression estimates identifies modules of coordinately

regulated genes, which are grouped using hierarchical clustering of module eigengenes. (**b**)

Pearson correlation of each module eigengene with parameters for both transcriptional subtype

and label. Significance of each correlation was determined (*) using the Pearson's product

moment test (p < 0.01; Benjamini-Hochberg corrected). (**c**) Pearson correlation of each module

eigengene with component rotations for principal components 1-5. (**d,e**) Hypergeometric test

for enrichment of the 286 genes differentially expressed between transcriptionally defined

L6CThN subtypes (**d**) and genes associated with neuronal activity (**e**) within each module (p <

0.01; Benjamini-Hochberg corrected).

Figure 4: Variation in the transcriptional profiles of layer 6 corticothalamic neurons is defined

by both subtype-specific genes and genes reflecting laminar location within layer 6. (**a,d**) tSNE

plots showing the eigenvalue for each cell for the two WGCNA modules most correlated with

the first principal component (**a**, Midnight Blue; **d**, Turquoise). (**b,c,e,f**) Gene expression tSNE

plots (*left*) showing the normalized gene expression level in each cell for representative genes

with significant weights on PC1. (**b**) and (**c**) are representative members of the Midnight Blue

module, and (**e**) and (**f**) belong to the Turquoise module. Single molecule fluorescence *in situ*

hybridization (smFISH, *middle*) showing images of mRNAs detected for each gene of interest

(magenta), tdTomato (green), NeuN (cyan), and DAPI (blue) in layer 6 of Ntsr1-Cre;tdTomato

mice. Quantitative gene expression analysis of smFISH (*right*) showing the number of mRNAs

expressed per neuron as a function of normalized vertical position in layer 6 and neuronal cell

type (L6CThNs: Ntsr1;tdTomato-positive;NeuN-positive neurons in green and non-L6CThNs:

Ntsr1;tdTomato-negative, NeuN-positive neurons in grey). Curves represent Loess fits to the

individual data points, grouped by cell type, and shaded areas correspond to 95% confidence

intervals. Statistics: (**b**, *Lamp5*) "Subtype specific" $p < 7.3231 \times 10^{-19}$; "CThN+ position specific" p

$< 1.175 \times 10^{-17}$; "CThN- position specific" $p < 0.0035$; (**c**, *Pantr1*) "Subtype specific" $p < 0.9931$;

"CThN+ position specific" $p < 9.243 \times 10^{-11}$; "CThN- position specific" $p < 1.021 \times 10^{-14}$; (**e**,

Serpini1) "Subtype specific" $p < 1.606 \times 10^{-06}$; "CThN+ position specific" $p < 1.045 \times 10^{-06}$; "CThN-

position specific" $p < 0.3342$; (**f**, *Gabra5*) "Subtype specific" $p < 0.1020$; "CThN+ position

specific" $p < 1.994 \times 10^{-3}$; "CThN- position specific" $p < 0.09873$.

Figure 5: A signature of neuronal activity significantly contributes to the transcriptional

identities of layer 6 corticothalamic neuron subtypes. (**a**) Module eigengenes for the four

modules with significant enrichment for genes associated with neuronal activity. (**b,c**) tSNE plots

showing the discretized expression of genes with the highest (**b**) and second highest (**c**) weights

in each direction of PC1. (**d,e**) tSNE plots showing the discretized expression of genes with the

highest (**d**) and second highest (**e**) weights in each direction of PC2. (**f,g**) tSNE plots showing the

discretized expression levels of four genes reflecting neuronal activity: *Bdnf* and *Camk2n1* (**f**),

and *Fos* and *Plk5* (**g**).

Figure 6: Sustained modulation of gene expression after sensory manipulation in layer 6

corticothalamic neurons. (**a**) Experimental design for sensory manipulation and single-cell

transcriptional analysis of L6CThNs from the barrel cortex at 1 and 7 days post manipulation. (**b**)

tSNE plot of all 1023 neurons obtained from baseline (Day 0), 1, and 7 days following sensory

manipulation using the 286 genes identified as differentially expressed between L6CThN

subtypes at day 0. Day 0 neurons are labeled by transcriptional subtype. Day 1 and Day 7

neurons are colored light gray. tSNE positions were fit to a Gaussian mixture model (black lines)

which was used to classify Day 1 and Day 7 neurons as members of the previously defined

L6CThN transcriptional subtypes. (**c**) tSNE plot colored by transcriptional subtype as assigned in

(**b**). 10 of 340 neurons (2.9%) from Day 0 were assigned to a different subtype than in **Fig. 2a**

and are labeled light green. (**d**) tSNE plot with neurons colored by projection label. (**e**) k-means

clustering analysis of mean-centered gene expression, aggregated by Day and transcriptional

subtype (Subtype 1: gold; Subtype 2: blue) for genes with significant differential expression after

sensory manipulation. Semi-transparent lines represent individual genes and bold lines

represent cluster centroids.

Figure 7: Pseudotemporal reconstruction of transcriptional responses to sensory manipulation

in layer 6 corticothalamic neurons. (**a**) Discriminative dimensionality reduction projection of

1023 L6CThNs using genes identified as significantly differentially expressed after sensory

manipulation. Neurons are colored by day relative to manipulation. (**b**) Density distribution of

L6CThNs across pseudotime, grouped by day following manipulation. (**c**) Heatmap of normalized

response curves for the 1507 genes with significant differential expression across pseudotime,

and significantly enriched gene sets identified for each cluster ($p < 1.0 \times 10^{-2}$, hypergeometric

test).

Figure 8: Sensory manipulation induces distinct cellular responses in layer 6 corticothalamic

neurons biased with respect to transcriptional subtype. (**a**) Distribution of the pairwise

Euclidean distances within each subtype for subtype 1 (*left*, gold) and subtype 2 (*right*, blue),

using variance-stabilized expression estimates for all expressed genes, showing a significant

increase in heterogeneity of gene expression for both subtype 1 and subtype 2. (**b**) Distribution

of pairwise inter-subtype Euclidean distances between transcriptionally defined L6CThN

subtypes across all expressed genes plotted for each day following sensory manipulation. There

is a significant divergence between these two subtypes across time points as indicated by a

positive shift in the distances after induction of experience-dependent plasticity. (**c**) Density

distributions of L6CThNs at each day plotted across pseudotime for the two transcriptional

subtypes of L6CThNs. (**d**) Discriminative dimensionality reduction projection of 1023 L6CThNs

shown in **Fig. 7a**, now colored by transcriptional subtype. Red and blue arrows indicate the first

major cellular response branch point following sensory manipulation. Grey arrow indicates the

tree root state and direction of response progression. Pie charts indicate the proportion of each

subtype in the population for each branch. (**e**) BEAM analyses of gene sets with significant

differential expression dependent on either major branch point showing all significant DE genes.

(**f-h**) BEAM heatmap for branch-dependent transcription factors not detected in the aggregate

pseudotime response (**f**), presynaptic proteins (**g**), and ligand-gated neurotransmitter receptors

(**h**).

SUPPLEMENTARY INFORMATION (provided separately)

REFERENCES

1. Junker, J.P. & van Oudenaarden, A. Every cell is special: genome-wide studies add a new

dimension to single-cell biology. *Cell* **157**, 8-11 (2014).

2. Liu, S. & Trapnell, C. Single-cell transcriptome sequencing: recent advances and remaining

challenges. *F1000Res* **5**, 182 (2016).

3. Molyneaux, B.J., Arlotta, P., Menezes, J.R. & Macklis, J.D. Neuronal subtype specification in

the cerebral cortex. *Nat Rev Neurosci* **8**, 427-437 (2007).

4. West, A.E. & Greenberg, M.E. Neuronal activity-regulated gene transcription in synapse

development and cognitive function. *Cold Spring Harb Perspect Biol* **3** (2011).

5. Lyons, M.R. & West, A.E. Mechanisms of specificity in neuronal activity-regulated gene

transcription. *Prog Neurobiol* **94**, 259-295 (2011).

6. Flavell, S.W. & Greenberg, M.E. Signaling mechanisms linking neuronal activity to gene

expression and plasticity of the nervous system. *Annu Rev Neurosci* **31**, 563-590 (2008).

7. Greig, L.C., Woodworth, M.B., Galazo, M.J., Padmanabhan, H. & Macklis, J.D. Molecular logic

of neocortical projection neuron specification, development and diversity. *Nat Rev Neurosci*

14, 755-769 (2013).

8. Molyneaux, B.J.*, et al.* DeCoN: genome-wide analysis of in vivo transcriptional dynamics

during pyramidal neuron fate selection in neocortex. *Neuron* **85**, 275-288 (2015).

9. Darmanis, S.*, et al.* A survey of human brain transcriptome diversity at the single cell level.

Proc Natl Acad Sci U S A **112**, 7285-7290 (2015).

10. Lake, B.B.*, et al.* Neuronal subtypes and diversity revealed by single-nucleus RNA sequencing

of the human brain. *Science* **352**, 1586-1590 (2016).

11. Pollen, A.A.*, et al.* Low-coverage single-cell mRNA sequencing reveals cellular heterogeneity

and activated signaling pathways in developing cerebral cortex. *Nat Biotechnol* **32**, 1053-

1058 (2014).

12. Tasic, B.*, et al.* Adult mouse cortical cell taxonomy revealed by single cell transcriptomics.

Nat Neurosci **19**, 335-346 (2016).

13. Zeisel, A.*, et al.* Brain structure. Cell types in the mouse cortex and hippocampus revealed by

single-cell RNA-seq. *Science* **347**, 1138-1142 (2015).

14. Hevner, R.F.*, et al.* Beyond laminar fate: toward a molecular classification of cortical

projection/pyramidal neurons. *Dev Neurosci* **25**, 139-151 (2003).

15. Sugino, K.*, et al.* Molecular taxonomy of major neuronal classes in the adult mouse

forebrain. *Nat Neurosci* **9**, 99-107 (2006).

16. Sorensen, S.A.*, et al.* Correlated gene expression and target specificity demonstrate

excitatory projection neuron diversity. *Cereb Cortex* **25**, 433-449 (2015).

17. Sanes, J.R. & Masland, R.H. The types of retinal ganglion cells: current status and

implications for neuronal classification. *Annu Rev Neurosci* **38**, 221-246 (2015).

18. Wagner, A., Regev, A. & Yosef, N. Revealing the vectors of cellular identity with single-cell

genomics. *Nat Biotechnol* **34**, 1145-1160 (2016).

19. Dueck, H., Eberwine, J. & Kim, J. Variation is function: Are single cell differences functionally

important?: Testing the hypothesis that single cell variation is required for aggregate

function. *Bioessays* **38**, 172-180 (2016).

20. Bourassa, J. & Deschenes, M. Corticothalamic projections from the primary visual cortex in

rats: a single fiber study using biocytin as an anterograde tracer. *Neuroscience* **66**, 253-263

(1995).

21. Bourassa, J., Pinault, D. & Deschenes, M. Corticothalamic projections from the cortical barrel

field to the somatosensory thalamus in rats: a single-fibre study using biocytin as an

anterograde tracer. *Eur J Neurosci* **7**, 19-30 (1995).

22. Briggs, F., Kiley, C.W., Callaway, E.M. & Usrey, W.M. Morphological substrates for parallel

streams of corticogeniculate feedback originating in both V1 and V2 of the macaque

monkey. *Neuron* **90**, 388-399 (2016).

23. Briggs, F. & Usrey, W.M. Parallel processing in the corticogeniculate pathway of the

macaque monkey. *Neuron* **62**, 135-146 (2009).

24. Katz, L.C. Local circuitry of identified projection neurons in cat visual cortex brain slices. *J*

Neurosci **7**, 1223-1249 (1987).

25. Killackey, H.P. & Sherman, S.M. Corticothalamic projections from the rat primary

somatosensory cortex. *J Neurosci* **23**, 7381-7384 (2003).

26. Kwegyir-Afful, E.E. & Simons, D.J. Subthreshold receptive field properties distinguish

different classes of corticothalamic neurons in the somatosensory system. *J Neurosci* **29**,

964-972 (2009).

27. Usrey, W.M. & Fitzpatrick, D. Specificity in the axonal connections of layer VI neurons in tree

shrew striate cortex: Evidence for distinct granular and supragranular systems. *J Neurosci*

16, 1203-1218 (1996).

28. Zhang, Z.W. & Deschenes, M. Intracortical axonal projections of lamina VI cells of the

primary somatosensory cortex in the rat: a single-cell labeling study. *J Neurosci* **17**, 6365-

6379 (1997).

29. Thomson, A.M. Neocortical layer 6, a review. *Front Neuroanat* **4**, 13 (2010).

30. Gong, S.*, et al.* Targeting Cre recombinase to specific neuron populations with bacterial

artificial chromosome constructs. *J Neurosci* **27**, 9817-9823 (2007).

31. Kim, J., Matney, C.J., Blankenship, A., Hestrin, S. & Brown, S.P. Layer 6 corticothalamic

neurons activate a cortical output layer, layer 5a. *J Neurosci* **34**, 9656-9664 (2014).

32. Bortone, D.S., Olsen, S.R. & Scanziani, M. Translaminar inhibitory cells recruited by layer 6

corticothalamic neurons suppress visual cortex. *Neuron* **82**, 474-485 (2014).

33. Brennecke, P*., et al.* Accounting for technical noise in single-cell RNA-seq experiments. *Nat*

Methods **10**, 1093-1095 (2013).

34. Hicks, S., Teng, M. & Irizarry, R.A. On the widespread and critical impact of systematic bias

and batch effects in single-cell RNA-seq data. *BioRxiv* **doi:10.1101/025528** (2015).

35. Chung, N.C. & Storey, J.D. Statistical significance of variables driving systematic variation in

high-dimensional data. *Bioinformatics* **31**, 545-554 (2015).

36. Kiselev, V.Y*., et al.* SC3: consensus clustering of single-cell RNA-seq data. *Nat Methods* **14**,

483-486 (2017).

37. Lin, P., Troup, M. & Ho, J.W. CIDR: Ultrafast and accurate clustering through imputation for

single-cell RNA-seq data. *Genome Biol* **18**, 59 (2017).

38. Lein, E.S*., et al.* Genome-wide atlas of gene expression in the adult mouse brain. *Nature* **445**,

168-176 (2007).

39. Sauvageau, M*., et al.* Multiple knockout mouse models reveal lincRNAs are required for life

and brain development. *Elife* **2**, e01749 (2013).

40. Joo, K.M*., et al.* Distribution of vasoactive intestinal peptide and pituitary adenylate cyclase-

activating polypeptide receptors (VPAC1, VPAC2, and PAC1 receptor) in the rat brain. *J Comp*

Neurol **476**, 388-413 (2004).

41. Sun, Q.Q., Prince, D.A. & Huguenard, J.R. Vasoactive intestinal polypeptide and pituitary

adenylate cyclase-activating polypeptide activate hyperpolarization-activated cationic

current and depolarize thalamocortical neurons in vitro. *J Neurosci* **23**, 2751-2758 (2003).

42. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 559 (2008).

43. Lacar, B.*, et al.* Nuclear RNA-seq of single neurons reveals molecular signatures of activation.

Nat Commun **7**, 11022 (2016).

44. Mardinly, A.R.*, et al.* Sensory experience regulates cortical inhibition by inducing IGF1 in VIP

neurons. *Nature* **531**, 371-375 (2016).

45. Cho, J.H., Huang, B.S. & Gray, J.M. RNA sequencing from neural ensembles activated during

fear conditioning in the mouse temporal association cortex. *Sci Rep* **6**, 31753 (2016).

46. Kim, T.K.*, et al.* Widespread transcription at neuronal activity-regulated enhancers. *Nature*

465, 182-187 (2010).

47. Wallace, H. & Fox, K. The effect of vibrissa deprivation pattern on the form of plasticity

induced in rat barrel cortex. *Somatosens Mot Res* **16**, 122-138 (1999).

48. Fox, K. *Barrel Cortex* (Cambridge University Press, Cambridge, 2008).

49. Fraley, C., Raftery, A.E., Murphy, T.B. & Scrucca, L. mclust Version 4 for R: Normal Mixture

Modeling for Model-Based Clustering, Classification, and Density Estimation Technical

Report No. 597. *Department of Statistics, University of Washington* (2012).

50. Fraley, C. & Raftery, A.E. Model-based clustering, discriminant analysis and density

estimation. *Journal of the American Statistical Association* **97**, 611-631 (2002).

51. Ehlers, M.D. Activity level controls postsynaptic composition and signaling via the ubiquitin-

proteasome system. *Nat Neurosci* **6**, 231-242 (2003).

52. Spiegel, I.*, et al.* Npas4 regulates excitatory-inhibitory balance within neural circuits through

cell-type-specific gene programs. *Cell* **157**, 1216-1229 (2014).

53. Whitney, O.*, et al.* Core and region-enriched networks of behaviorally regulated genes and

the singing genome. *Science* **346**, 1256780 (2014).

54. Frenkel, M.Y. & Bear, M.F. How monocular deprivation shifts ocular dominance in visual

cortex of young mice. *Neuron* **44**, 917-923 (2004).

55. Glazewski, S., Benedetti, B.L. & Barth, A.L. Ipsilateral whiskers suppress experience-

dependent plasticity in the barrel cortex. *J Neurosci* **27**, 3910-3920 (2007).

56. Madisen, L.*, et al.* A robust and high-throughput Cre reporting and characterization system

for the whole mouse brain. *Nat Neurosci* **13**, 133-140 (2010).

57. Schindelin, J.*, et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods*

9, 676-682 (2012).

58. Agmon, A. & Connors, B.W. Thalamocortical responses of mouse somatosensory (barrel)

cortex in vitro. *Neuroscience* **41**, 365-379 (1991).

59. Saxena, A.*, et al.* Trehalose-enhanced isolation of neuronal sub-types from adult mouse

brain. *Biotechniques* **52**, 381-385 (2012).

60. Picelli, S.*, et al.* Full-length RNA-seq from single cells using Smart-seq2. *Nat Protoc* **9**, 171-

181 (2014).

61. Mudge, J.M. & Harrow, J. Creating reference gene annotation for the mouse C57BL6/J

genome assembly. *Mamm Genome* **26**, 366-378 (2015).

62. Blackshaw, S. High-throughput RNA in situ hybridization in mouse retina. *Methods Mol Biol*

935, 215-226 (2013).

63. Goff, L.A.*, et al.* Spatiotemporal expression and transcriptional perturbations by long

noncoding RNAs in the mouse brain. *Proc Natl Acad Sci U S A* **112**, 6855-6862 (2015).

64. Mueller, F.*, et al.* FISH-quant: automatic counting of transcripts in 3D FISH images. *Nat*

Methods **10**, 277-278 (2013).

65. Li, X., Glazewski, S., Lin, X., Elde, R. & Fox, K. Effect of vibrissae deprivation on follicle

innervation, neuropeptide synthesis in the trigeminal ganglion, and S1 barrel cortex

plasticity. *J Comp Neurol* **357**, 465-481 (1995).

66. Kim, D., Langmead, B. & Salzberg, S.L. HISAT: a fast spliced aligner with low memory

requirements. *Nat Methods* **12**, 357-360 (2015).

67. Trapnell, C*., et al.* Differential analysis of gene regulation at transcript resolution with RNA-

seq. *Nat Biotechnol* **31**, 46-53 (2013).

68. Trapnell, C*., et al.* Differential gene and transcript expression analysis of RNA-seq

experiments with TopHat and Cufflinks. *Nat Protoc* **7**, 562-578 (2012).

69. Trapnell, C*., et al.* The dynamics and regulators of cell fate decisions are revealed by

pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381-386 (2014).

70. Huber, W*., et al.* Orchestrating high-throughput genomic analysis with Bioconductor. *Nat*

Methods **12**, 115-121 (2015).

71. Wood, S.N. Fast stable restricted maximum likelihood and marginal likelihood estimation of

semiparametric generalized linear models. *Journal of the Royal Statistical Society: Series B*

(Statistical Methodology) **73**, 3-36 (2011).

72. Krijthe, J. Rtsne: T-distributed stochastic neighbor embedding using Barnes-Hus

implementation. R package version 0.10. https://CRAN.R-project.org/package=Rtsne.

(2015).

73. Macosko, E.Z*., et al.* Highly parallel genome-wide expression profiling of individual cells

using nanoliter droplets. *Cell* **161**, 1202-1214 (2015).

74. Ashburner, M*., et al.* Gene ontology: tool for the unification of biology. The Gene Ontology

Consortium. *Nat Genet* **25**, 25-29 (2000).

75. Yu, G. & He, Q.Y. ReactomePA: an R/Bioconductor package for reactome pathway analysis

and visualization. *Mol Biosyst* **12**, 477-479 (2016).

76. Yu, G., Wang, L.G., Han, Y. & He, Q.Y. clusterProfiler: an R package for comparing biological

themes among gene clusters. *OMICS* **16**, 284-287 (2012).

77. Kolde, R. pheatmap: Pretty heatmaps. R package verion 1.0.8. https://CRAN.R-

project.org/package=pheatmap. (2015).

78. Subramanian, A.*, et al.* Gene set enrichment analysis: a knowledge-based approach for

interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550

(2005).

Figure 1
Chevée et al.

Figure 2
Chevée et al.

Figure 5
Chevée et al.

Figure 5
Chevée et al.

a

b

c

Figure 7
Chevée et al.

Figure 8
Chevée et al.