1

The tumor suppressor APC is an attenuator of spindle-pulling forces during *C. elegans* asymmetric cell division

-Author names and Affiliations

Kenji Sugioka[1,2,3], Lars-Eric Fielmich[4], Kota Mizumoto[2], Bruce Bowerman[3], Sander van den Heuvel[4], Akatsuki Kimura[5, 6] and Hitoshi Sawa[1,2,6]

[1]Multicellular Organization Laboratory, National Institute of Genetics, 1111 Yata, Mishima, 411-8540 Japan

[2]RIKEN Center for Developmental Biology, 2-2-3 Minatojima-minamimachi, Chuo-ku, Kobe 650-0047 Japan

[3]Institute of Molecular Biology, University of Oregon, Eugene, OR 97403 USA

[4]Developmental Biology, Biology Department, Science 4 Life, Utrecht University, Padualaan 8,

3584 CH, Utrecht, Netherlands

[5]Cell Architecture Laboratory, National Institute of Genetics, 1111 Yata, Mishima, 411-8540 Japan

[6]Department of Genetics, School of Life Science, Sokendai, 1111 Yata, Mishima, 411-8540 Japan

-Corresponding authors

Hitoshi Sawa

Multicellular Organization Laboratory, National Institute of Genetics, 1111 Yata,

Mishima, 411-8540 Japan

hisawa@nig.ac.jp

Phone: +81-55-981-6845

Fax: +81-55-981-6846

Akatsuki Kimura

Cell Architecture Laboratory, National Institute of Genetics, 1111 Yata,

Mishima, 411-8540 Japan

akkimura@nig.ac.jp

Sander van den Heuvel

Developmental Biology, Biology Department, Utrecht University, Padualaan 8,

3584 CH, Utrecht, Netherlands

S.J.L.vandenHeuvel@uu.nl

Present address:

Kota Mizumoto

Department of Zoology, the University of British Columbia, Vancouver, Canada,

V6T 1Z3

Abstract

The adenomatous polyposis coli (APC) tumor suppressor has dual functions in

Wnt/β-catenin signaling and accurate chromosome segregation, and is frequently

mutated in colorectal cancers. Although APC contributes to proper cell division, the

underlying mechanisms remain poorly understood. Here we show that *C. elegans*

APR-1/APC is an attenuator of the pulling forces acting on the mitotic spindle.

During asymmetric cell division of the *C. elegans* zygote, a LIN-5/NuMA protein

complex localizes dynein to the cell cortex to generate pulling forces on astral

microtubules that position the mitotic spindle. We found that APR-1 localizes to the

anterior cell cortex in a Par-aPKC polarity-dependent manner and suppresses anterior

centrosome movements. Our combined cell biological and mathematical analyses

support the conclusion that cortical APR-1 reduces force generation by stabilizing

microtubule plus ends at the cell cortex. Furthermore, APR-1 functions in

coordination with LIN-5 phosphorylation to attenuate spindle pulling forces. Our

results document a physical basis for spindle-pulling force attenuation, which may be

generally used in asymmetric cell division, and when disrupted potentially contributes

to division defects in cancer.

Significance Statement

APC (adenomatous polyposis coli) is a Wnt signaling component as well as a

microtubule-associated protein, and its mutations are frequently associated with

colorectal cancers in humans. Although APC stabilizes microtubules (MTs), its

mechanical role during cell division is largely unknown. Here we show that APC is an

attenuator of forces acting on the mitotic spindle during asymmetric cell division of

the *C. elegans* zygote. We performed live-imaging, laser-microsurgery, and numerical

simulation to show how APC suppresses spindle pulling force generation by

stabilizing microtubule plus-ends and reducing microtubule catastrophe frequency at

the cell cortex. Our study is the first to document a mechanical role for the APC

protein, and provides a physical basis for spindle-pulling force attenuation.

\body

Introduction

The mitotic spindle segregates chromosomes and determines the plane of cell

cleavage during animal cell division. Forces that act on the mitotic spindle regulate its

position to produce daughter cells of the proper size, fate and arrangement, thereby

playing a significant role in asymmetric cell division, tissue integrity and

organogenesis. In various organisms, cells regulate spindle positioning through

cortical force generators that pull on astral microtubules (1-5). An evolutionarily

conserved force generator complex, consisting of LIN-5/NuMA, GPR-1,2/LGN and

Gα, interacts with dynein and dynamic astral microtubules to position the mitotic

spindle during the asymmetric divisions of the *C. elegans* early embryo (4),

Drosophila and mammalian neuroblasts (1, 2), and skin stem cells (3). Although Par-

aPKC polarity and cell cycle regulators are known to control spindle positioning (4,

6), how the forces are regulated spatiotemporally to position the spindle in various

cell types during development remains poorly understood.

The tumor suppressor adenomatous polyposis coli (APC) is a widely

conserved multifunctional protein with two major roles. First, APC functions as part

of a degradation complex to down-regulate β-catenin-TCF dependent transcription,

thereby controlling cell fate and proliferation in various cell types (7). Second, APC

functions as a microtubule-associated protein to stabilize MTs. It has been suggested

that this function of APC regulates cell migration (8, 9), spindle orientation (10, 11),

and chromosome segregation (12, 13). In mammals, loss of the former function is

closely associated with colon cancer (14, 15). Loss of the latter function causes

spindle positioning defects (16, 17) and chromosome instability (CIN) (18-20), a

hallmark of metastatic tumors (21), suggesting that the cytoskeletal roles of APC

during mitosis are also relevant for oncogenesis. How APC regulates the mitotic

spindle remains poorly understood and is complicated by its multiple functions,

binding-partners and cellular locations (12, 22).

Yeast and fly studies have suggested that APC at the cell cortex contributes to

mitotic spindle positioning. Kar9, a yeast protein with limited homology to APC,

localizes asymmetrically to the cell cortex of budding daughter cells through type V

myosin-dependent transport of growing microtubule ends (23-25). Cortical Kar9

captures microtubules (MTs) by binding yeast EB1, and promotes alignment of the

spindle along the mother-bud axis (24-27). *Drosophila* APC2 predominantly localizes

to the cell cortex in syncytial embryos. APC2 mutants show a CIN phenotype,

presumably because APC2 is required for proper centrosome separation (28). The

forces that mediate centrosome separation have been proposed to depend on APC2

connecting astral MTs to cortical actin (28). However, the mechanism by which

cortical APC regulates spindle-pulling forces has not been directly addressed in any

organism.

We report here that loss of cortical APR-1/APC disrupts asymmetries in

spindle movements during mitotic division of the *C. elegans* zygote. In wild-type

embryos, the net pulling forces acting on the mitotic spindle become higher in the

posterior compared to the anterior, causing the spindle to move posteriorly during

metaphase and anaphase (spindle displacement) (29, 30). In anaphase, the posterior

spindle pole swings along the transverse axis (spindle oscillation), while the anterior

pole remains relatively stable. We found APR-1 to be enriched at the anterior cortex

in a PAR-polarity dependent manner. Depletion of APR-1 resulted in anterior pole

oscillations that resemble those of the posterior pole. Moreover, laser-mediated

spindle severing showed that the spindle-pulling forces acting on the anterior spindle

pole are increased in *apr-1(RNAi)* embryos. Using live imaging and numerical

simulation, we found that the APR-1 dependent stabilization of MT-cortex

interactions negatively regulates the pulling forces acting on the anterior centrosome

in wild-type zygotes. Our study identifies APR-1 as an attenuator of spindle-pulling

forces, and improves our understanding of how cortical polarity precisely regulates

spindle positioning during asymmetric cell division.

Results and Discussion

APR-1/APC localizes asymmetrically to the cell cortex in a PAR and Frizzled protein dependent manner

We have previously shown that APR-1 localizes asymmetrically to the anterior cortex in the EMS blastomere at the six-cell stage and in post-embryonic seam cells, in response to Wnt signals that regulate the asymmetry of these divisions (31, 32). While analyzing GFP::APR-1 localization in early embryos, we noticed that APR-1 is also asymmetrically localized in the zygote, called P0, where roles for Wnt signaling have not been reported. APR-1 formed dot-like particles that were enriched within the anterior cortex throughout P0 cell division (APR-1 asymmetry) (Figure 1A). We quantified the number of APR-1 dots by counting the fluorescent foci with a signal above a threshold (see Materials and methods). The foci numbers changed from prophase to metaphase, and from anaphase to telophase. Nevertheless, we observed anterior enrichment of APR-1 foci throughout mitosis (Figure 1A and 1D).

It is well-established that the Par-aPKC system generates anterior-posterior (A-P) cell polarity to regulate the asymmetric division of P0, through interactions between anterior (PAR-3, PAR-6, PKC-3) and posterior (PAR-2, PAR-1) *partitioning defective* (PAR) proteins at the cell cortex (Figure 1B; 33). We found that APR-1 asymmetry in P0 was disrupted after RNAi knockdown of *par-3*, *pkc-3* or *par-2* (Figure 1C, 1E, and Figure S1), suggesting that its asymmetry is established through the Par-aPKC system.

In EMS and seam cells, the establishment of APR-1 asymmetry depends on Wnt proteins (31, 32). In P0, MOM-2 is the only Wnt protein that is maternally provided as mRNA (34), although the mRNA appears not to be translated until the 4-cell stage (35). As expected, we found that APR-1 localization was not affected in

1   *mom-2(or309)* null mutants, suggesting that the APR-1 asymmetry in P0 does not

2   require Wnt ligands (Figure 1C, 1E, and Figure S1).

3   Despite the lack of a requirement for MOM-2/Wnt, we observed altered APR-

1 localization after RNAi knockdown of downstream Wnt signaling components.

Specifically, knockdown of the Frizzled receptor MOM-5 or simultaneous inhibition

of the Dishevelled homologs, DSH-2 and MIG-5, increased the numbers of APR-1

foci at metaphase/anaphase in both the anterior and posterior cortex without altering

APR-1 expression levels (Figure 1C, 1E, Figure S1, and Figure S2A). Inhibition of

WRM-1/β-catenin did not affect APR-1 localization, and *mom-5(RNAi)* as well as

dsh-2;mig-5(RNAi) embryos still showed APR-1 asymmetry (Figure 1C, 1E, and

Figure S1). DSH-2 localizes to the posterior cell cortex during Wnt-dependent

asymmetric cell divisions later in development (31, 36). In contrast, DSH-2

localization in P0 was not asymmetric (Figure S2B), consistent with the lack of

Dishevelled requirement in APR-1 asymmetry. Interestingly, inhibition of the Axin

homolog PRY-1 and Casein kinase homolog KIN-19 resulted in loss of APR-1

asymmetry only during meta-anaphase, suggesting their partial requirement in the

establishment or maintenance of APR-1 asymmetry (Figure S1B and S1C).These

results are consistent with observations at a later developmental stage (37). We

conclude that APR-1 asymmetry in P0 is established by the Par-aPKC system with

partial involvement of Axin and Casein kinase, while Frizzled and Dishevelled

negatively regulate the levels of cortical APR-1.

APR-1 asymmetrically suppresses centrosome movements during P0 cell division

The Par-aPKC system independently regulates two P0 asymmetries: the segregation

of cell fate determinants (e.g. PIE-1 and PGL-1) and posterior mitotic spindle

displacement and thereby asymmetric cell cleavage. In *apr-1(RNAi)* embryos,

GFP::PIE-1 segregated into the posterior daughter cell as in wild-type embryos,

indicating that APR-1 is not involved in cytoplasmic determinant localization (Figure

S2C). In contrast, *apr-1(RNAi)* embryos showed abnormal spindle oscillations. In

wild type P0, posterior spindle displacement (represented by centrosome movements

along the A-P axis) starts during metaphase and continues during anaphase when it

coincides with transverse oscillations (represented by centrosome movements along

the transverse axis) of the two spindle poles (Figure 2A, 2B, 2D, 2E). The posterior

spindle pole oscillates more vigorously than the anterior pole (Figure 2B, 2E and

Video 1), as a result of higher posterior than anterior cortical pulling forces (38). In

apr-1(RNAi) embryos, spindle movements were exaggerated: in some embryos, the

mitotic spindle moved back and forth along the A-P axis (Figure 2C, 2D, and Video

2), and in some cases, the anterior spindle pole exhibited excessive transverse

oscillations, visible by the increased frequency and amplitude of the spindle pole

tracks (Figure 2C, 2E and Video 2). As a result, the total distance traveled by the

anterior centrosome significantly increased compared to that in control embryos

(Figure 2F). These data indicate that APR-1 suppresses anterior spindle pole

movements and thereby control spindle positioning during anaphase.

In *mom-5(ne12)* null mutant embryos, in which APR-1 levels were increased

at both the anterior and posterior cortex, we observed reduced posterior spindle pole

oscillations (Figure S3A and S3B). However, spindle pole oscillations were not

restored in *apr-1(RNAi); mom-5(null)* embryos (Figure S3B). These results suggest

APR-1-independent functions of MOM-5 that influence spindle movements. Because

of this, we could not determine the effects of excess cortical APR-1 on spindle pole

movements in the *mom-5(null)* background. However, in other aspects of spindle

dynamics described below, elevated cortical APR-1 localization potentiated APR-1

function.

APR-1 asymmetrically stabilizes microtubule-cortex interactions

As mammalian APC (39) and *C. elegans* APR-1 in the EMS cell (32) can stabilize

MTs, we hypothesized that anteriorly enriched APR-1 in the P0 cell may also increase

MT stability at the cell cortex to regulate asymmetric spindle movements. To assess

this possibility, we analyzed the MT-cortex interactions using live imaging of

GFP::β-tubulin expressing embryos. In kymographs of midplane images, astral

microtubules appear to persist longer on the anterior cell cortex than on the posterior,

consistent with previous observations (Figure 3A; 40). To precisely quantify MT-plus

end residence time at the cortex, we measured the duration of GFP::β-tubulin foci on

the flattened cell surface (Figure 3B). Most of the GFP::β-tubulin foci initially co-

localized with the EB1-related plus-end binding protein EBP-2 (96.1%; n = 255),

confirming that the foci represent MT plus-ends. Shortly after the cortical attachment,

EB1 dissociates from MT plus-ends, while some MTs remained at the cortex after the

release of EB1 (Fig. 3B and 3D). The numbers of such long-lived microtubule plus-

ends were higher anteriorly, accounting for the asymmetry in cortical MT residence

time in wild-type zygotes (Figure 3B-3D; red arrows in 3C, Video 3 and Video 4).

Notably, the MT residence time at the anterior cortex was significantly lower

in *apr-1(RNAi)* embryos than in the wild type (Figure 3C, 3E and Video 5). In

contrast, *mom-5* mutants with excess cortical APR-1 showed an increased MT

residence time at both the anterior and posterior cell cortex (Figure 3C, 3E and Video

6). RNAi knockdown of *apr-1* overcame this *mom-5* phenotype, reducing MT cortical

residence throughout the cortex (Figure 3C, 3E and Video 7). Thus, APR-1 stabilizes microtubule-cortex interactions and acts downstream of MOM-5 (Figure 4D).

APR-1 asymmetrically attenuates pulling forces acting on the mitotic spindle

The exaggerated anterior spindle pole movements in *apr-1(RNAi)* embryos implicate APR-1 in spindle-pulling force regulation. We investigated this possibility using spindle severing assays (Figure 4A; 41). After cutting the spindle midzone with a UV laser, the average peak velocities of the anterior and posterior spindle poles moving toward the cell cortex were calculated (Figure 4A). In control embryos, the posterior spindle pole moved faster than the anterior pole, as expected (Figure 4A, 4B, and Video 8). In *apr-1(RNAi)* embryos, we observed an increased average peak velocity specifically for the anterior spindle pole (Figure 4A, 4B, and Video 8). In *mom-5(null)* embryos with excess cortical APR-1, both the anterior and posterior spindle poles showed reduced average peak velocities (Figure 4B and Video 8). Combined *apr-1(RNAi);mom-5(null)* embryos showed increased average peak velocities and resembled *apr-1(RNAi)* embryos (Figure 4B and Video 8). These results indicate that the cortical levels of APR-1 inversely correlate with spindle-pulling forces and suggest a role for APR-1 as cortical pulling force attenuator (Figure 4D).

APR-1-dependent stabilization of MTs accounts for reduced pulling forces on the anterior spindle pole

We have shown that APR-1 is enriched at the anterior cell cortex, promotes cortical MT residence times anteriorly, and suppresses both spindle-pulling forces and anterior spindle pole oscillations, raising the possibility that these processes are mechanistically linked. It has been shown that cortical pulling forces are generated

when MTs reaching the cortex meet dynein and undergo catastrophe (transition from

MT plus end growth to rapid shrinkage) (42). Therefore, we hypothesized that cortical

APR-1 reduces the MT catastrophe frequency and thereby attenuates force generation

and spindle movement. However, it is not clear whether the magnitude of APR-1-

dependent cortical MT stabilization is sufficient to suppress spindle movement.

We decided to examine this issue using numerical simulation. First, we

estimated MT catastrophe frequencies from their cortical residence time

(Supplementary Table 1, Figure S4). In control embryos, the estimated catastrophe

frequency at the anterior cortex was about half of that at the posterior cortex. Such a

reduced catastrophe frequency was not detected at the anterior cortex of *apr-1(RNAi)*

embryos, indicating that in wild type embryos the catastrophe frequency is suppressed

by APR-1.

We set the rescue frequency of all MTs high, so that soon after the MTs start

to shorten, they regrow to reach the cortex (Supplemental Table 2). This assumption

was introduced to make the number of MTs reaching the cortex almost constant

regardless of the differences in catastrophe frequencies between anterior and posterior,

which is the case in living embryos (Video 3). Without this assumption, the number

of MTs reaching the cortex should be ~2-fold higher at the anterior because the

catastrophe frequency is about half of the posterior catastrophe frequency. The

mechanistic basis of this assumption is provided by the *in vivo* observation that

individual microtubules appear to form bundles, and multiple EB1 tracks move along

a bundled fiber toward the cell cortex, making rescue frequency of the fiber higher

than individual MTs (Video 4), which is consistent to the previous observation (43).

We conducted 3-dimensional simulations of spindle movements. As in

previous simulations (44-47), the spindle moves as a result of three kinds of forces

acting on astral MTs that radiate from each spindle pole (Figure 2G). First, all MTs

generate pulling forces proportional to their length ("cytoplasmic pulling force"). This

force is important for positioning of the spindle in the cell center during mitotic

prophase (45, 48, 49), and is also critical for oscillation (38). Second, MTs that reach

the cell cortex generate the pulling force at their plus ends only when they undergo

catastrophe ("cortical pulling force"). The current theory for the basis of oscillation is

that when the spindle poles move toward one side, the pulling force from that side

becomes stronger ("positive feedback" or "negative friction"), while the opposing

centering force also increases (38, 50, 51). With this mechanism, the spindle is not

stabilized at the center but oscillates. In our model, the frequency of the force

generation depends on the number of active cortical force generators and the MT

residence time controlled by APR-1, both of which have A-P asymmetry. The third

force connects the anterior and posterior spindle poles. We assumed a spring-like

connection between the poles that was weakened after anaphase onset to mimic the

spindle elongation.

Numerical simulations were conducted for control, *apr-1(RNAi),* and *mom-5(null)* situations (Figure S5), by setting the catastrophe frequency to values estimated

from experimental data (e.g. 0.31 /s for the anterior and 0.72 /s for the posterior, see

Supplementary Table 1). The simulation results indicated that the APR-1-dependent

stabilization of MTs is sufficient to suppress oscillation of the anterior pole (Figure

2H). In wild-type simulations, the spindle moved toward the posterior and elongated

upon anaphase onset (Figure S5A and Video 9). The oscillations perpendicular to the

A-P axis were also reproduced for both spindle poles (Figure S5B). In *apr-1(RNAi)*

simulations, in which the catastrophe frequency at the anterior cortex was increased,

the amplitude of the anterior spindle pole oscillations was increased (Figure 2H,

Figure S5 and Video 9). Furthermore, the average peak velocities of anterior poles in

the severing experiments were also consistent with the forces acting on anterior

spindle poles in our simulations (Figure 4C). Overall, the numerical simulations

supported the hypothesis that the APR-1-dependent stabilization of MTs at the cortex

can suppress spindle pole oscillations through the reduction of force generation.

Anterior APR-1 and LIN-5 phosphorylation together attenuate spindle pulling forces

We investigated the significance of spindle pulling force attenuators in asymmetric

cell division. Along with APR-1, we focused on the LIN-5 protein. LIN-5 interacts

with cortical GPR-1/2 and dynein in cortical force generation (52). We have

previously reported that anteriorly-localized PKC-3/aPKC phosphorylates LIN-5 to

attenuate cortical-pulling forces (53). We edited the *lin-5* genomic locus to substitute

four aPKC phosphorylated serine residues with alanine by CRISPR/Cas9-mediated

homologous recombination (*lin-5 4A* mutation). In spindle severing experiments,

combining *apr-1(RNAi)* and the *lin-5 4A* mutation caused significantly enhanced

average peak velocities of the anterior poles as compared to *apr-1(RNAi)* embryos

(Figure 5A). Compared to *lin-5 4A* embryos, the increase in anterior peak velocity

was not significant (p= 0.07; Figure 5A). However, in contrast to the single mutants,

the ratio of anterior to posterior centrosome peak velocities in *apr-1(RNAi)*; *lin-5 4A*

double mutants was significantly reduced compared to wild-type controls (Figure 5B).

These data suggest that the Par-aPKC-dependent asymmetric localization of APR-1

and phosphorylation of LIN-5 together attenuate cortical pulling forces to generate

pulling force asymmetry that positions the mitotic spindle (Figure 5C-5E).

Conclusion

In this study, we investigated how the APR-1/APC protein regulates mitotic spindle movements in the *C. elegans* one-cell embryo, a well-established model for asymmetric cell division. We observed that APR-1/APC becomes asymmetrically enriched at the anterior cell cortex, dependent on the Par-PKC-3 polarity pathway. We found that APR-1 attenuates spindle pulling forces, most likely though stabilization of MTs at the anterior cell cortex. In concert, Wnt signaling components MOM-5/Frizzled and Disheveled proteins suppressed cortical accumulation of APR-1, thereby contributing to the correct levels of pulling forces. To test these assumptions, we performed numerical simulations, which closely mimicked the spindle movements in wild-type and mutant embryos. These combined data strongly support the conclusion that MT stabilization by APR-1 contributes to correct spindle positioning. Finally, we provide evidence to suggest that asymmetric APR-1 enrichment and PKC-3 phosphorylation of LIN-5 act in parallel to regulate asymmetric cell division. These conclusions are likely to apply broadly and improve our understanding of the microtubule-associated functions of APC.

Although APC is a component of Wnt signaling, its localization has been reported to be regulated by the Par-aPKC polarity pathway in migrating mammalian astrocytes (54), and during axonal differentiation of developing hippocampal neurons (55), as we observed in the *C. elegans* one-cell embryo. Scratching of astrocyte monolayers in wound-healing assays triggers APC localization to the cell cortex at the leading edge, in response to CDC42-induced Par-aPKC polarity and Wnt5a signaling (56). Interestingly, polarity establishment in this system is followed by centrosome re-orientation through APC-microtubule interactions (54). Thus, the mechanisms that control centrosome positioning through interactions between Par polarity, Wnt

signaling, and APC may be conserved across species. The dynamic change in cortical

APR-1 levels during P0 cell division is intriguing: this may reflect cell cycle

dependent activation of the Wnt signaling pathway as reported in fly and mammalian

cultured cells (57).

While the roles of cortical APC have been unclear, it was previously proposed

that APC stabilizes microtubules through microtubule plus-end binding protein EB1

(54, 58). Consistently, in the *C. elegans* EMS blastomere, cortical APC stabilizes MT

ends coated with EB1 (32). However, a few examples including the present study

indicate that cortical APC can stabilize microtubules independently of EB1. First,

truncated mammalian APC that lacks the EB1 interaction domain has been shown to

localize to the cell cortex and to MTs in epithelial cells (59). In addition, *Drosophila*

APC2, which lacks the C-terminal EB1 binding domain, interacts with microtubule

plus ends at the cortex and contributes to centrosome segregation (28). In our study,

APR-1 at the anterior cortex stabilizes MTs but the mean cortical residence time of

EBP-2/EB1 was symmetric. We also observed that the cortical residence time of EB1

is much shorter than that of MTs in P0, as reported previously (43). Therefore, APR-

1 at the anterior cortex of P0 likely stabilizes MTs independently of EB1 binding. We

observed recently that deleting all EB family members has limited effects on spindle

behavior and viability in *C. elegans* (60). Therefore, the microtubule stabilizing

effects of cortical APC probably do not depend on EB1 protein interactions.

Mitotic spindle positioning is tightly controlled during embryogenesis, in

various adult stem cell divisions, and in symmetric divisions (1, 3, 61). While many

studies have focused on the localization of cortical force generators that pull on

microtubule plus ends, attenuators of spindle pulling forces may be just as important

in creating asymmetry. In fact, a variety of molecular mechanisms appear to suppress

spindle pulling forces in the one-cell embryo, including PKC-3-mediated LIN-5

phosphorylation (53), cortical actin (62), and posterior-lateral LET-99 localization

(63). This study provides insight into and a physical basis of spindle pulling force

attenuation: we found that APC acts as an attenuator of spindle pulling forces,

through stabilization of microtubule plus ends at the cortex. Importantly, a similar

force attenuator function of APC is potentially used in oriented divisions of

Drosophila germline stem cells (11), as well as mouse embryonic stem cells (ES

cells) attached to Wnt-immobilized beads (64), as these systems exhibit asymmetric

APC localizations similar to what we have observed in the *C. elegans* zygote. Our

study also implies that not only APC but also other proteins involved in MT

stabilization are potential cortical spindle pulling force attenuators.

The observed pulling force attenuation function may be relevant for the

chromosomal instability (CIN) phenotype associated with APC loss in human colon

cancer (18, 20). Initial studies of cultured mammalian cells associated APC loss and

CIN with defective kinetochore-microtubule attachments, although abnormal spindle

structures were also observed in APC defective cells (18, 20). In *Drosophila* embryos,

APC2 was found to localize predominantly to the cell cortex (65). Chromosome

missegregation associated with APC2 loss in such embryos was linked to a

cytoskeletal function of APC in centrosome segregation (28). In our study, we found

that *C. elegans* APC localizes to the cell cortex where it negatively regulates spindle-

pulling forces. Consequently, the absence of APC results in increased pulling forces

exerted on the spindle poles. Interestingly, defective kinetochore attachments have

been shown to cause chromosome segregation defects in *C. elegans*, in a manner

dependent on cortical pulling forces (66). Thus, combining these data with our results

raises a new and testable hypothesis that increased cortical-pulling forces and

abnormal MT-kinetochore interactions synergistically elevate the risk of CIN in

developing tumors with APC mutations.

1 Materials and methods

2 *C. elegans* culture and strains

All strains used in this study were cultured by standard methods (67). Most worms

were grown at 20 ˚C or 22.5 ˚C and then incubated at 25 ˚C overnight before the

analysis. Worms used for anti-DSH-2 staining were grown at 22.5 ˚C. Worms

carrying PIE-1::GFP were grown at 15˚C and incubated at 25˚C overnight before the

analysis. The following alleles were used: *mom-2(or309), mom-5(ne12), par-2(it51)*.

We used *mom-5(ne12)* null mutants for all *mom-5* experiments except those in Figure

1. The following integrated transgenic lines were used: *osIs15* (32) for GFP::APR-1;

ruIs32 (68) for GFP::H2B; *ojIs1* (69) for GFP::β-tubulin; *axIs1462* (70) for GFP::PIE-

1; *axIs1720* (70) for GFP::PGL-1; *tjIs8* for GFP::EBP-1; *ruIs57* for GFP::tubulin;

ax1928 for mCherry::PAR-6 (71);. We also generated EBP-2::mKate2 fusion strain

ebp-2(*or1954*[*ebp-2*::mKate2]) and *lin-5 (he260[S729A,S734A,S737A,S739A])*strain

by CRISPR/Cas9 genome editing as described below.

16 Generation of CRISPR repair templates

For the generation of the *ebp-2*::mKate2 strain, CRISPR repair constructs containing

700 bp homologous arms were synthesized as gBlock fragments (Integrated DNA

Technologies, Coralville, IA) and assembled into pJET2.1 vector using in-house

Gibson Assembly reaction mix (72). For the generation of the *lin-5 4A* strain,

CRISPR repair constructs were inserted into the pBSK vector using Gibson Assembly

(New England Biolabs, Ipswich, MA). Homologous arms of at least 1500 bp

upstream and downstream of the CRISPR/Cas9 cleavage site were amplified from

cosmid C03G3 using KOD Polymerase (Novagen (Merck) Darmstadt, Germany).

Linkers containing the point mutations were synthesized (Integrated DNA

technologies, Coraville, IA). Mismatches were introduced in the sgRNA target site to

prevent cleavage of the repair template and knock-in alleles. All plasmids and primers

used for this study are available upon request.

CRISPR/ Cas9 genome editing

Young adults were injected with solutions containing the following injection mix. For

ebp-2::mKate2, 10 ng/µl pDD162 P*eft-3*::Cas9 with sgRNA targeting C-terminus of

ebp-2 locus (Addgene 47549; 73), 10 ng/µl repair template, and 65 ng/µl selection

marker pRF4 were used. For *lin-5* 4A, 50 ng/µl P*eft-3*::Cas9 (Addgene 46168; 74), 50

ng/µl of two PU6::sgRNAs targeting the region of the four serine residues to be

mutated to alanine, 50 ng/µl repair template and 2.5 ng/µl selection marker P*myo-*

2::tdTomato were used. Progeny of animals that carry selection markers were

transferred to new plates 3–4 days post injection. For *ebp-2*::mKate2, GFP positive

animals were crossed with a strain carrying GFP:: tubulin to obtain *ebp-2*::mKate2

with GFP::tubulin (EU3068; *ebp-2*(*or1954*[*ebp-2*::mKate2] II). For *lin-5* 4A, PCRs

with primers diagnostic for recombination products at the endogenous locus were

performed on F2-F3 populations, where one primer targeted the altered base pairs in

the sgRNA site and point mutations and the other just outside the homology arm. The

resulting strain (SV1689; *lin-5* (*he260*[S729A/S734A/S737A/S739A]) II) was crossed

with AZ244 (*unc-119(ed3)* III; *ruIs57*) to obtain the *lin-5* 4A strain with GFP::tubulin

(SV1690; *lin-5(he260); ruIs57*).

RNAi

DNA fragments corresponding to nucleotide 848-1547 of the *apr-1* cDNA were

amplified and used for the production of the dsRNA and feeding RNAi. For the

experiments in Figure 5, we injected the dsRNA into the gonad and worms were

subsequently cultured under feeding RNAi at 25 ˚C for over 16 hrs before dissecting

embryos. For the rest of experiments, after injection of the dsRNA into the gonad,

worms were incubated at 25 ˚C without feeding RNAi for over 30 hrs before

dissecting embryos.

Microscopy and analysis of living embryos

All embryos were dissected in an egg salt buffer from gravid hermaphrodites (75). For

live imaging except for the experiments in Figure 5, the embryos were mounted on

4 % agar pads under a coverslip and sealed with petroleum jelly. For most of the

experiments embryos were observed at room temperature by a CSU10 spinning-disc

confocal system (Yokogawa Electric, Musashino, Japan) mounted on an AxioPlan 2

microscope (Carl Zeiss, Oberkochen, Germany) with a Plan-Apochromat 100X 1.4

NA oil immersion lens. The specimens were illuminated with a diode-pumped solid-

state 488 nm laser (HPU50100, 20mW; Furukawa Electric, Tokyo, Japan). Images

were acquired with an Orca ER12-bit cooled CCD camera (Hamamatsu Photonics,

Hamamatsu, Japan), and the acquisition system was controlled by IP lab software (2

X 2 binning; Milwaukee, WI). Acquired images were processed with the Image J (76)

(NIH) and Adobe Photoshop (Adobe Systems, San Jose, CA). For the experiments in

Figure 3B, images were captured with a confocal unit CSU-W with Borealis (Andor

Technology, Belfast, Northern Ireland) and dual EMCCD cameras iXon Ultra 897

(Andor Technology) mounted on an inverted microscope Leica DMi8 (Leica

Microsystems, Wetzlar, Germany) controlled by Metamorph (Molecular Devices,

Sunnyvale, CA). Spindle severing experiments were performed with a Micropoint

system (Photonic instruments, St Charles, IL) equipped with a 2 mW pulsed nitrogen

laser (model VL-337; Laser Science Inc., Franklin, MA) exciting Coumarin 440 dye.

For the experiments in Figure 5, embryos were mounted on 4 % agarose pad

dissolved in egg salts buffer and observed by a Nikon Eclipse Ti microscope with

Perfect Focus System (Nikon, Tokyo, Japan) equipped with CSU-X1-A1 spinning

disk confocal head (Yokogawa Electric) and S Flour 100X 1.3 NA objectives. The

specimens were illuminated with Cobolt Calypso 491 nm laser (Cobolt, Solna,

Sweden). Spindle severing experiments were performed with 355 nm Q-switched

pulsed lasers (Teem Photonics, Meylan, France) with ILas system (Roper Scientific

France, Lisses, France/ PICT-IBiSA, Institut Curie). Temperature was maintained at

25°C by INUBG2E-ZILCS Stage Top Incubator (Tokai Hit, Fujinomiya, Japan) on

the motorized stage MS-2000-XYZ with Piezo Top plate (ASI, Eugene, OR). Images

were acquired with an Evolve 512 EMCCD camera (Photometrics, Tucson, AZ), and

the acquisition system was controlled by MetaMorph (Molecular Devices).

Immunostaining

For the analysis of GFP::APR-1 and mCherry::PAR-6 colocalization, we performed

the freeze-crack method to permeabilize embryos and fixed them in methanol at -

20°C for 5 min followed by acetone at -20°C for 5 min. After washing three times

with PBS supplemented with 1% tween-20, the embryos were incubated with rabbit

polyclonal anti-GFP antibody (1:1000, invitrogen) overnight. After incubation with

goat anti-rabbit Fluorescein (1:1000, Invitrogen), embryos were imaged for

Fluorescein and mCherry signal. Embryos were fixed and stained with rabbit anti-

DSH-2 antibody as described (77).

Measurement of embryo volumes

The volumes (V) of embryos were estimated from the measured embryo length (X)

and width (Y). When three semi-axes of ellipsoid (embryo) in the x, y and z axes are

defined as a, b and c, volume of ellipsoid $V = 4/3\pi abc$. With the assumption of equal

embryo width in the y and z axes, we estimated a, b and c as 0.5X, 0.5Y and 0.5Y and

calculated V.

Statistical analysis

For multiple comparisons, one-way ANOVA with Holm-Sidak's method and Kruskal-

Wallis test followed by Dunn's multiple comparison test were performed for the data

with normal distribution and skewed distribution (judged by F-test), respectively. No

statistical method was used to predetermine sample size. The experiments were not

randomized. The investigators were not blinded.

Quantification of the data from fluorescence images

For the quantification of the number of dots formed by GFP::APR-1, 8 bit images

were processed with Gaussian blur and segmented with the threshold that covers all

the visible dots using Fiji. Then number of segments were counted by the Image J

plug-in Analyze Particles. For the quantification of total APR-1 level in Figure S2A, 4

successive focal planes including cell center and cell surfaces (corresponding to the

upper half of the cell) were combined by the sum projection and average signal

intensity of cell region was subtracted by that in the area devoid of embryos. For the

generation of kymographs that show the centrosome movements along the anterior-

posterior axis, (Figure 2B and 2C, left panels), we drew lines passing through both

centrosomes (some centers are missing due to the transverse movements) and

generated kymographs using Image J function Multi Kymograph. For the generation

of kymographs that show centrosome movements along the transverse axis (Figure 2B

and 2C, right panels), we first adjusted the center of the centrosome manually and

drew a line that passes through the center of the anterior or posterior centrosome and

performed the same procedures. Note that kymographs are composed of linear pixels

of each frame for all time points that together show the centrosome trajectory over

time. For the quantification of spindle movements, the coordinates of the center of the

centrosomes were analyzed with the Image J plug-in Manual Tracking. For the

generation of kymographs of cortical microtubules, (Figure 3A), we extracted and

straightened cortical regions and performed photo-bleach corrections (exponential fit

method) by Image-J. The image color map was changed to mpi-inferno with Image-J.

For the quantification of cortical residence times of GFP::EB1 and GFP::β-tubulin,

the number of frames from appearance to disappearance of each dot were counted

manually. Note that some MT dots whose start and end of cortical localization were

unclear were not counted. The average peak velocity after spindle severing was

calculated from the distance traveled by the centrosome center.

3-dimensional simulation of spindle movement

Overview. The simulations included 2 spindle poles connected by a spring with

dynamic astral MTs inside a cell. The cell was simulated as an oval with a long axis

of 50 μm and two short axes of 30 μm. The initial position of the spindle poles was

set in the center of the cell and aligned along the long axis with the distance of 10μm,

which corresponds to the size of the spindle. The MTs grow and shrink from the

spindle poles stochastically according to the dynamic instability. Depending on the

length and configuration of the MTs, 3 kinds of forces act on spindle poles to move

them as explained below. From an initial configuration, the configuration of the MTs

and the spindle poles was calculated at successive time steps as conducted in previous

simulations (44-47). The parameters used are listed in Table S2.

Force 1, cytoplasmic pulling forces. All MTs generate pulling force proportional to

their length. This force is important to bring the spindle at the cell center (45, 48, 49),

and is also critical for oscillation (38). The cytoplasmic pulling force generated for an

i-th MT was modeled as $F_{cytoplasm}(i) = D \times L(i) \times F_{FG}(i)$, where D is the density of

active force generators in the cytoplasm and $L(i)$ is the length of the MT. $F_{FG}(i)$ is

same as in the cortical pulling force. The direction of the force is the same as the

direction of the MT. We note that the centering force required for oscillation can also

be provided by a force that microtubules produce when they push against the cortex

(78) instead of the cytoplasmic pulling force. The detailed mechanisms (i.e. pulling or

pushing) of the centering force do not affect the overall behavior of our model.

Force 2, cortical pulling forces. MTs that reached the cell cortex generate pulling

forces toward their direction only when they start to shrink. The cortical pulling force

generated for an i-th MT was modeled as $F_{cortex}(i) = N_{potential}(i) \times P_{active}(i) \times F_{FG}(i)$.

$N_{potential}$ is the number of force generators that can potentially interact with the MT.

We set this value at 30 for the posterior cortex and 15 for the anterior cortex. The

experimental value of this parameter has not been investigated, but this number is

consistent with a previous study estimating that the total number of force generators is

less than 50 and the density is double at the posterior cortex compared to anterior

cortex (79). P_{active} is the probability that the potentially interacting force generators are

active. A critical assumption to generate robust oscillation here is to model this value

high when the spindle pole is approaching the site of the force generator, and low

when the spindle pole is leaving (38, 50). In the previous study (38), P_{active} was

defined as $P_{active} = p_{mean} + (f'/f_c) \times p_{mean} \times (1-p_{mean}) \times v - \tau \times (f'/f_c) \times p_{mean} \times (1-p_{mean}) \times a$.

For simplicity, we neglected the acceleration term (a) and fixed the p_{mean} parameter to

0.5 to see the extensive oscillation (38). We set $f'/f_c = 4.0/ V_{max}$, and thus used $P_{active} =$

$0.5 + v/V_{max}$. Here v is the velocity of the spindle pole toward the direction of the force

generator on the cortex. When $v<0$, we set $P_{active} =0$. F_{FG} is formulated as $F_{FG} = F_{stall}$

$(1-v/V_{max})$ (38, 45). When $v>V_{max}$, we set $F_{FG} = 0$. In the simulation, force generation

for shrinking MTs lasts for 100 steps (1 s).

Force 3, forces connecting the two poles. To connect the anterior and posterior

spindle poles, which is done by spindle MTs *in vivo*, we treated the spindle as a

Hookean spring. The natural length increases proportionally from 10 μm at time zero

to 12 μm at $t = 100$ s, which is the onset of anaphase in the simulation. After the onset

of anaphase, the natural length increases proportionally to 22 μm at $t = 200$ s. The

spring constant is high (1 pN/μm) so that the length of spindle is almost maintained to

the natural length.

Acknowledgements

We thank Nancy Hawkins for the anti-DSH-2 antibody, the *Caenorhabditis* Genetics

Center (funded by the NIH Office of Research Infrastructure Programs; P40

OD010440) for strains. This work was supported by the Netherlands Organization for

Scientific Research (NWO) research program 821.02.001 to SvdH, NIH grant

R01GM049869 to B.B., by the Human Frontier Science Program and NIG-JOINT

(2013-A60) to K.S., by the Uehara Memorial Foundation to H.S., and Grants-in-Aid

for Scientific Research from the Ministry of Education, Culture, Sports, Science, and

Technology of Japan to H.S (JP22127005) and A.K. (JP15H04732 and JP15KT0083).

References

1. Siller KH, Doe CQ (2009) Spindle orientation during asymmetric cell division. *Nat Cell Biol* 11(4):365–74.

2. Knoblich JA (2010) Asymmetric cell division: recent developments and their implications for tumour biology. *Nat Rev Mol Cell Biol* 11(12):849–860.

3. Williams SE, Fuchs E (2013) Oriented divisions, fate decisions. *Curr Opin Cell Biol* 25(6):749–758.

4. Rose L, Gönczy P (2014) Polarity establishment, asymmetric division and segregation of fate determinants in early *C. elegans* embryos. *WormBook*, doi: 10.1895/wormbook.1.30.2

5. di Pietro F, Echard A, Morin X (2016) Regulation of mitotic spindle orientation: an integrated view. *EMBO Rep* 17(8):1106–1130.

6. Portegijs V, et al. (2016) Multisite Phosphorylation of NuMA-Related LIN-5 Controls Mitotic Spindle Positioning in *C. elegans*. *PLoS Genet* 12(10). doi:10.1371/journal.pgen.1006291.

7. Clevers H, Nusse R (2012) Wnt/beta-catenin signaling and disease. *Cell* 149(6):1192–1205.

8. Barth AI, Caro-Gonzalez HY, Nelson WJ (2008) Role of adenomatous polyposis coli (APC) and microtubules in directional cell migration and neuronal polarization. *Semin Cell Dev Biol* 19(3):245–251.

9. Etienne-Manneville S (2009) APC in cell migration. *Adv Exp Med Biol* 656:30–40.

10. Pereira G, Yamashita YM (2011) Fly meets yeast: checking the correct orientation of cell division. *Trends Cell Biol* 21(9):526–533.

11.   Yamashita YM, Jones DL, Fuller MT (2003) Orientation of asymmetric stem

cell division by the APC tumor suppressor and centrosome. *Science*

301(5639):1547–50.

12.   Bahmanyar S, Nelson WJ, Barth AI (2009) Role of APC and its binding

partners in regulating microtubules in mitosis. *Adv Exp Med Biol* 656:65–74.

13.   Rusan NM, Peifer M (2008) Original CIN: reviewing roles for APC in

chromosome instability. *J Cell Biol* 181(5):719–26.

14.   Moser AR, Dove WF, Roth KA, Gordon JI (1992) The Min (multiple

intestinal neoplasia) mutation: its effect on gut epithelial cell differentiation

and interaction with a modifier system. *J Cell Biol* 116(6):1517–1526.

15.   Su LK, et al. (1992) Multiple intestinal neoplasia caused by a mutation in the

murine homolog of the APC gene. *Science* 256(5057):668–670.

16.   Beamish H, et al. (2009) Cyclin A/cdk2 regulates adenomatous polyposis coli-

dependent mitotic spindle anchoring. *J Biol Chem* 284(42):29015–23.

17.   Green RA, Wollman R, Kaplan KB (2005) APC and EB1 Function Together

in Mitosis to Regulate Spindle Dynamics and Chromosome Alignment. *Mol*

Biol Cell 16(10):4609–4622.

18.   Fodde R, et al. (2001) Mutations in the APC tumour suppressor gene cause

chromosomal instability. *Nat Cell Biol* 3(4):433–8.

19.   Green RA, Kaplan KB (2003) Chromosome instability in colorectal tumor

cells is associated with defects in microtubule plus-end attachments caused by

a dominant mutation in APC. *J Cell Biol* 163(5):949–961.

20.   Kaplan KB, et al. (2001) A role for the Adenomatous Polyposis Coli protein in

chromosome segregation. *Nat Cell Biol* 3(4):429–32.

21.   Hanahan D, Weinberg RA (2011) Hallmarks of cancer: the next generation.

Cell 144(5):646–674.

22.   Nelson S, Näthke IS (2013) Interactions and functions of the adenomatous

polyposis coli (APC) protein at a glance. *J Cell Sci* 126(Pt 4):873–877.

23.   Hwang E, Kusch J, Barral Y, Huffaker TC (2003) Spindle orientation in

Saccharomyces cerevisiae depends on the transport of microtubule ends along

polarized actin cables. *J Cell Biol* 161(3):483–488.

24.   Korinek WS, Copeland MJ, Chaudhuri A, Chant J (2000) Molecular linkage

underlying microtubule orientation toward cortical sites in yeast. *Science*

287(5461):2257–2259.

25.   Lee L, et al. (2000) Positioning of the mitotic spindle by a cortical-

microtubule capture mechanism. *Science* 287(5461):2260–2262.

26.   Miller RK, Rose MD (1998) Kar9p is a novel cortical protein required for

cytoplasmic microtubule orientation in yeast. *J Cell Biol* 140(2):377–390.

27.   Siller KH, Cabernard C, Doe CQ (2006) The NuMA-related Mud protein

binds Pins and regulates spindle orientation in *Drosophila* neuroblasts. *Nat*

Cell Biol 8(6):594–600.

28.   Poulton JS, Mu FW, Roberts DM, Peifer M (2013) APC2 and Axin promote

mitotic fidelity by facilitating centrosome separation and cytoskeletal

regulation. *Development* 140(20):4226–4236.

29.   Galli M, van den Heuvel S (2008) Determination of the cleavage plane in

early *C. elegans* embryos. *Annu Rev Genet* 42:389–411.

30.   Gönczy P (2008) Mechanisms of asymmetric cell division: flies and worms

pave the way. *Nat Rev Mol Cell Biol* 9(5):355–66.

31.   Mizumoto K, Sawa H (2007) Cortical beta-catenin and APC regulate

asymmetric nuclear beta-catenin localization during asymmetric cell division

in *C. elegans*. *Dev Cell* 12(2):287–99.

32.   Sugioka K, Mizumoto K, Sawa H (2011) Wnt Regulates Spindle Asymmetry

to Generate Asymmetric Nuclear β-Catenin in C. elegans. *Cell* 146(6):942–

954.

33.   Munro E, Bowerman B (2009) Cellular symmetry breaking during

Caenorhabditis elegans development. *Cold Spring Harb Perspect Biol*

1(4):a003400.

34.   Harterink M, et al. (2011) Neuroblast migration along the anteroposterior axis

of *C. elegans* is controlled by opposing gradients of Wnts and a secreted

Frizzled-related protein. *Development* 138(14):2915–2924.

35.   Oldenbroek M, et al. (2013) Regulation of maternal Wnt mRNA translation in

C. elegans embryos. *Development* 140(22):4614–4623.

36.   Walston T, et al. (2004) Multiple Wnt signaling pathways converge to orient

the mitotic spindle in early *C. elegans* embryos. *Dev Cell* 7(6):831–41.

37.   Baldwin AT, Phillips BT (2014) The tumor suppressor APC differentially

regulates multiple β-catenins through the function of axin and CKIα during *C.*

elegans asymmetric stem cell divisions. *J Cell Sci* 127(Pt 12):2771–2781.

38.   Pecreaux J, et al. (2006) Spindle oscillations during asymmetric cell division

require a threshold number of active cortical force generators. *Curr Biol CB*

16(21):2111–22.

39.   Zumbrunn J, Kinoshita K, Hyman AA, Nathke IS (2001) Binding of the

adenomatous polyposis coli protein to microtubules increases microtubule

stability and is regulated by GSK3 beta phosphorylation. *Curr Biol* 11(1):44–

49.

40.   Labbé JC, Maddox PS, Salmon ED, Goldstein B (2003) PAR proteins regulate

microtubule dynamics at the cell cortex in *C. elegans*. *Curr Biol* 13(9):707–

714.

41.   Grill SW, Gönczy P, Stelzer EH, Hyman AA (2001) Polarity controls forces

governing asymmetric spindle positioning in the *Caenorhabditis elegans*

embryo. *Nature* 409(6820):630–3.

42.   Laan L, et al. (2012) Cortical dynein controls microtubule dynamics to

generate pulling forces that position microtubule asters. *Cell* 148(3):502–514.

43. Kozlowski C, Srayko M, Nedelec F (2007) Cortical microtubule contacts

position the spindle in *C. elegans* embryos. *Cell* 129(3):499–510.

44. Hara Y, Kimura A (2009) Cell-size-dependent spindle elongation in the

Caenorhabditis elegans early embryo. *Curr Biol CB* 19(18):1549–54.

45. Kimura A, Onami S (2005) Computer simulations and image processing

reveal length-dependent pulling force as the primary mechanism for *C.*

elegans male pronuclear migration. *Dev Cell* 8(5):765–75.

46. Kimura A, Onami S (2007) Local cortical pulling-force repression switches

centrosomal centration and posterior displacement in *C. elegans*. *J Cell Biol*

179(7):1347–54.

47. Kimura A, Onami S (2010) Modeling microtubule-mediated forces and

centrosome positioning in *Caenorhabditis elegans* embryos. *Methods Cell*

Biol 97:437–453.

48.   Hamaguchi MS, Hiramoto Y (1986) Analysis of the Role of Astral Rays in

Pronuclear Migration in Sand Dollar Eggs by the Colcemid-UV Method. *Dev*

Growth Differ 28(2):143–156.

49.   Kimura K, Kimura A (2011) Intracellular organelles mediate cytoplasmic

pulling force for centrosome centration in the *Caenorhabditis elegans* early

embryo. *Proc Natl Acad Sci U S A* 108(1):137–42.

50.   Grill S, Kruse K, Jülicher F (2005) Theory of mitotic spindle oscillations.

Phys Rev Lett 94(10):1–4.

51.   Vogel SK, Pavin N, Maghelli N, Julicher F, Tolic-Norrelykke IM (2009) Self-

organization of dynein motors generates meiotic nuclear oscillations. *PLoS*

Biol 7(4):e1000087.

52.   Nguyen-Ngoc T, Afshar K, Gönczy P (2007) Coupling of cortical dynein and

G alpha proteins mediates spindle positioning in *Caenorhabditis elegans*. *Nat*

Cell Biol 9(11):1294–302.

53.   Galli M, et al. (2011) aPKC phosphorylates NuMA-related LIN-5 to position

the mitotic spindle during asymmetric division. *Nat Cell Biol* 13(9):1132–

1138.

54.   Etienne-Manneville S, Hall A (2003) Cdc42 regulates GSK-3beta and

adenomatous polyposis coli to control cell polarity. *Nature* 421(6924):753–

756.

55.   Shi SH, Cheng T, Jan LY, Jan YN (2004) APC and GSK-3beta are involved in

mPar3 targeting to the nascent axon and establishment of neuronal polarity.

Curr Biol 14(22):2025–2032.

56. Schlessinger K, McManus EJ, Hall A (2007) Cdc42 and noncanonical Wnt

signal transduction pathways cooperate to promote cell polarity. *J Cell Biol*

178(3):355–61.

57. Davidson G, et al. (2009) Cell cycle control of wnt receptor activation. *Dev*

Cell 17(6):788–99.

58. Gundersen GG, Gomes ER, Wen Y (2004) Cortical control of microtubule

stability and polarization. *Curr Opin Cell Biol* 16(1):106–12.

59. Reilein A, Nelson WJ (2005) APC is a component of an organizing template

for cortical microtubule networks. *Nat Cell Biol* 7(5):463–473.

60. Schmidt R, et al. (2017) Two populations of cytoplasmic dynein contribute to

spindle positioning in *C. elegans* embryos. *J Cell Biol* 216(9):2777–2793.

61. Kiyomitsu T, Cheeseman IM (2012) Chromosome- and spindle-pole-derived

signals generate an intrinsic code for spindle position and orientation. *Nat Cell*

Biol 14(3):311–317.

62. Berends CWH, et al. (2013) F-actin asymmetry and the endoplasmic

reticulum-associated TCC-1 protein contribute to stereotypic spindle

movements in the *Caenorhabditis elegans* embryo. *Mol Biol Cell*

24(14):2201–2215.

63. Krueger LE, Wu J-C, Tsou M-FB, Rose LS (2010) LET-99 inhibits lateral

posterior pulling forces during asymmetric spindle elongation in *C. elegans*

embryos. *J Cell Biol* 189(3):481–495.

64. Habib SJ, et al. (2013) A localized Wnt signal orients asymmetric stem cell

division *in vitro*. *Science* 339(6126):1445–1448.

65. McCartney BM, et al. (2001) *Drosophila* APC2 and Armadillo participate in

tethering mitotic spindles to cortical actin. *Nat Cell Biol* 3(10):933–8.

66.   Cheeseman IM, MacLeod I, Yates JR, Oegema K, Desai A (2005) The CENP-

2   F-like proteins HCP-1 and HCP-2 target CLASP to kinetochores to mediate

3   chromosome segregation. *Curr Biol CB* 15(8):771–7.

4   67.   Brenner S (1974) The genetics of *Caenorhabditis elegans*. *Genetics* 77(1):71–

5   94.

6   68.   Praitis V, Casey E, Collar D, Austin J (2001) Creation of low-copy integrated

7   transgenic lines in *Caenorhabditis elegans*. *Genetics* 157(3):1217–26.

8   69.   Strome S, et al. (2001) Spindle dynamics and the role of gamma-tubulin in

9   early *Caenorhabditis elegans* embryos. *Mol Biol Cell* 12(6):1751–64.

70.   Merritt C, Rasoloson D, Ko D, Seydoux G (2008) 3' UTRs are the primary

regulators of gene expression in the *C. elegans* germline. *Curr Biol*

18(19):1476–1482.

71.   Zonies S, Motegi F, Hao Y, Seydoux G (2010) Symmetry breaking and

polarization of the *C. elegans* zygote by the polarity protein PAR-2.

Development 137(10):1669–1677.

72.   Gibson DG, et al. (2009) Enzymatic assembly of DNA molecules up to

several hundred kilobases. *Nat Methods* 6(5):343–345.

73.   Dickinson DJ, Ward JD, Reiner DJ, Goldstein B (2013) Engineering the

Caenorhabditis elegans genome using Cas9-triggered homologous

recombination. *Nat Methods* 10(10):1028–1034.

74.   Friedland AE, et al. (2013) Heritable genome editing in *C. elegans* via a

CRISPR-Cas9 system. *Nat Methods* 10(8):741–743.

75.   Edgar LG (1995) Blastomere culture and analysis. *Methods Cell Biol* 48:303–

321.

76.   Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25

2   years of image analysis. *Nat Methods* 9(7):671–675.

3   77.   Hawkins NC, Ellis GC, Bowerman B, Garriga G (2005) MOM-5 frizzled

4   regulates the distribution of DSH-2 to control *C. elegans* asymmetric

5   neuroblast divisions. *Dev Biol* 284(1):246–59.

6   78.   Garzon-Coral C, Fantana HA, Howard J (2016) A force-generating machinery

7   maintains the spindle at the cell center during mitosis. *Science*

8   352(6289):1124–1127.

79.   Grill SW, Howard J, Schäffer E, Stelzer EHK, Hyman A a (2003) The

distribution of active force generators controls mitotic spindle position.

Science 301(5632):518–21.

Figure Legends

Figure 1. The Par-aPKC system and Frizzled signaling regulate APR-1

asymmetric localization during zygote division

(A) GFP::APR-1 signals on the cell surface in different mitotic stages. In the right

panels, computationally detected APR-1 dots are shown (see Material and Methods).

(B) GFP::APR-1 and mCherry::PAR-6 localizations in the cell midplane during

asymmetric cell division. GFP signal was amplified by the anti-GFP immunostaining.

Schematic drawing shows polarized protein localizations. (C) GFP::APR-1 signals on

the cell surface in *mom-2(null)* mutants and *mom-5*, *par-2* or *par-3* RNAi embryos.

(D) Quantified numbers of GFP::APR-1 dots on the anterior and posterior cell cortex

of wild-type embryos in different mitotic stages. n = 5, 10, 5 from left to right. (E)

Quantified numbers of GFP::APR-1 dots at metaphase or anaphase in RNAi embryos.

n = 10, 7, 10, 9, 10, 10, from left to right. Ends of whiskers indicate minimum or

maximum values. Double asterisk, asterisk and n.s. indicates: $p < 0.01$, $p < 0.05$ and p

> 0.05 (One-way ANOVA with Holm-Sidak's multiple comparison test). Scale bars

are 10 μm.

Figure 2. APR-1 asymmetrically suppresses centrosome movements during the

P0 cell division

(A) Schematic drawings of spindle movements along the A-P and transverse axes.

Spindle displacement and oscillations contribute mainly to the movements along the

A-P and transverse axes, respectively. Blue arrows and red arrowhead indicate

centrosomes (gamma-tubulin) and chromosomes (Histone H2B), respectively. (B, C)

Centrosome movements in A-P (left panels) and transverse (right panels) axes in

control (B) and *apr-1(RNAi)* ± 40 second around anaphase onset (C). Kymographs

(stack of line images of each time point) were made to show centrosome movements

along the A-P and transverse axes separately. (D, E) Anterior centrosome position

during cell division along the A-P (D) and vertical axes (E). Cell centers are position

zero. (F, H) Total distances for movements of the anterior and posterior poles in

living embryos (F) and in 3D simulations (H). (G) Physical model used for 3D

simulations. A and P indicate the anterior and posterior spindle poles harboring

shrinking MTs (orange) and elongating MTs (blue). Red and black arrows indicate

centrosome movements and cortical force generation. For each MT catastrophe at the

cortex, the average pulling forces acting on a single MT at the posterior are stronger

than those at the anterior, due to the different probabilities of MT-force generator

interactions (see Materials and methods). Error bars show 95% CI. Double asterisk

and n.s. indicates: $p < 0.01$ and $p > 0.05$ compared to control (Kruskal-Wallis test

followed by Dunn's multiple comparison test). Scale bars indicate 5 μm.

Figure 3. APR-1 asymmetrically stabilizes microtubule-cortex interactions

(A) Cortical MT dynamics. Cortical area outlined by the solid line in top figure was extracted, straightened, and corrected for photobleaching. This cortical area, depicted by the dotted line (middle), was used to generate a kymograph (bottom). Color code of the kymograph was changed to highlight MTs. (B) Measurement of cortical MT residence. The embryos were mounted on agarose pads and flattened by coverslips to visualize cortical microtubule ends in a single focal plane. Examples of short and long-lived foci were shown below with simultaneous imaging of GFP::β-tubulin and EB1::mKate2. (C) Cortical microtubule dots in the indicated genotypes during metaphase-anaphase. Images are max projection of cortical GFP::β-tubulin for 60 frames (42 sec). Yellow and Magenta arrows indicate the MT dots whose residence time was shorter and longer than 2.1 sec, respectively. See also Video 3, 5-7. (D) Distribution of quantified cortical MT or EB1 residence time in wild-type animals. (E) Mean cortical MT residence time of indicated genotypes. n = 47, 42, 77, 67, 64, 61, 37, 44, from left to right. Error bars show 95% CI. Double asterisk and asterisk indicate: $p < 0.01$ and $p < 0.05$ compared to control (Kruskal-Wallis test followed by Dunn's multiple comparison test). Scale bars indicate 2.5 μm.

Figure 4 APR-1 asymmetrically attenuates pulling forces acting on the mitotic spindle

(A) Spindle severing experiments. The midzones of mitotic spindles were severed by laser irradiation around anaphase onset (upper left panel). Upon spindle severing, spindle remnants moved at different velocities depending on the net strength of pulling forces (upper right panel). Montages of dissected spindle dynamics were

shown in the bottom panels as DIC images; spindle poles devoid of yolk granules

were indicated by arrowheads. (B) Average peak velocity of spindle poles after

spindle severing. (C) The average of outward pulling forces over 5 sec from anaphase

onset (t = 100 s) for 20 independent simulations. Error bars show 95% CI. Double

asterisk and asterisk indicate: $p < 0.01$ and $p < 0.05$ compared to control (one-way

ANOVA with Holm-Sidak's method). (D) Summary of relationships between cortical

APR-1 level, cortical MT residence, cortical MT catastrophe frequencies, and spindle

pulling forces. Scale bars indicate 10 μm.

Figure 5 Anterior APR-1 enrichment and LIN-5 phosphorylation together

attenuate spindle pulling forces to generate pulling force asymmetry

(A, B) Average peak velocity of spindle poles (A) and their posterior/anterior ratio

(B) after spindle severing. Error bars show 95% CI. Double asterisk and asterisk

indicate: $p < 0.01$ and $p < 0.05$ compared to control (one-way ANOVA with Holm-

Sidak's method). (C) Three elementary processes used in the model described in the

panel C. (1) aPKC-dependent LIN-5 phosphorylation results in the inhibition of force

generation, (2) Cortical MT stabilization by APC reduces the MT catastrophe

frequency and (3) MT shrinkage-dependent force generation is suppressed by step (2).

(D) A schematic model of asymmetric spindle force regulation in P0 cell (see text).

(E) A diagram of spindle pulling force regulation pathways at the anterior cell cortex.

Figure 1

Figure 1. The Par-aPKC system and Frizzled signaling regulate APR-1 asymmetric localization during zygote division.

(A) GFP::APR-1 signals on the cell surface in different mitotic stages. In the right panels, computationally detected APR-1 dots are shown (see Material and Methods). (B) APR-1 and PAR-6 localizations in midplane during asymmetric cell division. GFP signal was amplified by the anti-GFP immunostaining. Schematic drawing show polarized protein localizations. (C) GFP::APR-1 signals on the cell surface in *mom-2(null)* mutants and *mom-5*, *par-2* or *par-3* RNAi embryos. (D) Quantified numbers of GFP::APR-1 dots on anterior and posterior cell cortex of wild-type embryos in different mitotic stages. n = 5, 10, 5 for each sample. (E) Quantified numbers of GFP::APR-1 dots at metaphase or anaphase in RNAi embryos. n = 10, 7, 10, 9, 10, 10, from left to right. Ends of whiskers indicate minimum or maximum values. Double asterisk, asterisk and n.s. indicates: p < 0.01, p < 0.05 and p > 0.05 (One-way ANOVA with Holm-Sidak's multiple comparison test). Scale barsare 10μm.

Figure 2. APR-1 asymmetrically suppresses centrosome movements during the P0 cell division
(A) Schematic drawings of spindle movements along the A-P and transverse axes. Spindle displacement and oscillations contribute mainly to the movements along the A-P and transverse axes, respectively. Blue arrows and red arrowhead indicate centrosomes (gamma-tubulin) and chromosomes (Histone H2B), respectively. (B, C) Centrosome movements in A-P (left panels) and transverse (right panels) axes in control (B) and *apr-1(RNAi)* (C). Kimographs (stack of line images of each time point) were made to show centrosome movements along the A-P and trasnverse axes separately. (D, E) Anterior centrosome position during cell division along the A-P (D) and vertical axes (E). Cell centers are position zero. (F, H) Total distances for movements of the anterior and posterior poles in living embryos (F) and in 3D simulations (H). (G) Physical model used for 3D simulations. A and P indicate the anterior and posterior spindle poles harboring shrinking MTs (orange) and elongating MTs (blue). Red and black arrows indicate centrosome movements and cortical force generation. For each MT catastrophe at the cortex, the average pulling forces acting on a single MT at the posterior are stronger than those at the anterior, due to the different probabilities of MT-force generator interactions (see Materials and methods). Error bars show 95% CI. Double asterisk and n.s. indicates: $p < 0.01$ and $p > 0.05$ compared to control (Kruskal-Wallis test followed by Dunn's multiple comparison test). Scale bars indicate 5 μm.

Figure 3

Figure 3. APR-1 asymmetrically stabilizes microtubule-cortex interactions.
(A) Cortical MT dynamics. Cortical area outlined by the solid line in top figure was extracted, straightened, and corrected for photobleaching. This cortical area, depicted by the dotted line (middle), was used to generate a kymograph (bottom). Color code of the kymograph was changed to highlight MTs. (B) Measurement of cortical MT residence. The embryos were mounted on agarose pads and flattened by coverslips to visualize cortical microtubule ends in a single focal plane. Examples of short and long-lived foci were shown below with simultaneous imaging of GFP::β-tubulin and EB1::mKate2. (C) Cortical microtubule dots in the indicated genotypes during metaphase-anaphase. Images are max projection of cortical GFP::β-tubulin for 60 frames (42 sec). Yellow and Magenta arrows indicate the MT dots whose residence time was shorter and longer than 2.1 sec, respectively. See also Video 3, 5-7. (D) Distribution of quantified cortical MT or EB1 residence time in wild-type animals. (E) Mean cortical MT residence time of indicated genotypes. n = 47, 42, 77, 67, 64, 61, 37, 44, from left to right. Error bars show 95% CI. Double asterisk and asterisk indicate: p < 0.01 and p < 0.05 compared to control (Kruskal-Wallis test followed by Dunn's multiple comparison test). Scale bars indicate 2.5 μm.

Figure 4 APR-1 asymmetrically attenuates pulling forces acting on the mitotic spindle.
(A) Spindle severing experiments. The midzones of mitotic spindles were severed by laser irradiation around anaphase onset (upper left panel). Upon spindle severing, spindle remnants moved at different velocities depending on the net strength of pulling forces (upper right panel). Montages of dissected spindle dynamics were shown in the bottom panels as DIC images; spindle poles devoid of yolk granules were indicated by arrowheads. (B) Average peak velocity of spindle poles after spindle severing. (C) The average of outward pulling forces over 5 sec from anaphase onset (t = 100 s) for 20 independent simulations. Error bars show 95% CI. Double asterisk and asterisk indicate: p < 0.01 and p < 0.05 compared to control (one-way ANOVA with Holm-Sidak's method). (D) Summary of relationships between cortical APR-1 level, cortical MT residence, cortical MT catastrophe frequencies, and spindle pulling forces. Scale bars indicate 10 μm.

Figure 5 Anterior APR-1 enrichment and LIN-5 phosphorylation together attenuate spindle pulling forces to generate pulling force asymmetry.

(A, B) Average peak velocity of spindle poles (A) and their posterior/anterior ratio (B) after spindle severing. Error bars show 95% CI. Double asterisk and asterisk indicate: $p < 0.01$ and $p < 0.05$ compared to control (one-way ANOVA with Holm-Sidak's method). (C) Three elementary processes used in the model described in the panel C. (1) aPKC-dependent LIN-5 phosphorylation results in the inhibition of force generation, (2) Cortical MT stabilization by APC reduces the MT catastrophe frequency and (3) MT shrinkage-dependent force generation is suppressed by step (2). (D) A schematic model of asymmetric spindle force regulation in P0 cell (see text). (E) A diagram of spindle pulling force regulation pathways at the anterior cell cortex.

Figure S1. Temporal and genetic regulation of cortical GFP::APR-1 localization
(**A, C**) Quantified numbers of GFP::APR-1 dots on the anterior and posterior cell cortex are shown for prophase, metaphase, anaphase and telophase of control and RNAi embryos. (B) APR-1 dots in the indicated RNAi experiments. Left and right panels are original and computationally segmented binary images, respectively. Ends of whiskers indicate minimum to maximum values. Double asterisk, asterisk and n.s. indicates: p < 0.01, p < 0.05 and p > 0.05 (One-way ANOVA with Holm-Sidak's multiple comparison test). Scale bars indicate 10 μm.

Figure S2. Roles of Wnt signaling in P0 cell division.
(A) APR-1 level after RNAi experiments. GFP fluorescence intensity per area of the whole embryo including the cell cortex and cytoplasm were measured and shown. Signal in wild-type indicates auto-fluorescence. (B) Immunofluorescence images of the DSH-2 protein during P0 and EMS cell division. Blue is DAPI staining. In EMS, the DSH-2 protein is enriched at the cell boundary between EMS and P2 (arrowheads) while no asymmetry was observed in P0. (C) Localizations of the cell fate determinant GFP::PIE-1 in the indicated genotypes. Control and *apr-1(RNAi)* shows PIE-1 enrichment in the posterior blastomere P1. In the *par-2* mutant, PIE-1 asymmetry was lost. Scale bars indicate 10 μm.

Figure S3. Effects of *mom-5(RNAi)* on spindle pole movements and embryo sizes.
(A) Kymographs of the spindle movements in *mom-5(RNAi)*. Kymographs were generated as in Figure 2.
(B) Distance traveled by the anterior or posterior spindle poles. Total distance centrosome traveled for ± 40 sec and ± 100 sec from anaphase onset were shown for *in vivo* measurements (left) and 3D simulations (right). Error bars show 95% CI. Asterisk indicate p < 0.05 compared to control (Kruskal-Wallis test followed by Dunn's multiple comparison test).

Figure S4. Estimation of microtubule catastrophe frequencies at the cortex.
(A) Frequencies of MT residence times at the cell cortex observed experimentally (histograms) and predicted from the estimated catastrophe frequencies (black lines). (B) Estimated catastrophe frequencies for indicated genotypes. The data is the same as in Supplementary Table 1.

Figure S5. Numerical simulation of spindle movements
(A-D) Representative trajectories of the spindle poles in the simulation. The trajectories of the anterior (green) and posterior (red) poles are shown. Their midpoint (black) is also shown in (A and D). (A, B) Control condition. (C, D) *apr-1(RNAi)* condition. (A and D) Trajectories along A-P axis (x axis). (C and E) and those along an axis perpendicular to the x axis (y axis) are shown.

Table 1

Estimated catastrophe frequencies of the microtubules at the cortex

	anterior cortex	posterior cortex
control (N2)	0.31 (/s)	0.72 (/s)
apr-1	0.53 (/s)	0.59 (/s)
mom-5	0.20 (/s)	0.44 (/s)
apr-1; mom-5	0.58 (/s)	0.51 (/s)

When catastrophe occurs stochastically with the frequency of λ, the probability distribution of the cortical residency time will be P(t) = λexp(-λt). Therefore, the probability of observing cortical residency time between t1 and t2 will be P(t1~t2) = exp(-λt1) - exp(-λt2). We fitted the experimentally obtained probability distribution of the cortical MT residency time to this equation to estimate the catastrophe frequencies of the MTs at the cortex

Table 2. Parameter values used in the simulation

		References
Microtubule (MT) dynamics		
Growth velocity (Vg) [μm/s]	0.328	(Srayko et al., 2005)
Shrinkage velocity (Vs) [μm/s]	0.537	(Kozlowski et al., 2007)
Catastrophe frequency (Fcat) at cytoplasm [/s][a]	0.046	(Kozlowski et al., 2007)
Rescue frequency (Fres) [/s][b]	1	
Number of fibers per pole	296	(Srayko et al., 2005)
Pulling force, motor mediated		
Stall force of motor (F_{stall}) [pN]	1.1	(Gross et al., 2000)
Maximum velocity of motor (V_{max}) [μm/s]	2.0	(Gross et al., 2000)
Pulling force, attachment of FG (cytoplasmic length dependent)		
Density of motors (D) [/μm]	0.2	
Pulling force, attachment of FG (cortical)		
Potential number of force generators at the cortex ($N_{potential}$, anterior, PAR-3 dependent)	15	
Potential number of force generators at the cortex ($N_{potential}$, posterior, PAR-2 dependent)	30	
The mean probability of the activation of the force generators (p_{mean}) [/s]	0.5	(Pecreaux et al., 2006)
Spindle as a spring		
Natural length [μm]	10-22	
Spring constant [pN/μm]	1	
Size of the cell		
Long axis [μm]	50	
Short axis [μm]	30	
Drag force of nucleus/spindle pole		
Drag coefficient, for translational movement (Γ_{trans}) [pN s/μm][c]	190	
Drag coefficient, for rotational movement (Γ_{rot}) [pN s μm][c]	25,000	
Model-specific parameter		
Time step [s]	0.01	

[a] See Table S1 for catastrophe frequency at the cortex

[b] A high frequency was used in this study. See text for a detailed explanation.

[c] $6\pi r\eta$ for translational movement and $8\pi r^3\eta$ for rotational movement. Here, we set r (Stokes' radius) to 10 μm and η (viscosity of the cytosol) to 1.0 pNs/μm^2.