# Widely used commercial ELISA does not detect preHP-2, but recognizes properdin as a potential second member of the zonulin family

Running Title: Commercial ELISA for human zonulin detects also properdin

Lucas Scheffler[1], Alyce Crane[1], Henrike Heyne[1], Anke Tönjes[2], Dorit Schleinitz[1], Christian

H. Ihling[3], Michael Stumvoll[2], Rachel Freire[4], Maria Fiorentino[4],

Alessio Fasano[4], Peter Kovacs[1*], John T. Heiker[5*]

[1]Leipzig University Medical Center, IFB Adiposity Diseases, Leipzig, Germany; [2]Divisions of Endocrinology and Nephrology, University of Leipzig, Leipzig, Germany; [3]Department of Pharmaceutical Chemistry and Bioanalytics, Institute of Pharmacy, Martin-Luther-University Halle-Wittenberg, Halle, Germany; [4]Mucosal Immunology And Biology Research Center, Massachusetts General Hospital – Harvard Medical School, Boston, MA U.S.A.; [5]Institute of Biochemistry, Faculty of Biosciences, Pharmacy and Psychology, University of Leipzig, Leipzig, Germany

**\*Corresponding authors:**

E-mail: jheiker@uni-leipzig.de

E-mail: peter.kovacs@medizin.uni-leipzig.de

**Keywords:** diabetes ‐ lipid metabolism ‐ pre-Haptoglobin 2 ‐ ELISA ‐ intestinal permeability ‐ properdin – obesity – metabolic diseases

**Nonstandard abbreviations**: HP, haptoglobin; T2D, type 2 diabetes; WHR, waist-to-hip ratio.

**Abstract**

**BACKGROUND.** There is increasing evidence for the role of impaired intestinal permeability in obesity and associated metabolic diseases. Zonulin is an established serum marker for intestinal permeability and identical to pre-haptoglobin2. Here, we aimed to investigate the relationship between circulating zonulin and metabolic traits related to obesity.

**METHODS.** Serum zonulin was measured by using a widely used commercial ELISA kit in 376 subjects from the metabolically well-characterized cohort of Sorbs from Germany. In addition, haptoglobin genotype was determined in DNA samples from all study subjects.

**RESULTS.** As zonulin concentrations did not correlate to the haptoglobin genotypes, we investigated the specificity of the zonulin ELISA assay using antibody capture experiments, mass spectrometry and Western blot analysis. Using serum samples that gave the highest or lowest ELISA signals, we detected several proteins that are likely to be captured by the antibody in the present kit. However, none of these proteins corresponds to pre-haptoglobin2. We used increasing concentrations of recombinant pre-haptoglobin 2 and complement C3 as one of the representative captured proteins and the ELISA kit did not detect either. Western blot analysis using both the polyclonal antibodies used in this kit and monoclonal antibodies rose against zonulin showed a similar protein recognition pattern but with different intensity of detection. The protein(s) measured using the ELISA kit was (were) significantly increased in patients with diabetes and obesity and correlated strongly with markers of the lipid and glucose metabolism. Combining mass spectrometry and Western blot analysis using the polyclonal antibodies used in the ELISA kit, we identified properdin as another member of the zonulin family.

**CONCLUSIONS.** Our study suggests that the zonulin ELISA does not recognize pre-haptoglobin 2, rather structural (and possibly functional) analogue proteins belonging to the mannose-associated serine protease family, with properdin being the most likely possible candidate.

## Introduction

The "intestinal barrier" is an established term, defined as a functional entity separating the gut lumen from the inner host, and consisting of mechanical, humoral, immunological, muscular and neurological elements. Intestinal barrier dysfunction is a characteristic feature of pathological states such as inflammatory bowel disease, celiac disease, nonalcoholic steatohepatitis and ulcerative colitis (1, 2). There is also emerging evidence for the role of impaired intestinal permeability in metabolic diseases including obesity and type 2 diabetes (T2D) (3–5). It has been hypothesized that gut bacteria and bacterial endotoxins may disrupt the intestinal barrier resulting in the so called "leaky gut" (4, 6). The leakage of toxins, bacterial components or even live bacteria and their transfer to target organs such as adipose tissue might contribute to the development of obesity and T2D (6, 7). Indeed, numerous studies in mouse models have demonstrated that changes in the gut microbiota can alter the gut permeability and lead to an endotoxemia-induced inflammation in adipose tissue, and ultimately to obesity (3, 8, 9). Results from experimental mouse models are supported by studies in humans by showing an increase in circulating endotoxin levels and circulating bacterial DNA in obese/diabetic patients, likely due to an increased intestinal permeability in affected subjects (10, 11).

Intestinal barrier transport is mainly regulated by structures of the paracellular pathway called tight junctions which form barriers between epithelial cells and regulate the transport of ions and small molecules across the intestinal lumen. Intestinal permeability is a functional feature of the intestinal barrier. It can be measured by analyzing flux rates of inert molecules across the intestinal wall as a whole or across wall components (1). The gold standard for assessment of intestinal permeability in vivo is an assay combining indigestible large and small oligosaccharides, such as lactulose and mannitol; the larger oligosaccharide, lactulose, is only transported via the paracellular pathway, whereas the smaller oligosaccharide, mannitol, is taken up freely over the intestinal barrier via the transcellular route. However, these

oligosaccharide assays are expensive, laborious, poorly reproducible, and time consuming.

Therefore, identifying appropriate biomarkers for intestinal permeability is highly desirable.

Zonulin has been identified as a tight junction regulating protein which is, functionally, the

human counterpart of the Vibrio cholera endotoxin zonula occludens toxin (12, 13). Precisely,

subsequent studies recognized zonulin as the precursor of haptoglobin 2 (pre-HP2) (14).

Haptoglobin is a well-known protein involved in scavenging hemoglobin, whereas the

function of its precursor is largely unknown. Haptoglobin is first synthesized into a single

chain precursor protein, which is cleaved into a light N-terminal α-chain and heavy C-

terminal β-chain. An exon duplication of exons 3 and 4 of the haptoglobin gene differentiates

the HP1 from the HP2 allele. Due to this exon duplication the HP2 α-chain is 1.7kb longer

than in the HP1 allele. Haptoglobin is active as tetramer consisting of 2 α- and 2 β-chains

linked by disulfide bonds, resulting in three possible genotypes: homozygous HP1/1 and

HP2/2 as well as heterozygous HP1/2 (15, 16). About 15 % of the Caucasian population is

homozygous for HP (16, 17). Zonulin as pre-HP2 reversibly opens tight junctions and is

upregulated in diseases such as celiac disease and type 1 diabetes (T1D) (14, 18). Serum

zonulin concentrations are also increased in T2D and obesity (19–21) and strong correlations

were observed with various metabolic markers, including fasting plasma glucose, IL-6, HDL,

and triglyceride (TG) levels (19–21).

Here, we aimed at characterizing the relationship between circulating serum zonulin and traits

related to obesity in a metabolically well-characterized cohort of Sorbs from Germany. To

measure zonulin, we used the commercially available ELISA kit (Immundiagnostik,

Bensheim, Germany). In addition, we determined the haptoglobin genotypes in the entire

cohort. Due to a lack of correspondence between the observed circulating zonulin

concentrations and the haptoglobin genotypes in our study cohort, we further investigated the

possible identity of the product captured by the commercial ELISA assay. We found that the

ELISA kit used in the present study does not detect purified pre-HP2 but rather targets one or

more proteins from a range of candidate molecules possibly structurally and functionally

related to zonulin. Our data also showed that protein concentrations measured by this ELISA

correlated with parameters of obesity and related metabolic traits.

**Materials and Methods**

*Study subjects*

All subjects are part of a sample from an extensively clinically characterized population from

Eastern Germany, the Sorbs (22–24). Extensive phenotyping included standardized

questionnaires to assess past medical history and family history, collection of anthropometric

data (weight, height, waist-to-hip ratio (WHR)), and an oral glucose tolerance test. Glucose

was assessed by the Hexokinase method (Automated Analyzer Modular, Roche Diagnostics,

Mannheim, Germany) and serum insulin was measured using the AutoDELFIA Insulin assay

(PerkinElmer Life and Analytical Sciences, Turku, Finland). Total serum cholesterol and TG

concentrations were measured by standard enzymatic methods (CHOD-PAP and GPO-PAP;

Roche Diagnostics). Serum LDL-C and HDL-C concentrations were determined using

commercial homogeneous direct measurement methods (Roche Diagnostics). All assays were

performed in an automated clinical chemistry analyzer (Hitachi/ Roche Diagnostics) at the

Institute of Laboratory Medicine, University Hospital Leipzig.

All blood samples were taken in the morning after an overnight fast and stored at -80°C until

analyses. From the 1040 Sorbs enrolled in the cohort, a subgroup of 376 subjects was

genotyped for haptoglobin and provided blood samples for zonulin measurements (Table 1).

Main metabolic characteristics of the study subjects are summarized in Table 2.

The study was approved by the ethics committee of the University of Leipzig and all subjects

gave written informed consent before taking part in the study.

*ELISA measurements*

Circulating zonulin was measured by a competitive ELISA (Immundiagnostik AG, Bensheim, Germany) in serum samples of 376 individuals according to the manufacturer's protocol. Inter-assay coefficient of variation was 6.5%. When purified proteins zonulin and properdin and the synthetic peptide AT1001 were tested by the ELISA, they were re-suspended in PBS and diluted in Diluent buffer (IDK kit) to reach a final concentration of 5 μg/ml.

*Genotyping*

Haptoglobin genotypes were determined by PCR using a method adapted from Koch et al. (17). Briefly, the following two primer pairs were used: A (5'-GAGGGGAGCTTGCCTTTCCATTG-3') and B (5'-GAGATTTTTGAGCCCTGGCTGGT-3'), as well as C (5'-CCTGCCTCGTATTAACTGCACC AT-3') and D (5'-CCGAGTGCTCCACATAGCCATGT-3'). The primer pair A/B generates two bands: a 1,757-bp allele 1 specific band and a 3,481-bp allele 2 specific band. The primer pair C/D produces one allele 2 specific band of 349 bp. The combination of the bands allows a reproducible typing of the two common haptoglobin genotypes HP1 and HP2. In contrast, no band is detectable for the rare HP deletion genotype, present in ~0.1% Caucasians (16).

*Antibody capturing experiment*

We aimed to isolate the target protein of the ELISA antibody from serum samples utilizing the immobilized anti-zonulin antibodies on the ELISA plates to perform antibody capturing experiments. Based to the manufacturing information, these polyclonal antibodies were raised against an octapeptide sequenced from the zonulin molecule (25). Equal amounts of undiluted serum samples with highest and lowest concentrations of zonulin, as measured using the same ELISA kit, were transferred to separate wells, incubated and washed according to the manufacturer's protocol. Afterwards we eluted the captured protein(s) by incubation with 50

µl of hot (95°C) SDS sample buffer with β-mercaptoethanol for 5 minutes. The captured

protein of high or low zonulin serum samples were pooled (N=8 for high (2 pooled groups)

and low (1 pooled group); measured protein concentrations using the zonulin ELISA of the

serum samples is given in Supplementary Table 1). Twenty µl of these elution samples (high

or low zonulin) were separated by SDS-PAGE using precast Bolt 4-12% Bis-Tris Plus gels

(ThermoFisher, Waltham, MA, USA). Proteins were stained using the Pierce silver stain for

mass spectrometry (ThermoFisher) or detected by Western Blot.

*Mass spectrometry*

To identify proteins isolated from serum samples as described above, bands were excised

from silver-stained gels and in-gel digested with trypsin following a standard protocol (26).

After enzymatic digestion, the peptide mixtures were immediately analyzed by LC/MS/MS on

an U3000 RSLC nano-HPLC system (Thermo Fisher Scientific) coupled to an Orbitrap

Fusion Tribrid mass spectrometer (Thermo Fisher Scientific). Samples were loaded onto a

pre-column (RP-C8, 300 µm * 5 mm, 5 µm, 100 Å, Thermo Fisher Scientific) and washed

with water containing 0.1% (v/v) TFA for 15 min, before the peptides were separated on the

separation column (RP-C18, 75 µm * 250 mm, 2 µm, 100 Å, Thermo Fisher Scientific) using

gradients from 1% to 40% (v/v) B (45 min), 40% to 85% (v/v) B (5 min) followed by 85% B

(5 min), with solvent A: 0.1% (v/v) formic acid (FA) in water and solvent B: 0.08% (v/v) FA

in acetonitrile. Data were acquired using data-dependent MS/MS mode where each high-

resolution full-scan in the orbitrap (m/z 198 to 1,500; R = 120,000) was followed by high-

resolution product ion scans in the orbitrap (higher energy collision-induced dissociation

(HCD), 27% normalized collision energy, R = 15,000, isolation window 2 Th) within 5 s,

starting with the most intense signal in the full-scan mass spectrum.

Data analysis was performed using the Proteome Discoverer 2.1 (Thermo Fisher Scientific).

MS/MS data of precursor ions (m/z range 350-5,000) were searched against the Swissprot

Database (version 11/2016, taxonomy human, 20,082 entries) and a contaminant database

using Sequest HT. Mass accuracy was set to 5 ppm and 20 mmu for precursor and fragment

ions, respectively. Carbamidomethylation of cysteines was set as fixed modification;

oxidation of methionines and N-terminal acetylation were set as variable modifications, two

missed cleavages of trypsin were allowed. Results were filtered for non-contaminant proteins

identified by at least three unique highly confident peptides (peptide FDR ≤ 1%).

*Western blot analysis*

Western blot experiments were performed to validate the results of mass spectrometric data

analysis and to compare the serum target proteins identified by the polyclonal antibodies used

by this ELISA kit compared to monoclonal anti-zonulin antibodies. Gels were blotted on a

PVDF membrane and Western blots were probed with anti-C3 β-chain (1:2000) (Biozol,

Eching, Germany), anti-haptoglobin (1:1000) (Abcam, Cambridge, UK), polyclonal anti-

zonulin (1:500) (kindly provided by Immundiagnostik), monoclonal anti-zonulin (1:5000)

(BioRad, Hercules, CA USA) antibodies. Purified C3c from plasma (Athens Research,

Athens, Georgia, USA) and recombinant zonulin were used as positive controls. Properdin

(R&D Systems Minneapolis, MN, U.S.A.) was also used to validate the potential protein

candidate identified by the ELISA.  Incubation with primary and secondary antibodies (HRP-

conjugated) was done for 90 minutes at room temperature. Blots were visualized by enhanced

chemiluminescence using Pierce ECL Western Blotting Substrate (ThermoFisher).

*Statistical analysis*

Statistical analysis was performed using with SPSS 24 (IBM). All non-normally distributed

metric parameters were log transformed to generate a Gaussian normal distribution.

Spearman's rank correlation method was used to assess the relationship between metabolic

traits. To test for significant differences in distribution for ordinal values, the Kruskal-Wallis-

test was used. Exact differences between two groups were tested by both the Mann‑Whitney‑

U test and unpaired student's t‑test. In addition, multiple linear regression analyses were done

to assess the linear relationship between continuous variables and genotypes. For all tests, a p-

value <0.05 was considered to be statistically significant.

**Results**

*Haptoglobin genotype*

Haptoglobin genotype HP1/1 was present in 15.8% of the subjects, HP1/2 in 47.6% and

HP2/2 in 36.6%. These frequencies are comparable to the distribution of HP genotypes in

cohorts of European ancestry reported by others (16, 17). We tested the association of the HP

genotypes with various anthropometric and metabolic traits in all study subjects. The analysis

revealed that blood hemoglobin levels significantly increase with the presence of at least one

HP2 allele (p=0.004 over all three groups, p=$4.2x10^{-4}$ between HP1 homozygote and HP2

carriers). Furthermore, we observed that the total protein concentration in the urine

significantly differed between the three groups, with an increase in the HP2 carriers

(p=0.027). Interestingly, mean triiodothyronine (fT3) levels were lower in the HP1/1 group

than in the HP2/2 group (p=0.012) and in accordance, an increase in administered thyroid

gland hormones (p=0.023) was observed.

*Zonulin ELISA data do not match HP genotype distribution*

Strikingly, there were no significant differences in levels of the zonulin ELISA signal

between the three haptoglobin genotype groups (Figure 1; p=0.153 using ANOVA, p=0.07 for

the t-test comparing log transformed zonulin signals between HP1/1 vs. HP1/2 + HP2/2).

Assuming that the protein measured by the kit is zonulin (i.e. pre-HP2), subjects with the

HP1/1 genotype were expected to have no detectable zonulin levels. As the zonulin

concentrations measured in patient sera using the zonulin ELISA kit clearly did not reflect the

HP genotype distribution we therefore concluded that the protein measured by the kit is not

identical to zonulin as pre-HP2 or that, beside pre-HP2, the kit detects other unrelated

proteins. Consequently, we aimed at identifying the protein(s) detected by the alleged zonulin

ELISA kit.

*The zonulin ELISA does not detect recombinant pre-HP2, but targets multiple proteins*

To attempt identifying proteins bound and quantified by the capturing antibody deployed in

the zonulin ELISA kit, we performed an immune-capturing experiment using patient sera and

the immobilized antibody of the ELISA kit as supplied. After incubation of the immobilized

ELISA kit antibodies with selected patient sera representing the highest and lowest measured

ELISA signals in the cohort, the captured proteins were separated by SDS-PAGE. Notably,

we could not measure the protein content of the eluted samples, but given that equal amounts

of serum were used, that the same washing and elution procedure was performed for all

samples and that 20µl of the pooled elution samples were used for SDS-PAGE and Western

blot, the detected amount of "captured" protein should resemble the amount of protein that

was present in the initial serum sample.

Silver staining revealed multiple bands, with the most intense band at ~70 kDa and further

prominent bands at ~55 kDa, ~180 kDa and >180kDA (Figure 2A). This band pattern was

incompatible with a band pattern that would be expected for pre-HP2 or haptoglobin-derived

proteins and further supported the results demonstrating the lack of correspondence of the

captured protein with HP genotypes. To further characterize major proteins captured by the

ELISA kit, protein bands were cut (Figure 2A) and subjected to MS analysis after tryptic

digestion (Supplementary Table 2). Mass spectrometry demonstrated that bands 1 (>180kDa),

2 (~150 kDa), 3 (~70 kDa) were all very likely representing the C3 protein or cleavage

products derived from the C3 protein, such as the C3 precursor (187 kDa), C3c (144 kDa) and

the C3 β-chain (71 kDa). Furthermore, the 55 kDa band was identified as properdin or factor

P (MW 53 kDa). To validate results from mass spectrometry, we performed Western blot

analyses. The major band at 70 kDa was clearly detected by an anti-C3 β-chain antibody

(Figure 2B).

Consequently, we tested several C3c proteins from different suppliers (Abcam, Cambridge,

UK; Athens Research, Athens, Georgia, USA; mybiosource, San Diego, California, USA), in

a range from 0.1 mg/ml to 0.1 ng/ml under native and denatured conditions, as well as diluted

in serum, with the respective ELISA kit. All results were negative (data not shown),

indicating that C3 might represent a contaminant only. Additionally, we tested C3,

recombinant zonulin, HP1, and HP2 at increasing concentrations (range 1-50 μg/ml), along

with sera from celiac patients (both HP1-1 and HP2-2), healthy controls (both HP1-1 and

HP2-2), and our standard control (AF, HP2-2). The results showed in Figure 3 demonstrated

that while this kit does not recognize C3, zonulin, or mature haptoglobin (both HP1 and HP2),

it does recognize protein(s) both in HP1-1 and HP2-2 genotype subjects, irrespective of their

disease status.

**Anti-zonulin monoclonal antibodies and Immundiagnostik polyclonal antibodies**

**recognize same target proteins in Western blot.**

Our ELISA results clearly showed that this kit does not detect zonulin (alias, pre-HP2 as

originally claimed by the manufacturer) or complement C3, the protein captured by our

experiments described above. In order to further characterize the target protein(s) of this kit,

we performed Western blotting analysis of sera from a HP2-2 subject either under baseline

condition or after deglycosylation as we have previously described (14), using recombinant

zonulin as control. As anticipated, the zonulin monoclonal antibodies recognized recombinant

zonulin as well as a variety of bands in the serum sample, including bands with a MW of ~70,

52, 37, 27, and 16 kDa (Figure 4A). Based on similar patterns we detected when zonulin was

originally cloned (14), we predicted that the ~72 kDa corresponded to the glycosylated HP β-

chain, the 52 kDa zonulin, and the 16 kDa the HP2 α-chain. To confirm this interpretation, we

performed deglycosylation experiments showing the shift of the 72 kDa β-chain to a lower

MW, while, as anticipated, the zonulin band and the HP2 α-chain remained unchanged

(Figure 4A). Interestingly enough, the Immundiagnostik polyclonal antibodies raised against

the zonulin synthetic peptide inhibitor GGVLVQPG (AT1001) (25) recognized the same main

bands detected by the monoclonal antibodies but with different intensity, being the

recombinant zonulin and serum α-chain bands fainter compared to the monoclonal antibody

signal, while the serum β-chain and serum zonulin-like signals stronger (Figure 4A). When

combined to our ELISA results, these data suggest that, while the zonulin monoclonal

antibodies specifically detect in serum samples only zonulin at its predicted molecular weight,

most likely the polyclonal antibodies used in this kit are possibly detecting "zonulin-like"

protein(s) (as suggested by the much more intense signal of the ~52 kDa band compared to

the monoclonal antibodies) with similar molecular weight, structure, and possibly function.

**Immundiagnostik polyclonal antibodies recognize both zonulin as well as properdin as**
**an additional target in Western blot**

Among all the proteins we identified with our mass spec analysis, properdin fulfills the

structural-functional characteristics mentioned above and, therefore, may represent the most

likely candidate target detected by this ELISA kit.  To explore this hypothesis, we performed

both Western blotting analysis using the Immunodiagnostik polyclonal antibodies and ELISA

test using commercially available properdin. The Western blot showed that beside

recombinant zonulin, the Immunodiagnostik polyclonal antibodies also detect properdin (Fig

4B) that migrated at the same molecular weight of zonulin and serum bands recognized by the

antibodies in both HP2-2 and HP 1-1 subjects (Fig 4B). Similar results were obtained with

polyclonal antibodies raised against recombinant zonulin (data not shown). The same samples

used for Western blotting analysis, including zonulin, properdin and AT1001 at a

concentration of 5 μg/ml were tested using the ELISA kit. The results showed that properdin

and AT1001 were both detected by the ELISA test, however, their amounts were substantially

underestimated by 914 (5.47 ng/ml) and 40 folds (126.04 ng/ml), respectively. However,

despite zonulin being detected in Western blot, it was not detected with the ELISA kit (Figure

3). In comparison, serum from subjects with HP 1-1 and HP 2-2 genotype were measured at a

concentration of 67.54 and 56.41 ng/ml, respectively (Table 3).

*Correlations of measured protein concentrations using the ELISA with metabolic traits*

In a sample of 376 subjects tested using the purchased zonulin ELISA, the product was

measured in a mean concentration of $72.2 \pm 27.2$ ng/ml (mean $\pm$ standard deviation), ranging

from 17.8 to 207.1 ng/ml. The ELISA signal was significantly increased in subjects with T2D

$(81.78 \pm 25.31$ ng/ml) compared to subjects with normal glucose tolerance $(67.25 \pm 25.45$

ng/ml, Mann-Whitney-U test; $p=2.1 \times 10^{-8}$) or impaired glucose tolerance $(71.88 \pm 29.36$

ng/ml, $p=0.0017$) (Figure 5A). Additionally, lean subjects had significantly lower values

$(65.64 \pm 25.23$ ng/ml) than subjects with overweight $(74.20 \pm 30.68$ ng/ml, $p=0.0082$) or

obesity $(76.24 \pm 24.17$ ng/ml, $p=7.0 \times 10^{-5}$) (Figure 5B). We observed no gender differences or

any correlations with age (data not shown). The ELISA signal correlated with traits related to

glucose and lipid metabolism (Spearman's rank correlation test, adjusted for age, sex and

BMI; Table 4). It was positively correlated with BMI, HOMA-IR and -IS and fasting plasma

glucose (Table 4). Strong correlations were also observed for lipid metabolism parameters,

such as TG levels, total cholesterol, LDL and apolipoprotein B (Table 4). Correlations were

tested for a total of 95 accessible traits. After Bonferroni correction for multiple testing

(adjusted p-value for significance $p < 5.2 \times 10^{-4}$), correlations for BMI $(p=1.2 \times 10^{-5})$, fasting

glucose $(p=9.0 \times 10^{-6})$, TG $(p=4.2 \times 10^{-10})$, total cholesterol $(p=3.3 \times 10^{-5})$ and apolipoprotein B

$(p=2.3 \times 10^{-5})$ remained statistically significant.

**Discussion**

The role of impaired intestinal permeability in metabolic diseases including obesity and T2D

has recently been acknowledged in multiple studies (3, 4, 27). The tight junction regulator

zonulin, which was identified as pre-HP2 by Tripathi et al. (14), is an established circulating

marker of intestinal permeability in humans (28, 29). Here, we aimed to investigate the

relationship between intestinal permeability, represented by circulating zonulin, and metabolic

traits related to obesity and T2D. We set out to measure zonulin with a widely used

commercially available ELISA kit in a metabolically well-characterized cohort of Sorbs from

Germany. Considering the identity of zonulin as pre-HP2, we also genotyped the haptoglobin

gene in DNA samples from all subjects. The genotype frequency of the HP1/1 genotype

corresponded to previously reported frequencies of 15% in populations of European ancestry

(16, 17). Assuming that the measured zonulin is identical with pre-HP2 as reported previously

(13), we expected subjects with the HP1/1 genotype to have no detectable, or if taking into

account cross-reactions, at least significantly lower zonulin levels. However, concentrations

measured in the HP1/1 homozygous subjects were comparable with those found in HP2/2 and

HP1/2 carriers.

The epitope used to generate the capture antibody in the zonulin ELISA is based on the

sequence GGVLVQPG published by Wang et al. (25) (communication with customer

support; Immundiagnostik AG, Bensheim, Germany), which was initially thought to represent

the N-terminal sequence of fetal zonulin (25). However, this sequence is not present in pre-

HP2, which has since been proposed to be zonulin by the same authors (14). According to the

authors, the discrepancies between the previously reported zonulin sequence and the pre-HP2

sequence may be due to intraspecies variability associated with a high zonulin mutation rate

or due to sequencing errors at that time (14). In summary, the commercially available

competitive ELISA does very likely not detect preHP2 or zonulin, but rather unknown targets

bound by the antibody raised against the sequence reported by Wang et al (25).

Based on database searches, the epitope could correspond to Glu-Rich2, a protein which

shares 7 out of 8 amino acids. The zonulin/preHP2 binding partner CD163 shows some

conformity with the epitope (30). Our antibody capture experiment and subsequent mass

spectrometry analysis did not provide any evidence for either protein. The most abundant

protein identified by MS, C3, is evidently an unspecific product overshadowing the real

targets. Indeed, the respective ELISA kit did not detect any complement C proteins obtained

from different suppliers when tested under native and denatured conditions, as well as diluted

in serum. Also, the same kit did not detect recombinant zonulin, mature HP1 or mature HP2.

Considering the additional MS hits, a few proteins stand out, although, without further

validation, we interpret these data with caution, since only more abundant proteins may be

identified by MS analysis, while our protein(s) of interest may be in low abundance in serum

samples and, therefore, not identifiable with this approach. The van Willebrand factor (vWF,

band A, Figure 2) is involved in the intrinsic coagulation pathway and the acute phase

response and known to be increased in inflammatory bowel disease and bacterial diarrhea

(31). Inter-alpha-inhibitor heavy chain 4 (band B, Figure 2) a large glycoportein cleaved into

smaller fragments by Kallikrein which is also involved in the intrinsic coagulation pathway.

One of these fragments, called urinary trypsin-inhibitor, attenuates LPS-induced endothelial

barrier dysfunction by upregulation of vascular endothelial-cadherin expression (32).

Complement component 9 (C9, band C, Figure 2) is an important component of the

membrane attacking complex within the complement cascade and is required for complement-

mediated lipopolysaccharide release and outer membrane damage in bacteria (33). Protein

S100-A8, or Calprotectin, (band D, Figure 2) is a calcium- and zinc-binding protein which

plays a prominent role in the regulation of inflammatory processes and immune response (34).

However, based on the data presented in Figure 4, our most likely candidate(s) should be in

the ~50 kDa range, were the polyclonal antibody signal was stronger. Therefore, the most

interesting candidate protein we have identified is properdin or factor P (band D, Figure 2), a

member of the complement alternative pathway that has a molecular weight (53 kDa), within

the range of those proteins recognized by the Immundiagnostik anti-AT1001 polyclonal

antibodies (Figures 4A and B) and serum levels (~25 ng/ml) (35) similar to the range of

detection of the kit. Our combined Western blot analysis (Figure 4B) and ELISA test

confirmed that the polyclonal antibodies raised against AT1001 detect properdin amongst

other proteins. However, when purified proteins/peptides, including the AT1001 peptide used

to raise the polyclonal antibodies, which is also used as internal control in the ELISA kit,

were tested by ELISA, they were highly under-estimated by the test. One possible explanation

for these results is that zonulin and also properdin are not the main targets detected by the

ELISA, however the fact that even AT1001 was under-estimated seems to suggest that this

hypothesis cannot entirely explain our results.  Alternatively, it is possible that tertiary and

quaternary (multimers) structure arrangements present in sera samples but not in recombinant

proteins are necessary in order to properly detect both zonulin and properdin by this ELISA.

If this is the case and/or that the main target of this ELISA is/are additional proteins in the

~50kDa range present in human serum remains to be established.

Once released from neutrophils, T cells and macrophages in response to acute microbial

exposure, properdin causes production of chemotactic anaphylatoxin C3a and C5a with

subsequent formation of immune complexes that cause increased endothelial permeability

(35). Intriguingly, zonulin as pre-HP2 also causes generation of C3a and C5a, with

subsequent increased vascular permeability in several districts, including the lung, with

subsequent onset of acute lung injury (36). Another striking similarity between zonulin and

properdin is the fact that both are associated to viral respiratory tract infections (37, 38).

Notably, the peptide sequence used for the generation of the antibody is also not present in

any of the discussed proteins. The five proteins mentioned above, including properdin, have

~50% similarity to this epitope. Zonulin as preHP-2 is a member of a larger family of tight

junction regulating proteins.  Indeed, phylogenetic analyses suggest that haptoglobins evolved

from mannose-associated serine protease (MASP), a complement-associated protein (like

properdin), with their alpha-chain containing a complement control protein (CCP) (this

domain activates complement similarly to properdin), while the β-chain is related to

chymotrypsin-like serine proteases (SP domain) (39, 40). However, the SP domain of HP

lacks the essential catalytic amino acid residues required for protease function; structure-

function analyses have implicated this domain in receptor recognition and binding (41).

Although not a serine protease, zonulin shares approximately 19% amino acid sequence

homology with chymotrypsin, and their genes both map on chromosome 16. Alignment of the

β-chain sequence of zonulin to that of several serine proteases is remarkably consistent except

for an insertion of 16 residues in the region corresponding to the methionyl loop of the serine

proteases. Comparison of the zonulin α-β junction region with the heavy-light-chain junction

of tissue-type plasminogen activator strengthens the evolutionary homology of zonulin and

serine proteases. The active-site residues typical of the serine proteases, His57 and Ser195,

are replaced in zonulin by lysine and alanine, respectively. Because of these mutations, during

evolution zonulin most likely lost its protease activity despite that zonulin and serine

proteases evolved from a common ancestor (18). Therefore, zonulin, and the serine proteases

represent a striking example of homologous proteins with different biological functions but

with the common characteristic of complement activation. Beside zonulin and properdin,

other members of the MASP family include a series of plasminogen-related growth factors

(epidermal growth factor (EGF), hepatocyte growth factor (HGF), etc.) involved in cell

growth, proliferation, differentiation and migration, and disruption of intercellular junctions.

In light of these considerations, other MASP members identified in our capturing experiments

in general, and properdin in particular, are intriguing possible targets (36).

Analyzing the protein concentrations measured using this ELISA in subjects who have been

extensively characterized for metabolic phenotypes, our data suggest that it is upregulated

both in diabetic and obese patients. This is in line with previously reported findings using this

ELISA kit (19–21). Our data shows, that the ELISA target is potentially involved in the lipid

metabolism by showing in various linear stepwise regression models that triglyceride levels

and fasting glucose are the strongest independent available variables explaining the observed

variance in measured protein concentrations (Supplementary Table 3).

It is important to note, that our study, as it has not been initially designed to address the

question of ELISA specificity, has clear limitations. To obtain amounts of isolated protein in

the antibody capture experiments, we needed to change the experimental protocol provided by

the manufacturer. We used undiluted serum samples instead of a 50-fold dilution, which very

likely increased the risk of non-specific binding. Yet, we used sera from patient that exhibited

the highest and the lowest concentrations measured by the ELISA kit using the

manufacturer's protocol. Thus, any non-specific binding should be detected as equally strong

bands in the silver stained gels after the antibody capture experiment. Yet, we observed band

intensities of affinity purified protein that clearly correlated with the concentrations measured

using the ELISA, in the silver stained gels (total protein) and in the Western blot using the

anti-C3-β-chain antibody, indicating a specific isolation of proteins recognized by the kit

antibody. Nevertheless, ELISA results that failed to recognize C3 disputed the notion that

complement C3 is the target of this kit. Also, we have performed this experiment using two

different Lot no. of the ELISA, using sera from eight different patients of each high and low

concentration and obtained the same results.

In conclusion, based on our data we suggest that the Immundiagnostik ELISA kit supposedly

testing serum zonulin (pre-HP2) levels could identify a variety of proteins structurally and

possibly functionally related to zonulin, suggesting the existence of a family of zonulin

proteins as previously hypothesized (42), rather than a single member of permeability-

regulating proteins. Additional studies are necessary to establish the primary target proteins

(zonulin, properdin and/or other structurally similar proteins) detected by this commercially

available ELISA.

**Acknowledgements**

We thank all those who participated in the studies, in particular our study subjects. We thank

Dr. Ingo Bechmann for helpful advice and discussions.

**Funding**

This work was supported by grants from the German Research Council (SFB-1052 ''Obesity

mechanisms'' B03, C01, C07), from the German Diabetes Association and from the DHFD

(Diabetes Hilfs- und Forschungsfonds Deutschland). IFB Adiposity Diseases is supported by

the Federal Ministry of Education and Research (BMBF), Germany, FKZ: 01EO1501 (AD2-

060E, AD2-6E99), and by the National Institutes of Health (NIH) U.S.A. grants R01-

DK104344 and P30-DK040461.

**Author contributions**

LS, PK and JTH conceived the study, designed and conducted experiments, analyzed data and

wrote the paper. AT recruited patients. CHI performed mass spectrometry experiments. AC,

HH, DS and MS interpreted and analyzed data. RF performed the zonulin ELISA test, MF

performed the Western blotting analysis, and AF critically revised the manuscript, contributed

to the study design of some of the performed experiments, and provided critical interpretation

of the data. All authors discussed results, edited and commented on the manuscript. All the

authors have accepted responsibility for the entire content of this submitted manuscript and

approved submission.

**Disclosure statement:** The authors have no conflicts of interest to declare.

**References**

1.  Bischoff SC, Barbara G, Buurman W, Ockhuizen T, Schulzke J-D, Serino M, et al. Intestinal

permeability – a new target for disease prevention and therapy. *BMC gastroenterology* (2014)

14:189. doi:10.1186/s12876-014-0189-7.

2.  Arrieta MC, Bistritz L, Meddings JB. Alterations in intestinal permeability. *Gut* (2006) 55:1512–20.

doi:10.1136/gut.2005.085373.

3.  Cani PD, Bibiloni R, Knauf C, Waget A, Neyrinck AM, Delzenne NM, et al. Changes in gut

microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity

and diabetes in mice. *Diabetes* (2008) 57:1470–81. doi:10.2337/db07-1403.

4.  Teixeira TF, Collado MC, Ferreira, Celia L L F, Bressan J, Peluzio, Maria do Carmo G. Potential

mechanisms for the emerging link between obesity and increased intestinal permeability. *Nutrition*

*research (New York, N.Y.)* (2012) 32:637–47. doi:10.1016/j.nutres.2012.07.003.

5.  Frazier TH, DiBaise JK, McClain CJ. Gut microbiota, intestinal permeability, obesity-induced

inflammation, and liver injury. *JPEN. Journal of parenteral and enteral nutrition* (2011) 35:14–20.

doi:10.1177/0148607111413772.

6.  Harris K, Kassis A, Major G, Chou CJ. Is the gut microbiota a new factor contributing to obesity

and its metabolic disorders? *Journal of obesity* (2012) 2012:879151. doi:10.1155/2012/879151.

7.  Burcelin R, Serino M, Chabo C, Garidou L, Pomie C, Courtney M, et al. Metagenome and

metabolism: the tissue microbiota hypothesis. *Diabetes, obesity & metabolism* (2013) 15 Suppl

3:61–70. doi:10.1111/dom.12157.

8.  Cani PD, Possemiers S, Van de Wiele, T, Guiot Y, Everard A, Rottier O, et al. Changes in gut

microbiota control inflammation in obese mice through a mechanism involving GLP-2-driven

improvement of gut permeability. *Gut* (2009) 58:1091–103. doi:10.1136/gut.2008.165886.

9.  Turnbaugh PJ, Backhed F, Fulton L, Gordon JI. Diet-induced obesity is linked to marked but

reversible alterations in the mouse distal gut microbiome. *Cell host & microbe* (2008) 3:213–23.

doi:10.1016/j.chom.2008.02.015.

10. Ortiz S, Zapater P, Estrada JL, Enriquez P, Rey M, Abad A, et al. Bacterial DNA translocation

holds increased insulin resistance and systemic inflammatory levels in morbid obese patients. *The*

*Journal of clinical endocrinology and metabolism* (2014) 99:2575–83. doi:10.1210/jc.2013-4483.

11. Sato J, Kanazawa A, Ikeda F, Yoshihara T, Goto H, Abe H, et al. Gut dysbiosis and detection of
"live gut bacteria" in blood of Japanese patients with type 2 diabetes. *Diabetes care* (2014)
37:2343–50. doi:10.2337/dc13-2817.

12. Fasano A, Baudry B, Pumplin DW, Wasserman SS, Tall BD, Ketley JM, et al. Vibrio cholerae
produces a second enterotoxin, which affects intestinal tight junctions. *Proceedings of the*
*National Academy of Sciences of the United States of America* (1991) 88:5242–6.

13. Fasano A, Fiorentini C, Donelli G, Uzzau S, Kaper JB, Margaretten K, et al. Zonula occludens
toxin modulates tight junctions through protein kinase C-dependent actin reorganization, in vitro.
*The Journal of clinical investigation* (1995) 96:710–20. doi:10.1172/JCI118114.

14. Tripathi A, Lammers KM, Goldblum S, Shea-Donohue T, Netzel-Arnett S, Buzza MS, et al.
Identification of human zonulin, a physiological modulator of tight junctions, as prehaptoglobin-2.
*Proceedings of the National Academy of Sciences of the United States of America* (2009)
106:16799–804. doi:10.1073/pnas.0906773106.

15. Wobeto, VÃ\textcentnia Peretti de Albuquerque, Zaccariotto TR, Sonati Md. Polymorphism of
human haptoglobin and its clinical importance. *Genetics and Molecular Biology* 31:602–20.

16. Langlois MR, Delanghe JR. Biological and clinical significance of haptoglobin polymorphism in
humans. *Clinical chemistry* (1996) 42:1589–600.

17. Koch W, Latz W, Eichinger M, Roguin A, Levy AP, Schomig A, et al. Genotyping of the common
haptoglobin Hp 1/2 polymorphism based on PCR. *Clinical chemistry* (2002) 48:1377–82.

18. Fasano A. Zonulin and its regulation of intestinal barrier function: the biological door to
inflammation, autoimmunity, and cancer. *Physiological reviews* (2011) 91:151–75.
doi:10.1152/physrev.00003.2008.

19. Zhang D, Zhang L, Zheng Y, Yue F, Russell RD, Zeng Y. Circulating zonulin levels in newly
diagnosed Chinese type 2 diabetes patients. *Diabetes research and clinical practice* (2014)
106:312–8. doi:10.1016/j.diabres.2014.08.017.

20. Żak-Gołąb A, Kocełak P, Aptekorz M, Zientara M, Juszczyk Ł, Martirosian G, et al. Gut
Microbiota, Microinflammation, Metabolic Profile, and Zonulin Concentration in Obese and Normal
Weight Subjects. *International Journal of Endocrinology* (2013) 2013:9. doi:10.1155/2013/674106.

21. Moreno-Navarrete JM, Sabater M, Ortega F, Ricart W, Fernandez-Real JM. Circulating zonulin, a
marker of intestinal permeability, is increased in association with obesity-associated insulin
resistance. *PloS one* (2012) 7:e37160. doi:10.1371/journal.pone.0037160.

22. Tonjes A, Zeggini E, Kovacs P, Bottcher Y, Schleinitz D, Dietrich K, et al. Association of FTO
variants with BMI and fat mass in the self-contained population of Sorbs in Germany. *European*
*journal of human genetics EJHG* (2010) 18:104–10. doi:10.1038/ejhg.2009.107.

23. Tonjes A, Koriath M, Schleinitz D, Dietrich K, Bottcher Y, Rayner NW, et al. Genetic variation in
GPR133 is associated with height: genome wide association study in the self-contained
population of Sorbs. *Human molecular genetics* (2009) 18:4662–8. doi:10.1093/hmg/ddp423.

24. Veeramah KR, Tönjes A, Kovacs P, Gross A, Wegmann D, Geary P, et al. Genetic variation in the
Sorbs of eastern Germany in the context of broader European genetic diversity. *European Journal*
*of Human Genetics* (2011) 19:995–1001. doi:10.1038/ejhg.2011.65.

25. Wang W, Uzzau S, Goldblum SE, Fasano A. Human zonulin, a potential modulator of intestinal
tight junctions. *Journal of cell science* (2000) 113 Pt 24:4435–40.

26. Shevchenko A, Tomas H, Havlis J, Olsen JV, Mann M. In-gel digestion for mass spectrometric
characterization of proteins and proteomes. *Nature protocols* (2006) 1:2856–60.
doi:10.1038/nprot.2006.468.

27. Damms-Machado A, Louis S, Schnitzer A, Volynets V, Rings A, Basrai M, et al. Gut permeability
is related to body weight, fatty liver disease, and insulin resistance in obese individuals
undergoing weight reduction. *The American journal of clinical nutrition* (2017) 105:127–35.
doi:10.3945/ajcn.116.131110.

28. Sapone A, Magistris L de, Pietzak M, Clemente MG, Tripathi A, Cucca F, et al. Zonulin
upregulation is associated with increased gut permeability in subjects with type 1 diabetes and
their relatives. *Diabetes* (2006) 55:1443–9.

29. Drago S, El Asmar R, Di Pierro M, Grazia Clemente M, Tripathi A, Sapone A, et al. Gliadin,
zonulin and gut permeability: Effects on celiac and non-celiac intestinal mucosa and intestinal cell
lines. *Scandinavian    journal    of    gastroenterology*    (2006)    41:408–19.
doi:10.1080/00365520500235334.

30. Thomsen JH, Etzerodt A, Svendsen P, Moestrup SK. The Haptoglobin-CD163-Heme Oxygenase-
1 Pathway for Hemoglobin Scavenging. *Oxidative Medicine and Cellular Longevity* (2013)
2013:11. doi:10.1155/2013/523652.

31. Sadler JE. Biochemistry and Genetics of von Willebrand Factor. *Annual Review of Biochemistry*
(1998) 67:395–424. doi:10.1146/annurev.biochem.67.1.395.

32. Chen J, Wang J, Su C, Qian W, Sun L, Sun H, et al. Urinary trypsin inhibitor attenuates LPS-

induced endothelial barrier dysfunction by upregulation of vascular endothelial-cadherin

expression. *Inflammation Research* (2016) 65:213–24. doi:10.1007/s00011-015-0907-9.

33. Tegla CA, Cudrici C, Patel S, Trippe R, Rus V, Niculescu F, et al. MEMBRANE ATTACK BY

COMPLEMENT: THE ASSEMBLY AND BIOLOGY OF TERMINAL COMPLEMENT

COMPLEXES. *Immunologic research* (2011) 51:45–60. doi:10.1007/s12026-011-8239-5.

34. Yui S, Nakatani Y, Mikami M. Calprotectin (S100A8/S100A9), an inflammatory protein complex

from neutrophils with a broad apoptosis-inducing activity. *Biological & pharmaceutical bulletin*

(2003) 26:753–60.

35. Kouser L, Abdul-Aziz M, Nayak A, Stover CM, Sim RB, Kishore U. Properdin and Factor H:

Opposing Players on the Alternative Complement Pathway "See-Saw". *Frontiers in Immunology*

(2013) 4:93. doi:10.3389/fimmu.2013.00093.

36. Rittirsch D, Flierl MA, Nadeau BA, Day DE, Huber-Lang MS, Grailer JJ, et al. Zonulin as

prehaptoglobin2 regulates lung permeability and activates the complement system. *American

journal of physiology. Lung cellular and molecular physiology* (2013) 304:72.

doi:10.1152/ajplung.00196.2012.

37. Shirey KA, Lai W, Patel MC, Pletneva LM, Pang C, Kurt-Jones E, et al. Novel strategies for

targeting innate immune responses to influenza. *Mucosal immunology* (2016) 9:1173–82.

doi:10.1038/mi.2015.141.

38. Ahout IM, Brand KH, Zomer A, van den Hurk WH, Schilders G, Brouwer ML, et al. Prospective

observational study in two Dutch hospitals to assess the performance of inflammatory plasma

markers to determine disease severity of viral respiratory tract infections in children. *BMJ open*

(2017) 7:e014596. doi:10.1136/bmjopen-2016-014596.

39. Wicher KB, Fries E. Haptoglobin, a hemoglobin-binding plasma protein, is present in bony fish

and mammals but not in frog and chicken. *Proceedings of the National Academy of Sciences of

the United States of America* (2005) 103:4168–73. doi:10.1073/pnas.0508723103.

40. Kurosky A, Barnett DR, Lee TH, Touchstone B, Hay RE, Arnott MS, et al. Covalent structure of

human haptoglobin: A serine protease homolog. *Proceedings of the National Academy of

Sciences of the United States of America* (1980) 77:3388–92.

41. Nielsen MJ, Petersen SV, Jacobsen C, Thirup S, Enghild JJ, Graversen JH, et al. A unique loop

extension in the serine protease domain of haptoglobin is essential for CD163 recognition of the

haptoglobin-hemoglobin complex. *The Journal of biological chemistry* (2007) 282:1072–9.

doi:10.1074/jbc.M605684200.

42. Fasano A. Regulation of intercellular tight junctions by zonula occludens toxin and its eukaryotic

analogue zonulin. *Annals of the New York Academy of Sciences* (2000) 915:214–22.

## Tables

**Table 1: Composition of the study cohort included in zonulin measurements.**

|  | *all* | *female* | *male* |
|---|---|---|---|
| *n* | 376 | 233 | 143 |
| *Age (years)* | 53.3 ± 16.2 | 52.7 ± 15.8 | 54.2 ± 16.8 |
| *BMI (kg/m²)* | 28.1 ± 5.6 | 27.9 ±5.9 | 28.5 ± 5.1 |
| *NGT* | 191 (50.8%) | 124 (53.2%) | 67 (46.8%) |
| *IGT* | 79 (21.0%) | 49 (21.0%) | 30 (21.0%) |
| *T2D* | 106 (28.2 %) | 60 (25.8%) | 46 (32.2%) |

Age and BMI are mean ± SD, NGT, IGT, T2D as absolute number (percentage); NGT: normal glucose tolerance; IGT: impaired glucose tolerance; IFG: impaired fasting glucose

**Table 2: Main characteristics of the study participants.**

|  | *NGT* | *IGT* | *T2D* |
|---|---|---|---|
| *n* | 191 | 79 | 106 |
| *Age (years)* | 45.4 ± 15.6 | 59.7 ± 29.4** | 62.9 ± 11.2** |
| *BMI (kg/m²)* | 25.8 ± 5 | 29.7 ± 4.6** | 30.97 ± 5.53** |
| *WHR* | 0.84 ± 0.10 | 0.92 ± 0.09 | 0.94 ± 0.09** |
| *FPG (mmol/l)* | 5.15 ± 0.50 | 5.68 ± 0.56** | 7.72 ± 2.43** °° |
| *120 min PG (mmol/l)* | 5.00 ± 1.40 | 8.95 ± 0.98** | 5.78 ± 6.86 °° |
| *Triglycerides (mmol/l)* | 1.29 ± 0.94 | 1.56 ± 0.92* | 1.87 ± 1.24** |
| *Fasting Insulin (pmol/l)* | 35.90 ± 25.98 | 53.56 ± 25.98** | 62.38 ± 43.07** |
| *120 min Insulin (pmol/l)* | 148.39 ± 133.58 | 409.28 ± 232.81** | 182.22 ± 272.34 °° |
| *HDL (mmol/l)* | 1.69 ± 0.40 | 1.56 ± 0.36* | 1.43 ± 0.39** ° |
| *LDL (mmol/l)* | 3.20 ± 0.96 | 3.70 ± 1.06** | 3.33 ± 0.95 ° |
| *Total cholesterol (mmol/l)* | 5.12 ± 1.03 | 5.57 ± 1.18* | 5.27 ± 0.99 |
| *HOMA IR (pmol/l*mmol/l)* | 1.34 ± 1.06 | 2.30 ± 1.1** | 3.58 ± 2.87** °° |
| *HOMA IS (ratio)* | 1.07 ± 0.76 | 0.67 ± 0.78** | 0.68 ± 1.63** |
| *CRP (mg/l)* | 2.22 ± 4.20 | 3.19 ±4.54 | 3.87 ± 4.76** |
| *Zonulin (ng/ml)* | 67.25 ± 25.45 | 71.88 ± 29.36 | 81.78 ± 25.31** °° |

*: p<0.05 vs NGT; **: p<0.01 vs NGT
°: p<0.05 vs IGT; °°: p<0.01 vs IGT
All data given as mean ± standard deviation, NGT: normal glucose tolerance, IGT: impaired glucose tolerance, T2D: type 2 diabetes, CRP: C-reactive protein, HDL: High density lipoprotein, LDL: Low density lipoprotein, FPG: Fasting plasma glucose, WHR: Waist to hip ratio

**Table 3: Correlation of ELISA signal with metabolic phenotypes.**

|  | *Non-adjusted* |  | *Adjusted for age, sex, BMI* |  |
|---|---|---|---|---|
|  | r | p-value | r | p-value |
| *Anthropometric trait* |  |  |  |  |
| *Age* | 0.31 | 0.55 | - | - |
| *Sex* | -0.005 | 0.93 | - | - |
| *BMI* | **0.221** | **$1.2 \times 10^{-5}$** | - | - |
| *WHR* | **0.172** | **$7.3 \times 10^{-4}$** | **0.118** | **$2.1 \times 10^{-2}$** |
| *Glucose metabolism* |  |  |  |  |
| *Glucose (0 min)* | **0.294** | **$6.6 \times 10^{-9}$** | **0.227** | **$9.0 \times 10^{-6}$** |
| *Glucose (120 min)* | -0.039 | 0.45 | -0.046 | 0.38 |
| *Insulin (0 min)* | **0.173** | **$7.7 \times 10^{-4}$** | 0.079 | 0.13 |
| *Insulin (120 min)* | -0.039 | 0.46 | -0.044 | 0.39 |
| *HOMA IR* | **0.244** | **$2.0 \times 10^{-6}$** | **0.149** | **$4.0 \times 10^{-3}$** |

| | | | | |
|---|---|---|---|---|
| *HOMA IS* | **-0.243** | **2.0x10$^{-6}$** | **-0.145** | **4.9 x10$^{-3}$** |
| *Lipid metabolism* | | | | |
| *Triglycerides* | <u>**0.370**</u> | <u>**6.5x10$^{-14}$**</u> | <u>**0.312**</u> | <u>**4.2 x10$^{-10}$**</u> |
| *Total cholesterol* | <u>**0.219**</u> | <u>**1.5x10$^{-5}$**</u> | <u>**0.211**</u> | <u>**3.3x10$^{-5}$**</u> |
| *LDL* | **0.182** | **3.4x10$^{-4}$** | **0,160** | **1.7 x10$^{-3}$** |
| *HDL* | **-0.136** | **7.7x10$^{-3}$** | -0.063 | 0.22 |
| *Apolipoprotein B* | <u>**0.247**</u> | <u>**9.9x10$^{-7}$**</u> | <u>**0.215**</u> | <u>**2.3x10$^{-5}$**</u> |
| *Adipokines* | | | | |
| *Adiponectin* | **-0.215** | **2.2x10$^{-6}$** | **-0.176** | **5.4x10$^{-4}$** |
| *Progranulin* | **0.151** | **3.0x10$^{-3}$** | **0.129** | **1.1 x10$^{-2}$** |
| *Vaspin* | 0.027 | 0.6 | 0.05 | 0.33 |
| *Chemerin* | **0.103** | **4.3x10$^{-2}$** | 0.065 | 0.21 |
| *FGF21* | **0.165** | **2.1x10$^{-3}$** | **0.152** | **4.7x10$^{-3}$** |
| *Other* | | | | |
| *C-reactive protein* | **0.232** | **4.0x10$^{-6}$** | **0.166** | **1.1x10$^{-3}$** |
| *Total protein* | **0.124** | **1.5x10$^{-2}$** | **0,134** | **8.6x10$^{-3}$** |
| *Hemoglobin* | **0.201** | **7.4x10$^{-5}$** | **0.143** | **5.1x10$^{-3}$** |
| *Uric acid* | **0.176** | **5.2x10$^{-4}$** | **0.110** | **3.1 x10$^{-2}$** |

639 **bold**: significant correlations after adjustment, <u>underlined:</u> significant after Bonferroni
640 adjustment for multiple testing ($p < 5.2x10^{-4)}$; r: Spearman rank correlation coefficient, p:
641 significance level; BMI: Body Mass Index; HDL: High density lipoprotein, LDL: Low
642 density lipoprotein, WHR: Waist to hip ratio, FGF21: Fibroblast growth factor 21, HOMA-
643 IR/IS: Homeostasis Model Assessment Insulin Resistance/ Insulin Sensitivity.
644

     **Figure legends**

647
**Fig 1. Distribution of zonulin ELISA values according to haptoglobin genotypes.**

Data is presented as boxplots with Turkey-Whiskers and outliers.

**Fig 2. A) Silver stain of immunoprecipitated ELISA products and B) Western blot**
**analyses using an anti-C3 β-chain antibody.**

Commercially available C3c protein isolated from human plasma was run as positive control.
high: pooled IP samples of sera that gave highest ELISA signals; low: pooled IP samples of
sera that gave lowest ELISA signals; neg: negative control using dilution buffer from the
ELISA kit; tracer: competitive tracer reagent from the ELISA kit. Letters in (A) indicated
bands analyzed by MS after tryptic in gel digestion (Supplementary Table 2).

Fig 3. Mean zonulin ELISA values obtained by testing specific candidate target proteins and patients' serum samples. This ELISA kit did not detect increasing concentrations (range 1-50 μg/ml) of complement, recombinant zonulin (pre-HP2), mature HP1 or mature HP2. Conversely, a strong signal was detected in sera of both celiac disease patients (CD) and healthy controls (non-CD, internal control AF with HP2-2), irrespective of their HP genotype. All samples are technical triplicates, kit controls are shown for comparison.

670
671

Fig 4. Western blot analyses of a prototype human serum sample (AF HP2-2) using both anti-zonulin monoclonal antibodies and Immundiagnostik polyclonal antibodies. (A) Prototype serum sample of a HP2-2 homozygous subject, either untreated (lane 1) or after deglycosylation (lane 2) was resolved and then immunoblotted using either anti-zonulin monoclonal antibodies (left panel) or Immundiagnostik polyclonal antibodies (right panel). Recombinant zonulin was added as control (lane 3). As anticipated, zonulin monoclonal antibodies recognize the recombinant protein as well as a serious of serum bands, with the strongest signal being related to the 16kDa zonulin α2-chain. The Immundiagnostik polyclonal antibodies also recognize recombinant zonulin but with a much weaker signal

compared to the monoclonal antibodies. These antibodies also recognize the same serum
bands detected by the zonulin monoclonal antibodies, with the ~70 and 52 kDa bands being
highlighted with the strongest signal. Sample's deglycosyltation showed the shift of the 70
kDa band to a lower MW, suggesting that this may represent the zonulin β-chain as we have
previously shown (14). (B) Prototype serum samples of a HP1-1 and a HP2-2 homozygous
subject were resolved and immunoblotted using the Immundiagnostik polyclonal antibodies.
Recombinant zonulin and properdin were added as control. The antibodies also detect
properdin that migrated at the same molecular weight of zonulin and serum bands recognized
by the antibodies in both HP2-2 and HP 1-1 subjects.

**Fig 5. A) Mean zonulin ELISA values according to glucose tolerance groups B) mean**
**ELISA values according to BMI groups.** \*\*: $p<0.01$; \*\*\*: $p<0.001$
NGT: normal glucose tolerance, IGT: impaired glucose tolerance, T2D: type 2 diabetes

## Supporting Information

698    **S1 Table: Characteristics of serum samples used for antibody capture experiments**

699    **S2 Table: Mass spectrometry results of tryptic digests**

700    **S3 Table: Multiple stepwise linear regression analysis**