The Latent Semantic Space and Corresponding Brain Regions of the Functional

Neuroimaging Literature via NeuroSynth

Fahd Alhazmi

School of Behavioral and Brain Sciences

The University of Texas at Dallas

MS: GR4.1

800 West Campbell Rd.

Richardson, TX 75080 USA

Derek Beaton

Rotman Research Institute

Baycrest Health Sciences

3560 Bathurst Street

Toronto, ON, Canada M6A 2E1

Hervé Abdi

School of Behavioral and Brain Sciences

The University of Texas at Dallas

MS: GR4.1

800 West Campbell Rd.

Richardson, TX 75080 USA

Abstract

Because terminology varies widely from study-to-study, and even more so from discipline-to-discipline, the neuroimaging literature is becoming increasingly complex to navigate. For examples: (1) terminology changes over time (e.g., alcoholism to alcohol use disorders), (2) a single term can have many—or even amorphous—definitions (e.g., MVPA as multi-voxel or multivariate pattern analysis, which itself spans numerous different techniques) and (3) multiple terms describe the same concept (e.g., in vision studies "striate cortex," "calcarine sulcus," "V1," "primary visual cortex," and "Brodmann area 17," all describe, essentially, the same brain region in functional neuroimaging). Such a diversity of terminologies makes interpretations, reviews, and meta-analyses of the literature difficult to perform and consume. As the volume of scientific questions across neuroimaging has grown—and with this growth, the evolution of terminology across neuroimaging—it is important to clarify: What are the primary domains of neuroimaging and what are their most commonly reported brain regions?

INTRODUCTION

In recent years, several meta-analytic approaches have been developed that link reported brain activations with keywords and topics such as coordinate-based meta-analysis (CBMA). CBMA was specifically developed for aggregating and synthesizing neuroimaging data reported in a standard format (P. T. Fox et al. 2014). Some of the most prominent CBMA tools used in research are BrainMap (Laird et al. 2005), SumsDB (Van Essen et al. 2009), Brede (Nielsen 2003) and NeuroSynth (Yarkoni et al. 2011)—the database of interest in this paper. The main functionalities of many CBMA tools are to (a) store coordinate information by study, and (b) provide spatially consistent meta-analytic activation maps. For example, Nielsen and colleagues (Nielsen et al. 2004) analyzed 121 neuroimaging papers with 2,655 reported activations using probability models followed by a non-negative matrix factorization-latent semantic analysis to associate brain coordinates with the words used in the papers' abstract (e.g., "pain" was strongly associated with the anterior cingulate). More recently, Poldrack and colleagues (Poldrack et al. 2012) analyzed more than 5,800 papers to model the associations between topics derived from the full text of studies and the reported peak coordinates via "topic mapping." Poldrack and colleagues showed that—with a "topic mapping" approach to semantic analysis—CBMA could reveal new relationships between brain activation and cognitive processes or psychiatric disorders (for different flavors of CBMA, see e.g., Rubin et al. 2016; de la Vega et al. 2016). Furthermore, some extensions of CBMA can link additional data types (e.g., gene expression) with brain regions and keywords (Mesmoudi et al. 2015; A. Fox et al. 2014; Gorgolewski et al. 2014; Rizzo et al. 2016).

Meta analytics tools can better facilitate interdisciplinary communication because they integrate data from different sources. However, there is currently no characterization of the primary domains of neuroimaging and no characterization of the brain regions most associated with these domains. In this work, we define the primary domains of neuroimaging

based on the semantics of the literature (i.e., abstracts). Our study is designed to achieve three major goals: (1) define the "semantic space" of the neuroimaging literature, (2) identify semantically defined domains within the literature, and (3) map brain activations onto these domains. To do so, we used correspondence analysis (CA)—a technique similar to principal components analysis (PCA) that was originally created for analyzing the co-occurrences of words in a corpus (Lebart et al. 1998; Benzecri 1976; Escofier-Cordier 1965, Abdi & Béra, 2014)—to identify neuroimaging domains from co-occurrences of words in the neuroimaging literature as identified in the databases in Neurosynth (Yarkoni et al. 2011).

METHODS

Data and preprocessing

Neurosynth is an open source and open science initiative—hosted via the website www.neurosynth.org—that facilitates meta-analyses and reviews of the functional neuroimaging literature (Yarkoni et al. 2011). Neurosynth, at the time of this writing, contains more than 11,406 articles from the functional neuroimaging literature. When we began this work, Neurosynth contained 10,903 articles (from 43 journals), which were the basis of this study. As an aside, some articles in our data set no longer appear in Neurosynth because Neurosynth periodically updates its content for exclusion (e.g., to remove structural only studies) and public release. See http://github.com/neurosynth/neurosynth-data and http://www.neurosynth.org/ for details

Neurosynth uses automated webcrawlers to fetch data (e.g., abstract text, peak coordinates) and metadata (e.g., PubMed ID, title, year published, journal) of neuroimaging studies. For our study, we created and analyzed two data tables derived from Neurosynth data: (1) a "studies × words" matrix and (2) a "studies × voxels" matrix, where studies are identified by their PubMed ID (PMID) number. To achieve our three goals, our study

comprised three major parts that correspond to each goal, wherein each major part has several steps. All analyses were conducted with a variety of publicly available packages (noted in relevant sections) or in-house scripts written in Matlab (MathWorks, Natick, MA), R (R Core Team 2014), and Python (Python Software Foundation, https://www.python.org/) languages and environments.

To create a studies-by-words matrix for analysis, we required information from Neurosynth and PubMed (http://www.ncbi.nlm.nih.gov/pubmed/). With PMIDs from the Neurosynth database, we obtained from PubMed the text in all abstracts associated with the studies in the Neurosynth database. Next, we used the `tm` package in R (Feinerer 2011) to conduct several preprocessing steps that were used in previous works (Ailem et al. 2016) and that consisted in: (1) converting all text to lower-case, (2) removing all punctuations, numbers, emails and web addresses, (3) removing all words of length one or two, (4) removing step words, meta-words and words that describe numbers, quantities, nationalities, cities, or names (e.g., "publisher," "article," "date," "ten," "zero," "weeks," "european," "montreal," "welcome"), (5) converting British English to American English, (e.g., "behaviour" to "behavior") and finally (6) stripping out white spaces. Once the data were cleaned, some words with different meanings were updated so they did not have the same "stems." For example, "posit," "positive," "positively," and "position," "positioning" would correspond to the same stem "posit;" therefore some words were altered so they would only have the same stem if they had (in general) the same meaning. In a final step, we went through all remaining words individually to identify words that were potentially missed in the previous steps (e.g., "science," "academic," "publishing"). With a final set of words, we created a matrix of studies (rows) by words (columns). Each cell of this matrix contained the frequency of a word used in the abstract of a study. For example, the abstract of PMID 17360197 used the word "cold" 28 times. Finally, the studies-by-words matrix went through

one more cleaning step: we eliminated infrequent words. Words with frequencies below the 3rd quartile (in our case: less than 16 occurrences) were removed. This step was followed by the removal of two studies that had withdrawal notices from the publisher. The final studies-by-words matrix contained 10,898 studies and 3,114 words (the full data table is provided in Supplemental Material and at https://github.com/fahd09/neurosynth_semantic_map).

Correspondence Analysis

Because our data are the number of occurrences (i.e., counts), we required correspondence analysis (CA)—a technique designed specifically to analyze co-occurrences and often described as a version PCA tailored for qualitative data. Like PCA, CA decomposes a matrix into orthogonal components rank-ordered by explained variance (Greenacre 1984; Abdi & Béra 2014; Lebart et al. 1984) . CA is a bi-factor analysis that analyzes the relationship between the row and the column of a contingency table data matrix. Each row (study) and column (word) item is assigned a "component score" (a.k.a. factor score) that reflects the amount of variance this item contributes to a given component. CA places emphasis on rare items so that they contribute a high amount of variance, while frequent items contribute little variance (see Greenacre 2017); this is particularly useful for our study because frequent words (e.g., "brain") will be near the origin (i.e., zero) of the components whereas rare words (e.g., "polymorphism") will be far from the origin and thus are the sources of variance for the components. CA operates under the same assumptions as χ^2 that is how far the observed data deviate from independence. Finally, because both rows and columns are represented in the same space (with the same variance), we can interpret the relationships within row items and within column items as well as the relative relationships between row and column items. Finally, because we wanted to identify brain regions most associated with semantically-defined domains, we used a technique called *supplementary projection* (Greenacre 2017; Abdi & Béra 2014) that allows us to predict a *supplementary*

data set (i.e., studies × voxels) from the component structure of the *active* data set (i.e., studies × words).

We applied CA to studies-by-words matrix to define the semantic space of the neuroimaging literature. We used in-house MATLAB code, as well as the `ExPosition` (Beaton et al. 2014) and `ggplot2` (Wickham 2009) packages in R to perform CA as well as visualizations and additional analyses (i.e., visualizations, resampling-based inference tests, clustering, and supplementary projections; see following sections).

Split-half Resampling

Split-half resampling (SHR, Strother et al. 2002; Churchill et al. 2012) is a cross-validation (CV) technique that evaluates the stability of the results of a statistical analysis performed on a data set by randomly splitting this data set into two (approximately) equal sized non overlapping data sets, and then performing the same analysis on each data set. The similarity (e.g., correlation) between the results obtained from these two data sets is then used to evaluate the reliability of the results (i.e., replicable effects). SHR is performed many times to create a distribution of reliability estimates.

We used SHR to identify the most replicable components in two ways: (1) split the data by rows, and (2) split the data by columns; in both approaches, we performed CA on each split set, and then computed the absolute correlation[1] between the component scores of each split. SHR was performed 1,000 times to create a distribution of (absolute) correlations between components for both the (1) row component scores conditional to the columns, and (2) the column sets of scores conditional to the rows. We then computed the average (absolute) correlations to detect which components (after 1,000 splits) were most replicable between splits in order to identify a low rank approximation of the semantic space (i.e., component selection via SHR).

---

[1] We used absolute correlation because there can be trivial sign flips between sub-samples of data, wherein the sign is irrelevant, but the magnitude of the correlation is relevant.

Clustering of Studies and Assignment of Words

We performed hierarchical clustering (HC), with squared Ward linkage (Murtagh &
Legendre, 2014), on the component scores for the studies (rows). We chose squared Ward
linkage because its objective function minimizes the error sums of squares (and thus aims for
an optimal ANOVA-like configuration). The component scores take into account the
explained variance per component (i.e., Component 1 explains more variance than
Component 2). After HC, we performed cluster stability analysis via Calinski-Harabasz index
(Calinski & Harabasz, 1974) in order to identify a stable number of clusters. Once studies
were divided into clusters, we then used distance-based classification in order to assign each
word (column) to a study cluster. Hierarchal clustering and cluster stability were conducted
in R via `hclust` and `clusterCrit` (Desgraupes 2015), respectively.

Producing Brain Maps

Activation maps are represented in Neurosynth as peak activations of individual
studies as centers of a sphere with a radius of 6mm. Voxels inside the sphere have a value of
1 and the other voxels have a value of 0. The voxels-by-studies matrix then uses a vectorized
(flattened) version of the peak activation maps with reference to a 3D brain. The voxels-by-
studies matrix initially contained 10,898 studies and 228,453 voxels (i.e., voxels within MNI
space). For our analyses, infrequently reported voxels (i.e., less than 10% of studies) were
removed. The final studies-by-voxels matrix contained 10,898 studies and 206,077 voxels.
We computed two different brain activation maps from the semantic space. The first type of
activation map was a component-wise map. Brains were projected onto (i.e., predicted by)
the semantic space—per replicable component—via supplementary projections. The second
type of activation map was simply the sum of peak activations *per study cluster*.

Supplementary Projections

Supplementary—a.k.a. out of sample—observations (or variables) can be integrated into an existing analysis performed on a different set of observations (or variables) referred to as the active data set. Supplementary data are assigned component scores by computing the least square projection for observations (or variables) onto the space defined by the active observations (or variables). We used supplementary projection to predict component scores for voxels *from* the component scores of studies defined in the in the semantic space (i.e., CA of studies × words). Predicted activation maps (from the supplementary scores) were projected back into MNI space. The predicted maps were then thresholded to 2% (to only show extreme voxels on either side of the distribution of factor scores) and smoothed with a Gaussian 10mm kernel (FWHM). Smoothing and glass brain visualizations were conducted with `nilearn` (Abraham et al. 2014) in python. All resulting maps (5 components maps and 6 cluster maps) are shared publicly in a Neurovault repository (http://neurovault.org/collections/2002/, last accessed June 12, 2017; Gorgolewski et al. 2016).

RESULTS

CA was applied to the 10,898 studies × 3,114 words matrix and produced 3,112 components (see Figure 1a for the Scree plot). SHR was applied 1,000 times to split the studies and then again 1,000 times to split the words of the studies × words matrix. SHR of both sets revealed that the first three components were very reliable (see Figure 1b and c). Additionally, Components 4 and 5 show a moderate-to-strong reliability via correlation of words component scores (via splitting of studies; see Fig 1b). Hierarchical clustering and cluster stability analysis were performed only on the first five components. Cluster stability analysis revealed 6 reliable study clusters (see SFig. 1). Words were then assigned to the closest study cluster.

To help interpret the components we used the words at the extremes (i.e., highest contributing variance) for each component (Table 1). Table 2 shows the total and relative number of studies and words per cluster. As with the components, the most frequent words within each cluster help us interpret the cluster's meaning (Table 3). Furthermore, we also identified the words that were closest to the barycenter (i.e., multidimensional mean) of each cluster (across all five dimensions; Table 4). Finally, we also provide the titles and PubMed IDs of the twenty studies closest to the barycenter of each cluster in supplemental Tables S1-S6. Component maps are presented in Figure 2. We present component maps of the words and studies separately. In each map, we color each dot (i.e., a study or word) by its associated cluster. Components 1, 2, and 3 are visualized in Figures 2a–d. We show Components 4 and 5 separately from the other components (Figures 2e and f) because studies on Components 4 and 5 comprise a single cluster (see next section). Finally, brain maps for the components are presented in Figure 3, and brain maps for clusters are presented in Figure 4. In *Results* the components and clusters are referred to by numbers: The component number reflects its rank order (by variance), but the cluster numbers are arbitrary.

Components

Component 1

Component 1 separates words associated with cognitive (especially linguistic) studies from words associated with clinical or translational studies (see Fig. 2a–b). Words at extreme positive and negative side of Component 1 are listed in Table 1. The projected brain map for Component 1 (Fig. 3a) shows that: (1) the positive side of Component 1 is associated with the left temporal lobe, bilateral occipito-temporal, and parietal regions, and (2) the negative side of Component 1 is associated with many subcortical structures.

Component 2

Component 2 separates words often associated with diffusion imaging studies from words often associated with affective and decision-making studies (see Fig. 2a–b, and Table 1 for words at positive and negative extremes). The projected brain maps for Component 2 (shown in Fig. 3b) show that: (1) the positive side of Component 2 is associated with bilateral somatosensory areas and the right cerebellum, and (2) the negative side of Component 2 is associated with subcortical structures as well as medial prefrontal cortex.

Component 3

Component 3 separates words associated with language studies from words associated with somatosensory studies and pain (see Fig. 2c–d, and Table 1 for words at positive and negative extremes). The projected brain maps for Component 3 (shown in Fig. 3c) show that: (1) the positive side of Component 3 is associated with the left lateralized language-related areas (e.g., temporal and frontal areas known as Broca's Area and Wernicke's Area), and (2) the negative side of Component 3 is associated with somatosensory cortex in addition to the brainstem.

Components 4 & 5

Components 4 and 5 (together) identify a highly distinct set of words and studies related to molecular, genetic, and genomic neuroimaging studies (i.e., "imaging genetics;" see Fig. 2e–f, and Table 1 for words at positive and negative extremes). The brain maps for Component 4 (Fig. 3d) show that: (1) the positive side of Component 4 is associated with medial structures of parietal areas (precuneus), and (2) the negative side of Component 4 is associated with bilateral somato-sensory areas as well as the insular cortex and brainstem. The brain maps for Component 5 (in Fig. 3e) show that: (1) the positive side of Component 5 is associated with the posterior cingulate and medial prefrontal cortices, and (2) the negative side of Component 5 is associated with bilateral somato-sensory areas and the brainstem.

Component summaries

Component 1 reflects the research spectrum of neuroimaging: from basic science research on the positive side to clinical/translational neuroimaging research on the negative side (Figs 2a-b; Table 1). While the basic science research is more associated with cortical structures, the clinical/translational research is more linked with subcortical structures (Fig 3a). Component 2 reflects a methodological spectrum that ranges from cognitive tasks on the negative side to multi-modal imaging (i.e., structural-functional association) on the positive side. The negative—and much more densely populated—side of Component 2 is more associated with studies on affect and emotion (Fig. 2a–b and Table 1) with dense projections to subcortical and prefrontal cortex, whereas the positive side of Component 2 is linked to studies that rely on multi-modal MRI and other related methodologies with somatosensory, temporal and cerebellar projections (Fig. 3b). Component 3 reflects low-to-high level of cognition; studies about higher-order cognitive processes (especially linguistics) are on the positive side of Component 3 (with cortical projections to the bilateral temporal and frontal regions; Figs 2c–d; Table 1; Fig. 3c), whereas studies on lower-order cognitive processes (such as sensation, perception, and direct sensory stimulation) are on the negative side of Component 3 (and also projects to the bilateral somatosensory areas and the brainstem in addition to the left cerebellum; Figs 2c–d; Table 1; Fig. 3c). Finally, Components 4 and 5 showed a distribution of words and studies that makes a near perfect 45° angle between Components 4 and 5 that extended out from the origin. These words and studies were almost entirely molecular, genetic, and genomic neuroimaging (i.e., "imaging genetics") studies. While studies on Component 4 are more associated with genetic contributions to cognition in healthy populations, with projections to parietal and frontal regions studies on Component 5 were more associated with genetic contributions in disordered populations with projections to bilateral somatosensory regions.

Clusters

Cluster 1

Cluster 1 contains words primarily associated with language/speech production, comprehension, and disorders, as well as knowledge processing. Some examples include: *decod, word, superior, auditori, languag, semant, percept, speech, recognit, complex* (see Tables 3 and 4; a full list is provided in the Supplemental Materials). Figures. 2a-b show that this cluster primarily lies on positive sides of Components 1 and 3. Summed peak activations of individual studies in this cluster showed in the bilateral frontal and temporal regions—which are often associated with language and knowledge representation—in addition to the anterior cingulate cortex (see Fig 4a).

Cluster 2

Cluster 2 contains words associated with developmental, lifespan, and aging studies, as well as their respective disorders. Some examples include: *patient, differ, chang, healthi, age, structur, breakdown, degen, ecnp, epidemiolog* (see also Tables 3 and 4; a full list is provided in the Supplemental Materials). Figs. 2a–b shows that this cluster primarily loads on the negative side of Component 1 and on the positive side of Component 2. Summed peak activations of studies in this cluster (see Fig 4b) showed a diffuse pattern of activations in the frontal and parietal areas, as well as in the subcortical regions.

Cluster 3

Cluster 3 contains words primarily associated with sensation (cutaneous and olfaction) and movement. Some examples include: *include motor, pain, movement, hand, stimul, sensori, thalamus, somatosensori, reflex, anesthet* (see also Tables 3 and 4; the full list is provided in the Supplemental Materials). Figs. 2c-d shows that this cluster loads primarily on negative side of Component 3. Summed peak activations of studies in this cluster (Fig. 4c) showed in the bilateral somato-sensory areas and the thalamus.

Cluster 4

Cluster 4 contains words associated with more "traditional" aspects of human cognitive neuroscience: those rooted in cognitive and experimental psychology (i.e., they rely primarily on behavioral tasks to examine neural correlates). Tables 3 and 4 show words that characterize Cluster 4, and some examples include: *activ, function, task, area, fmri, network, memori, effect, visual, decay*. Figures 2a–c show Cluster 4 is closest to the origin point across all components with no apparent tendency toward any axis. Summed peak activations of studies in this cluster (shown in Fig. 4d) showed in the mid-line and bilateral frontal regions, in addition to bilateral occipito-temporal region.

Cluster 5

Cluster 5 contains studies/words that describe affective processes, such as emotional responses and decision-making, but also includes a number of studies and words related to substance use disorders and mood disorders. Cluster 5 includes the words: *emot, prefront, reson, cingul, examin, medial, amygdala, negat, social, diminish, take* (see Tables 3 and 4). Figs. 2a–b show that this cluster lays mostly on the negative side of Component 2. Summed peak activations of Cluster 5 (Fig. 4e) appeared in subcortical areas and medial frontal regions.

Cluster 6

Cluster 6 loads almost entirely and exclusively on both Components 4 and 5 (see Figs 2e–f). Cluster 6 contains words such as: *variat, genet, dopamin, gene, carrier, allel, genotyp, receptor, polymorph, dopaminerg, comt, serotonin, apo, norepinephrine* (see Tables 3 and 4). Summed peak activations of Cluster 6 (Fig. 4f) showed almost entirely in the subcortical areas.

Cluster Summaries

Clustering revealed semantically homogenous subgroups within neuroimaging. Cluster 1 (henceforth: *Knowledge Representation & Language Processing*) represents studies

that mainly investigate knowledge representation and language processing. Cluster 2 (henceforth: *Developmental, Lifespan, and Disorders*) represents studies that investigate developmental and adult lifespan research in addition to brain disorders. Cluster 3 (henceforth: *Sensation, Movement, and Action*) represents studies that investigate sensation, movement and action. Cluster 4 (henceforth: *Cognition and Psychology*) represents the majority of cognition and psychological research. Cluster 5 (henceforth: *Decision, Emotion, and Substance Use*) represents studies that investigate decision-making, emotions and substance use (or abuse). Cluster 6 (henceforth: *"Imaging Genetics"*) represents a unique dimension (i.e., Components 4 and 5) of molecular, genetic, and genomic neuroimaging ("imaging genetics") studies.

Follow up Analyses

Upon completion of analyses, there were two clusters that stood out: (1) Cluster 4 (*Cognition and Psychology*)—which is essentially the "average" neuroimaging study because it is centered roughly on the origin of the components—and (2) Cluster 6 (*"Imaging Genetics"*)—which is comprised of the studies that define Components 4 and 5). Notably, Cluster 4 (*Cognition and Psychology*) reflects the origins of neuroimaging use (i.e., cognitive psychology), whereas Cluster 6 (*"Imaging Genetics"*) reflects the current state-of-the-art (i.e., translational and interdisciplinary work).

Furthermore, we wanted to compare our (components and clusters) brain maps with existing functional networks, especially those derived from thousands of fMRI studies such as Yeo et al.'s 12-component ontology (Yeo et al. 2015). In Yeo et al. (2015), a hierarchal Bayesian model was applied to 10,449 experimental contrasts in the BrainMap database in order to estimate the probability that each pre-defined task category would engage a specific cognitive component, and the probability that each cognitive component would engage brain regions (represented by voxels).

Therefore we followed up with two additional sets of analyses: (1) a breakdown of our clusters over time, and (2) a comparison of our predicted brain maps to those in Yeo et al., (2015).

Temporal Properties of Clusters

Figure 5 shows the relative frequency of the number of studies in each cluster sorted by year. Cluster 4 (*Cognition and Psychology*) accounts for a substantial amount of studies in the earlier years. For example, approximately 50% of all neuroimaging studies (in Neurosynth) in 2000 were in Cluster 4 (*Cognition and Psychology*). On the other hand, Cluster 5 (*Decision, Emotion, & Substance Use*) started as a small proportion of all neuroimaging studies in the earlier years, but now accounts for nearly 33% of all studies. We discuss the temporal properties of these clusters further in the Discussion.

Comparisons with Yeo et al., (2015)

Figure 6 shows the correlations between our maps and the maps from Yeo et al (2015). In order to use only the valid voxels that lie within the brain, we only included non-zero voxels from the component maps. Correlations were performed with Yeo et al.'s (2015) maps available from NeuroVault (http://neurovault.org/collections/866/, last accessed June 7, 2017). We refer to Yeo et al.'s (2015) components as, e.g., Yeo Component 1 (YC1) or Yeo Component 6 (YC6) while we refer to our own components as "Component 1" or "Component 6." There were several correlations of note for both the components (Figure 6a) and the clusters (Figure 6b). To note, although the magnitudes of those correlations are interpretable, the sign (or direction) of the correlation are not easily interpretable.

Figure 6a shows strong correlations between our Component 2 and YC11 and YC12. Both maps show strong association with bilateral subcortical structures (e.g., amygdala and striatum) in addition to their relationship with subcortical-related functions such as emotions and affect. Also, there is a strong correlation between our Component 3 and YC5 because

both maps show strong association with temporal and frontal activations in addition to their relationships with semantic knowledge and language processing. Furthermore, there is a strong correlation between our Component 4 and YC6 because both maps show strong associations with medial parietal and frontal areas (commonly known as the frontal-parietal network; Smith et al. 2009).

Similar to the components, correlations between our clusters and Yeo components are illustrated in Figure 6b. Our Cluster 1 is most correlated with YC 5 followed by YC3, all of which have activations in the temporal and frontal regions and are generally involved with knowledge representation and higher-order semantic processing. Our Cluster 3 is correlated with YC1 followed by YC7, both of which have activations in somatosensory areas and are involved in sensation and movement processing. Our Clusters 5 are both correlated with YC11 and YC12, all of which are associated with activations in subcortical structures and are associated with tasks that involve some aspect of affective or emotional processing. Although our Cluster 6 also share the same regions (and is most similar to YC11 and YC12), it comes from entirely different set of molecular and genetic studies.

DISCUSSION

In recent years, there have been many meta-analyses, mega-analyses (analyses of pooled data across many studies), and other large-scale analyses of data within neuroimaging. In general, the aims of such analyses are to (1) test or refute findings and hypotheses (Wager et al. 2007), (2) build a consensus around particular models, hypotheses, or theories (Salimi-Khorshidi et al. 2009), (3) estimate consistency of findings (Wager et al. 2009), (4) help define related brain regions and networks (Toro et al. 2008; Mesmoudi et al. 2013), (5) interpret functional maps (Laird et al. 2011), or (6) segment the brain in new ways with

resting-state *f*MRI measurements (Yeo et al. 2011; Power et al. 2011) or using massive multi-modal data (Glasser et al. 2016).

Relationship with other functional maps and meta-analyses

Using the meta-analytic cognitive component maps from Yeo et al (2015), we showed a substantial overlap between many of our maps and maps from Yeo. The major difference is that our meta-analytic maps were defined conditionally through the language used (in abstracts) from the functional neuroimaging literature, whereas others authors took a more brain-centric approach, for examples: network- and meta-maps generated by with resting state fMRI (Yeo et al. 2011; Power et al. 2011) or via meta-analysis of data from hundreds or even thousands of studies (Yang et al. 2016; Yeo et al. 2015; Poldrack et al. 2012; de la Vega et al. 2016).

Our components explain the primary sources of variance of language used: in the field at large (i.e., Component 1), for methodological tools (i.e., Component 2), in various aspects of cognition (i.e., Component 3), and in relatively new studies with highly-specific terminology (i.e., Components 4 and 5). With supplementary projections we also showed that these language-based components are frequently associated with particular brain regions. While the components indicate language variation and gradients, our clusters define the boundaries of functional neuroimaging into specific—albeit large—sub-domains. Furthermore, our analyses revealed that there are, perhaps, biases or preferentially studied brain areas based on domains (i.e., clusters).

Some of our components also reflect, to a degree, current debates such as the distinctions between "neurological vs. psychiatric brain disorders" (White et al. 2012). For example, Crossley et al. (2015) recently used CBMA of voxel-based morphometric (VBM) studies to show a "neuroimaging-based" evidence for the biological distinctions between neurological vs. psychiatric disorders (Crossley et al. 2015). However, our components show

that neuroimaging studies in neurology and psychiatry do not use the same terminology and

thus could be a source of the "versus" argument. As an illustration of this contention, we

have selected some of the same neurology and psychiatry related terms used by Crossley et

al., (2015) to highlight particular features of our components. First, all the words related to

psychiatric or neurological disorders (Table 5) appear on the negative side of Component 1—

a configuration that supports our interpretation of a spectrum from basic science to applied

and clinical neuroimaging. Furthermore, the neurological and psychiatric terms from

Crossley et al., (2015) oppositely load on both Components 2 and 4 (Table 5): a configuration

that reflects overall differences in *patterns* of terminology between neurological and

psychiatric studies and thus expresses a dissociation of neurological studies and their regions

(such as sensorimotor cortices and insula; in red) from psychiatric studies and their regions

(such as limbic and prefrontal areas; in blue) as see in Figure 3b.

Clusters: corresponding brains and temporal effects

The positions—and contents—of our clusters reveal a broad configuration of the

neuroimaging literature. Cluster 4 (*Cognition and Psychology*) is the closest to the barycenter

(origin of the axes across all components) and thus represents the average or most common

neuroimaging study. This interpretation is supported by Cluster 4 (*Cognition and

Psychology*) because it contains a substantial proportion of words and studies (~33% of

words and ~29% of studies, see Table 3). Thus, much of the neuroimaging literature has

been—and appears to still be—rooted in the approaches from cognitive and psychological

domains. Summed peak activations of studies in this cluster (shown in Fig. 4d) show a high

association with a wide set of cortical areas in the medial and bilateral frontal, occipital and

subcortical regions that are associated with task performance. We also see opposition of

clusters and thus suggests these are the sources of variance for our components. For example,

Cluster 5 (*Decision, Emotion and Substance Use*) opposes all other clusters on Component 2

(Figs. 2a-b)—a pattern that further supports the neurological vs. psychiatric dissociation of Component 2. Summed peak activations of studies in this cluster (shown in Fig. 4e) show high association with the subcortical areas and medial frontal regions that are generally associated with emotional processing and decision-making process. Similarly, Cluster 3 (*Sensation, Movement and Action*) is opposed to all other clusters on Component 3 (Figs. 2c-d)—a component, as we previously noted, expresses a spectrum from low-to-high level processing. Summed peak activations of studies in this cluster (shown in Fig. 4c) show high association with the bilateral somatosensory areas and the thalamus. Furthermore, Cluster 6 (*"Imaging Genetics"*) is almost entirely defined by the unique configuration of both Components 4 and 5 (Figs. 2e–f). Not only does Cluster 6 reflects a unique subfield of neuroimaging, but also indicates that "imaging genetics" uses an almost exclusive set of words different from the vocabulary of the rest of neuroimaging (cf., the 45 degree angle from Components 4 and 5). Summed peak activations of Cluster 6 (Fig. 4f) are almost entirely associated with subcortical areas. Finally, both Clusters 4 (*Cognition and Psychology*) and 5 (*Decision, Emotion and Substance Use*) proportionally explain over half of the literature at any given time (Fig. 5).

Our clusters and their respective brain maps are consistent with results of other meta-analysis. The activation map of Cluster 1 (*Knowledge Representation and Language Processing*; Fig. 4a) is similar to other published meta-analytic maps and reviews of language processing and semantic representation (Binder et al. 2009; Price 2010; Price 2012; Fedorenko & Thompson-Schill 2014; Bookheimer 2002). The activation map of Cluster 3 (*Sensation, Movement and Action*; Fig 4c) is similar to other maps from studies investigating pain localization (Perini et al. 2013; Amanzio et al. 2013; Schomers & Pulvermüller 2016; Friebel et al. 2011; Vierck et al. 2013) in addition to the somatosensory co-activation network (Smith et al. 2009). Finally, the activation map of Cluster 5 (*Decision, Emotion, and*

Substance Use; Fig. 4e) is also highly similar to the map of the structures involved in different aspects of emotional processing and decision-making (Bartra et al. 2013; Lindquist 2010; Etkin & Wager 2007; Buhle et al. 2014; Phan et al. 2002).

General discussion

Many meta-analyses and meta-analytic tools for neuroimaging have a common (even if unstated) goal: to help define ontologies (Poldrack & Yarkoni 2016; Poldrack et al. 2011) so that we can bridge the gaps in our understanding of brain function and cognition. However, even with many tools at our disposal, there are known biases in neuroimaging (Jennings & Van Horn 2012) and the language we use can make building such ontologies difficult. However, with well defined language and homogenization of reporting results, fields such as genomics can provide a more robust assessment of the relationship between studies and the roles of particular genetic effects (Ailem et al. 2016).

Based on the analysis of term co-occurrences in the abstracts of 10,898 neuroimaging article, we have identified a highly reliable set of dimensions and subfields that define the underlying semantic space of the neuroimaging field. Most researchers tend to stay *within* their specialized domain (by using specific key terms common to their field) and this behavior may restrict what they can conclude and how they report their findings, because they use a preferred or required terminology. In fact, Clusters 2 (*Development, Lifespan and Disorders*) and 5 (*Decision, Emotion and Substance Use*), as well as Components 1 and 2 show that there are language barriers between different types of clinical and experimental studies that could preclude thorough reviews of relevant literature.

Because such diverse terminologies and highly specialized fields could cause researchers to overlook relevant work in domains related and unrelated to their own, several recent tools have been proposed, including one that uses Neurosynth. Recently, (Yuan et al. 2017) introduced MAPBOT, which helps researchers navigate relevant studies in

Neurosynth, but conditional to a region of interest. For example, in their paper, Yuan et al., (2017) use a thalamic mask to generate a study × term matrix and then decompose that matrix with non-negative matrix factorization. MAPBOT helps researchers navigate the literature and relevant terms conditional to *a priori* masks. Additionally, a tool called Papr (Chawla 2017; Leek 2017) was released to help researchers find preprints on bioRxiv that may be of interest to them. Papr is based on an approach similar to the approach that we took here: Users can move through the components-based subspace to find articles with similar abstracts as well as other users with similar interest. Papr provides for bioarxiv some of the mechanisms that our approach could provide for the neuroimaging field, namely the ability to function as a recommendation engine and allow for users to interact with other users that have similar profiles. While both Papr and MAPBOT provide some tools to navigate and search the literature—especially MAPBOT with Neurosynth—both are lacking the key features and information we provided here. We believe that our approach to structuring the functional neuroimaging literature is critical to both help organize the field and to help researchers navigate the literature.

Limitations

There are limitations of our work: The first limitation is that Neurosynth only searches a specific set of journals. Although it samples from 43 journals, there is a possible sampling bias that may exclude certain neuroimaging studies; these could be highly specific studies published in journals that are not highly focused on neuroimaging. Thus our characterization of neuroimaging is a characterization of the field by proxy of Neurosynth. A second limitation is the fact that abstracts need to obey many constraints and rules that may introduce a systematic bias. While analyzing full texts of studies could provide a different configuration, abstracts are the primary entry point for readers, and are considered the

primary message authors wish to convey. A third limitation is that we only relied on single words to represent literature instead of 2-words or *N*-word model. For example, both "working memory" and "post-traumatic stress disorder" are concepts that are expressed in many words and should be treated as mutli-word concepts.

Conclusion

Our work shows that different domains use different words, and that studies within these domains also report (or only study) common brain areas. Finally, we believe that neuroimaging would benefit from a broader harmonization of language (à la the COBIDAS appendix on how to report routine *f*MRI analyses; Nichols et al., 2016) to put the field on the path towards formal ontologies (Poldrack & Yarkoni 2016). Formal and more rigorous ontologies—such as those in genomics—will be required as our field moves forward and connects brain imaging to a variety of other modalities (e.g., genetics; Cioli et al. 2014; Rizzo et al. 2016).

Figures

Figure 1

(a) Scree plot of the explained variance for all components

(b) Heatmap of correlations between words component scores through SHR

(c) Heatmap of correlations between studies component scores through SHR

Figure 1: Variance explained and reproducibility (via split-half resampling; SHR) of latent semantic components. (a) The Scree plot shows the explained variance per component for all components. Figures 1 (b) and (c) show heatmaps of correlations between component scores after SHR where (b) shows average correlation after SHR for the words component scores and (c) shows the average correlation after SHR for the studies component scores. Both the Scree plot and the heatmap for the studies component scores suggest three high variance and highly reproducible components. The heatmap for the words component scores also show that the first three components are highly reproducible, but also that Components 4 and 5 are reproducible in the words component scores.

Figure 2

(a)

(b)

(c)

(d)

(e)

(f)

Figure 2: Visualization of component scores for both the words and the studies on Components 1 through 5. Axes are components and individual dots represent either a particular study or particular word. Words and studies are colored by which cluster they belong to and thus illustrate the large sub-domains within fMRI. Figures 2 (a) and (b) show the words and studies (respectively) component scores for Components 1 (horizontal) and 2 (vertical). Figures 2 (c) and (d) show the words and studies (respectively) component scores for Components 1 (horizontal) and 3 (vertical). Figures 2 (e) and (f) show the words and studies (respectively) component scores for Components 4 (horizontal) and 5 (vertical). While most words and studies form large groups within the axes, Components 4 and 5 show a highly specific subset of words and studies, of which nearly all are assigned to Cluster 6 (typically fMRI studies that include genetic and molecular terms).

Figure 3

(a) Voxel supplementary projections onto Component 1 (left) and some positive and negative

words associated with Component 1 (right).

(b) Voxel supplementary projections onto Component 2 (left) and some positive and negative

words associated with Component 2 (right).

(c) Voxel supplementary projections onto Component 3 (left) and some positive and negative

words associated with Component 3 (right).

(d) Voxel supplementary projections onto Component 4 (left) and some positive and negative

words associated with Component 4 (right).

(e) Voxel supplementary projections onto Component 5 (left) and some positive and negative words associated with Component 5 (right).

Figure 3: Visualization of brain maps, per component, as predicted (via supplementary projections) by the words × studies component scores and a word cloud that shows some words that either loads on the positive or negative axis. (a) The projected map for Component 1 shows that the positive side (marked in red) is associated with the left temporal lobe, bilateral occipito-temporal, and parietal regions, while the negative side (marked in blue) is associated with many subcortical structures. (b) The projected maps for Component 2 show that the positive side is associated with bilateral somatosensory areas and the right cerebellum, while the negative side is associated with subcortical structures as well as medial prefrontal cortex. (c) The projected brain maps for Component 3 show that the positive side is associated with the left lateralized language-related areas while the negative side is associated with somatosensory cortex in addition to the brainstem. (d) The projected map for Component 4 show that the positive side is associated with medial structures of parietal areas (precuneus) while the negative side is associated with bilateral somato-sensory areas as well as the insular cortex and brainstem. (e) The projected brain maps for Component 5 show that

the positive side is associated with the posterior cingulate and medial prefrontal cortices

while the negative side is associated with bilateral somato-sensory areas and the brainstem.

Figure 4

(a) Sum of peak activations for Cluster 1

(b) Sum of peak activations for Cluster 2

(c) Sum of peak activations for Cluster 3

(d) Sum of peak activations for Cluster 4

(e) Sum of peak activations for Cluster 5

(f) Sum of peak activations for Cluster 6

Figure 4: Visualization of brain maps, per cluster, computed as the sum of peak activations for all studies within a particular cluster. (a) Summed activations of individual studies in cluster 1 (*Knowledge Representation & Language Processing*) showed in the bilateral peaks in the frontal and temporal lobes— which are often associated with language and knowledge representation—in addition to a peak area in the anterior cingulate cortex. (b) Summed peak activations of studies in cluster 2 (*Development, Lifespan and Disorders*) showed a diffuse pattern of reported activations in frontal and parietal areas, as well as subcortical regions. (c) Summed peak activations of studies in cluster 3 (*Sensation, Movement and Action*) showed in bilateral somatosensory areas and the thalamus. (d) Summed peak activations of studies in cluster 4 (*Cognition & Psychology*) showed in the mid-line and bilateral areas in frontal regions, in addition to bilateral occipito-temporal regions. (e) Summed peak activations of cluster 5 (*Decision, Emotion, & Substance Use*) appeared in subcortical areas and medial frontal regions. (f) Summed peak activations of Cluster 6 (*Imaging Genetics*) showed almost entirely subcortical areas.

Figure 5

Figure 5: Shows the proportion of studies within each cluster over time. Cluster 4 (Cognition and Psychology) was, and still generally is the core of of fMRI research and as such comprises a substantial proportion of the literature. Though Cluster 4 (Cognition and Psychology) remains very large, it has decreased over time. Both Cluster 5 (Decision, Emotion, Substance Use) and Cluster 2 (Developmental, Lifespan, Disorders) have shown a considerable increases over time and now comprise, respectively, comparable proportions of the literature as Cluster 4 (Cognition and Psychology).

Figure 6

(a) Correlation between the 5 components and Yeo et al. (2015) maps.

(b) Correlation between the 6 clusters and Yeo et al. (2015) maps.

Figure 6: Shows the correlations between the maps generated by Yeo et al., (2015) and (a) our components or (b) our clusters. There are some notably high similarities between our brain maps (which were generated conditional to the latent semantic space) and the Yeo et al., (2015) maps, such as the Yeo components 11 and 12 with our Component 2 (see a), and the Yeo components 1 and 7 with our Cluster 3 (see b).

Tables

Table 1

Extreme words per component

Component	Top 20 Positive Extreme Words	Top 20 Negative Extreme Words
1	imageability, icon, phonet, haptic, inflect, aip, signer, sublex, themat, inanimat, nonspeech, sententi, orthographi, vowel, verb, gestur, textur, syllab, grasp, metaphor	tbss, corona, fornix, radiata, csf, peduncl, panic, cerebrospin, comt, capsul, genu, splenium, hypertens, ventricular, anisotropi, fasciculus, bdnf, fasciculi, uncin, mutat
2	tle, ictal, arcuat, interict, mni, epileptogen, tbss, presurg, peduncl, seizur, signer, talairach, corticospin, spike, deaf, postop, atlas, fasciculi, fibr, tractographi	trust, oxytocin, unfair, receipt, httlpr, aggress, phobic, appetit, ultimatum, punish, spider, monetari, phobia, incent, reapprais, methyltransferas, catechol, comt, anxious, val
3	english, orthographi, imageability, inflect, orthograph, reader, transpar, bilingu, metaphor, rhyme, spell, monolingu, alphabet, nativ, signer, liter, grammat, profici, syntact, sententi	allodynia, innocu, cold, brush, neuropath, cutan, hyperalgesia, nocicept, nonpain, noxious, trigemin, forearm, heat, thermal, sii, laser, supraspin, analgesia, emg, temperatur
4	val, comt, catechol, methyltransferas, valmet, met, dat, bdnf, enzym, polymorph, homozyg, genotyp, tandem, allel, homozygot, neurotroph, drd, carrier, httlpr, protein	cold, allodynia, innocu, brush, neuropath, hyperalgesia, thermal, trigemin, signer, noxious, heat, temperatur, nocicept, warm, cutan, combat, deaf, intranas, english, pain
5	unfair, moral, dilemma, ultimatum, intertempor, valuat, discount, amnes, beauti, belief, amnesia, ration, option, bad, fornix, choic, norm, protagonist, aesthet, certainti	comt, val, catechol, methyltransferas, valmet, met, bdnf, dat, polymorph, httlpr, enzym, homozyg, genotyp, allel, neurotroph, homozygot, drd, tandem, carrier, gene

Note. The 20 most extreme positive and negatively loading words from the five reliable components we identified. These extreme words help identify the largest sources of variance for our components.

Table 2

Clusters

Cluster	# Studies (%)	# Terms (%)	Brief description
1	1569 (14.4%)	351 (11.3%)	Knowledge Representation & Language Processing
2	2272 (20.8%)	720 (23.1%)	Development, Lifespan and Disorders
3	728 (6.7%)	347 (11.1%)	Sensation, Movement and Action
4	3159 (28.9%)	1019 (32.7%)	Cognition & Psychology
5	2927 (26.9%)	612 (19.7%)	Decision, Emotion, & Substance Use
6	243 (2.2%)	65 (2.1%)	Imaging Genetics

Note. The cluster solution indicated six clusters. The total number of studies and terms per cluster are presented. Furthermore, we provide a brief description that helps characterize the contents (i.e., studies and terms) of each cluster.

Table 3

Extreme words per cluster

Cluster	Most 20 frequent words	Least 20 frequent words
1	process, left, tempor, frontal, gyrus, inferior, word, superior, auditori, languag, semant, hemispher, middl, percept, speech, recognit, complex, read, mean, sulcus	master, focuss, fruit, innat, singular, strip, anteced, kept, learnt, mapp, posterolater, quiet, anterolater, nonmusician, orbitali, cogn, colleg, holist, inferotempor, nonlinguist
2	brain, control, imag, patient, subject, correl, use, group, compar, connect, signific, cognit, differ, chang, healthi, analysi, bilater, age, base, structur	breakdown, checklist, degen, ecnp, inadequ, custom, cut, epidemiolog, eventu, inpati, plateau, subsampl, coexist, indistinguish, mortem, nation, proof, suppli, algebra, clinician
3	cortic, motor, pain, movement, bold, hand, stimul, sensori, primari, induc, thalamus, bodi, blood, somatosensori, central, appli, evok, intens, sensorimotor, record	electromyograph, exacerb, neglig, collater, transmit, innerv, allevi, amplif, bowel, cerebrum, debilit, genicul, fingertip, first, peripheri, cyclic, subthreshold, territori, width, desynchron
4	activ, function, region, cortex, task, studi, relat, area, fmri, result, network, suggest, perform, memori, effect, visual, magnet, involv, pariet, investig	favour, host, vast, disput, expens, obscur, truli, wealth, abolish, accentu, instantan, necessit, perfect, prescrib, succeed, sudden, convolv, decay, necess, paid
5	respons, associ, neural, increas, emot, prefront, reson, particip, anterior, find, behavior, stimuli, face, individu, cingul, examin, medial, amygdala, negat, social	foster, operation, philosoph, schedul, suboptim, pituitari, problemat, assay, evoc, height, idiosyncrat, avenu, cautious, endocrin, give, salivari, deeper, laden, told, tryptophan
6	variat, genet, dopamin, gene, carrier, allel, genotyp, receptor, polymorph, dopaminerg, variant, met, dose, comt, serotonin, suscept, val, mutat, phenotyp, apo	norepinephrin, enzym, ethnic, neuromodul, tandem, composit, methyltransferas, neurotroph, methyl, uptak, catechol, glutam, monozygot, molecular, serum, commiss, apolipoprotein, neurotransmitt, asymptomat, delet

Note. The 20 most and least frequent words per cluster. The most frequent words help

characterize the subdomain within functional neuroimaging. The least frequent words also

help us interpret which words are rarely used in that cluster, which helps characterize the

types of studies not within the subdomains of functional neuroimaging.

Table 4

Terms at the cluster barycenters.

Cluster	Top 20 Words Closest to the Center of words clusters
1	decod, nonsens, class, occipitotempor, regular, ifg, heteromod, anterolater, inner, pseudo, specialis, passag, figur, embed, smg, meaningless, letter, broca, artifici, learner
2	homogen, regress, heterogen, widespread, mri, predictor, disrupt, entorhin, patient, covari, ancova, incid, statist, onset, lower, reduct, clinic, subtl, enlarg, priori
3	peripher, transmit, primari, subthreshold, secondari, contact, sharp, cyclic, tip, flow, effer, analogu, rtms, threshold, territori, reflex, genicul, anesthet, calibr, burst
4	necessari, varieti, seri, propos, step, separ, causal, allow, permit, accord, exact, driven, temporari, manner, equival, path, necessarili, assumpt, hybrid, model
5	foster, experienc, laboratori, dorsomedi, negat, tendenc, subjected, psycholog, acc, positiv, pleas, detriment, prone, diminish, take, ventromedi, introspect, style, remind, paracingul
6	gene, methyl, nucleotid, snp, transport, protein, delet, carrier, variat, genet, dopamin, glutam, uptak, composit, suscept, mutat, ethnic, variant, fmr, genom

Note. The 20 terms closest to the center of each cluster. These terms help characterize the

expected "average" or typical study within each cluster.

Table 5

Psychiatric and neurological terms

Word	Cluster	Frq	Class	C 1	C 2	C 3	C 4	C 5
adhd	2	580	psychiatric	*-1.22*	*-0.23*	0.45	0.48	0.08
autism	2	468	psychiatric	*-0.41*	*-0.12*	0.56	*-0.53*	0.21
bipolar	2	382	psychiatric	*-1.7*	*-0.04*	0.68	*-0.4*	0.02
unipolar	2	38	psychiatric	*-1.41*	*-0.61*	0.28	*-0.36*	0.07
anorexia	5	56	psychiatric	*-0.77*	*-0.74*	*-0.12*	*-0.98*	0.24
depress	2	1487	psychiatric	*-1.36*	*-0.66*	0.31	*-0.27*	*-0.17*
antidepress	2	112	psychiatric	*-1.42*	*-0.81*	*-0.12*	*-0.12*	*-0.33*
obsess	2	113	psychiatric	*-1.5*	*-0.26*	0.19	*-0.17*	0.3
schizophrenia	2	2250	psychiatric	*-1.12*	0.17	0.58	*-0.01*	0.02
posttraumat	5	72	psychiatric	*-1.3*	*-0.99*	0.21	*-1.11*	0.01
traumat	2	151	psychiatric	*-1.24*	*-0.42*	0.22	*-0.78*	0.18
alzheim	2	409	neurological	*-1.28*	0.67	0.52	0.72	0.28
dementia	2	250	neurological	*-1.27*	0.76	0.76	0.29	0.31
frontotempor	2	131	neurological	*-0.76*	0.63	0.8	*-0.27*	*-0.12*
dystonia	6	62	neurological	*-1.05*	0.75	*-0.61*	1.01	*-1.64*
parkinson	2	232	neurological	*-0.88*	0.47	*-0.42*	0.64	0.04
huntington	2	29	neurological	*-1.15*	0.33	0.03	0.22	*-0.13*
epilepsi	2	277	neurological	*-1.21*	1.35	*-0.3*	0.33	0.76

Note. Psychiatric vs. neurological terms (as defined by Crossley et al., 2015) within our components solution. Negative loadings are *italicized*. Components 2 and 4 show a clear dissociation between psychiatric and neurological disorders within the semantic space of Neurosynth.

References

Abdi, H. & Béra, M., 2014. Correspondence Analysis. In *Encyclopedia of Social Network Analysis and Mining*. New York, NY: Springer New York, pp. 275–284. Available at: http://oxfordhandbooks.com/view/10.1093/oxfordhb/9780199934898.001.0001/oxfordhb-9780199934898-e-007.

Abraham, A. et al., 2014. Machine learning for neuroimaging with scikit-learn. *Frontiers in neuroinformatics*, 8(February), p.14. Available at: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3930868&tool=pmcentrez&rendertype=abstract.

Ailem, M. et al., 2016. Unsupervised text mining for assessing and augmenting GWAS results. *Journal of Biomedical Informatics*, 60(February), pp.252–259.

Amanzio, M. et al., 2013. Activation likelihood estimation meta-analysis of brain correlates of placebo analgesia in human experimental pain. *Human Brain Mapping*, 34(3), pp.738–752.

Bartra, O., McGuire, J.T. & Kable, J.W., 2013. The valuation system: A coordinate-based meta-analysis of BOLD fMRI experiments examining neural correlates of subjective value. *NeuroImage*, 76, pp.412–427. Available at: http://dx.doi.org/10.1016/j.neuroimage.2013.02.063.

Beaton, D., Fatt, C.R.C. & Abdi, H., 2014. An ExPosition of multivariate analysis with the singular value decomposition in R. *Computational Statistics & Data Analysis*, 72, pp.176–189. Available at: http://dx.doi.org/10.1016/j.csda.2013.11.006%5Cnpapers3://publication/doi/10.1016/j.csda.2013.11.006.

Benzecri, J.P., 1976. History and prehistory of data analysis. III: Era piscatoria (French). *Les*

Cahiers de l'Analyse des Données, 1, pp.221–241.

Binder, J.R. et al., 2009. Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. *Cerebral Cortex*, 19(12), pp.2767–2796.

Bookheimer, S., 2002. Functional MRT of Language: New Approaches to Understanding the Cortical Organization of Semantic Processing. *Annual review of neuroscience*, 25, pp.151–188.

Buhle, J.T. et al., 2014. Cognitive reappraisal of emotion: A meta-analysis of human neuroimaging studies. *Cerebral Cortex*, 24(11), pp.2981–2990.

Calinski, T. & Harabasz, J., 1974. A dendrite method for cluster analysis. *Communications in Statistics - Theory and Methods*, 3(1), pp.1–27. Available at: http://www.tandfonline.com/doi/abs/10.1080/03610927408827101.

Chawla, D., 2017. Great paper? Swipe right on the new "Tinder for preprints" app. *Science/AAAS*. Available at: https://www.sciencemag.org/news/2017/06/great-paper-swipe-right-new-tinder-preprints-app.

Churchill, N.W. et al., 2012. Optimizing preprocessing and analysis pipelines for single-subject fMRI. I. Standard temporal motion and physiological noise correction methods. *Human Brain Mapping*, 33(3), pp.609–627.

Cioli, C. et al., 2014. Differences in human cortical gene expression match the temporal properties of large-scale functional networks. *PLoS ONE*, 9(12), pp.1–28.

Crossley, N.A. et al., 2015. Neuroimaging distinction between neurological and psychiatric disorders. *British Journal of Psychiatry*, 207(5), pp.429–434.

Desgraupes, B., 2015. clusterCrit: Clustering Indices. *R package version 1.2.6*. Available at: http://cran.r-project.org/package=clusterCrit.

Escofier-Cordier, B., 1965. L'Analyse des Correspondences. *Unver{ö}ffentlichte Dissertation: Universit{é} de Rennes.*

Van Essen, D. et al., 2009. Mining the neuroimaging literature using the SumsDB database:

stereotaxic coordinates and more! *Frontiers in Neuroinformatics*. Available at:

http://www.frontiersin.org/10.3389/conf.neuro.11.2009.08.021/event_abstract.

Etkin, A. & Wager, T.D., 2007. Functional Neuroimaging of Anxiety: A Meta-Analysis of

Emotional Processing in PTSD, Social Anxiety Disorder, and Specific Phobia. *American

Journal of Psychiatry*, 164(10), pp.1476–1488. Available at:

http://psychiatryonline.org/doi/abs/10.1176/appi.ajp.2007.07030504.

Fedorenko, E. & Thompson-Schill, S.L., 2014. Reworking the language network. *Trends in

Cognitive Sciences*, 18(3), pp.120–127. Available at:

http://dx.doi.org/10.1016/j.tics.2013.12.006.

Feinerer, I., 2011. Introduction to the tm Package Text Mining in R. *R vignette*, pp.1–8.

Fox, A. et al., 2014. Bridging psychology and genetics using large-scale spatial analysis of

neuroimaging and neurogenetic data. *bioRxiv*. Available at:

http://biorxiv.org/content/early/2014/12/09/012310.

Fox, P.T. et al., 2014. Meta-Analysis in Human Neuroimaging: Computational Modeling of

Large-Scale Databases. *Annual Review of Neuroscience*, 37(1), pp.409–434. Available

at: http://www.annualreviews.org/doi/abs/10.1146/annurev-neuro-062012-170320.

Friebel, U., Eickhoff, S.B. & Lotze, M., 2011. Coordinate-based meta-analysis of

experimentally induced and chronic persistent neuropathic pain. *NeuroImage*, 58(4),

pp.1070–1080. Available at: http://dx.doi.org/10.1016/j.neuroimage.2011.07.022.

Glasser, M.F. et al., 2016. A multi-modal parcellation of human cerebral cortex. *Nature*,

536(7615), pp.171–8. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/27437579%5Cnhttp://www.pubmedcentral.nih.go

v/articlerender.fcgi?artid=PMC4990127.

Gorgolewski, K. et al., 2014. Tight fitting genes: Finding relations between statistical maps

and gene expression patterns. *Organization for Human Brain Mapping. Hamburg, Germany*.

Gorgolewski, K.J. et al., 2016. NeuroVault.org: A repository for sharing unthresholded statistical maps, parcellations, and atlases of the human brain. *NeuroImage*, 124, pp.1242–1244.

Greenacre, M., 2017. *Correspondence analysis in practice*, CRC press.

Greenacre, M.J., 1984. *Theory and Applications of Correspondence Analysis*, Academic Press. Available at: https://books.google.com/books?id=LsPaAAAAMAAJ.

Jennings, R.G. & Van Horn, J.D., 2012. Publication bias in neuroimaging research: Implications for meta-analyses. *Neuroinformatics*, 10(1), pp.67–80.

de la Vega, A. et al., 2016. Large-Scale Meta-Analysis of Human Medial Frontal Cortex Reveals Tripartite Functional Organization. *The Journal of Neuroscience*, 36(24), pp.6553–6562. Available at: http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.4402-15.2016.

Laird, A.R. et al., 2011. Behavioral Interpretations of Intrinsic Connectivity Networks. *Journal of Cognitive Neuroscience*, 23(12), pp.4022–4037.

Laird, A.R., Lancaster, J.L. & Fox, P.T., 2005. BrainMap: the social evolution of a human brain mapping database. *Neuroinformatics*, 3(1), pp.65–78. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15897617.

Lebart, L., Morineau, A. & Warwick, K.M., 1984. Multivariate descriptive statistical analysis; correspondence analysis and related techniques for large matrices.

Lebart, L.L. et al., 1998. Exploring Textual Data. *Computational Linguistics*, 4, pp.165–166. Available at: http://link.springer.com/10.1007/978-94-017-1525-6.

Leek, J., 2017. Paper app. Available at: https://github.com/jtleek/papr [Accessed June 21, 2017].

Lindquist, K.A., 2010. The brain basis of emotion: A meta-analytic review. *Dissertation Abstracts International, B: Sciences and Engineering*, 71(4), p.2744.

Mesmoudi, S. et al., 2015. LinkRbrain: Multi-scale data integrator of the brain. *Journal of Neuroscience Methods*, 241, pp.44–52. Available at: http://dx.doi.org/10.1016/j.jneumeth.2014.12.008.

Mesmoudi, S. et al., 2013. Resting State Networks' Corticotopy: The Dual Intertwined Rings Architecture. *PLoS ONE*, 8(7).

Murtagh, F. & Legendre, P., 2014. Ward's Hierarchical Clustering Method: Clustering Criterion and Agglomerative Algorithm. *Journal of Classification*, 31(October), pp.274–295. Available at: http://arxiv.org/abs/1111.6285.

Nichols, T.E. et al., 2016. Best Practices in Data Analysis and Sharing in Neuroimaging using MRI. *bioRxiv*, p.54262. Available at: http://biorxiv.org/lookup/doi/10.1101/054262.

Nielsen, F.Å.A., 2003. The Brede database: a small database for functional neuroimaging. In *Neuroimage*. pp. 2000–2001.

Nielsen, F.A., Hansen, L.K. & Balslev, D., 2004. Mining for associations between text and brain activation in a functional neuroimaging database. *Neuroinformatics*, 2(4), pp.369–80. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15800369.

Perini, I., Bergstrand, S. & Morrison, I., 2013. Where Pain Meets Action in the Human Brain. *Journal of Neuroscience*, 33(40), pp.15930–15939. Available at: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.3135-12.2013.

Phan, K.L. et al., 2002. Functional neuroanatomy of emotion: a meta-analysis of emotion activation studies in PET and fMRI. *NeuroImage*, 16(2), pp.331–48. Available at: http://www.ncbi.nlm.nih.gov/pubmed/12030820.

Poldrack, R.A. et al., 2012. Discovering Relations Between Mind, Brain, and Mental

Disorders Using Topic Mapping. *PLoS Computational Biology*, 8(10), p.e1002707.

Available at: http://dx.plos.org/10.1371/journal.pcbi.1002707.

Poldrack, R.A. et al., 2011. The cognitive atlas: toward a knowledge foundation for cognitive

neuroscience. *Frontiers in neuroinformatics*, 5(September), pp.1–11. Available at:

http://journal.frontiersin.org/article/10.3389/fninf.2011.00017/abstract.

Poldrack, R.A. & Yarkoni, T., 2016. From Brain Maps to Cognitive Ontologies : Informatics

and the Search for Mental Structure. *Annual Review of Psychology*, 67, pp.1–26.

Power, J.D. et al., 2011. Functional network organization of the human brain. *Neuron*, 72(4),

pp.665–78. Available at:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3222858&tool=pmcentrez&

rendertype=abstract [Accessed July 15, 2014].

Price, C.J., 2012. A review and synthesis of the first 20years of PET and fMRI studies of

heard speech, spoken language and reading. *NeuroImage*, 62(2), pp.816–847. Available

at: http://dx.doi.org/10.1016/j.neuroimage.2012.04.062.

Price, C.J., 2010. The anatomy of language: A review of 100 fMRI studies published in 2009.

Annals of the New York Academy of Sciences, 1191, pp.62–88.

R Core Team, 2014. R: a language and environment for statistical computing. *R: A Language

and Environment for Statistical Computing*, 55, p.http://www.R-project.org/. Available

at: http://www.mendeley.com/research/r-language-environment-statistical-computing-

96/%5Cnpapers2://publication/uuid/A1207DAB-22D3-4A04-82FB-D4DD5AD57C28.

Rizzo, G. et al., 2016. Menga: A new comprehensive tool for the integration of neuroimaging

data and the allen human brain transcriptome atlas. *PLoS ONE*, 11(2), pp.1–20.

Rubin, T.N. et al., 2016. *Decoding brain activity using a large-scale probabilistic functional-

anatomical atlas of human cognition*, Available at:

http://biorxiv.org/content/early/2016/06/18/059618.abstract.

Salimi-Khorshidi, G. et al., 2009. Meta-analysis of neuroimaging data: A comparison of image-based and coordinate-based pooling of studies. *NeuroImage*, 45(3), pp.810–823. Available at: http://dx.doi.org/10.1016/j.neuroimage.2008.12.039.

Schomers, M.R. & Pulvermüller, F., 2016. Is the sensorimotor cortex relevant for speech perception and understanding? An integrative review. *Frontiers in Human Neuroscience*, 10(September).

Smith, S.M. et al., 2009. Correspondence of the brain' s functional architecture during activation and rest. *Proceedings of the National Academy of Sciences of the United States of America*, 106, pp.13040–13045.

Strother, S.C. et al., 2002. The quantitative evaluation of functional neuroimaging experiments: the NPAIRS data analysis framework. *NeuroImage*, 15(4), pp.747–771.

Toro, R., Fox, P.T. & Paus, T., 2008. Functional coactivation map of the human brain. *Cerebral Cortex*, 18(11), pp.2553–2559.

Vierck, C.J. et al., 2013. Role of primary somatosensory cortex in the coding of pain. *Pain*, 154(3), pp.334–344. Available at: http://dx.doi.org/10.1016/j.pain.2012.10.021.

Wager, T.D. et al., 2009. Evaluating the consistency and specificity of neuroimaging data using meta-analysis. *NeuroImage*, 45(1 Suppl), pp.S210–S221. Available at: http://dx.doi.org/10.1016/j.neuroimage.2008.10.061.

Wager, T.D., Lindquist, M. & Kaplan, L., 2007. Meta-analysis of functional neuroimaging data: Current and future directions. *Social Cognitive and Affective Neuroscience*, 2(2), pp.150–158.

White, P.D., Rickards, H. & Zeman, a. Z.J., 2012. Time to end the distinction between mental and neurological illnesses. *BMJ*, 344(may24 1), pp.e3454–e3454. Available at: http://www.bmj.com/cgi/doi/10.1136/bmj.e3454.

Wickham, H., 2009. *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag New

York. Available at: http://link.springer.com/10.1007/978-0-387-98141-3%5Cnpapers3://publication/doi/10.1007/978-0-387-98141-3.

Yang, Y. et al., 2016. Identifying functional subdivisions in the human brain using meta-analytic activation modeling-based parcellation. *NeuroImage*, 124, pp.300–309.

Yarkoni, T. et al., 2011. Large-scale automated synthesis of human functional neuroimaging data. *Nature methods*, 8(8), pp.665–70. Available at: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3146590&tool=pmcentrez&rendertype=abstract.

Yeo, B.T. et al., 2011. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol*, 106, pp.1125–1165. Available at: http://www.ncbi.nlm.nih.gov/pubmed/21653723.

Yeo, T. et al., 2015. Functional Specialization and Flexibility in Human Association Cortex. *Cerebral Cortex*, 25(10), pp.3654–3672. Available at: http://cercor.oxfordjournals.org/cgi/doi/10.1093/cercor/bhu217.

Yuan, R. et al., 2017. MAPBOT Meta-Analytic Parcellation Based On Text, and its application to the human thalamus. *NeuroImage*. Available at: http://linkinghub.elsevier.com/retrieve/pii/S1053811917305037.