A systematic survey of regional multitaxon biodiversity: evaluating strategies and coverage

Ane Kirstine Brunbjerg[1*], Hans Henrik Bruun[2], Lars Brøndum[3], Aimée T. Classen[4,5], Lars Dalby[1], Kåre Fog[6], Tobias G. Frøslev[7], Irina Goldberg[2], Anders Johannes Hansen[8], Morten D.D. Hansen[3], Toke T. Høye[1,9], Anders A. Illum[5], Thomas Læssøe[2], Gregory S. Newman[5], Lars Skipper[3], Ulrik Søchting[2], Rasmus Ejrnæs[1]

[1] *Section for Biodiversity & Conservation, Department of Bioscience, Aarhus University, DK-8410 Rønde, Denmark (present address)*

[2] *Section for Ecology and Evolution, Department of Biology, University of Copenhagen, DK-2100 Copenhagen, Denmark / Natural History Museum of Denmark, Universitetsparken 15, University of Copenhagen, DK-2100 Copenhagen, Denmark*

[3] *Natural History Museum Aarhus, DK-8000 Aarhus C, Denmark*

[4] *Rubenstein School of Environment and Natural Resources, University of Vermont, Burlington, VT 05405, USA,*

[5] *Center for Macroecology, Evolution and Climate, Natural History Museum of Denmark, University of Copenhagen, DK-1350 Copenhagen, Denmark.*

[6] *Hesselholm 107, DK-3670 Veksø, Denmark*

[7] *Center for GeoGenetics, University of Copenhagen, DK-2100 Copenhagen, Denmark*

[8] *Natural History Museum of Denmark, University of Copenhagen, DK-1153 Copenhagen, Denmark*

[9] *Arctic Research Centre, Aarhus University, Ny Munkegade 114, building 1540, DK-8000 Aarhus C, Denmark*

* Corresponding author: Tel.: +45 87158865, E-mail address: akb@bios.au.dk

<u>E-mail-addresses (co-authors)</u>: AKB: akb@bios.au.dk, HHB: hhbruun@bio.ku.dk, LB:

lars@nathist.dk, ATC: atclassen@gmail.com, LD: lars@bios.au.dk, KF: kaarefog@teliamail.dk,

TGF: tobiasgf@bio.ku.dk, IG: irina.goldberg@gmail.com, AJH: ajhansen@bio.ku.dk , MDDH:

mortendd@molslab.dk, TTH: tth@bios.au.dk, AAI: andersaillum@snm.ku.dk, TL:

thomsal@bio.ku.dk, GSN: gsnewm@gmail.com, LS: lars.skipper@get2net.dk, US:

ulriks@bio.ku.dk, RE: rasmus@bios.au.dk

Abstract

Background: In light of the biodiversity crisis and our limited ability to explain variation in biodiversity, tools to quantify spatial and temporal variation in biodiversity and its underlying drivers are critically needed. Inspired by the recently published ecospace framework, we developed and tested a protocol for environmental and biotic mapping that is scalable to habitats, ecosystems and biomes. We selected study sites (40×40m) across Denmark using stratified random sampling along the major environmental gradients underlying biotic variation. Using standardized methods, we collected site species data on vascular plants, bryophytes, macrofungi, lichens, gastropods and arthropods. To evaluate sampling efficiency, we calculated regional coverage (relative to the known species number per taxonomic group), and site scale coverage (i.e., sample completeness per taxonomic group at each site). To extend taxonomic coverage to organisms that are difficult to sample by classical inventories (e.g., nematodes and non-fruiting fungi), we collected soil for metabarcoding. Finally, to assess site conditions, we mapped abiotic conditions, biotic resources and habitat continuity.

Results: Despite the 130 study sites only covering a minute fraction (0.0005 %) of the total Danish terrestrial area, we found 1774 species of macrofungi (54% of the Danish fungal species pool), 663 vascular plant species (42%), 254 bryophyte species (41%) and 200 lichen species (19%). For arthropods, we observed 330 spider species (58%), 123 carabid beetle species (37%) and 99 hoverfly species (33%). Overall, sample coverage was remarkably high across taxonomic groups and sufficient to capture substantial spatial variation in biodiversity across Denmark. This inventory is unprecedented in detail and resulted in the discovery of 143 species with no previous record for Denmark. Comparison between plant OTUs detected in soil DNA and observed plant species confirmed the usefulness of carefully curated environmental DNA-data.

Species richness did not correlate well among taxa suggesting differential and complex biotic

responses to environmental variation.

Conclusions: We successfully and adequately sampled a wide range of mega-diverse taxa along

key environmental gradients across a large region using an approach that includes multi-taxon

biodiversity assessment and ecospace mapping. Our approach is applicable to assessments of

biodiversity in other regions and biomes.

Keywords: Abiotic gradients, Biotic factors, Continuity, Denmark, Disturbance, eDNA,

Moisture, Productivity

Background

The vast number of species on Earth have yet to be described, challenging our understanding of biodiversity [1]. For a deeper understanding of what determines the distribution of species across the planet, comprehensive data on species occurrence and environmental conditions are required. While some progress has been made in understanding the distribution of biodiversity at coarse spatial resolution, our knowledge of biodiversity at high spatial resolution is deficient [2]. In this study, we consider biodiversity as the richness and turnover of taxonomic units, whether species or operational taxonomic units (OTUs) derived by eDNA (environmental DNA) metabarcoding. While progress has been made in the interpretation and prediction of richness and turnover of vascular plants and vertebrates, various types of bias, e.g. temporal, spatial, and taxonomic bias [3], have constrained similar advances for less well-known, but mega-diverse groups such as fungi and insects [1]. As a result, conservation management is typically based on biodiversity data from a non-random subset of taxa [4].

Recent developments in molecular techniques – in particular the extraction and sequencing of eDNA – hold the promise of more time-efficient sampling and identification of species [5, 6]. Further, eDNA enables the exploration of communities and organisms not easily recorded by traditional biodiversity assessment, such as soil-dwelling nematodes [7]. In fact, PCR-based methods combined with DNA sequencing have already provided valuable insight into the taxonomic diversity within complex environmental samples, such as soil [8-10] and water [e.g. 11, 12]. Due to the ongoing rapid development in DNA sequencing technologies, with the emergence of next generation sequencing (NGS) techniques – generating billions of DNA sequences [13] – an environmental sample could now be analyzed to a molecular depth that gives an almost exhaustive picture of the species composition at the site of collection. Despite

this potential, rigorous assessments with complete taxonomic coverage from eDNA samples are

still missing [6, 12, 14]. To assess the suitability and potential of eDNA data in complementing –

or even replacing – traditional field survey data, tests on comprehensive data sets are needed.

Undertaking an ambitious biodiversity field study across a wide geographical space comes

with major logistical and methodological challenges. It is not clear what environmental gradients

structure biodiversity across the tree of life and for most taxa standardized field protocols to

sample species occurrences are non-existent. The recently developed ecospace framework

suggests that biodiversity varies in relation to its position along environmental gradients

(position), the availability of biotic resources, such as organic matter and structures e.g. trees for

epiphytes (expansion), and spatio-temporal extent of biotopes (continuity) [15]. Environmental

conditions and local processes can be a template shaping local biodiversity (e.g. through

environmental filtering) [16, 17]. This template is highlighted by the ecospace position of

sampled biotopes in abiotic environmental space. In addition to the physico-chemical conditions

shaping abiotic gradients – particularly important to autotrophic organisms – the presence and

abundance of specific biological resources, crucial to heterotrophic organisms, such as specialist

herbivores, detritivores and saproxylic species are likely important and thus should be considered

[18]. The quantification of biotic resources and structures, e.g. dead wood, dung and carcasses, is

not often included in community studies, despite the limited knowledge in the area [15, 19]

speaks for further studies. Spatial and temporal processes at regional extent, such as extinction,

speciation and migration, shape species pools and thereby set the limits to local richness and

species composition [16, 17, 20]. In order to improve our understanding of biodiversity patterns,

local and regional factors should be considered concurrently [17, 21].

In this study, we used the ecospace framework to develop a comprehensive protocol for

large-scale mapping of variation in biodiversity and to evaluate the efficacy of ecospace in

capturing environmental variation across Denmark. The protocol development and evaluation

was carried out as part of a research project (called *Biowide*). The project aimed to cover all of

the major environmental gradients, including natural variation in moisture, soil fertility and

succession, as well as habitats under cultivation. Within this environmental space, we performed

a systematic and comprehensive sampling of the environment and biodiversity. We combined

traditional species observation and identification with modern methods of biodiversity mapping

in the form of massive parallel sequencing of eDNA extracted from soil samples.

Methods

Study area and site selection

We aimed to characterize biodiversity across the country of Denmark (Fig. 1a) – a lowland area

of 42,934 km^2 and an elevational range of 0-200 meters above sea level. While there are some

limestone and chalk outcrops, there is no exposed bedrock in the investigated area. Soil texture

ranges from coarse sands to heavy clay and organic soils of various origins [22]. Land use is

dominated by arable land (61 %), most of which is in annual rotation, while forests are mostly

plantations established during the 19th and 20th centuries. Scrubs cover approximately 17 %,

natural and semi-natural terrestrial habitats some 10 %, and freshwater lakes and streams 2 %.

The remaining 10 % is made up of urban areas and infrastructure [23, 24].

When selecting sites, we considered major environmental gradients, the potential size of the

sampling units (sites), as well as practicalities of sampling across the large geographical space

within the same season. The sites were 40×40 m which was a compromise between within-site

homogeneity and the representativeness of a particular habitat type. We stratified site selection

according to the identified major environmental gradients, including the intensity of human land

use. We measured 30 sites that were cultivated habitats and 100 sites that were natural and semi-

natural habitats. This balance between natural and cultivated habitat was chosen, because we

expected cultivated habitats to have shorter environmental gradients. The cultivated subset

represented major land-use categories and the natural subset was stratified across natural

gradients in soil fertility, soil moisture, and successional stage from sparsely vegetated to closed

canopy forest, (Appendix A). We deliberately excluded linear features, such as hedgerows and

road verges, urban areas with predominantly exotic plants as well as saline and aquatic habitats,

but included temporarily inundated heath, dune depressions and wet mires.

The final set of 25 sampling classes consisted of six cultivated habitat types; three types of

fields (rotational, leys, and oldfield) and three types of plantations (beech, oak, and spruce). 18

natural classes consisted of all factorial combinations of natural soil fertility (fertile or infertile),

moisture (dry, moist, or wet), and successional stage (low vegetation with bare soil, closed

herb/scrub, or forest) (Appendix A). Finally, we included a class of perceived hotspots for

species richness [25] in Denmark. These sites were selected subjectively by performing a public

poll among active natural history volunteers in the Danish nature conservation and nature

management societies. The 25 classes were replicated in each of five geographical regions within

Denmark (Fig. 1a). The result was 130 sites with 18 natural, 6 cultivated, and two perceived

species richness hotspot sites evenly distributed across each of five geographic regions of

Denmark (Table 1). For logistical reasons, we did not place any sites on Bornholm although we

acknowledge that this island is geologically different than the rest of Denmark.

For the 18 natural habitat classes, site selection through stratified random sampling was

guided by a large nation-wide dataset of vegetation plots in semi-natural habitats distributed

across the entire country (n = 96,400 plots of 78.5 m^2 each, www.naturdata.dk) from a national

monitoring and mapping project [26] and in accordance with the EU Habitats Directive [27]. We

used environmental conditions computed from plant indicator values to select candidate sites for

each class. First, we calculated plot mean values for Ellenberg indicator values based on vascular

plants species lists [28] and Grime CSR-strategy allocations of recorded plants [29], the latter

were recoded to numeric values following Ejrnæs & Bruun [30]. We excluded saline and

artificially fertilized habitats by excluding plots with Ellenberg S > 1 or Ellenberg N > 6. We

then defined stratification categories as: fertile (Ellenberg N 3.5-6.0), infertile (Ellenberg N <

3.5), dry (Ellenberg F < 5.5), moist (Ellenberg F 5.5-7.0), wet (Ellenberg F > 7.0), early

succession (Grime R > 4 and Ellenberg L > 7 or > 10 % of annual plants), late succession

(mapped as forest), mid succession (remaining sites).

To reduce transport time and costs, all 26 sites within each region were grouped into three

geographic clusters (Fig. 1a). The nested sampling design allowed us to take spatially structured

species distributions into account [31]. The procedure for site selection involved the following

steps:

1) Designation of three geographic clusters within each region with the aim to cover all

natural classes while a) keeping the cluster area below 200 km^2 and b) ensuring high

between-cluster dispersion in order to represent the geographic range of the region. In

practice, species richness hotspots were chosen first, then clusters were placed with

reference to the highest ranking hotspots and in areas with a wide range of classes

represented in the national vegetation plot data [32].

2) Representing the remaining 24 classes in each region by selecting 8-9 potential sites in

each cluster. Sites representing natural classes were selected from vegetation plot data.

Cultivated classes were assumed omnipresent and used as buffers in the process of

completing the non-trivial task of finding all classes within each of three cluster areas of

< 200 km^2 in each region.

3) Negotiating with land owners and, in case of disagreement, replacing the preferred site

with an alternative site from the same class.

After each of the 130 sites were selected using available data, we established each 40×40 m

site in a subjectively selected homogenous area that accounted for topography and vegetation

structure. Each site was divided into four 20×20 m quadrants, and from the center of each

quadrant a 5 m radius circle (called a plot) was used as a sub-unit for data collection to

supplement the data collected at site level (40 × 40 m) (Fig. 1b).

Collection of biodiversity data

For each of the 130 sites, we aimed at making an unbiased and representative assessment of

multi-taxon species richness. Data on vascular plants, bryophytes, lichens, macrofungi,

arthropods and gastropods were collected using standard field inventory methods (Appendix B).

For vascular plants, bryophytes and gastropods, we collected exhaustive species lists. For the

remaining taxonomic groups that are more demanding to find, catch, and identify, we aimed at

collecting a reproducible and unbiased sample through a standardized level of effort (typically

one hour). Multiple substrates (soil, herbaceous debris, wood, stone surfaces and bark of trees up

to 2 m) were carefully searched for lichens and macrofungi at each site. For fungi, we visited

each site twice during the main fruiting season in 2014 – in August and early November – and

once during the main fruiting season in 2015 – between late August and early October.

Specimens that were not possible to identify with certainty in the field were sampled and, when

possible, identified in the laboratory. For arthropod sampling, a standard set of pitfall traps

(including meat-baited and dung-baited traps), yellow Möricke pan traps and Malaise traps were

operated during a fixed period of the year. In addition, we used active search and collection

methods, including sweep netting and beating as well as expert searches for plant gallers, miners

and gastropods. Finally, we heat-extracted collembolas and oribatid mites from soil cores. Due to

the limited size of the sites relative to the mobility of mammals, birds, reptiles and amphibians,

data on these groups were not recorded. Records of arthropods were entered in https://www.

naturbasen.dk. Records of fungi were entered in https://svampe.databasen.org/. All species

occurrence data and environmental data has been made available at the project home page

http://bios.au.dk/om-instituttet/organisation/biodiversitet/projekter/biowide/. Species data will be

made available for GBIF (www.gbif.org) through the above-mentioned web portals. Specimens

are stored at the Natural History Museum Aarhus (fungal specimens at the fungarium at the

Natural History Museum of Denmark). For further details on the methods used for collection of

biodiversity data see Appendix B.

Collection of eDNA data

We used soil samples collected from all 130 sites for the eDNA inventory. At each site, we

sampled 81 soil cores in a 9×9 grid covering the entire 40 m $\times$ 40 m plot and pooled the

collected samples after removal of coarse litter. We homogenized the soil by mixing with a

mixing paddle mounted on a drilling machine. A subsample of soil was sampled from the

homogenized sample and DNA was extracted for marker gene amplification and sequencing

[14]. We chose the MiSeq platform by Illumina for DNA sequencing. MiSeq is adapted to

amplicon sequencing [33]. For further details on methods for eDNA data generation and

considerations on eDNA species richness and community composition measures see Appendix

B.

Site environmental data

223 We have followed the suggestion in Brunbjerg et al. [15] to describe the fundamental

224 requirements for biodiversity in terms of the ecospace (position, expansion and spatio-temporal

225 continuity of the biotope).

Position

To assess the environmental variation across the 130 sites, we measured a core set of site factors

that described the abiotic conditions at each site. Environmental recordings and estimates

included soil pH, total soil carbon (C, g/m^2), total soil nitrogen (N, g/m^2) and total soil

phosphorus (P, g/m^2), soil moisture (% volumetric water content), leaf CNP (%), soil surface

temperature (°C) and humidity (vapour pressure deficit), air temperature (°C), light intensity

(Lux), and boulder density. For further details on methods used to collect abiotic data see

Appendix B.

Expansion

We collected measurements that represent the expansion or biotic resources which some species

consume and the organic and inorganic structures which some species use as habitat. Although

many invertebrates are associated with other animals, for practical reasons, we restricted our

quantification of biotic resources to the variation in live and dead plant tissue, including dung.

We measured litter mass (g/m^2), plant species richness, vegetation height (of herb layer, cm),

cover of bare soil (%), bryophyte cover (%) and lichen cover (%), dead wood volume (m^3/site),

dominant herbs, the abundance of woody species, the number of woody plant individuals, flower

density (basic distance abundance estimate, [34]), density of dung (basic distance abundance

estimate), number of carcasses, fine woody debris density (basic distance abundance estimate),

ant nest density (basic distance abundance estimate), and water puddle density (basic distance

abundance estimate). For further details on methods used to collect expansion data see Appendix

B.

Mapping of temporal and spatial continuity

For each site, we inspected a temporal sequence of aerial photos (from 1945 to 2014) and

historical maps (1842-1945) starting with the most recent photo taken. We defined temporal

continuity as the number of years since the most recent major documented land use change. The

year in which a change was identified was recorded as a 'break in continuity'. To estimate spatial

continuity, we used ArcGIS to construct four buffers for each site (500 m, 1000 m, 2000 m, 5000

m). Within each buffer we estimated the amount of habitat similar to the site focal habitat by

visual inspection of aerial photos with overlays representing nation-wide mapping of semi-

natural habitat. For further details on methods for collection of continuity data see Appendix B.

Analyses

To illustrate the coverage of the three main gradients (moisture, fertility, and successional stage)

spanned by the 130 sites, Ellenberg mean site values (mean of mean Ellenberg values for the

four 5m radius quadrats within each site) for soil moisture (Ellenberg F), soil nutrients

(Ellenberg N) and light conditions (Ellenberg L) were plotted relative to Ellenberg F, N and L

values for a reference data set of 5 m radius vegetation quadrats (47202 from agricultural, semi-

natural and natural open vegetation and 12014 from forests (www.naturdata.dk) [26]. Mean

Ellenberg values were only calculated for quadrats with more than five species and 95 percentile

convex hull polygons where drawn for the reference data set as well as the Biowide data set.

We assessed the coverage for each taxonomic group across sites as well as within each site for

spiders, harvestmen, and insect orders represented by at minimum of 75 species, for which we

had abundance data by comparing the number of species found to the estimated species richness

of the sample using rarefaction in the iNEXT R-package [35].

To further evaluate the turnover component of biodiversity and how well we covered the

environmental gradient for our inventory, we related community composition to the measured

environmental variables (abiotic and biotic) based on a Nonmetric Multidimensional Scaling

(NMDS) analyses in R v. 3.2.3 [36] using the vegan R-package [37] and the plant species $\times$ site

matrix as well as the macrofungi species $\times$ site matrix. Abiotic and biotic variables were

correlated with ordination axes to facilitate interpretation.

To illustrate and substantiate the adequacy of the eDNA sampling protocol and subsequent

laboratory protocols, we correlated basic biodiversity measures of community composition

(NMDS axes) and richness for plant eDNA (ITS2 marker region) with the same measures for our

observed plant data (see Appendix B for detailed methods).

Results

Results

The 130 sites were distributed in 15 clusters nested within five regions across Denmark (Fig. 1a).

The measured variables differed according to the initial stratification of sites based on simple

indicators (Table 1, Fig. 2a, b, ranges of measured variables in Appendix C). Managed sites

(plantations and agricultural fields) revealed little variation in soil moisture (Fig. 2b). The

species richness hotspots spanned the full variation of natural sites regarding fertility, moisture

and successional stage (Fig. 2b).

The selected 130 sites covered the main gradients reflected by a huge reference dataset from a

national monitoring program (Fig. 3) as judged from a vegetation-based calibration of site

conditions regarding moisture, fertility and succession (light intensity). Biowide data seemed to

increase the upper range of the fertility gradient, which can be explained by the inclusion in

Biowide of rotational fields that were not included in reference data (Fig. 2b, 3).

The environmental expansion of ecospace, which was measured as the amount and

differentiation of organic carbon sources, varied among habitat types with high litter mass in tree

plantations and late successional habitats, high plant species richness in early and mid-

successional habitats, high dung density in open habitats (early successional and fields) and high

amounts of dead wood in late successional habitats (Fig. 4). Spatial and temporal continuity

varied for the 130 sites with less spatial continuity at larger buffer sizes (Appendix D). The

number of species found per site differed with taxonomic group with the highest number for

macrofungi and lowest for bryophytes and lichens (Appendix E).

We collected 1774 species of macrofungi (corresponding to 54 % of the number of

macrofungi recorded in Denmark), 200 lichens (19 %), 663 vascular plants (42 %) and 254

bryophytes (41 %) during the study period. We collected 75 species of gastropods (75 %), 330

spiders (58 %), 99 hoverflies (33 %), 123 carabid beetles (37 %) and 203 gallers and miners

species (21 %). For all groups except macrofungi, the number of species found was higher in

natural (n = 90) than in cultivated (n = 40) sites, but across taxonomic groups, plantations and

agricultural fields harbored unique species – plantations were particularly important in harboring

unique species of macrofungi (Table 2). The taxonomic sample coverage calculated by

rarefaction within the 130 sites was high overall (range: 0.86-0.99), but highest for gastropods

and spiders and lowest for gallers and miners (Table 2).

Data from the fungal eDNA community matrix was mapped to the Darwin Core data standard

(http://rs.tdwg.org/dwc/) and wrapped in a DwC archive for publication to the Global

Biodiversity Information Facility. The 'dataGeneralizations' field was used to indicate the

identity of OTUs towards the UNITE species hypothesis concept [38], Sampling sites were

included as WKT polygons in the 'footprintWKT' field and sampling site names were included in

the 'eventID' field. The representative sequences (OTUs) were included using the GGBN

amplification extension. The dataset is available from gbif.org (https://doi.org/10.15468/nesbvx)

The inventory was unprecedented in detail for Denmark and possibly for any region of the

same size globally and resulted in a total of 110 new macrofungi, 1 new lichen and 32 new

invertebrate species (of which 12 were gallers and miners and 3 spiders) that had not previously

been documented in Denmark (Table 2).

Turnover of plant communities among sites was adequately described by the NMDS

ordination, which accounted for 81 % of the variation in plant species composition (when

correlating the original distance matrix with distances in ordination space, 3-dimensional, final

stress = 0.102) of which 26%, 26%, and 11% could be attributed to axis 1, 2 and 3, respectively.

Likewise for macrofungal communities the NMDS ordination accounted for 72 % of the

variation in species composition (3-dimensional, final stress = 0.146) of which 35%, 21% and

14% could be attributed to axis 1, 2, and 3, respectively. The major gradients in plant species

composition of the 130 sites correlated strongly with soil fertility (NMDS axis 1 strong

correlation with soil N, P and pH), successional stage (NMDS axis 2 strong correlation with light

intensity and opposite correlation with litter mass and number of large trees) and soil moisture

(NMDS axis 3 strong correlation with measured soil moisture), reflecting the gradients that the

sites were selected to cover (Fig. 5, see correlation matrix for the rest of the environmental

variables in Appendix F). Macrofungal species composition showed the same gradients, however

succession and fertility swapped with succession as primary gradient (NMDS1) and fertility as

secondary gradient (NMDS2). NMDS axis 3 reproduced a strong correlation with soil moisture.

Spearman Rho correlations between observational plant species richness and eDNA OTU

'richness' as well as observational plant community composition (as represented by NMDS axes

1-3) and eDNA OTU composition were both strong and confirmative for a recovery of plant

diversity by metabarcoding of soil-derived DNA ($R^2_{richness}$ = 0.652, $R^2_{composition}$ = 0.577-697, Fig.

6). Plant diversity (richness and composition) inferred from soil derived DNA thus resembled

similar metrics derived from direct observation of plant communities, which has also been

investigated in more detail in [39]. We found cross-correlations among species richness of

different taxonomic groups to be predominantly positive or non-significant (Fig. 7). Negative

correlations typically involved insect taxa like Diptera, Lepidoptera, and Orthoptera and e.g.

Fungi.

Discussion

Using ecospace as a conceptual framework [15], we developed a protocol for mapping terrestrial

biodiversity across Denmark represented by numerous, mega-diverse taxa. Across the 130

surveyed sites, covering a tiny fraction (0.0005 %) of the total land area of Denmark, we

observed approximately 5500 species, of which 143 represented new species records for the

country. Our stratification procedures allowed us to cover the local and national environmental

variation across Denmark using only 130 sites of 40 m × 40 m each and provided a good across

and within site coverage of diverse groups of invertebrates and fungi. Finally, the study

demonstrates that eDNA data, once properly curated [39], can be used as an important

supplement to classical biodiversity surveys.

Since, environmental filtering is an important process in community assembly [40], the most

obvious design principle for a biodiversity inventory is to stratify sampling according to major

abiotic and biotic environmental gradients [e.g. 41]. In strongly human-dominated landscapes,

such stratification should incorporate both cultivated and non-cultivated areas and since

environmental gradients are often narrower in cultivated areas, this needs to be taken into

account. We found a close correspondence between the variation in average Ellenberg values at

our sites and those extracted from a very large vegetation database comprising vascular plant

species lists from a national monitoring program. This indicates that we managed to cover the

main environmental gradients found across Denmark. Turnover of plant and macrofungi

communities was significantly linked to moisture, light and fertility and allows us to generalize

relationships between environment and biodiversity derived from local measurements to a large

spatial extent. We note that the use of stratified random sampling implies a biased representation

of rare and common environmental conditions. On the other hand, a completely random

sampling would have led to limited representation of natural biotopes and their disproportionate

contribution to the total biodiversity may have been missed.

While the ecospace framework helped structure our sampling, it also proved challenging with

respect to trade-offs between site size and homogeneity (related to ecospace position), methods

to quantify biotic resources (assessing ecospace expansion) and definitions of temporal and

spatial continuity. Ideally, abiotic and biotic conditions should be homogenous across a site in

order to ensure that site measurements reflect the abiotic position and biotic expansion [15]. A

smaller area would be more likely to be homogenous, but would be less representative. Across

long environmental gradients, homogeneity and representativeness may also vary among for

example, grassland, heathland, and forest. Similarly, while counting the number of different

plant species is easy, accounting for the relative contribution of each species to total biomass and

measuring the availability of different biotic resources such as dead wood, woody debris, litter,

dung, flowers and seeds is much harder. Finally, spatial and temporal continuity is hard to

quantify due to data limitations and because past soil tillage, fertilization, or other land

management or disturbance regimes have not been recorded and must be inferred indirectly. In

addition, an unambiguous definition of continuity breaks is impossible given that most land use

changes and derived community turnover occur gradually over time. We estimated spatial

continuity using broad habitat classes at a range of scales (500 m, 1000 m, 2000 m, 5000 m)

acknowledging that the dependency on spatio-temporal continuity depend on the mobility, life

history and habitat specificity of different species. Our estimate of temporal continuity were also

limited by the availability of aerial photographs and maps, which while not perfect, is good

relative to other parts of the world. Despite these constraints, our estimates of spatial and

temporal continuity varied among sites and were uncorrelated, which allowed us to statistically

test for their relative roles.

We aimed at equal sampling effort per site in terms of trapping and searching time. However,

this was challenged by an array of practicalities. The preferred species sampling methods varied

among taxonomic groups [42, 43] and despite our application of a suite of methods, including

passive sampling in pitfall traps and Malaise traps, baited traps, soil core sampling and active

search, our taxonomic coverage was still incomplete (e.g. aphids, phorid flies and other species-

rich groups living in the canopy are inevitably under-sampled). Our budget also forced us to be

selective with the morphology-based identification of the most difficult species groups, in

particular within Hymenoptera and Diptera. Among identified groups, across-site sample

coverage was consistently high (>0.86) and typically close to 1, which indicates that very few

unseen species remain to be recorded in each community. Invertebrate sampling and

identification is extremely time consuming and relies on rare taxonomic expertise. The within

site sample coverage could only be calculated for spiders and insect orders for which abundance

data were available. Median values of within site sample coverage were also consistently above

0.5, which we consider adequate for cross-site comparisons. We spent more than half of the

inventory budget on invertebrate sampling and identification. Invertebrates constitute by far the

largest fraction of the total biota and, for many species, the adult life stage is short-lived, highly

mobile, and the range of active species varies with season [44, 45]. Trapping also implies a

certain risk of suboptimal placement or vandalism by visiting humans, domestic livestock or wild

scavengers. The resulting number of invertebrate species per site is relatively high and revealed a

considerable variation, which gives ample opportunity for comparative analyses. The high

number of new species for Denmark, particularly macrofungi, can most likely be attributed to the

effort, but also to the inclusion of habitat types that would otherwise have been avoided or

overlooked during opportunistic field surveys [3].

Although methods for DNA extraction, amplification, sequencing and bioinformatics

processing are continuously improved and may lead to better biodiversity metrics from

environmental samples, collecting representative samples from larger areas with unevenly

distributed species remains a challenge. We pooled and homogenized large amounts of soil,

followed by extraction of intracellular as well as extracellular DNA, from a large subsample, to

maximize diversity coverage within a manageable manual workload. Biodiversity metrics based

on plant DNA were correlated to the same metrics for observational plant data. This indicates

that the procedure for sampling, DNA extraction and amplification can be assumed to be

adequate for achieving amplicon data to quantify variation in biodiversity across wide ecological

and environmental gradients for plants, but most likely also for other organisms present in the

soil. These methods are promising for biodiversity studies of many organism groups that are

otherwise difficult to sample and identify (e.g. nematodes, fungi, protists, and arthropods). High

throughput sequencing (HTS) methods produce numerous errors [e.g. 46, 47] and it has been

suggested that richness measures should be avoided altogether for HTS studies [48]. Despite the

remaining challenge of relating genetic units to well-known taxonomic entities, our results along

with those presented in [39] indicate that reliable metrics of α-diversity and community

composition are achievable. With respect to taxonomic annotation, reference databases are far

from complete and the taxonomic annotation of reference sequences are often erroneous.

Furthermore, for many groups of organisms, we have still only described and named a fraction of

the actual species diversity, and the underlying genetic diversity within and between species is

largely unknown for most taxa, leading to uncertainties in OTU/species delimitation and

taxonomic assignment of sequence data. This also means that ecological interpretation of

OTU/species assemblages assessed by eDNA is largely impossible as there is little ecological

knowledge that can be linked to OTUs. Thus, for eDNA-based biodiversity assessment to further

mature, molecular biologists, ecologists, and taxonomists need to work closely together to

produce well-annotated reference databases. Our environmental samples for eDNA, including

soil and litter samples as well as extracted DNA will be preserved for the future. This material

represents a unique resource for the further development of methods within ecology and eDNA.

As more efficient technologies become available in the future, it will be possible to process this

material at an affordable cost and derive further insights on the relationship between traditional

species occurrence, OTU data and environmental variation.

Conclusion

We have presented a comprehensive protocol to obtain a representative, unbiased sample of

multi-taxon biodiversity stratified with respect to the major abiotic gradients. By testing and

evaluating the protocol, we conclude that it is operational and that observed biodiversity

variation may be attributed to measured abiotic and biotic variables. We developed our sampling

protocol based on the ecospace concept, and with this study, we took the first step towards

general models and model inferences with transferability to terrestrial ecosystems and biotas in

other parts of the world. We believe the protocol is also useful for monitoring biodiversity i.e.

tracking changes in biodiversity through time. Meta-barcoding of environmental DNA offers a

promising alternative to traditional inventories (economically and logistically), but barcode

reference libraries are still far from complete. Thus, combining classical taxonomic identification

with metabarcoding of environmental DNA currently appears to offer the most promising

approach to biodiversity research.

Additional files

Additional file 1: Appendix A: Site characteristics for each of the 130 40 × 40 m sites.

Additional file 2: Appendix B: Protocols for data collection**.**

Additional file 3: Appendix C: Ranges of environmental (abiotic and biotic) variables

measured within the 130 sites as well as species richness of various taxonomic groups.

Additional file 4: Appendix D: Temporal and spatial continuity for the 130 Biowide sites.

Additional file 5: Appendix E: Relative richness of arthropods, bryophytes, gastropods, lichens,

macrofungi and vascular plants across all 130 sites

Additional file 6: Appendix F: Correlation matrix for NMDS axes 1, 2 and 3 and environmental

variables

Additional file 7: Appendix G: Number of species in each arthropod family for natural habitats,

species richness hotspots, arable land and plantations.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

RE, LB, IG, TL, TGF, AAI, and AKB were supported by a grant from VILLUM foundation (Biowide, VKR-023343). VILLUM foundation did not play a role in the design of the study, data collection, analyses, interpretation of data or writing of the manuscript.

Authors' contributions

AKB, HHB, AC, TGF, TL, MDDH and RE conceived and designed the study. AKB, HHB, LB, KF, TGF, IG, TL, GN, LS, US, and RE conducted field work. AKB, RE, LD, TTH, and TGF analyzed the data and prepared the figures. LB, KF, IG, MDDH, TL, LS, US, AAI, and HHB sorted and identified specimens. AKB, HHB, LB, ATC, KF, IG, MDDH, TTH, TGF, TL, GSN, LS, US, and RE wrote the manuscript. All authors have read and approved the final version of the manuscript.

Acknowledgements

We thank Ako O. Mirza for plant and soil lab work, Vagn Alstrup and Roar Skovlund Poulsen

for lichen surveys, volunteers that have helped in data collection, land owners, Karl-Henrik

Larsson for aid in identifying critical corticioid fungi, Leif Örstadius for identifying Psathyrella

collections. In regard to identification of invertebrate specimens, we would like to thank Henning

Petersen for identifying Springtails (Collembola), Hjalte Kjærby for identifying Grasshoppers

(Orthoptera) and Harvestmen (Opilliones), Kåre Fog for identifying snails (Gastropoda), Kåre

Würtz Sørensen for identifying various Wasps (Symphyta, Spechidae, Crabronidae, and

Vespidae), Lars Dyhrberg Bruun for identifying spiders (Araneae), Lars Brøndum for identifying

Hoverflies (Syrphidae), Carrion Beetles (Silphidae) as well as Scarabs (Scarabidae), Lars

Skipper for identifying True bugs (Heteroptera), Maja Møholt for identifying Dung beetles

(Aphodius, Onthophagus and Geotrupidae) and Cantharidae, Marianne Graversen for identifying

Longhorn beetles (Cerambycidae) and Ladybugs (Coccinellidae), Peter Wiberg-Larsen for

identifying Caddisflies (Trichoptera), Mathias Holm for identifying True weevils and Seed

weevils (Curculionidae and Apionidae), Monica Aimeé Harlund Oyre for identifying various

Dipterans (Syrphida, Tachinadae, Stratiomyidae, Acroceridae, Rhagionidae, Tephritidae,

Plastytomatidae, Asilidae) as well as Strepsipterans (Strepsiptera) and Book- and Barklice

(Psocoptera), Morten D. D. Hansen for identifying Bees (Apoidea), Carrion beetles (Silphidae),

Click beetles (Elateridae), Scarabs (Scarabaeidea) and Dung beetles (*Aphodius*, *Onthophagus*

and Geotrupidae), Ole Fogh Nielsen for identifying net-winged insects (Neuroptera) and

Strepsipterans (Strepsiptera), Oskar Liset Pryds Hansen and Emil Skovgaard Brandtoft for

identifying Ground beetles (Carabidae), Sofie Amund Kjeldgaard and Steffen Kjeldgaard for

identifying Owlet moths (Noctuidae), Mathias G. Skytte for identifying Rove beetles

(Staphyllinidae), Simon Haarder for identifying galling and mining arthropods, and Ulrik Hasle

Nielsen for identifying Cicadas (Cicadoidea) as well as numerous other volunteers. In regard to

carrying out the eDNA lab work we would like to thank Anne Aagaard Lauridsen, Sarah Mak,

Stine Raith Richter, Carlotta Pietroni and Ida Broman Nielsen. Thomas Stjernegaard Jeppesen

(GBIF) is thanked for making the fungal eDNA dataset publicly available through the GBIF

portal.

Tables

Table 1: Stratification of sites in the survey.

Category	Class	Successional stage	Moisture	Fertility	Number of sites
Arable	Rotational	-	-	-	5
Arable	Ley	-	-	-	5
Arable	Old field	-	-	-	5
Plantation	Beech	-	-	-	5
Plantation	Oak	-	-	-	5
Plantation	Spruce	-	-	-	5
Hotspots	Hotspots	-	-	-	10
Natural	Early/Dry/Rich	Early	Dry	Rich	5
Natural	Mid/Dry/Rich	Mid	Dry	Rich	5
Natural	Late/Dry/Rich	Late	Dry	Rich	5
Natural	Early/Moist/Rich	Early	Moist	Rich	5
Natural	Mid/Moist/Rich	Mid	Moist	Rich	5
Natural	Late/Moist/Rich	Late	Moist	Rich	5
Natural	Early/Wet/Rich	Early	Wet	Rich	5
Natural	Mid/Wet/Rich	Mid	Wet	Rich	5
Natural	Late/Wet/Rich	Late	Wet	Rich	5
Natural	Early/Dry/Poor	Early	Dry	Poor	5
Natural	Mid/Dry/Poor	Mid	Dry	Poor	5
Natural	Late/Dry/Poor	Late	Dry	Poor	5
Natural	Early/Moist/Poor	Early	Moist	Poor	5
Natural	Mid/Moist/Poor	Mid	Moist	Poor	5
Natural	Late/Moist/Poor	Late	Moist	Poor	5
Natural	Early/Wet/Poor	Early	Wet	Poor	5
Natural	Mid/Wet/Poor	Mid	Wet	Poor	5
Natural	Late/Wet/Poor	Late	Wet	Poor	5

525   The sites are sub-divided into four categories (arable, plantations, species richness hotspots, and

526   natural). The natural sites were stratified across specific levels of succession (early, mid, and

527   late), soil moisture (wet, moist, and dry) and soil fertility (rich and poor), while this was the case

528   for the other classes of sites. The number of sites within each of the 25 classes is given.

Table 2: Species richness and sample coverage across habitats per taxonomic group

		Habitat type				Coverage		
	Total	**Natural**	**Hotspots**	**Arable**	**Plantations**	**Across sites**	**Within sites**	**New records in DK**
Vascular plants	719	601 (225)	330 (21)	192 (47)	131 (2)	0.97	-	
Mosses	254	221 (106)	96 (11)	20 (3)	78 (4)	0.97	-	
Lichens	200	183 (92)	76 (9)	19 (5)	58 (3)	0.96	-	1
Fungi	1774	1532 (995)	615 (128)	146 (18)	557 (131)	0.92	-	110
Gallers/miners	203	169 (108)	48 (10)	19 (6)	41 (16)	0.86	-	12
Gastropods	75	72 (18)	42 (0)	19 (1)	38 (2)	0.99	-	
Araneae	335	313 (102)	147 (5)	126 (4)	127 (12)	1	0.87	3
Coleoptera	554	473 (215)	154 (23)	203 (49)	135 (17)	1	0.91	
Carabidae	123	104 (43)	34 (3)	51 (15)	35 (1)	1	0.93	
Hemiptera	446	470 (188)	192 (13)	168 (9)	107 (8)	1	0.79	7
Diptera	196	181 (89)	63 (9)	77 (12)	35 (4)	0.99	0.79	1
Syrphidae	98	89 (42)	31 (2)	42 (6)	20 (2)	0.98	0.81	
Hymenoptera	186	180 (104)	53 (14)	40 (6)	28 (5)	0.98	0.71	1
Lepidoptera	127	127 (71)	31 (3)	33 (3)	16 (2)	0.87	0.67	
Trichoptera	80	77 (39)	23 (1)	24 (2)	16 (1)	0.99	0.92	
Psocoptera	37	41 (8)	23 (0)	26 (0)	18 (1)	-	-	
Neuroptera	23	21 (8)	9 (1)	7 (0)	7 (3)	-	-	
Orthoptera	20	20 (5)	11 (0)	10 (1)	3 (0)	-	-	
Opiliones	18	17 (3)	11 (0)	9 (0)	14 (0)	-	-	
Prostigmata	5	4 (4)	2 (1)	-	-	-	-	
Strepsiptera	2	2 (1)	-	1 (0)	1 (0)	-	-	1
Raphidioptera	2	2 (2)	-	-	-	-	-	
Plecoptera	1	1 (1)	-	-	-	-	-	

Number of species per taxonomic group found in natural sites (n=90), species richness hotspots (n=10), plantations (n=15) arable land

(n=15), and plantations (n=15). Gallers/miners represent multiple insect taxa (see appendix B for a full list). Data for insects are given

per order with additional rows for the species-rich families of Carabidae and Syrphidae. The number of unique species for each habitat

type and taxonomic group is given in brackets. Across sites coverage is the proportion of species likely to be found across all 130

sites, which were actually observed as estimated by extrapolation using the iNEXT package. Within sites coverage is the mean of the

site specific coverage values across the 130 sites for invertebrates with abundance data. The number of new species for Denmark

found during the project is also given for each taxonomic group.

Figures

Figure 1: a) Map of Denmark showing the location of the 130 sites grouped into 15 clusters

within five regions (Njut: Northern Jutland, Wjut: Western Jutland, Ejut: Eastern Jutland, FLM:

542   Funen, Lolland, Møn, Zeal: Zealand). b) Site layout with four 20×20 m quadrants each

543   containing a 5 m radius circle (plot).

545
Figure 2: Validation of the stratification scheme used in site selection. Boxplots of measured

values of nutrient levels (soil N g/m^2), moisture levels (trimmed site mean % Volumetric Water

Content (VWC)), and vegetation height (mean LIDAR canopy height (m)) for the a) 90 natural

sites of different fertility levels (infertile, fertile), moisture levels (dry, moist, wet), and

successional stages (early (open), mid (herb/scrub), late (forest)) and b) the 15 plantations, 15

fields and 10 species richness hotspots.

552
553 **Figure 3:** 95 percentile convex hull plots of Ellenberg F, L and N values from a reference data

554 set (www.naturdata.dk) of open and forest habitat types (blue, n= 59 227) as well as the data set

555 used in this study, Biowide (red, n=130). Black dots represent Ellenberg values of the 130

556 Biowide sites.

557
558 **Figure 4**: Habitat mean values for various carbon resources in the 130 40×40 m sites. Volume

559 of dead wood (m^3/ha), density of dung (cow, sheep, deer, horse, hare) (number/m^2), summed

560 flower density in April, June and August (number/m^2), litter mass (g/m^2) and plant species

561 richness per site are depicted for natural habitat types (early, mid and late successional stage),

562 fields and plantations.

563
Figure 5: Three dimensional NMDS plots for plants with a) showing axis 2 against axis 1 and b)

showing axis 3 against axis 2 and fungi with c) showing axis 2 against axis 1 and d) showing

axis 3 against axis 1. The three main gradients used for selecting the 130 sites (fertility, moisture,

successional stage) are overlaid as arrows (from an envfit analyses in the R package Vegan). The

ordinations are based on plant species lists from the 130 sites a) & b) or macrofungi species lists

from the 124 sites with more than five species c & d) and the arrows reflect soil moisture

measured using a soil moisture meter, fertility measured as soil N and light measured as light

intensity using HOBO loggers. The ordination plots illustrate that the community composition of

vascular plants and macrofungi actually reflect the main gradients the sites were selected to

cover. The scatter of dots shows the variation in abiotic conditions across the 130 sites.

Correlations and p-values can be seen in Appendix D.

Figure 6: Correlation between a) observed site plant species richness and OTU site 'richness' for the 130 sites (Spearman Rho: R^2=0.652, S = 70457, p-value < 0.001), b-d) observed site plant community composition and OTU community composition for the 130 sites b) NMDS axes 1 (Spearman Rho: R^2=0.576, S = 644210, p-value < 0.001), c) NMDS axes 2 (Spearman Rho: R^2=0.594, S = 648480, p-value < 0.001), and d) NMDS axes 3 (Spearman Rho: R^2=0.697, S = 671850, p-value < 0.001).

583
584 **Figure 7**: Cross correlation among the main taxonomic groups included in the study. The colour

585 and shape of the symbol is scaled according to spearman rank correlation coefficients and non-

586 significant (p>0.05) correlations are indicated by a cross.

References

1.   Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B: **How many species are there on Earth and**
in the ocean? *PLoS Biol* 2011, **9**(8):e1001127.
2.   Jetz W, McPherson JM, Guralnick RP: **Integrating biodiversity distribution knowledge: toward a**
global map of life. *Trends Ecol Evol* 2012, **27**(3):151-159.
3.   Isaac NJB, van Strien AJ, August TA, de Zeeuw MP, Roy DB: **Statistics for citizen science:**
extracting signals of change from noisy ecological data. *Methods in Ecology and Evolution* 2014,
5(10):1052-1060.
4.   Nichols JD, Williams BK: **Monitoring for conservation**. *Trends Ecol Evol* 2006, **21**(12):668-673.
5.   Taberlet P, Coissac E, Pompanon F, Brochmann C, Willerslev E: **Towards next-generation**
biodiversity assessment using DNA metabarcoding. *Mol Ecol* 2012, **21**(8):2045-2050.
6.   Thomsen PF, Willerslev E: **Environmental DNA – An emerging tool in conservation for**
monitoring past and present biodiversity. *Biological Conservation* 2015, **183**:4-18.
7.   Porazinska DL, Giblin-Davis RM, Esquivel A, Powers TO, Sung WAY, Thomas WK:
Ecometagenetics confirm high tropical rainforest nematode diversity. *Mol Ecol* 2010,
19(24):5521-5530.
8.   Andersen K, Bird KL, Rasmussen M, Haile J, Breuning-Madsen H, Kjær KH, Orlando L, Gilbert
MTP, Willerslev E: **Meta-barcoding of 'dirt' DNA from soil reflects vertebrate biodiversity**. *Mol*
Ecol 2012, **21**(8):1966-1979.
9.   Taberlet P, Prud'Homme SM, Campione E, Roy J, Miquel C, Shehzad W, Gielly L, Rioux D, Choler
P, CIÉMent J-C *et al*: **Soil sampling and isolation of extracellular DNA from large amount of**
starting material suitable for metabarcoding studies. *Mol Ecol* 2012, **21**(8):1816-1820.
10.   Yoccoz NG, Brathen KA, Gielly L, Haile J, Edwards ME, Goslar T, von Stedingk H, Brysting AK,
Coissac E, Pompanon F *et al*: **DNA from soil mirrors plant taxonomic and growth form diversity**.
Mol Ecol 2012, **21**(15):3647-3655.
11.   Ficetola GF, Miaud C, Pompanon F, Taberlet P: **Species detection using environmental DNA**
from water samples. *Biology Letters* 2008, **4**(4):423-425.
12.   Thomsen PF, Kielgast JOS, Iversen LL, Wiuf C, Rasmussen M, Gilbert MTP, Orlando L, Willerslev E:
Monitoring endangered freshwater biodiversity using environmental DNA. *Mol Ecol* 2012,
21(11):2565-2573.
13.   Shokralla S, Spall JL, Gibson JF, Hajibabaei M: **Next-generation sequencing technologies for**
environmental DNA research. *Mol Ecol* 2012, **21**(8):1794-1805.
14.   Taberlet P, Coissac E, Hajibabaei M, Rieseberg LH: **Environmental DNA**. *Mol Ecol* 2012,
21(8):1789-1793.
15.   Brunbjerg AK, Bruun HH, Moeslund JE, Sadler JP, Svenning J-C, Ejrnæs R: **Ecospace: A unified**
framework for understanding variation in terrestrial biodiversity. *Basic and Applied Ecology*
2017, **18**:86-94.
16.   Belyea LR, Lancaster J: **Assembly rules within a contingent ecology**. *Oikos* 1999, **86**(3):402-416.
17.   Ricklefs RE: **Community diversity - relative roles of local and regional processes**. *Science* 1987,
235(4785):167-171.
18.   Horák J, Kout J, Vodka Š, Donato DC: **Dead wood dependent organisms in one of the oldest**
protected forests of Europe: Investigating the contrasting effects of within-stand variation in a
highly diversified environment. *Forest Ecology and Management* 2016, **363**:229-236.
19.   Seibold S, Bässler C, Brandl R, Gossner MM, Thorn S, Ulyshen MD, Müller J: **Experimental**
studies of dead-wood biodiversity — a review identifying global gaps in knowledge. *Biological*
Conservation 2015, **191**:139-149.
20.   Laliberté E, Zemunik G, Turner BL: **Environmental filtering explains variation in plant diversity**
along resource gradients. *Science* 2014, **345**(6204):1602-1605.

21. Houseman GR, Gross KL: **Linking grassland plant diversity to species pools, sorting and plant**
traits. *Journal of Ecology* 2011, **99**(2):464-472.
22. Rubæk GH, Kristensen K, Olesen SE, Østergaard HS, Heckrath G: **Phosphorus accumulation and**
spatial distribution in agricultural soils in Denmark. *Geoderma* 2013, **209–210**:241-250.
23. Arler F, Jørgensen MS, Galland D, Sørensen EM: **Kampen om m^2 - Prioritering af fremtidens**
arealanvendelse i Danmark. Fonden Teknologirådet. http://www.tekno.dk/wp-
content/uploads/2015/08/Prioritering-af-fremtidens-arealanvendelse-i-Danmark.pdf. In.;
2015.
24. Nygaard B, Juel A, Fredshavn JR: **Ændringer i det § 3-beskyttede naturareal 1995-2014.**
Resultater fra Naturstyrelsens opdateringsprojekt. In. Edited by Aarhus Universitet DNCfMoE,
106 s. - Teknisk rapport fra DCE - Nationalt Center for Miljø og Energi nr. 79; 2016.
25. Williams P, Gibbons D, Margules C, Rebelo A, Humphries C, Pressey R: **A Comparison of**
Richness Hotspots, Rarity Hotspots, and Complementary Areas for Conserving Diversity of
British Birds. *Conserv Biol* 1996, **10**(1):155-174.
26. **Terrestriske Naturtyper 2004 – 2015. NOVANA. Aarhus Universitet, DCE – Nationalt Center for**
Miljø og Energi. www.novana.au.dk.
27. Council Directive 92/43/EEC: **On the conservation of natural habitats and of wild fauna and**
flora. European Commission. 1992.
28. Hill MO, Mountford JO, Roy DB, Bunce RGH: **Ellenberg's indicator values for British plants :**
ECOFACT volume 2 : Technical Annex. Natural Environment Research Council: Centre for
Ecology and Hydrology; 1999.
29. Grime JP, Hodgson JG, Hunt R: **Comparative plant ecology: a functional approach to common**
British species. London: Unwin Hyman; 1989.
30. Ejrnæs R, Bruun HH: **Gradient analysis of dry grassland vegetation in Denmark**. *Journal of*
Vegetation Science 2000, **11**(4):573-584.
31. Lomolino MV, Riddle BR, Brown JH, Brown JH: **Biogeography**. In.: Sinauer Associates Sunderland,
MA; 2006: 65-96.
32. Bijl Lvd, Boutrup S, Jensen PN: **NOVANA. Det nationale program for overvågning af vandmiljøet**
og naturen. Programbeskrivelse 2007-2009 - del 2. Danmarks Miljøundersøgelser, Aarhus
Universitet. *Faglig rapport fra DMU nr 615* 2007.
33. Schmidt P-A, Bálint M, Greshake B, Bandow C, Römbke J, Schmitt I: **Illumina metabarcoding of a**
soil fungal community. *Soil Biology and Biochemistry* 2013, **65**:128-132.
34. White N, Engeman R, Sugihara R, Krupa H: **A comparison of plotless density estimators using**
Monte Carlo simulation on totally enumerated field data sets. *BMC Ecology* 2008, **8**(1):6.
35. Hsieh TC, Ma KH, Chao A: **iNEXT: an R package for rarefaction and extrapolation of species**
diversity (Hill numbers). *Methods Ecol Evol* 2016, **7**(12):1451-1456.
36. R Core team: **R: a language and environment for statistical computing. R Foundation for**
Statistical Computing, Vienna, Austria. In.; 2015.
37. Oksanen J, Blanchet FG, Kindt R, Legendre P, O'Hara RB, Simpson GL, Solymos P, Stevens MHH,
Wagner H: **Package 'vegan': Community Ecology Package. Version 1.17-2.** http://cran.r-
project.org/web/packages/vegan/vegan.pdf. 2010.
38. Koljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AFS, Bahram M, Bates ST, Bruns TD,
Bengtsson-Palme J, Callaghan TM *et al*: **Towards a unified paradigm for sequence-based**
identification of fungi. *Mol Ecol* 2013, **22**(21):5271-5277.
39. Frøslev TG, Kjoller R, Bruun HH, Ejrnæs R, Brunbjerg AK, Pietroni C, Hansen AJ: **Algorithm for**
post-clustering curation of DNA amplicon data yields reliable biodiversity estimates. *Nature*
Comm 2017, **8**:1188.

40. Kraft NJB, Adler PB, Godoy O, James EC, Fuller S, Levine JM: **Community assembly, coexistence**
and the environmental filtering metaphor. *Functional Ecology* 2015, **29**(5):592-599.
41. Gillison AN, Brewer KRW: **The use of gradient directed transects or gradsects in natural**
resource surveys. *Journal of Environmental Management* 1985, **20**:103-127.
42. Popic TJ, Davila YC, Wardle GM: **Evaluation of common methods for sampling invertebrate**
pollinator assemblages: net sampling out-perform pan traps. *PLoS One* 2013, **8**(6):e66665.
43. Standen V: **The adequacy of collecting techniques for estimating species richness of grassland**
invertebrates. *Journal of Applied Ecology* 2000, **37**(5):884-893.
44. CaraDonna PJ, Petry WK, Brennan RM, Cunningham JL, Bronstein JL, Waser NM, Sanders NJ:
Interaction rewiring and the rapid turnover of plant–pollinator networks. *Ecology Letters* 2017,
20(3):385-394.
45. Valverde J, Gómez JM, Perfectti F: **The temporal dimension in individual-based plant**
pollination networks. *Oikos* 2016, **125**(4):468-479.
46. Brown SP, Veach AM, Rigdon-Huss AR, Grond K, Lickteig SK, Lothamer K, Oliver AK, Jumpponen
A: **Scraping the bottom of the barrel: are rare high throughput sequences artifacts?** *Fungal*
Ecology 2015, **13**:221-225.
47. Kunin V, Engelbrektson A, Ochman H, Hugenholtz P: **Wrinkles in the rare biosphere:**
pyrosequencing errors can lead to artificial inflation of diversity estimates. *Environmental*
Microbiology 2010, **12**(1):118-123.
48. Bálint M, Bahram M, Eren AM, Faust K, Fuhrman JA, Lindahl B, O'Hara RB, Öpik M, Sogin ML,
Unterseher M *et al*: **Millions of reads, thousands of taxa: microbial community structure and**
associations analyzed via marker genes. *FEMS Microbiol Rev* 2016, **40**(5):686-700.