Diverse *Streptococcus pneumoniae* strains drive a MAIT cell response through MR1-dependent and cytokine-driven pathways

Ayako Kurioka[1][¶][*], Bonnie van Wilgenburg[1][*][¶], Reza Rezaei Javan[1][*][¶], Ryan Hoyle[1][*][¶],

Andries J. van Tonder[1], Caroline L. Harrold[1], Tianqi Leng[1], Lauren J. Howson[2],

Dawn Shepherd[2], Vincenzo Cerundolo[2], Angela B. Brueggemann[1,3], Paul

Klenerman[1]

[1]The Peter Medawar Building for Pathogen Research, Nuffield Department of

Medicine, University of Oxford, Oxford, UK

[2]MRC Human Immunology Unit, Weatherall Institute of Molecular Medicine,

University of Oxford, Oxford, UK

[3]Department of Medicine, Imperial College London, London, UK

*Corresponding author

E-mail: ayako.kurioka@ndm.ox.ac.uk

[¶]These authors contributed equally

Abstract

Mucosal Associated Invariant T (MAIT) cells represent an innate T cell population of emerging significance. These abundant cells can recognize ligands generated by microbes utilizing the riboflavin synthesis pathway, presented via the major histocompatibility complex (MHC) class I-related molecule MR1 and binding of specific T cell receptors (TCR). They also possess an innate functional programme allowing microbial sensing in a cytokine-dependent, TCR-independent manner. *Streptococcus pneumoniae* is a major human pathogen that is also associated with commensal carriage, thus host control at the mucosal interface is critical. The recognition of *S. pneumoniae* strains by MAIT cells has not been defined, nor have the genomics and transcriptomics of the riboflavin operon (Rib genes). We examined the expression of Rib genes in *S. pneumoniae* at rest and in response to metabolic stress and linked this to MAIT cell activation *in vitro*. We observed robust recognition of *S. pneumoniae* strains at rest and following stress, using both TCR-dependent and TCR-independent pathways. The pathway used was highly dependent on the antigen-presenting cell, but was maintained across a wide range of clinically-relevant strains. The riboflavin operon was highly conserved across a range of 571 *S. pneumoniae* from 39 countries dating back to 1916, and different versions of the riboflavin operon were also identified in related *Streptococcus* species. These data indicate an important functional relationship between MAIT cells and *S. pneumoniae*, which may be tuned by local factors, including the metabolic state of the organism and the antigen-presenting cell that it encounters.

Author Summary

Streptococcus pneumoniae is the leading cause of bacterial pneumonia, causes invasive diseases such as meningitis and bacteraemia, and is associated with significant morbidity and mortality, particularly in children and the elderly. Here, we demonstrate that a novel T cell population called Mucosal-associated invariant T (MAIT) cells is able to respond to a diverse range of *S. pneumoniae* strains. We found that this response was dependent on the T cell receptor (which recognises metabolites of the bacterial riboflavin biosynthesis pathway), cytokines, and the type of antigen-presenting cell. A population genomics approach was also used to assess the prevalence and diversity of the genes encoding the riboflavin biosynthesis pathway among a large and diverse collection of *S. pneumoniae*. These genes were highly conserved across a range of 571 *S. pneumoniae* from 39 countries dating back to 1916, and was also present in other related *Streptococcus* species. Given the low levels of MAIT cells in neonates and MAIT cell decline in the elderly, both of whom are at the highest risk of invasive pneumococcal disease, further understanding of the functional role of MAIT cells in host defense against this major pathogen may allow novel therapeutics or vaccines to be designed.

Introduction

Streptococcus pneumoniae (the 'pneumococcus') is the most common cause of community-acquired pneumonia and is associated with significant morbidity and mortality, especially among young children and older adults (1,2). *S. pneumoniae* also causes invasive diseases like meningitis and bacteraemia, and upper respiratory tract infections like otitis media and sinusitis, among all age groups (3). Antimicrobial-resistant strains are widespread and pose problems in the treatment of infections, which led the World Health Organisation to include *S. pneumoniae* on their recent list of priority pathogens (4). The currently available pneumococcal conjugate vaccines prompt an immune response to polysaccharide capsules (differentiated as serotypes) and they are highly effective at preventing invasive pneumococcal disease due to vaccine-serotype strains; however, current vaccines only protect against a small number of the possible serotypes which has led to increases in the rates of disease caused by non-vaccine-serotype pneumococci are problematic (5,6). Therefore, pneumococcal disease remains a serious problem and a better understanding of the host defense against *S. pneumoniae* may allow for the design of novel therapeutics or vaccines.

There is an increasing appreciation of the role played by unconventional T cells in orchestrating early cellular events in response to invading pathogens (7). Mucosal-associated invariant T (MAIT) cells are a recently described innate T cell population, which are abundant in mucosal tissues including the lung, as well as the blood and liver (8–10). These cells conventionally express a semi-invariant T cell receptor

(TCR) consisting of a Vα7.2-Jα33/Jα12/Jα20 chain, paired with a limited repertoire

of Vβ chains (11). This TCR is able to recognize ligands presented by the conserved

major histocompatibility complex (MHC)-related protein 1 (MR1) (8). MR1 binds

vitamin-B based precursors derived from the riboflavin biosynthesis pathway, which

is conserved across various bacteria and fungi, but does not occur in humans (10).

The ligands presented by MR1 are derived from either early intermediates in

riboflavin synthesis (5-A-RU) that form adducts with other small metabolites (5-OP-

RU), or the direct precursors of riboflavin (e.g. ribityllumazine (RL)-6,7-diMe)

(10,12,13). Human MAIT cells are also characterized by a high expression of the C-

type lectin-like receptor CD161 and the interleukin (IL)-18 receptor, and are able to

respond to innate cytokines even in the absence of prior activation or TCR signaling

(14,15). Upon activation, these cells produce various immunomodulatory cytokines

including IFNγ, TNFα, IL-17, and Granzyme B.

MAIT cells are critical for the control of bacterial infections in mice, particularly in the

lungs (16–18). For instance, aerosol infection models with *Mycobacterium bovis*

Bacillus Calmette-Guérin (BCG) and *Francisella tularensis* live vaccine strain (LVS)

demonstrated that MAIT cells were essential for the early control of the bacterial

burden in the lung (18,19). Indeed, MAIT cells are required to coordinate both innate

and adaptive arms of the immune response during pulmonary infections, as early

GM-CSF production from lung MAIT cells in response to *F. tularensis* was required

for the differentiation of dendritic cells, without which there was a delayed

recruitment of activated CD4+ T cells (20). Thus, the rapid and innate activation of

MAIT cells in response to pulmonary bacteria is critical for bridging the innate and

adaptive system, shaping the local adaptive immune response.

Despite their role in pulmonary infection models, it remains unclear whether MAIT

cells play a role in the defense against *S. pneumoniae* infection. Our first aim was to

determine whether MAIT cells can respond to *S. pneumoniae,* and if so, to elucidate

the mechanism driving such activation. We found that MAIT cells responded to *S.*

pneumoniae in an MR1-dependent manner in the presence of macrophages but not

monocytes, and was dependent on co-stimulation provided by innate cytokines. *In*

vitro experiments and RNA-sequencing experiments also suggested that

temperature and riboflavin availability affected *S. pneumoniae*, which in turn affected

the responsiveness of MAIT cells. Secondly, using a population-level genomics

approach, we found that the riboflavin synthesis pathway is ubiquitous and highly

conserved amongst *S. pneumoniae*, which may have implications in the design of

immunotherapy approaches or vaccines that target MAIT cell activity. Riboflavin

operon genes were also found among other non-pneumococcal *Streptococcus*

species (spp.), including *S. agalactiae* (group B streptococci), which suggests that

perhaps the immunological observations made here could be extended to other

human-associated *Streptococcus* spp*.* infections.

Results

S. pneumoniae possess a highly-conserved riboflavin synthesis operon that is upregulated with heat stress

The genes encoding the riboflavin biosynthetic enzymes of S. pneumoniae (ribD, ribE, ribA and ribH) were found to be clustered together in the same orientation in a predicted 3.4 Kb operon structure (Figure 1A). The prevalence and sequence diversity of the coding regions of the riboflavin genes were investigated in a large, global and historical genome dataset of S. pneumoniae isolated between 1916 and 2008 from people of all ages residing in 36 different countries. 561 (98.2%) of the S. pneumoniae genomes contained the riboflavin operon. Nine of the ten genomes that lacked the operon were of a single multilocus sequence type (note: ST[serotype]), ST13[14/nontypable], and the other belonged to ST695[4] (Supplementary Table 1). All genes in the riboflavin operon were found to be highly conserved: nucleotide and amino acid sequence identity were >99% (Table 1). The dN/dS analysis revealed a higher prevalence of synonymous versus non-synonymous mutations, supporting the importance of maintaining the riboflavin operon (Table 1).

Sequencing was performed on RNA extracted from a pneumococcus that was incubated at a higher temperature than normal (40°C vs. 37°C). Differential expression analysis revealed that all the riboflavin operon genes were significantly up-regulated after 2-4 hours of incubation under heat stress as compared to the control (Figure 1B). Subsequently, the riboflavin operon was found to be significantly

down-regulated after 5–6 hours of incubation. The concurrent increase and decrease

in the expression of the four riboflavin genes suggested that these genes are

transcriptionally coupled.

MAIT cells are activated by *S. pneumoniae*

To determine whether MAIT cells were able to respond to *S. pneumoniae,* 10

Pneumococcal Molecular Epidemiological Network (PMEN) reference strains were

used to probe the activation of MAIT cells in the presence of the monocytic cell line,

THP1. Peripheral blood mononuclear cells (PBMCs) were cultured with

paraformaldehyde (PFA)-fixed *S. pneumoniae* and THP1 cells overnight, or

Escherichia coli (E. coli) as a positive control (Figure 2A). There was a clear

production of interferon-γ (IFNγ) from MAIT cells across all strains although there

was variability in the responses: 7 out of 10 strains reached statistical significance

(PMEN2, PMEN13, PMEN14, PMEN34, PMEN35, PMEN36, and PMEN39).

Similarly, CD69 expression was induced by all 10 strains as measured by geometric

mean fluorescence intensity (gMFI), and reached significance for 7 strains (PMEN9,

PMEN12, PMEN13, PMEN34, PMEN35, PMEN36, and PMEN39). In comparison,

there was negligible activation of non-MAIT cells (CD161−CD8+ T cells; Figure 2B)

across all tests as measured by IFNγ or CD69 expression, suggesting that *S.*

pneumoniae specifically activated MAIT cells.

MAIT cell activation by *S. pneumoniae* in the presence of monocytes is not MR1-dependent

We have previously found that the response of MAIT cells to *E. coli* is dependent on both MR1 as well as innate cytokines IL-12 and IL-18 (14). To investigate whether the response of MAIT cells to *S. pneumoniae* was dependent on MR1, cytokines, or both, we cultured PFA-fixed *S. pneumoniae* with PBMCs and THP1 cells in the presence of anti-MR1, anti-IL-12, and anti-IL-18 blocking antibodies (Figure 3A). As expected, IFNγ expression in response to *E. coli* could be blocked significantly by anti-IL-12 and anti-IL-18 blocking antibodies, which was further reduced and abrogated by the addition of an anti-MR1 blocking antibody, showing that both the TCR and cytokine signaling pathways contribute to MAIT cell activation by *E. coli*. Surprisingly, we found that blockade of MR1 had no effect on pneumococcal MAIT cell activation. Instead, blocking IL-12 and IL-18 completely abrogated MAIT cell activation across all strains tested. This suggested that despite the fact that *S. pneumoniae* possesses the riboflavin synthesis pathway (Table 1, Figure 1A), activation of MAIT cells by *S. pneumoniae* in the presence of THP1 cells was dependent on cytokines, and not activation through antigen presentation by MR1.

Next, to investigate whether *S. pneumoniae* can activate MAIT cells through the MR1 pathway alone, Jurkat cells engineered to express the MAIT cell TCR Vα7.2-Jα33 were cultured with fixed *S. pneumoniae* strains and cultured overnight in the presence of THP1 cells (Figure 3B). There was no significant change in the expression of CD69 by MAIT-Jurkats in the presence of any of the *S. pneumoniae*

strains, although the cells were able to respond to the *E. coli* positive control. Thus,

in the presence of monocytic cells, there was very little activation of MAIT cells by *S.*

pneumoniae through the MR1 pathway.

Given the upregulation of the riboflavin synthesis pathway in *S. pneumoniae* upon

heat stress at 40°C (Figure 1B), we tested whether changing environmental factors

such as temperature and availability of riboflavin in the growth media would trigger

activation of MAIT cells through the MR1 pathway. *S. pneumoniae* strain PMEN34

was grown for 16 hours in Todd Hewitt Broth (THB) media supplemented with yeast

(THB-Y) at 36°C and then either transferred to 40°C incubation, or transferred to

riboflavin-free assay media (RAM) at 36°C or 40°C for 4 hours, before the bacteria

were fixed (Figure 3C). Although there was a slight increase in the fraction of MAIT

cells expressing IFNγ when bacteria were cultured in the absence of riboflavin in

RAM, regardless of temperature, this increase was not dependent on MR1 as the

addition of an anti-MR1 blocking antibody did not reduce the frequency of IFNγ-

expressing MAIT cells. Thus, the complete absence of riboflavin, increased

incubation temperature or the combination of both did not lead to MR1-dependent

activation of MAIT cells.

We also tested whether using live *S. pneumoniae* strain PMEN34 or the supernatant

of *S. pneumoniae* growth culture, instead of fixed bacteria, would stimulate MAIT

cells through the MR1-pathway (Figure 3D); however, these responses were small

and could not be significantly blocked by an anti-MR1 blocking antibody. We

enriched for CD8+ T cells and added these to the assay instead of PBMCs in case

other cells were interfering with MAIT cell activation through MR1, but these

responses were similar to those using PBMCs and not affected by MR1-blocking.

Thus, in the presence of monocytes, MAIT cells are activated mainly through innate

cytokines rather than MR1, regardless of temperature or riboflavin availability.

MR1-dependent activation of MAIT cells by *S. pneumoniae* in the presence of

macrophages

We next tested whether monocyte-derived macrophages may be able to present the

MR1-ligand to activate MAIT cells more effectively through MR1, as alveolar

macrophages play an important role in the immune response to *S. pneumoniae*

(21,22). Furthermore, we investigated whether temperature or the abundance of

riboflavin in the media affects MR1-dependent activation of MAIT cells in the

presence of macrophages. For this, *S. pneumoniae* strain PMEN34 was grown for

16 hours in THB media, with or without yeast extract (THB-Y and THB, respectively)

at either 36°C or 40°C. Given that the riboflavin synthesis pathway was upregulated

by a short incubation at 40°C (Figure 1), we also transferred half of the bacteria

grown overnight at 36°C to 40°C for 4 hours. The bacteria were fixed immediately

and added to PBMCs and monocyte-derived macrophages overnight (Figure 4A, B).

Surprisingly, we found that when using monocyte-derived macrophages, *S.*

pneumoniae induced IFNγ expression from MAIT cells that was reduced by an anti-

MR1 blocking antibody, suggesting that the response was MR1-dependent, in

contrast to the response in the presence of monocytes. The addition of the anti-MR1

blocking antibody significantly reduced the frequency of IFNγ-expressing MAIT cells

regardless of the temperature and media in which the *S. pneumoniae* were grown.

Interestingly, there was a clear increase in activation induced by bacteria grown in

THB media without the addition of yeast, compared to bacteria grown in the

presence of yeast. Degranulation was also induced by *S. pneumoniae* grown in THB

media and was blocked by the anti-MR1-blocking antibody to a varying degree

(Figure 4C, D).

Next, in order to compare the dependency of MAIT cell activation by *S. pneumoniae*

on MR1, IL-12 and IL-18 in the context of macrophages, *S. pneumoniae* strain

PMEN34 was grown for 16 hours in THB or THB-Y media at either 36˚C or 40˚C.

Half of the bacteria grown overnight at 36˚C were transferred to 40˚C for an

additional 4 hours. Then the bacteria were fixed and added to PBMCs and

macrophages overnight in the presence of anti-MR1, anti-IL-12, and anti-IL-18

antibodies (Figure 4E, F). We found that there was a significant effect of blocking

MR1 on IFNγ production from MAIT cells in the presence of *S. pneumoniae* cultured

in THB media, regardless of temperature, but this was almost completely abrogated

in the presence of anti-IL-12 and anti-IL-18 blocking antibodies. There was a similar

effect of blocking antibodies on MAIT cell IFNγ production in response to *S.*

pneumoniae cultured in THB-Y media overnight at 40˚C. Furthermore, although we

have previously found that degranulation in response to *E. coli* was completely

dependent on MR1 and not on cytokines (23), degranulation of MAIT cells in

response to *S. pneumoniae* was, in each case, further reduced by the addition of

anti-IL-12 and anti-IL-18 compared to blocking MR1 alone. In particular,

degranulation in response to *S. pneumoniae* grown at 40°C for the last 4 hours in

THB media was significantly reduced by anti-MR1 blocking, which was further

reduced by the addition of anti-IL-12 and anti-IL-18 blocking antibodies (Figure 4F).

This suggests that degranulation in response to *S. pneumoniae* requires co-

stimulation from cytokines.

Finally, we set out to confirm that *S. pneumoniae* were able to activate MAIT cells in

the presence of macrophages in an MR1-dependent manner. We cultured Jurkat

cells that express the MAIT cell TCR Vα7.2-Jα33 with fixed *S. pneumoniae* grown in

THB or THB-Y media at different temperatures in the presence of monocyte-derived

macrophages (Figure 4G). This showed that *S. pneumoniae* grown in THB media

significantly increased the expression of CD69 in MAIT-Jurkat cell lines, compared to

the Jurkat cells cultured with the sterility control, although the response was

relatively weak. Thus, in the presence of monocyte-derived macrophages, *S.*

pneumoniae were able to activate MAIT cells in an MR1-dependent manner, but

were highly dependent on the presence of cytokines to boost the response.

Riboflavin operons are also present in non-pneumococcal *Streptococcus* spp.

A bioinformatic investigation of 824 genomes of 69 different *Streptococcus* spp.

revealed that the riboflavin operon was also present in other streptococci. 11

different versions of the riboflavin operon were identified among 13 non-

pneumococcal *Streptococcus* spp. (Supplementary Table 2). The majority of these

riboflavin operons were located between genes involved in arginine biosynthesis

(*argC*, *argJ*, *argB*, *argD*) and ribonucleotide reduction (*nrdF2, nrdE2, nrdH*) (Figure

5A). Despite identical gene synteny between different versions of the riboflavin

operon (Figure 5A), they differed greatly in nucleotide sequence identity (Figure 5B).

The *S. pneumoniae* version of the riboflavin operon was found among all 16

genomes of *S. pseudopneumoniae* and two genomes each of *S. mitis* and *S. oralis*

(Figure 5C-D). No riboflavin operon genes were identified among the remaining 48 *S.*

mitis and 48 *S. oralis* genomes. *S. pneumoniae, S pseudopneumoniae, S. mitis* and

S. oralis are all closely-related commensal streptococcal species that can exchange

DNA with one another; therefore the limited number of riboflavin operons present in

S. mitis and *S. oralis* suggest that the examples identified here were the result of

horizontal genetic exchange (24). Some versions of riboflavin operons identified

among other *Streptococcus* spp. were exclusively present in only one species, e.g.

version 2 was identified in all 50 genomes of *S. agalactiae* but no other species;

whereas in contrast, *S. equinus* contained three different versions of the riboflavin

operon, one of which (version 5) was also found in *S. infantarius* (Figure 5D;

Supplementary Table 2). Furthermore, over 2400 *S. pyogenes* genomes were

investigated for the presence of Rib genes. The riboflavin operon could not be found

in the genomes of *S. pyogenes* (data not shown).

Discussion

Our study is the first to demonstrate that MAIT cells can respond to and recognize *S. pneumoniae*. Given the urgent global need to tackle antimicrobial-resistant strains and pneumococci not covered by the currently available vaccines, understanding the mechanism by which MAIT cells are activated by *S. pneumoniae* provides a method of targeting a metabolic pathway that is highly conserved amongst *S. pneumoniae*.

We found that MAIT cell activation by *S. pneumoniae* through the MR1-restricted pathway was dependent on the type of antigen-presenting cell. There was a significant effect of blocking MR1 recognition by MAIT cells only when the *S. pneumoniae* was presented by macrophages, but not monocytes (either those amongst PBMCs or by the monocytic cell line THP1). Indeed, the lack of MAIT cell activation by *S. pneumoniae* in a previous paper (25) is in line with our finding that monocytes are not able to sufficiently activate MAIT cells solely through the presentation of ligands from *S. pneumoniae* by MR1. The reason for the difference in the ability of these antigen-presenting cells to activate MAIT cells through MR1 antigen presentation in the case of *S. pneumoniae* is not clear. It has been suggested that monocytes are poor antigen presenters (26) and instead function as antigen-transporting cells that deliver antigen to the draining lymph nodes (27). Although THP1 cells and monocytes have been used repeatedly to activate MAIT cells in an MR1-dependent manner with numerous other bacteria, the pathogenesis of *S. pneumoniae* is related to its many virulence factors, some of which are associated with minimizing phagocytosis and thereby antigen presentation (further

discussed below). Macrophages may be a more powerful antigen-presenting cell that

are able to efficiently phagocytose the bacteria, provide more co-stimulatory signals,

or resist interference with antigen presentation and loading of MR1. Indeed, mice

lacking alveolar macrophages, located at the interphase between air and lung tissue,

show a delayed and impaired bacterial clearance as compared with control mice

(21,22). Thus, alveolar macrophages provide the first line of cellular defense against

microbes and may be an important presenter of *S. pneumoniae* antigen to MAIT

cells.

Even in the presence of macrophages, the MAIT cell response to *S. pneumoniae*

was highly co-dependent on cytokines. This is most likely due to the weak TCR

signal induced by *S. pneumoniae*, as seen by the significant but weak activation of

MAIT-Jurkat cell lines even when using macrophages. MAIT cells are characterized

by downregulated TCR signaling components and a high dependency on co-

stimulation provided by cytokines for activation and expansion (28–30). The

particularly weak TCR signal provided by *S. pneumoniae* may be because *S.*

pneumoniae is a bacterium highly specialized to evade the host immune system by

circumventing phagocytosis and antigen presentation. The efficiency of phagocytosis

can be impeded by the presence of the thick polysaccharide capsule, while another

virulence-associated trait is the ability of this bacterium to undergo autolysis.

Autolysis of *S. pneumoniae* has been shown to specifically reduce the phagocytosis

of intact bacteria, even in the absence of a polysaccharide capsule, and reduce the

production of phagocyte-activating cytokines such as TNFα, IFNγ and IL-12 (31).

The uniquely high sensitivity of MAIT cells to cytokines such as IL-12 (32) allows

these cells to boost the immune response and provide early IFNγ production.

The biggest disease burden caused by *S. pneumoniae* by far is pneumonia, which is

the leading infectious cause of mortality in children under five years of age (1), as

well as accounting for a large part of community-acquired pneumonia in the elderly

(2). Given that we show the mechanism by which MAIT cells respond to *S.*

pneumoniae, and the critical role that pulmonary MAIT cells play *in vivo* against lung

infections (18,20), it would be reasonable to suggest that MAIT cells may play a role

in pneumococcal pneumonia. These cells may also be a critical factor in secondary

bacterial pneumonia that follow influenza infection, which are most commonly

caused by *S. pneumoniae* and associated with high mortality (33). MAIT cells are

reduced in frequency in the blood of patients with acute influenza infection,

particularly in those who succumbed to this disease (15,34). The loss of MAIT cells

due to primary influenza infection may leave the patient vulnerable to secondary

infection by *S. pneumoniae*, due to the lack of early control and immunomodulation

provided by these cells, as has been shown with γδ T cells (35). Whether the low

numbers of MAIT cells in neonates (36,37), or the decline in MAIT cell numbers in

the elderly (38) and in influenza (15) affects the susceptibility of these patients to

pneumococcal pneumonia will be important to investigate in *in vivo* models of

pneumococcal infection (39).

Interestingly, we found that temperature and riboflavin availability for *S. pneumoniae*

have the potential to affect riboflavin synthesis and MAIT cell activation. In our

models using macrophages, there was a difference in MR1-dependent MAIT cell

activation induced by *S. pneumoniae* cultured in THB media compared to THB-Y

media, the latter of which is supplemented with riboflavin-rich yeast extract. Although

the THB media alone supports the growth of riboflavin-auxotrophs such as

Streptococcus pyogenes, this has been shown to be due to the ability of *S.*

pyogenes to utilize THB-derived components as a substitute for riboflavin, and

mutant strains without this ability require the addition of riboflavin or riboflavin-rich

yeast in order to grow in THB media (40). Thus, THB media on its own does not

contain riboflavin. We found that *S. pneumoniae* grown in THB media consistently

induced greater MAIT cell activation, which may be partly due to the increased

production of riboflavin in bacteria grown in riboflavin-deficient THB media,

compared to those grown in riboflavin-rich THB-Y media. We also found that

riboflavin synthesis among *S. pneumoniae* was influenced by temperature: RNA-seq

transcriptomic analyses of *S. pneumoniae* revealed that incubation at 40°C

upregulated the riboflavin synthesis genes. A relationship between riboflavin

production and temperature has been demonstrated before in *Lactococcus lactis*: it

suffers from oxidative stress caused by riboflavin starvation at high temperatures and

results in severe growth inhibition, which can be improved by the addition of

riboflavin to the medium (41). Although no consistent increase in MR1-dependent

MAIT cell activation could be detected in response to pneumococci grown at higher

temperatures in our short-term activation models, it may be possible that MAIT cells

are able to react to increased riboflavin production of *S. pneumoniae* at higher body

temperatures during a fever. Thus, MAIT cells may be able to not only recognize the

presence of *S. pneumoniae* but respond to the environment and condition in which

the bacteria is growing - an important indicator of the pathogenic potential of a

commensal pathogen such as *S. pneumoniae.*

We used a population genomics approach to assess the prevalence and diversity of

the riboflavin operon among a large and diverse collection of *S. pneumoniae*. This

revealed that the riboflavin genes are nearly ubiquitous and highly conserved at a

nucleotide level among *S. pneumoniae* that had been recovered over the past

century. This provides confidence that the findings presented here are generalizable

among *S. pneumoniae*, with rare exceptions.

We also assessed the presence of riboflavin synthesis genes among non-

pneumococcal *Streptococcus* spp. and found that a number of other species do

possess these genes, including other human-associated commensal streptococci

like *S. agalactiae*. Notably though, not all *Streptococcus* spp. possessed a riboflavin

operon (at least one that was detectable by our screening method) and there was

variation among the genomes that did, either in that more than one version of

riboflavin operon was detected within a bacterial species, or that the presence of a

riboflavin operon was not ubiquitous among genomes of a particular bacterial

species. Hence, caution must be exercised when extrapolating findings based on a

small number of bacterial strains to the population as a whole since they may not be

representative, and ideally a population-based approach is used whenever possible.

However, the fact that other *Streptococcus* spp. also possess a riboflavin operon

presents opportunities for future studies. For example, given that MAIT cells reside in

the female genital mucosa (42) it would be important to explore whether there is a

MAIT cell response in the context of vaginal colonization of *S. agalactiae* (group B

streptococci) among pregnant women and invasive neonatal infections (43).

Overall these data show a robust response of MAIT cells to *S. pneumoniae*, and

conservation of the relevant biosynthetic pathway in this organism and other closely-

related *Streptococcus* spp. Given the low levels of MAIT cells in early life and their

decline in old age – the highest risk populations for invasive pneumococcal disease

– further understanding of the functional role of MAIT cells *in vivo* in host defense

against this major pathogen should be of substantial interest.

Materials and Methods

Flow cytometry

For immunofluorescence staining, dead cells were excluded with the Live/Dead Fixable near-IR dead-cell stain (Invitrogen). For internal staining, cells were fixed with 1% formaldehyde (Sigma Aldrich) and permeabilised with permeabilisation buffer (eBioscience). Antibodies used were: CD3 Pacific Orange, Granzyme B APC (Life Technologies), CD8 PE-Cy7, TNFα PerCP-Cy5.5, Vα7.2 APC, FITC, or PE, CD107a PE-Cy7 (BioLegend), CD3 efluor 450, CD8 PerCP-Cy5.5, CD69 FITC or Pacific Blue, IFNγ PerCP-Cy5.5 (eBiosciences), CD161 PE, CD4 VioGreen, IFNγ FITC (Miltenyi Biotec).

Cells

Whole blood was obtained from leukocyte cones (NHS Blood and Transplant). PBMCs were isolated by standard density gradient centrifugation (Lymphoprep^{TM} Axis Shield Diagnostics). All samples were collected with appropriate patient consent and local research ethics committee approval (COREC 04.OXA.010). Where indicated, CD8+ cells were positively enriched using CD8 Microbeads (Miltenyi Biotech) before *in vitro* stimulation. Monocyte-derived macrophages were generated by enriching for monocytes using CD14 Microbeads (Miltenyi Biotech) before culturing with 50ng/ml GM-CSF (Miltenyi Biotech) in RPMI media, penicillin/streptomycin, L-glutamine, and 10% human serum for 6-8 days.

The Jurkat-MAIT cell line was produced by cloning into a pHR-IRES vector the

TRAV1-2–TRAJ33 (CDR3α: CAVMDSNYQLIW) MAIT cell α chain and then cloning

in the TRBV6-1–TRBJ2-3 β chain (CDR3β-CASSETSGSPDTQYF). The vector was

then co-transfected into 293T cells with the HIV gag-pol and VSV-G expression

plasmids using X-tremeGENE™ 9 DNA Transfection Reagent (Sigma) according to

the manufacturer's instructions. The supernatant from this culture containing the

lentiviral particles was then used to transduce J.RT3-T3.5 (JRT3) cells (which lack

an endogenous TCRβ chain). TCR expression and pairing with MAIT cell TRAV1-2

(Vα7.2) chain was confirmed by flow cytometry and cells sorted based on CD3 and

Vα7.2 expression.

Pneumococcal reference strains used in experiments

Ten PMEN reference strains of *S. pneumoniae* were tested in this study (note:

country of first detection[serotype]; multilocus sequence type): PMEN2 (Spain[6B]-2; ST90),

PMEN3 (Spain[9V]-3; ST156), PMEN9 (England[14]-9; ST9), PMEN12 (Finland[6B]-12;

ST270), PMEN13 (S.Africa[19A]-13; ST41), PMEN14 (Taiwan[19F]-14; ST236), PMEN34

(Denmark[12F]-34; ST218), PMEN35 (Netherlands[14]-35; ST124), PMEN36

(Netherlands[18C]-36; ST113), PMEN39 (Netherlands[7F]-39; ST191)(44–46). Eight

serotypes were represented, all of which are vaccine serotypes apart from 12F

(PMEN34).

PMEN strains were cultured from freezer stocks to Columbia blood agar plates

(Oxoid) and incubated overnight at 37°C+5% CO_2, then transferred to THB media

(Sigma Aldrich) with 0.5% yeast extract (THB-Y; Sigma Aldrich) and incubated at

$37°C + 5\%$ CO_2 overnight, unless indicated otherwise. In specific experiments,

bacteria were grown in Riboflavin Assay Media (RAM; BD Difco). *E. coli* (DH5a,

Invitrogen) were cultured in Lysogeny Broth (LB) medium and incubated overnight in

a shaking incubator.

S. pneumoniae or *E. coli* were fixed in 1% PFA for 15 minutes, and washed

extensively in sterile, filtered phosphate-buffered saline (PBS). A negative control

was prepared in identical fashion and used as a sterility control. Following fixing,

bacteria were immediately used in *in vitro* stimulation assays. Alternatively, filtered

bacterial culture supernatant or unfixed live bacteria was added to assays where

indicated. Live pneumococci were added to PBMCs and antigen-presenting cells for

30 minutes at $37°C$ before cells were extensively washed and cultured for the

remainder of the overnight assay in $100\mu g/ml$ gentamycin.

***In vitro* stimulation of MAIT cells**

THP1 cells (ECACC, UK) were incubated overnight with PFA-fixed *S. pneumoniae* or

E. coli at a ratio of 30 bacteria per cell (BpC), or the sterility control (without bacteria,

see above). THP1 cells were washed and PBMCs or enriched CD8+ T cells were

added to THP1 cells overnight. Brefeldin A (eBioscience) was added for the final 4

hours of the stimulation. Alternatively, for the assessment of degranulation, anti-

CD107a PE-Cy7 (BioLegend) was added from the start of the stimulation. For

blocking experiments, anti-MR1 (BioLegend), anti-IL-12p40/70, anti-IL-18 antibody

(both BioLegend), or the appropriate isotype controls, were added for the duration of

the experiment.

Data acquisition and statistical analysis

Data on MAIT cell activation were collected on the MACSQuant Analyser (Miltenyi Biotech) and were analysed using FlowJo v9.8 (TreeStar). All graphs and statistical analyses were completed using GraphPad Prism software version 6. All data are presented as means with standard error of the mean (S.E.M.) unless otherwise indicated.

Compilation of the genome datasets

Two large genome datasets were compiled for this study and data were stored in a BIGSdb database (47). The *S. pneumoniae* dataset consisted of 571 historical and modern genomes isolated from 1916–2009 from people of all ages residing in 39 different countries. The *S. pneumoniae* isolates were recovered from both carriage and disease, and 89 serotypes and 296 multilocus sequence types were represented in this dataset (Supplementary Table 1). 486 *S. pneumoniae* genome sequences were compiled from previously published studies or were downloaded from GenBank (48). The remaining 85 *S. pneumoniae* genomes were recently sequenced. *S. pneumoniae* cultures were prepared and incubated as described above, before DNA was extracted using the Promega Maxwell® 16 Instrument and Buccal Swab LEV DNA purification kits following the manufacturer's protocol. DNA extracts were sent to the Oxford Genomics Centre where libraries were made and DNA was sequenced on the Ilumina® platform. Velvet was used to make *de novo* genome assemblies, which were further improved using SSPACE and GapFiller (49–51).

The non-pneumococcal *Streptococcus* spp. dataset contained 834 genomes of 69

different streptococcal species (Supplementary Table 2). 34 genomes were newly

sequenced as above and the rest were downloaded from the ribosomal multilocus

sequence typing (rMLST) database (52). For each species, the number of genomes

included in this study dataset was capped at 50: if fewer than 50 genomes were

available in the rMLST database for a given species, then all available genomes

were included, but if more than 50 sequenced genomes were available then

genomes were manually selected for inclusion. In these instances, the population

structure of the species was depicted using PHYLOViZ and 50 genomes were

selected with the aim of maximising the population-level diversity of that species

from the available genomes (53).

Genomes were annotated using both RAST (54) and Prokka (55). Genome

sequences can be accessed from the European Nucleotide Archive (56), PubMLST

website(57), GenBank(58) and/or the rMLST database(59) (see Supplementary

Tables 1 and 2).

RNA-sequencing experiments and analyses

S. pneumoniae isolate 2/2 was cultured in seven 10 ml tubes of brain-heart infusion

broth and incubated at 40°C + 5% CO_2 for 6 hours to mimic heat shock. The

experimental control was the same isolate 2/2, also cultured in seven 10 ml tubes of

brain-heart infusion broth but incubated at standard conditions of 37°C + 5% CO_2.

Broth cultures at five time points (2, 3, 4, 5 and 6 hours of incubation) were removed

and 19 ml of RNAprotect Bacteria Reagent (Qiagen) was added to stabilise the RNA.

RNA was extracted from the samples using the Promega Maxwell® 16 Instrument

and LEV simplyRNA Cells purification kit, following the manufacturer's protocol.

Extracted RNA samples were sent to the Oxford Genomics Centre for processing.

Library preps were made using RNA-Seq Ribozero kits (Illumina, Inc) and

sequencing was performed on the MiSeq (Illumina, Inc).

The sequenced forward and reverse reads were paired and mapped to the *S.*

pneumoniae strain 2/2 genome using Bowtie2 with the highest sensitivity option (60).

Differential gene expression was analysed in Geneious version 9.1 (Biomatters Ltd)

using the DESeq method (61). Genes with an adjusted p-value < 0.05 were

considered to be differentially expressed. RNA sequencing data used in this paper

have been deposited in the Gene Expression Omnibus (GEO) database with

accession number GSMXXXXXXX (pending).

Genomic analyses of riboflavin operon genes

Genes involved in riboflavin metabolism were identified using the KEGG pathway

database (KEGG entry number: snp00740) and previously published experimental

work (12,62). Individual BLAST searches of *ribD*, *ribE, ribA* and *ribH* sequences

among all 571 *S. pneumoniae* genomes were performed via the BIGSdb database.

There were two instances each for *ribD* and *ribH* where the gene sequences were

split over multiple assembly contigs and these sequences were excluded from

further analyses. Individually, multiple nucleotide sequence alignments for *ribD*, *ribE*,

ribA and *ribH* were performed in Geneious using the ClustalW algorithm with default

parameters (Gap open cost=15, Gap extend cost=6.66) (63). To compute the dN/dS

ratio, the number of synonymous (dS) and non-synonymous (dN) substitutions per

site was determined on codon-aligned sequences using a maximum likelihood

method, conducted through HyPhy (64) on the Datamonkey server (65,66). For the

pairwise positive selection analysis, amino acid sequences were aligned in Geneious

using the ClustalW algorithm (Cost matrix: BLOSUM, Gap open cost=10, Gap

extend cost=0.1).

Identification and categorisation of the riboflavin operon genes in non-

pneumococcal *Streptococcus* spp.

The *ribD*, *ribE*, *ribA* and *ribH* sequences in *S. pneumoniae* strain 2/2 genome were

used as the query to BLAST against the non-pneumococcal *Streptococcus* spp.

genome dataset (parameters: word size 11; reward 5; penalty -4; gap open 8; gap

extend 6). All BLAST hits were manually inspected to confirm the presence of the

riboflavin genes. Protein domains in the identified genes were annotated using the

Conserved Domain search feature at NCBI (67) to confirm the presence of riboflavin

biosynthesis genes. To categorise the different versions of the riboflavin operon, all

sequences were clustered using CD-Hit (68) at a ≥90% similarity threshold, and a

representative sequence from each cluster was selected. One 'cluster' contained a

single riboflavin operon sequence that was disrupted by a transposon and this single

sequence was excluded from further analyses. Multiple nucleotide sequence

alignments of different versions of the riboflavin operons was performed in Geneious

using the ClustalW algorithm with default parameters (Gap open cost=15, Gap

extend cost=6.66). The multiple nucleotide sequence alignment output was used

within the Geneious environment to calculate the percentage identity matrix.

Assessment of the relationships among *Streptococcus* spp.

A phylogenetic tree was built using concatenated sequence data from the ribosomal

loci using the neighbour-joining method as implemented using the BIGSdb

PhyloTree plugin (52,69). The tree was annotated using iTOL (70) and Inkscape (71).

Acknowledgements

The authors gratefully acknowledge Professor Regine Hakenbeck at the University

of Kaiserslautern for the stock cultures of streptococci that were newly sequenced in

this study.

Figure legends

Figure 1. Genetic and transcriptomic data related to the riboflavin operon in *S. pneumoniae*. A) The riboflavin operon is depicted with riboflavin genes *ribD, ribE, ribA* and *ribH* (red) and flanking genes (grey). B) RNA expression data at five timepoints (2-6 hours after initial incubation) are illustrated for each riboflavin gene and the flanking genes. Genes marked with differential expression levels in green were upregulated and those in red were downregulated during 40°C incubation, as compared to normal 37°C incubation. An asterisk (*) to the left of a cell indicates a statistically significant result.

Figure 2. MAIT cells are activated by *S. pneumoniae*. 10 Pneumococcal Molecular Epidemiological Network (PMEN) reference strains were used to probe the activation of MAIT cells in the presence of the monocytic cell line, THP1. *Escherichia coli (E. coli)* were added as a positive control. (A-B) Frequency of cells expressing IFNγ in MAIT cells (A) or CD161- CD8+ T cells (B) are shown (n=9). (C-D) CD69 expression measured by geometric mean fluorescence intensity (gMFI) in MAIT cells (C) or CD161- CD8+ T cells (D) are shown (n=6). **P<0.01, *P<0.05 by repeated measures one-way ANOVA with Dunnett's multiple comparisons test, compared to the sterility control (SC). Numbers indicate the PMEN reference strains. SC= sterility control.

Figure 3. MAIT cell activation by *S. pneumoniae* in presence of monocytes is not MR1-dependent. (A) Paraformaldehyde (PFA)-fixed *S. pneumoniae* PMEN reference strains or *E. coli* were cultured with PBMCs and THP1 cells in the

presence of anti-MR1, anti-IL-12, and anti-IL-18 blocking antibodies. IFNγ

expression from MAIT cells are shown. ****P<0.0001, ***P<0.001, **P<0.01 by

repeated measures one-way ANOVA with Dunnett's multiple comparisons test,

compared to the no antibody control (n=7). (B) Jurkat cells expressing the MAIT cell

TCR (white bars) or control cells not expressing the MAIT cell TCR (black bars) were

cultured with THP1 cells overnight with the indicated PMEN strains of *S.*

pneumoniae, or *E. coli* as a positive control. Activation was measured as the

geometric mean fluorescence intensity (gMFI) of CD69 expressed by Jurkat cells.

SC=sterility control baseline activation, as indicated by the dotted line.

Representative of three independent experiments. (C) The *S. pneumoniae* PMEN34

strain was grown overnight at 36°C or 40°C in THB-Y media and either cultured in

THB-Y media for the last 4 hours or transferred to riboflavin-free media (RAM). The

bacteria were then fixed and cultured with PBMCs and THP1 cells overnight.

Frequency of IFNγ expressing MAIT cells is shown, in the presence or absence of

anti-MR1 blocking antibody. ns=non-significant by two-way ANOVA with Sidak's

multiple comparisons test (n=3). (D) The *S. pneumoniae* PMEN34 strain was grown

overnight and the supernatant, fixed or live bacteria were added to THP1 cells

overnight with either PBMCs (left) or enriched CD8+ T cells (right), in the presence

or absence of anti-MR1 blocking antibody. Frequency of IFNγ expressing MAIT cells

are shown. SC=sterility control, ns=non-significant by two-way ANOVA with Sidak's

multiple comparisons test (n=3).

Figure 4. MR1-dependent activation of MAIT cells by *S. pneumoniae* in the

presence of macrophages. A-D) The *S. pneumoniae* PMEN34 strain was grown

for 16 hours in THB media, with or without yeast extract (THB-Y and THB,

respectively) at either 36˚C (36) or 40˚C (40 overnight). Half of the bacteria grown

overnight at 36˚C were transferred to 40˚C for 4 hours (40 4 hr). The bacteria were

fixed immediately and added to PBMCs and monocyte-derived macrophages

overnight, in the presence or absence of anti-MR1 blocking antibody. Frequency of

MAIT cells expressing IFNγ (A) and representative example of IFNγ expression from

MAIT cells by FACS (B) are shown with isotype control or anti-MR1 blocking

antibody. ****$P<0.0001$, ***$P<0.001$, and * $P<0.05$ by two-way ANOVA with Sidak's

multiple comparisons test (n=6). Frequency of MAIT cells expressing CD107a (C)

and representative example of CD107a expression from MAIT cells by FACS (D) are

shown with isotype control or anti-MR1 blocking antibody. ***$P<0.001$, **$P<0.01$, or

non-significant (ns) by two-way ANOVA with Sidak's multiple comparisons test (n=4).

36=36°C, 40=40°C. E-F) The *S. pneumoniae* PMEN34 strain was grown for 16

hours in THB-Y or THB media, at either 36˚C or 40˚C. Half of the bacteria grown

overnight at 36˚C were transferred to 40˚C for 4 hours. The bacteria were fixed

immediately and added to PBMCs and monocyte-derived macrophages overnight, in

the presence or absence of indicated combinations of anti-MR1, anti-IL-12, and anti-

IL-18 blocking antibodies. Frequency of MAIT cells expressing IFNγ (E) and CD107a

(F) are shown. 36=36°C, 40=40°C. ****$P<0.0001$, ***$P<0.001$, **$P<0.01$, *$P<0.05$ or

non-significant (ns) by two-way ANOVA with Dunnett's multiple comparisons test

(n=3). (G) Jurkat cells expressing the MAIT cell TCR were cultured with monocyte-

derived macrophages overnight with the PMEN34 strain of *S. pneumoniae*.

Activation was measured as frequency of Jurkat cells expressing CD69. Dotted line

indicates CD69 expression by Jurkats in the presence of sterility control (SC).

THB=Todd Hewitt Broth, THB-Y=THB with yeast extract. *$P<0.05$ by repeated

measures one-way ANOVA with Dunnett's multiple comparisons test. All

experiments were performed in duplicates, and representative of two independent

experiments.

Figure 5. Evidence for different versions of riboflavin operons in other

***Streptococcus* spp.** A) The riboflavin operon found in *S. pneumoniae* (version 1)

and its flanking genes are depicted and compared to 10 additional representative

versions of the riboflavin operon found among other *Streptococcus* spp. B) Matrix of

pairwise comparisons of nucleotide similarity among the 11 different versions of the

riboflavin operon. C) Table summarizing the riboflavin operons found in 13 non-

pneumococcal *Streptococcus* spp. D) Phylogenetic tree constructed based upon the

concatenated sequences of 53 ribosomal MLST loci among 571 *S. pneumoniae* and

824 *Streptococcus* spp. genomes. Branches of the tree were coloured grey if no

riboflavin operon was identified within the genome, whereas other colours represent

genomes in bacterial species that did possess a version of a riboflavin operon. The

coloured outer ring indicates the version of riboflavin operon that was identified in

each genome or set of genomes. The rectangular box contains an expanded view of

the circled area of the phylogenetic tree.

References

1. Liu L, Oza S, Hogan D, Perin J, Rudan I, Lawn JE, et al. Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis. Lancet. 2015;385(9966):430–40.

2. Drijkoningen JJC, Rohde GGU. Pneumococcal infection in adults: burden of disease. Clin Microbiol Infect. 2014;20:45–51.

3. Mehr S, Wood N. Streptococcus pneumoniae – a review of carriage, infection, serotype replacement and vaccination. Paediatr Respir Rev. 2012;13(4):258–64.

4. WHO | Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics. WHO. 2017;

5. Huang SS, Platt R, Rifas-Shiman SL, Pelton SI, Goldmann D, Finkelstein JA. Post-PCV7 Changes in Colonizing Pneumococcal Serotypes in 16 Massachusetts Communities, 2001 and 2004. Pediatrics. 2005;116(3):e408–13.

6. Singleton RJ, Hennessy TW, Bulkow LR, Hammitt LL, Zulz T, Hurlburt DA, et al. Invasive Pneumococcal Disease Caused by Nonvaccine Serotypes Among Alaska Native Children With High Levels of 7-Valent Pneumococcal Conjugate Vaccine Coverage. JAMA. 2007;297(16):1784.

7. Godfrey DI, Uldrich AP, McCluskey J, Rossjohn J, Moody DB. The burgeoning family of unconventional T cells. Nat Immunol. 2015;16(11):1114–23.

8. Treiner E, Duban L, Bahram S, Radosavljevic M, Wanner V, Tilloy F, et al. Selection of evolutionarily conserved mucosal-associated invariant T cells by MR1. Nature. 2003;422(6928):164–9.

9. Dusseaux M, Martin E, Serriari N, Péguillet I, Premel V, Louis D, et al. Human MAIT cells are xenobiotic-resistant, tissue-targeted, CD161hi IL-17-secreting T cells. Blood. 2011;117(4):1250–9.

10. Kjer-Nielsen L, Patel O, Corbett AJ, Le Nours J, Meehan B, Liu L, et al. MR1 presents microbial vitamin B metabolites to MAIT cells. Nature. 2012;491(7426):717–23.

11. Gherardin NA, Keller AN, Woolley RE, Le Nours J, Ritchie DS, Neeson PJ, et al. Diversity of T Cells Restricted by the MHC Class I-Related Molecule MR1 Facilitates Differential Antigen Recognition. Immunity. 2016;44(1):32–45.

12. Corbett AJ, Eckle SBG, Birkinshaw RW, Liu L, Patel O, Mahony J, et al. T-cell activation by transitory neo-antigens derived from distinct microbial pathways. Nature. 2014;509(7500):361–5.

13. Eckle SBG, Birkinshaw RW, Kostenko L, Corbett AJ, McWilliam HEG, Reantragoon R, et al. A molecular basis underpinning the T cell receptor heterogeneity of mucosal-associated invariant T cells. J Exp Med. 2014;211(8):1585–600.

14. Ussher JE, Bilton M, Attwod E, Shadwell J, Richardson R, Lara C De, et al. CD161++CD8+ T cells, including the MAIT cell subset, are specifically activated by IL-12+IL-18 in a TCR-independent manner. Eur J Immunol. 2014;44(1):195–203.

15. van Wilgenburg B, Scherwitzl I, Hutchinson EC, Leng T, Kurioka A, Kulicke C,

et al. MAIT cells are activated during human viral infections. Nat Commun.
2016;7:11653.

16.　Le Bourhis L, Martin E, Péguillet I, Guihot A, Froux N, Coré M, et al.
Antimicrobial activity of mucosal-associated invariant T cells. Nat Immunol.
2010;11(8):701–8.

17.　Georgel P, Radosavljevic M, Macquin C, Bahram S. The non-conventional
MHC class I MR1 molecule controls infection by Klebsiella pneumoniae in
mice. Mol Immunol. 2011;48(5):769–75.

18.　Meierovics A, Yankelevich WC, Cowley SC. MAIT cells are critical for optimal
mucosal immune responses during in vivo pulmonary bacterial infection. Proc
Natl Acad Sci U S A. 2013;110(33):E3119–28.

19.　Chua W-J, Truscott SM, Eickhoff CS, Blazevic A, Hoft DF, Hansen TH.
Polyclonal mucosa-associated invariant T cells have unique innate functions in
bacterial infection. Infect Immun. 2012;80(9):3256–67.

20.　Meierovics AI, Cowley SC. MAIT cells promote inflammatory monocyte
differentiation into dendritic cells during pulmonary intracellular infection. J Exp
Med. 2016;

21.　Broug-Holub E, Toews GB, van Iwaarden JF, Strieter RM, Kunkel SL, Paine R,
et al. Alveolar macrophages are required for protective pulmonary defenses in
murine Klebsiella pneumonia: elimination of alveolar macrophages increases
neutrophil recruitment but decreases bacterial clearance and survival. Infect
Immun. 1997;65(4):1139–46.

22.　Kooguchi K, Hashimoto S, Kobayashi A, Kitamura Y, Kudoh I, Wiener-Kronish
J, et al. Role of alveolar macrophages in initiation and regulation of
inflammation in Pseudomonas aeruginosa pneumonia. Infect Immun.
1998;66(7):3164–9.

23.　Kurioka A, Ussher JE, Cosgrove C, Clough C, Fergusson JR, Smith K, et al.
MAIT cells are licensed through granzyme exchange to kill bacterially
sensitized targets. Mucosal Immunol. 2015;8(2):429–40.

24.　Werno AM, Christner M, Anderson TP, Murdoch DR. Differentiation of
Streptococcus pneumoniae from nonpneumococcal streptococci of the
Streptococcus mitis group by matrix-assisted laser desorption ionization-time
of flight mass spectrometry. J Clin Microbiol. 2012;50(9):2863–7.

25.　Liuzzi AR, Kift-Morgan A, Lopez-Anton M, Friberg IM, Zhang J, Brook AC, et al.
Unconventional Human T Cells Accumulate at the Site of Infection in
Response to Microbial Ligands and Induce Local Tissue Remodeling. J
Immunol. 2016;197(6):2195–207.

26.　Kamphorst AO, Guermonprez P, Dudziak D, Nussenzweig MC. Route of
Antigen Uptake Differentially Impacts Presentation by Dendritic Cells and
Activated Monocytes. J Immunol. 2010;185(6):3426–35.

27.　Samstein M, Schreiber HA, Leiner IM, Sušac B, Glickman MS, Pamer EG.
Essential yet limited role for CCR2 $^{+}$ inflammatory monocytes during
Mycobacterium tuberculosis -specific T cell priming. Elife. 2013;2:e01086.

28.　Chen Z, Wang H, D'Souza C, Sun S, Kostenko L, Eckle SBG, et al. Mucosal-
associated invariant T-cell activation and accumulation after in vivo infection
depends on microbial riboflavin synthesis and co-stimulatory signals. Mucosal
Immunol. 2016;doi: 10.1038/mi.2016.39.

29.　Turtle CJ, Delrow J, Joslyn RC, Swanson HM, Basom R, Tabellini L, et al.

Innate signals overcome acquired TCR signaling pathway regulation and
govern the fate of human CD161(hi) CD8α⁺ semi-invariant T cells. Blood.
2011;118(10):2752–62.

30. Slichter CK, McDavid A, Miller HW, Finak G, Seymour BJ, McNevin JP, et al.
Distinct activation thresholds of human conventional and innate-like memory T
cells. JCI insight. 2016;1(8).

31. Martner A, Skovbjerg S, Paton JC, Wold AE. Streptococcus pneumoniae
autolysis prevents phagocytosis and production of phagocyte-activating
cytokines. Infect Immun. 2009;77(9):3826–37.

32. Ussher JE, Bilton M, Attwod E, Shadwell J, Richardson R, Lara C De, et al.
CD161 ++ CD8 + T cells , including the MAIT cell subset , are specifically
activated by IL-12 + IL-18 in a TCR-independent manner. 2013;1–9.

33. Morens DM, Taubenberger JK, Fauci AS. Predominant Role of Bacterial
Pneumonia as a Cause of Death in Pandemic Influenza: Implications for
Pandemic Influenza Preparedness. J Infect Dis. 2008;198(7):962–70.

34. Loh L, Wang Z, Sant S, Koutsakos M, Jegaskanda S, Corbett AJ, et al. Human
mucosal-associated invariant T cells contribute to antiviral influenza immunity
via IL-18-dependent activation. Proc Natl Acad Sci U S A.
2016;113(36):10133–8.

35. Li W, Moltedo B, Moran TM. Type I interferon induction during influenza virus
infection increases susceptibility to secondary Streptococcus pneumoniae
infection by negative regulation of γδ T cells. J Virol. 2012;86(22):12304–12.

36. Walker LJ, Kang YH, Smith MO, Tharmalingham H, Ramamurthy N, Fleming
VM, et al. Human MAIT and CD8αα cells develop from a pool of type-17
precommitted CD8 + T cells. Blood. 2012;119(2):422–33.

37. Koay H-F, Gherardin NA, Enders A, Loh L, Mackay LK, Almeida CF, et al. A
three-stage intrathymic development pathway for the mucosal-associated
invariant T cell lineage. Nat Immunol. 2016;17(11):1300–11.

38. Novak J, Dobrovolny J, Novakova L, Kozak T. The decrease in number and
change in phenotype of mucosal-associated invariant T cells in the elderly and
differences in men and women of reproductive age. Scand J Immunol.
2014;80(4):271–5.

39. Smith NM, Wasserman GA, Coleman FT, Hilliard KL, Yamamoto K, Lipsitz E,
et al. Regionally compartmentalized resident memory T cells mediate naturally
acquired protection against pneumococcal pneumonia. Mucosal Immunol.
2017;

40. Liu S, Sela S, Cohen G, Jadoun J, Cheung A, Ofek I. Insertional inactivation of
Streptolysin S expression is associated with altered riboflavin metabolism
inStreptococcus pyogenes. Microb Pathog. 1997;22(4):227–34.

41. Chen J, Shen J, Solem C, Jensen PR. Oxidative stress at high temperatures in
Lactococcus lactis due to an insufficient supply of Riboflavin. Appl Environ
Microbiol. 2013;79(19):6140–7.

42. Gibbs A, Leeansyah E, Introini A, Paquin-Proulx D, Hasselrot K, Andersson E,
et al. MAIT cells reside in the female genital mucosa and are biased towards
IL-17 and IL-22 production in response to bacterial stimulation. Mucosal
Immunol. 2016;doi: 10.1038/mi.2016.30.

43. Martins ER, Pessanha MA, Ramirez M, Melo-Cristino J, Portuguese Group for
the Study of Streptococcal Infections and the PG for the S of S. Analysis of

group B streptococcal isolates from infants and pregnant women in Portugal
revealing two lineages with enhanced invasiveness. J Clin Microbiol.
2007;45(10):3224–9.

44.   McGee L, McDougal L, Zhou J, Spratt BG, Tenover FC, George R, et al.
Nomenclature of major antimicrobial-resistant clones of Streptococcus
pneumoniae defined by the pneumococcal molecular epidemiology network. J
Clin Microbiol. 2001;39(7):2565–71.

45.   PMEN :: Pneumococcal Molecular Epidemiology Network [Internet]. Available
from: http://www.pneumogen.net/pmen/

46.   MLST allelic profiles, pbp gene profiles and macrolide resistance determinants
for clones [Internet]. Available from:
http://web1.sph.emory.edu/PMEN/pmen_table2.html

47.   Jolley KA, Maiden MC. BIGSdb: Scalable analysis of bacterial genome
variation at the population level. BMC Bioinformatics. 2010;11(1):595.

48.   Brueggemann AB, Harrold CL, Rezaei Javan R, van Tonder AJ, McDonnell AJ,
Edwards BA. Pneumococcal prophages are diverse, but not without structure
or history. Sci Rep. 2017;7:42976.

49.   Zerbino DR, Birney E. Velvet: Algorithms for de novo short read assembly
using de Bruijn graphs. Genome Res. 2008;18(5):821–9.

50.   Boetzer M, Pirovano W. SSPACE-LongRead: scaffolding bacterial draft
genomes using long read sequence information. BMC Bioinformatics.
2014;15(1):211.

51.   Boetzer M, Pirovano W. Toward almost closed genomes with GapFiller.
Genome Biol. 2012;13(6):R56.

52.   Jolley KA, Bliss CM, Bennett JS, Bratcher HB, Brehony C, Colles FM, et al.
Ribosomal multilocus sequence typing: universal characterization of bacteria
from domain to strain. Microbiology. 2012;158(Pt 4):1005–15.

53.   Francisco AP, Vaz C, Monteiro PT, Melo-Cristino J, Ramirez M, Carrio JA.
PHYLOViZ: phylogenetic inference and data visualization for sequence based
typing methods. BMC Bioinformatics. 2012;13(1):87.

54.   Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, et al. The SEED
and the Rapid Annotation of microbial genomes using Subsystems Technology
(RAST). Nucleic Acids Res. 2014;42:D206-14.

55.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics.
2014;30(14):2068–9.

56.   European Nucleotide Archive [Internet]. Available from:
http://www.ebi.ac.uk/ena

57.   Streptococcus pneumoniae MLST website [Internet]. Available from:
http://pubmlst.org/spneumoniae/

58.   GenBank [Internet]. Available from: https://www.ncbi.nlm.nih.gov/genbank/

59.   Keith Jolley. Ribosomal Multilocus Sequence Typing (rMLST) [Internet].
Available from: https://pubmlst.org/rmlst/

60.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat
Methods. 2012;9(4):357–9.

61.   Anders S, Huber W. Differential expression analysis for sequence count data.
Genome Biol. 2010;11(10):R106.

62.   Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M. KEGG: Kyoto
Encyclopedia of Genes and Genomes. Nucleic Acids Res. 1999;27(1):29–34.

63. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al.
Geneious Basic: An integrated and extendable desktop software platform for
the organization and analysis of sequence data. Bioinformatics.
2012;28(12):1647–9.
64. Pond SLK, Frost SDW, Muse S V. HyPhy: hypothesis testing using
phylogenies. Bioinformatics. 2005;21(5):676–9.
65. Delport W, Poon AFY, Frost SDW, Kosakovsky Pond SL. Datamonkey 2010: a
suite of phylogenetic analysis tools for evolutionary biology. Bioinformatics.
2010;26(19):2455–7.
66. Pond SLK, Frost SDW. Datamonkey: rapid detection of selective pressure on
individual sites of codon alignments. Bioinformatics. 2005;21(10):2531–3.
67. Marchler-Bauer A, Lu S, Anderson JB, Chitsaz F, Derbyshire MK, DeWeese-
Scott C, et al. CDD: a Conserved Domain Database for the functional
annotation of proteins. Nucleic Acids Res. 2011;39(Database):D225–9.
68. Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets
of protein or nucleotide sequences. Bioinformatics. 2006;22(13):1658–9.
69. Arnaoudova E, Jaromczyk JW, Moore N, Schardl CL, Yoshida R. Phylotree - a
toolkit for computing experiments with distance-based methods for genome
coevolution. BMC Bioinformatics. 2010;11(Suppl 4):P6.
70. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display
and annotation of phylogenetic and other trees. Nucleic Acids Res.
2016;44(W1):W242–5.
71. Inkscape [Internet]. Available from: https://inkscape.org/en/
975
976

977 **Supporting information captions**

978 Supplementary Table 1. Descriptive data for the 571 *Streptococcus pneumoniae*

979 genomes included in this study.

980 Supplementary Table 2. Descriptive data for the 824 non-pneumococcal

981 *Streptococcus* species genomes included in this study.

Table 1. Description of the four riboflavin operon gene sequences within 571 pneumococcal genomes.

Gene	Present (%)[a]	Nucleotide % pairwise identity	Amino acid % pairwise identity	Mean dN/dS	Gene annotation
ribD	559 (98.2%)	99.6%	99.8%	0.36	Diaminohydroxyphosphoribosylamino-pyrimidine deaminase
ribE	561 (98.2%)	99.6%	99.9%	0.28	riboflavin synthase
ribA	561 (98.2%)	99.7%	99.9%	0.42	3,4-dihydroxy-2-butanone-4-phosphate synthase
ribH	559 (98.2%)	99.5%	99.1%	0.12	6,7-dimethyl-8-ribityllumazine synthase

a. Two genomes possessed sequence assembly gaps in *ribD* and *ribH,* and 10 genomes were missing all four of the riboflavin genes (see Results).

Figure 1

A

B

Figure 2

Figure 3

Figure 4