Affiliative social relationships but not dominance rank affects endo-parasite

product excretion in a cooperatively breeding bird species

Claudia A.F. Wascher[1], Daniela Canestrari[2] & Vittorio Baglione[3]

[1] School of Life Sciences, Anglia Ruskin University, United Kingdom

[2] Unidad Mixta de Investigación en Biodiversidad, Campus de Mieres, University of

Oviedo, Spain

[3] Departamento de Biodiversidad y Gestión Ambiental, Universidad de León, Spain

14 Short title: Social relationships reduce parasite burden in crows

16 Corresponding author (present address):

17 Claudia A.F. Wascher: Department of Biology, Anglia Ruskin University, United

18 Kingdom; Phone: ++4369912381419; e-mail: claudia.wascher@gmail.com

Abstract

In group living animals behavioural interactions with conspecifics strongly modulate an

individual's physiological stress response. A secure social embedding, with stable social

relationships is expected to dampen an individual's stress response which is in turn will

affect the immune system and health. Ultimately, positive health effects of stable social

bonds may contribute to maintain group living. We investigated whether, in

cooperatively breeding carrion crows (*Corvus corone*), the quality of social

relationships have measurable effects on patterns of gastrointestinal parasite product

excretion. We repeatably collected behavioural data on dyadic social interactions and

individual droppings to quantify parasite products, from 34 individuals in a captive

population of carrion crows in northern Spain. Individuals with strong social bonds and

living with more related individuals in a group, excreted less samples containing

coccidian oocysts. Sex, group size, number of social bonds and dominance rank had no

significant effect on number of samples containing coccidian oocysts. Also, the

likelihood to excrete samples containing nematode eggs was not affected by any of the

factors we considered. Our data presents evidence for affiliative social relationships to

be linked to coccidian oocyst excretion, indicating that the social environment has

strong fitness implications. This suggests that next to humans and other primate species,

social relationships may be important in driving the evolution of complex sociality in a

larger variety of taxa, including birds.

Key words: carrion crows, coccidian oocysts, dominance rank, group size, parasites,

social bonds.

Background

In group living animals the social environment directly affects an individual's physiology and health (1–4). Adverse effects are caused for example by increased competition and aggressive behaviour (5,6), which affect especially low-ranking individuals (7,8). Further, large group sizes increase the exposure to directly transmitted parasites (9,10) and social interaction patterns can also facilitate pathogen transmission (11,12).

Affiliative behaviour and reliable social allies on the other hand, have a stress-reducing effect (13–16). In several mammalian species, stable social bonds have a fitness enhancing effect (17–19). Social relationships in female chacma baboons (*Papio hamadryas ursinus*) help coping with stressful events and increase longevity (20), whereas social bonds in male Assamese macaques (*Macaca assamensis*) and feral horses (*Equus ferus caballus*) are directly linked to the number of offspring sired (18).

In humans, a rich social network and social support yields measurable positive consequences for health (21–27). Recent research, however, has shown that complex sociality associated with the formation of strong, long-lasting social bonds, is not unique to humans and closely related primate species (28,29). This suggests the fascinating hypothesis that positive health effects of social bonds may contribute to maintain stable group living in a large variety of taxa and types of social organizations (30).

Among birds, corvids show complex social relationships. Common ravens (*Corvus corax*), for example, are capable of forming valuable relationships, not only within

reproductive pairs but also with social partners in their groups (31,32). Such bonds are

characterized by low levels of aggression and high levels of affiliative behaviours

exchanged in a reciprocal way over extended periods of time (33). Within such valuable

relationships, individuals support each other in agonistic encounters (32,34) and share

information and resources (32,35,36), whereas they typically act competitively with

other non-bonded conspecifics (37,38). In the present study we investigated the benefits

of social bonds in carrion crows (*Corvus corone*). In most European populations carrion

crows form socially monogamous pairs during the breeding season (39,40) whereas in

northern Spain crows live in stable social groups of up to nine individuals, consisting of

the breeding pair and retained offspring as well as male immigrants (41). Group living

in carrion crows is based on kinship (41) and cooperation in nestling provisioning (42)

and territory defence (Hillemann, et al. in prep.). Dominant males are more aggressive

towards same sex group members compared to females (43) and are less aggressive

towards related individuals than to unrelated ones (44). In captive crows, males are

more attentive towards non-kin compared to kin individuals (45) and, unlike females,

show less avoidance towards the scent of a stressed familiar individual, compared to a

stressed unfamiliar one, which might reflect a stronger willingness to provide social

support (46).

In this study, we investigated whether the social environment and the quality of an

individual's social relationships affects prevalence of gastrointestinal parasites in

captive groups of carrion crows that varied in size and composition. In particular, we

focused on the pattern of excretion of oocysts of coccidian protozoans and different

nematode species, mainly *Syngamus trachea* and *Capillaria* sp.. Coccidian oocysts are

among the most common endoparasites of birds (47,48) and are widespread among

corvids, including carrion crows (49–51). Infections with large numbers of coccidia

clinically manifest as 'coccidiosis', which damages the intestinal tract (52) and severely

affects individual body condition, longevity and fecundity by inhibiting the uptake of

essential dietary components (53,54). The pathogenicity is well documented in poultry

(55–57), and further evidence is accumulating in wild bird species (e.g. 58,59). Coccidia

are transmitted via the faecal-oral route, when oocysts undergo sporogony and become

infectious (typically within 24 hours after excretion) (56). Similarly, different nematode

species are described as regular endo-parasites of different corvid species (60,61).

Syngamus trachea is usually located in the trachea of birds and can lead to severe

respiratory distress and pulmonary lesions (62,63). *Capillaria* sp. infect the intestinal

tract and clinical symptoms are weight loss, diarrhoea, regurgitation, anaemia and oral

necrotic plaques (64). Different nematode species infect their avian hosts via a direct

life cycle, when embryonated eggs are ingested. Nematode eggs have a prepatent period

of several weeks (65).

Here we address three hypotheses: 1. Group size hypothesis: parasite product

excretion depends on the size of the group, with infections spreading more easily in

crowded conditions; 2. Social bonds hypothesis: parasite product excretion depend on

the quality of the social relationships that an individual maintains, with individuals with

strong social bonds being less likely to be infected than individuals with weak social

bonds. The presence of relatives in the group might also show a positive effect by

reducing the occurrence of stressful conflicts and competition (44) that can impair the

ability to fight against parasites (66). 3. Dominance rank hypothesis: parasite product

patterns depend on the position along the hierarchy. Indeed, dominance rank is known

to affect individual social stress, weakening the immune system either in subordinates

or dominants, depending the relative allostatic load of social status in the respective

species (67). In addition, we expect sex differences, because males are involved in more

competitive interactions than females within their groups (43) and are probably more

exposed to the negative consequences of social stress.

Results

Social bonds and parasitological examination

We observed 50 bonded dyads (out of 312 dyads), 23 of which were male-female dyads (4 between related individuals and 19 between unrelated individuals), 12 were male-male (7 between related and 5 between unrelated individuals) and 15 female-female (5 related, 10 unrelated). On average, males and females had 1.136 ± 1.206 and 1.315 ± 0.945 bonded partners respectively. Mean CSI between bonded dyads was 2.698 ± 2.182 for male-female dyads, 2.299 ± 1.373 for female-female bonds and 3.035 ± 3.388 for male-male bonds.

Occurrence of coccidian oocysts

Overall, 104 out of 416 samples from captive crows contained coccidian oocysts. Coccidian oocysts occurred in 69 out of 220 samples in males (31 %) and 35 out of 196 samples in females (18 %). The maximum CSI value and the number of related individuals in the group had a significant influence on the occurrence of coccidian oocysts. Crows with a strong affiliative relationship (maximum CSI value: estimate $\pm$ SE $= -1.25 \pm 0.557$, $z = 2.237$, $p = 0.025$) and living with a larger number of related individuals (estimate $\pm$ SE $= -1.229 \pm 0.467$, $z = 2.623$, $p = 0.008$) were less likely to excrete samples containing coccidian oocysts (Figures 1 and 2), whereas the size of the group (estimate $\pm$ SE $= -0.384 \pm 0.5$, $z = 0.765$, $p = 0.443$), number of social bonds (estimate $\pm$ SE $= 0.723 \pm 0.512$, $z = 1.412$, $p = 0.157$) and dominance rank (estimate $\pm$ SE $= 0.326 \pm 0.463$, $z = 0.702$, $p = 0.482$) did not have a significant effect. Males showed a weak trend to excrete more samples containing coccidian oocysts compared to females (estimate $\pm$ SE $= -0.811 \pm 0.461$, $z = 1.753$, $p = 0.079$).

155   *Occurrence of nematode eggs*

156   Overall, 52 out of 416 samples contained nematode eggs. None of the factors

157   investigated significantly affected excretion patterns of nematode eggs (supplementary

158   Table 4), in fact the null model was amongst the best models in the candidate set (Table

159   1).

Discussion

In the present study, we have shown that carrion crows with closely bonded social partners (high CSI scores) proved less likely to excrete coccidian oocysts, which represent an important threat for health in birds (53,54). Health correlates of high quality social ties have been reported in humans (*e.g.* 76) and a large body of research has focused on the causal psychological and biological processes that underlie this strong link (21,69). Similar evidence has been recently found in non-human primates (18,20), suggesting that the tendency to form strong social ties may have deep phylogenetic roots (70). Social complexity, however, is not limited to primate species. Indeed, coalition formation has been reported in several mammalian and bird species, e.g. African elephants (*Loxodonta africanus)* (71), spotted hyenas (*Crocutta crocutta)* (72), bottlenose dolphins (*Tursiops aduncus)* (73), ravens (74), and carrion crows (41). Our study uncovered, for the first time, that high quality social relationships may reduce the occurrence of parasites, showing immediate positive effects that parallel those of primates (including humans) and suggesting a route towards advanced sociality that may be common to a variety of taxa.

Cooperatively breeding groups of carrion crows form through two different mechanisms, namely delayed dispersal of offspring, which remain in the natal territory with their parents and siblings (75), and formation of long lasting social bonds among distant relatives, most often males, that share all-purpose territories and frequently mate polyandrously (41,76). These bonds typically form when an immigrant male joins an established family group and allies with the same sex resident breeder, to whom it is related (41). Cooperative breeding, with group members working together to raise the brood, can only arise once stable groups have formed (77). Both offspring delayed

dispersal and bonding between adults males are therefore necessary preceding steps that

eventually lead to cooperation, which has to be considered a consequence rather than

the cause for the formation of the social group (78–80). In other words, to understand

why carrion crows, as well as any other bird species, form groups we need to

understand the advantages that sociality conveys, independently of the payoff derived

from cooperation at the nest. In carrion crows, as well as in many other cooperatively

breeding species, the benefits of delayed dispersal for offspring are well studied (81),

but little is known about the advantages of forming long lasting social bonds outside the

nuclear family. We argue that the health benefits of high-quality relationships extend

beyond the bonds between parents and offspring or reproductive partners, and may be

associated with a reduction of harmful endo-parasites and may be an important factor

for establishing stable relationships in groups.

Besides the positive effect of strong social bonds, we found that the number of

relatives in the group also affected parasite product excretion in carrion crows, with

individuals living in groups with more related individuals being less likely to excrete

droppings containing coccidian oocysts. Sociality is based on kinship in wild

cooperatively breeding carrion crows. Offspring remain in the natal territory with their

parents for years and more distant relatives are actively recruited to form cooperative

alliances with the resident breeders (Baglione et al. 2003). As a result, social groups in

cooperatively breeding carrion crow are extended families, comprising members with

different degree of relatedness. Indirect fitness benefits are known to be a primary

driver of kin-based sociality in many taxa of animals (83). However, our results now

show that living with kin can also accrue immediate direct benefits to carrion crows

through reduced infection by coccidia.

Increased exposure to parasites and disease transmission is considered as one of

the major disadvantages of group living. Group size is usually positively related to

parasite burden, which is seen as major selective force in group living animals (10). In

our study, group size did not have a significant effect on frequency of samples

containing coccidian oocysts. Therefore, contrary to the expectation that probability of

pathogen transmission increases with group size, we found group composition, *i.e.*

number of related individuals and the quality of social relationships to mediate an

individual's parasite product excretion, likely via dampening of the physiological stress

response which might suppress an individual's immune system. This is also supported

by a recent study in Japanese macaques (*Macaca fuscata*), which shows that individuals

with more affiliative social partners to have fewer ectoparasites (84).

We only found effects of social relationships and number of related individuals

in coccidia, but not in nematode species. One reason could be that the lower occurrence

of nematode eggs compared to coccidian oocysts in the samples (12 % of samples

contained nematode eggs and 24 % of samples contained coccidian oocysts) limited the

power of our analysis. However, the quicker life cycle of coccidians, with a prepatent

period of approximately one week, compared to nematodes, *e.g.* capillaria, with a pre-

patent period of 3 - 4 weeks (64) could also play a role, as short-term changes in the

immune system, probably in response to suppressive effects of glucocorticoids in

response to social interactions, might have stronger effects on coccidia compared to

nematodes.

Conclusions

Social complexity has deep phylogenetic roots, being widespread in a wide

variety of mammals and birds. Understanding its evolution, however, is difficult

because of our scant knowledge of the fitness consequences of sophisticated social

behaviour across taxa. We are just starting to fill the gaps, and this study shows a

measurable positive effect of the quality of social relationships in a bird species that

may trigger social bonding, setting the stage for a complex form of cooperation

(cooperative breeding). Importantly, our results parallel previous findings in primate

species, suggesting a general pattern in vertebrates for the evolution of sociality.

Methods

Study subjects and ethics statement

We collected data over four different periods (November 2008 - January 2009,

January - February 2013, May - June 2013, January - February 2014), from a total of 34

crows (16 females, 18 males), which were housed in a large outdoor aviary (30 x 12 x 6

m) in Northern Spain (Navafría, León, 42°36'33 N 5°26'56 W). The aviary comprised

four major compartments (12 x 6 x 3 m) connected through eight smaller testing

compartments (3 x 3 x 3 m). The major compartments were equipped with wooden

perches, natural vegetation and rocks. An enriched diet consisting of fruit, vegetables,

bread, meat and milk products was provided on a daily basis. Water and dry food were

available *ad libitum*. Keeping crows in captivity was authorized by Junta de Castilla y

León (núcleo zoológico 005074).

Birds were kept in groups that mirrored the social aggregations that naturally occur

in the wild: (A) 'Flock': three or more juvenile individuals; in 2008 some juveniles

were initially kept in 'sibling pairs' and subsequently merged into a juvenile flock; (B)

'Pairs': adult male and adult female; in some cases adults were also kept in trios; (C)

'Cooperative Family': a reproductive pair with its own already independent offspring.

Group composition (supplementary Table 1), however, varied during the course of the

study due to the death of some individuals.

Behavioural data

A total of 1045 individual focal observations have been recorded. Each observation

lasted 5 minutes, and all occurring behaviours were recorded. For this study, we focused

on frequencies of agonistic behaviour (threat, chase flight, and fight) and affiliative

behaviours (approach, allopreening and sitting in contact). The identity of the

interacting individuals was noted, as well as their role (initiator/receiver). All

behavioural observations were recorded on video and analysed by CAFW

(supplementary Table 1).

Composite sociality index

For each period of data recording, we calculated a composite sociality index (CSI)

for each crow dyad within a group according to Silk et al. 2010 (20). A CSI was

constructed from three affiliative behaviours: approach, contact sit and allopreen. Note

that we used mean values per observation instead of absolute numbers because the

number of focal observations varied among individuals (supplementary Table 1). For

each period of data collection, we also calculated for each individual the maximum CSI

value among all the dyads that involved that particular bird. Dyads with CSI higher than

the average of the entire sample were classified as 'bonded', and we counted the

number of bonded group members for each individual.

Normalized David's Score

We calculated the success of individuals in agonistic encounters using the corrected

normalized David's Score (nDS), which considers the proportion of wins and losses,

corrected for the chance occurrence (85). In order to compare individuals between

different groups, we calculated the relative difference in nDS among individuals in each

group (ΔnDS), with 0 being the most dominant individual (supplementary Table 1).

Parasitological examination

During the entire study period, 441 individual droppings were collected, directly

after defecation, so that each sample could be assigned to a particular individual

(supplementary Table 1). Samples were examined for eggs and oocysts of intestinal

parasites. In 2008, we used a modified version of the flotation method (64) to examine

the occurrence of parasite products in the droppings. Fresh droppings (0.1 g) were

suspended in a 2 ml collecting tube with 1 ml saturated saline. Collection tubes were

shaken for 10 seconds and afterwards centrifuged for 5 minutes at 3000 rpm. After

centrifugation, the collection tubes were filled with saline solution and a cover slip (18 x

18 mm) was positioned onto the tube. The high density of the saline solution causes the

parasite eggs and oocysts to float up and be caught on the cover slip (86). After 10

minutes, the cover slip was moved onto an object slide and the parasite eggs and

oocysts were identified and counted. From January 2013 onwards, we used a McMaster

counting chamber to examine the occurrence of parasite products in the droppings. The

entire sample was weighed, then diluted with 3 ml saturated NaCl solution per 0.1 g of

droppings and thoroughly mixed. Afterwards, the solution was filled into both

McMaster counting chambers. After 10 minutes of resting period we identified the

parasite eggs and oocysts in both chambers. We used a binocular with 100-fold and

400-fold amplification to identify parasite products. We found Coccidian oocysts, eggs

of several nematode species (*Capillaria sp*., *Ascarida sp*., *Syngamus sp., Heterakis sp.,*

Trichostrongylus tenius) and cestode eggs to a varying degree. The proportion of

positive samples was highest for coccidian oocysts (24 %, n = 441). Nematode eggs

were found in 12 % of samples and cestode eggs where only found in less than 1 % of

samples. We therefore limited our statistical analysis to coccidian occysts and

nematodes.

Data analysis

We analysed the occurrence (presence/absence) of coccidian oocysts and

nematode eggs in the droppings. We used generalized linear mixed models (GLMMs)

with binomial error distributions and logit link functions. GLMMs were fitted by

maximum likelihood. All models were calculated in R v. 3.0.2 (87) using the

glmmADMB package (version 0.7.7; 88). We based our model selection on second-

order Akaike's Information Criterion values (AICc; 89). We calculated the difference

between the best model and each other possible model (ΔAICc) and ranked the model

combinations according to their ΔAICc, which provides an evaluation of the overall

strength of each model in the candidate set. Multiple models qualified as the similarly

good models, *i.e.* ΔAICc ≤ 2 (90,91) and therefore we applied a model averaging

approach, which calculates model averaged parameters using the MuMIn package

(version 1.15.6). Full statistical models are presented in the supplementary materials

(supplementary Table 2 and 4). Maximum CSI value, group size, number of bonds,

number of related individuals, normalized David's score (ΔnDS) and sex were included

as explanatory variables. Further, we ran a model including two-way interactions

between sex and the fixed factors. All interactions were non-significant (supplementary

Table 3 and 5) and subsequently removed from the model (92). For each model, we

fitted individual identity as a random term to control for the potential dependence

associated with multiple samples from the same individuals.

Author contributions

C.A.F.W. designed the study, collected the data, performed lab identification of

parasites and analysed the data; all the authors wrote the paper.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the

corresponding author on reasonable request.

Acknowledgments

We thank Kurt Kotrschal for feedback and discussion, Hugo Robles for statistical

advice and Friederike Hillemann for help with droppings collection. This work was

supported by the Spanish Plan Nacional I+D, project CGL2011-27260 and CGL2016-

77636-P, to VB and a KWA ('kurzfristiges wissenschaftliches Auslandsstipendium')

stipend of the University of Vienna to CAFW.

358
359

References

1.   DeVries AC, Glasper ER, Detillion CE. Social modulation of stress responses. Physiol Behav. 2003;79(3):399–407.
2.   Holst D Von. The concept of stress and its relevance for animal behavior. Adv Study Behav. 1998;37:1–131.
3.   Hanley KA, Stamps JA. Does corticosterone mediate bidirectional interactions between social behaviour and blood parasites in the juvenile black iguana, Ctenosaura similis? Anim Behav. 2002;63(2):311–22.
4.   Sheridan JF, Dobbs C, Brown D, Zwilling B. Psychoneuroimmunology: stress effects on pathogenesis and immunity during infection. Clin Microbiol Rev. 1994;7(2):200–12.
5.   Hawley DM, Lindström K, Wikelski M. Experimentally increased social competition compromises humoral immune responses in house finches. Horm Behav. 2006;49(4):417–24.
6.   Chester EM, Bonu T, Demas GE. Social defeat differentially affects immune responses in Siberian hamsters (Phodopus sungorus). Physiol Behav. 2010;101(1):53–8.
7.   Ungerfeld R, Correa O. Social dominance of female dairy goats influences the dynamics of gastrointestinal parasite eggs. Appl Anim Behav Sci. 2007;105(1):249–53.
8.   Zuk M, Kim T, Robinson S, Johnsen T. Parasites influence social rank and morphology, but not mate choice, in female red junglefowl, Gallus gallus. Anim Behav. 1998;56(2):493–9.
9.   Loehle C. Social barriers to pathogen transmission in wild animal populations. Ecology. 1995;76(2):326–35.
10.   Côté IM, Poulin R. Parasitism and group size in social animals: a meta-analysis. Behav Ecol. 1995;6(2):159.165.
11.   Drewe JA. Who infects whom? Social networks and tuberculosis transmission in wild meerkats. Proc R Soc B Biol Sci. 2010;277(1681):633–42.
12.   Fouchet D, Verrier D, Ngoubangoye B, Souquiere S, Makuwa M, Kazanji M, et al. Who infects whom? Social networks and tuberculosis transmission in wild meerkats. Proc R Soc B Biol Sci. 2012;279(1742):3426–35.
13.   Frigerio D, Weiss B, Dittami J, Kotrschal K. Social allies modulate corticosterone excretion and increase success in agonistic interactions in juvenile hand-raised graylag geese (Anser anser). Can J Zool. 2003;81:1746–54.
14.   Stöwe M, Bugnyar T, Schloegl C, Heinrich B, Kotrschal K, Möstl E. Corticosterone excretion patterns and affiliative behavior over development in ravens (Corvus corax). Horm Behav. 2008;53(1):208–16.
15.   Sachser N, Dürschlag M, Hirzel D. Social relationships and the management of stress. Psychoneuroendocrinology. 1998;23(8):891–904.
16.   Young C, Majolo B, Schülke O, Ostner J. Male social bonds and rank predict supporter selection in cooperative aggression in wild Barbary macaques. Anim Behav; 2014;95:23–32.
17.   Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, et al. The benefits of social capital: close social bonds among female baboons enhance offspring survival. Proc Biol Sci. 2009;276(1670):3099–104.
18.   Schülke O, Bhagavatula J, Vigilant L, Ostner J. Social bonds enhance reproductive success in male macaques. Curr Biol. 2010;20(24):2207–10.
19.   Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, et al. Strong and consistent social bonds enhance the longevity of female baboons.

410   Curr Biol. 2010;20(15):1359–61.

411   20.   Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, et al.
412   Female chacma baboons form strong, equitable, and enduring social bonds.
413   Behav Ecol Sociobiol. 2010;64(11):1733–47.

414   21.   Berkman LF, Glass T. Social integration, social networks, social support, and
415   health. In: Berkman LF, Kawachi I, editors. Social Epidemiology. Oxford, UK:
416   Oxford University Press; 2000. p. 137.

417   22.   Theorell T, Blomkvist V, Jonsson H, Schulman S, Berntorp E, Stigendal L.
418   Social support and the development of immune function in human
419   immunodeficiency virus infection. Psychosom Med. 1995;57(1):200–12.

420   23.   Kiecolt-Glaser JK, McGuire L, Robles TF, Glaser R. Psychoneuroimmunology:
421   Psychological influences on immune function and health. J Consult Clin Psychol.
422   2002;70(3):537–47.

423   24.   Cobb S. The role of social relations in health promotion. Psychosom Med.
424   1976;38(5):300–14.

425   25.   Uchino BN, Cacioppo JT, Kiecolt-Glaser JK. The relationship between social
426   support and physiological processes: a review with emphasis on underlying
427   mechanisms and implications for health. Psychol Bull. 1996;119(3):488–531.

428   26.   Berkman LF. The role of social relations in health promotion. Psychol Bull.
429   1995;57(3):245–54.

430   27.   Cassel J. The contribution of the social environment to host resistance.
431   Psychosom Med. 1976;104(2):107–23.

432   28.   Shultz S, Dunbar RIM. Social bonds in birds are associated with brain size and
433   contingent on the correlated evolution of life-history and increased parental
434   investment. Biol J Linn Soc. 2010;100(1):111–23.

435   29.   Pollen AA, Dobberfuhl AP, Scace J, Igulu MM, Renn SCP, Shumway CA, et al.
436   Environmental complexity and social organization sculpt the brain in Lake
437   Tanganyikan cichlid fish. Brain Behav Evol. 2007;70(1):21–39.

438   30.   Silk JB. The adaptive value of sociality in mammalian groups. Philos Trans R
439   Soc B. 2007; 362(1480):539–59.

440   31.   Heinrich B. Conflict, cooperation, and cognition in the common raven. Adv
441   Study Behav. 2011;43:189–237.

442   32.   Fraser ON, Bugnyar T. Reciprocity of agonistic support in ravens. Anim Behav.
443   2012;83(1):171–7.

444   33.   Fraser ON, Bugnyar T. The quality of social relationships in ravens. Anim
445   Behav. 2010;79(4):927–33.

446   34.   Emery NJ, Seed AM, von Bayern AMP, Clayton NS. Cognitive adaptations of
447   social bonding in birds. Philos Trans R Soc B. 2007;362(1480):489–505.

448   35.   Bugnyar T, Kijne M, Kotrschal K. Food calling in ravens: are yells referential
449   signals? Anim Behav. 2001;61(5):949–58.

450   36.   de Kort SR, Emery NJ, Clayton NS. Food sharing in jackdaws, Corvus
451   monedula: what, why and with whom? Anim Behav. 2006;72(2):297–304.

452   37.   Bugnyar T, Heinrich B. Pilfering ravens, Corvus corax, adjust their behaviour to
453   social context and identity of competitors. Anim Cogn. 2006;9(4):369–76.

454   38.   Bugnyar T, Heinrich B. Ravens, Corvus corax, differentiate between
455   knowledgeable and ignorant competitors. Proc R Soc B Biol Sci.
456   2005;272(1573):1641–6.

457   39.   Glutz von Blotzheim UN. Handbuch der Vögel Mitteleuropas. Glutz von
458   Blotzheim UN, editor. Wiesbaden: Aula-Verlag; 1985.

40.   Meide M. Raben- und Nebelkrähe. Magdeburg: Westarp Wissenschaften; 1984.

41. Baglione V, Canestrari D, Marcos JM, Ekman J. Kin selection in cooperative
alliances of carrion crows. Science. 2003;300(5627):1947–9.
42. Canestrari D, Chiarati E, Marcos JM, Ekman J, Baglione V. Helpers but not
breeders adjust provisioning effort to year-round territory resource availability in
carrion crows. Anim Behav. 2008;76(3):943–9.
43. Chiarati E, Canestrari D, Vera R, Marcos JM, Baglione V. Linear and Stable
Dominance Hierarchies in Cooperative Carrion Crows. Ethology.
2010;116(4):346–56.
44. Chiarati E, Canestrari D, Vila M, Vera R, Baglione V. Nepotistic access to food
resources in cooperatively breeding carrion crows. Behav Ecol Sociobiol.
2011;65:1791-1800.
45. Wascher CAF, Núñez Cebrián C, Valdez JW, Canestrari D, Baglione V. Social
factors modulating attention patterns in carrion crows. Behaviour.
2014;151(5):555–72.
46. Wascher CAF, Heiss RS, Baglione V, Canestrari D. Behavioural responses to
olfactory cues in carrion crows. Behav Proc. 2015;111:1–5.
47. Page CD, Haddad K. Coccidial infections in birds. Semin Avian Exot Pet Med.
1995;4(3):138–44.
48. López G, Figuerola J, Soriguer R. Time of day, age and feeding habits influence
coccidian oocyst shedding in wild passerines. Int J Parasitol. 2007;37(5):559–64.
49. Poon SK, Chew WK. Isospora corvi Ray, Shivnani, Oommen and Bhaskaran,
1952 from the common house crow (Corvus splendens Vieillot) of Selangor,
peninsular Malaysia. Folia Parasitol. 1991;38(3):201–7.
50. Upton SJ, Langen TA, Wright TF. A new species of Isospora Schneider, 1881
(Apicomplexa: Eimeriidae) from the white-throated magpie jay Calocitta
formosa (Passeriformes: Corvidae) from Costa Rica. Syst Parasitol.
1995;31(3):195–9.
51. Cawthorn RJ, Wobeser GA. Exogenous and endogenous stages of Isospora
brachyrhynchi sp. n.(Protozoa: Eimeriidae) from the American crow Corvus
brachyrhynchos Brehm. Can J Zool. 1985;63(11):2639–45.
52. Conway DP, McKenzie ME. Poultry Coccidiosis. Diagonistic Testing and
Procedures. Iowa: Blackwell Scientific Publications; 2007.
53. Hõrak P, Saks L, Karu U, Ots I, Surai PF, McGraw KJ. How coccidian parasites
affect health and appearance of greenfinches. J Anim Ecol. 2004;73(5):935–47.
54. Stenkewitz U, Nielsen ÓK, Skírnisson K, Stefánsson G. Host-parasite
interactions and population dynamics of rock ptarmigan. PLoS One.
2016;11(11):1–19.
55. Allen PC. Physiological responses of chicken gut tissue to coccidial infection:
Comparative effects of Eimeria acervulina and Eimeria mitis on mucosal mass,
carotenoid content, and brush border enzyme activity. Poult Sci.
1987;66(8):1306–15.
56. Allen PC, Fetterer RH. Recent Advances in Biology and Immunobiology of.
Cytokine. 2002;15(1):58–65.
57. Allen PC, Lydon J, Danforth HD. Effects of components of Artemisia annua on
coccidia infections in chickens. Poult Sci. 1997;76(8):1156–63.
58. Brawner WR, Hill GE, Sundermann CA. Effects of coccidial and mycoplasmal
infections on carotenoid-based plumage pigmentation in male house finches.
Auk. 2000;117(4):952–63.
59. McGraw KJ, Hill GE. Differential effects of endoparasitism on the expression of
carotenoid- and melanin-based ornamental coloration. Proc R Soc B Biol Sci.

2000;267(1452):1525–31.

60.   Halajian A, Eslami A, Mobedi I, Amin O, Mariaux J, Mansoori J, et al.
Gastrointestinal Helminths of Magpies (Pica pica), Rooks (Corvus frugilegus)
and Carrion Crows (Corvus corone) in Mazandaran Province, North of Iran. Iran
J Parasitol. 2011;6(2):38–44.

61.   Loman J. Reproduction in a Population of the Hooded Crow Corvus cornix.
Holarct Ecol. 1980;3:26–35.

62.   Nevarez JG, Gamble KC, Aczm D, Tully TN, Avian DA. Syngamus trachea
infection in two red-and-yellow barbets (Trachyphonus erythrocephalus). J Avian
Med Surg. 2002;16:31–3.

63.   Fernando AMA, Stockdale PHG, Remmler O. The route of migration,
development, and pathogenesis of syngamus trachea (Montagu, 1811) Chapin
1925, in pheasants. J Parasitol. 1971;57:107–16.

64.   Schnieder T, Boch J, Supperer R. Veterinärmedizinische Parasitologie. 6th ed.
Schnieder T, Boch J, Supperer R, editors. Berlin: Parey; 2006.

65.   Ortlepp RJ. The life-history of Syngamus trachealis (Montagu) v. Siebold, the
gape-worm of chickens. J Helminthol. 1923;1(3):119–40.

66.   Schat KA, Skinner MA. Avian immunosuppressive diseases and immune
evasion. In: Davison F, Kaspers B, Schat KA, editors. Avian Immunology.
Oxford, UK: Elsevier Academic Press; 2008. p. 299–322.

67.   Goymann W, Wingfield JC. Allostatic load, social status and stress hormones:
the costs of social status matter. Anim Behav. 2004;67:591–602.

68.   Seeman TE. Social ties and health: The benefits of social integration. Ann
Epidemiol. 1996;6(5):442–51.

69.   Cacioppo JT, Hawkley LC. Perceived social isolation and cognition. Trends
Cogn Sci. 2009;13(10):447–54.

70.   Shultz S, Dunbar R. Bondedness and sociality. Behaviour. 2010;147(7):775–803.

71.   Bates LA, Lee PC, Njiraini N, Poole JH, Sayialel K, Sayialel S, et al. Do
elephants show empathy? J Conscious Stud. 2008;15(10):204–25.

72.   Holekamp KE, Sakai ST, Lundrigan BL. Social intelligence in the spotted hyena
(Crocuta crocuta). Philos Trans R Soc Lond B. 2007;362(1480):523–38.

73.   Connor RC, Krützen M. Male dolphin alliances in Shark Bay: Changing
perspectives in a 30-year study. Anim Behav. 2015;103:223–35.

74.   Braun A, Bugnyar T. Social bonds and rank acquisition in raven nonbreeder
aggregations. Anim Behav. 2012;84(6):1507–15.

75.   Baglione V, Marcos JM, Canestrari D. Cooperatively breeding groups of carrion
crow (Corvus corone corone) in Northern Spain. Auk. 2002;119(3):790–9.

76.   Baglione V, Marcos M, Canestrari D, Ekman JAN. Direct fitness benefits of
group living in a complex cooperative society of carrion crows , Corvus corone
corone. Anim Behav. 2002;64:887–93.

77.   Canestrari D, Marcos J, Baglione V. Effect of parentage and relatedness on the
individual contribution to cooperative chick care in carrion crows Corvus corone
corone. Behav Ecol Sociobiol. 2005;116(5):422–8.

78.   Ekman J, Baglione V, Egger S, Griesser M. Delayed dispersal: living under the
reign of nepotistic parents. Auk. 2001;118(1):1–10.

79.   Ekman J, Dickinson JL, Hatchwell BJ, Griesser M. Delayed dispersal. In:
Ecology and Evolution of Cooperative Breeding in Birds. Cambridge, UK:
Cambridge University Press; 2004. p. 35–47.

80.   Hatchwell BJB, Komdeur J. Ecological constraints, life history traits and the
evolution of cooperative breeding. Anim Behav. 2000;59(6):1079–86.

81.   Chiarati E, Canestrari D, Vera R, Baglione V. Subordinates benefit from
exploratory dominants: response to novel food in cooperatively breeding carrion
crows. Anim Behav. 2012;83(1):103–9.
82.   Baglione V, Canestrari D. Carrion crows: Family living and helping in a flexible
social system. In: Koenig WD, Dickinson JL, editors. Cooperative Breeding in
Vertebrates: Studies of Ecology, Evolution, and Behavior. Cambridge, UK:
Cambridge University Press; 2016. p. 97–114.
83.   Clutton-Brock T. Breeding Together: Kin Selection and Mutualism in
Cooperative Vertebrates. Science. 2002;296(5565):69–72.
84.   Duboscq J, Romano V, Sueur C, MacIntosh A. Pick your lousy friends wisely:
Network centrality and seasonality interact to predict lice load in Japanese. Sci
Rep; 2015; 22095.
85.   De Vries H, Stevens JMG, Vervaecke H. Measuring and testing the steepness of
dominance hierarchies. Anim Behav. 2006;71(3):585–92.
86.   Carta LK, Carta DG. Nematode specific gravity profiles and applications to
flotation extraction and taxonomy. Nematology. 2000;2(2):201–10.
87.   Team RC. R: a language and environment for statistical computing. [Internet].
Vienna, Austria: R foundation for statistical computing; 2016. Available from:
http://www.r-project.org/
88.   Skaug H, Fournier D, Nielsen A, Magnusson A, Bolker B. Generalized Linear
Mixed Models using AD Model Builder. 2013.
89.   Hurvich C, Tsai C, Chih-Ling T. Regression and time series model selection in
small samples. Biometrika. 1989;76:297–307.
90.   Burnham KP, Anderson DR. Model selection and multimodel inference: a
practical information-theoretic approach. New York: Springer; 2002.
91.   Burnham KP. Multimodel Inference: Understanding AIC and BIC in Model
Selection. Sociol Methods Res. 2004;33(2):261–304.
92.   Engqvist L. The mistreatment of covariate interaction terms in linear model
analyses of behavioural and evolutionary ecology studies. Anim Behav.
2005;70(4):967–71.
590
591

 Table 1. Model selection of analyses that examined the occurrence of coccidian
oocysts. The occurrence of oocysts was related to variables associated with two
hypotheses: the 'group size' hypothesis and the 'social bonds' hypothesis: maximum
CSI value (max CSI), number of social bonds (nr bonds), number of related individuals
(nr related), normalized David's score (ΔnDS). Individual identity was fitted as a
random term in all models. LogLik: log-likelihood, df: degrees of freedom, ΔAICc:
difference in AICc to the best model. Models are ranked according to their Akaike
weight (Weight). N = 216 samples of 17 males and 200 samples of 16 females.

	Models	df	LogLik	AICc	ΔAICc	Weight
(a) Coccidia	Max CSI, nr bonds, nr related, sex	6	-209.815	431.835	0.00	0.115
	Max CSI, nr related, sex	5	-210.905	431.956	0.12	0.108
	Max CSI, nr bonds, nr related	5	-211.286	432.718	0.882	0.074
	Max CSI, nr related	4	-212.456	433.009	1.173	0.064
	Group size, max CSI, nr bonds, nr related, sex	7	-209.39	433.054	1.219	0.062
	ΔnDS, max CSI, nr bonds, nr related, sex	7	-209.544	433.362	1.527	0.053
	ΔnDS, max CSI, nr related, sex	6	-210.692	433.589	1.754	0.047
(b) Nematodes	ΔnDS, max CSI	4	-131.953	272.003	0	0.067
	Null model	2	-134.043	272.115	0.111	0.063
	ΔnDS	3	-133.048	272.154	0.15	0.062
	Max CSI	3	-133.054	272.166	0.162	0.061
	Max CSI, nr related	4	-133.714	273.525	1.522	0.031
	Nr related	3	-133.742	273.542	1.538	0.031
	Group size	3	-133.841	273.74	1.736	0.028
	ΔnDS, group size, max CSI	5	-131.804	273.754	1.751	0.027
	ΔnDS, group size	4	-132.84	273.777	1.774	0.027
	ΔnDS, max CSI, nr related	5	-131.84	273.826	1.823	0.026
	Group size, max CSI	4	132.909	273.915	1.191	0.025
	ΔnDS, max CSI, sex	5	-131.9	273.946	1.943	0.025

ΔnDS, sex	4	-132.932	273.961	1.958	0.252
ΔnDS, max CSI, nr bonds	5	-131.917	273.98	1.977	0.024
ΔnDS, nr related	4	-132.944	273.985	1.982	0.024

600
601
602

603 **Table 2.** Results of model averaged generalized mixed linear model investigating
604 factors affecting patterns of coccidian oocyst excretion. Significant values ($p \leq 0.05$) are
605 highlighted in bold.
606

Parameters	Estimate ± SE	z	p	Relative importance
Max CSI	**-1.25 ± 0.557**	**2.237**	**0.025**	**0.82**
Nr related	**-1.229 ± 0.467**	**2.623**	**0.008**	**0.93**
Sex	-0.811 ± 0.461	1.753	0.079	0.59
Nr bonds	0.723 ± 0.512	1.412	0.157	0.51
Group size	-0.384 ± 0.5	0.765	0.443	0.34
ΔnDS	0.326 ± 0.463	0.702	0.482	0.32

Figure 1.
Occurrence of coccidan oocysts in carrion crow droppings in relation to the maximum
CSI. Occurrence probabilities were calculated as the predicted values (solid line) and
95% CI (shaded area).

Figure 2.
Occurrence of coccidan oocysts in carrion crow droppings in relation to the number of
related individuals in the group. Occurrence probabilities were calculated as the
predicted values (solid line) and 95% CI (shaded area).

617