Title page

Quartz-Seq2: a high-throughput single-cell RNA-sequencing method that effectively uses

limited sequence reads

Yohei Sasagawa[1†], Hiroki Danno[1†], Hitomi Takada[2†], Masashi Ebisawa[1], Tetsutaro Hayashi[1],

Akira Kurisaki[2*], and Itoshi Nikaido[1*]

[†]Contributed equally

[*]Corresponding author

[1]Bioinformatics Research Unit, Advanced Center for Computing and Communication, RIKEN,

Hirosawa 2-1, Wako, Saitama, Japan

[2]Laboratory of Stem Cell Technology, Graduate School of Biological Sciences, Nara Institute of

Science and Technology, Takayama-cho 8916-5, Ikoma, Nara, Japan

[†]Contributed equally

[*]Corresponding author

Yohei Sasagawa: youhei.sasagawa@riken.jp

Hiroki Danno: hiroki.danno@riken.jp, redgrapefruit@mac.com

Hitomi Takada: htakada@bs.naist.jp

Masashi Ebisawa: masashi.ebisawa@riken.jp

Tetsutaro Hayashi: tetsutaro.hayashi@riken.jp

Akira Kurisaki: akikuri@bs.naist.jp

Itoshi Nikaido: itoshi.nikaido@riken.jp

Abstract

High-throughput single-cell RNA-seq methods assign limited unique molecular identifier (UMI) counts as gene expression values to single cells from shallow sequence reads and detect limited gene counts. We thus developed a high-throughput single-cell RNA-seq method, Quartz-Seq2, to overcome these issues. Our improvements in several of the reaction steps of Quartz-Seq2 allow us to effectively convert initial reads to UMI counts (at a rate of 30%–50%). To demonstrate the power of Quartz-Seq2, we analyzed transcriptomes from a cell population of *in vitro* embryonic stem cells and an *in vivo* stromal vascular fraction with a limited number of sequence reads.

Keywords

High-throughput single-cell RNA-seq

Flow-cytometry

Cell sorter

Quartz-Seq

Poly-A tagging

Stromal vascular fraction

Mesenchymal stem cell

Background

Single-cell transcriptome analysis allows us to identify nongenetic cellular heterogeneity, which includes differences in cell type due to differentiation and differences in cell state within a cell population. In previous studies, various methods for single-cell RNA-seq were developed [1-19]. Some of these that generate read coverage across all transcripts have been exploited to detect alternative transcription splicing isoforms [14,17] and others using unique molecular identifier (UMI) have been applied to quantify the number of transcripts expressed in a cell [1-3,6,7,9,11-13,20]. To extract substantial information on a cell population, such as the composition of different cell types or the distribution of cell states, it is necessary to analyze hundreds or thousands of cells. Cell barcoding is a key technology for this, which enables us to deal with samples from numerous cells in a single tube. Cell barcoding technology, which tags nucleotides unique to each cell to target RNA molecules from that cell, is a key technology for increasing the throughput of single-cell RNA-seq [16,18]. Mixing cDNA tagged with cell barcodes before whole-transcript amplification decreases the cost of reaction reagents and the laboriousness of experimental steps. There are two types of cell barcoding technology according to the method of cell sampling used. One method involves single cells being selectively sorted to multi-well plates using flow cytometry, which allows us to remove dead or aggregated cells. Besides, transcriptome data can be linked to cellular information measured in flow cytometry. The other method involves single cells and barcoded beads being captured in water-in-oil droplets using droplet-generation microfluidic devices [6,7]. In this latter method, thousands of cells can probabilistically be captured in half an hour. However, the total number of sequence reads generated by a deep sequencer is still limited. To increase the number of analyzed cells, each cell is assigned a limited number of initial sequence reads. For example, approximately 400 million initial reads were sequenced for 3,000–4,000 cells in several previous studies [6,7]. UMI counts were converted from initial sequence reads for each cell, and the conversion ratio of that was limited to approximately one-tenth [21]. Ideally, greater UMI counts should be generated from limited sequence reads because the increase in UMI count assigned to each cell leads to the detection of low-copy genes and the identification of cell-type-specific genes using statistical tests.

In this study, we developed a novel high-throughput single-cell RNA-seq method,

Quartz-Seq2. As Quartz-Seq is a sensitive and reproducible single-cell RNA-seq method,

Quartz-Seq2 was developed based on it [15]. Quartz-Seq is based on a poly-A tagging strategy.

By the combination of molecular biological improvements including major improvement of

poly-A tagging, Quartz-Seq2 resulted in an increase in the effectiveness with which the initial

sequence reads were converted to the expression UMI counts (UMI conversion efficiency:

30%–50%). To demonstrate the highly effective use of initial reads in Quartz-Seq2, we

analyzed a population of mouse embryonic stem (ES) cells as *in vitro* cells and the cell

population from the stromal vascular fraction (SVF) as *in vivo* cells.

Results

Outline of Quartz-Seq2 experiment

To increase the UMI conversion efficiency, we improved several steps in the preparation of the single-cell RNA-seq library (Additional file 1: Figure S1), resulting in the development of Quartz-Seq2. Below, we explain the five steps of the Quartz-Seq2 procedure (Figure 1).

(1) The first step is single-cell collection using a cell sorter. We selectively sort living single cells into lysis buffer in a 384-well PCR plate without dead cells. In cell sorting, various types of channel information, such as the intensity of fluorescence, are measured for each cell. This enables us to link the transcriptome to cellular information from a cell sorter for each cell.

(2) The second step is cell barcoding. Each well contains lysis buffer and reverse-transcription (RT) primer, which includes a cell barcode sequence (14- or 15-mer), a UMI sequence (8-mer), and an oligo-dT sequence (24-mer). Using these RT primers, respective RNA from single cells is converted to cDNA with unique cell barcodes. Note that a long RT primer resulted in a severe problem regarding the synthesis of byproducts at the downstream reaction in our system (Additional file 1: Figure S2 and Supplemental Note). Therefore, we use a relatively short RT primer (73- or 74-mer), which allows us to skip the step of removing byproducts using exonuclease I. We design two types of RT primer set (v3.1: 384 barcodes; v3.2: 1536 barcodes). Within each primer set, barcode sequences are designed such that the minimum Sequence–Levenshtein distance between two sequences should be greater than 5, which leads to the correction of two nucleotides of mismatch, insertion and deletion in sequence reads [22]. We also optimize the buffer and temperature in the RT reaction, leading to an improvement of RT efficiency from that of the original Quartz-Seq (Additional file 1: Figure S3 and Supplemental Note).

(3) The third step involves the pooling of cell-barcoded cDNA. By cell barcoding, Quartz-Seq2 can pool cDNA of up to 1,536 individual cells into one mixture. We developed a rapid and high-throughput method for collecting small volumes of cDNA in multiwell plates. This method also achieved higher efficiency of collection than dispensing with pipettes and tips (Additional file 1: Figure S4 and Supplemental Note). As the efficiency of cDNA purification

after pooling was 93.77%, we estimated that approximately 80% of cell-barcoded cDNA could

be used for subsequent whole-transcript amplification in our system (Additional file 1: Figure

S3a and Supplemental Note).

(4) The fourth step is whole-transcript amplification based on an improved poly-A tagging

strategy. Poly-A tagging is one of the methods of converting first-strand cDNA to amplifiable

cDNA. First-strand cDNA is extended with a poly-A tail by the terminal transferase.

Subsequently, second-strand cDNA is synthesized with tagging primers that contain a poly-dT

sequence, followed by PCR amplification. Here, we improve the efficiency of poly-A tagging by

3.6-fold. This improvement is a crucial point in the development of Quartz-Seq2. We describe

the details of this in the next subsection.

(5) The fifth step is library preparation for deep sequencing. Amplified cDNA is

fragmented, ligated with the sequence adapter, and amplified by PCR. In sequencing using

Illumina sequencers, a sequence for a cell barcode and a UMI is read in Read1, while a

sequence for a region of a transcript (mRNA) is read in Read2.

Improvement of poly-A tagging efficiency

We previously reported Quartz-Seq based on the poly-A tagging strategy, which has a

significant potential for detecting a large number of genes expressed in a cell [15]. However,

the efficiency of poly-A tagging itself for single-cell RNA-seq has not been improved. We

hypothesized that the improvement of poly-A tagging would lead to high UMI conversion

efficiency. Therefore, we attempted to improve the efficiency of this tagging step. Poly-A

tagging is composed of two processes: (1) The first-strand cDNA is modified with a poly-A tail

by terminal deoxynucleotidyl transferase. (2) Next, the poly-A-tailed cDNA is annealed with a

tagging primer, which has a poly-dT sequence. Then, the second-strand cDNA is extended.

The resulted second-strand cDNA is amplifiable cDNA, which has a PCR primer sequence at

both ends of it.

It is known that the DNA yield of amplified cDNA generally reflects the quantitative

performance of single-cell RNA-seq methods [3,13,14]. Thus, we determined the effects of

various buffers for the poly-A tailing step on the amplified cDNA yield. We performed the poly-A

tailing reaction with various buffers using purified first-strand cDNA from 1 ng of total RNA

(Figure 2a, Additional file 1: Figure S5). Finally, we obtained amplified cDNA. We found that the

use of T55 buffer and RH55 buffer in the poly-A tailing reaction efficiently improved the cDNA

yield (Figure 2a). The amount of amplified cDNA increased 2.88-fold and 2.55-fold using T55

buffer and RH55 buffer, respectively, compared with the level using Quartz-Seq buffer (Figure

2a). In these buffer conditions, we did not observe any obvious byproducts derived from the RT

primer (Figure 2a, Additional file 1: Figures S2 and S5, and Supplemental Note).

Next, we added an "Increment" temperature condition for the tagging and second-strand

synthesis steps (see Material and Methods). In this condition, the reaction temperature of these

steps was gradually increased. As a result, the amount of cDNA tended to increase, by

approximately 1.2-fold (Figure 2a). Moreover, upon combining T55 buffer and the Increment

condition, the amount of cDNA increased approximately 3.6-fold. We also confirmed the

reproducibility of this phenomenon of cDNA increment in the additional experiments (Additional

file 1: Figure S5). Moreover, we confirmed the amplified cDNA yield of various genes by qPCR

analysis as another assay. Specifically, we determined the qPCR scores of eight genes from

amplified cDNA and nonamplified cDNA (Figure 2b). Spearman's rank correlation coefficients

(SCCs) between amplification and nonamplification were approximately 0.79 in both conditions

(T55+Increment, RH55+Increment). The SCC was approximately 0.66 in Quartz-Seq-like

conditions. We also observed clear increments of qPCR scores for almost all genes. These

results show that the combination of T55 buffer and this temperature condition improved the

efficiency of the poly-A tagging step. We also found that other conditions (NBF40+Increment)

improved the cDNA yield. However, under these conditions, byproducts were clearly

synthesized (Additional file 1: Figures S4c and S5b). Moreover, the cDNA amount with T55

buffer was slightly greater than the cDNA amount with RH55 (Figure 2 and Additional file 1:

Figure S5a). Therefore, we used the combination of T55 buffer and the "Increment"

temperature condition for the poly-A tagging strategy.

Evaluation of the quantitative performance of Quartz-Seq2 using single-cell total RNA

Finally, we adopted three molecular-biological improvements [modified RT condition, poly-A

tailing buffers (T55), and the "Increment" temperature condition] for Quartz-Seq2 (Additional file

1: Figure S1). To determine the technical variability and specificity of Quartz-Seq2, we

performed whole-transcript amplification using 10 pg of diluted mouse total RNA as a

single-cell-like averaged sample with the v3.1 384 cell-barcode RT primer in a 384-well plate.

In this experiment, we used the "RT100" enzyme concentration for reverse transcription. The

effect of enzyme concentration is specifically described in the next subsection of this paper. We

analyzed 10 pg of total RNA in all wells at approximately 0.19 M fastq reads on average per

well. In the case of Quartz-Seq-like conditions, we detected 19,614±3516 UMI counts and

5726±494 gene counts (n=384 wells). In the case of Quartz-Seq2, we achieved a UMI count of

35,391±8719 and a gene count of 6621±711 (n=384 wells). We confirmed the reproducibility of

the quantitative performance by a further batch experiment with other 384-well plates, which

had 10 pg of total RNA in all wells (Additional file 1: Figure S6). These results indicated that the

combination of the three molecular-biological improvements for Quartz-Seq2 clearly improved

the conversion ratio from target RNA to sequence library DNA. We also confirmed the

correlation between single-cell RNA-seq data and bulk RNA-seq data. SCC between

Quartz-Seq2 and bulk RNA-seq was 0.85 (Figure 2d). This result indicates that Quartz-Seq2

data are comparable to conventional bulk RNA-seq data. We also determined the variability of

expression between technical replicates to investigate the UMI filtering effect. The mean-CV

(coefficient of variation) plot of Quartz-Seq2 clearly showed that UMI filtering reduced

expression variability derived from PCR bias (Figure 2e). Moreover, technical gene expression

variability of Quartz-Seq2 became fairly close to the theoretical variability of a Poisson

distribution (Figure 2e).

We also focused on the relationship between sequencing platform and Quartz-Seq2.

We did not observe clear differences between the NextSeq500 and HiSeq2500 sequencing

platforms for determination of the UMI count and the gene count for Quartz-Seq2 (Figure S7a).

On the other hand, the unique mapping ratio and UMI count clearly depended on the Read2

length for transcript mapping (Figure S7c, d). In this regard, the number of detected genes

slightly depended on Read2 length (Figure 7d). Long Read2 length led to better UMI counts,

but high sequencing cost (Figure 7b). Therefore, in this study, we mainly used the NextSeq500

sequence platform and a Read2 length of 62 due to the cost performance of this combination.

Moreover, we mainly used the adaptor ligation method for sequence library preparation of

Quartz-Seq2. We did not observe a clear difference between the two sequence library

preparation methods (ligation and Nextera/Tn5 transposase) regarding determination of the

UMI count and the gene count. However, we could easily identify specific familiar genes (such

as *Pou5f1* and *Gapdh*) using the ligation method (Figure S7e). Therefore, we mainly used the

ligation method for Quartz-Seq2.

Reduction of enzyme concentration in reverse transcription decreased cost and

technical noise of Quartz-Seq2

The cost of experimental preparation per cell was approximately ¥2600 ($23) for our previously

reported Quartz-Seq, which does not use cell barcoding (Figure S8c). CEL-seq2 is one of the

PCR plate-based single-cell RNA-seq methods using a cell sorter. This method has the

advantage of reducing costs via cell barcoding. Indeed, the experimental cost of CEL-seq2 is

lower (approximately ¥372 or $3.35) than the cost of Quartz-Seq. On the other hand, the

experimental cost per cell is extremely low (approximately ¥10 or $0.08) for droplet-based

high-throughput single-cell RNA-seq methods, such as Drop-seq ([6] and Figure S8c). Ideally,

the experimental cost for high-throughput single-cell RNA-seq methods using a cell sorter

should be reduced to approach the cost of droplet-based methods. As mentioned earlier, we

used the "RT100" enzyme condition in reverse transcription for Quartz-Seq2. This enzyme

concentration in RT reaction is broadly used for various molecular biological application

including single-cell RNA-seq methods. By using the cell barcoding strategy, the cost of

experimental preparation for Quartz-Seq2 under the "RT100" condition was reduced (¥121 or

$1.09 per cell). We found that 65% of the cost in experimental steps is derived from reverse

transcription in Quartz-Seq2 under the "RT100" condition (Figure 3a). Therefore, we

investigated the effect of a low enzyme concentration in reverse transcription in Quartz-Seq2.

In the assessment assay, we noticed that a low enzyme concentration did not markedly affect

the efficiency of reverse transcription and the amount of amplified cDNA yield (Figure S8a right

panel and S8b). Therefore, we prepared three technical replicates of a 384-well PCR plate with

10 pg of total RNA with the "RT100" condition or the "RT25" low-enzyme condition. In the

"RT25" condition, the amount of cDNA yield showed a tendency for a slight increase of

approximately 1.17-fold, which although not being a major improvement did at least not involve

a decrease. The cost of experimental preparation per cell became approximately ¥45–¥62

($0.40–$0.55) in the "RT25" condition.

We validated the quantitative performance with 10 pg of total RNA in the "RT100" and

"RT25" conditions at various input data sizes. Unexpectedly, we found that the "RT25"

condition improved the quantitative performance (Figure 3c and Additional file 1: Figure S9). In

a typical pattern, approximately 4,000 single cells were analyzed by the NextSeq500

sequencer in the high-throughput single-cell RNA-seq method [6,7,21]. In this case, the input

data size for a single cell involves shallow initial reads (0.1 M fastq reads) (Figure 3d). We

compared RT25 with RT100 at approximately 0.096 M fastq reads on average per well. In the

case of Quartz-Seq2 in the "RT25" condition, we achieved a UMI count of 30,117±789 and a

gene count of 6,320±35 (n=3 plates). In the case of Quartz-Seq2 in the "RT100" condition, we

achieved a UMI count of 25,671±1020 and a gene count of 5,889±35 (n=3 plates). The capture

ratios of ERCC spike were 11.49±0.46% and 10.28±0.31% in the "RT25" and "RT100"

conditions, respectively (Figure 3d and e). In addition, the capture ratio of ERCC spike was

16.38±0.66% in the "RT25" condition at approximately 0.27 M fastq reads on average per well

(Figure 3c, e). We also investigated the technical variability of UMI count and gene count. The

well-to-well technical variability for UMI count and gene count clearly decreased in the "RT25"

condition (Figure 3). These results showed that the "RT25" low-enzyme condition clearly

reduced the experimental cost and improved the quantitative performance, including that for

the UMI count. We thus applied the "RT25" condition in subsequent experiments using real

single cells.

Quartz-Seq2 shows high efficiency of UMI conversion

The increase of UMI counts generated from limited initial sequence reads leads to the

detection of low-copy genes and the identification of more marker genes using statistical tests.

We calculated the UMI conversion efficiency, which indicates how effectively initial reads can

be converted to UMI counts (Figure 4a and Additional file 1: Figure S10). The UMI conversion

efficiency of Quartz-Seq2 with 10 pg of purified total RNA (from mouse ES cells) ranged from

31% to 39% at approximately 0.096 M initial reads depending on the Read2 length. These

results show that major improvements (such as poly-A tagging) in Quartz-Seq2 increase the

efficiency of UMI conversion.

We performed Quart-Seq2 and Drop-seq on a mixture of mouse ES cells and the

ES-derived primitive endoderm (PrE) cells, and calculated the UMI conversion efficiency

(Figure 4b and Additional file 1: Figure S11a). Drop-seq is one of the high-throughput

single-cell RNA-seq methods, which can capture thousands of cells and barcoded beads in half

an hour. However, there are many barcoded beads that are not co-captured with a cell in a

droplet, leading that 10%–40% of initial sequence reads could be lost [6]. We performed

Drop-seq experiments, which were validated by species-mixing analysis (Figure S11b, c). The

effectiveness of Quartz-Seq2 ranged from 25% to 35% depending on the initial read depth

(Figure 4), which was higher than that of Drop-seq. In previous studies, average UMI counts

and gene counts were compared among several single-cell RNA-seq methods [1,6,21]. Using

such data, we calculated the UMI conversion efficiency of each method and compared them

with that of Quartz-Seq2 (Additional file 1: Figure S12). UMI conversion efficiency depended on

the length of Read2 that corresponds to the transcript sequence because the rate of unique

mapping depends on it (Additional file 1: Figure S7). As Figure S12 shows, the UMI conversion

efficiency of Quartz-Seq2 was higher than those of the other single-cell RNA-seq methods.

As Figure 4b shows, Quartz-Seq2 detected more genes with high UMI conversion

efficiency. To examine the power for identifying marker genes, we performed principal

component analysis (PCA) and clustering. The number of genes differentially expressed

between two distinct cell types was calculated (ES and PrE cells) (Additional file 1: Figure

S11d). The results showed that more genes that were differentially expressed between the two

cell types were identified with Quartz-Seq2 (Figure 4b). In addition, more biological pathways

particularly associated with the differentially expressed genes were detected in Quartz-Seq2

(Figure 4c). Furthermore, we calculated genes with highly variable expression in Quartz-Seq2

and Drop-seq, which potentially include not only genes that are differentially expressed

between cell types but also genes of which the expression changes depending on the cell state

in a cell type. The results also showed that more Gene Ontology (GO) terms enriched among

the genes with highly variable expression were identified with Quartz-Seq2. Terms on cell cycle

state were only enriched in the genes calculated with Quartz-Seq2 (Figure 4d). These results

suggest that high UMI conversion efficiency with limited initial reads leads to more biological

information being revealed, such as functional terms and biological pathways.

Quartz-Seq2 analysis on 4,484 mouse embryonic stem cells in maintenance and

differentiated conditions

We did not observe any clear differences in UMI counts and gene counts between the v3.1 and

v3.2 RT primer sets (Additional file 2: Table S1). To demonstrate the capability of Quartz-Seq2

for quantifying the transcriptome of a large number of cells and identifying rare cell populations,

we analyzed approximately 4,484 cells from a mixture of mouse ES cells and

differentiation-induced cells. Cells were prestained with Hoechst 33342 and/or Calcein-AM as

an indicator of DNA content and culture condition (ES or PrE cells), respectively (Figure 5a).

Calcein-AM-positive and -negative cells were sorted to 12 of 384-well plates in a checkered

pattern (Figure 5a). In this condition, the cell doublets caused by mis-sorting can be detected at

the highest probability. In this analysis, as an average for cells, 0.1 M initial fastq reads were

effectively converted to 35,915±1,188 UMI counts in a final gene expression matrix. The UMI

conversion efficiency was 35.91±1.18%.

Dimensionality reduction of the expression UMI matrix resulting from Quartz-Seq2

showed clear separation of six clusters, including the main populations of ES cells and PrE

cells and four small populations. We did not observe a clear batch effect between 12 of the

384-well plates (Additional file 1: Figure S13). One cluster contained a relatively high ratio of

mitochondrial RNA (Additional file 1: Figure S14), which was judged to reflect low-quality cells

[23]. Another cluster showed high values of detected UMI counts and gene counts, and

expressed both ES and PrE marker genes, which were judged to represent doublets of an ES

cell and a PrE cell. To characterize the identified populations, we determined the genes that

were specific or enriched for each cluster using binomial tests (Figure 5c and Additional file 1:

Figure S15). Although cells in cluster 3 shared gene expression with the main population of ES

cells (cluster 1), some genes including *Stmn2* and *Rhox5* were additionally expressed,

suggesting that cluster 1 and cluster 3 shared the same cell type but had different cell states.

Cluster 5 was a small population (0.17%, Figure 5b), but was characterized by many specific

marker genes (Figure 5b), including *Zscan4c* and *Zscan4d*, which are known to be expressed

in a subpopulation of mouse ES cells [24]. However, among our culture conditions, these

genes were more expressed in the Dex-treatment condition than in the ES-maintenance

condition. Flow cytometry information on cells in cluster 5 also showed no fluorescence of

Calcein-AM, indicating that these cells were from the suspension of Dex-treated cells (Figure

5d). These observations were consistent with a previous study demonstrating that *Zscan4* was

more expressed in a differentiation condition induced by withdrawal of leukemia inhibitory

factor [7].

390   In sampling of cells using a cell sorter, flow cytometry information including the intensity

of Hoechst 33342 staining for each cell was collected. Using this information, we reconstructed

the distribution of Hoechst 33342 staining intensity for each cluster (Figure 5e). The main

population of ES cells (cluster 1) showed an embryonic pattern of distribution of DNA content,

which is characterized by a high ratio of cells in G2 and M phases to cells in the G1 phase. On

the other hand, the main population of PrE cells (cluster 4) showed a somatic pattern of

distribution of DNA content. This was consistent with PrE cells being differentiated from ES

cells. The cells in cluster 3, the gene expression of which was similar to that of the ES main

population (cluster 1), showed an embryonic pattern rather than a somatic one. As the number

of cells in cluster 5 that expressed *Zscan4c/d* was small, it was difficult to classify the observed

pattern as the embryonic or somatic type. These findings indicated the usefulness of single-cell

RNA-seq using flow cytometry for the reconstruction of population information after

transcriptome-based clustering. For good interpretation of the distribution, a large number of

cells for each cluster are required.

Quartz-Seq2 revealed two types of cell state: cell cycle-dependent and -independent

heterogeneity

When cells were plotted on t-SNE (t-distributed stochastic neighbor embedding) space using

transcriptome analysis and the intensity of Hoechst 33342 staining was colored, the gradient

pattern was easily observed for clusters 1, 3, and 4 (Figure 6a), suggesting that Quartz-Seq2

was highly sensitive for detection of the cell cycle state. To examine this, we plotted the gene

expression of several cell cycle markers against the intensity of Hoechst 33342 staining, which

indicated the strong dependence of such expression on DNA content (Figure 6b). Using the

intensity of Hoechst 33342 staining, cell cycle dependence was calculated for all genes

detected in Quartz-Seq2 experiments, resulting in the identification of numerous genes for

which the expression level changed depending on the DNA content (Figure 6c, d and

Additional file 1: Figure S16). We called these genes cell cycle-associated genes. Again, it was

confirmed that the ratio of cells in G1 and S phases to those in G2 and M phases was different

between ES and PrE cells (Figure 6c). As we showed that the efficiency of UMI conversion was

high in the Quartz-Seq2 method, we examined whether 10,000 initial reads are sufficient for

cell cycle analysis (Additional file 1: Figure S17). In this analysis, final UMI count was 4,774±62

and the UMI conversion efficiency was 47.74±0.66%. Plotting cells on t-SNE space with

coloring the intensity of Hoechst 33342 staining revealed a gradient pattern, which was

positively correlated with G2/M marker genes including *Ccnb1* and *Top2a* and was negatively

correlated with G1-phase marker genes such as *Ccne1* (Additional file 1: Figure S17). These

observations demonstrated Quartz-Seq2 with few initial sequence reads could analyze cell

cycle due to the high UMI conversion efficiency.

Within a single cell type, there are two types of cell state. One is the different stages of

the cell cycle as mentioned above, and the other refers to cellular heterogeneity that is only

remotely related to the cell cycle phases. This means that genes with variable expression in a

cell type include cell cycle-associated genes and gene with variable expression that is

independent of the cell cycle phases (cell cycle-phase independent variable genes). We

determined the number of this latter group of genes by subtracting the cell cycle-associated

genes from the genes with variable expression (Figure 6e). For the main population of ES cells

(cluster 1), we identified tens of cell cycle-phase independent variable genes. These genes

included *Spp1*, *Rhox5*, and *S100a6*, which were previously identified as genes with highly

variable expression by scRNA-seq (Figure 6) [7,13,15]. We also identified *Stmn2*, *Dnmt3l*,

Tmsb4x, and several genes as novel cell cycle phase-independent variable genes. Single-cell

RT-qPCR also showed high variability for *Stmn2* and *Rhox5* compared with that for *Nanog*

(Figure 6f). In previous studies, *Sgk1* and *Actb* were identified as genes with variable

expression by scRNA-seq that analyzed cells only in G1 phase or cells without measuring the

intensity of Hoechst 33342 [7,15]. In this study, these were classified as cell cycle-associated

genes (Additional file 1: Figure S16).

These results show that Quartz-Seq2 globally classified genes with variable expression

within a cell type into cell cycle-associated genes and cell cycle phase-independent genes with

variable expression by using Hoechst 33342 staining intensity data obtained in flow cytometry.

The global picture of cell-type composition in stromal vascular fraction as revealed by

Quartz-Seq2

Next, Quartz-Seq2 was applied to cells from the mouse stromal vascular fraction (SVF), which

was taken from adipose tissue (Figure 7a). SVF is thought to be one of the most important

sources of mesenchymal stem cells (MSC) due to its potential for cell therapies and the

stimulation of endogenous repair [25-28]. However, the global picture of cell-type composition

in SVF has not been clarified. To do so, we first observed the distribution of cell size in the SVF

population and found it to be broad (5–13 μm, average 6.43 ± 1.35 μm; n=200) (Figure 7a).

This was confirmed by flow cytometry data that were obtained using a cell sorter. It is known

that the typical amount of total RNA in a mammalian cell is approximately 10 pg [15,17]. Thus,

the average amount of total RNA in the SVF population of approximately 1.7 pg is relatively

small.

We collected cells in multiple 384-well PCR plates and analyzed their transcriptome

using Quartz-Seq2. In this analysis, 38,450±3513 initial fastq reads were converted to UMI

counts in the final digital expression matrix. The UMI conversion efficiency was 26.85±2.70%

(n=3). We observed broader distributions of UMI counts and gene counts in SVF than in ES

and 10 pg of total RNA. The gene counts correlated well with side scatter (SSC) values, which

were associated with cell size (Additional file 1: Figure S19).

Dimensionality reduction and clustering of single-cell transcriptome data showed the

clear separation of 11 clusters (Figure 7b). To annotate the cell type on each cluster, we

calculated marker genes and functional terms that were either specific or enriched for each

cluster (Figure 7c, d and Additional file 1: Figures S18 and S20). As Figure 7c shows, clusters 1,

2, 5, and 8-11 expressed cluster-specific genes, and clusters 3, 4, 6, 7, and 9 shared

expressed genes (marker class C) as well as differentially expressed genes of classes D-G.

Taking these markers and functional terms as well as previous knowledge together, each

cluster was identified as follows: two types of *Cd34*-positive MSC (cluster 1: 7.8%, cluster 8:

6.4%), two types of *Cd4*-positive T cells (cluster 3: 45.0%, cluster 4: 1.0%), *Cxcr6*-positive T

cells (cluster 9: 1.71%), B cells (cluster 5: 15.9%), killer T cells (cluster 6: 11.0%), natural killer

cells (cluster 7: 3.0%), macrophages (cluster 2: 5.5%), dendritic cells (cluster 10: 0.86%), and

plasmacytoid dendritic cells (cluster 11: 0.57%). We noted that the cell sizes of the T cells and

B cells from clusters 3, 4 and 5, which were measured using a cell sorter, were smaller than

those of MSCs and macrophages from clusters 1, 2, and 8 (Additional file 1: Figure S19). We

confirmed that the cell sizes of CD4-positive T cells and CD79-positive B cells were smaller in

SVF via immunofluorescence observation (Additional file 1: Figure S19). These results show

that Quartz-Seq2 with flow-cytometry data is useful for defining respective cell clusters with cell

size information.

Subsequently, we investigated the gene expression of receptors and ligands in stromal

vascular fractions. The variety of receptors expressed in each cluster was similar (Additional

file 1: Figure S21). However, we found a wide variety of ligands expressed in clusters 1 and 8

(Additional file 1: Figure S21). It is known that MSCs secrete many paracrine mediators, which

have a therapeutic effect [29,30]. These results suggest that clusters 1 and 8 are MSCs, which

secretes many paracrine mediators.

We focused on two potential MSC clusters (1 and 8) because the heterogeneity of the

MSC population has been discussed in many papers [31,32]. By using Quartz-Seq2, clusters 1

and 8 had the *Cd31*-/*Cd34*+/*Cd45*- phenotype (Figure 7, Additional file 1: Figure S22). In a

previous study, qPCR-based single-cell transcript analysis of the *Cd31-/Cd34+/Cd45-*

population of SVF was reported. By analyzing 140 genes, the authors identified the

Cd55+/Dpp4+/Cd31-/Cd34+/Cd45- cell population and showed that administration of this

population was effective for normalizing diabetic wound healing in mice [31]. They identified 13

marker genes for the *Cd55+/Dpp4+/Cd31-/Cd34+/Cd45* population in mouse SVF. We

compared the expression of those 13 genes between clusters 1 and 8 (Additional file 1: Figure

S23). The results showed that almost all the genes were more highly expressed in cluster 1

than in cluster 8, suggesting that the cluster 1 population was similar to the

Cd55+/Dpp4+/Cd31-/Cd34+/Cd45- population. To further analyze the relationship between our

identified clusters and previously reported heterogeneity of the MSC population, we examined

the expression of typical MSC markers (*CD90*, *CD105*, *PDGFRa*, and *Sca-1*) [31,33] as well as

subpopulation specific markers (*Pou5f1*, *Nanog*, *Sox2*, *Tnnt2*, and *Myog*) [34,35].

PDGFRa (*Pdgfra*) and *Sca1* were expressed in both clusters 1 and 8, whereas *Cd90* was

expressed at a relatively higher level in cluster 8 than in cluster 1, and *Cd105* was not strongly

expressed in either cluster. Neither pluripotent markers (*Pou5f1*, *Nanog*, and *Sox2*) nor skeletal

and cardiomyogenic markers (*Tnnt2* and *Myog*) were detected in either cluster (Additional file

1: Figure S22). Collectively, our transcriptome analysis showed that the MSC population is

divided into two clusters, suggesting that there is less heterogeneity of MSC in the SVF than

expected.

Next, we investigated the difference between the two MSC clusters in SVF. We detected

182 differentially expressed genes between clusters 1 and 8 (Additional file 1: Figure S24,

Additional file 5: Table S4). Cluster 8 was characterized by the enriched expression of genes

that encode extracellular matrix proteins, including collagens, *Hspg2*, and *Bgn* (Additional file

1: Figure S22). By contrast, cluster 1 was characterized by the expression of *IGFBPs* (*Igfbp4*,

Igfbp5, *Igfbp6*), prostaglandin endoperoxidase synthases (*Ptgs2*), and secreted factors *Fgf2*

and Bmp2. Of note, the expression of stem cell characteristic genes *Aldh1a3*, *Fgf2*, *Bmp2*, and

Tgfbr2 were identified as important modules in cluster 1 (Additional file 1: Figure S22),

suggesting that cluster 1 had more stem-like phenotypes that were secreting with medicinal

factors.

To leverage the power of the entire transcriptome analysis, we performed an enrichment

analysis of biological pathways using the 182 differentially expressed genes between the two

clusters (Figure 7e, Additional file 6: Table S5). Cluster 1 was characterized by elastic fiber

pathways and IGF transport pathways. Elastic fibers contribute various connective tissues and

are related to vascular function [36]. In cluster 8, a tightly linked network of collagen

metabolism pathways was highly enriched. These results show that two MSC clusters of SVF

can have different functions in extracellular matrix formation.

Taken together, our transcriptome analysis by Quart-seq2 successfully identified cell

populations that consisted of various types of both large and small cells, which were isolated

from *in vivo* tissue, and demonstrated that fresh SVF contains two closely related types of

MSCs that have distinct characteristics.

Discussion

The rate of conversion from initial reads to UMI counts was low, especially in high-throughput single-cell RNA-seq. An increase in UMI counts assigned to each cell theoretically leads to the detection of low-copy genes and the identification of cell-type-specific genes using statistical tests. Therefore, in this study, we developed a novel high-throughput single-cell RNA-seq method, Quartz-Seq2. We calculated the UMI conversion efficiency, which indicated how effectively initial reads can be converted to UMI counts. The UMI conversion efficiency of Quartz-Seq2 was higher than those of other single-cell RNA-seq methods (Additional file 1: Figure S12). In other words, Quartz-Seq2 could detect more genes from limited fastq reads and for a lower cost. By cell barcoding, Quartz-Seq2 can pool cDNA of up to 1,536 individual cells into one mixture. Streamlined and simplified experimental processes of Quartz-Seq2 allowed us to handle thousands of single cells within a few days. Quartz-Seq2 also showed high accuracy and high reproducibility, leading to precise measurement of the transcriptome from single cells (Figure 2d and e). We demonstrated Quartz-Seq2 analyses on mouse ES cells as an i*n vitro* sample, and mouse SVF including MSC as an *in vivo* sample. Moreover, we decreased the cDNA preparation cost per cell of Quartz-Seq2 by 97.6%-98.2% compared with that of Quartz-Seq (Figure S8c). The total cost (cDNA preparation cost and sequence cost per cell) of Quartz-Seq2 as a plate-based method approached that of a droplet-based method (Additional file 1: Figure S8d). Furthermore, the use of evaporation-preventing oil might reduce the cDNA preparation cost of Quartz-Seq2 by 75% [2].

We proposed that UMI conversion efficiency could be a useful variable for evaluating performance for the further development of high-throughput single-cell RNA-seq methods with shallow reads. The increase of initial sequence reads by additional sequencing does not cancel out the low efficiency of UMI conversion at equal rate. For example, approximately 24,000 fastq reads were converted to approximately 11,000 UMI counts in Quartz-Seq2, while Drop-seq required approximately 169,000 fastq reads to generate the same UMI counts. At this condition, the difference of UMI conversion efficiency was 2.5-fold but the difference of required fastq reads to generate the same UMI counts was 7-fold. This is because UMI counts do not linearly increase with an increase of initial sequence reads due to UMI filtering. Unfortunately, the rate of

increase of sequence throughput was lower than that of the processing ability for single cells. It

will thus continue to be important to utilize limited initial fastq reads for high-throughput

single-cell RNA-seq methods.

Quartz-Seq2 is based on the poly-A tagging, which is one of the strategies of

converting first-strand cDNA to amplified cDNA [19,37]. However, the poly-A tagging efficiency

itself has not been improved for single-cell RNA-seq. In this study, three molecular biological

improvements on Quartz-Seq2 contributed to the increase of amplified cDNA, which led to high

UMI conversion efficiency (Figure 2 and Additional file1: Figure S1). In addition, improvement of

poly-A tagging steps was most efficient to increase amplified cDNA for Quartz-Seq2 (up 360%,

Additional file 1: Figure S1). Several previous studies improved the efficiency of conversion from

an mRNA molecule to amplifiable cDNA, and showed that the increase of amplified cDNA is a

good guidepost to improve UMI counts or quantitative performance for single-cell RNA-seq

[3,14,15]. These results suggest that improved poly-A tagging was the most important point for

high UMI conversion efficiency of Quartz-Seq2.

Several improvements proposed in this study could contribute to the further

development of other single-cell RNA-seq methods, as follows. (1) We showed that a decrease

of enzyme concentration in RT solution led to decreases in technical error and cDNA preparation

cost. (2) We applied the Sequence–Levenshtein distance for the design of cell barcodes

containing RT primer sets. The use of primer sets designed in this study allows the user to

correct mutations of at most two nucleotides of cell barcode, including substitution, insertion, or

deletion; the correction capability of these primer sets is higher than that of barcode sets used in

previous studies [1,3]. Such correction of the cell barcode increased the average UMI count by

3%–5%. (3) We hope that the spin-down collection system developed in this study can be

applied to other plate-based single-cell RNA-seq methods [2,3,11,12]. (4) In Quartz-Seq2, we

improved the efficiency of poly-A tagging itself (Figure 2). Several single-cell RNA-seq methods

including Quartz-Seq were based on poly-A tagging [4,5,10,15,19,38]. Such improvement may

directly contribute to these methods having increased quantitative performance.

We achieved a marked improvement in UMI conversion efficiency from initial reads to

UMI counts, allowing us to analyze 3- to 10-fold more single cells in limited sequence

experiments. However, it is difficult to sort more than 20,000 single cells using one cell-sorter

machine in a day. While there is greater scalability for cell sampling in droplet-based single-cell

RNA-seq methods than in cell sorter-based methods, these latter methods can utilize additional

information obtained in flow cytometry [39,40], which cannot be determined from transcriptome

data alone. These two types of method provide complementary approaches for investigating

complex biological phenomena.

In this study, we showed that Quartz-Seq2 has advantages in gene detection and the

identification of biological pathways via high UMI conversion efficiency (Figures 4, 5, and 7).

Moreover, we analyzed thousands of single cells from all cell cycle phases within a cell type

(Figure 6). These specifications with Quarz-Seq2 allow us to perform the global classification of

genes with variable expression in a cell type into those whose expression varies in association

with the cell cycle or independently of it.

A previous study utilizing the FUCCI fluorescent reporter system reported that

competence to respond to specific differentiation signaling was limited to only an early or a late

window of the G1 phase in human ES cells [41]. In the future, combining Quartz-Seq2 and

FUCCI/FUCCI2 or other fluorescent reporter systems will lead to an understanding of the

global picture of differentiation dynamics on competence, response, transition, and

commitment because Quartz-Seq2 can analyze changes of cell state for thousands of cells.

In summary, Quartz-Seq2 can be used to detect continuous cell states because of the

large number of cells with which it deals and the high efficiency of use of initial sequence reads.

Quartz-Seq2 can facilitate investigation of the cell state within a cell type, such as gradient or

stochastic changes of the cell population in organism development and disease progression.

Conclusions

In this study, we developed a high-throughput single-cell RNA-seq method, Quartz-Seq2, which

can analyze cells numbering up to 1,536 that are pooled together in a single sample.

Quartz-Seq2 allows us to effectively utilize initial sequence reads from a next-generation

sequencer. The UMI conversion efficiency in Quartz-Seq2 ranged from 30% to 50%, which is

much higher than for other single-cell RNA-seq methods (5%–20%). This was caused by the

improvements in several molecular biological steps including poly-A tagging. The technical gene

expression variability of Quartz-Seq2 was sufficiently low, and that was close to the theoretical

variability of a Poisson distribution. As we showed in the analysis of SVF and the ES/PrE mixture,

cell types in the population were identified with marker genes and functional terms that

characterized each cell type. We identified two types of *Cd34*-positive MSC in SVF, namely,

those that express numerous transcription factor-coding and secreted protein-coding genes

specific to each. Quartz-Seq2 can also be used to provide continuous data on cell states

because of the large number of cells with which it deals and the high efficiency in the use of initial

sequence reads. Quartz-Seq2 should facilitate investigations of the cell state within a cell type,

such as gradient or stochastic changes of cell populations in organism development and disease

progression.

Methods

Cell culture

Mouse embryonic stem cells were cultured as described previously [15]. Briefly, 1.0 x 10^5 cells

were seeded on a 60-mm dish coated with gelatin (0.1%). Cells were maintained in

GMEM-based medium containing 10% FBS and 1000 units/ml leukemia inhibitory factor

(Millipore ESGRO). The cell line used in this study was G6GRGFP, which was established in a

previous study [42]. To differentiate ES cells into primitive endoderm-like cells, 1.0 x 10^5 cells

were seeded on a 60-mm dish coated with gelatin and cultured in GMEM-based medium

containing 10% FBS, 1000 units/ml LIF, and 100 mM dexamethasone for 72 h.

Cell staining

To identify dead or damaged cells, propidium iodide (PI) was added to the cell suspension (final

concentration 1–2 µg/ml). As an indicator of an undifferentiated state in a suspension containing

a mixture of cells, Calcein-AM was used. Suspensions for cells cultured under maintenance and

differentiation conditions were prepared independently, and cells in the maintenance condition

were treated with 1 µg/ml Calcein-AM for 10 min on ice. After washing with PBS, cell

suspensions of ES cells and PrE cells at the same concentration were mixed. Hoechst 33342

was used as an indicator of the DNA content in a cell. The procedures were performed as

described previously [15].

Single-cell preparation for stromal vascular fraction

These experiments were carried out in accordance with the protocol approved by the National

Institute of Advanced Industrial Science and Technology (AIST) Animal Care and Use

Committee and the RIKEN Animal Experiment Committee. Subcutaneous fat tissues from three-

to four-month-old ICR male mice (n=3 per sample, two biological replicate samples) were

minced into small pieces and incubated with 0.4 U/mL collagenase NB4G (Serva) at 37°C for 35

min in a shaking water bath. The digested solution was sequentially filtered through 100-µm and

40-µm cell strainers (Corning), followed by centrifugation at 250 × g for 5 min to remove mature

adipocytes. The pellet was treated with erythrocyte lysis buffer (BD Biosciences) and centrifuged

at 180 × g for 5 min. The nucleated cells were suspended in HBSS with 0.1% BSA, filtered

through a 20-μm cell strainer (pluriSelect), and then kept on ice (Cell solution A). The cell

aggregates that did not pass through the 20-μm strainer were further treated with Accutase

(Thermo Fisher Scientific) at 37°C for 15 min to dissociate them into single cells, centrifuged at

180 × g for 5 min, and suspended in HBSS with 0.1% BSA (Cell solution B). Cell solutions A and

B were mixed and again filtered through the 20-μm cell strainer, followed by centrifugation at 180

× g for 5 min. The pellet was resuspended with HBSS with 0.1% BSA, stained with PI, and used

for single-cell analysis.

RNA preparation

Total RNA was purified from cultured cells using Direct-zol RNA MiniPrep kit (Zymo Research)

with TRIzol RNA Isolation Reagents (Thermo). We measured the concentration of purified total

RNA using a NanoDrop 1000 Spectrophotometer (Thermo). We confirmed that the RNA integrity

number of total RNA was over 9.5 using an Agilent RNA 6000 Nano Kit (Agilent). For

high-throughput single-cell RNA-seq, we prepared total RNA from a single cell with ERCC

spike-in mix I. First, we diluted the ERCC spike mix I by tenfold. We then added 6 μL of 1:10

diluted ERCC spike-in mix I per 10 μg of total RNA. We used diluted 10 pg of total RNA with

ERCC spike mix I for the technical validation of Quartz-Seq2.

Bulk RNA-seq methods for populations of cells

We prepared sequence library DNA with 1 μg of total RNA using NEBNext Poly(A) mRNA

Magnetic Isolation Module and NEBNext Ultra Directional RNA Library Prep Kit. The total RNA

did not contain ERCC spin-in mix I. In addition, we used SuperScript III instead of ProtoScript in

the reverse-transcription step and KAPA HiFi DNA polymerase instead of NEBNext High-Fidelity

PCR DNA polymerase in the PCR step. The resulting sequence library DNA was analyzed by

HiSeq2500.

Design of cell barcodes

The selection of 384 or 1,536 sequences for v3.1 and v3.2 barcode primer sets was performed

as described below. First, 1,582 or 4,714 candidate sequences were created using the

DNABarcodes package of R Bioconductor for the v3.1 set with 14-mer and the v3.2 set with

15-mer, respectively (version: 1.0.0). To reduce the loss of reads converted into UMI counts, we

applied the Sequence–Levenshtein distance as an edit distance in order to maximize the ability

to correct errors that occur during the synthesis of oligonucleotides or sequencing [22]. The

minimum distance between any two sequences was controlled to 5, leading to the correction of a

maximum of two errors of substitution, insertion, or deletion. As the base composition of the

created sequences was not uniform, we selected 384 or 1,536 sequences from among 1,582 or

4,714 created sequences so that the variance of base composition decreased. These sequences

are listed in Additional file 3: Table S2.

Single-cell collection using flow cytometry

Stained cells were analyzed using flow cytometry SH800 (Sony) or MoFlo Astrios EQ (Beckman

Coulter). For SH800, we used 130-μm microfluidic sorting chips. For MoFlo Astrios EQ, we used

100-μm nozzle sizes. Each cell sorter was equipped with a custom-made splash-guard to

prevent contamination from unexpected droplets between the target well and neighboring wells.

We used two sets of RT primer for Quartz-Seq2. The set of v3.1 RT primers has 384 kinds of

unique cell barcodes, with a length of 14 nucleotides (OPC purification, FASMAC). The set of

v3.2 RT primers has 1,536 kinds of unique cell barcodes, with a length of 15 nucleotides (OPC

purification, Sigma). The set of v3.1 RT primers corresponds to one set of the 384-well PCR

plate with lysis buffer, the wells of which have unique barcodes. The set of v3.2 RT primers

corresponds to four sets of the 384-well PCR plate with lysis buffer. The RT primer position in the

384-well plate and the sequence were as described in Additional file 3: Table S2. Single cells

were isolated in the 384-well PCR plate with 1 μL of lysis buffer (0.1111 μM respective RT

primers, 0.12 mM dNTP mix, 0.3% NP-40, 1 unit/μL RNasin plus) containing ERCC spike mix I.

During single-cell sorting, the 384-well PCR plate was kept on a 384 aluminum stand at 4°C. For

Moflo Astrios EQ, we used G5498B-060 Insert 384 Eppendorf twin.tec PCR as a 384 aluminum

stand (Agilent). For SH800, we used an SH800 384 aluminum stand (Sony). Immediately after

the cell collection, the plate was temporarily sealed with LightCycler 480 Sealing Foil (Roche)

and the sealed 384-well PCR plate was centrifuged at 10,000 g and 4ºC for 1 min using TOMY

MX307, equipped with a Rack-in-Rotor and centrifugation rack PCR96-02. These steps were

very important to collect the droplet with a single cell in lysis buffer in an efficient manner. By

altering the volume of lysis buffer from 0.4 µL (as described in the original Quartz-Seq paper) to

1 µL (Quartz-Seq2), bubbling of lysis buffer was not required before single-cell sorting and the

lysis buffer could easily be handled. We agitated the plate at 2,600 rpm and 4ºC for 1 min using

ThermoMixer C (Eppendorf), after which we centrifuged the plate again. We then peeled open

the temporary seal and re-sealed it with Agilent PlateLoc Thermal Microplate Sealer (Agilent).

The resulting 384-well plate was then immediately cryopreserved at −80ºC and maintained

under such conditions until subsequent reverse transcription for cell barcoding. After

cryopreservation, we performed subsequent whole-transcript amplification using the

cryopreserved 384-well PCR plate within a few months.

Whole-transcript amplification of Quartz-Seq2

Cryopreserved 384-well plates with single-cell lysate were centrifuged at 10,000 g and 4ºC for 1

min. Subsequently, we denatured total RNA in each 384 plate at 70ºC for 90 s and hybridized the

RT primer to poly-adenylated RNA at 35ºC for 15 s using the C1000/S1000 thermal cycler. The

resulting plates were again centrifuged at 10,000 g and 4ºC for 1 min. Next, the plates were

placed on the 384 aluminum plate at 0ºC. We peeled away the seal and added 1 µL of RT

premix (2x Thermopol buffer, 5 units/µL SuperScript III, 0.55 units/µL RNasin plus) to 1 µL of

lysis buffer for each well using a Mantis microfluidic dispensing system (Formulatrix) or a 384

Transfer Plate system (1859-384S, Watson). The above RT solution was used for the RT25

condition. For the RT100 condition, we used the following RT solution: 2x Thermopol buffer, 20

units/µL SuperScript III, and 2.2 units/µL RNasin plus. We sealed the plates again and agitated

them at 2,600 rpm and 4ºC for 1 min. The plates were then centrifuged at 10,000 g and 4ºC for 1

min. We then performed reverse transcription at 35ºC for 5 min and 50ºC for 50 min. The reverse

transcription was stopped at 70ºC for 15 min. Then, the plates were placed on a prechilled

aluminum block, after which we peeled off their seals. Subsequently, we turned the plates upside

down on the assembled collector type A or type B (supplemental figure). We mainly used type A.

We centrifuged the plates with an assemble collector at 3,010 g and 4ºC for 3 min with

swing-bucket rotors. Subsequently, we collected the cDNA solution into a disposable reservoir.

Typically, we obtained 650–700 µL of cDNA solution from one 384-well PCR plate. We purified

and concentrated the cDNA solution using the DNA Clean & Concentrator™-5 kit (Zymo

Research). We used three purification columns for one 384-well PCR plate in the case of the

v3.1 RT primer system (384-cell barcode). Purified cDNA was extracted into 20 µL of

nuclease-free water from one column purification and transferred into an eight-linked PCR tube

(TaKaRa). The PCR tubes were placed on an aluminum PCR stand at 0ºC. We added 25 µL of

TdT solution [1x Thermopol buffer, 2.4 mM dATP, 0.0384 units/µL RNase H (Invitrogen), 26.88

units/µL terminal transferase (Roche)] into 20 µL of extracted cDNA using a pipette at 0ºC. The

resulting 45 µL of TdT solution was mixed with a pipette at 0ºC or ThermoMixer at 2,000 g and

0ºC for 1 min. Immediately thereafter, the PCR tubes were centrifuged at 10,000 g and 0ºC for 1

min. We used a C1000/S1000 thermal cycler equipped with the 96-Deep Well Reaction Module

for the following steps. The PCR tubes were placed on the block of the thermal cycler, which had

been prechilled to 0ºC. We then performed a poly-A tailing reaction at 37ºC for 75 s. The solution

was inactivated at 65ºC for 10 min. The PCR tubes were placed on an aluminum PCR stand at

0ºC. We then dispensed approximately 11 µL of solution into four wells from 45 µL of TdT

solution. We added 46.16 µL of PCR I premix (1.08492x MightyAmp Buffer version 2, 0.06932

µM Tagging primer, 0.05415 units/µL MightyAmp DNA polymerase) to 11 µL of TdT solution for

the respective wells of the PCR tube. We performed gentle inversion mixing on the resulting

solution in the PCR tube. The tubes were then centrifuged at 10,000 g and 4ºC for 1 min.

Subsequently, the solution was mixed with ThermoMixer at 2,000 rpm and 4ºC for 2 min. Then,

we spun down the tube again. Next, we denatured the solution at 98ºC for 130 s and hybridized

Tagging primer to poly-A-tailed cDNA at 40ºC for 1 min. After that, we performed the "Increment

step" by heating to 68ºC at 0.2ºC every second and performed second-strand synthesis at 68ºC

for 5 min. The tubes were placed on an aluminum PCR stand at 0ºC. We added 50.232 µL of

PCR II premix (0.99697x MightyAmp Buffer version.2, 1.8952 µM gM primer) to 56.16 µL of PCR

I solution. We performed gentle inversion mixing on the resulting solution in the PCR tube. The

tubes were then centrifuged at 10,000 g and 4ºC for 1 min. Subsequently, the solution was

mixed with ThermoMixer at 2,000 rpm and 4ºC for 2 min, after which we spun down the tube

again. We then placed it on the block of the thermal cycler at 68ºC. Subsequently, we amplified

the cDNA for 11 cycles under the following conditions: 98ºC for 10 s, 65ºC for 15 s, and 68ºC 5

min. We then incubated the tube at 68ºC for an additional 5 min. Finally, we transferred all of the

PCR solution, derived from one 384-well PCR plate, to a 50-mL Polypropylene Centrifuge Tube

(Watson). Typically, we obtained approximately 1.2 mL of PCR solution per 384-well PCR plate.

We added 32 µL of 3 M sodium acetate (pH 5.2) and 6420 µL PB-Buffer (Qiagen) to the PCR

solution. The mixture was then purified using a MinElute Spin Column (Qiagen). Purified cDNA

was extracted into 40 µL of nuclease-free water. We additionally purified the cDNA with 32 µL of

Ampure XP beads. Finally, we obtained 32 µL of purified cDNA. We checked the length

distribution of amplified cDNA with an Agilent High Sensitivity DNA Kit (Agilent). The typical

average size of the amplified cDNA in Quartz-Seq2 was approximately 1,400 bp (Additional file

1: Figure S2c). Primer sequences are listed in Additional file 3: Table S2.

In the case of the usage of the v3.2 RT primer in Quartz-Seq2, we modified the above

steps as follows. After reverse transcription, we collected cDNA solution into a disposable

reservoir from four sets of 384-well plates, which corresponded to 1,536 wells. We purified and

concentrated the cDNA solution using eight purification columns for four 384-well PCR plates in

the case of the v3.2 RT primer system. In the PCR step, we amplified cDNA for nine cycles with

the following conditions: 98ºC for 10 s, 65ºC for 15 s, and 68ºC for 5 min. Finally, we transferred

all of the PCR solution derived from four 384-well plates to a 50-mL Polypropylene Centrifuge

Tube (Watson). Typically, we obtained approximately 3.5 mL of PCR solution per four 384-well

PCR plates. We added 88 µL of 3 M sodium acetate (pH 5.2) and 17.6 mL of PB-Buffer (Qiagen)

to the PCR solution. The mixture was purified using the MinElute Spin Column (Qiagen).

Subsequently, cDNA was again purified using Ampure XP magnetic beads.

For the Quartz-Seq1-like reaction, we performed the following procedure in

accordance with our previous study. We added 1 µL of RT premix (2x PCR buffer, 5 units/µL

SuperScript III, 0.55 units/µL RNasin plus) to 1 µL of lysis buffer. We then performed reverse

transcription at 35ºC for 5 min and 45ºC for 20 min. This reverse transcription was stopped at

70ºC for 15 min. We added 5 µL of ExoIB solution (1.6x Exonuclease I buffer, 3.2x PCR buffer,

16 mM DTT) and 20 µL of TdT solution [1x PCR buffer, 3 mM dATP, 0.0384 units/µL RNase H

(Invitrogen), 33.6 units/µL terminal transferase (Roche)] into 20 µL of extracted cDNA using a

pipette at 0ºC. The PCR tubes were placed on the block of the thermal cycler that had been

prechilled to 0ºC. We performed a poly-A tailing reaction at 37ºC for 75 s. We then denatured the

solution at 98ºC for 130 s and hybridized Tagging primer to poly-A-tailed cDNA at 40ºC for 1 min.

After that, we performed second-strand synthesis at 68ºC for 5 min. Subsequently, we amplified

the cDNA via a PCR reaction for 12 cycles.

Preparation of truncated sequence adaptor

The truncated sequence adaptor was composed of an rYshapeP5 primer (HPLC-purified) and

rYshapeP7LT primers (HPLC-purified), which had a TruSeqLT-compatible pool barcode. We

prepared 100 µM respective primers with adaptor buffer (10 mM Tris-HCl pH7.8, 0.1 mM EDTA

pH8.0, 50 mM NaCl). We added 5 µL of 100 µM rYshapeP5 primer and rYshapeP7LTxx primer

into a single PCR tube. We denatured the solution at 90ºC for 90 s. After that, we achieved

annealing by cooling to 10ºC by 0.5ºC every 30 s and then maintaining the sample at 4ºC. We

then placed the tube on an aluminum PCR stand at 0ºC. Subsequently, we added adaptor buffer,

which was prechilled at 0ºC, to 10 µL of 50 µM truncated adaptor. Finally, we obtained about 50

µL of 10 µM truncated adaptor. We cryopreserved 1 µL of 10 µM truncated adaptor in a PCR

tube at −80ºC until usage in the adaptor ligation step. Primer sequences are listed in Additional

file 3: Table S2.

Sequence library preparation of Quartz-Seq2

We added 130 µL of nuclease-free water including 5–10 ng of amplified cDNA into a Crimp-Cap

microTUBE with AFA Fiber. We then performed cDNA fragmentation using an LE220

Focused-ultrasonicator (Covaris) under the following conditions: duty factor 15%, peak incident

power 450 W, cycles per burst 200, and treatment time 80 s. We purified and concentrated the

cDNA solution using the DNA Clean & Concentrator™-5 kit. Purified cDNA was extracted into 10

μL of nuclease-free water from one column purification and transferred into an eight-linked PCR

tube (TaKaRa).

We then added 2 μL of End-Repair premix [1.4 μL of End repair & A-tailing buffer and

0.6 μL of End repair & A-tailing Enzyme (KAPA Biosystems)] to 10 μL of fragmented cDNA

solution. Subsequently, we mixed the solution by pipetting on an aluminum PCR stand at 0°C.

The PCR tubes were placed on the block of the thermal cycler, which had been prechilled to

20°C. We then incubated the tubes at 37°C for 60 min and 65°C for 30 min. After this, we added

2 μL of adaptor buffer (1.5 μM truncated adaptor, 10 mM Tris-HCl pH7.8, 0.1 mM EDTA pH8.0,

50 mM NaCl) and 8 μL of ligation premix [6 μL of ligation buffer, 2 μL of DNA ligase (KAPA

Biosystems)] at 4°C. The 22 μL of ligation solution was well mixed by pipetting at 4°C. Then, we

performed adaptor ligation at 20°C for 15 min. After ligation, we added 18 μL of Ampure XP

beads to 22 μL of adaptor ligation solution and mixed them well. Adaptor-ligated cDNA was

extracted into 20 μL of nuclease-free water. We added 32 μL of PCR premix [25 μL of 2xKAPA

HiFi ready mix, 1.75 μL of 10 μM TPC2 primer (HPLC-purified), 10 μM P5-gMac hybrid primer

(HPLC-purified)] to 18 μL of adaptor-ligated cDNA. We denatured the solution at 98°C for 45 s.

Subsequently, we amplified cDNA for eight cycles under the following conditions: 98°C for 15 s,

60°C for 30 s, and 72°C for 1 min. Finally, we additionally incubated the tube at 72°C for 5 min.

We added 40 μL of Ampure XP beads to 50 μL of PCR solution. Purified

sequence-library DNA was eluted into 20–30 μL of nuclease-free water. We checked the DNA

concentration and DNA size of sequence library DNA using the Agilent High Sensitivity DNA Kit

(Agilent) and QuantiFluor® dsDNA System (Promega). We also prepared sequence library DNA

with Tn5 transposase, in accordance with a modified version of a procedure used in a previous

study [6]. Specifically, the modifications were as follows. We used 0.75 ng of amplified cDNA for

the Nextera XT library preparation kit. We amplified sequence library DNA with the P5-gMac

hybrid primer and the Nextera XT primer, which has the P7 sequence.

Deep sequencing for Quartz-Seq2

We analyzed the sequence library DNA from Quartz-Seq2 using NextSeq500 (Illumina) and

HiSeq2500 (Illumina). For the Read1 sequence, we used a custom sequence primer named

Read1DropQuartz primer (HPLC-purified). In the case of the v3.1 RT primer, sequence

specification was as follows (Read1: 22 cycles, Index1: 6 cycles, Read2: 64–118 cycles). In the

case of the v3.2 RT primer, sequence specification was as follows (Read1: 23 cycles, Index1: 6

cycles, Read2: 63 cycles). When Read1 length was within 64 nt we mainly used the NextSeq

500/550 High Output v2 Kit (75 cycles), which can be used for sequencing up to 92 nt. For long

sequences (over 64 nt), we analyzed sequence library DNA using the HiSeq Rapid SBS Kit v2.

Sequence analysis with HiSeq2500 was supported by the Genomic Network Analysis Support

facility (GeNAS) in RIKEN.

Amplification-free single-cell qPCR analysis

Amplification-free single-cell qPCR was performed in accordance with our previous study with

the following modifications [15]. We collected a single cell into 1 µL of lysis buffer (2.5 µM

random hexamer, 1 mM dNTP mix, 0.3% NP40, 2 units/µL RNasin plus) of a 384-well PCR plate

(Eppendorf) using SH800. The resulting 384-well PCR plate was cryopreserved at −80ºC. We

denatured total RNA in the 384-well PCR plate at 70ºC for 1.5 min using a C1000/S1000 thermal

cycler and hybridized the random primer to RNA at 0ºC for 2 min. We added 1 µL of

reverse-transcription solution (2x SSIV buffer, 10 mM DTT, 2 units/µL RNasin plus, 20 units/µL

SuperScript IV) to the lysis buffer. Subsequently, we performed reverse transcription at 23ºC for

10 min and 50ºC for 10 min. The reverse transcription was stopped at 80ºC for 10 min. The

resulting solution was diluted with qPCR solution (10 mM Tris-HCl pH 8.0, 0.05% Tween-20).

The obtained diluted solution was then used for qPCR detection with QuantiTect SYBR Green

PCR Master Mix and the LightCycler480 system. For primer sets for each gene, see Additional

file 3: Table S2.

Drop-seq experiments and data analysis

Drop-seq was performed as reported previously [6] and in line with an online protocol

(http://mccarrolllab.com/dropseq/), but with the following modifications: flow rates for oil and

aqueous suspensions were 14,000 µl/h and 4,000 µl/h, respectively. The diameter of droplets

was 95–100 µm. Microfluidic devices were fabricated by Fluidware Technologies (Japan). The

lot number of barcoded beads was 051415 (ChemGenes). Data analysis for Drop-seq was

performed as described online (http://mccarrolllab.com/dropseq/). The versions of the software

and databases were as follows: STAR: v2.5.1b; mouse genome: mm10; genome annotation:

gencode GRCm38.p4 vM9; and Drop-seq tools: v1.11.

Quartz-Seq2 read alignment and generation of digital expression data

The structure of the sequence library and data processing was designed based on those of

Drop-seq ([6] and online as referenced above). BCL files generated by Illumina NextSeq500

were converted to fastq files by bcl2fastq2 (v2.17.1.14) with demultiplexing pool barcodes. The

--mask-short-adapter-reads parameter was set to 20. If needed, fastq reads were randomly

downsampled by seqtk software (version: sgdp). The last two bases of Read2, the sequence

quality of which was relatively low, were removed by FASTX-Toolkit (version: 0.0.14). Fastq files

for Read1 and Read2 were converted to a bam file by FastqToSam of Picard tools (version:

1.134). Extracting cell barcodes and UMI (equal to molecular barcode) and filtering out reads

with low barcode quality were performed using Drop-seq tools (version: 1.11). The resulting bam

files were re-converted to fastq files by SamToFastq of Picard tools, and mapped to the mouse

genome (mm10) using STAR (version: 2.5.1b). After sorting the resulting bam files by SortSam

of Picard tools, the unaligned bam and aligned bam were merged by MergeBamAlignment of

Picard tools. Then, gene names were assigned to each record using TagReadWithGeneExon of

Drop-seq tools and gtf file (version: gencode GRCm38.p4 vM9). For the correction of errors of

cell barcodes considering the Sequence–Levenshtein distance, a custom-made python program

was used, which enabled the correction of up to two errors of substitution, insertion, or deletion.

This program used python2 (version: 2.7.11+), PypeR (version: 1.1.2), R (version: 3.2.3),

Bioconductor (version: 3.2), and Bioconductor package DNABarcodes (version: 1.0.0). Finally,

the UMI for each gene for each cell was counted using DigitalExpression of Drop-seq tools,

which generated the digital expression matrix. To generate a non-UMI-filtered matrix, a

custom-made python program that counts reads for each gene for each cell was used.

Dimensionality reduction, clustering, and term analysis

Cells with low detected gene counts were removed for further analysis (Quartz-Seq2 on ES/PrE

mixture: 4,000 genes; Quartz-Seq2 on SVF: 500 genes). Total counts of each cell were

normalized to 10,000 or the mean of total UMI counts for cells. UMI counts had 1 added to them

and were then log-transformed (natural logarithm), after which they were scaled to have mean

and variance for each gene of 0 and 1, respectively. For PCA, UMI counts for all detected genes

(for ES/PrE mixture) or genes with highly variable expression (for SVF) were used. For t-SNE,

the top 10 to 40 principal components of matrices produced by PCA were used. For clustering,

the DBSCAN algorithm was used. The values of the parameter epsilon were 0.69 for

Quartz-Seq2 on the 4,500 mouse ES/PrE mixture, 5 for Quartz-Seq2 on the 384 mouse ES/PrE

mixture, 3 for Drop-seq on the 500 mouse ES/PrE mixture, and 2.2 for SVF analysis.

Significantly enriched GO terms with the top principal components were calculated using the

GO-PCA package [43].

Identification of differentially expressed genes for each cluster

Marker genes for each cluster were identified based on a generalized linear model. The

difference in deviance between two models, in which the gene was or was not differentially

expressed between two clusters, was calculated for the genes. To filter out noise, pseudocount 1

was added to the averaged gene expression in clusters and only genes with fold change of 2 or

more between two clusters were further analyzed. P-values were calculated as 1 − cumulative

density function of a chi-squared continuous random variable for difference in deviance. After

corrections for multiple testing, genes with FDR of less than 0.05 were identified as differentially

expressed genes for the cluster. For comparison between differentially expressed genes

identified by scRNA-seq methods and those identified by bulk RNA-seq, this pseudocount was

not added.

Identification of cell cycle-associated genes and cell cycle-phase independent variable

genes

To identify genes for which the expression changes depending on the phase of the cell cycle, we

used the intensity of Hoechst 33342 staining measured using a cell sorter. First, cells were

discretized into 40 equal-sized buckets based on the rank of Hoechst 33342 staining intensity.

Then, the CV of averaged UMI counts for a gene in each bin was calculated. After z-scaling of

this, genes with high z-scores were identified. To identify genes whose expression fluctuates in a

manner not dependent on the cell cycle phase, we calculated the CV of UMI counts for a gene in

each cell and the CV of averaged UMI counts for a gene in each bin. After z-scaling of such data,

the genes of which difference between two scaled CV was large were identified as cell

cycle-phase independent variable genes.

Enrichment analysis on pathway and Gene Ontology (GO) terms

Pathways that were particularly enriched for the differentially expressed genes were calculated

using the ReactomePA package of R Bioconductor (version: 1.14.4, [44]) and Metascape

(http://metascape.org/ [45]). The cut-off parameter for the q value was 0.05. Terms that were

enriched for genes with highly variable expression were calculated using DAVID, the Database

for Annotation, Visualization and Integrated Discovery (version: 6.8, [46,47]). Ontology used for

calculation was GOTERM_BP_FAT, GOTERM_MF_FAT, and GOTERM_CC_FAT. The terms

with FDR < 0.05 were identified as enriched terms.

1000

1001   **Data resources**

1002   Raw and processed data files for Quartz-Seq2 and Drop-seq experiments are available under

1003   GEO: GSE99866 for single-cell RNA-seq data and DRA002954 for bulk-RNA-seq data.

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013 **List of abbreviations**

1014

1015 UMI: unique molecular identifier

1016 PCR: polymerase chain reaction

1017 SCC: Spearman's rank correlation coefficient

1018 CV: coefficient of variation

1019 ES cell: embryonic stem cell

1020 PrE cell: primitive endoderm cell

1021 SVF: stromal vascular fraction

1022 RT: reverse-transcription

1023 Dex: dexamethasone

1024 PCA: principal component analysis

1025 MSC: mesenchymal stem cell

1026 TdT: terminal deoxynucleotidyl transferase

1027 GO: Gene Ontology

1028 FDR: false discovery rate

1029 t-SNE: t-distributed stochastic neighbor embedding

1030 nt: nucleotide

1031 bp: base pair

1032 SSC: side scatter

1033

1034

1035

1036

1037

1038

1039

1040

1041

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

The data have been deposited under GEO accession number GSE99866 and DRA accession

number: DRA002954.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Projects for Technological Development, Research Center

Network for Realization of Regenerative Medicine by Japan, the Japan Agency for Medical

Research and Development (AMED), and the Japan Science and Technology Agency (JST).

The sequence analyses of bulk RNA-seq were supported by MEXT KAKENHI (No. 221S0002)

and JSPS KAKENHI Grant Number JP24651218 and JSPS KAKENHI Grant number

JP15K16329. The sequence operations of Quartz-Seq2 using HiSeq2500 were supported by the

Platform Project for Supporting Drug Discovery and Life Science Research (Platform for Drug

Discovery, Informatics, and Structural Life Science) from AMED.

Author's contributions

YS, HD and IN designed the study. YS designed and performed experiments of single-cell

RNA-seq. HD designed and performed data analysis. HT and AK designed and performed

experiments of mouse stromal vascular fraction. ME assisted the experiments. TH assisted the

1071 development of spin-down collection system. YS, HD, HT, AK and IN wrote the manuscript. All

1072 authors read and approved the final manuscript.

1073

Acknowledgements

We greatly appreciate the Drop-seq team for useful comments on our technical introduction to

Drop-seq. We thank Mr. Akihiro Matsushima and Mr. Manabu Ishii for assistance with the

infrastructure for the data analysis. We thank Dr. Yutaka Suzuki for assistance with the bulk

RNA-seq. We are also grateful to Dr. Mika Yoshimura and Dr. Haruka Ozaki (RIKEN) for

assistance with the bulk RNA-seq analysis. Thanks also go to Sony's cell sorter team for special

customization of the 384-well sorting mode of SH800.

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

References

1100 **References**

1101

1. Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, et al. Massively parallel
digital transcriptional profiling of single cells. Nat Commun. 2017;8:14049.
2. Muraro MJ, Dharmadhikari G, Grün D, Groen N, Dielen T, Jansen E, et al. A Single-Cell
Transcriptome Atlas of the Human Pancreas. Cell Syst. 2016;3:385–394.e3.
3. Hashimshony T, Senderovich N, Avital G, Klochendler A, de Leeuw Y, Anavy L, et al.
CEL-Seq2: sensitive highly-multiplexed single-cell RNA-Seq. Genome Biol. 2016;17:892.
4. Fan X, Zhang X, Wu X, Guo H, Hu Y, Tang F, et al. Single-cell RNA-seq transcriptome
analysis of linear and circular RNAs in mouse preimplantation embryos. Genome Biol.
2015;16:236.
5. Nakamura T, Yabuta Y, Okamoto I, Aramaki S, Yokobayashi S, Kurimoto K, et al. SC3-seq: a
method for highly parallel and quantitative measurement of single-cell gene expression. Nucleic
Acids Res. 2015;43:e60–0.
6. Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, et al. Highly Parallel
Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. Cell.
2015;161:1202–14.
7. Klein AM, Mazutis L, Akartuna I, Tallapragada N, Veres A, Li V, et al. Droplet Barcoding for
Single-Cell Transcriptomics Applied to Embryonic Stem Cells. Cell. 2015;161:1187–201.
8. Fan HC, Fu GK, Fodor SPA. Combinatorial labeling of single cells for gene expression
cytometry. Science. 2015;347:1258367–7.
9. Bose S, Wan Z, Carr A, Rizvi A, Vieira G, Pe'er D, et al. Scalable microfluidics for single-cell
RNA printing and sequencing. Genome Biol. 2015;16:120.
10. Streets AM, Zhang X, Cao C, Pang Y, Wu X, Xiong L, et al. Microfluidic single-cell
whole-transcriptome sequencing. Proc. Natl. Acad. Sci. 2014;111:7048–53.
11. Jaitin DA, Kenigsberg E, Keren-Shaul H, Elefant N, Paul F, Zaretsky I, et al. Massively
Parallel Single-Cell RNA-Seq for Marker-Free Decomposition of Tissues into Cell Types.
Science. 2014;343:776–9.

1128 12. Soumillon M, Cacchiarelli D, Semrau S, van Oudenaarden A, Mikkelsen TS. Characterization

1129 of directed differentiation by high-throughput single-cell RNA-Seq. 2014.

1130 13. Islam S, Zeisel A, Joost S, La Manno G, Zajac P, Kasper M, et al. Quantitative single-cell

1131 RNA-seq with unique molecular identifiers. Nat. Methods. 2013;11:163–6.

1132 14. Picelli S, Björklund ÅK, Faridani OR, Sagasser S, Winberg G, Sandberg R. Smart-seq2 for

1133 sensitive full-length transcriptome profiling in single cells. Nat. Methods. 2013;10:1096–8.

1134 15. Sasagawa Y, Nikaido I, Hayashi T, Danno H, Uno KD, Imai T, et al. Quartz-Seq: a highly

1135 reproducible and sensitive single-cell RNA sequencing method, reveals non-genetic

1136 gene-expression heterogeneity. Genome Biol. 2013;14:R31.

1137 16. Hashimshony T, Wagner F, Sher N, Yanai I. CEL-Seq: Single-Cell RNA-Seq by Multiplexed

1138 Linear Amplification. Cell Rep. 2012;2:666–73.

1139 17. Ramsköld D, Luo S, Wang Y-C, Li R, Deng Q, Faridani OR, et al. Full-length mRNA-Seq

1140 from single-cell levels of RNA and individual circulating tumor cells. Nat. Biotechnol.

1141 2012;30:777–82.

1142 18. Islam S, Kjallquist U, Moliner A, Zajac P, Fan JB, Lonnerberg P, et al. Characterization of the

1143 single-cell transcriptional landscape by highly multiplex RNA-seq. Genome Res. 2011;21:1160–

1144 7.

1145 19. Tang F, Barbacioru C, Wang Y, Nordman E, Lee C, Xu N, et al. mRNA-Seq

1146 whole-transcriptome analysis of a single cell. Nat. Methods. 2009;6:377–82.

1147 20. Grün D, Kester L, van Oudenaarden A. Validation of noise models for single-cell

1148 transcriptomics. Nat. Methods. 2014;11:637–40.

1149 21. Ziegenhain C, Vieth B, Parekh S, Reinius B, Guillaumet-Adkins A, Smets M, et al.

1150 Comparative Analysis of Single-Cell RNA Sequencing Methods. Mol. Cell. 2017;65:631–4.

1151 22. Buschmann T. DNABarcodes: an R package for the systematic construction of DNA sample

1152 tags. Bioinformatics. 2017;:btw759.

1153 23. Ilicic T, Kim JK, Kolodziejczyk AA, Bagger FO, McCarthy DJ, Marioni JC, et al. Classification

1154 of low quality cells from single-cell RNA-seq data. Genome Biol. 2016;17:57.

1155 24. Zalzman M, Falco G, Sharova LV, Nishiyama A, Thomas M, Lee S-L, et al. Zscan4 regulates

1156 telomere elongation and genomic stability in ES cells. Nature. 2010;464:858–63.

25. Pak J, Lee JH, Park KS, Park M, Kang L-W, Lee SH. Current use of autologous adipose

tissue-derived stromal vascular fraction cells for orthopedic applications. J. Biomed. Sci.

2017;24:1496.

26. Trounson A, McDonald C. Stem Cell Therapies in Clinical Trials: Progress and Challenges.

Cell Stem Cell. 2015;17:11–22.

27. Baer PC, Geiger H. Adipose-Derived Mesenchymal Stromal/Stem Cells: Tissue Localization,

Characterization, and Heterogeneity. Stem Cells Int. 2012;2012:1–11.

28. Lindroos B, Suuronen R, Miettinen S. The Potential of Adipose Stem Cells in Regenerative

Medicine. Stem Cell Rev. 2010;7:269–91.

29. Kwon HM, Hur S-M, Park K-Y, Kim C-K, Kim Y-M, Kim H-S, et al. Multiple paracrine factors

secreted by mesenchymal stem cells contribute to angiogenesis. Vascular Pharmacology.

2014;63:19–28.

30. Madrigal M, Rao KS, Riordan NH. A review of therapeutic effects of mesenchymal stem cell

secretions and induction of secretory modification by different culture methods. J Transl Med.

2014;12:71.

31. Rennert RC, Januszyk M, Sorkin M, Rodrigues M, Maan ZN, Duscher D, et al. Microfluidic

single-cell transcriptional analysis rationally identifies novel surface marker profiles to enhance

cell-based therapies. Nat Commun. 2016;7:11945.

32. Baer PC. Adipose-derived mesenchymal stromal/stem cells: An update on their phenotype in

vivo and in vitro. World J Stem Cells. 2014;6:256.

33. Mabuchi Y, Houlihan DD, Akazawa C, Okano H, Matsuzaki Y. Prospective Isolation of

Murine and Human Bone Marrow Mesenchymal Stem Cells Based on Surface Markers. Stem

Cells Int. 2013;2013:1–7.

34. Dromard CC, Barreau C, Andr M, Berger-M ller S, Casteilla L, Planat-Benard V. Mouse

adipose tissue stromal cells give rise to skeletal and cardiomyogenic cell sub-populations. Front

Cell Dev Biol. 2014;2:697.

35. Heneidi S, Simerman AA, Keller E, Singh P, Li X, Dumesic DA, et al. Awakened by Cellular

Stress: Isolation and Characterization of a Novel Population of Pluripotent Stem Cells Derived

from Human Adipose Tissue. Ljubimov AV, editor. PLoS ONE. 2013;8:e64752.

36. Wachi H. Role of Elastic Fibers on Cardiovascular Disease. J. Health Sci. The

Pharmaceutical Society of Japan; 2011;57:449–57.

37. Kurimoto K. An improved single-cell cDNA amplification method for efficient high-density

oligonucleotide microarray analysis. Nucleic Acids Res. 2006;34:e42–2.

38. Yang L, Ma Z, Cao C, Zhang Y, Wu X, Lee R, et al. MR-Seq: measuring a single cell's

transcriptome repeatedly by RNA-seq. Science Bulletin. 2017.

39. Velten L, Haas SF, Raffel S, Blaszkiewicz S, Islam S, Hennig BP, et al. Human

haematopoietic stem cell lineage commitment is a continuous process. Nat. Cell Biol.

2017;Complex Sy:1695.

40. Kowalczyk MS, Tirosh I, Heckl D, Rao TN, Dixit A, Haas BJ, et al. Single-cell RNA-seq

reveals changes in cell cycle and differentiation programs upon aging of hematopoietic stem

cells. Genome Res. 2015;25:1860–72.

41. Pauklin S, Vallier L. The Cell-Cycle State of Stem Cells Determines Cell Fate Propensity.

Cell. 2013;155:135–47.

42. Shimosato D, Shiki M, Niwa H. Extra-embryonic endoderm cells derived from ES cells

induced by GATA Factors acquire the character of XEN cells. BMC Dev. Biol. 2007;7:80.

43. Wagner F. GO-PCA: An Unsupervised Method to Explore Gene Expression Data Using Prior

Knowledge. Li X, editor. PLoS ONE. 2015;10:e0143196.

44. Yu G, He Q-Y. ReactomePA: an R/Bioconductor package for reactome pathway analysis

and visualization. Mol. BioSyst. 2016;12:477–9.

45. Tripathi S, Pohl MO, Zhou Y, Rodriguez-Frandsen A, Wang G, Stein DA, et al. Meta- and

Orthogonal Integration of Influenza ?OMICs? Data Defines a Role for UBR4 in Virus Budding.

Cell Host Microbe. 2015;18:723–35.

46. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene

lists using DAVID bioinformatics resources. Nat Protoc. 2008;4:44–57.

47. Huang DW, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the

comprehensive functional analysis of large gene lists. Nucleic Acids Res. 2008;37:1–13.

1213

1214

 Figure titles and legends

 Figure 1. Overview of Quartz-Seq2 experimental processes

 Quartz-Seq2 consisted of five steps. (1) Each single cell in a droplet was sorted into lysis buffer

 in each well of a 384-well PCR plate using flow cytometry analysis data. (2) Poly-adenylated

 RNA in each well was reverse-transcribed into first-strand cDNA with reverse transcription

 primer, which had a unique cell barcode. We prepared 384 or 1,536 kinds of cell barcode (CB)

 with a unique sequence based on the Sequence–Levenshtein distance (SeqLv). The edit

 distance of SeqLv was 5. RT primer also had a UMI sequence for reduction of PCR bias (MB)

 and a poly-dT sequence for binding to poly-A RNA. (3) Cell barcode-labeled cDNAs from all 384

 wells were promptly collected by centrifugation using assembled collectors. (4) Collected

 first-strand cDNAs were purified and concentrated for subsequent whole-transcript amplification.

 In the poly-A tailing step, purified cDNA was extended with a poly-A tail by terminal

 deoxynucleotidyl transferase (TdT). Subsequently, second-strand cDNA was synthesized with a

 tagging primer, which had a poly-dT sequence. The resulting second-strand cDNA had a PCR

 primer sequence (M) at both ends of the cDNA. The cDNA was amplifiable at subsequent PCR

 amplification. (5) For conversion from amplified cDNA to sequence library DNA, we fragmented

 the amplified cDNA using the ultrasonicator Covaris. Such fragmented cDNA was ligated with a

 truncated Y-shaped sequence adaptor, which had an Illumina flow-cell binding sequence (P7)

 and a pool barcode sequence (PB). PB made it possible to mix different sets of

 cell-barcode-labeled cDNA. Ligated cDNA, which had CB and MB sequences, was enriched by

 PCR amplification. The resulting sequence library DNA contained P7 and P5 flow-cell binding

 sequences at respective ends of the DNA. We sequenced the cell-barcode site and the UMI site

 at Read1, the pool-barcode site at Index1, and the transcript sequence at Read2.

 Figure 2. Sequence performance of Quartz-Seq2 with molecular biological improvements

 a) Improvement of poly-A tagging efficiency. Barplot represents the relative DNA yield in various

 poly-A tagging conditions using purified first-strand cDNA from 1 ng of total RNA. T55 buffer as

 the terminal deoxynucleotidyl transferase (TdT) buffer and the temperature condition "Increment"

for the poly-A tagging step improved the cDNA yield of whole-transcript amplification. Buffer

compositions were indicated in Additional file 4: Table S3. QuartzB represents use of a

Quartz-Seq-like buffer as a positive control, in accordance with the approach described in the

original Quartz-Seq paper. Finally, we quantified cDNA yield (300–9,000 bp) and byproduct DNA

yield (50–300 bp) using Bioanalyzer (Agilent). (b) The improvement of cDNA yield with the

combination of T55 buffer and "Increment" (T55+Inc) was confirmed by qPCR assay. We

detected eight genes (*Eef1b2*, *Nanog*, *Pou5f1*, *Sox2*, *Utf1*, *Spp1*, *Ywhae*, *Tbp*) with amplified

cDNA and nonamplified cDNA. We prepared nonamplified cDNA with 200 pg of total RNA and

random hexamer. SCC stands for Spearman's rank correlation coefficient. c) Comparison

between Quartz-Seq2 and Quartz-Seq-like conditions regarding sequence performance. We

analyzed 384 wells with 10 pg of total RNA and used approximately 0.19 M fastq reads per well

on average. We show the UMI count and gene count in barplots. We also show a diagrammatic

depiction of the relationship between the cumulative number of detected genes and the gene

expression level. d) Gene expression reproducibility between bulk poly-A-RNA-seq (1 μg of total

RNA) and Quartz-Seq2 (10 pg of total RNA, averaged over 384 wells). e) The panels show a

scatter plot between the mean of gene expression and the variability of gene expression with 10

pg of total RNA in 384 wells. Red lines represent the theoretical variability of gene expression in

the form of a Poisson distribution. The left and right panels represent the mean-CV plots before

and after UMI filtering. The term "RT100" represents the condition of enzyme concentration in

reverse transcription for Quartz-Seq2 (see Material and Methods).

1264

Figure 3. Cost reduction of Quartz-Seq2 upon improvement in quantitative performance

a) Reverse-transcription step has a highest share in single-cell experimental part of Quartz-Seq2.

We present the different proportions of costs associated with each of the experimental steps.

The reverse-transcription step accounts for approximately 65% of the total cost in the "RT100"

condition. We thus established the "RT25" condition by reducing the enzyme concentration to

25% in reverse transcription (for details, see Material and Methods). b) Barplots present the

amount of amplified cDNA from three technical replicates of a 384-well plate with 10 pg of total

RNA in all wells for the "RT100" and "RT25" conditions. The y-axis represents the yield of

amplified cDNA. The cDNA yields from 10 pg of total RNA were 55.9 ± 5.5 ng for "RT100" and

65.7 ± 2.5 ng for "RT25". c) We analyzed amplified cDNA from 10 pg of total RNA by

Quartz-Seq2 with various initial data sizes (approximately 0.27 M, 0.19 M, 0.171 M, 0.152 M,

0.133 M, 0.114 M, 0.095 M, 0.076 M, 0.057 M, 0.038 M and 0.019 M fastq reads per 10 pg of

total RNA on average) from NextSeq500. The Read2 length was 62 nt. In all panels, the x-axis

represents the initial data size for one sample. We present the UMI count, gene count, and spike

capture ratio in the upper panels. We also show the technical variability of the UMI count or gene

count between wells in the lower panels. The y-axis presents the coefficient of variation of the

UMI count or the gene count. d) Relationship between initial data size and the number of single

cells for sequence analysis in NextSeq500 runs. Typically, one sequence run with NextSeq

500/550 High Output v2 Kit reads out 400–450 M fastq reads. The x-axis represents the input

cell number for one sequence run. The y-axis represents the initial data size (fastq reads) per

cell on average. If we analyzed 4,000 single cells in one sequence run, approximately 0.1 M

fastq reads on average would be assigned to one single cell. Red circle represents the typical

range of shallow input read depth for a single cell. e) Capture ratio of ERCC spike mix I in the

"RT25" condition. The capture ratio of ERCC spike-in mix I was 16.38±0.66% (n=3) at

approximately 0.27 M fastq reads on average per well. The x-axis represents the copy number of

input spike RNAs. The y-axis represents the averaged UMI count for ERCC spike RNAs.

1291

Figure 4. Quartz-Seq2 with high UMI conversion efficiency identified more marker genes

and biological pathways

a) We defined the UMI counts divided by the raw fastq reads as the UMI conversion efficiency. b)

UMI counts, gene counts, and UMI conversion efficiency for Quartz-Seq2 and Drop-seq2

experiments. These values depended on the initial fastq reads. c) Venn diagram of genes that

were differentially expressed between the ES cluster and the Dex-treated ES (PrE) cluster, as

identified by Quartz-Seq2, Drop-seq, and bulk RNA-seq (left). The number of genes that differed

in expression levels between ES and PrE cells by at least twofold was counted (FDR < 0.05). We

also present a Venn diagram of the Reactome pathway (right). d) Venn diagram of GO terms for

genes with highly variable expression, as identified by Quartz-Seq2 and Drop-seq.

1302

Figure 5. Quartz-Seq2 analysis of 4,484 cells from mouse embryonic stem cells and

differentiated cells

a) We successfully analyzed 97.3% of 4,608 wells. The procedures for cell suspension as used

in this assay are shown. Cells cultured under ES-maintenance and Dex-treatment conditions

were separately dissociated into single cells, stained with Hoechst 33342 and/or Calcein-AM,

and mixed evenly. Calcein-AM-positive and -negative cells were sorted to 384-well plates in a

checkered pattern. White scale bars represent 100 μm. b) Clustering of 4,484 single cells

according to the transcriptome. Plotting of cells on t-SNE space with color labeling for each

cluster. The percentage indicates the proportion of cells for each cluster relative to all cells

analyzed. Numbers in parentheses indicate the numbers of cells making up the cluster. c)

Marker genes for each cluster identified by Quart-Seq2. Cluster-specific or cluster-enriched

genes were calculated for each cluster, and their expression is displayed as a color in a heatmap.

No more than 50 cells are shown for simplicity. d) Reconstructed distribution of Calcein-AM

intensity for each cluster. The x-axis represents the intensity of Calcein-AM dye staining. e)

Reconstructed distribution of Hoechst 33342 intensity for each cluster. The y-axis represents the

density of cells. The x-axis represents the intensity of Hoechst 33342 dye staining.

1319

Figure 6 Quartz-Seq2 analysis on two types of cell state: cell cycle-dependent and

-independent cellular heterogeneity

a) Plotting cells on t-SNE space with the colors representing the intensity of Hoechst 33342

staining. b) Dependency of expression of several genes on the intensity of Hoechst 33342

staining [G2/M phase: *Ccnb1*; G1 phase: *Ccne1*; housekeeping gene: *Gnb2l1*; upper panels: ES

main population (cluster 1); lower panels: PrE main population (cluster 4)]. c) Identification of cell

cycle-associated genes. Average gene expression of cells in a bin is colored in the heatmap.

Bins were arranged in order of Hoechst 33342 staining intensity. d) The number of cell

cycle-associated genes identified for each z-threshold. e) Several examples for genes whose

expression was variable, but not associated with the phase of the cell cycle. f) Single-cell qPCR

detection of genes with variable expression.

1331

Figure 7 Quartz-Seq2 analysis of stromal vascular fraction (SVF) from mouse adipose

tissue

a) Morphology of SVF cells. Adipose tissue from a cell suspension of SVF was prepared. Upper

panels present a photograph of adipose tissues and dissociated SVF samples. Yellow scale bar

represents 1 cm. White scale bar represents 10 μm. Lower panels represent the distribution of

cell size information with different platforms (left: diameter of cell size using photography, right:

flow cytometry information using a cell sorter). The diameter of cell size for SVF samples was

6.43 ± 1.35 μm (n=200). b) Clustering of cells included in SVF. The transcriptome of

approximately 1,000 cells was quantified by Quartz-Seq2 and clustering on t-SNE space was

performed. In accordance with the genes and functional terms enriched in each cluster, the cell

type was annotated. The percentage indicates the proportion of cells for each cluster relative to

all cells analyzed. Numbers in parentheses indicate the numbers of cells constituting the cluster.

c) Marker genes for each cluster were identified by Quart-Seq2. Cluster-specific or

cluster-enriched genes were calculated for each cluster, with their expression being displayed as

color in a heatmap. No more than 50 cells are shown for simplicity. d) The results of GO-PCA

analysis. Functional terms enriched in the genes with high factor loadings of PCA were

calculated and the enrichment is displayed as color in the heatmap. No more than 50 cells are

shown for simplicity. e) Reactome pathway with genes differentially expressed between cluster 1

and cluster 8.

1. Single-cell collection (Cell sorter)

2. Cell barcoding

3. Spindown collection

4. Whole transcript amplification based on poly-A tagging

5. Sequence library preparation

a)

ES cells → Dissociation → Hoechst33342 Calcein AM → Mixed cells evenly → Cell-sorter (MoFlo) → Quartz-Seq2 (1,536 indexes) (384 plate x4) x3 sets

+Dex 72 hours

Treated cells (PrE cells) → Dissociation → Hoechst33342 → Mixed cells evenly

Checkered pattern
CalceinAM (+ ●, - ○)

b)

PrE (main)   ES (*Stmn2* high)

97.3% (4,484cells/4,608wells)

- ES (main), 44.6% (2,000)
- ES (*Stmn2 high*), 3.3% (148)
- PrE (main), 49.6% (2,225)
- *Zscan4c/d* high, 0.17% (8)

- unclassified, 0.33% (15)
- ES/PrE doublets, 1.38% (62)
- ES (low-quality cell), 0.57% (26)

Zscan4c/d high

ES (main)

c)

ES (main) ES (*Stmn2* high) PrE (main) *Zscan4c/d* high

Class A

Class B

Class C

Genes

Class D

d)

Mixture of cell-types | ES (main), 2,000 cells | ES (*Stmn2 high*), 148 cells | PrE (main), 2,225 cells | *Zscan4c/d* high, 8 cells

Calcein AM

e)

Mixture of cell-types | ES (main), 2,000 cells | ES (*Stmn2 high*), 148 cells | PrE (main), 2,225 cells | *Zscan4c/d* high, 8 cells

Hoechst33342

a)

Diameter of cell (um)

Density (Cell)

488-SSC-Height

488-FSC1-Height

L

S

b)

Dim2

Dim1

91.31% (1,052cell/1,152well)

unclassified, 1.33% (14)
1: *Cd34+/Efemp1+* mesenchymal stem cell, 7.79% (82)
2: Macrophage, 5.51% (58)
3: *Cd4+* T-cell, 44.96% (473)
4: *Cd4+* T-cell (proliferation), 0.95% (10)
5: B-cell, 15.87% (167)
6: Killer T-cell, 11.02% (116)
7: Natural killer cell, 3.04% (32)
8: *Cd34+/Col15a1+* mesenchymal stem cell, 6.36% (67)
9: *Cxcr6+* T-cell, 1.71% (18)
10: Dendritic cell, 0.85% (9)
11: Plasmacytoide dendritic cell, 0.57% (6)

c)

Cell-clusters

Genes

Class A
Class B
Class C
Class D
Class E
Class F
Class G
Class H
Class I
Class J
Class K

d)

Cell-clusters

GO terms

EMT regulation
Organ formation
Embryonic pattern specification
Collagen trimer
Osteoblast differentiation
Cell-cell junction
Angiogenesis
Mesenchymal cell differentiation
Ribosome biogenesis
T cell receptor signal. pathway
MHC class I complex
Response to IFN-gamma
snoRNA binding
Lamellipodium organization
Dendritic cell differentiation
Mast cell activation
Chemokine production
VEGF pathway
Antigen processing and presentation
Mitochondrial repiratory chain
Cell division
DNA replication

e)

-Log(P-value)

[Reactome pathways]

[Differentially expressed genes]

Cluster 1: *Cd34+/Efemp1+*

Reactome pathway	Genes
Extracellular matrix organization	*Bmp2,Col14a1,Fbn1,Fgf2,Fn1,Itgb7,Mfap5,Ntn4,Pcolce2,Efemp1*
Molecules associated with elastic fibres	*Bmp2,Fbn1,Fn1,Mfap5,Efemp1*
Elastic fibre formation	*Bmp2,Fbn1,Fn1,Mfap5,Efemp1*
IGF transport	*Igfbp4,Igfbp5,Igfbp6*
Non-integrin membrane-ECM interactions	*Fgf2,Fn1,Ntn4*

-Log(P-value)

[Reactome pathways]

[Differentially expressed genes]

Cluster 8: *Cd34+/Col15a1+*

Reactome pathway	Genes
Extracellular matrix organization	*Bgn,Serpinh1,Col15a1,Col3a1,Col4a1,Col4a2,Col4a4,Col5a1,Col5a2,Col6a1,Col6a2, Col1a1,Col1a2,Ctsk,Lama2,Lamb1,Lox,Lum,Mmp2,Nid2,Pcolce,Sparc,Jam2*
Collagen formation	*Serpinh1,Col15a1,Col3a1,Col4a1,Col4a2,Col4a4,Col5a1,Col5a2,Col6a1,Col6a2, Col1a1,Col1a2,Lox,Pcolce*
Collagen degradation	*Col15a1,Col3a1,Col4a1,Col4a2,Col4a4,Col5a1,Col5a2,Col6a1,Col6a2,Col1a1,Col1a2, Ctsk,Mmp2*
Collagen biosynthesis and modifying enzymes	*Serpinh1,Col15a1,Col3a1,Col4a1,Col4a2,Col4a4,Col5a1,Col5a2,Col6a1,Col6a2,Col1a1, Col1a2,Pcolce*
Collagen chain trimerization	*Col15a1,Col3a1,Col4a1,Col4a2,Col4a4,Col5a1,Col5a2,Col6a1,Col6a2,Col1a1,Col1a2*
Assembly of collagen fibrils and other multimeric structures	*Col3a1,Col4a1,Col4a2,Col4a4,Col5a1,Col5a2,Col6a1,Col6a2,Col1a1,Col1a2,Lox*
Degradation of the extracellular matrix	*Col15a1,Col3a1,Col4a1,Col4a2,Col4a4,Col5a1,Col5a2,Col6a1,Col6a2,Col1a1,Col1a2, Ctsk,Lamb1,Mmp2*
Integrin cell surface interactions	*Col3a1,Col4a1,Col4a2,Col4a4,Col5a1,Col5a2,Col6a1,Col6a2,Col1a1,Col1a2,Lum,Jam2*
Non-integrin membrane-ECM interactions	*Col3a1,Col4a1,Col4a2,Col4a4,Col5a1,Col5a2,Col1a1,Col1a2,Lama2,Lamb1*
ECM proteoglycans	*Bgn,Col3a1,Col5a1,Col5a2,Col6a1,Col6a2,Col1a1,Col1a2,Sparc*