Improving the accuracy of two-sample summary data Mendelian randomization: moving beyond the NOME assumption

Jack Bowden[1,2*], Fabiola Del Greco M[3], Cosetta Minelli[4], Qingyuan Zhao[5], Debbie A Lawlor[1,2], Nuala A Sheehan[6], John Thompson[6] & George Davey Smith[1,2].

[1] *MRC Integrative Epidemiology Unit at the University of Bristol, U.K.*
[2] *Population Health Sciences, University of Bristol, U.K.*
[3] *Institute for Biomedicine, Eurac Research, Bolzano, Italy*
[4] *Population Health and Occupational Disease, NHLI, Imperial College, London, U.K.*
[5] *Department of Statistics, The Wharton School, University of Pennsylvania, U.S.A.*
[6] *Department of Health Sciences, University of Leicester, Leicester, U.K.*

[*]Address for correspondence:
Jack Bowden
MRC Integrative Epidemiology Unit
Oakfield House, Bristol, BS8 2BN, U.K
jack.bowden@bristol.ac.uk

Abstract

Background: Two-sample summary data Mendelian randomization (MR) incorporating multiple genetic variants within a meta-analysis framework is a popular technique for assessing causality in epidemiology. If all genetic variants satisfy the instrumental variable (IV) and necessary modelling assumptions, then their individual ratio estimates of causal effect should be homogeneous. Observed heterogeneity signals that one or more of these assumptions could have been violated.

Methods: Causal estimation and heterogeneity assessment in MR requires an approximation for the variance, or equivalently the inverse-variance weight, of each ratio estimate. We show that the most popular '1st order' weights can lead to an inflation in the chances of detecting heterogeneity when in fact it is not present. Conversely, ostensibly more accurate '2nd order' weights can dramatically increase the chances of failing to detect heterogeneity, when it is truly present. We derive modified weights to mitigate both of these adverse effects.

Results: Using Monte Carlo simulations, we show that the modified weights outperform 1st and 2nd order weights in terms of heterogeneity quantification and causal estimation. The added benefit is most noticeable when the genetic instruments are weak, the causal effect is large and the number of instruments is large. We illustrate the utility of the new method using data from a recent two-sample summary data MR analysis to assess the causal role of systolic blood pressure on coronary heart disease risk.

Conclusions: We propose the use of modified weights within two-sample summary data MR studies for model fitting, quantifying heterogeneity and outlier detection.

Key words: Two-sample summary data Mendelian randomization; Inverse-variance weighted estimate; Cochran's Q statistic; Outlier detection.

Total words in main body of paper: 4,210.

Introduction

Mendelian randomization (MR) [1] is an instrumental variable approach that uses genetic data, typically in the form of single nucleotide polymorphisms (SNPs), to assess whether a modifiable exposure exerts a causal effect on a health outcome in the presence of unmeasured confounding. A particular MR study design gaining in popularity instead combines publically available summary data on SNP-exposure and SNP-outcome associations from two separate studies for large numbers of uncorrelated variants within the framework of a meta-analysis. These studies should contain no overlapping individuals (to ensure independence) but should also originate from the same source population. This is referred to as two-sample summary data MR [2]. Providing the necessary modelling assumptions are met and the chosen set of SNPs are all valid instrumental variables, an inverse-variance weighted (IVW) average of their individual causal ratio estimates provides an efficient and consistent estimate for the causal effect. This is referred to as the IVW estimate (see **Box 1**). Cochran's Q statistic, which is derived from the IVW estimate, should follow a χ^2 distribution with degrees of freedom equal to the number of SNPs minus 1. Excessive heterogeneity is an indication that either the modelling assumptions have been violated, or that some of the genetic variants violate the IV assumptions, by exerting a direct effect on the outcome not through the exposure [3]. This is termed 'horizontal pleiotropy' [4, 5]. For brevity we will refer to problematic horizontal pleiotropy simply as pleiotropy from now on.

The presence of heterogeneity due to pleiotropy does not necessarily invalidate an MR study. If across all variants: (i) the amount of pleiotropy is independent of instrument strength (the InSIDE assumption [6]) and (ii) it has a zero mean, then a standard random effects meta-analysis will still yield reliable inferences [6, 7]. Although many MR methods now exist which offer robustness to pleiotropy, in this paper we focus solely on the standard IVW estimate.

Choice of weights in two-sample summary data MR

Typically, '1st order' inverse-variance weights are used to calculate both the IVW estimate and Cochran's Q. 1st order weights ignore uncertainty in the denominator of the ratio estimate, which is equivalent to making the 'NO Measurement Error' (NOME) assumption, as defined in [7, 8]. This nomenclature is chosen to remind the practitioner that the SNP-exposure association estimates are only equal to the true associations when measured with infinite precision (or without error). The NOME assumption does not relate to absence of measurement error in the exposure itself, which can also be problematic for MR studies [9]. Although the NOME assumption is never completely satisfied, strong violation (via the use of weak genetic instruments) induces classical regression dilution bias in the IVW estimate towards the null. So called '2nd order' weights, attempt to better acknowledge the full uncertainty in the ratio estimate of causal effect from each SNP [10, 11] (see **Box 1**). It may appear obvious that 2nd order weights should used as standard within an MR study to cal-

culate the IVW estimate and Cochran's Q. In fact, Thompson et al. [12] showed that 2nd order weighting produces causal estimates which are generally more biased than 1st order weighting. The ability of 1st and 2nd order weighting to furnish reliable Q statistics has yet to be fully explored.

Methods

It is possible to view Cochran's Q statistic not just as a method for quantifying heterogeneity, but as a tool for directly estimating the causal effect. That is, the IVW estimate actually minimises Cochran's Q. It can therefore be viewed as an estimating equation for the causal effect. We use this fact to derive a generalised estimating equation based on an extended version of Cochran's Q statistic (see **Box 2**), where its weight term is allowed to be a function of the causal effect parameter. We show that 1st order and 2nd order weighting are special cases of this general weight function. Using this formulation we propose two new procedures for causal effect estimation and heterogeneity quantification.

Our first procedure is termed the 'iterative' approach. It iteratively updates the weight term with improved guesses for the causal parameter, using the 1st order IVW estimate as a starting point. This procedure is closely related to the 'two-step GMM' estimator [13] used in econometrics. Our contribution has been to describe how it can be implemented using Cochran's Q statistic in the two-sample summary data MR setting. It will be shown that the iterative IVW approach improves causal effect estimation and heterogeneity detection compared to 1st and 2nd order weighting. However, regardless of the number of iterations performed, this procedure will not in general yield the same results as that obtained from directly minimising a more general version of Cochran's Q, where in addition the weight term is allowed to be a proper function of the causal effect parameter β. We refer to this second procedure as the 'exact' approach. The exact IVW estimate can be viewed as analogous to the limited information maximum likelihood (LIML) estimate, translated to the two-sample summary data MR setting [14]. For further details see **Box 2**.

Estimation and inference after detection of pleiotropy

The Q statistics can be used to calculate the IVW estimate under a fixed effect (or no pleiotropy) model and to test for the presence of heterogeneity (see **Box 2**). If however, the Q statistics derived using either the 1st order, 2nd order, iterative or exact weighting correctly detect substantial heterogeneity, inferences about the causal effect need to be adjusted to take this additional uncertainty into account, by assuming an additive or multiplicative random effects model [15, 16]. In Appendix 1 of *Online Supplementary Material* we describe in detail how to generalise the Q statistics in **Box 2** to obtain point estimates, standard errors and confidence intervals for the 1st order, 2nd order, iterative and exact IVW estimate under both fixed and multiplicative random effects models (the multiplicative model is currently preferred

for MR studies). This task is straightforward for the 1st order, 2nd order and iterative weighting approaches because they can be fitted using standard regression software. Bespoke methods are needed for exact weighting, however. Specifically, in the fixed effect case we describe how to invert the exact Q statistic to get a 95% confidence interval for the exact weighted IVW estimate. In the random effect case, we describe how to jointly estimate the causal effect and multiplicative over-dispersion parameter using a system of two estimating equations. A non-parametric bootstrap algorithm is then proposed to obtain a confidence interval for the causal effect.

Performance of the Q statistics under no pleiotropy

We now assess the extent to which Q statistics derived using 1st order, 2nd order, iterative and exact weighting erroneously detect heterogeneity due pleiotropy when it is not present (i.e. its type I error rate). To assess this, two-sample summary data MR studies comprising 25 SNP-exposure and SNP outcome association estimates were generated from models with no heterogeneity due to pleiotropy. This furnished a set of ratio estimates between which no additional variation should exist as their instrument strength grows large (because NOME is satisfied), or if the causal effect (β) equals zero. To highlight this we simulated MR studies with a range of instrument strengths - from weak (a mean F-statistic of 10) to strong (a mean F-statistic of 100). Further details of the simulation study set up are described in Appendix 2 of *Online Supplementary Material*.

Table 1 (columns 2-9) show the mean Q statistic and the probability of the Q statistic detecting heterogeneity at the 5% significance level (the type I error rate), when using 1st order, 2nd order, iterative and exact weights. Five different mean F-statistic values were considered for β=0 (no causal effect), β=0.05 and β=0.1, giving 15 scenarios in total. Four iterations were used for the iterative weighting method, as this was sufficient to ensure convergence. We note that in the absence of a causal effect (β=0), 1st order weights are exactly correct. Furthermore, in the presence of a causal effect, when the mean F-statistic is 100 all weighting methods are near-exact. Under the causal null, all weighting schemes control the type I error rate for detecting heterogeneity. 2nd order weighting is extremely conservative in this respect with weak instruments, however (e.g. a type I error rate near zero when F=10).

In the presence of a causal effect, 1st order weights under-estimate the true variability amongst the ratio estimates as the mean F-statistic reduces. The associated Q statistics are then too large on average (i.e. positively biased beyond their expected value of 24). This inflates the type I error rate for detecting pleiotropy beyond nominal levels (e.g. a type I error rate of $\approx$ 80% when F=10 and β=0.1). 2nd order weighting continues to over-correct the Q statistic so that it is negatively biased, thereby removing *any* ability to detect heterogeneity at all. In contrast, iterative weights are much more effective at preserving the type I error rate of the Q statistic at its nominal level, unless the mean F-statistic is very low (indicating weak instruments). Exact weighting perfectly controls the type I error rate of Cochran's Q across all the

scenarios considered. Appendix 2 of *Online Supplementary Material* shows equivalent results for MR studies of 10 and 100 variants, with highly similar results.

Figure 1 (left and right) shows the distribution of Q statistics using 1st order, 2nd order and exact weights for β=0.1 and when the mean F-statistic is 100 and 10. This illustrates how exact weighting ensures Cochran's Q statistic is faithful to its correct null distribution.

Power to detect pleiotropy

In Table 1 the type I error rate of Cochran's Q statistic for detecting heterogeneity using 2nd order weights was below its nominal level. This is detrimental if it translates into a low statistical power to detect heterogeneity when it *is* truly present. Figure 2 (left) shows the power of Cochran's Q to detect heterogeneity at the 5% significance level as a function of 1st order, 2nd order, iterative and exact weights when data are simulated under a multiplicative random effects model with heterogeneity due to pleiotropy of increasing magnitude (specifically, equation (2) in Appendix 1 of *Online Supplementary Material* was used).

The simulation is repeated for MR analyses with 10, 25 and 100 SNPs. For all simulations, the causal effect equalled 0.05 and the mean F-statistic equalled 61. We see that the power of Cochran's Q to detect heterogeneity approaches 100% for all weighting schemes as the pleiotropy variance increases. Power also increases with the number of SNPs. The power of iterative or exact weights is near identical, therefore we only show results for the exact weights for clarity. The most striking result in this plot is that the power of 2nd order weighting always lags considerably behind that of 1st order or exact weights.

Figure 2 (right) shows the results of a near identical simulation for the case L=25, except that the causal effect is set to 0.1 and the mean F-statistic is equal to 25. We see that the power to detect heterogeneity is always greatest when using 1st order weights, but only because its power curve starts at a baseline level of 28% when there is no pleiotropy. This corresponds to the type I error rate observed in row 14 of Table 1. The power of iterative and exact weighting starts at the correct 5% level, and rapidly increases to 100% as the pleiotropy variance increases. The two implementations of our modified weights can be differentiated in this simulation, with the iterative approach being slightly more powerful. The power of 2nd order weighting, unsurprisingly, lags considerably behind the rest. Equivalent plots for data simulated under an additive pleiotropy model are shown in Appendix 3 of *Online Supplementary Material*, and are highly similar.

Detecting outliers using individual components of Q

When heterogeneity is detected by the IVW model, it is interesting to investigate whether this is contributed to by all SNPs, or if instead a small number of SNPs

are responsible. Under the null hypothesis of no heterogeneity, Q should follow an appropriate χ^2_{L-1} distribution, with L being the number of SNPs. Likewise, each individual component of Q can be approximated by a χ^2_1 distribution. If an individual SNP's Q contribution is extreme (for example above the 5% threshold of 3.84 or instead a bonferroni corrected threshold), then it may be desirable to exclude the SNP in a sensitivity analysis. Although we do not want to advocate a rigid, blanket policy of outlier removal, in Appendix 4 of *Online Supplementary Material* we illustrate via simulation how the reliability of such a procedure depends on the choice of weights. The simulation (with 26 SNPs and a single larger outlier) is motivated by the real data example in the following section. In this instance, our simulation suggests that iterative rather than exact weights are best at correctly identifying outliers due to pleiotropy.

Estimator performance with and without pleiotropy

Table 2 shows the performance of the 1st order, 2nd order, iterative and exact weigthing in providing accurate point estimates, standard errors and confidence intervals for the causal effect under a fixed effect (no heterogeneity) model for MR analyses of 25 variants. For exact weighting we show the empirical coverage using two different methods: A symmetric 95% confidence interval assuming a normality (labelled 'CF$_1$') and a 95% confidence interval obtained from inverting its Q statistic (labelled 'CF$_2$'). Importantly, all methods give reliable unbiased estimates with correct coverages under the causal null hypothesis. In the presence of a non-zero causal effect 1st order and 2nd order IVW estimates are increasingly affected by regression dilution bias (and consequently worsening coverage) as the instrument strength decreases. Iterative weights also produce IVW estimates that suffer from regression dilution bias and sub-optimal coverage, but to a lesser extent than 1st or 2nd order weighting. Exact weighting perfectly removes the effect of regression dilution bias (although the precision of the estimate is reduced) and confidence intervals obtained via the inversion method have the correct coverage. Equivalent results for MR studies with 10 and 100 SNPs are shown in Appendix 5 of *Online Supplementary Material*. When only 10 SNPs are available, they are all weak and the causal effect is large, the coverage of the inverted confidence interval for the exact IVW estimate is slightly conservativie (e.g. 98% instead of 95%). As the number of SNPs increases to 100, coverage is very close to the nominal 95% level.

Table 3 show equivalent results when summary data sets of 25 SNPs are simulated under a multiplicative random effect model allowing for pleiotropy. The data are simulated so that the variability of the ratio estimates is twice that expected in the absence of pleiotropy (i.e the variance inflation parameter $\phi = 2$). The performance of each approach follows a similar pattern to that presented for the fixed effect case in Table 2, with 1st order, 2nd order and iterative weights adversely affected by weak instrument bias and under coverage. The exact IVW estimate and its corresponding variance parameter estimate are approximately unbiased. The non-parametric bootstrap procedure yields confidence intervals with approximately correct coverage.

As before, confidence intervals have a tendency to be slightly conservative when the instruments are weak. Equivalent results for MR studies with 10 and 100 SNPs are shown in Appendix 6 of *Online Supplementary Material*. As the number of SNPs increases, the coverage of the exact IVW estimate's confidence interval is increasingly closer to the nominal level.

Power to detect a causal effect

In Appendix 7 of *Online Supplementary Material*, we show the power of 1st order, 2nd order, iterative and exact weighting to detect a causal effect for MR studies of 10, 25 and 100 SNPs when the data are generated from the same multiplicative random effects model. These simulations highlight the one downside of exact weighting for causal estimation: when there are only a small number of weak instruments, its power can be considerably lower. For example, when $F = 10$ and the causal effect is 0.05 its power is just under half that of the 1st order IVW estimate (29% vs 13%). However, the power difference reduces considerably for 25 SNPs (e.g 60% vs 40%) and is effectively zero for 100 SNPs. The power of iterative weighting is much more similar to that of 1st order weighting, but always slightly lower.

Applied example

Figure 3 (top) shows a scatter plot of summary data estimates for the associations of 26 genetic variants with systolic blood pressure (SBP, the exposure) and coronary heart disease (CHD, the outcome). SNP-exposure association estimates were obtained from the International Consortium for Blood Pressure consortium (ICBP) [17]. SNP-CHD association odds ratios were collected from Coronary ARtery Disease Genome-Wide Replication And Meta-Analysis (CARDIoGRAM) consortium [18], which are plotted (and subsequently modelled) on the log odds ratio scale by making a normal approximation. These data have previously been used in a two-sample summary data MR analysis by Ference et al. [19] and Lawlor et al. [20], but we extend their original analysis here by applying our modified weights and conducting a more in depth inspection of each variant's contribution to the overall heterogeneity. The mean F-statistic for these data is 61. Using 1st order weights the IVW estimate, which represents the causal effect of a 1mmHg increase in SBP on the log-odds ratio of CHD, is 0.053. This is shown as the slope of a solid black line passing through the origin (note: the origin is not visible because of a truncated x-axis). Cochran's Q statistic based on 1st order weights is equal to 67.1, indicating the presence of substantial heterogeneity.

Table 4 shows the results of further IVW analyses using all weighting schemes. All schemes detect significant heterogeneity. As expected, the observed heterogeneity is largest when using 1st order weights, smallest when using 2nd order weights, and in between the two when using modified weights. Point estimates and standard errors are in good agreement across the different weights, because the mean instrument strength is high. Exact weighting gives the largest point estimate of 0.057 under a

fixed effect model and 0.054 under a random effects model. This is followed by 1st order and then 2nd order weights respectively. This ordering is as expected, given their relative susceptibility to regression dilution bias. Although the large amount of heteroeneity in these data make a fixed effect model innapropriate, we report both fixed and multiplicative random effects estimates under exact weighting to highlight that their point estimates are truly distinct (they are identical for all other weighting schemes).

For comparison, we also report the Weighted Median [21], $\hat{\beta}_{WM}$, that can identify the causal effect when up to (but not including) half of the information in the analysis stems from genetic variants that are invalid IVs. Its estimate, which is calculated using 1st order weights, is 0.063. Although all approaches provide strong evidence in favour of a non-zero causal effect, the exact random effects IVW estimate is the least precise of all estimates. Consequently its p-value for testing the causal null hypothesis is the largest of all.

Figure 3 (bottom-left) shows the individual contribution to Cochran's Q statistic under each weighting scheme. Horizontal lines have been drawn to indicate the location of the 5th, 1st and 0.19th percentile of a χ^2_1 in order to help assess the magnitude of the contributions. The 0.19th percentile is derived as a 0.05 threshold adjusted for multiple testing using the Bonferroni correction. We see that the eighth SNP in our list (rs17249754) is responsible for the vast majority of the excess heterogeneity. Its contribution, Q_8, ranges from approximately 24.5 to 28 depending on weighting. Variant rs17249754 sits in the ATPase plasma membrane Ca2+ transporting 1 (*ATP2B1*) gene, which is involved in intracellular calcium homeostasis, and is strongly associated with higher SBP. However, in the CARDIoGRAM consortium it is associated with reduced CHD.

Since rs17249754 is also a strong instrument, and is potentially pleiotropic, its presence in the data could lead to the InSIDE assumption being violated. We therefore opt to remove it in a further sensitivity analysis, and Table 4 show the results. All IVW estimates increase by around 20% (lying between 0.063 and 0.067), but are ordered as before. Removal of rs17249754 leads to a dramatic reduction in the amount of heterogeneity present in the data, as referenced by Q statistics between 30 and 35 for all methods. Figure 3 (bottom-right) shows the updated contributions of each SNP to the various Q statistics after removing rs17249754. If only 1st order weighting were available, it might be tempting to exclude further variants from the analysis, but this signal is appropriately tempered when using exact weights. The Weighted Median estimate without rs17249754 is 0.065 (compared to 0.063 with). This highlights its inherent robustness to outliers, which is a major strength.

Discussion

In this paper we have demonstrated the limitations of 1st and 2nd order weighting when used for IVW analysis in two-sample summary data Mendelian randomization. Most importantly, we highlight the potential for serious type I error inflation of Cochran's Q statistic when using standard 1st order weights with weak instruments. In recent work, Verbank et al. [22] also noted this same tendency and proposed a simulation-based alternative to 1st order weighting named 'MR-PRESSO'. Our simulations show that our modified weights can deliver much more accurate causal estimates and reliable tests for heterogeneity than either 1st or 2nd order weighting, offer a simple alternative to MR-PRESSO.

Modified weights were also shown to be a more reliable tool for the detection and removal of outliers in a given data set, as apposed to 1st order weights (which may detect too many outliers) and 2nd order weights (that may detect too few). Our simulations suggest that the exact weights should be used when testing for the overall presence of heterogeneity (referred to as the 'global' test by Verbank et al. [22]) but that iterative weights are preferable if looking at the individual outliers. We suspect this is because exact weighting makes a more aggressive correction for regression dilution bias than iterative weighting. Its resulting estimate then makes more variants appear as outliers, because their ratio estimates are further away from the corrected slope. In effect, exact weighting leads to the detection of SNPs that are weak or pleiotropic.

An exciting finding of this paper is that the exact weighting yields causal estimates that are remarkably robust to regression dilution bias. This opens up the potential for the significance threshold used to select SNPs as instruments to be substantially dropped. However, there are two downsides to the use of exact weights. Firstly, it can produce causal estimates with significantly reduced precision compared to simple 1st order weighting when there are only a small number of instruments. Secondly, if weak instruments are used, then SNP-exposure estimates are more susceptible to the 'winner's curse' than strong instruments. In preliminary work conducted in tandem with this paper, Zhao et al. [14] investigate the use of exact weighting for causal estimation and attempt to address both these issues. Specifically, they incorporate a penalized weight function within the exact weights. This reduces the effect of outliers (as apposed to explicit outlier removal) and increases the precision of the causal estimate. Sampling splitting is proposed to remove the affect of winner's curse. The methods laid out in this paper differ from that of Zhao et al. [14] in four important ways. Firstly, we focus on the case of a multiplicative random effects pleiotropy commonly used in summary data MR whereas Zhao et al assume an additive random effects model. Secondly, Zhao et al derive and implement their method using profile-likelihood theory, whereas as our approach is motivated and implemented using Cochran's Q statistic. Thirdly, we propose two forms of modified weighting (iterative and exact). Fourthly, we describe how both iterative and exact weighting can be used to test for heterogeneity as well as for causal estimation. For further details see

Appendix 1 *Online Supplementary Material.*

Limitations

Our conclusions regarding the use of modified weights are limited to the two-sample summary setting where SNP-outcome and SNP-exposure associations are estimated in independent but homogeneous samples. Further research would be required to decide if modified weights should be used in MR analyses of summary data estimates when there is partial overlap between samples, or in the single sample (total overlap) setting.

When Cochran's Q statistic detects significant amounts of heterogeneity, it is prudent to test whether it is meaningfully biasing the analysis. This would indeed be the case if the heterogeneity were caused in part by directional pleiotropy with a non-zero mean. This would lead to bias in the IVW estimte, unless of course it was caused by a small number of SNPs that could be identified and removed from the analysis. MR-Egger regression [6, 7] could instead be used to address this. This approach simply regresses SNP-outcome associations on the SNP-exposure associations, tests for bias via its intercept, and estimates a bias-adjusted causal effect via its slope. Observed heterogeneity around the MR-Egger fit can then be quantified using an extended version of Cochran's Q statistic, Rücker's Q' [7, 23], and each variant's contribution to Q' can be used as the basis for outlier detection. Currently MR-Egger and Rücker's Q' statistic use 1st order weights. Preliminary work suggests that modified weighting can be applied to MR-Egger regression to improve its performance - both in terms of causal effect estimation and heterogeneity quantification - just as for an IVW analysis, but further development and validation of this method is required.

Software to implement all of the methods introduced in this paper can be found within the `RadialMR` package to perform two-sample summary data MR, which can be downloaded from `https://github.com/WSpiller/RadialMR`.

> **Key messages:**
>
> - Two-sample summary data Mendelian randomization requires the specification of inverse-variance weights for model fitting, heterogeneity quantification and outlier detection amongst a set of causal estimates.
>
> - Heterogeneity indicates a possible violation of the necessary IV or modelling assumptions of which pleiotropy is a likely major cause.
>
> - 1st order weights can inflate the type I error rate of Cochran's Q statistic for detecting heterogeneity about the IVW estimate when the NOME assumption is strongly violated (as judged by a low F-statistic), and the true causal effect of interest is non-zero.
>
> - 2nd order weights can reduce the power of Cochran's Q statistic for detecting heterogeneity about the IVW estimate when the NOME assumption is violated.
>
> - Modified weights (developed in this paper) preserve the type I error rate of Cochran's Q statistic, whilst maintaining its statistical power, and naturally correct for regression dilution bias due to NOME violation.
>
> - 'Exact' weights should be used for global tests of heterogeneity. 'Iterative' weights should be used to assess the outlier status of individual SNPs.
>
> - Exact IVW estimates can be imprecise relative to other weighting schemes with small numbers of instruments, but perform best with large numbers of (potentially weak) instruments.

Acknowledgements

We thank the reviewers and associate editor for providing helpful comments and suggestions that greatly improved this paper. Jack Bowden, Debbie A Lawlor and George Davey Smith all work in a Unit that receives support from the University of Bristol and UK Medical Research Council (MCUU00011/1, MCUU00011/2 and MCUU00011/6). The authors declare no conflict of interest.

References

[1] Davey Smith G, Ebrahim S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *International Journal of Epidemiology* 2003; **32**:1–22.

[2] Burgess S, Butterworth A, Thompson SG. Mendelian randomization analysis with multiple genetic variants using summarized data. *Genetic Epidemiology* 2013; **37**:658–665.

[3] Del Greco M F, Minelli C, Sheehan NA, Thompson JR. Detecting pleiotropy in Mendelian randomisation studies with summary data and a continuous outcome. *Statistics in Medicine* 2015; **34**:2926–2940.

[4] Davey Smith G, Hemani, G. Mendelian randomization: genetic anchors for causal inference in epidemiological studies. *Human molecular genetics* 2014, **23**: 89–98.

[5] Stearns FW. One hundred years of pleiotropy: a retrospective. *Genetics* 2010; **186**:767–773.

[6] Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *International Journal of Epidemiology* 2015; **44**:512–525.

[7] Bowden J, Del Greco M F, Minelli C, Davey Smith G, Sheehan NA, Thompson JR. A framework for the investigation of pleiotropy in two-sample summary data Mendelian randomization *Statistics in Medicine* 2017; **36**: 1783–1802

[8] Bowden J, Del Greco M F, Minelli C, Davey Smith G, Sheehan NA, Thompson JR. Assessing the suitability of summary data for Mendelian randomization analyses using MR-Egger regression: the role of the I^2 statistic. *International Journal of Epidemiology* 2016; **45**: 1961–74.

[9] Pierce B, VanderWeele T. The effect of non-differential measurement error on bias, precision and power in Mendelian randomization studies. *International Journal of Epidemiology.* 2012 **41**:1383–93.

[10] Burgess S, Bowden J. Integrating summarized data from multiple genetic variants in Mendelian randomization: bias and coverage properties of inverse-variance weighted methods Technical report, University of Cambridge 2015. `https://arxiv.org/pdf/1512.04486.pdf`

[11] Thomas D, Lawlor, D, Thompson J. Re: Estimation of Bias in Nongenetic Observational Studies Using Mendelian Triangulation by Bautista et al. *Annals of Epidemiology* 2007; **17**: 511–513.

[12] Thompson J, Minelli C, Del Greco M F. Mendelian randomization using public data from genetic consortia. *International Journal of Biostatistics* 2016; **12**:2

[13] Cumby, RE, Huizinga J, Obstfeld M. Two-step two-stage least squares estimation in models with rational expectations. *Journal of Econometrics* 1982; **21**: 333–355.

[14] Zhao Q, Wang J, Hemani G, Bowden J, Small D. Statistical inference in two-sample summary data Mendelian randomization using a robust adjusted profile score. Technical report, University of Pennsylvania 2018. `http://www-stat.wharton.upenn.edu/~qyzhao/papers/mr_raps.pdf`

[15] DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clinical Trials* 1986; **7**:177–188.

[16] Thompson SG, Sharp S. Explaining heterogeneity in meta-analysis: a comparison of methods. *Statistics in Medicine* 1999; **18**:2693–2708.

[17] International Consortium for Blood Pressure Genome-Wide Association Studies. Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature* 2011; **478**:103-9.

[18] CARDIoGRAMplusC4D. Large-scale association analysis identifies new risk loci for coronary artery disease. *Nature Genetics* 2003; **45**:25–33.

[19] Ference BA, Julius S, Mahajan N, Levy PD, Williams KASr, Flack JM. Clinical effect of naturally random allocation to lower systolic blood pressure beginning before the development of hypertension. *Hypertension* 2014; **63**: 1182–88.

[20] Lawlor DA, Tilling K, Davey Smith G. Triangulation in aetiological epidemiology. *International Journal of Epidemiology* 2016; **45**:1866–86.

[21] Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent estimation in Mendelian randomization with some invalid instruments using a weighted median estimator. *Genetic Epidemiology* 2016; **40**: 304–14.

[22] Verbanck M, Chen CY, Neale B, Do R. Widespread pleiotropy confounds causal relationships between complex traits and diseases inferred from Mendelian randomization. bioRxiv. 2017 `http://www.biorxiv.org/content/biorxiv/early/2017/06/30/157552.full.pdf`.

[23] Rücker G, Schwarzer G, Carpenter J, Binder H, Schumacher M. Treatment-effect estimates adjusted for small study effects via a limit meta-analysis. *Biostatistics* 2011; **12**:122–142.

Box 1: Standard two-sample summary data MR

The IV assumptions: The canonical approach to MR assumes that group of SNPs are valid IVs for the purposes of inferring the causal effect of an exposure, X, on an outcome, Y. That is they are: associated with X (IV1); not associated with any confounders of X and Y (IV2); and can only be associated with Y through X (IV3). The IV assumptions are represented by the solid lines in the causal diagram below for a SNP G_j, with unobserved confounding represented by U. Dotted lines represent dependencies between G and U, and G and Y that are prohibited by the IV assumptions. The causal effect of a unit increase in X on the outcome Y, denoted by β, is the quantity we are aiming to estimate.

The Ratio estimate: Assume that exposure X causally affects outcome Y linearly across all values of X, so that a hypothetical intervention which induced a 1 unit increase in X would induce a β increase in Y. Suppose also that all L SNPs predict the exposure via an additive linear model with no interactions. If SNP j is a valid IV, and the two study samples are homogeneous, then the underlying SNP-outcome association from sample 1, Γ_j, should be a scalar multiple of the underlying SNP-exposure association estimate from sample 2, γ_j, the scalar multiple being the causal effect β. That is:

$$\Gamma_j = \beta\gamma_j.$$

The ratio estimate for the causal effect of X on Y using SNP j (out of L), $\hat{\beta}_j = \hat{\Gamma}_j/\hat{\gamma}_j$, where $\hat{\Gamma}_j$ is the estimate for SNP j's association with the outcome (with standard error σ_{Yj}) and $\hat{\gamma}_j$ is the estimate for SNP j's association with the exposure (with standard error σ_{Xj}).

The IVW estimate: The overall inverse-variance weighted (IVW) estimate for the causal effect obtained across L uncorrelated SNPs is then given by

$$\hat{\beta}_{IVW} = \frac{\sum_{j=1}^{L} w_j \hat{\beta}_j}{\sum_{j=1}^{L} w_j}$$

where w_j is the inverse-variance of $\hat{\beta}_j$. Cochran's Q statistic:

$$Q = \sum_{j=1}^{L} Q_j = \sum_{j=1}^{L} w_j (\hat{\beta}_j - \hat{\beta}_{IVW})^2, \tag{1}$$

can then be used to test for the presence of heterogeneity. If heterogeneity is detected, this provides evidence of horizontal pleiotropy. Two popular choices for the inverse-variance weights used to calculate the IVW estimate and Cochran's Q statistic are:

$$\text{1st order (fixed effect) weights:} \quad w_j \quad = \quad \frac{\hat{\gamma}_j^2}{\sigma_{Yj}^2}$$

$$\text{2nd order (fixed effect) weights:} \quad w_j \quad = \quad \left(\frac{\sigma_{Yj}^2}{\hat{\gamma}_j^2} + \frac{\hat{\Gamma}_j^2 \sigma_{Xj}^2}{\hat{\gamma}_j^4} \right)^{-1}$$

In the two-sample setting, 2nd order weights are simplified because it is not necessary to include terms involving the covariance of $\hat{\gamma}_j$ and $\hat{\Gamma}_j$, since they are obtained from independent samples. For a more detailed description of the assumptions required by two-sample summary data MR, see Bowden et al. [7].

Box 2: Two-sample summary data MR using a generalized Q statistic: fixed effect model

We start by writing down two models: firstly the underlying data generating model for the SNP-outcome association estimates under the assumption of no pleiotropy, which is a function of the causal effect and the true SNP-exposure association; and secondly the model that we actually *fit* to the data, which is a function of the causal effect and the SNP-exposure association *estimate*:

$$\text{Underlying model:} \quad \hat{\Gamma}_j \quad = \quad \beta\gamma_j + \sigma_{Yj}\epsilon_j, \tag{2}$$

$$\text{Fitted model given (2):} \quad \hat{\Gamma}_j \quad = \quad \beta\hat{\gamma}_j + \sqrt{\beta^2\sigma_{Xj}^2 + \sigma_{Yj}^2}\epsilon_j'. \tag{3}$$

Note that the variance of the error term in the fitted model has been inflated by a factor of $\beta^2\sigma_{Xj}^2$ by virtue of replacing γ_j with its estimate in (3). Dividing both sides of the fitted model through by $\hat{\gamma}_j$ we can obtain a model for the jth ratio estimate, and from that an expression for its variance:

$$\hat{\beta}_j = \beta + \sqrt{\frac{\beta^2\sigma_{Xj}^2 + \sigma_{Yj}^2}{\hat{\gamma}_j^2}}\epsilon_j' \quad \Rightarrow Var(\hat{\beta}_j) = \frac{\beta^2\sigma_{Xj}^2 + \sigma_{Yj}^2}{\hat{\gamma}_j^2}. \tag{4}$$

The variance term $Var(\hat{\beta}_j)$ in (4) is a function of the true causal effect β. Let its reciprical inverse-variance weight be denoted as $w_j(\beta) = 1/Var(\hat{\beta}_j)$. Using this weight, we now define the following modfied Q statistic:

$$Q_m(w(\beta),\beta) = \sum_{j=1}^{L} w_j(\beta)(\hat{\beta}_j - \beta)^2, \tag{5}$$

which can be used to explain all of the methods introduced in this paper.

Firstly, it is easy to see that the IVW estimate obtained using 1st order weights is the value of β that minimises $Q_m(w(0),\beta)$. Secondly, it can be shown that the IVW estimate obtained using 2nd order weights is the value that minimises $Q_m(w(\hat{\beta}_j),\beta)$. We now introduce two new IVW estimates as direct modifications of equation (5).

The 'iterative' IVW estimate

Let $\hat{\beta}_{IVW0}$ be the IVW estimate obtained using 1st order weights. That is, the value which minimises $Q_m(w(0),\beta)$. Now define $\hat{\beta}_{IVW1}$ as the value which minimises $Q_m(w(\hat{\beta}_{IVW0}),\beta)$. Lastly, define the $\hat{\beta}_{IVWi}$ as the value which minimises $Q_m(w(\hat{\beta}_{IVW(i-1)}),\beta)$. We call $\hat{\beta}_{IVWi}$ the ith 'iterative' IVW estimate.

The 'exact' IVW estimate

The exact IVW estimate is that obtained from directly minimising the generalised Q statistic, $Q_m(w(\beta),\beta)$, where in addition the weight term is allowed to be a proper function of β.

Extension to random effects models

In Appendix 1 of *Online Supplementary Material* we describe in detail how to generalise the modified Q statistic in formula (5) to obtain point estimates, standard errors and confidence intervals for the 1st order, 2nd order, iterative and exact IVW estimate allowing for heterogeneity due to pleiotropy.

Mean	1st order w_j		2nd order w_j		Modified w_j			
					Iterative		Exact	
F	Q	T1E(Q)	Q	T1E(Q)	Q	T1E(Q)	Q	T1E(Q)
			No heterogeneity, $\beta{=}0$					
100	23.9	0.044	22.8	0.022	23.9	0.044	23.9	0.044
61	24.1	0.052	21.9	0.016	24.1	0.051	24.1	0.051
40	23.9	0.049	20.3	0.006	23.9	0.048	23.9	0.048
25	24.0	0.052	17.7	0.002	23.9	0.051	23.9	0.051
10	24.0	0.052	12.3	0.000	23.6	0.047	23.4	0.042
			No heterogeneity, $\beta{=}0.05$					
100	24.2	0.053	22.9	0.028	24.0	0.049	24.0	0.049
61	24.4	0.058	21.9	0.017	24.0	0.051	24.0	0.051
40	24.7	0.064	20.3	0.007	23.9	0.050	23.9	0.049
25	25.9	0.092	17.8	0.002	24.1	0.052	23.9	0.048
10	31.4	0.272	13.4	0.000	25.6	0.095	23.7	0.043
			No heterogeneity, $\beta{=}0.1$					
100	24.7	0.065	22.8	0.027	23.9	0.052	23.9	0.051
61	25.6	0.084	21.8	0.017	23.9	0.048	23.9	0.047
40	27.3	0.132	20.5	0.009	24.1	0.053	24.0	0.050
25	31.7	0.282	18.2	0.003	24.4	0.060	23.9	0.048
10	53.9	0.792	15.8	0.004	27.8	0.166	23.9	0.051

Table 1: *Mean Q statistic and type I error rate (T1E) of 1st order, 2nd order, iterative (four iterations were performed) and exact weighting. Results are the average of 10,000 simulated data sets. Type I error rate (T1E(Q)) refers to the proportion of times Q is greater than the upper 95th percentile of a χ^2_{24} distribution.*

Figure 1: *Distribution of Q statistics (with 25 degrees of freedom) using 1st order, 2nd order and exact weights. The causal effect $\beta{=}0.1$ and the mean F statistic equals 100 (left) and 10 (right) respectively.*

Figure 2: *Left: Power of Cochran's Q statistic to detect heterogeneity as a function of the pleiotropy variance and number of SNPs (L) using 1st order, 2nd order and exact weights. Pleiotropy is simulated under a multiplicative random effects model. The causal effect is equal to 0.05 and the mean F-statistic is 61. Right: Equivalent power plot except the causal effect is equal to 0.1 and the mean F-statistic is 25.*

Mean	1st order w_j	2nd order w_j	Modified w_j		
			Iterative	Exact	
F	$\hat{\beta}_{IVW}$(SE); CF	$\hat{\beta}_{IVW}$(SE); CF	$\hat{\beta}_{IVW}$(SE); CF	$\hat{\beta}_{IVW}$(SE);CF_1	CF_2
		No Heterogeneity, $\beta=0$			
100	0.000 (0.011); 0.952	0.000 (0.011); 0.951	0.000 (0.011); 0.952	0.000 (0.011); 0.961	0.948
61	0.000 (0.011); 0.947	0.000 (0.011); 0.947	0.000 (0.011); 0.948	0.000 (0.011); 0.956	0.946
40	0.000 (0.011); 0.954	0.000 (0.010); 0.952	0.000 (0.011); 0.955	0.000 (0.011); 0.957	0.946
25	0.000 (0.011); 0.947	0.000 (0.010); 0.941	0.000 (0.011); 0.949	0.000 (0.011); 0.942	0.949
10	0.000 (0.009); 0.952	0.000 (0.007); 0.928	0.000 (0.009); 0.958	0.000 (0.010); 0.836	0.958
		No Heterogeneity, $\beta=0.05$			
100	0.049 (0.011); 0.952	0.049 (0.011); 0.951	0.049 (0.011); 0.954	0.050 (0.011); 0.962	0.952
61	0.049 (0.011); 0.948	0.047 (0.011); 0.944	0.049 (0.011); 0.952	0.050 (0.011); 0.961	0.953
40	0.048 (0.011); 0.939	0.045 (0.011); 0.918	0.048 (0.011); 0.943	0.050 (0.012); 0.951	0.946
25	0.046 (0.011); 0.910	0.041 (0.010); 0.819	0.046 (0.011); 0.923	0.050 (0.012); 0.940	0.954
10	0.033 (0.010); 0.589	0.027 (0.008); 0.286	0.034 (0.011); 0.670	0.051 (0.012); 0.868	0.957
		No Heterogeneity, $\beta=0.1$			
100	0.099 (0.011); 0.945	0.097 (0.011); 0.945	0.099 (0.012); 0.950	0.100 (0.012); 0.963	0.946
61	0.098 (0.011); 0.932	0.095 (0.011); 0.920	0.098 (0.012); 0.944	0.100 (0.012); 0.956	0.947
40	0.097 (0.012); 0.911	0.091 (0.011); 0.859	0.097 (0.012); 0.933	0.100 (0.013); 0.954	0.951
25	0.092 (0.012); 0.844	0.083 (0.011); 0.649	0.092 (0.013); 0.896	0.101 (0.014); 0.947	0.955
10	0.065 (0.013); 0.348	0.055 (0.010); 0.094	0.072 (0.014); 0.518	0.102 (0.016); 0.895	0.964

Table 2: *Mean causal estimate $\hat{\beta}_{IVW}$, standard error (SE) and coverage frequency (CF) of the 95% confidence interval when using 1st order, 2nd order, iterative and exact weights. Number of variants L=25. CF_1 = coverage of a symmetric 95% confidence interval, CF_2 = coverage of inverted Q statistic confidence interval.*

| Mean | 1st order w_j | 2nd order w_j | Modified w_j | | |
| | | | Iterative | Exact | |
F	$\hat{\beta}_{IVW}$(SE); CF	$\hat{\beta}_{IVW}$(SE); CF	$\hat{\beta}_{IVW}$(SE); CF	$\hat{\beta}_{IVW}$(SE);CF	$\hat{\phi}$
		Heterogeneity, $\beta=0$			
100	0.000(0.016); 0.949	0.000 (0.015); 0.950	0.000 (0.016); 0.950	0.000 (0.016); 0.939	2.000
61	0.000 (0.016); 0.950	0.000 (0.015); 0.951	0.000 (0.016); 0.951	0.000 (0.016); 0.940	2.004
40	0.000 (0.016); 0.953	0.000 (0.014); 0.951	0.000 (0.016); 0.955	0.000 (0.016); 0.944	1.999
25	0.000 (0.015); 0.949	0.000 (0.013); 0.945	0.000 (0.015); 0.954	0.000 (0.017); 0.945	2.003
10	0.000 (0.013); 0.952	0.000 (0.009); 0.924	0.000 (0.013); 0.960	0.000 (0.037); 0.970	1.943
		Heterogeneity, $\beta=0.05$			
100	0.050 (0.016); 0.948	0.048 (0.015); 0.947	0.050 (0.016); 0.949	0.050 (0.016); 0.938	2.002
62	0.049 (0.016); 0.951	0.046 (0.015); 0.943	0.049 (0.016); 0.954	0.050 (0.016); 0.943	1.998
40	0.048 (0.016); 0.949	0.044 (0.014); 0.924	0.048 (0.016); 0.953	0.050 (0.017); 0.943	1.995
25	0.046 (0.015); 0.933	0.039 (0.013); 0.839	0.046 (0.016); 0.940	0.051 (0.018); 0.944	1.987
10	0.033 (0.014); 0.719	0.025 (0.010); 0.378	0.034 (0.015); 0.778	0.051 (0.037); 0.960	1.967
		Heterogeneity, $\beta=0.1$			
100	0.099 (0.016); 0.947	0.096 (0.016); 0.942	0.099 (0.016); 0.952	0.100 (0.016); 0.942	2.005
61	0.098 (0.016); 0.941	0.092 (0.016); 0.922	0.098 (0.017); 0.951	0.100 (0.017); 0.941	2.004
40	0.097 (0.016); 0.932	0.088 (0.015); 0.862	0.097 (0.017); 0.947	0.101 (0.017); 0.940	2.003
25	0.092 (0.016); 0.888	0.078 (0.015); 0.676	0.093 (0.018); 0.924	0.101 (0.019); 0.942	2.003
10	0.065 (0.016);0.456	0.051 (0.012); 0.131	0.072 (0.018); 0.639	0.101 (0.042); 0.956	2.023

Table 3: *Mean causal estimate $\hat{\beta}_{IVW}$, standard error (SE) and coverage frequency (CF) of the 95% confidence interval when using 1st order, 2nd order, iterative and exact weights. L=25. $\hat{\phi}$ equals the variance inflation factor estimate (true value = 2).*

Figure 3: *Top: Scatter plot of SNP-outcome associations $\hat{\Gamma}_j$ versus SNP-exposure associations $\hat{\gamma}_j$. IVW estimate shown as a black slope. Bottom-left: Q contribution plots for the same data. Bottom-right: Q contributions after removal of rs17249754.*

Method (weights)	Estimate	S.E.	P-value	Het. Stat (p)	$\hat{\phi}$
		All 26 SNPs			
		Causal estimate			
IVW (1st -R.E)	$\hat{\beta}_{IVW}$: 0.053	0.010	3.01×10^{-5}	$Q = 67.1$ (1.03×10^{-5})	2.68
IVW (2nd -R.E)	$\hat{\beta}_{IVW}$: 0.050	0.010	4.54×10^{-5}	$Q = 58.8$ (1.54×10^{-4})	2.35
IVW (Iterative -R.E)	$\hat{\beta}_{IVW}$: 0.054	0.010	2.40×10^{-5}	$Q = 62.7$ (4.43×10^{-5})	2.51
IVW (Exact - F.E)	$\hat{\beta}_{IVW}$: 0.057	0.007	4.87×10^{-9}	$Q = 62.4$ (4.84×10^{-5})	1
IVW (Exact -R.E)	$\hat{\beta}_{IVW}$: 0.054	0.014	4.60×10^{-4}	$Q = 62.4$ (4.84×10^{-5})	2.61
		Weighted median (1st order weights)			
Weighted Median	$\hat{\beta}_{WM}$: 0.063	0.011	4.90×10^{-6}	-	-
		SNP rs17249754 removed			
		Causal estimate			
IVW (1st)	$\hat{\beta}_{IVW}$: 0.066	0.008	2.63×10^{-8}	$Q = 35.0$ (0.068)	1.46
IVW (2nd)	$\hat{\beta}_{IVW}$: 0.063	0.008	4.06×10^{-8}	$Q = 30.6$ (0.164)	1.27
IVW (Iterative)	$\hat{\beta}_{IVW}$: 0.066	0.008	2.90×10^{-8}	$Q = 32.8$ (0.107)	1.37
IVW (Exact - FE)	$\hat{\beta}_{IVW}$: 0.068	0.007	1.07×10^{-9}	$Q = 32.8$ (0.108)	1
IVW (Exact - RE)	$\hat{\beta}_{IVW}$: 0.067	0.009	8.37×10^{-8}	$Q = 32.8$ (0.108)	1.39
		Weighted median (1st order weights)			
Weighted Median	$\hat{\beta}_{WM}$: 0.065	0.010	1.81×10^{-6}	-	

Table 4: *IVW and Weighted Median analyses of the causal effect of SBP on CHD risk using 1st order, 2nd order, iterative and exact weights for the complete data (top) and with SNP rs17249754 removed (bottom). $\hat{\beta}_{IVW}$ is the IVW estimate. $\hat{\beta}_{WM}$ is the Weighted Median estimate. 1st order, 2nd order and iterative IVW estimates fitted under a multiplicative random effects model (R.E), where $\hat{\phi}$ refers to the variance inflation factor estimate. Exact IVW estimate fitted under both fixed (F.E) and multiplicative random effects models, because their estimates are distinct.*