Reverse immunodynamics: a new method for identifying targets of protective immunity

José Lourenço[1*], Katrina J Spensley[1*], Paul S Wikramaratna[2*], Bridget S Penman[1], Andrew Walker[1], Adrian L Smith[1], Oliver G Pybus[1], Létitia Jean[3] & Sunetra Gupta[1**]

1. Department of Zoology, University of Oxford, Oxford OX1 3PS
2. Institute of Evolutionary Biology, University of Edinburgh, Edinburgh EH9 3JT
3. The Sir William Dunn School of Pathology, South Parks Road, Oxford OX1 3RE

*these authors contributed equally to the manuscript
**Correspondence to: sunetra.gupta@zoo.ox.ac.uk (SG)

Abstract

Despite a dramatic increase in our ability to catalogue variation among pathogen genomes, we have made far fewer advances in using this information to identify targets of protective immunity. We propose a novel methodology that combines predictions from epidemiological models with phylogenetic and structural analyses to identify such targets. Mathematical models predict that strong immune selection can cause antigenic variants to exist in non-overlapping combinations. A corollary of this theory is that targets of immunity may be identified by searching for non-overlapping associations among antigenic variants. We applied this concept to the AMA-1 protein of the malaria parasite *Plasmodium falciparum* and found strong signatures of immune selection among certain regions of low variability which could render them ideal vaccine candidates.

Apical membrane antigen 1 (AMA-1) is a trans-membrane protein common to all

Plasmodium species with a putative role in invasion at erythrocytic and pre-

erythrocytic stages of the parasite life-cycle. Antibodies to AMA-1 appear to inhibit

erythrocyte invasion and are thus thought to play a role in clinical protection against

both *P. falciparum* [1-7] and *P. vivax* [8]. Cellular responses to AMA-1 may provide

immunity against both erythrocytic [9] and pre-erythrocytic forms of *P. falciparum*

[10]. The potential of AMA-1 as a component of a multi-stage malaria vaccine is

compromised by the polymorphism of many of the targets of natural immunity [11,12]

as well as by difficulties in inducing lasting immune responses through vaccination

[13-16]. Identifying variable regions of AMA-1 that are under strong immune

selection would afford the possibility of focusing vaccine-induced immune responses

on these regions and ensure full coverage of variability at these critical epitopes.

The extracellular region of AMA-1 can be structurally resolved into three domains:

DI, DII and DIII. The highest level of polymorphism is seen among residues in DI

and DII surrounding a conserved hydrophobic trough [17,18] which plays a vital role

in the invasion process [19-21]. Antibodies against this region have been shown to

block invasion in a strain-specific manner; however, it is also clear that these

antibodies constitute only a small proportion of the total inhibitory humoral response

[12,22,23]. *Pf* AMA1 also contains a number of less variable residues across the three

domains, and our aim was to identify which of these were also located within epitopes

eliciting protective immune responses.

We attempted to identify these additional epitopes under strong immune selection by

using a novel approach based on the predictions of a multilocus mathematical model

of pathogen evolution [24-26]. The central premise is that regions or epitopes under

strong immune selection will self-organise into non-overlapping combinations of

variants [24-26] and supported by empirical observations within a variety of host-

pathogen systems [27-32]. We inverted this result to search for signatures of immune

selection among polymorphic loci by identifying those pairs which exhibit a high

degree of 'non-overlap' but excluding those that may have arisen as a consequence of

shared ancestry or due to structural interactions. Using this "reverse immunodynamic"

method, we found strong signatures of immune selection among a small number of

dimorphic residues in Domain III of *Pf*AMA-1, suggesting that epitopes containing

these residues are targets of protective immunity.

We implemented the reverse immunodynamic method by performing pairwise

comparisons of all dimorphic sites among 1198 unique *Pf*AMA-1 sequences obtained

from GenBank, to determine whether any of these pairs exhibited non-overlapping

associations between amino acid residues. We used an Information Theory statistic

known as Mutual Information (MI), scaled to its maximum value, to measure the

degree of non-overlap in amino acid combinations between pairs of sites. Most pairs

of sites showed low non-overlap, with only 10 pairs exhibiting a scaled MI in excess

of 0.3 (Tables 1 & S1). Of these, pairs 404/405, 448/451 and 283/285 were

eliminated on grounds of structural proximity, leaving 7 pairs as shown by the bars in

Fig 1A.

To assess whether high non-overlap between sites could have arisen by neutral

evolutionary processes, we estimated a phylogenetic tree from the sequences and

calculated the parsimony score (PS) for each site, which indicates the minimum

number of mutations (in this case, back and forth between the two possible amino

acids at each site) required to explain the observed ancestral relationships. A new

statistic S, the product of MI and the geometric mean of PS for each site, was found to

preserve the same order as the scaled MI scores of the 7 sites under consideration,

except in the case of the 206/225 (Table S2). To obtain an appropriate null

distribution of S for hypothesis testing, we simulated 1000 sets of sequences which

were (a) consistent with the empirical phylogenetic tree and (b) had the same PS score

at each dimorphic site as the empirical data. The null distribution of the top-scoring

measures of S (Fig 1C) shows that neutral evolution can generate sites with high MI

or high PS values, but not both. In contrast, the pairs of sites indicated in red in Fig

1A lie well outside the null distribution of high-scoring site-pairs (Fig 1C). Only site-

pair 206/225 (shown in green) did not meet this criterion. Reasons for this pattern

become evident upon inspection of the distribution of the amino acid variation on the

phylogenetic tree (Fig 2). The dominant non-overlapping pairs of amino acids at

206/225 are highly segregated on the phylogeny and not improbable under the null

hypothesis, while those of 451/485 exhibit a more interdigitated pattern, indicating

multiple independent and convergent paired amino acid changes that are

exceptionally unlikely in the absence of immune selection.

Figure 1: (A) Degree of non-overlap in associations between amino acid residues at pairs of dimorphic sites. Each site on the x-axis is compared with all other dimorphic sites: those with reciprocally high scores are shown by bars spanning both sites. Red bars show significant associations. Green bars represent associations that do not achieve significance when phylogenetic relationships are taken into account (B) Surface representation of the crystal structure of AMA-1 with specific amino acids highlighted [Lysine (Lys) 206 = Cyan; Isoleucine (Ile) 225 = Magenta; Asparagine (Asn) 439 = Green; Methionine (Met) 451 = Red, Lys 485 = Blue, Met 496 = Orange, Arginine (Arg) 503 = Yellow]. (C) Pairs of sites shown by red and green bars in Fig1A are shown in relation to a null distribution of site-pairs from 1000 simulations that scored most highly on the product of scaled Mutual Information (measuring non-overlap) and Geometric Mean of Parsimony Scores (indicating the minimum number of mutations required to explain the observed ancestral relationships). (D) Phylogenetic tree of AMA-1 with outside ring coloured on the basis of observed combination of amino acid variants found at tree tips for (A) sites 206 and 225 (pink = NE, blue = IK, orange = IE), (B) sites 451 and 485 (pink = IK, blue = KM, orange = KK), and (C) sites 496 and 503 (pink = NI, blue = RM, grey = rare variants).

The group of sites which scored most highly on the conservative test (described above)

comprised 496, 503 and 512. However, mapping these onto the tertiary structure of

AMA-1 (Fig 1B) shows that they are in very close proximity of each other and

therefore we were unable to exclude biochemical interactions as the primary cause of

this striking association. It is still possible that variation at these sites is held in strong

non-overlap through immune selection. Antibodies specific to epitopes containing

amino acids I/N/K may cross-react with all other epitopic variants except those

containing M/R/R, thereby promoting the co-circulation of these non-overlapping

combinations of residues at these sites; however, because of their structural adjacency,

it is impossible to privilege this explanation over intrinsic steric advantages to these

combinations

High non-overlap was also observed among sites 439/451, 439/485 and

451/485 which were sufficiently far enough apart in the tertiary structure to represent

potential examples of immune selection forcing non-structurally related residues to

co-segregate. We cannot, of course, completely eliminate long-range physical

interactions as a cause of these associations, and whether it is indeed immune

selection that is responsible for these patterns can only be established by functional

studies. It is clear that there is no fundamental structural impediment to the existence

of intermediate forms since they do occur at low frequencies, but whether they are

disadvantaged due to lower intrinsic fitness or immune-mediated selection remains to

be resolved by further experimental work.

We next investigated the associations between positions 439, 451 and 485 on a

smaller geographical scale to confirm that these patterns did not arise due to certain

combinations of alleles dominating by chance in particular areas. In all countries

studied, both of the major sequence variant associations were present, with the

secondary variants present at a lower frequency (Table 2).

A dataset of 660 unique sequences was compiled for Mali and showed high

non-overlap for each pair of associations. A smaller dataset comprising 79 sequences

from the Gambia also yielded MI values in excess of those obtained from the global

data set. Slightly weaker associations were found in 86 unique sequences from Kenya;

however these improved significantly when the data were analysed as a collection of

isolates rather than sequences: i.e. without excluding identical sequences present in

different individuals (Table 3).

As the model predictions pertain to population level frequencies of allele

combinations, rather than their presence in a collection of unique sequences, we

believe that Table 3 presents a more valid test (than when only unique sequences are

included) of the model's capabilities to pick out regions under strong immune

selection; it is reassuring that these patterns are also evident under the more stringent

conditions where only unique sequences are included.

Our results suggest that Domain III (DIII) of the AMA-1 protein is under strong

immune selection, and may play an important role in protective immunity alongside

the more polymorphic C1L region of Domain I. At a functional level, Domain III has

been shown to bind to the erythrocyte membrane protein Kx [33]. Immune responses

that interrupt this process may, therefore, have a significant inhibitory effect on

parasite growth but, so far, the evidence that Domain III elicits protective antibody

responses is limited. Nair *et al.* [34] have shown that antibodies from a single blood

donor affinity purified on DIII are inhibitory. In a study by Polley *et al.* [2], it was

found that antibody responses against recombinant full-length ectodomains were more

strongly associated with protection than responses to those which did not express DIII,

even though naturally acquired antibodies to recombinant DIII alone were rare. A

virosomal formulation of loop I of DIII, comprising residues 446-490, has been

shown to elicit growth-inhibitory antibodies [35] and volunteers within a vaccine trial

containing this AMA1 peptide exhibited both reduced blood stage growth rates and a

higher rate of morphological parasite abnormalities [36]. A recent analysis of allele-

specific efficacy of a vaccine against AMA1 [11] showed that the relative risk ratio

(RRR) of re-infection with strains containing vaccine-type amino acids among

vaccinees vs controls was high (although not significant) at residues 439, 451 and 485.

Residues associated with the polymorphic sites surrounding the hydrophobic trough

displayed significant RRR, as might be expected; sites 308 and 332 also had high

scores but no evidence of sustained immune selection at these sites was available in

our analysis. In the same study, the likelihood of a shift from a vaccine-type to non-

vaccine-type residue among sequences obtained following vaccination was found to

be significantly increased at position 485 and non-significantly increased at position

451. Significant changes in residue composition were also found at highly

polymorphic site 197 but not at any of the other polymorpic sites in the C1L region.

Significant change was also observed at dimorphic site 435, but we found no non-

overlapping associations in any of the pairwise comparisons between 435 and other

sites. Several population genetic studies [37-41] based on site frequency spectrum

analysis techniques (such as Tajima's D) strongly indicate that DIII is under

balancing selection; our "reverse immunodynamic" method complements and extends

these analyses by showing that we can use a measure of non-overlap between site-

pairs in addition to standard population genetic measures to finely map regions under

immune selection within DIII.

An interesting feature of our analyses is that while particular groups of sites exhibit

strong non-overlap, the associations between these groups appear to be random. For

example, the triad 439-451-485 is in complete linkage equilibrium with triad 496-

503-512. Furthermore, there are no non-overlapping associations present between the

highly polymorphic residues in the C1L region of Domain I and the 439-451-485 triad

(although the former exhibit strong non-overlap: Table S1). If the analysis we present

here is correct, then it would be sufficient to present the dominant "non-overlapping"

variants of 439-451-485 (i.e. HKI and NMK) in a manner that induced strong immune

responses to the epitopes containing these residues, rather than to try and cover the

entire haplotypic diversity of AMA-1 [42]. Immune responses to the epitopes

containing residues 439, 451 and 485 appear to develop more slowly under natural

conditions than antibodies to the highly polymorphic regions shielding the

hydrophobic groove [2], but should be easier to induce by vaccination than immune

responses to completely conserved epitopes [43].

Our analysis of AMA-1 highlights the role of less polymorphic epitopes in

population-level immune evasion. At a more general level, our analyses emphasize

that targets of natural immunity are not polarised between the completely invariant

and the highly polymorphic but also include a range of epitopes of intermediate

variability that have, so far, been largely overlooked. Among pathogens where

protective immune responses are directed against conserved targets or epitopes, a

single infection ensures lifelong immunity and the goal of vaccination is to elicit

natural immunity against such conserved targets. Most infectious diseases (such as

measles) against which we have effective vaccines fall into this category. By contrast,

the primary targets of protective immunity in several important pathogens such as *P.*

falciparum, are highly diverse and present a challenge to the development of vaccines.

One solution to this problem that is currently being explored by several research

groups is the artificial boosting of immunity towards conserved regions that are not

sufficiently naturally immunogenic to prevent further infection upon single exposure

but do eventually confer protection after multiple exposures. Another strategy is to

develop multi-allelic vaccines in which immunogenic epitopes that confer protection

against a sub-population of strains are combined to provide broad coverage. However,

many of the known targets of protective immunity are too diverse for this strategy to

be successfully implemented. Our work in this paper highlights that between these

two extremes, there exist epitopes of intermediate variability which are nonetheless

potentially under strong immune selection and thus offer alternative targets of

vaccination.

Methods

229
SEQUENCE SELECTION: *Pf*AMA-1 sequences were obtained by searching the

NCBI Entrez nucleotide and protein databases. On 04/07/2014 this yielded 2835

protein sequences and 2926 nucleotide sequences. In cases when the GenBank

nucleotide record did not include the corresponding protein sequence, these were

translated using Biopython, an open source set of tools (biopython.org).

The analysis was restricted to sequences between 400 and 622 amino acids in length.

The lower limit aims to ensure that all the polymorphic regions across the three

domains are included; the upper limit corresponds to the entire processed protein, and

therefore removes any entries which contain the non-coding sequence. To avoid

pseudoduplication, all identical sequences were removed, leaving 1198 unique

sequences. These were then aligned using Clustal W [40].

MEASURING NON-OVERLAP AMONG DIMORPHIC SITES: A frequency

analysis of all residues within the three domains (q.v.) was used to identify any

polymorphic sites. Only variants with a frequency exceeding 2.5% were included in

the subsequent analysis. This identified 44 dimorphic sites (Fig S1). We searched for

signatures of immune selection by analysing the mutual information (MI) scores of all

pairs of sites (Table S1).

GEOGRAPHIC SUB-ANALYSES: We repeated the calculation of MI for all

isolates whose country of origin could be identified (either from the Entrez database

or with reference to published papers).

PHYLOGENETIC ANALYSIS: We first truncated our protein alignment to the

region where we had >99% coverage [namely, from residue 149 to residue 534]. This

alignment was then used to construct a Maximum Likelihood Tree in Mega6 under

the JTT model (with also: 4 discrete categories of gamma-distributed rates among

sites; the nearest-neighbour-interchange heuristic; and a very strong branch swap

filter) and this tree was used to calculate the maximum parsimony score (PS) for each

dimorphic site (Fig S2; Table S1). We defined a new statistic S as the product of the

MI and the geometric mean of the PS for each site (Table S2). We used Seq-Gen to

simulate 1000 sets of protein sequences with the same phylogenetic characteristics,

restricting variation at each of the sites to two amino acids. We were able to establish

that the pairs of interest (see Table 1) lay significantly outside the distribution of the

identically ranked (by S-score) sites from each simulation (Fig S3).

IMAGES: From the protein data bank (PDB) the sequences and tertiary structures of

Plasmodium vivax AMA-1 (*Pv*AMA-1, PDB 1W81) and *Pf*AMA-1 domain III (PDB

1HN6) were downloaded. As the tertiary structure of 1HN6 was mostly unstructured,

it could not be used on its own. Therefore a Fold and Function Assignment System

(FFAS) was used to align the amino acid sequences of 1W81 and 1HN6. From this

alignment a 'hybrid' structural model with 94% confidence was generated using

SCWRL. This resulting model gave a single tertiary structure with the residues within

domain III from *Pv*AMA-1 (1W81) substituted by those from *Pf*AMA-1 (1HN6)

using the *P. vivax* structural coordinates. Pymol (http://www.pymol.org/) was used to

visualise the 'hybrid' tertiary structure and to highlight the relevant residues.

Supplementary files

279   Table S1: parsimony score and MI for pairs of sites
280   Table S2: statistic S (product of MI and the geometric mean of the PS for each site)

Acknowledgements

282   We thank Dr. CRC Spensley for assistance with data manipulation and programming,

283   and Dr. Pietro Roversi for assistance with generating the structure.

284
285

References

1. Remarque EJ, Faber BW, Kocken CHM & Thomas AW (2008) Apical membrane antigen 1: A malaria vaccine candidate in review. *Trends Parasitol* 24(2): 74-84.
2. Polley SD, et al (2004) Human antibodies to recombinant protein constructs of plasmodium falciparum apical membrane antigen 1 (AMA1) and their associations with protection from malaria. *Vaccine* 23(5): 718-728.
3. Fowkes FJI, Richards JS, Simpson JA & Beeson JG (2010) The relationship between anti-merozoite antibodies and incidence of plasmodium falciparum malaria: A systematic review and meta-analysis. *Plos Medicine* 7(1): e1000218.
4. Richards JS, et al (2013) Identification and prioritization of merozoite antigens as targets of protective human immunity to plasmodium falciparum malaria for vaccine and biomarker development. *Journal of Immunology* 191(2): 795-809.
5. Mugyenyi CK, et al (2013) Antibodies to polymorphic invasion-inhibitory and non-inhibitory epitopes of plasmodium falciparum apical membrane antigen 1 in human malaria. *Plos One* 8(7): e68304.
6. Stanisic DI, et al (2015) Acquisition of antibodies against plasmodium falciparum merozoites and malaria immunity in young children and the influence of age, force of infection, and magnitude of response. *Infect Immun* 83(2): 646-660.
7. Osier FH, et al (2014) New antigens for a multicomponent blood-stage malaria vaccine. *Science Translational Medicine* 6(247)
8. Vicentin EC, et al (2014) Invasion-inhibitory antibodies elicited by immunization with plasmodium vivax apical membrane antigen-1 expressed in pichia pastoris yeast. *Infect Immun* 82(3): 1296-1307.
9. Biswas S, et al (2012) Recombinant viral-vectored vaccines expressing plasmodium chabaudi AS apical membrane antigen 1: Mechanisms of vaccine-induced blood-stage protection. *Journal of Immunology* 188(10): 5041-5053.
10. Schussek S, et al (2013) Immunization with apical membrane antigen 1 confers sterile infection-blocking immunity against plasmodium sporozoite challenge in a rodent model. *Infect Immun* 81(10): 3586-3599.
11. Ouattara A, et al (2013) Molecular basis of allele-specific efficacy of a blood-stage malaria vaccine: Vaccine development implications. *J Infect Dis* 207(3): 511-519.
12. Drew DR, et al (2012) Defining the antigenic diversity of plasmodium falciparum apical membrane antigen 1 and the requirements for a multi-allele vaccine against malaria. *Plos One* 7(12): e51023.
13. Dicko A, et al (2007) Impact of a plasmodium falciparum AMA1 vaccine on antibody responses in adult malians. *Plos One* 2(10): e1045.
14. Thera MA, et al (2008) Safety and immunogenicity of an AMA-1 malaria vaccine in malian adults: Results of a phase 1 randomized controlled trial. *Plos One* 3(1): e1465.
15. Laurens MB, et al (2013) Extended safety, immunogenicity and efficacy of a blood-stage malaria vaccine in malian children: 24-month follow-up of a randomized, double-blinded phase 2 trial. *Plos One* 8(11): e79323
16. Biswas S, et al (2014) Assessment of humoral immune responses to blood-stage malaria antigens following ChAd63-MVA immunization, controlled human malaria infection and natural exposure. *Plos One* 9(9): e107903.
17. Bai T, et al (2005) Structure of AMA1 from plasmodium falciparum reveals a clustering of polymorphisms that surround a conserved hydrophobic pocket. *Proc Natl Acad Sci U S A* 102(36): 12736-12741.
18. Lim SS, et al (2014) Structure and dynamics of apical membrane antigen 1 from plasmodium falciparum FVO. Biochemistry (N Y) 53(46): 7310-7320.

19. Richard D, et al (2010) Interaction between plasmodium falciparum apical membrane antigen 1 and the rhoptry neck protein complex defines a key step in the erythrocyte invasion process of malaria parasites. *J Biol Chem* 285(19): 14815-14822.

20. Lamarque M, et al (2011) The RON2-AMA1 interaction is a critical step in moving junction-dependent invasion by apicomplexan parasites. *Plos Pathogens* 7(2): e1001276.

21. Srinivasan P, et al (2011) Binding of plasmodium merozoite proteins RON2 and AMA1 triggers commitment to invasion. *Proc Natl Acad Sci U S A* 108(32): 13275-13280.

22. Healer J, et al (2004) Allelic polymorphisms in apical membrane antigen-1 are responsible for evasion of antibody-mediated inhibition in plasmodium falciparum. *Mol Microbiol* 52(1): 159-168.

23. Cortes A, et al (2005) Allele specificity of naturally acquired antibody responses against plasmodium falciparum apical membrane antigen 1. *Infect Immun* 73(1): 422-430.

24. Gupta S, et al (1996) The maintenance of strain structure in populations of recombining infectious agents. *Nat Med* 2(4): 437-442.

25. Gupta S, Ferguson N & Anderson R (1998) Chaos, persistence, and evolution of strain structure in antigenically diverse infectious agents. *Science* 280(5365): 912-915.

26. Lourenço J, Wikramaratna SP, Gupta S (2015) MANTIS: an R package that simulates multilocus models of pathogen evolution. *BMC Bioinformatics*. 16:176.

27. Hinnebusch BJ, Barbour AG, Restrepo BI, Schwan TG. (1998) Population structure of the relapsing fever spirochete *Borrelia hermsii* as indicated by polymorphism of two multigene families that encode immunogenic outer surface lipoproteins. *Infect Immun* 66:432–40.

28. Johnson DR, Kaplan EL, VanGheem A, Facklam RR, Beall B (2006) Characterization of group A streptococci (Streptococcus pyogenes): Correlation of M-protein and emm-gene type with T-protein agglutination pattern and serum opacity factor. J Med Microbiol 55:157e164.

29. Areschoug, T., Stålhammar-Carlemalm, M., Larsson, C., & Lindahl, G. (1999). Group B Streptococcal Surface Proteins as Targets for Protective Antibodies: Identification of Two Novel Proteins in Strains of Serotype V. *Infection and Immunity*, *67*(12), 6350–6357.

30. Buckee CO, Gupta S, Kriz P, Maiden MC & Jolley KA (2010) Long-term evolution of antigen repertoires among carried meningococci. *Proc Biol Sci* 277:1635-41

31. Callaghan MJ, Buckee C, Jolley, KA, Kriz, P, Maiden MCJ , Gupta S. (2008) The Effect of Immune Selection on the Structure of the Meningococcal Opa Protein Repertoire. *PLoS Pathogens* 14;4(3):e1000020.

32. Polley SD, et al. (2007) Plasmodium falciparum merozoite surface protein 3 is a target of allele-specific immunity and alleles are maintained by natural selection. J Infect Dis 195:279e287.

33. Kato K, Mayer DCG, Singh S, Reid M & Miller LH (2005) Domain III of plasmodium falciparum apical membrane antigen 1 binds to the erythrocyte membrane protein kx. *Proc Natl Acad Sci U S A* 102(15): 5552-5557.

34. Nair M, et al (2002) Structure of domain III of the blood-stage malaria vaccine candidate, plasmodium falciparum apical membrane antigen 1 (AMA1). *J Mol Biol* 322(4): 741-753.

35. Mueller MS, et al (2003) Induction of parasite growth-inhibitory antibodies by a virosomal formulation of a peptidomimetic of loop I from domain III of plasmodium falciparum apical membrane antigen 1. *Infect Immun* 71(8): 4749-4758.

36. Thompson FM, et al (2008) Evidence of blood stage efficacy with a virosomal malaria vaccine in a phase IIa clinical trial. *Plos One* 3(1): e1493.

37. Polley SD, Conway DJ (2001) Strong diversifying selection on domains of the *Plasmodium falciparum* apical membrane antigen 1 gene. Genetics 158: 1505–1512.

38. Tetteh KK, Stewart LB, Ochola LI, Amambua-Ngwa A, Thomas AW, et al. (2009) Prospective identification of malaria parasite genes under balancing selection. PLoS ONE 4: e5568 doi:10.1371/journal.pone.0005568. doi: 10.1371/journal.pone.0005568

39. Osier FH, Weedall GD, Verra F, Murungi L, Tetteh KK, et al. (2010) Allelic diversity
and naturally acquired allele-specific antibody responses to *Plasmodium falciparum*
apical membrane antigen 1 in Kenya. Infect Immun 78: 4625–4633. doi:
10.1128/iai.00576-10
40. Amambua-Ngwa A, Tetteh KKA, Manske M, Gomez-Escobar N, Stewart LB, Deerhake
ME, et al. (2012) Population Genomic Scan for Candidate Signatures of Balancing
Selection to Guide Antigen Characterization in Malaria Parasites. PLoS Genet 8(11):
e1002992. doi:10.1371/journal.pgen.1002992
41. Arnott A, et al (2014) Distinct patterns of diversity, population structure and evolution in
the AMA1 genes of sympatric plasmodium falciparum and plasmodium vivax
populations of Papua New Guinea from an area of similarly high transmission. *Malaria*
Journal 13: 233.
42. Duan J, et al (2008) Population structure of the genes encoding the polymorphic
plasmodium falciparum apical membrane antigen 1: Implications for vaccine design.
Proc Natl Acad Sci U S A 105(22): 7857-7862.
43. Dutta S, et al (2013) Overcoming antigenic diversity by enhancing the immunogenicity of
conserved epitopes on the malaria vaccine candidate apical membrane antigen-1. *Plos*
Pathogens 9(12): e1003840.
44. Larkin MA, et al (2007) Clustal W and clustal X version 2.0. *Bioinformatics* 23(21):
2947-2948.
408